

A shallow water approximation for water waves

Dedicated to the late Professor Alexandre V. Kazhikhov

By

Tatsuo IGUCHI

1. Introduction

In this paper we are concerned with the initial value problem for water waves in arbitrary space dimensions. The water wave is a model system for irrotational flow of an incompressible ideal fluid with a free surface under the gravitational field. The analysis of this problem is very hard because of the nonlinearity of the equations together with the presence of an unknown free surface. In order to understand various phenomena of water waves, one has approximated the equations by simple ones and analyzed the approximated equations. The simplest approximation is the linear one around the trivial flow by assuming that the amplitude of the free surface and the motion of the fluid are infinitesimal. However, this approximation could not explain the existence of solitary waves nor the breaking of water waves. In order to explain such phenomena we have to include nonlinear effects of the waves in the approximation. The shallow water equations are one of such approximations and derived from the water wave by assuming that the water depth is sufficiently small compared to the wave length. The aim of this paper is to give a mathematically rigorous justification of the shallow water approximation for water waves in Sobolev spaces.

Rewriting the equations for water waves in an appropriate non-dimensional form, we have two non-dimensional parameters δ and ε the ratio of the water depth h to the wave length λ and the ratio of the maximum vertical amplitude of the free surface a to the water depth h , respectively, in the equations. The shallow water equations are derived from the water wave in the limit $\delta \rightarrow +0$ by keeping $\varepsilon \simeq 1$. In the case of a flat bottom, they are of the same form as the compressible Euler equation for a barotropic gas and the solution generally has a singularity in finite time even if the initial data are sufficiently smooth. Therefore, this approximation is used to explain the breaking of the waves. The derivation of the shallow water equations goes back to G. B. Airy [1]. Then, K. O. Friedrichs [6] derived systematically the equations from the water wave

2000 *Mathematics Subject Classification(s)*. 76B07, 76B15, 35Q35.

Received December 8, 2006

Revised October 30, 2008

problem by using an expansion of the solution with respect to δ^2 , which is called the Friedrichs expansion. See also H. Lamb [14] and J. J. Stoker [22]. A mathematically rigorous justification of the shallow water approximation for two-dimensional water waves was given by L. V. Ovsjannikov [18, 19] under the periodic boundary condition with respect to the horizontal spatial variable, and then by T. Kano and T. Nishida [10]. A mathematical justification of the Friedrichs expansion was investigated by T. Kano and T. Nishida [11] and the justification in the three-dimensional case by T. Kano [9]. In order to guarantee the existence of solutions for water waves, they used an abstract Cauchy-Kowalevski theorem in a scale of Banach spaces so that analyticity of the initial data was required. It is natural to ask if the approximation is valid in Sobolev spaces. However, this question was not resolved.

On the other hand, the Korteweg-de Vries (KdV) equation is also derived from the two-dimensional water wave in the limit $\varepsilon \simeq \delta^2 \rightarrow +0$. It is well-known that the solution of the KdV equation exists globally in time and the equation has solitary wave solutions. The derivation of the KdV equation goes back to D. J. Korteweg and G. de Vries [13]. A mathematically rigorous justification of the KdV equation for the water wave was investigated by T. Kano and T. Nishida [12] in a class of analytic functions. Concerning this KdV approximation, a justification in Sobolev spaces was given by W. Craig [3] under a restriction that the wave is almost one-directional. Then, G. Schneider and C. E. Wayne [20] gave a justification without assuming one-directional motion of the wave. In the case with the surface tension on the free surface, G. Schneider and C. E. Wayne [21] and the author [7] gave justifications. We also refer to W. Craig, P. Guyenne, D. P. Nicholls, and C. Sulem [4], who adopted a Hamiltonian formulation of the problem and derived systematically the KdV and the KP equations. Important parts of the analysis in [3, 7] are to approximate a non-local operator, such as the Dirichlet-to-Neumann map for Laplace's equation or the Dirichlet-to-Dirichlet map for the Cauchy-Riemann equations, in terms of Fourier multipliers by expanding it with respect to a function which represents the surface elevation and to give a precise estimate for the remainder part. However, in a shallow water regime $\delta \ll 1$ and $\varepsilon \simeq 1$ we cannot obtain a good estimate for the remainder part so that we have to use another method in order to give a justification of the shallow water approximation.

In connection with the well-posedness of the initial value problem for water waves, the solvability in Sobolev spaces was given by several authors. In his pioneering work [17], V. I. Nalimov investigated the initial value problem in the case where the motion of the fluid is two-dimensional and the fluid has infinite depth. He showed that if the initial data are sufficiently small in a Sobolev space, that is, if the initial surface is almost flat and the initial movement of the fluid is sufficiently small, then there exists a unique solution of the problem locally in time in a Sobolev space. H. Yosihara [26] extended this result to the case of presence of an almost flat bottom. S. Wu [24] studied the problem in exactly the same situation as Nalimov's and gave the existence theorem locally in time without assuming the initial data to be small. It is

known that the well-posedness of the problem may be broken unless a generalized Rayleigh-Taylor sign condition $-\partial p/\partial N \geq c_0 > 0$ on the free surface is satisfied, where p is the pressure and N is the unit outward normal to the free surface. She showed that this condition always holds for any smooth nonself-intersecting surface. S. Wu [25] also succeeded in giving an existence theory in Sobolev spaces for three-dimensional water waves of infinite depth. Note that all of the three authors mentioned above used the Lagrangian coordinates. D. Lannes [15] studied the initial value problem for water waves of finite depth in arbitrary space dimensions. One of interesting features of his paper is that he did not use the Lagrangian coordinates but the Euler coordinates although the surface tension on the free surface was neglected. Another interesting feature is that he obtained a good expression of the Fréchet derivative of the Dirichlet-to-Neumann map for Laplace's equation with respect to a function which represents the surface elevation. As a result, he derived nice linearized equations and succeeded in giving an existence theory in Sobolev spaces.

The existence theories in Sobolev spaces were based on the energy method. In calculation of the time evolution of an energy function, we need to estimate commutators of the Dirichlet-to-Neumann map and differential operators. S. Wu [25] obtained precise commutator estimates by using the theory of singular integral operators and Clifford analysis, whereas D. Lannes [15] used the theory of pseudo-differential operators and obtained commutator estimates by imposing much differentiability on the coefficients. This is one of the reasons why a Nash-Moser implicit function theorem was used to obtain the solution of the nonlinear equations in [15]. A relation between the generalized Rayleigh-Taylor sign condition and the bottom topography was also analyzed in [15]. Under a shallow water regime $\delta \ll 1$ and $\varepsilon \simeq 1$, such techniques in [25, 15] in estimating commutators do not give nice uniform estimates with respect to small δ . In this paper, to obtain the uniform estimates, we only use the standard technique in estimating the solution of a boundary value problem for elliptic differential equations, so that the proof may become much simpler and more elementary than the previous ones. We adopt the formulation of the problem used in [15]. However, thanks to a precise energy estimate for linearized equations and a quasi-linearization of the full nonlinear equations, we do not use the Nash-Moser implicit function theorem to obtain the solution of the nonlinear equations.

After writing this manuscript, the author knew results of Y. A. Li [16], where he considered a shallow water approximation for two-dimensional water waves over a flat bottom and gave a mathematical justification of the approximation by the Green-Naghdi equations in Sobolev spaces, and of B. Alvarez-Samaniego and D. Lannes [2], where they gave mathematical justifications of several asymptotic models for three-dimensional water waves including the shallow water and KP equations. However, in [2] they still used the Nash-Moser implicit function theorem, whereas we do not use the theorem in this paper.

The contents of this paper are as follows. In Section 2 we formulate the problem, rewrite it in a non-dimensional form, transform it into an equivalent problem on the free surface, and give one of our main results, which asserts the

existence of the solution with uniform bounds in a Sobolev space. In Section 3 we formally derive the shallow water equations from the water wave and give another main result, which justifies rigorously the shallow water approximation. In Section 4 we define and analyze the Dirichlet-to-Neumann map for Laplace's equation. In the analysis, we transform a boundary value problem for Laplace's equation in the fluid domain $\Omega(t)$ to a problem on the simple fixed domain $\Omega_0 = \mathbf{R}^n \times (0, 1)$ by using a suitable diffeomorphism $\Theta : \Omega_0 \rightarrow \Omega(t)$. This is one of the crucial parts of this paper. In Sections 5 and 6 we derive uniform estimates with respect to small δ in a Sobolev space for the Dirichlet-to-Neumann map and its Fréchet derivatives with respect to the function which represents the surface elevation. In Section 7, according to D. Lannes [15] we first linearize the full equations, and then we evaluate an energy function for the linearized problem uniformly with respect to small δ . In Section 8 we reduce the full nonlinear equations to quasi-linear equations. Finally, in Section 9, by applying the energy estimates established in Section 7 to the quasi-linear equations derived in Section 8 we prove main theorems.

Notation. For a real number s , we denote by H^s the Sobolev space of order s on \mathbf{R}^n equipped with the inner product $(u, v)_s = (2\pi)^{-n} \int_{\mathbf{R}^n} (1 + |\xi|)^{2s} \hat{u}(\xi) \overline{\hat{v}(\xi)} d\xi$, where $\hat{u} = \mathcal{F}[u]$ is the Fourier transform of u , that is, $\hat{u}(\xi) = \int_{\mathbf{R}^n} u(x) e^{-ix \cdot \xi} dx$. We put $\|u\|_s = \sqrt{(u, u)_s}$, $(u, v) = (u, v)_0$, and $\|u\| = \|u\|_0$. For $1 \leq p \leq \infty$, we denote by $|\cdot|_p$ the norm of the Lebesgue space $L^p = L^p(\mathbf{R}^n)$. The norm of a Banach space X is denoted by $\|\cdot\|_X$. For $0 < T < \infty$, a non-negative integer j , and a Banach space X , we denote by $C^j([0, T]; X)$ the Banach space of all functions of C^j -class on the interval $[0, T]$ with the value in X . We put $\partial_j = \partial/\partial x_j$, $\partial_{ij} = \partial_i \partial_j$, and $\partial_{ijk} = \partial_i \partial_j \partial_k$. A pseudo-differential operator $P(D)$, $D = (D_1, \dots, D_n)$ and $D_j = -i\partial_j$, with a symbol $P(\xi)$ is defined by $P(D)u(x) = (2\pi)^{-n} \int_{\mathbf{R}^n} P(\xi) \hat{u}(\xi) e^{ix \cdot \xi} d\xi$. We put $J = 1 + |D|$, so that $\|u\|_s = \|J^s u\|$. For operators A and B , we denote by $[A, B] = AB - BA$ the commutator. Throughout this paper, we denote inessential constants by the same symbol C .

2. Formulation of the problem

Let $x = (x_1, x_2, \dots, x_n)$ be the horizontal spatial variables and x_{n+1} the vertical spatial variable. We denote by $X = (x, x_{n+1}) = (x_1, \dots, x_n, x_{n+1})$ the whole spatial variables. We will consider a water wave in $(n + 1)$ -dimensional space and assume that the domain $\Omega(t)$ occupied by the fluid at time $t \geq 0$, the free surface $\Gamma(t)$, and the bottom Σ are of the forms

$$\begin{aligned} \Omega(t) &= \{X = (x, x_{n+1}) \in \mathbf{R}^{n+1}; b(x) < x_{n+1} < h + \eta(x, t)\}, \\ \Gamma(t) &= \{X = (x, x_{n+1}) \in \mathbf{R}^{n+1}; x_{n+1} = h + \eta(x, t)\}, \\ \Sigma &= \{X = (x, x_{n+1}) \in \mathbf{R}^{n+1}; x_{n+1} = b(x)\}, \end{aligned}$$

where h is the mean depth of the fluid. The functions b and η represent the bottom topography and the surface elevation, respectively. In this paper b is

a given function, while η is the unknown. In fact, our main interest is the behavior of the free surface.

We assume that the fluid is incompressible and inviscid, and that the flow is irrotational. Then, the fluid motion is described by the velocity potential $\Phi = \Phi(X, t)$ satisfying the equation

$$(2.1) \quad \Delta_X \Phi = 0 \quad \text{in } \Omega(t), \quad t > 0,$$

where Δ_X is the Laplacian with respect to X , that is, $\Delta_X = \Delta + \partial_{n+1}^2$ and $\Delta = \partial_1^2 + \dots + \partial_n^2$. The boundary conditions on the free surface are given by

$$(2.2) \quad \begin{cases} \eta_t + \nabla \Phi \cdot \nabla \eta - \partial_{n+1} \Phi = 0, \\ \Phi_t + \frac{1}{2} |\nabla_X \Phi|^2 + g\eta = 0 \quad \text{on } \Gamma(t), \quad t > 0, \end{cases}$$

where $\nabla = (\partial_1, \dots, \partial_n)^T$ and $\nabla_X = (\partial_1, \dots, \partial_n, \partial_{n+1})^T$ are the gradients with respect to $x = (x_1, \dots, x_n)$ and to $X = (x, x_{n+1})$, respectively, and g is the gravitational constant. The first equation is the kinematical condition and the second one is derived from Bernoulli's law. The boundary condition on the bottom is given by

$$(2.3) \quad N \cdot \nabla_X \Phi = 0 \quad \text{on } \Sigma, \quad t > 0,$$

where N is the normal vector to the bottom Σ . Finally, we impose the initial conditions

$$(2.4) \quad \eta(x, 0) = \eta_0(x), \quad \Phi(X, 0) = \Phi_0(X).$$

It should be assumed that the initial data satisfy the compatibility conditions, that is, $\Delta_X \Phi_0 = 0$ in $\Omega(0)$ and $N \cdot \nabla_X \Phi_0 = 0$ on Σ .

Remark 2.1. In a derivation of the second equation in (2.2) we first integrate the conservation of momentum, that is, the Euler equation $0 = \rho(v_t + (v \cdot \nabla_X)v) + \nabla_X p + \rho g \mathbf{e}_{n+1} = \rho \nabla_X (\Phi_t + \frac{1}{2} |\nabla_X \Phi|^2 + \frac{1}{\rho}(p - p_0) + g(x_{n+1} - h))$ and obtain

$$\Phi_t + \frac{1}{2} |\nabla_X \Phi|^2 + \frac{1}{\rho}(p - p_0) + g(x_{n+1} - h) = f(t) \quad \text{in } \Omega(t), \quad t > 0,$$

where $v = \nabla_X \Phi$ is a velocity, ρ is a constant density, p_0 is a constant atmospheric pressure, \mathbf{e}_{n+1} is the unit vector in the vertical direction, and $f(t)$ is an arbitrary function of time t . This equation expresses what is called Bernoulli's law. Replacing Φ by $\Phi + \int f(t) dt$, restricting the above equation on the free surface $\Gamma(t)$, and using the dynamical boundary condition $p = p_0$ on $\Gamma(t)$, we get the second equation in (2.2).

We proceed to rewrite the equations (2.1)–(2.4) in an appropriate non-dimensional form. Let λ be the typical wave length and h the mean depth. We

introduce a non-dimensional parameter δ by $\delta = h/\lambda$ and rescale the independent and dependent variables by

$$(2.5) \quad \begin{cases} x = \lambda\tilde{x}, & x_{n+1} = h\tilde{x}_{n+1}, & t = \frac{\lambda}{\sqrt{gh}}\tilde{t}, \\ \Phi = \lambda\sqrt{gh}\tilde{\Phi}, & \eta = h\tilde{\eta}, & b = h\tilde{b}. \end{cases}$$

Here, we note that the function η of the free surface is not rescaled by the maximum vertical amplitude of the free surface a but by the mean depth h . This means that we will consider the water waves in the regime $\varepsilon \simeq 1$. Putting these into (2.1)–(2.4) and dropping the tilde sign in the notation we obtain

$$(2.6) \quad \delta^2 \Delta \Phi + \partial_{n+1}^2 \Phi = 0 \quad \text{in } \Omega(t), \quad t > 0,$$

$$(2.7) \quad \begin{cases} \delta^2 (\eta_t + \nabla \Phi \cdot \nabla \eta) - \partial_{n+1} \Phi = 0, \\ \delta^2 (\Phi_t + \frac{1}{2} |\nabla \Phi|^2 + \eta) + \frac{1}{2} (\partial_{n+1} \Phi)^2 = 0 \end{cases} \quad \text{on } \Gamma(t), \quad t > 0,$$

$$(2.8) \quad \partial_{n+1} \Phi - \delta^2 \nabla b \cdot \nabla \Phi = 0 \quad \text{on } \Sigma, \quad t > 0,$$

$$(2.9) \quad \eta(x, 0) = \eta_0^\delta(x), \quad \Phi(X, 0) = \Phi_0^\delta(X),$$

where

$$\begin{aligned} \Omega(t) &= \{X = (x, x_{n+1}) \in \mathbf{R}^{n+1}; b(x) < x_{n+1} < 1 + \eta(x, t)\}, \\ \Gamma(t) &= \{X = (x, x_{n+1}) \in \mathbf{R}^{n+1}; x_{n+1} = 1 + \eta(x, t)\}, \\ \Sigma &= \{X = (x, x_{n+1}) \in \mathbf{R}^{n+1}; x_{n+1} = b(x)\}. \end{aligned}$$

Since we are interested in asymptotic behavior of the solution when $\delta \rightarrow +0$, we always assume $0 < \delta \leq 1$ in the following.

As in the usual way, we transform equivalently the initial value problem (2.6)–(2.9) to a problem on the free surface. To this end, we introduce new unknown function ϕ by

$$(2.10) \quad \phi(x, t) = \Phi(x, 1 + \eta(x, t), t),$$

which is the trace of the velocity potential on the free surface. Then, we see that

$$(2.11) \quad \begin{cases} \phi_t = \Phi_t|_{\Gamma(t)} + \partial_{n+1} \Phi|_{\Gamma(t)} \eta_t, \\ \nabla \phi = \nabla \Phi|_{\Gamma(t)} + \partial_{n+1} \Phi|_{\Gamma(t)} \nabla \eta. \end{cases}$$

It follows from (2.6), (2.8), and (2.10) that

$$(2.12) \quad \Lambda(\eta, b, \delta) \phi = (\delta^{-2} \partial_{n+1} \Phi - \nabla \eta \cdot \nabla \Phi)|_{\Gamma(t)},$$

where $\Lambda = \Lambda(\eta, b, \delta)$ is a linear operator called the Dirichlet-to-Neumann map for Laplace's equation. In Section 4 we will give the definition and basic properties of this map $\Lambda = \Lambda(\eta, b, \delta)$. This and the second equation in (2.11) imply that

$$(2.13) \quad \begin{cases} \partial_{n+1} \Phi|_{\Gamma(t)} = \delta^2 (1 + \delta^2 |\nabla \eta|^2)^{-1} (\Lambda \phi + \nabla \eta \cdot \nabla \phi), \\ \nabla \Phi|_{\Gamma(t)} = \nabla \phi - \delta^2 (1 + \delta^2 |\nabla \eta|^2)^{-1} (\Lambda \phi + \nabla \eta \cdot \nabla \phi) \nabla \eta. \end{cases}$$

It follows from the first equation in (2.7) and (2.12) that $\eta_t - \Lambda\phi = 0$, so that by the first equation in (2.11) we get

$$\Phi_t|_{\Gamma(t)} = \phi_t - \delta^2(1 + \delta^2|\nabla\eta|^2)^{-1}(\Lambda\phi + \nabla\eta \cdot \nabla\phi)\Lambda\phi.$$

Putting this and (2.13) into the second equation in (2.7) we obtain

$$(2.14) \quad \begin{cases} \eta_t - \Lambda(\eta, b, \delta)\phi = 0, \\ \phi_t + \eta + \frac{1}{2}|\nabla\phi|^2 \\ - \frac{1}{2}\delta^2(1 + \delta^2|\nabla\eta|^2)^{-1}(\Lambda(\eta, b, \delta)\phi + \nabla\eta \cdot \nabla\phi)^2 = 0 \quad \text{for } t > 0, \end{cases}$$

$$(2.15) \quad \eta = \eta_0^\delta, \quad \phi = \phi_0^\delta \quad \text{at } t = 0,$$

where $\phi_0^\delta = \Phi_0^\delta(\cdot, 1 + \eta_0^\delta(\cdot))$. This is the initial value problem that we are going to investigate in this paper. The following theorem is one of the main results in this paper and asserts the existence of the solution for the above initial value problem with uniform bounds of the solution on a time interval independent of small $\delta > 0$.

Theorem 2.1. *Let $M_0, c_0 > 0$ and $s > n/2 + 1$. There exist a time $T > 0$ and constants $C_0, \delta_0 > 0$ such that for any $\delta \in (0, \delta_0]$, $\nabla\phi_0^\delta \in H^{s+3}$, $\eta_0^\delta \in H^{s+3+1/2}$, and $b \in H^{s+4+1/2}$ satisfying*

$$\begin{cases} \|\nabla\phi_0^\delta\|_{s+3} + \|\eta_0^\delta\|_{s+3+1/2} + \|b\|_{s+4+1/2} \leq M_0, \\ 1 + \eta_0^\delta(x) - b(x) \geq c_0 \quad \text{for } x \in \mathbf{R}^n, \end{cases}$$

the initial value problem (2.14) and (2.15) has a unique solution $(\eta, \phi) = (\eta^\delta, \phi^\delta)$ on the time interval $[0, T]$ satisfying

$$\begin{cases} \|\eta^\delta(t)\|_{s+3} + \|\nabla\phi^\delta(t)\|_{s+2} + \|(\eta_t^\delta(t), \phi_t^\delta(t))\|_{s+2} \leq C_0, \\ 1 + \eta^\delta(x, t) - b(x) \geq c_0/2 \quad \text{for } x \in \mathbf{R}^n, 0 \leq t \leq T, 0 < \delta \leq \delta_0. \end{cases}$$

Remark 2.2. We cannot expect that the velocity potential Φ and its trace ϕ on the free surface vanish at spatial infinity even if so does the velocity $v = \nabla_X\Phi$. Hence, it is natural to consider the initial value problem (2.14) and (2.15) in a class $\nabla\phi \in H^s$ (not a class $\phi \in H^s$). However, if we impose additional conditions $\phi_0^\delta \in L^2$ and $\|\phi_0^\delta\| \leq M_0$, then we have $\phi^\delta \in C([0, T]; H^{s+3})$ with a uniform estimate $\|\phi^\delta(t)\|_{s+3} \leq C_0$.

3. A shallow water approximation

In this section we study formally asymptotic behavior of the solution $(\eta^\delta, \phi^\delta)$ to the initial value problem (2.14) and (2.15) when $\delta \rightarrow +0$ and derive the shallow water equations, whose solution approximates $(\eta^\delta, \nabla\phi^\delta)$ in a suitable sense. Then, we will give another main theorem which ensures a rigorous approximation of the water wave by the shallow water equations.

It follows from the second equation in (2.14) that

$$\phi_t + \eta + \frac{1}{2}|\nabla\phi|^2 = O(\delta^2).$$

By (2.6) and (2.8),

$$\begin{aligned} & (\partial_{n+1}\Phi)(x, x_{n+1}, t) \\ (3.1) \quad &= (\partial_{n+1}\Phi)(x, b(x), t) + \int_{b(x)}^{x_{n+1}} (\partial_{n+1}^2\Phi)(x, y, t)dy \\ &= \delta^2\nabla b(x) \cdot \nabla\Phi(x, b(x), t) - \delta^2 \int_{b(x)}^{x_{n+1}} (\Delta\Phi)(x, y, t)dy, \end{aligned}$$

which implies that $(\partial_{n+1}\Phi)(X, t) = O(\delta^2)$. Therefore,

$$\begin{aligned} \nabla\Phi(x, x_{n+1}, t) &= \nabla\Phi(x, 1 + \eta(x, t), t) + \int_{1+\eta(x,t)}^{x_{n+1}} (\nabla\partial_{n+1}\Phi)(x, y, t)dy \\ &= \nabla\Phi(x, 1 + \eta(x, t), t) + O(\delta^2). \end{aligned}$$

Moreover, by the definition (2.10) it holds that

$$\begin{aligned} \nabla\phi(x, t) &= \nabla\Phi(x, 1 + \eta(x, t), t) + \nabla\eta(x)(\partial_{n+1}\Phi)(x, 1 + \eta(x), t) \\ &= \nabla\Phi(x, 1 + \eta(x, t), t) + O(\delta^2) \\ &= \nabla\Phi(X, t) + O(\delta^2). \end{aligned}$$

Similarly, we have

$$\Delta\phi(x, t) = \Delta\Phi(X, t) + O(\delta^2).$$

These relation and (3.1) imply that

$$\begin{aligned} & (\partial_{n+1}\Phi)(x, 1 + \eta(x, t), t) \\ &= \delta^2\nabla b(x) \cdot \nabla\phi(x, t) - \delta^2 \int_{b(x)}^{1+\eta(x,t)} \Delta\phi(x, t)dy + O(\delta^4) \\ &= -\delta^2(1 + \eta(x, t))\Delta\phi(x, t) + \delta^2\nabla \cdot (b(x)\nabla\phi(x, t)) + O(\delta^4). \end{aligned}$$

Hence, by (2.12) we have

$$(3.2) \quad (\Lambda\phi)(x, t) = -\nabla \cdot ((1 + \eta(x, t) - b(x))\nabla\phi(x, t)) + O(\delta^2).$$

This and the first equation in (2.14) imply that

$$\eta_t + \nabla \cdot ((1 + \eta - b)\nabla\phi) = O(\delta^2).$$

To summarize, we have derived the partial differential equations

$$\begin{cases} \eta_t + \nabla \cdot ((1 + \eta - b)\nabla\phi) = O(\delta^2), \\ \phi_t + \eta + \frac{1}{2}|\nabla\phi|^2 = O(\delta^2), \end{cases}$$

which approximate the equations in (2.14) up to order δ^2 . Letting $\delta \rightarrow 0$ in the above equations we obtain

$$\begin{cases} \eta_t^0 + \nabla \cdot ((1 + \eta^0 - b)\nabla\phi^0) = 0, \\ \phi_t^0 + \eta^0 + \frac{1}{2}|\nabla\phi^0|^2 = 0. \end{cases}$$

Finally, putting $u^0 := \nabla\phi^0$ and taking the gradient of the second equation, we are led to the shallow water equations

$$(3.3) \quad \begin{cases} \eta_t^0 + \nabla \cdot ((1 + \eta^0 - b)u^0) = 0, \\ u_t^0 + (u^0 \cdot \nabla)u^0 + \nabla\eta^0 = 0. \end{cases}$$

Moreover, u^0 satisfies the irrotational condition

$$(3.4) \quad \text{rot } u^0 = 0,$$

where $\text{rot } u$ is the rotation of $u = (u_1, \dots, u_n)^T$ defined by $\text{rot } u = (\partial_j u_i - \partial_i u_j)_{1 \leq i, j \leq n}$.

The following theorem is another main result in this paper and gives a mathematically rigorous justification of the shallow water approximation for water waves.

Theorem 3.1. *In addition to hypothesis of Theorem 2.1 we assume that as $\delta \rightarrow +0$ the initial data $(\eta_0^\delta, \nabla\phi_0^\delta)$ converge to (η_0^0, u_0^0) in $H^{s+3} \times H^{s+2}$. Then, as $\delta \rightarrow +0$ the solution obtained in Theorem 2.1 satisfies*

$$\begin{aligned} (\eta^\delta, \nabla\phi^\delta) &\rightarrow (\eta^0, u^0) \quad \text{weakly}^* \text{ in } L^\infty(0, T; H^{s+3} \times H^{s+2}), \\ &\text{strongly in } C([0, T]; H^{s+3-\epsilon} \times H^{s+2-\epsilon}) \end{aligned}$$

for each $\epsilon > 0$, where (η^0, u^0) is a unique solution of the shallow water equations (3.3) with initial conditions $(\eta^0, u^0)|_{t=0} = (\eta_0^0, u_0^0)$ and u^0 satisfies the irrotational condition (3.4).

Moreover, if we also assume that $\|\eta_0^\delta - \eta_0^0\|_s + \|\nabla\phi_0^\delta - u_0^0\|_s = O(\delta^2)$, then for any $\delta \in (0, \delta_0]$ and $t \in [0, T]$ we have

$$\|\eta^\delta(t) - \eta^0(t)\|_s + \|\nabla\phi^\delta(t) - u^0(t)\|_s \leq C\delta^2$$

with a constant C independent of δ and t .

4. The Dirichlet-to-Neumann map Λ

Throughout this and following two sections the time t is arbitrarily fixed, so that $\Omega(t)$, $\Gamma(t)$, and $\eta(x, t)$ are simply denoted by Ω , Γ , and $\eta(x)$, respectively. Introducing a $(n + 1) \times (n + 1)$ matrix I_δ by

$$I_\delta = \begin{pmatrix} E_n & 0 \\ 0 & \delta^{-1} \end{pmatrix},$$

where E_n is the $n \times n$ unit matrix, we consider the boundary value problem

$$(4.1) \quad \begin{cases} \nabla_X \cdot I_\delta^2 \nabla_X \Phi = 0 & \text{in } \Omega, \\ \Phi = \phi & \text{on } \Gamma, \\ N \cdot I_\delta^2 \nabla_X \Phi = 0 & \text{on } \Sigma. \end{cases}$$

We note that the first and the third equations in (4.1) are nothing but those in (2.6) and (2.8), respectively. Under suitable assumptions on η and b , for any function ϕ on Γ in some class there exists a unique solution Φ of the boundary value problem (4.1).

Definition 4.1. Using the solution Φ we define a linear operator $\Lambda = \Lambda(\eta, b, \delta)$ by

$$\Lambda(\eta, b, \delta)\phi = (-\nabla\eta, 1)^T \cdot I_\delta^2 \nabla_X \Phi(x, 1 + \eta(x)),$$

which is called the Dirichlet-to-Neumann map. The solution Φ will be denoted by $\phi^{\mathring{h}}$.

The following lemma was given by D. Lannes [15]. For the completeness, we will give the proof.

Lemma 4.1. *The Dirichlet-to-Neumann map $\Lambda = \Lambda(\eta, b, \delta)$ is symmetric in L^2 , that is, for any $\phi, \psi \in H^1$ it holds that*

$$(\Lambda\phi, \psi) = (\phi, \Lambda\psi).$$

Proof. Set $\Phi := \phi^{\mathring{h}}$ and $\Psi := \psi^{\mathring{h}}$. By Green's formula we have

$$\begin{aligned} 0 &= \int_{\Omega} ((\nabla_X \cdot I_\delta^2 \nabla_X \Phi)\Psi - \Phi(\nabla_X \cdot I_\delta^2 \nabla_X \Psi)) dX \\ &= \int_{\Gamma} ((N \cdot I_\delta^2 \nabla_X \Phi)\Psi - \Phi(N \cdot I_\delta^2 \nabla_X \Psi)) dS, \end{aligned}$$

where N is the unit outward normal to the boundary $\partial\Omega$. In the above calculation we used the boundary condition on the bottom Σ . Since $\Phi = \phi$, $\Psi = \psi$, $\sqrt{1 + |\nabla\eta|^2} N \cdot I_\delta^2 \nabla_X \Phi = \Lambda\phi$, $\sqrt{1 + |\nabla\eta|^2} N \cdot I_\delta^2 \nabla_X \Psi = \Lambda\psi$, and $dS = \sqrt{1 + |\nabla\eta|^2} dx$ on Γ , we obtain the desired identity. \square

Lemma 4.2. *For any $\phi \in H^1$, it holds that $(\Lambda\phi, \phi) = \|I_\delta \nabla_X \Phi\|_{L^2(\Omega)}^2$, where $\Phi = \phi^{\mathring{h}}$.*

Proof. By Green's formula we see that

$$0 = \int_{\Omega} (\nabla_X \cdot I_\delta^2 \nabla_X \Phi)\Phi dX = \int_{\partial\Omega} (N \cdot I_\delta^2 \nabla_X \Phi)\Phi dS - \int_{\Omega} |I_\delta \nabla_X \Phi|^2 dX.$$

This together with the boundary conditions yields the desired identity. \square

In the case where the free surface and the bottom are both flat, that is, the case $(\eta, b) = 0$, the Dirichlet-to-Neumann map can be written explicitly in terms of the Fourier multiplier as $\Lambda_0 = \Lambda(0, 0, \delta) = \frac{1}{\delta}|D| \tanh(\delta|D|)$ so that we can easily handle the map. In order to obtain uniform estimates of the map Λ in the general case, it would be better to make a connection between the maps Λ and Λ_0 . In the two-dimensional case we have a conformal map from the strip $\mathbf{R} \times (0, 1)$ onto the fluid region Ω to make the connection. In fact, L. V. Ovsjannikov [18, 19], T. Kano and T. Nishida [10]–[12], S. Wu [24], and Y. A. Li [16] used the conformal map. In the case where the spatial dimension is greater than two, unfortunately, we do not have such a nice map. D. Lannes [15] used a simple diffeomorphism which stretches or compresses only on vertical line segments. In this paper, in place of the conformal map or the simple diffeomorphism we will use an appropriate diffeomorphism $\Theta = (\Theta_1, \dots, \Theta_n, \Theta_{n+1}) : \overline{\Omega}_0 = \mathbf{R}^n \times [0, 1] \rightarrow \overline{\Omega}$, which is conformal in the tangential and the normal directions on the boundary in some sense, so that the Neumann boundary condition on the bottom is transformed into again the Neumann condition with a very simple normal vector $N = (0, \dots, 0, -1)^T$. Thanks of this choice of the diffeomorphism Θ and standard elliptic estimates we can obtain important uniform estimates of the Dirichlet-to-Neumann map Λ with respect to small δ .

We take functions $\theta = (\theta_1, \dots, \theta_n, \theta_{n+1})$ satisfying the conditions

$$(4.2) \quad \begin{cases} \theta_j(x, 0) = \theta_j(x, 1) = 0, & \partial_{n+1}\theta_j(x, 0) = -\partial_j b(x), \\ \partial_{n+1}\theta_j(x, 1) = -\partial_j \eta(x) & \text{for } 1 \leq j \leq n, \\ \theta_{n+1}(x, 0) = b(x), & \theta_{n+1}(x, 1) = \eta(x), \\ \partial_{n+1}\theta_{n+1}(x, 0) = \partial_{n+1}\theta_{n+1}(x, 1) = 0, \end{cases}$$

and define the diffeomorphism Θ by

$$(4.3) \quad \begin{cases} \Theta_j(X) = x_j + \delta^2 \theta_j(X) & \text{for } 1 \leq j \leq n, \\ \Theta_{n+1}(X) = x_{n+1} + \theta_{n+1}(X). \end{cases}$$

It is easy to see that

$$(4.4) \quad \frac{\partial \Theta}{\partial X} = \begin{pmatrix} E_n + \delta^2 \frac{\partial(\theta_1, \dots, \theta_n)}{\partial(x_1, \dots, x_n)} & \delta^2 \partial_{n+1}(\theta_1, \dots, \theta_n)^T \\ (\nabla \theta_{n+1})^T & 1 + \partial_{n+1} \theta_{n+1} \end{pmatrix}$$

and that

$$\begin{cases} \Theta(x, 0) = (x, b(x)), & \Theta(x, 1) = (x, 1 + \eta(x)), \\ \frac{\partial \Theta}{\partial X}(x, 0) = \begin{pmatrix} E_n & -\delta^2 \nabla b(x) \\ (\nabla b(x))^T & 1 \end{pmatrix}, \\ \frac{\partial \Theta}{\partial X}(x, 1) = \begin{pmatrix} E_n & -\delta^2 \nabla \eta(x) \\ (\nabla \eta(x))^T & 1 \end{pmatrix}. \end{cases}$$

We put $\tilde{\Phi} := \Phi \circ \Theta$ and

$$(4.5) \quad \begin{aligned} P &:= \det\left(\frac{\partial\Theta}{\partial X}\right) I_\delta^{-1} \left(\frac{\partial\Theta}{\partial X}\right)^{-1} I_\delta^2 \left(\left(\frac{\partial\Theta}{\partial X}\right)^{-1}\right)^T I_\delta^{-1} \\ &= \det\left(\frac{\partial\Theta}{\partial X}\right) \left(\left(I_\delta^{-1} \frac{\partial\Theta}{\partial X} I_\delta\right)^T \left(I_\delta^{-1} \frac{\partial\Theta}{\partial X} I_\delta\right)\right)^{-1}. \end{aligned}$$

Then, the boundary value problem (4.1) is transformed into

$$(4.6) \quad \begin{cases} \nabla_X \cdot I_\delta P I_\delta \nabla_X \tilde{\Phi} = 0 & \text{in } 0 < x_{n+1} < 1, \\ \tilde{\Phi} = \phi & \text{on } x_{n+1} = 1, \\ \partial_{n+1} \tilde{\Phi} = 0 & \text{on } x_{n+1} = 0. \end{cases}$$

We refer to §6.3.2 in the textbook of L. C. Evans [5] for the transformation of the equation. Here, we have

$$(4.7) \quad I_\delta^{-1} \frac{\partial\Theta}{\partial X} I_\delta = \begin{pmatrix} E_n + \delta^2 \frac{\partial(\theta_1, \dots, \theta_n)}{\partial(x_1, \dots, x_n)} & \delta \partial_{n+1}(\theta_1, \dots, \theta_n)^T \\ \delta(\nabla_{\theta_{n+1}})^T & 1 + \partial_{n+1} \theta_{n+1} \end{pmatrix}.$$

On the upper boundary $x_{n+1} = 1$, we have $\det\left(\frac{\partial\Theta}{\partial X}\right) = 1 + \delta^2 |\nabla\eta|^2$ so that

$$\begin{aligned} P &= (1 + \delta^2 |\nabla\eta|^2) \left(\begin{pmatrix} E_n & \delta \nabla\eta \\ -\delta(\nabla\eta)^T & 1 \end{pmatrix} \begin{pmatrix} E_n & -\delta \nabla\eta \\ \delta(\nabla\eta)^T & 1 \end{pmatrix} \right)^{-1} \\ &= (1 + \delta^2 |\nabla\eta|^2) \begin{pmatrix} E_n + \delta^2 \nabla\eta(\nabla\eta)^T & 0 \\ 0 & 1 + \delta^2 |\nabla\eta|^2 \end{pmatrix}^{-1}. \end{aligned}$$

Similar identity holds for the lower boundary $x_{n+1} = 0$. Therefore, we see that

$$(4.8) \quad P(x, 0) = \begin{pmatrix} * & 0 \\ 0 & 1 \end{pmatrix}, \quad P(x, 1) = \begin{pmatrix} * & 0 \\ 0 & 1 \end{pmatrix}.$$

Particularly, it holds that

$$(4.9) \quad \mathbf{e}_{n+1} \cdot I_\delta P I_\delta \nabla_X \tilde{\Phi} = \mathbf{e}_{n+1} \cdot I_\delta^2 \nabla_X \tilde{\Phi} = \delta^{-2} \partial_{n+1} \tilde{\Phi} \quad \text{on } x_{n+1} = 0, 1.$$

We also have the relation

$$(4.10) \quad \begin{aligned} &I_\delta \nabla_X \tilde{\Phi} \\ &= \begin{pmatrix} E_n + \delta^2 \left(\frac{\partial(\theta_1, \dots, \theta_n)}{\partial(x_1, \dots, x_n)}\right)^T & \delta \nabla\theta_{n+1} \\ \delta \partial_{n+1}(\theta_1, \dots, \theta_n) & 1 + \partial_{n+1} \theta_{n+1} \end{pmatrix} I_\delta (\nabla_X \Phi) \circ \Theta. \end{aligned}$$

Assumption 4.1. Let $r > n/2$.

(A1) There exists a C^1 -diffeomorphism $\Theta : \overline{\Omega}_0 \rightarrow \overline{\Omega}$ satisfying (4.2), (4.3), and the conditions $\det\left(\frac{\partial\Theta}{\partial X}(X)\right) \geq c > 0$ and $|\nabla_X \theta(X)| \leq M$ for $X \in \Omega_0$.

(A2) $\|\nabla_X \theta(\cdot, x_{n+1})\|_{r+1} \leq M$ for $0 \leq x_{n+1} \leq 1$.

The construction of a diffeomorphism Θ satisfying the above conditions will be given later. By (4.10) we can easily obtain the following lemma.

Lemma 4.3. *Under Assumption 4.1 (A1), there exists a constant $C = C(M, c) \geq 1$ independent of δ such that*

$$C^{-1} \|I_\delta \nabla_X \Phi\|_{L^2(\Omega)} \leq \|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} \leq C \|I_\delta \nabla_X \Phi\|_{L^2(\Omega)}.$$

The next lemma is one of the crucial part of this paper and leads uniform estimates of the Dirichlet-to-Neumann map Λ .

Lemma 4.4. *Under Assumption 4.1 (A1), there exists a constant $C = C(M, c) \geq 1$ independent of δ such that for any $\phi \in H^1$ we have*

$$C^{-1} \|\Lambda_0^{1/2} \phi\|^2 \leq (\Lambda \phi, \phi) \leq C \|\Lambda_0^{1/2} \phi\|^2,$$

where $\Lambda_0 = \Lambda(0, 0, \delta) = \frac{1}{8} |D| \tanh(\delta |D|)$.

Remark 4.1. It was shown by S. Wu [25] in the case of infinite depth and by D. Lannes [15] in the case of finite depth that $(\Lambda \phi, \phi) + \|\phi\|^2$ is equivalent to $\|\phi\|_{1/2}^2$. However, this equivalence depends strongly on the parameter δ . The most important point of the above lemma is that $(\Lambda \phi, \phi)$ is equivalent to $\|\Lambda_0^{1/2} \phi\|^2$ uniformly with respect to δ .

Proof. We set $\Phi := \phi^h$ and $\tilde{\Phi} := \Phi \circ \Theta$, and decompose $\tilde{\Phi} = \tilde{\Phi}_1 + \tilde{\Phi}_2$, where $\tilde{\Phi}_1$ and $\tilde{\Phi}_2$ are solutions of the boundary value problems

$$\begin{cases} \nabla_X \cdot I_\delta^2 \nabla_X \tilde{\Phi}_1 = 0 & \text{in } 0 < x_{n+1} < 1, \\ \tilde{\Phi}_1 = \phi & \text{on } x_{n+1} = 1, \\ \partial_{n+1} \tilde{\Phi}_1 = 0 & \text{on } x_{n+1} = 0 \end{cases}$$

and

$$\begin{cases} \nabla_X \cdot I_\delta^2 \nabla_X \tilde{\Phi}_2 = \nabla_X \cdot I_\delta (I_1 - P) I_\delta \nabla_X \tilde{\Phi} & \text{in } 0 < x_{n+1} < 1, \\ \tilde{\Phi}_2 = 0 & \text{on } x_{n+1} = 1, \\ \partial_{n+1} \tilde{\Phi}_2 = 0 & \text{on } x_{n+1} = 0, \end{cases}$$

respectively. Then, it holds that

$$\begin{aligned} \Lambda \phi &= \delta^{-2} \partial_{n+1} \tilde{\Phi}(\cdot, 1) = \delta^{-2} \partial_{n+1} \tilde{\Phi}_1(\cdot, 1) + \delta^{-2} \partial_{n+1} \tilde{\Phi}_2(\cdot, 1) \\ &= \Lambda_0 \phi + \delta^{-2} \partial_{n+1} \tilde{\Phi}_2(\cdot, 1) \end{aligned}$$

and, by Lemma 4.2, that

$$(\Lambda \phi, \phi) = \|I_\delta \nabla_X \Phi\|_{L^2(\Omega)}^2, \quad \|\Lambda_0^{1/2} \phi\|^2 = (\Lambda_0 \phi, \phi) = \|I_\delta \nabla_X \tilde{\Phi}_1\|_{L^2(\Omega_0)}^2.$$

By Green's formula we see that

$$\begin{aligned}
(\delta^{-2}\partial_{n+1}\tilde{\Phi}_2(\cdot, 1), \phi) &= (\delta^{-2}\partial_{n+1}\tilde{\Phi}_2(\cdot, 1), \tilde{\Phi}_1(\cdot, 1)) \\
&= \int_{\Omega_0} I_\delta \nabla_X \tilde{\Phi}_2 \cdot I_\delta \nabla_X \tilde{\Phi}_1 dX + \int_{\Omega_0} (\nabla_X \cdot I_\delta^2 \nabla_X \tilde{\Phi}_2) \tilde{\Phi}_1 dX \\
&= \int_{\Omega_0} I_\delta \nabla_X \tilde{\Phi}_2 \cdot I_\delta \nabla_X \tilde{\Phi}_1 dX + \int_{\Omega_0} (\nabla_X \cdot I_\delta(I_1 - P)I_\delta \nabla_X \tilde{\Phi}) \tilde{\Phi}_1 dX \\
&= \int_{\Omega_0} I_\delta \nabla_X \tilde{\Phi}_2 \cdot I_\delta \nabla_X \tilde{\Phi}_1 dX - \int_{\Omega_0} (I_1 - P)I_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \nabla_X \tilde{\Phi}_1 dX,
\end{aligned}$$

where we used (4.8). Therefore,

$$\begin{aligned}
|(\delta^{-2}\partial_{n+1}\tilde{\Phi}_2(\cdot, 1), \phi)| \\
\leq C(\|I_\delta \nabla_X \tilde{\Phi}_2\|_{L^2(\Omega_0)} + \|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)})\|I_\delta \nabla_X \tilde{\Phi}_1\|_{L^2(\Omega_0)}.
\end{aligned}$$

Similarly, by the equations for $\tilde{\Phi}_2$ we see that

$$\begin{aligned}
&\|I_\delta \nabla_X \tilde{\Phi}_2\|_{L^2(\Omega_0)}^2 \\
&= - \int_{\Omega_0} (\nabla_X \cdot I_\delta^2 \nabla_X \tilde{\Phi}_2) \tilde{\Phi}_2 dX = - \int_{\Omega_0} (\nabla_X \cdot I_\delta(I_1 - P)I_\delta \nabla_X \tilde{\Phi}) \tilde{\Phi}_2 dX \\
&= \int_{\Omega_0} (I_1 - P)I_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \nabla_X \tilde{\Phi}_2 dX \leq C\|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}\|I_\delta \nabla_X \tilde{\Phi}_2\|_{L^2(\Omega_0)},
\end{aligned}$$

so that

$$\|I_\delta \nabla_X \tilde{\Phi}_2\|_{L^2(\Omega_0)} \leq C\|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} \leq C\|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega)},$$

where we used Lemma 4.3. Summarizing the above estimates we obtain

$$\begin{aligned}
|(\Lambda\phi, \phi) - (\Lambda_0\phi, \phi)| &\leq C\|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega)}\|I_\delta \nabla_X \tilde{\Phi}_1\|_{L^2(\Omega_0)} \\
&\leq C\sqrt{(\Lambda\phi, \phi)}\sqrt{(\Lambda_0\phi, \phi)},
\end{aligned}$$

which easily yields the desired inequalities. \square

Now, let us construct the diffeomorphism Θ satisfying the conditions in Assumption 4.1.

Lemma 4.5. *Let $r > n/2$, $c_1, M_1 > 0$ and suppose that $\eta, b \in H^{1+r}$ satisfy the conditions*

$$\begin{cases} \|\eta\|_{1+r} + \|b\|_{1+r} \leq M_1, \\ 1 + \eta(x) - b(x) \geq c_1 \quad \text{for } x \in \mathbf{R}^n. \end{cases}$$

Then, there exists a constant $\delta_1 = \delta_1(M_1, c_1, r) > 0$ such that for any $\delta \in (0, \delta_1]$ we can construct a diffeomorphism Θ satisfying the conditions in Assumption 4.1 (A1). Moreover, for any $s > 0$ and $k \in \mathbf{N}$ we have

$$(4.11) \quad \begin{cases} \|J^s \nabla_X \theta\|_{L^2(\Omega_0)} \leq C_1(\|\eta\|_{s+1/2} + \|b\|_{s+1/2}), \\ \sup_{0 \leq x_{n+1} \leq 1} \|\partial_{n+1}^k \theta(\cdot, x_{n+1})\|_s \leq C_2(\|\eta\|_{s+k} + \|b\|_{s+k}), \end{cases}$$

where $C_1 = C_1(c_1) > 0$ and $C_2 = C_2(c_1, k) > 0$. In the case where η depends also on the time t , for any $l \in \mathbf{N}$ we have

$$(4.12) \quad \begin{cases} \|J^s \nabla_X \partial_t^l \theta(t)\|_{L^2(\Omega_0)} \leq C_1 \|\partial_t^l \eta(t)\|_{s+1/2}, \\ \sup_{0 \leq x_{n+1} \leq 1} \|\partial_{n+1}^k \partial_t^l \theta(\cdot, x_{n+1}, t)\|_s \leq C_2 \|\partial_t^l \eta(t)\|_{s+k}. \end{cases}$$

Proof. Without loss of generality we can assume that $0 < c_1 < 1$. Since $\frac{1}{2}(1 + \frac{1}{1-c_1}) > 1$, we can take $\varphi \in C^\infty(\mathbf{R})$ satisfying the conditions

$$\varphi(x_{n+1}) = \begin{cases} 0 & \text{for } x_{n+1} \leq 0, \\ 1 & \text{for } x_{n+1} \geq 1 \end{cases} \quad \text{and} \quad 0 \leq \varphi'(x_{n+1}) \leq \frac{1}{2} \left(1 + \frac{1}{1-c_1}\right).$$

Then, it is easy to check that $1 + (\eta(x) - b(x))\varphi'(x_{n+1}) \geq c_1/2$ holds for any $X = (x, x_{n+1}) \in \overline{\Omega_0}$. Define the functions θ_j , $1 \leq j \leq n+1$, by the relations

$$\begin{cases} \hat{\theta}_j(\xi, x_{n+1}) = \varphi(x_{n+1})e^{-|\xi|(1-x_{n+1})}i\xi_j(1-x_{n+1})\hat{\eta}(\xi) \\ \quad - (1-\varphi(x_{n+1}))e^{-|\xi|x_{n+1}}i\xi_jx_{n+1}\hat{b}(\xi) \quad \text{for } 1 \leq j \leq n, \\ \hat{\theta}_{n+1}(\xi, x_{n+1}) = \varphi(x_{n+1})e^{-\epsilon|\xi|(1-x_{n+1})}(1+\epsilon|\xi|(1-x_{n+1}))\hat{\eta}(\xi) \\ \quad + (1-\varphi(x_{n+1}))e^{-\epsilon|\xi|x_{n+1}}(1+\epsilon|\xi|x_{n+1})\hat{b}(\xi), \end{cases}$$

where $\epsilon > 0$ will be determined later. Obviously, (4.2) is satisfied. It is easy to see that

$$\begin{aligned} |\partial_{n+1}^k \hat{\theta}(\xi, x_{n+1})|^2 &\leq C|\xi|^2(1+|\xi|)^{2(k-1)}(|\hat{\eta}(\xi)|^2 + |\hat{b}(\xi)|^2), \\ \int_0^1 |\widehat{\nabla_X \theta}(\xi, x_{n+1})|^2 dx_{n+1} &\leq C|\xi|(|\hat{\eta}(\xi)|^2 + |\hat{b}(\xi)|^2), \end{aligned}$$

which yield (4.11). In the same way as above, we can show (4.12). It remains to show the estimates in Assumption 4.1 (A1). The latter estimate in (A1) comes from (4.11) and the Sobolev inequality. In view of the relation

$$\begin{aligned} &\hat{\theta}_{n+1}(\xi, x_{n+1}) - \varphi(x_{n+1})\hat{\eta}(\xi) - (1-\varphi(x_{n+1}))\hat{b}(\xi) \\ &= \epsilon \{ \varphi(x_{n+1})e^{-\epsilon|\xi|(1-x_{n+1})}|\xi|(1-x_{n+1})\hat{\eta}(\xi) \\ &\quad + (1-\varphi(x_{n+1}))e^{-\epsilon|\xi|x_{n+1}}|\xi|x_{n+1}\hat{b}(\xi) \} \\ &\quad + \varphi(x_{n+1})(e^{-\epsilon|\xi|(1-x_{n+1})} - 1)\hat{\eta}(\xi) + (1-\varphi(x_{n+1}))(e^{-\epsilon|\xi|x_{n+1}} - 1)\hat{b}(\xi), \end{aligned}$$

we obtain

$$\begin{aligned} &|\partial_{n+1}\theta_{n+1}(x, x_{n+1}) - (\eta(x) - b(x))\varphi'(x_{n+1})| \\ &\leq \epsilon C \int_{\mathbf{R}^n} |\xi|(|\hat{\eta}(\xi)| + |\hat{b}(\xi)|)d\xi \leq \epsilon C(\|\eta\|_{1+r} + \|b\|_{1+r}) \leq \epsilon CM_1. \end{aligned}$$

Therefore, if we take $\epsilon > 0$ so small that $\epsilon CM_1 \leq c_1/4$, then

$$\begin{aligned} 1 + \partial_{n+1}\theta_{n+1}(x, x_{n+1}) &\geq 1 + (\eta(x) - b(x))\varphi'(x_{n+1}) \\ &\quad - |\partial_{n+1}\theta_{n+1}(x, x_{n+1}) - (\eta(x) - b(x))\varphi'(x_{n+1})| \\ &\geq \frac{c_1}{2} - \frac{c_1}{4} = \frac{c_1}{4}. \end{aligned}$$

On the other hand, it follows from (4.4) that

$$(4.13) \quad \det\left(\frac{\partial\Theta}{\partial X}\right) = 1 + \partial_{n+1}\theta_{n+1} + \delta^2 J_1,$$

where J_1 is a polynomial of $\nabla_X\theta$ with coefficients which are polynomials of δ^2 . Hence, we have

$$\det\left(\frac{\partial\Theta}{\partial X}\right) \geq \frac{c_1}{4} - \delta^2 C.$$

Therefore, if we take $\delta_1 > 0$ so small that $\delta_1^2 C \leq c_1/8$, then we obtain the former estimate in (A1). Particularly, we see that $\Theta : \overline{\Omega}_0 \rightarrow \overline{\Omega}$ is a C^1 -diffeomorphism. \square

5. Estimates of the Dirichlet-to-Neumann map

In this section we will give operator norms of the Dirichlet-to-Neumann map $\Lambda = \Lambda(\eta, b, \delta)$ and its commutator $[J^s, \Lambda]$ in Sobolev spaces. Such estimates have already been given by S. Wu [25] in the case of infinite depth and by D. Lannes [15] in the case of finite depth. However, if we apply directly their estimates or techniques, then the estimates depend on the parameter δ . It should be noted that our estimates in this section are valid uniformly with respect to δ .

Lemma 5.1. *Let $r > n/2$. There exists a constant $C = C(r) > 0$ independent of δ such that we have*

$$\|[\Lambda_0^{1/2}, a]u\| \leq C\|\nabla a\|_r\|u\|.$$

Proof. Put $v := [\Lambda_0^{1/2}, a]u$. Then, we have

$$\hat{v}(\xi) = \frac{1}{(2\pi)^n} \int_{\mathbf{R}^n} (\sqrt{\delta^{-1}|\xi| \tanh(\delta|\xi|)} - \sqrt{\delta^{-1}|\eta| \tanh(\delta|\eta|)}) \hat{a}(\xi - \eta) \hat{u}(\eta) d\eta.$$

It is easy to see that $|\sqrt{\alpha \tanh \alpha} - \sqrt{\beta \tanh \beta}| \leq C|\alpha - \beta|$ for $\alpha, \beta \geq 0$, so that

$$(5.1) \quad |\sqrt{\delta^{-1}|\xi| \tanh(\delta|\xi|)} - \sqrt{\delta^{-1}|\eta| \tanh(\delta|\eta|)}| \leq C|\xi - \eta|,$$

and that

$$|\hat{v}(\xi)| \leq C \int_{\mathbf{R}^n} |\xi - \eta| |\hat{a}(\xi - \eta)| |\hat{u}(\eta)| d\eta.$$

This and Hausdorff-Young's inequality give the desired estimate. \square

Lemma 5.2. *For any real s , we have*

$$\left\{ \begin{array}{l} \|\nabla\phi\|_s \leq \sqrt{2(1+\delta)} \|\Lambda_0^{1/2}\phi\|_{s+1/2}, \\ \|\Lambda_0^{1/2}\phi\|_s \leq \min\{\|\nabla\phi\|_s, \delta^{-1/2}\|\phi\|_{s+1/2}\}. \end{array} \right.$$

Proof. By the inequalities $(1 + \sqrt{\alpha})^{-1}\alpha \leq \sqrt{\alpha \tanh \alpha} \leq \min\{\alpha, \sqrt{\alpha}\}$ for $\alpha \geq 0$, it holds that

$$(1 + \sqrt{\delta|\xi|})^{-1}|\xi| \leq \sqrt{\delta^{-1}|\xi| \tanh(\delta|\xi|)} \leq \min\{|\xi|, \delta^{-1/2}|\xi|^{1/2}\}$$

for $\xi \in \mathbf{R}^n$ and $\delta > 0$, which yields the desired estimates. \square

Lemma 5.3. *Under Assumption 4.1 (A1) and (A2), there exists a constant $C = C(M, c, r) > 0$ independent of δ such that we have*

$$\|\Lambda\phi\| \leq C(\|\Lambda_0\phi\| + \|\Lambda_0^{1/2}\phi\|).$$

Proof. Set $\Phi := \phi^{\hbar}$ and $\tilde{\Phi} := \Phi \circ \Theta$. We take $\psi \in H^0$ arbitrarily and define $\tilde{\Psi}$ by $\tilde{\Psi}(\cdot, x_{n+1}) = e^{-\delta|D|(1-x_{n+1})}\psi$. By Green's formula, we see that

$$\begin{aligned} (\Lambda\phi, \psi) &= \int_{\Omega_0} PI_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \nabla_X \tilde{\Psi} dX \\ (5.2) \quad &= \int_{\Omega_0} \Lambda_0^{1/2} PI_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \Lambda_0^{-1/2} \nabla_X \tilde{\Psi} dX. \end{aligned}$$

In view of the relations

$$\begin{aligned} |D|\Lambda_0^{-1/2}\tilde{\Psi}(\cdot, x_{n+1}) &= \delta^{-1}\partial_{n+1}\Lambda_0^{-1/2}\tilde{\Psi}(\cdot, x_{n+1}) \\ &= \sqrt{\frac{\delta|D|}{\tanh(\delta|D|)}} e^{-\delta|D|(1-x_{n+1})}\psi, \end{aligned}$$

we have

$$\int_{\Omega_0} |I_\delta \Lambda_0^{-1/2} \nabla_X \tilde{\Psi}|^2 dX \leq C\|\psi\|^2.$$

This and (5.2) imply that

$$\begin{aligned} \|\Lambda\phi\|^2 &\leq C \int_{\Omega_0} |\Lambda_0^{1/2} PI_\delta \nabla_X \tilde{\Phi}|^2 dX \\ (5.3) \quad &\leq C \left(\int_{\Omega_0} |PI_\delta \nabla_X \Lambda_0^{1/2} \tilde{\Phi}|^2 dX + \int_{\Omega_0} |[\Lambda_0^{1/2}, P] I_\delta \nabla_X \tilde{\Phi}|^2 dX \right). \end{aligned}$$

Set $\Phi_1 := (\Lambda_0^{1/2}\phi)^{\hbar}$ and $\tilde{\Phi}_1 := \Phi_1 \circ \Theta$. Then, it holds that

$$\begin{cases} \nabla_X \cdot I_\delta PI_\delta \nabla_X (\Lambda_0^{1/2} \tilde{\Phi} - \tilde{\Phi}_1) = -\nabla_X \cdot I_\delta [\Lambda_0^{1/2}, P] I_\delta \nabla_X \tilde{\Phi}, \\ (\Lambda_0^{1/2} \tilde{\Phi} - \tilde{\Phi}_1)(\cdot, 1) = 0, \\ \mathbf{e}_{n+1} \cdot I_\delta^2 \nabla_X (\Lambda_0^{1/2} \tilde{\Phi} - \tilde{\Phi}_1)(\cdot, 0) = 0. \end{cases}$$

Therefore, by Green's formula we see that

$$\begin{aligned}
& \int_{\Omega_0} PI_\delta \nabla_X (\Lambda_0^{1/2} \tilde{\Phi} - \tilde{\Phi}_1) \cdot I_\delta \nabla_X (\Lambda_0^{1/2} \tilde{\Phi} - \tilde{\Phi}_1) dX \\
&= - \int_{\Omega_0} (\nabla_X \cdot I_\delta PI_\delta \nabla_X (\Lambda_0^{1/2} \tilde{\Phi} - \tilde{\Phi}_1)) (\Lambda_0^{1/2} \tilde{\Phi} - \tilde{\Phi}_1) dX \\
&= \int_{\Omega_0} (\nabla_X \cdot I_\delta [\Lambda_0^{1/2}, P] I_\delta \nabla_X \tilde{\Phi}) (\Lambda_0^{1/2} \tilde{\Phi} - \tilde{\Phi}_1) dX \\
&= - \int_{\Omega_0} [\Lambda_0^{1/2}, P] I_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \nabla_X (\Lambda_0^{1/2} \tilde{\Phi} - \tilde{\Phi}_1) dX,
\end{aligned}$$

where we used (4.8). This implies that

$$\int_{\Omega_0} |I_\delta \nabla_X (\Lambda_0^{1/2} \tilde{\Phi} - \tilde{\Phi}_1)|^2 dX \leq C \int_{\Omega_0} |[\Lambda_0^{1/2}, P] I_\delta \nabla_X \tilde{\Phi}|^2 dX.$$

Hence, by (5.3) we obtain

$$\|\Lambda\phi\|^2 \leq C \left(\int_{\Omega_0} |I_\delta \nabla_X \tilde{\Phi}_1|^2 dX + \int_{\Omega_0} |[\Lambda_0^{1/2}, P] I_\delta \nabla_X \tilde{\Phi}|^2 dX \right).$$

Here, by Lemma 5.1 and the hypothesis on P we have $\|[\Lambda_0^{1/2}, P]u\| \leq C\|u\|$, so that

$$\begin{aligned}
\|\Lambda\phi\|^2 &\leq C \left(\int_{\Omega_0} |I_\delta \nabla_X \tilde{\Phi}_1|^2 dX + \int_{\Omega_0} |I_\delta \nabla_X \tilde{\Phi}|^2 dX \right) \\
&\leq C((\Lambda\Lambda_0^{1/2}\phi, \Lambda_0^{1/2}\phi) + (\Lambda\phi, \phi)) \\
&\leq C(\|\Lambda_0\phi\|^2 + \|\Lambda_0^{1/2}\phi\|^2),
\end{aligned}$$

where we used Lemmas 4.2–4.4. This shows the desired estimate. \square

Lemma 5.4. *Let $s > n/2 + 1$. Under Assumption 4.1 (A1) and*

$$\sup_{0 \leq x_{n+1} \leq 1} \|\nabla_X \theta(\cdot, x_{n+1})\|_{s+1} \leq M,$$

there exists a constant $C = C(M, c, s) > 0$ independent of δ such that we have

$$\| [J^s, \Lambda] \phi \| \leq C \|\Lambda_0^{1/2} \phi\|_s.$$

Proof. Set $\Phi := \phi^{\hbar}$, $\Phi_s := (J^s \phi)^{\hbar}$, $\tilde{\Phi} := \Phi \circ \Theta$, and $\tilde{\Phi}_s := \Phi_s \circ \Theta$. Then, we have

$$(5.4) \quad \begin{cases} \nabla_X \cdot I_\delta PI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s) = -\nabla_X \cdot I_\delta [J^s, P] I_\delta \nabla_X \tilde{\Phi}, \\ (J^s \tilde{\Phi} - \tilde{\Phi}_s)(\cdot, 1) = 0, \\ \mathbf{e}_{n+1} \cdot I_\delta^2 \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)(\cdot, 0) = 0. \end{cases}$$

and

$$[J^s, \Lambda] \phi = \mathbf{e}_{n+1} \cdot I_\delta^2 \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)(\cdot, 1).$$

We take $\psi \in H^0$ arbitrarily and define $\tilde{\Psi}$ by $\tilde{\Psi}(\cdot, x_{n+1}) = e^{-\delta|D|(1-x_{n+1})}\psi$. Taking the inner product of the equation in (5.4) and $\tilde{\Psi}$ in $L^2(\Omega_0)$ and using Green's formula, we see that

$$\begin{aligned} ([J^s, \Lambda]\phi, \psi) &= \int_{\Omega_0} [J^s, P]I_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \nabla_X \tilde{\Psi} dX \\ &\quad + \int_{\Omega_0} P I_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s) \cdot I_\delta \nabla_X \tilde{\Psi} dX. \end{aligned}$$

In view of $\|J^{-1}I_\delta \nabla_X \tilde{\Psi}\|_{L^2(\Omega_0)}^2 \leq C\|\psi\|^2$, we obtain

$$\begin{aligned} (5.5) \quad &\| [J^s, \Lambda]\phi \| \\ &\leq C(\|J[J^s, P]I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} + \|JPI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)}) \\ &\leq C(\|J^s I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} + \|JI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)}) \\ &\leq C(\|I_\delta \nabla_X \tilde{\Phi}_s\|_{L^2(\Omega_0)} + \|JI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)}). \end{aligned}$$

In the above calculation we used a well-known commutator estimate $\|[J^s, a]u\|_1 \leq C\|a\|_{s+1}\|u\|_s$.

On the other hand, taking the inner product of the equation in (5.4) and $J^2(J^s \tilde{\Phi} - \tilde{\Phi}_s)$ in $L^2(\Omega_0)$ and using Green's formula we obtain

$$\begin{aligned} &\int_{\Omega_0} PJI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s) \cdot JI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s) dX \\ &= - \int_{\Omega_0} [J, P]I_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s) \cdot JI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s) dX \\ &\quad - \int_{\Omega_0} J[J^s, P]I_\delta \nabla_X \tilde{\Phi} \cdot JI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s) dX, \end{aligned}$$

which implies that

$$\begin{aligned} &\|JI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)} \\ &\leq C(\|I_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)} + \|J^s I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}) \\ &\leq C(\|I_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)} + \|I_\delta \nabla_X \tilde{\Phi}_s\|_{L^2(\Omega_0)}). \end{aligned}$$

Here, by the interpolation inequality for any $\epsilon > 0$ we have

$$\begin{aligned} &\|I_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)} \\ &\leq \epsilon \|JI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)} + C_\epsilon \|J^{-s} I_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)} \\ &\leq \epsilon \|JI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)} + C_\epsilon (\|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} + \|I_\delta \nabla_X \tilde{\Phi}_s\|_{L^2(\Omega_0)}). \end{aligned}$$

Therefore,

$$\|JI_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)} \leq C(\|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} + \|I_\delta \nabla_X \tilde{\Phi}_s\|_{L^2(\Omega_0)}),$$

which together with (5.5) implies that

$$\|[J^s, \Lambda]\phi\| \leq C(\|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} + \|I_\delta \nabla_X \tilde{\Phi}_s\|_{L^2(\Omega_0)}).$$

This and Lemmas 4.2–4.4 show the desired estimate. \square

In view of $\|\Lambda\phi\|_s \leq \|\Lambda J^s\phi\| + \|[J^s, \Lambda]\phi\|$ and Lemmas 5.3–5.4, we can obtain the following lemma.

Lemma 5.5. *Under the hypothesis of Lemma 5.4, we have*

$$\|\Lambda\phi\|_s \leq C(\|\Lambda_0\phi\|_s + \|\Lambda_0^{1/2}\phi\|_s),$$

where $C = C(M, c, s) > 0$. Particularly, it holds that $\|\Lambda\phi\|_s \leq C\delta^{-1}\|\phi\|_{s+1}$.

Lemma 5.6. *Let $s \geq 0$ and set $\Phi := \phi^{\hbar}$ and $\tilde{\Phi} := \Phi \circ \Theta$. Under Assumption 4.1 (A1), (A2), and*

$$\|J^s \nabla_X \theta\|_{L^2(\Omega_0)} + \sup_{0 \leq x_{n+1} \leq 1} \|\nabla_X \theta(\cdot, x_{n+1})\|_{s-1/2} \leq M,$$

there exists a constant $C = C(M, c, r, s) > 0$ such that we have

$$\begin{cases} \|J^s I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} \leq C(\|\Lambda_0^{1/2}\phi\|_s + \|I_\delta \nabla_X \tilde{\Phi}\|_{L^\infty(\Omega_0)}), \\ \|J^{r+1} I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} \leq C\|\Lambda_0^{1/2}\phi\|_{r+1}. \end{cases}$$

Proof. Set $\Phi_s := (J^s \phi)^{\hbar}$ and $\tilde{\Phi}_s := \Phi_s \circ \Theta$. Then, we have (5.4). Taking the inner product of (5.4) with $J^s \tilde{\Phi} - \tilde{\Phi}_s$ in $L^2(\Omega_0)$ and using Green's formula, we see that

$$\begin{aligned} & \int_{\Omega_0} P I_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s) \cdot I_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s) dX \\ &= - \int_{\Omega_0} [J^s, P] I_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s) dX, \end{aligned}$$

which together with a commutator estimate $\|[J^s, a]u\| \leq C(|\nabla a|_\infty \|J^{s-1}u\| + \|J^s a\| \|u\|_\infty)$ implies that

$$\begin{aligned} (5.6) \quad & \|I_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)} \leq C\|[J^s, P] I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} \\ & \leq C(\|J^s P\|_{L^2(\Omega_0)} \|I_\delta \nabla_X \tilde{\Phi}\|_{L^\infty(\Omega_0)} \\ & \quad + \|\nabla P\|_{L^\infty(\Omega_0)} \|J^{s-1} I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}). \end{aligned}$$

Note that Assumption 4.1 (A2) and the Sobolev inequality imply the estimate $\|\nabla \nabla_X \theta\|_{L^\infty(\Omega_0)} \leq C$. Hence, it holds that $\|J^s P\|_{L^2(\Omega_0)} + \|\nabla P\|_{L^\infty(\Omega_0)} \leq C$, so that

$$\begin{aligned} & \|J^s I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} \\ & \leq \|I_\delta \nabla_X \tilde{\Phi}_s\|_{L^2(\Omega_0)} + \|I_\delta \nabla_X (J^s \tilde{\Phi} - \tilde{\Phi}_s)\|_{L^2(\Omega_0)} \\ & \leq \|I_\delta \nabla_X \tilde{\Phi}_s\|_{L^2(\Omega_0)} + C(\|I_\delta \nabla_X \tilde{\Phi}\|_{L^\infty(\Omega_0)} + \|J^{s-1} I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}). \end{aligned}$$

This and the interpolation inequality yields that

$$(5.7) \quad \begin{aligned} & \|J^s I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} \\ & \leq C(\|I_\delta \nabla_X \tilde{\Phi}_s\|_{L^2(\Omega_0)} + \|I_\delta \nabla_X \tilde{\Phi}\|_{L^\infty(\Omega_0)} + \|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}). \end{aligned}$$

It follows from Lemmas 4.2–4.4 that $\|I_\delta \nabla_X \tilde{\Phi}_s\|_{L^2(\Omega_0)}$ and $\|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}$ are equivalent to $\|\Lambda_0^{1/2} \phi\|_s$ and $\|\Lambda_0^{1/2} \phi\|$, respectively, so that we obtain the first estimate of the lemma.

Set $\Phi_{r+1} := (J^{r+1} \phi)^h$ and $\tilde{\Phi}_{r+1} := \Phi_{r+1} \circ \Theta$. Since $\|J^{r+1} P(\cdot, x_{n+1})\| \leq C$ for $0 \leq x_{n+1} \leq 1$, in place of (5.6) we have

$$\begin{aligned} \|I_\delta \nabla_X (J^{r+1} \tilde{\Phi} - \tilde{\Phi}_{r+1})\|_{L^2(\Omega_0)} & \leq C\|[J^{r+1}, P]I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} \\ & \leq C\|J^r I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}, \end{aligned}$$

where we used a commutator estimate $\|[J^{r+1}, a]u\| \leq C\|J^{r+1} a\|\|J^r u\|$. Therefore, in place of (5.7) we obtain

$$\|J^{r+1} I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} \leq C(\|I_\delta \nabla_X \tilde{\Phi}_{r+1}\|_{L^2(\Omega_0)} + \|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}),$$

which together with Lemmas 4.2–4.4 yields the second estimate of the lemma. \square

We proceed to give a L^∞ -estimate of $I_\delta \nabla_X \tilde{\Phi}$. To this end, we use (4.6), where the matrix P is defined by (4.5). It follows from (4.7) that

$$A := \left(I_\delta^{-1} \frac{\partial \Theta}{\partial X} I_\delta \right)^T \left(I_\delta^{-1} \frac{\partial \Theta}{\partial X} I_\delta \right) = A_1 + \delta^2 A_2,$$

where A_2 is a matrix whose elements are polynomials of $\nabla_X \theta$ with coefficients which are also polynomials of δ^2 , and

$$A_1 = \begin{pmatrix} E_n & \delta \mathbf{a} \\ \delta \mathbf{a}^T & b \end{pmatrix},$$

where

$$\begin{cases} \mathbf{a} = \partial_{n+1}(\theta_1, \dots, \theta_n)^T + (1 + \partial_{n+1} \theta_{n+1}) \nabla \theta_{n+1}, \\ b = (1 + \partial_{n+1} \theta_{n+1})^2. \end{cases}$$

By the definition of P we have $P = (\det(\frac{\partial \Theta}{\partial X}))^{-1} \tilde{A}$, where \tilde{A} is the adjoint matrix of A and has the form

$$\tilde{A} = \tilde{A}_1 + \delta^2 A_3,$$

where A_3 is a matrix whose elements are polynomials of $\nabla_X \theta$. Moreover, we see that

$$\tilde{A}_1 = \begin{pmatrix} (b - \delta^2 |\mathbf{a}|^2) E_n + \delta^2 \mathbf{a} \mathbf{a}^T & -\delta \mathbf{a} \\ -\delta \mathbf{a}^T & 1 \end{pmatrix}.$$

By these relations and (4.13) we see that the matrix P has the form

$$(5.8) \quad P = \begin{pmatrix} (1 + \partial_{n+1}\theta_{n+1})E_n + \delta^2 P_{11} & \delta \mathbf{p}_{12} \\ \delta \mathbf{p}_{12}^T & (1 + \partial_{n+1}\theta_{n+1})^{-1} + \delta^2 p_{22} \end{pmatrix},$$

where P_{11} , \mathbf{p}_{12} , and p_{22} are $n \times n$, $1 \times n$, and 1×1 matrixes whose elements are polynomials of $\nabla_X \theta$. Moreover, it follows from (4.8) that

$$(5.9) \quad \mathbf{p}_{12}(x, 0) = \mathbf{p}_{12}(x, 1) = \mathbf{0}, \quad p_{22}(x, 0) = p_{22}(x, 1) = 0.$$

Using these notations we can rewrite the first equation in (4.6) as

$$(5.10) \quad \begin{aligned} & \partial_{n+1}((\delta^{-2}(1 + \partial_{n+1}\theta_{n+1})^{-1} + p_{22})\partial_{n+1}\tilde{\Phi}) \\ &= -\nabla \cdot (((1 + \partial_{n+1}\theta_{n+1})E_n + \delta^2 P_{11})\nabla\tilde{\Phi}) \\ & \quad - \nabla \cdot (\mathbf{p}_{12}\partial_{n+1}\tilde{\Phi}) - \partial_{n+1}(\mathbf{p}_{12} \cdot \nabla\tilde{\Phi}). \end{aligned}$$

It follows from this, the boundary condition on $x_{n+1} = 0$, (4.2), and (5.9) that

$$(5.11) \quad \begin{aligned} \partial_{n+1}\tilde{\Phi} &= \int_0^{x_{n+1}} \partial_{n+1}(((1 + \partial_{n+1}\theta_{n+1})^{-1} + \delta^2 p_{22})\partial_{n+1}\tilde{\Phi})dx_{n+1} \\ &= -\delta^2 \int_0^{x_{n+1}} \nabla \cdot (((1 + \partial_{n+1}\theta_{n+1})E_n + \delta^2 P_{11})\nabla\tilde{\Phi})dx_{n+1} \\ & \quad - \delta^2 \int_0^{x_{n+1}} \nabla \cdot (\mathbf{p}_{12}\partial_{n+1}\tilde{\Phi})dx_{n+1} - \delta^2 \mathbf{p}_{12} \cdot \nabla\tilde{\Phi}. \end{aligned}$$

We also have

$$(5.12) \quad \nabla\tilde{\Phi} = \nabla\phi - \int_{x_{n+1}}^1 \nabla\partial_{n+1}\tilde{\Phi}dx_{n+1}.$$

Lemma 5.7. *Let $\Phi = \phi^h$ and $\tilde{\Phi} = \Phi \circ \Theta$. Under Assumption 4.1 (A1) and (A2), there exists a constant $C = C(M, c, r) > 0$ independent of δ such that we have*

$$\|I_\delta \nabla_X \tilde{\Phi}\|_{L^\infty(\Omega_0)} \leq C(\|\nabla\phi\|_r + \delta\|\Lambda_0^{1/2}\phi\|_{r+1}).$$

Proof. Note that the assumptions imply the uniform boundedness of P_{11} , p_{22} , \mathbf{p}_{12} , and their first derivatives with respect to x . It follows from (5.12), the Sobolev inequality, and Lemma 5.6 that

$$\|\nabla\tilde{\Phi}\|_{L^\infty(\Omega_0)} \leq \|\nabla\phi\|_r + \delta\|J^{r+1}I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} \leq \|\nabla\phi\|_r + C\delta\|\Lambda_0^{1/2}\phi\|_{r+1}.$$

Similarly, it follows from (5.11) that

$$\begin{aligned} \delta^{-1}\|\partial_{n+1}\tilde{\Phi}\|_{L^\infty(\Omega_0)} &\leq C\delta(\|J^{r+1}I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} + \|\nabla\tilde{\Phi}\|_{L^\infty(\Omega_0)}) \\ &\leq C\delta\|\Lambda_0^{1/2}\phi\|_{r+1}, \end{aligned}$$

where we used Lemma 5.2. These yield the desired estimate. \square

Remark 5.1. In the case of a flat bottom, applying the maximal principle to the subharmonic function $|I_\delta \nabla_X \Phi|^2$ we see that $\|I_\delta^2 \nabla_X \Phi\|_{L^\infty(\Omega)} = \|I_\delta^2 \nabla_X \Phi\|_{L^\infty(\Gamma)} \leq \sqrt{|\nabla \phi|_\infty^2 + (\delta |\Lambda \phi|_\infty)^2}$.

6. Fréchet derivatives of the Dirichlet-to-Neumann map

The following lemma was obtained by D. Lannes [15].

Lemma 6.1. *The Fréchet derivative of $\Lambda(\eta, b, \delta)$ with respect to η has the form*

$$D_\eta \Lambda(\eta, b, \delta)[\zeta] \phi = -\delta^2 \Lambda(\eta, b, \delta)(Z\zeta) - \nabla \cdot (v\zeta),$$

where

$$(6.1) \quad \begin{cases} Z = (1 + \delta^2 |\nabla \eta|^2)^{-1} (\Lambda(\eta, b, \delta) \phi + \nabla \eta \cdot \nabla \phi), \\ v = \nabla \phi - \delta^2 Z \nabla \eta. \end{cases}$$

We proceed to give estimates of the Fréchet derivatives of Λ in the Sobolev spaces.

Lemma 6.2. *Let $s > n/2$. Under Assumption 4.1 (A1) and*

$$\sup_{0 \leq x_{n+1} \leq 1} \|\nabla_X \theta(\cdot, x_{n+1})\|_{s+1} \leq M,$$

there exists a constant $C = C(M, c, s) > 0$ independent of δ such that we have

$$\|D_\eta^n \Lambda[\zeta_1, \dots, \zeta_n] \phi\|_s \leq C \|\zeta_1\|_{s+3/2} \cdots \|\zeta_n\|_{s+3/2} \|\Lambda_0^{1/2} \phi\|_{s+1}.$$

Similar estimate holds for the Fréchet derivative of Λ with respect to b .

Proof. We only show the estimate in the case $n = 1$, and the general case can be proved in the same way. Set $\Phi := \phi^h$ and $\tilde{\Phi} := \Phi \circ \Theta$. Then, it holds that

$$\begin{cases} \nabla_X \cdot I_\delta P I_\delta \nabla_X \tilde{\Phi} = 0, \\ \tilde{\Phi}(\cdot, 1) = \phi, \quad \mathbf{e}_{n+1} \cdot I_\delta^2 \nabla_X \tilde{\Phi}(\cdot, 1) = \Lambda \phi, \\ \mathbf{e}_{n+1} \cdot I_\delta^2 \nabla_X \tilde{\Phi}(\cdot, 0) = 0. \end{cases}$$

For simplicity, we write $\Lambda_\eta \phi = D_\eta \Lambda[\zeta] \phi$, $\tilde{\Phi}_\eta = D_\eta \tilde{\Phi}[\zeta]$, and $P_\eta = D_\eta P[\zeta]$. Taking the Fréchet derivative of the above equations, we obtain

$$(6.2) \quad \begin{cases} \nabla_X \cdot I_\delta P I_\delta \nabla_X \tilde{\Phi}_\eta = -\nabla_X \cdot I_\delta P_\eta I_\delta \nabla_X \tilde{\Phi}, \\ \tilde{\Phi}_\eta(\cdot, 1) = 0, \quad \mathbf{e}_{n+1} \cdot I_\delta^2 \nabla_X \tilde{\Phi}_\eta(\cdot, 1) = \Lambda_\eta \phi, \\ \mathbf{e}_{n+1} \cdot I_\delta^2 \nabla_X \tilde{\Phi}_\eta(\cdot, 0) = 0. \end{cases}$$

We take $\psi \in H^0$ arbitrarily and define $\tilde{\Psi}$ by $\tilde{\Psi}(\cdot, x_{n+1}) = e^{-\delta|D|(1-x_{n+1})} \psi$. Taking the inner product of the above equation and $J^s \tilde{\Psi}$ in $L^2(\Omega_0)$ and using Green's formula, we see that

$$(J^s \Lambda_\eta \phi, \psi) = \int_{\Omega_0} J^s P I_\delta \nabla_X \tilde{\Phi}_\eta \cdot I_\delta \nabla_X \tilde{\Psi} dX + \int_{\Omega_0} J^s P_\eta I_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \nabla_X \tilde{\Psi} dX.$$

In view of $\|\Lambda_0^{-1/2} I_\delta \nabla_X \tilde{\Psi}\|_{L^2(\Omega_0)} \leq C\|\psi\|$, we obtain

$$(6.3) \quad \begin{aligned} \|\Lambda_\eta \phi\|_s &\leq C(\|\Lambda_0^{1/2} J^s P I_\delta \nabla_X \tilde{\Phi}_\eta\|_{L^2(\Omega_0)} + \|\Lambda_0^{1/2} J^s P_\eta I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}) \\ &\leq C(\|J^{s+1} P I_\delta \nabla_X \tilde{\Phi}_\eta\|_{L^2(\Omega_0)} + \|J^{s+1} P_\eta I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}). \end{aligned}$$

On the other hand, taking the inner product of the first equation in (6.2) and $J^{2(s+1)} \tilde{\Phi}_\eta$ in $L^2(\Omega_0)$ and using Green's formula, we see that

$$\begin{aligned} &\int_{\Omega_0} P J^{s+1} I_\delta \nabla_X \tilde{\Phi}_\eta \cdot J^{s+1} I_\delta \nabla_X \tilde{\Phi}_\eta dX \\ &= - \int_{\Omega_0} [J^{s+1}, P] I_\delta \nabla_X \tilde{\Phi}_\eta \cdot J^{s+1} I_\delta \nabla_X \tilde{\Phi}_\eta dX \\ &\quad - \int_{\Omega_0} J^{s+1} P_\eta I_\delta \nabla_X \tilde{\Phi} \cdot J^{s+1} I_\delta \nabla_X \tilde{\Phi}_\eta dX, \end{aligned}$$

which implies that

$$(6.4) \quad \begin{aligned} &\|J^{s+1} I_\delta \nabla_X \tilde{\Phi}_\eta\|_{L^2(\Omega_0)} \\ &\leq C(\|[J^{s+1}, P] I_\delta \nabla_X \tilde{\Phi}_\eta\|_{L^2(\Omega_0)} + \|J^{s+1} P_\eta I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}) \\ &\leq C(\|J^s I_\delta \nabla_X \tilde{\Phi}_\eta\|_{L^2(\Omega_0)} + \|J^{s+1} P_\eta I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}). \end{aligned}$$

Similarly, taking the inner product of the first equation in (6.2) and $\tilde{\Phi}_\eta$ in $L^2(\Omega_0)$ and using Green's formula, we obtain the estimate $\|I_\delta \nabla_X \tilde{\Phi}_\eta\|_{L^2(\Omega_0)} \leq C\|P_\eta I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}$. Therefore, by the interpolation inequality we obtain

$$\begin{aligned} \|J^{s+1} I_\delta \nabla_X \tilde{\Phi}_\eta\|_{L^2(\Omega_0)} &\leq C\|J^{s+1} P_\eta I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)} \\ &\leq C(\|J^{s+1} P_\eta\|_{L^2(\Omega_0)} \|I_\delta \nabla_X \tilde{\Phi}\|_{L^\infty(\Omega_0)} + \|\nabla P_\eta\|_{L^\infty(\Omega_0)} \|J^{s+1} I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}) \\ &\leq C\|\zeta\|_{s+3/2} \|\Lambda_0^{1/2} \phi\|_{s+1}, \end{aligned}$$

where we used Lemmas 5.2, 5.6, and 5.7. Hence, we obtain the desired estimate. \square

Lemma 6.3. *Let $s > (n+1)/2$. Under Assumption 4.1 (A1) and*

$$\sup_{0 \leq x_{n+1} \leq 1} \|\nabla_X \theta(\cdot, x_{n+1})\|_{s+1/2} \leq M,$$

there exists a constant $C = C(M, c, s) > 0$ independent of δ such that we have

$$\|D_\eta^n \Lambda[\zeta_1, \dots, \zeta_n] \phi\|_s \leq C \delta^{-1/2} \|\zeta_1\|_{s+1} \cdots \|\zeta_n\|_{s+1} \|\Lambda_0^{1/2} \phi\|_{s+1/2}.$$

Similar estimate holds for the Fréchet derivative of Λ with respect to b .

Proof. By (6.3) we have

$$\|\Lambda_\eta \phi\|_s \leq C \delta^{-1/2} (\|J^{s+1/2} P I_\delta \nabla_X \tilde{\Phi}_\eta\|_{L^2(\Omega_0)} + \|J^{s+1/2} P_\eta I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}).$$

Therefore, by the same argument as in the proof of the previous lemma we obtain the desired estimate. \square

Lemma 6.4. *Let $s > n/2 + 2$. Under Assumption 4.1 (A1) and*

$$\|(\eta, b)\|_{s+1} + \sup_{0 \leq x_{n+1} \leq 1} \|\nabla_X \theta(\cdot, x_{n+1})\|_s \leq M,$$

there exists a constant $C = C(M, c, s) > 0$ independent of δ such that we have

$$\|\Lambda\phi\|_s \leq C\delta^{-1/2} \|\Lambda_0^{1/2}\phi\|_{s+1/2}.$$

Proof. It is sufficient to evaluate $\|\Lambda\phi\|_{s-1}$ and $\|\nabla\Lambda\phi\|_{s-1}$. By Lemmas 5.5 and 5.2 we have $\|\Lambda\phi\|_{s-1} \leq C\delta^{-1/2} \|\Lambda_0^{1/2}\phi\|_{s-1/2}$. By the relation $\nabla\Lambda\phi = \Lambda\nabla\phi + D_\eta\Lambda[\nabla\eta]\phi + D_b\Lambda[\nabla b]\phi$ and Lemma 6.3, we see that

$$\begin{aligned} \|\nabla\Lambda\phi\|_{s-1} &\leq C\delta^{-1/2} (\|\Lambda_0^{1/2}\nabla\phi\|_{s-1/2} + \|(\nabla\eta, \nabla b)\|_s \|\Lambda_0^{1/2}\phi\|_{s-1/2}) \\ &\leq C\delta^{-1/2} \|\Lambda_0^{1/2}\phi\|_{s+1/2}. \end{aligned}$$

Therefore, we obtain the desired estimate. \square

Lemma 6.5. *Let $s > n/2 + 1$. Under Assumption 4.1 (A1) and*

$$\|\eta\|_{s+3} + \sup_{0 \leq x_{n+1} \leq 1} \|\nabla_X \theta(\cdot, x_{n+1})\|_{s+2} \leq M,$$

there exists a constant $C = C(M, c, s) > 0$ independent of δ such that we have

$$\|D_\eta^2\Lambda[\zeta_1, \zeta_2]\phi\|_s \leq C\|\zeta_1\|_{s+2}\|\zeta_2\|_{s+1}(\|\nabla\phi\|_{s+1} + \delta^{1/2}\|\Lambda_0^{1/2}\phi\|_{s+3/2}).$$

Proof. By Lemma 6.1 we have

$$D_\eta\Lambda[\zeta_2]\phi = -\delta^2\Lambda(Z\zeta_2) - \nabla \cdot (v\zeta_2),$$

where $Z = (1 + \delta^2|\nabla\eta|^2)^{-1}(\Lambda\phi + \nabla\eta \cdot \nabla\phi)$ and $v = \nabla\phi - \delta^2 Z\nabla\eta$, so that

$$D_\eta^2\Lambda[\zeta_1, \zeta_2]\phi = -\delta^2 D_\eta\Lambda[\zeta_1](Z\zeta_2) - \delta^2\Lambda((D_\eta Z[\zeta_1])\zeta_2) - \nabla \cdot ((D_\eta v[\zeta_1])\zeta_2).$$

By Lemmas 5.5, 6.3, and 5.2 we see that

$$\begin{aligned} &\|D_\eta^2\Lambda[\zeta_1, \zeta_2]\phi\|_s \\ &\leq C(\delta^{3/2}\|\zeta_1\|_{s+1}\|\Lambda_0^{1/2}(Z\zeta_2)\|_{s+1/2} \\ &\quad + \delta\|(D_\eta Z[\zeta_1])\zeta_2\|_{s+1} + \|(D_\eta v[\zeta_1])\zeta_2\|_{s+1}) \\ &\leq C\|\zeta_2\|_{s+1}(\delta(\|\zeta_1\|_{s+1}\|Z\|_{s+1} + \|D_\eta Z[\zeta_1]\|_{s+1}) + \|D_\eta v[\zeta_1]\|_{s+1}). \end{aligned}$$

Here, by Lemmas 5.5 and 5.2 it holds that

$$\|Z\|_{s+1} \leq C(\|\Lambda\phi\|_{s+1} + \|\nabla\phi\|_{s+1}) \leq C\delta^{-1}\|\nabla\phi\|_{s+1}.$$

The Fréchet derivative of Z can be written as

$$\begin{aligned} D_\eta Z[\zeta_1] &= -2\delta^2(1 + \delta^2|\nabla\eta|^2)^{-2}\nabla\eta \cdot \nabla\zeta_1(\Lambda\phi + \nabla\eta \cdot \nabla\phi) \\ &\quad + (1 + \delta^2|\nabla\eta|^2)^{-1}(D_\eta\Lambda[\zeta_1]\phi + \nabla\zeta_1 \cdot \nabla\phi), \end{aligned}$$

so that by Lemmas 5.5, 6.3, and 5.2 we get

$$\begin{aligned} \|D_\eta Z[\zeta_1]\|_{s+1} &\leq C(\delta^2 \|\zeta_1\|_{s+2} (\|\Lambda\phi\|_{s+1} + \|\nabla\phi\|_{s+1}) \\ &\quad + \|D_\eta \Lambda[\zeta_1]\phi\|_{s+1} + \|\zeta_1\|_{s+2} \|\nabla\phi\|_{s+1}) \\ &\leq C\|\zeta_1\|_{s+2} (\|\nabla\phi\|_{s+1} + \delta^{-1/2} \|\Lambda_0^{1/2}\phi\|_{s+3/2}). \end{aligned}$$

Similarly, we see that

$$\begin{aligned} \|D_\eta v[\zeta_1]\|_{s+1} &= \delta^2 \|(D_\eta Z[\zeta_1])\nabla\eta + Z\nabla\zeta_1\|_{s+1} \\ &\leq C\delta \|\zeta_1\|_{s+2} (\|\nabla\phi\|_{s+1} + \delta^{1/2} \|\Lambda_0^{1/2}\phi\|_{s+3/2}). \end{aligned}$$

Therefore, we obtain the desired estimate. \square

7. Energy estimates of a linear system

Following D. Lannes [15], we linearize the equations in (2.14) around (η, ϕ) . Taking a derivative ∂ of the second equation in (2.14), we see that

$$\begin{aligned} 0 &= \partial\phi_t + \partial\eta + \nabla\phi \cdot \nabla\partial\phi \\ &\quad + \delta^4 \nabla\eta \cdot \nabla\partial\eta (1 + \delta^2 |\nabla\eta|^2)^{-2} (\Lambda\phi + \nabla\eta \cdot \nabla\phi)^2 \\ &\quad - \delta^2 (1 + \delta^2 |\nabla\eta|^2)^{-1} (\Lambda\phi + \nabla\eta \cdot \nabla\phi) (\partial\Lambda\phi + \partial(\nabla\eta \cdot \nabla\phi)) \\ (7.1) \quad &= \partial\phi_t + \partial\eta + \nabla\phi \cdot \nabla\partial\phi + \delta^4 Z^2 \nabla\eta \cdot \nabla\partial\eta \\ &\quad - \delta^2 Z (\partial\Lambda\phi + \nabla\partial\eta \cdot \nabla\phi + \nabla\eta \cdot \nabla\partial\phi) \\ &= \partial\phi_t + \partial\eta + (\nabla\phi - \delta^2 Z \nabla\eta) \cdot \nabla\partial\phi \\ &\quad - \delta^2 Z (\nabla\phi - \delta^2 Z \nabla\eta) \cdot \nabla\partial\eta - \delta^2 Z \partial\Lambda\phi, \end{aligned}$$

so that

$$(\partial\phi - \delta^2 Z \partial\eta)_t + (\nabla\phi - \delta^2 Z \nabla\eta) \cdot \nabla(\partial\phi - \delta^2 Z \partial\eta) + (1 + \delta^2 Z_t + \delta^2 v \cdot \nabla Z) \partial\eta = 0,$$

where we used the equation $\partial\eta_t = \partial\Lambda\phi$, which comes from the first equation in (2.14). By Lemma 6.1 we also have

$$\partial\eta_t = \Lambda(\partial\phi - \delta^2 Z \partial\eta) - \nabla \cdot (v\partial\eta) + D_b \Lambda[\partial b]\phi.$$

Introducing new functions ζ and ψ by

$$\zeta := \partial\eta, \quad \psi := \partial\phi - \delta^2 Z \partial\eta,$$

we obtain

$$(7.2) \quad \begin{cases} \zeta_t + \nabla \cdot (v\zeta) - \Lambda\psi = D_b \Lambda[\partial b]\phi, \\ \psi_t + v \cdot \nabla\psi + (1 + \delta^2 Z_t + \delta^2 v \cdot \nabla Z)\zeta = 0. \end{cases}$$

Taking these equations into account, we will consider the following system of linear equations

$$(7.3) \quad \begin{cases} \zeta_t + b_1 \cdot \nabla\zeta - \Lambda\psi = f_1, \\ \psi_t + b_2 \cdot \nabla\psi + a\zeta = f_2, \end{cases}$$

where $a, b_1 = (b_{11}, \dots, b_{1n}), b_2 = (b_{21}, \dots, b_{2n}), f_1, f_2$ are given functions of x and t and may depend on δ , and $\Lambda = \Lambda(\eta, b, \delta)$ is the Dirichlet-to-Neumann map. We assume the function a to be positively definite and define an energy function $E(t)$ by

$$(7.4) \quad E(t) := (a\zeta(t), \zeta(t)) + (\Lambda\psi(t), \psi(t)).$$

Remark 7.1. This energy function is natural, because the water wave problem has a conserved energy defined by

$$H = \int_{\Omega(t)} \frac{1}{2} |\nabla_X \Phi(X)|^2 dX + \int_{\mathbf{R}^n} \frac{g}{2} |\eta(x)|^2 dx = \frac{1}{2} (\Lambda\phi, \phi) + \frac{g}{2} \|\eta\|^2.$$

We mention that the water wave problem has a Hamiltonian structure whose Hamiltonian is H and the canonical variables are η and ϕ . The Hamiltonian formulation of water waves goes back to the work of V. E. Zakharov [27] in the case of deep water.

Let (ζ, ψ) be a solution of (7.3). Then, it holds that

$$(7.5) \quad \begin{aligned} \frac{d}{dt} E(t) &= (a_t \zeta, \zeta) + 2(a\zeta_t, \zeta) + ([\partial_t, \Lambda]\psi, \psi) + 2(\psi_t, \Lambda\psi) \\ &= (a_t \zeta, \zeta) + ((\nabla \cdot (ab_1))\zeta, \zeta) + 2(a f_1, \zeta) \\ &\quad + ([\partial_t, \Lambda]\psi, \psi) - 2(b_2 \cdot \nabla \psi, \Lambda\psi) + 2(f_2, \Lambda\psi). \end{aligned}$$

Lemma 7.1. Under Assumption 4.1 (A1) and

$$(7.6) \quad \|\nabla_X \theta_t(\cdot, t)\|_{L^\infty(\Omega_0)} \leq M,$$

there exists a constant $C = C(M, c) > 0$ independent of δ such that we have

$$|([\partial_t, \Lambda]\phi, \phi)| \leq C(\Lambda\phi, \phi).$$

Proof. Set $\Phi := \phi^{\tilde{h}}$ and $\tilde{\Phi} := \Phi \circ \Theta$. Then, by Lemma 4.2 we have

$$(\Lambda\phi, \phi) = \int_{\Omega} |I_\delta \nabla_X \Phi|^2 dX = \int_{\Omega_0} P I_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \nabla_X \tilde{\Phi} dX,$$

so that

$$\begin{aligned} ([\partial_t, \Lambda]\phi, \phi) &= \frac{d}{dt} (\Lambda\phi, \phi) \\ &= 2 \int_{\Omega_0} P I_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \nabla_X \tilde{\Phi}_t dX + \int_{\Omega_0} P_t I_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \nabla_X \tilde{\Phi} dX. \end{aligned}$$

Since $\tilde{\Phi}(\cdot, 1) = \phi$, we have $\tilde{\Phi}_t(\cdot, 1) = 0$. Therefore, by Green's formula we see that

$$\begin{aligned} &\int_{\Omega_0} P I_\delta \nabla_X \tilde{\Phi} \cdot I_\delta \nabla_X \tilde{\Phi}_t dX \\ &= - \int_{\Omega_0} (\nabla_X \cdot I_\delta P I_\delta \nabla_X \tilde{\Phi}) \tilde{\Phi}_t dX \\ &\quad + (\mathbf{e}_{n+1} \cdot I_\delta^2 \nabla_X \tilde{\Phi}(\cdot, 1), \tilde{\Phi}_t(\cdot, 1)) - (\mathbf{e}_{n+1} \cdot I_\delta^2 \nabla_X \tilde{\Phi}(\cdot, 0), \tilde{\Phi}_t(\cdot, 0)) \\ &= 0. \end{aligned}$$

Hence, we obtain

$$|([\partial_t, \Lambda]\phi, \phi)| \leq \|P_t\|_{L^\infty(\Omega_0)} \|I_\delta \nabla_X \tilde{\Phi}\|_{L^2(\Omega_0)}^2.$$

This together with Lemmas 4.2 and 4.3 implies the desired estimate. \square

The following lemma was given by S. Wu [25]. For the completeness, we will give the proof.

Lemma 7.2. *For the Dirichlet-to-Neumann map $\Lambda = \Lambda(\eta, b, \delta)$ it holds that*

$$|(\phi, \Lambda\psi)| \leq \sqrt{(\phi, \Lambda\phi)} \sqrt{(\psi, \Lambda\psi)}.$$

Proof. Set $\Phi := \phi^h$ and $\Psi := \psi^h$. By Green's formula we see that

$$\begin{aligned} (\Lambda\phi, \psi) &= \int_\Gamma (N \cdot I_\delta^2 \nabla_X \Phi) \Psi dS = \int_\Omega \nabla_X \cdot ((I_\delta^2 \nabla_X \Phi) \Psi) dX \\ &= \int_\Omega I_\delta \nabla_X \Phi \cdot I_\delta \nabla_X \Psi dX. \end{aligned}$$

Therefore, by Lemma 4.2 we obtain

$$|(\Lambda\phi, \psi)| \leq \|I_\delta \nabla_X \Phi\|_{L^2(\Omega)} \|I_\delta \nabla_X \Psi\|_{L^2(\Omega)} = \sqrt{(\phi, \Lambda\phi)} \sqrt{(\psi, \Lambda\psi)}.$$

This shows the desired estimate. \square

Lemma 7.3. *Let $r > n/2$. Under Assumption 4.1 (A1) and*

$$(7.7) \quad \|(\eta, b)\|_{r+2} \leq M,$$

then we have

$$|(\Lambda\phi, v \cdot \nabla\phi)| \leq C_1 \|v\|_{r+1} (\Lambda\phi, \phi).$$

Proof. Set $\Phi := \phi^h$ and let $V = (V_1, \dots, V_n, V_{n+1})^T$ be a vector field on Ω satisfying

$$(7.8) \quad \begin{cases} V_j|_\Gamma = v_j & (1 \leq j \leq n), & V_{n+1}|_\Gamma = \delta v \cdot \nabla\eta, \\ V_{n+1}|_\Sigma = \delta(V_1|_\Sigma, \dots, V_n|_\Sigma)^T \cdot \nabla b. \end{cases}$$

Such a vector field V will be constructed later. Then, it is easy to see that

$$V \cdot I_\delta \nabla_X \Phi|_\Gamma = v \cdot \nabla\phi, \quad V \cdot I_\delta N|_\Gamma = V \cdot I_\delta N|_\Sigma = 0.$$

By these relations and Green's formula we see that

$$\begin{aligned} &(\Lambda\phi, v \cdot \nabla\phi) \\ &= \int_\Gamma (N \cdot I_\delta^2 \nabla_X \Phi) (V \cdot I_\delta \nabla_X \Phi) dS = \int_\Omega \nabla_X \cdot ((I_\delta^2 \nabla_X \Phi) (V \cdot I_\delta \nabla_X \Phi)) dX \\ &= \int_\Omega I_\delta \nabla_X \Phi \cdot (I_\delta \nabla_X V) I_\delta \nabla_X \Phi dX + \frac{1}{2} \int_\Omega V \cdot I_\delta \nabla_X |I_\delta \nabla_X \Phi|^2 dX \\ &= \int_\Omega \left(I_\delta \nabla_X \Phi \cdot (I_\delta \nabla_X V) I_\delta \nabla_X \Phi - \frac{1}{2} (I_\delta \nabla_X \cdot V) |I_\delta \nabla_X \Phi|^2 \right) dX, \end{aligned}$$

where $I_\delta \nabla_X V = (I_\delta \nabla_X V_1, \dots, I_\delta \nabla_X V_{n+1})$. Therefore, we obtain

$$|(\Lambda \phi, v \cdot \nabla \phi)| \leq C \|I_\delta \nabla_X V\|_{L^\infty(\Omega)} \|I_\delta \nabla_X \Phi\|_{L^2(\Omega)}^2 = C \|I_\delta \nabla_X V\|_{L^\infty(\Omega)} (\Lambda \phi, \phi).$$

It remains to construct a vector field V satisfying (7.8) and $\|I_\delta \nabla_X V\|_{L^\infty(\Omega)} \leq C \|v\|_{r+1}$. Such V can be constructed in the form $\tilde{V} = V \circ \Theta$. In view of (4.10) and (7.8), \tilde{V} should satisfy

$$\begin{cases} \tilde{V}_j(\cdot, 1) = v_j \quad (1 \leq j \leq n), & \tilde{V}_{n+1}(\cdot, 1) = \delta v \cdot \nabla \eta, \\ \tilde{V}_{n+1}(\cdot, 0) = \delta(\tilde{V}_1(\cdot, 0), \dots, \tilde{V}_n(\cdot, 0))^T \cdot \nabla b. \end{cases}$$

and $\|I_\delta \nabla_X \tilde{V}\|_{L^\infty(\Omega_0)} \leq C \|v\|_{r+1}$. We take $\varphi \in C^\infty(\mathbf{R})$ satisfying $\varphi(x_{n+1}) = 1$ for $x_{n+1} \geq 1$ and $\varphi(x_{n+1}) = 0$ for $x_{n+1} \leq \frac{1}{2}$, and define \tilde{V} by

$$\begin{aligned} \mathcal{F}[\tilde{V}_j(\cdot, x_{n+1})](\xi) &= e^{-\delta|\xi|(1-x_{n+1})} \hat{v}_j(\xi) \quad (1 \leq j \leq n), \\ \mathcal{F}[\tilde{V}_{n+1}(\cdot, x_{n+1})](\xi) &= \varphi(x_{n+1}) e^{-\delta|\xi|(1-x_{n+1})} \delta \mathcal{F}[v \cdot \nabla \eta](\xi) \\ &\quad + \varphi(1-x_{n+1}) e^{-\delta|\xi|x_{n+1}} \delta \mathcal{F}[(e^{-\delta|D|} v) \cdot \nabla b](\xi). \end{aligned}$$

Then, it is easy to check that \tilde{V} satisfies the required conditions. The proof is complete. \square

Lemma 7.4. *Let $r > n/2$. In addition to Assumption 4.1 (A1), (7.6), and (7.7), we assume that*

$$(7.9) \quad M^{-1} \leq a(x, t) \leq M, \quad \|(a_t, \nabla a)\|_r + \|b_1\|_{r+1} + \|b_2\|_{r+2} \leq M.$$

Then, there exists a constant $C = C(M, c, r) > 0$ independent of δ such that for any smooth solution (ζ, ψ) of (7.3) we have

$$E(t) \leq e^{Ct} E(0) + \int_0^t e^{C(t-\tau)} (\|f_1(\tau)\|^2 + \|\Lambda_0^{1/2} f_2(\tau)\|^2) d\tau.$$

Proof. By (7.5) and Lemmas 7.1–7.3 we obtain

$$\frac{d}{dt} E(t) \leq C E(t) + \|f_1(t)\|^2 + \|\Lambda_0^{1/2} f_2(t)\|^2,$$

so that the desired energy estimate comes from Gronwall's inequality. \square

We proceed to estimate a high order energy function $E_s(t)$ defined by

$$(7.10) \quad E_s(t) := (a J^s \zeta(t), J^s \zeta(t)) + (\Lambda J^s \psi(t), J^s \psi(t)).$$

Let (ζ, ψ) be a solution of (7.3). Then, it holds that

$$(7.11) \quad \begin{aligned} \frac{d}{dt} E_s(t) &= (a_t J^s \zeta, J^s \zeta) + 2(a J^s \zeta_t, J^s \zeta) \\ &\quad + ([\partial_t, \Lambda] J^s \psi, J^s \psi) + 2(\Lambda J^s \psi_t, J^s \psi) \\ &\leq |a^{-1} a_t|_\infty (a J^s \zeta, J^s \zeta) + C(\Lambda J^s \psi, J^s \psi) \\ &\quad - 2(a J^s b_1 \cdot \nabla \zeta, J^s \zeta) + 2(a J^s \Lambda \psi, J^s \zeta) + 2(a J^s f_1, J^s \zeta) \\ &\quad - 2(\Lambda J^s b_2 \cdot \nabla \psi, J^s \psi) - 2(\Lambda J^s a \zeta, J^s \psi) + 2(\Lambda J^s f_2, J^s \psi). \end{aligned}$$

Lemma 7.5. *Let $s > n/2 + 1$. Then, there exists a constant $C = C(s) > 0$ independent of δ such that for any $j = 1, \dots, n$ we have*

$$\|\Lambda_0^{1/2}[J^s, \psi]\partial_j\phi\| \leq C\|\psi\|_{s+1}\|\Lambda_0^{1/2}\phi\|_s.$$

Proof. Put $u := \Lambda_0^{1/2}[J^s, \psi]\partial_j\phi$. Then, we have

$$\begin{aligned} \hat{u}(\xi) &= \frac{\sqrt{\delta^{-1}|\xi| \tanh(\delta|\xi|)}}{(2\pi)^n} \int_{\mathbf{R}^n} \hat{\psi}(\xi - \eta) ((1 + |\xi|)^s - (1 + |\eta|)^s) i\eta_j \hat{\phi}(\eta) d\eta \\ &= \frac{1}{(2\pi)^n} \int_{\mathbf{R}^n} \hat{\psi}(\xi - \eta) ((1 + |\xi|)^s - (1 + |\eta|)^s) \\ &\quad \times i\eta_j \sqrt{\delta^{-1}|\eta| \tanh(\delta|\eta|)} \hat{\phi}(\eta) d\eta \\ &\quad + \frac{1}{(2\pi)^n} \int_{\mathbf{R}^n} (\sqrt{\delta^{-1}|\xi| \tanh(\delta|\xi|)} - \sqrt{\delta^{-1}|\eta| \tanh(\delta|\eta|)}) \\ &\quad \times \hat{\psi}(\xi - \eta) ((1 + |\xi|)^s - (1 + |\eta|)^s) i\eta_j \hat{\phi}(\eta) d\eta. \end{aligned}$$

Therefore, by (5.1) we obtain

$$\begin{aligned} |\hat{u}(\xi)| &\leq \frac{1}{(2\pi)^n} \int_{\mathbf{R}^n} |\hat{\psi}(\xi - \eta)| |(1 + |\xi|)^s - (1 + |\eta|)^s| |\eta| \widehat{\Lambda_0^{1/2}\phi}(\eta) d\eta \\ &\quad + C \int_{\mathbf{R}^n} |\xi - \eta| |\hat{\psi}(\xi - \eta)| |(1 + |\xi|)^s - (1 + |\eta|)^s| |\widehat{\nabla\phi}(\eta)| d\eta. \end{aligned}$$

In view of the inequality $|(1 + |\xi|)^s - (1 + |\eta|)^s| \leq C|\xi - \eta|((1 + |\xi - \eta|)^{s-1} + (1 + |\eta|)^{s-1})$ and the first estimate in Lemma 5.2, the above estimate and Hausdorff-Young's inequality give the desired estimate. \square

Lemma 7.6. *Let $s > n/2 + 1$. In addition to Assumption 4.1 (A1), (7.6), (7.7), and (7.9), we assume that*

$$(7.12) \quad \|(\nabla a, b_1)\|_s + \|b_2\|_{s+1} + \sup_{0 \leq x_{n+1} \leq 1} \|\nabla_X \theta(\cdot, x_{n+1})\|_{s+1} \leq M.$$

Then, there exists a constant $C = C(M, c, s) > 0$ independent of δ such that for any smooth solution (ζ, ψ) of (7.3) we have

$$E_s(t) \leq e^{Ct} E_s(0) + \int_0^t e^{C(t-\tau)} (\|f_1(\tau)\|_s^2 + \|\Lambda_0^{1/2} f_2(\tau)\|_s^2) d\tau.$$

Proof. We will evaluate each term in the right hand side of (7.11). It is easy to see that

$$\begin{aligned} |(aJ^s b_1 \cdot \nabla \zeta, J^s \zeta)| &= \left| -\frac{1}{2} ((\nabla \cdot ab_1) J^s \zeta, J^s \zeta) + (a[J^s, b_1] \cdot \nabla \zeta, J^s \zeta) \right| \\ &\leq \frac{1}{2} |\nabla \cdot (ab_1)|_\infty \|\zeta\|_s^2 + C|a|_\infty \|b_1\|_s \|\zeta\|_s^2. \end{aligned}$$

By Lemma 7.2, we have

$$\begin{aligned} & |(aJ^s\Lambda\psi, J^s\zeta) - (\Lambda J^s a\zeta, J^s\psi)| \\ &= |([J^s, \Lambda]\psi, aJ^s\zeta) - ([J^s, a]\zeta, \Lambda J^s\psi)| \\ &\leq |a|_\infty \|\zeta\|_s \|[J^s, \Lambda]\psi\| + \sqrt{(\Lambda J^s\psi, J^s\psi)} \sqrt{(\Lambda [J^s, a]\zeta, [J^s, a]\zeta)}. \end{aligned}$$

Here, by Lemmas 5.4 and 4.4

$$\|[J^s, \Lambda]\psi\|^2 \leq C\|\Lambda_0^{1/2}\psi\|_s^2 = C(\Lambda_0 J^s\psi, J^s\psi) \leq C(\Lambda J^s\psi, J^s\psi),$$

by Lemmas 4.4 and 5.2

$$\sqrt{(\Lambda [J^s, a]\zeta, [J^s, a]\zeta)} \leq C\|\Lambda_0^{1/2}[J^s, a]\zeta\| \leq C\|[J^s, a]\zeta\|_1 \leq C\|\nabla a\|_s \|\zeta\|_s,$$

so that we obtain

$$|(aJ^s\Lambda\psi, J^s\zeta) - (\Lambda J^s a\zeta, J^s\psi)| \leq C(|a|_\infty + \|\nabla a\|_s) \|\zeta\|_s \sqrt{(\Lambda J^s\psi, J^s\psi)}.$$

By Lemmas 7.2 and 7.3 we have

$$\begin{aligned} & |(\Lambda J^s b_2 \cdot \nabla\psi, J^s\psi)| \\ &= |(\Lambda [J^s, b_2] \cdot \nabla\psi, J^s\psi) + (b_2 \cdot \nabla J^s\psi, \Lambda J^s\psi)| \\ &\leq \sqrt{(\Lambda J^s\psi, J^s\psi)} \sqrt{(\Lambda [J^s, b_2] \cdot \nabla\psi, [J^s, b_2] \cdot \nabla\psi)} + C(\Lambda J^s\psi, J^s\psi). \end{aligned}$$

Here, by Lemmas 4.4 and 7.7 we get

$$\begin{aligned} (\Lambda [J^s, b_2] \cdot \nabla\psi, [J^s, b_2] \cdot \nabla\psi) &\leq C\|\Lambda_0^{1/2}[J^s, b_2] \cdot \nabla\psi\|^2 \\ &\leq C\|b_2\|_{s+1}^2 (\Lambda J^s\psi, J^s\psi). \end{aligned}$$

By the above estimates and Lemma 7.2, it follows from (7.11) that

$$\frac{d}{dt} E_s(t) \leq C E_s(t) + \|f_1(t)\|_s^2 + \|\Lambda_0^{1/2} f_2(t)\|_s^2,$$

so that the desired energy estimate comes from Gronwall's inequality. \square

8. Reduction to a quasi-linear system

In this section we reduce the equations

$$(8.1) \quad \begin{cases} \eta_t - \Lambda\phi = 0, \\ \phi_t + \eta + \frac{1}{2}|\nabla\phi|^2 - \frac{1}{2}\delta^2(1 + \delta^2|\nabla\eta|^2)^{-1}(\Lambda\phi + \nabla\eta \cdot \nabla\phi)^2 = 0 \end{cases}$$

to a quasi-linear system of equations. In the same way as in (7.1), differentiating the second equation in (8.1) with respect to x_i we obtain

$$\partial_i\phi_t + \partial_i\eta + (\nabla\phi - \delta^2 Z\nabla\eta) \cdot (\nabla\partial_i\phi - \delta^2 Z\nabla\partial_i\eta) - \delta^2 Z\partial_i\Lambda\phi = 0.$$

Differentiating this with respect to x_j and x_k , we see that

$$\begin{aligned}
& \partial_{ijk}\phi_t + \partial_{ijk}\eta \\
& + v \cdot \{ \nabla \partial_{ijk}\phi - \delta^2 (Z \nabla \partial_{ijk}\eta + (\partial_{jk}Z) \nabla \partial_i\eta + (\partial_j Z) \nabla \partial_{ki}\eta + (\partial_k Z) \nabla \partial_{ij}\eta) \} \\
& + (\partial_j v) \cdot \{ \nabla \partial_{ik}\phi - \delta^2 (Z \nabla \partial_{ik}\eta + (\partial_k Z) \nabla \partial_i\eta) \} \\
& + (\partial_k v) \cdot \{ \nabla \partial_{ij}\phi - \delta^2 (Z \nabla \partial_{ij}\eta + (\partial_j Z) \nabla \partial_i\eta) \} \\
& + \{ \nabla \partial_{jk}\phi - \delta^2 (Z \nabla \partial_{jk}\phi \\
& \quad + (\partial_j Z) \nabla \partial_k\eta + (\partial_k Z) \nabla \partial_j\eta + (\partial_{jk}Z) \nabla \eta) \} \cdot (\nabla \partial_i\phi - \delta^2 Z \nabla \partial_i\eta) \\
& - \delta^2 \{ (\partial_j Z) \partial_{ik}\Lambda\phi + (\partial_k Z) \partial_{ij}\Lambda\phi + (\partial_{jk}Z) \partial_i\Lambda\phi + Z \partial_{ijk}\Lambda\phi \} = 0.
\end{aligned}$$

Here, by Lemma 6.1 we have

$$\partial_{ik}\Lambda = \partial_k(\Lambda(\partial_i\phi - \delta^2 Z \partial_i\eta) - (\nabla \cdot v) \partial_i\eta + D_b \Lambda[\partial_i b] \phi) - (\partial_k v) \cdot \nabla \partial_i\eta - v \cdot \nabla \partial_{ik}\eta,$$

so that

$$\begin{aligned}
& (\partial_{ijk}\phi - \delta^2 Z \partial_{ijk}\eta)_t + v \cdot \nabla (\partial_{ijk}\phi - \delta^2 Z \partial_{ijk}\eta) + (1 + \delta^2 Z_t + \delta^2 v \cdot \nabla Z) \partial_{ijk}\eta \\
& = \delta^2 (\partial_{jk}Z) v \cdot \nabla \partial_i\eta - (\partial_j v) \cdot \{ (\nabla \partial_{ik}\phi - \delta^2 Z \nabla \partial_{ik}\eta) - \delta^2 (\partial_k Z) \nabla \partial_i\eta \} \\
& \quad - (\partial_k v) \cdot \{ (\nabla \partial_{ij}\phi - \delta^2 Z \nabla \partial_{ij}\eta) - \delta^2 (\partial_j Z) \nabla \partial_i\eta \} \\
& \quad - (\nabla \partial_i\phi - \delta^2 Z \nabla \partial_i\eta) \cdot \{ (\nabla \partial_{jk}\phi - \delta^2 Z \nabla \partial_{jk}\eta) \\
& \quad \quad - \delta^2 ((\partial_j Z) \nabla \partial_k\eta + (\partial_k Z) \nabla \partial_j\eta + (\partial_{jk}Z) \nabla \eta) \} + \delta^2 (\partial_{jk}Z) \partial_i\Lambda\phi \\
& \quad + \delta^2 (\partial_j Z) \{ \partial_k(\Lambda(\partial_i\phi - \delta^2 Z \partial_i\eta) - (\nabla \cdot v) \partial_i\eta + D_b \Lambda[\partial_i b] \phi) - (\partial_k v) \cdot \nabla \partial_i\eta \} \\
& \quad + \delta^2 (\partial_k Z) \{ \partial_j(\Lambda(\partial_i\phi - \delta^2 Z \partial_i\eta) - (\nabla \cdot v) \partial_i\eta + D_b \Lambda[\partial_i b] \phi) - (\partial_j v) \cdot \nabla \partial_i\eta \}.
\end{aligned}$$

Now, we write $u = (\eta, b)$ and denote by Λ_n the n -th Fréchet derivative of the Dirichlet-to-Neumann map Λ with respect to u . Differentiating the first equation in (8.1) yields that

$$\begin{aligned}
\partial_{ijk}\Lambda\phi & = \Lambda \partial_{ijk}\phi + \Lambda_1[\partial_{ijk}u]\phi + \Lambda_1[\partial_i u] \partial_{jk}\phi + \Lambda_1[\partial_j u] \partial_{ki}\phi + \Lambda_1[\partial_k u] \partial_{ij}\phi \\
& \quad + \Lambda_1[\partial_{ij}u] \partial_k\phi + \Lambda_1[\partial_{jk}u] \partial_i\phi + \Lambda_1[\partial_{ki}u] \partial_j\phi \\
& \quad + \Lambda_2[\partial_i u, \partial_j u] \partial_k\phi + \Lambda_2[\partial_j u, \partial_k u] \partial_i\phi + \Lambda_2[\partial_k u, \partial_i u] \partial_j\phi \\
& \quad + \Lambda_2[\partial_{ij}u, \partial_k u] \phi + \Lambda_2[\partial_{jk}u, \partial_i u] \phi + \Lambda_2[\partial_{ki}u, \partial_j u] \phi \\
& \quad + \Lambda_3[\partial_i u, \partial_j u, \partial_k u] \phi.
\end{aligned}$$

Here, by Lemma 6.1 we have

$$\Lambda_1[\partial_{ijk}u]\phi = -\delta^2 \Lambda(Z \partial_{ijk}\eta) - \nabla \cdot (v \partial_{ijk}\eta) + D_b \Lambda[\partial_{ijk}b] \phi.$$

Therefore, introducing new functions ζ_{ijk} and ψ_{ijk} by

$$\zeta_{ijk} := \partial_{ijk}\eta, \quad \psi_{ijk} := \partial_{ijk}\phi - \delta^2 Z \partial_{ijk}\eta,$$

we obtain

$$(8.2) \quad \begin{cases} \partial_t \zeta_{ijk} + v \cdot \nabla \zeta_{ijk} - \Lambda \psi_{ijk} = f_1^{ijk}, \\ \partial_t \psi_{ijk} + v \cdot \nabla \psi_{ijk} + a \zeta_{ijk} = f_2^{ijk}, \end{cases}$$

where $a = 1 + \delta^2 Z_t + \delta^2 v \cdot \nabla Z$ and

$$\begin{aligned}
f_1^{ijk} &= -(\nabla \cdot v)\zeta_{ijk} + D_b \Lambda[\partial_{ijk} b]\phi \\
&\quad + \Lambda_1[\partial_i u]\partial_{jk}\phi + \Lambda_1[\partial_j u]\partial_{ki}\phi + \Lambda_1[\partial_k u]\partial_{ij}\phi \\
&\quad + \Lambda_1[\partial_{ij} u]\partial_k\phi + \Lambda_1[\partial_{jk} u]\partial_i\phi + \Lambda_1[\partial_{ki} u]\partial_j\phi \\
&\quad + \Lambda_2[\partial_i u, \partial_j u]\partial_k\phi + \Lambda_2[\partial_j u, \partial_k u]\partial_i\phi + \Lambda_2[\partial_k u, \partial_i u]\partial_j\phi \\
&\quad + \Lambda_2[\partial_{ij} u, \partial_k u]\phi + \Lambda_2[\partial_{jk} u, \partial_i u]\phi + \Lambda_2[\partial_{ki} u, \partial_j u]\phi \\
&\quad + \Lambda_3[\partial_i u, \partial_j u, \partial_k u]\phi, \\
f_2^{ijk} &= \delta^2(\partial_{jk} Z)v \cdot \nabla \partial_i \eta - (\partial_j v) \cdot \{(\nabla \partial_{ik}\phi - \delta^2 Z \nabla \partial_{ik}\eta) - \delta^2(\partial_k Z)\nabla \partial_i \eta\} \\
&\quad - (\partial_k v) \cdot \{(\nabla \partial_{ij}\phi - \delta^2 Z \nabla \partial_{ij}\eta) - \delta^2(\partial_j Z)\nabla \partial_i \eta\} \\
&\quad - (\nabla \partial_i \phi - \delta^2 Z \nabla \partial_i \eta) \cdot \{(\nabla \partial_{jk}\phi - \delta^2 Z \nabla \partial_{jk}\eta) \\
&\quad \quad - \delta^2((\partial_j Z)\nabla \partial_k \eta + (\partial_k Z)\nabla \partial_j \eta + (\partial_{jk} Z)\nabla \eta)\} + \delta^2(\partial_{jk} Z)\partial_i \Lambda \phi \\
&\quad + \delta^2(\partial_j Z)\{\partial_k(\Lambda(\partial_i \phi - \delta^2 Z \partial_i \eta) - (\nabla \cdot v)\partial_i \eta + D_b \Lambda[\partial_i b]\phi) \\
&\quad \quad - (\partial_k v) \cdot \nabla \partial_i \eta\} \\
&\quad + \delta^2(\partial_k Z)\{\partial_j(\Lambda(\partial_i \phi - \delta^2 Z \partial_i \eta) - (\nabla \cdot v)\partial_i \eta + D_b \Lambda[\partial_i b]\phi) \\
&\quad \quad - (\partial_j v) \cdot \nabla \partial_i \eta\}.
\end{aligned}$$

Setting $\zeta := (\zeta_{ijk})$ and $\psi := (\psi_{ijk})$, we can rewrite (8.2) as

$$(8.3) \quad \begin{cases} \partial_t \zeta + v \cdot \nabla \zeta - \Lambda \psi = f_1, \\ \partial_t \psi + v \cdot \nabla \psi + a \zeta = f_2, \end{cases}$$

where f_1 and f_2 can be written symbolically as

$$\begin{aligned}
f_1 &= -(\nabla \cdot v)\zeta + D_b \Lambda[\partial^3 b]\phi + 3\Lambda_1[\partial u]\partial^2 \phi + 3\Lambda_1[\partial^2 u]\partial \phi + 3\Lambda_2[\partial u, \partial u]\partial \phi \\
&\quad + 3\Lambda_2[\partial^2 u, \partial u]\phi + \Lambda_3[\partial u, \partial u, \partial u]\phi, \\
f_2 &= \delta^2(\partial^2 Z)v \cdot \nabla \partial \eta - 2(\partial v) \cdot (\psi - \delta^2(\partial Z)\nabla \partial \eta) \\
&\quad - (\nabla \partial \phi - \delta^2 Z \nabla \partial \eta) \cdot \{\psi - \delta^2(2(\partial Z)\nabla \partial \eta + (\partial^2 Z)\nabla \eta)\} + \delta^2(\partial^2 Z)\partial \Lambda \phi \\
&\quad + 2\delta^2(\partial Z)\{\partial(\Lambda v - (\nabla \cdot v)\partial \eta + D_b \Lambda[\partial b]\phi) - (\partial v) \cdot \nabla \partial \eta\}.
\end{aligned}$$

We proceed to give a uniform estimate of the coefficients v and a , and the remainder terms f_1 and f_2 .

Lemma 8.1. *Let $s > n/2$. There exists a constant $C = C(s) > 0$ independent of δ such that we have*

$$\|\Lambda_0^{1/2}(\phi\psi)\|_s \leq C(\|\phi\|_s \|\Lambda_0^{1/2}\psi\|_s + \|\Lambda_0^{1/2}\phi\|_s \|\psi\|_s).$$

Proof. In view of the fact that $\frac{\sqrt{\alpha \tanh \alpha}}{\alpha}$ is equivalent to $\frac{1}{\sqrt{1+\alpha}}$ for $\alpha \geq 0$, we easily obtain $\frac{\sqrt{(\alpha+\beta) \tanh(\alpha+\beta)}}{\alpha+\beta} \leq C(\frac{\sqrt{\alpha \tanh \alpha}}{\alpha} + \frac{\sqrt{\beta \tanh \beta}}{\beta})$ for $\alpha, \beta \geq 0$. Therefore, for any $\xi, \eta \in \mathbf{R}^n$ we have

$$\sqrt{\delta^{-1}|\xi| \tanh(\delta|\xi|)} \leq C(\sqrt{\delta^{-1}|\xi - \eta| \tanh(\delta|\xi - \eta|)} + \sqrt{\delta^{-1}|\eta| \tanh(\delta|\eta|)}).$$

Put $u := \Lambda_0^{1/2}(\phi\psi)$. Then, we have

$$\begin{aligned} |\hat{u}(\xi)| &= \left| \frac{1}{(2\pi)^n} \int_{\mathbf{R}^n} \sqrt{\delta^{-1}|\xi| \tanh(\delta|\xi|)} \hat{\phi}(\xi - \eta) \hat{\psi}(\eta) d\eta \right| \\ &\leq C \int_{\mathbf{R}^n} (|\widehat{\Lambda_0^{1/2}\phi}(\xi - \eta)| |\hat{\psi}(\eta)| + |\hat{\phi}(\xi - \eta)| |\widehat{\Lambda_0^{1/2}\psi}(\eta)|) d\eta. \end{aligned}$$

Therefore, Hausdorff-Young's inequality gives the desired estimate. \square

In the following we will use the notation $\partial\phi = (\partial_j\phi)$, $\partial^2\phi = (\partial_{ij}\phi)$, $\partial^3\phi = (\partial_{ijk}\phi)$, etc.

Lemma 8.2. *Let $s > (n+1)/2$, $M, c_1 > 0$ and suppose that*

$$(8.4) \quad \begin{cases} E \equiv \|\eta\|_{s+3} + \|\nabla\phi\|_{s+2} \\ \quad + \|(\partial^3\phi - \delta^2 Z \partial^3\eta, \Lambda_0^{1/2}(\partial^3\phi - \delta^2 Z \partial^3\eta))\|_s \\ \leq M, \\ \|b\|_{s+7/2} \leq M, \quad 1 + \eta(x) - b(x) \geq c_1 \text{ for } x \in \mathbf{R}^n, \end{cases}$$

where $\partial^3\phi - \delta^2 Z \partial^3\eta = (\partial_{ijk}\phi - \delta^2 Z \partial_{ijk}\eta)$. Then, there exist positive constants $\delta_2 = \delta_2(M, c_1, s)$ and $C = C(M, c_1, s)$ such that for any $\delta \in (0, \delta_2]$ we have

$$\begin{cases} \|Z\|_{s+1} + \delta^{1/2}\|Z\|_{s+2} + \delta\|\Lambda_0^{1/2}Z\|_{s+2} \leq CE, \\ \|v\|_{s+2} + \|\Lambda_0^{1/2}v\|_{s+2} \leq CE. \end{cases}$$

Proof. By Lemma 4.5, for any $\delta \in (0, \delta_1]$ we can construct a diffeomorphism Θ satisfying the Assumption 4.1 (A1)–(A2) and

$$\|J^{s+5/2}\nabla_X\theta\|_{L^2(\Omega_0)} + \sup_{0 \leq x_{n+1} \leq 1} \|\nabla_X\theta(\cdot, x_{n+1})\|_{s+2} \leq C.$$

Therefore, by the definition (6.1) of (v, Z) and Lemmas 5.5 and 5.2 we easily obtain $\|(v, Z)\|_{s+1} \leq CE$. By Lemmas 6.4 and 5.2 we obtain $\delta^{1/2}\|Z\|_{s+2} \leq CE$. Therefore,

$$\|v\|_{s+2} \leq \|v\|_s + \|\partial^2 v\|_s \leq \|v\|_s + \|\partial^3\phi - \delta^2 Z \partial^3\eta\|_s + C\delta^2\|Z\|_{s+2}\|\eta\|_{s+2} \leq CE.$$

It remains to evaluate $\|\Lambda_0^{1/2}v\|_{s+2}$ and $\delta\|\Lambda_0^{1/2}\nabla Z\|_{s+1}$. By the definition of Z and Lemma 6.1, we see that

$$\begin{aligned} \partial_j Z &= -2\delta^2(\nabla\eta \cdot \nabla\partial_j\eta)(1 + \delta^2|\nabla\eta|^2)^{-2}(\Lambda\phi + \nabla\eta \cdot \nabla\phi) \\ &\quad + (1 + \delta^2|\nabla\eta|^2)^{-1}(\Lambda(\partial_j\phi - \delta^2 Z \partial_j\eta) \\ &\quad \quad - \nabla \cdot (v\partial_j\eta) + D_b\Lambda[\partial_j b]\phi + \partial_j(\nabla\eta \cdot \nabla\phi)) \\ &= (1 + \delta^2|\nabla\eta|^2)^{-1}\{\Lambda(\partial_j\phi - \delta^2 Z \partial_j\eta) - \nabla \cdot (v\partial_j\eta) \\ &\quad \quad - 2\delta^2 Z \nabla\eta \cdot \nabla\partial_j\eta + D_b\Lambda[\partial_j b]\phi + \partial_j(\nabla\eta \cdot \nabla\phi)\} \\ &= (1 + \delta^2|\nabla\eta|^2)^{-1}\{\delta^2|\nabla\eta|^2\partial_j Z + \Lambda(\partial_j\phi - \delta^2 Z \partial_j\eta) \\ &\quad \quad + D_b\Lambda[\partial_j b]\phi + \nabla\eta \cdot \partial_j v - (\nabla \cdot v)\partial_j\eta\}, \end{aligned}$$

which implies the expression

$$(8.5) \quad \partial_j Z = \Lambda(\partial_j \phi - \delta^2 Z \partial_j \eta) + D_b \Lambda[\partial_j b] \phi + \nabla \eta \cdot \partial_j v - (\nabla \cdot v) \partial_j \eta.$$

Hence, by Lemma 8.1 we obtain

$$\begin{aligned} \delta \|\Lambda_0^{1/2} \nabla Z\|_{s+1} &\leq C \delta (\|\Lambda_0^{1/2} \Lambda v\|_{s+1} + \|\Lambda_0^{1/2} D_b \Lambda[\nabla b] \phi\|_{s+1} \\ &\quad + \|\Lambda_0^{1/2} \nabla v\|_{s+1} \|\nabla \eta\|_{s+1} + \|\nabla v\|_{s+1} \|\Lambda_0^{1/2} \nabla \eta\|_{s+1}). \end{aligned}$$

Here, by Lemmas 5.2 and 6.4 $\delta \|\Lambda_0^{1/2} \Lambda v\|_{s+1} \leq \delta^{1/2} \|\Lambda v\|_{s+3/2} \leq C \|\Lambda_0^{1/2} v\|_{s+2}$, and by Lemmas 5.2 and 6.3 $\delta \|\Lambda_0^{1/2} D_b \Lambda[\nabla b] \phi\|_{s+1} \leq \delta^{1/2} \|D_b \Lambda[\nabla b] \phi\|_{s+3/2} \leq C \|\delta\|_{s+7/2} \|\nabla \phi\|_{s+2}$. Therefore, we get

$$\delta \|\Lambda_0^{1/2} \nabla Z\|_{s+1} \leq C (\|\Lambda_0^{1/2} v\|_{s+2} + E).$$

Similarly, we see that

$$\begin{aligned} \|\Lambda_0^{1/2} v\|_{s+2} &\leq \|\Lambda_0^{1/2} v\|_s + \|\Lambda_0^{1/2} \partial^2 v\|_s \\ &\leq C (\|v\|_{s+1} + \|\Lambda_0^{1/2} (\partial^3 \phi - \delta^2 Z \partial^3 \phi)\|_s \\ &\quad + \delta^2 (\|\Lambda_0^{1/2} \nabla Z\|_{s+1} \|\eta\|_{s+2} + \|Z\|_{s+2} \|\Lambda_0^{1/2} \eta\|_{s+2})) \\ &\leq C (\delta^2 \|\Lambda_0^{1/2} \nabla Z\|_{s+1} + E). \end{aligned}$$

These two estimates imply that if we take $\delta_2 \in (0, \delta_1]$ sufficiently small, then for any $\delta \in (0, \delta_2]$ we have $\delta \|\Lambda_0^{1/2} \nabla Z\|_{s+1} + \|\Lambda_0^{1/2} v\|_{s+2} \leq CE$. The proof is complete. \square

Lemma 8.3. *In addition to hypothesis of Lemma 8.2 we assume that $\|b\|_{s+9/2} \leq M$. Then, there exists a constant $C = C(M, c, s) > 0$ independent of δ such that we have*

$$\|f_1\|_s + \|\Lambda_0^{1/2} f_2\|_s \leq CE.$$

Proof. In view of Lemmas 6.2 and 5.2 we easily obtain $\|D_b \Lambda[\partial^3 b] \phi\|_s \leq C \|\delta\|_{s+9/2} \|\nabla \phi\|_{s+1}$. By Lemmas 6.2 and 6.3 we see that

$$\begin{aligned} \|\Lambda_1[\partial u] \partial^2 \phi\|_s &\leq \|\Lambda_1[\partial u] (\partial^2 \phi - \delta^2 Z \partial^2 \eta)\|_s + \delta^2 \|\Lambda_1[\partial u] (Z \partial^2 \eta)\|_s \\ &\leq C (\|\partial u\|_{s+3/2} \|\Lambda_0^{1/2} (\partial^2 \phi - \delta^2 Z \partial^2 \eta)\|_{s+1} \\ &\quad + \delta^{3/2} \|\partial u\|_{s+1} \|\Lambda_0^{1/2} (Z \partial^2 \eta)\|_{s+1/2}) \\ &\leq C \|u\|_{s+5/2} (\|\Lambda_0^{1/2} \psi\|_s + \|\nabla \phi\|_{s+2} + \delta \|Z\|_{s+1} \|\eta\|_{s+3}), \end{aligned}$$

where $\psi = \partial^3 \phi - \delta^2 Z \partial^3 \eta$. By Lemma 6.1 we have

$$\Lambda_1[\partial^2 u] \partial \phi = -\delta^2 \Lambda(Z_1 \partial^2 \eta) - \nabla \cdot (v_1 \partial^2 \eta) + D_b \Lambda[\partial^2 b] \partial \phi,$$

where

$$\begin{cases} Z_1 = (1 + \delta^2 |\nabla \eta|^2)^{-1} (\Lambda \partial \phi + \nabla \eta \cdot \nabla \partial \phi), \\ v_1 = \nabla \partial \phi - \delta^2 Z_1 \nabla \eta. \end{cases}$$

Hence, by Lemmas 5.5, 5.2, and 6.2 we obtain

$$\begin{aligned} \|\Lambda_1[\partial^2 u] \partial \phi\|_s &\leq C(\delta \|Z_1 \partial^2 \eta\|_{s+1} + \|v_1 \partial^2 \eta\|_{s+1} + \|\partial^2 b\|_{s+3/2} \|\Lambda_0^{1/2} \partial \phi\|_{s+1}) \\ &\leq C(\delta \|\Lambda \partial \phi\|_{s+1} + \|\nabla \phi\|_{s+2}) \leq C \|\nabla \phi\|_{s+2}. \end{aligned}$$

We can directly evaluate $\Lambda_2[\partial u, \partial u] \partial \phi$ and $\Lambda_3[\partial u, \partial u, \partial u] \phi$ by Lemma 6.2, and $\Lambda_2[\partial^2 u, \partial u] \phi$ by Lemma 6.5. Combining the above estimates and those obtained in Lemma 8.2 yields the estimate for f_1 . By Lemmas 8.1, 6.2–6.4 and 5.2, and the estimates obtained in Lemma 8.2, we see that

$$\begin{aligned} \|\Lambda_0^{1/2} f_2\|_s &\leq C\{\delta^2 \|\Lambda_0^{1/2} Z\|_{s+2} + \|\Lambda_0^{1/2} v\|_{s+1} + \|\Lambda_0^{1/2} \psi\|_s + \|\Lambda_0^{1/2} (\partial^2 \phi - \delta^2 Z \partial^2 \eta)\|_s \\ &\quad + \delta^2 (\|Z\|_{s+2} \|\Lambda_0^{1/2} \Lambda \phi\|_{s+1} + \|\Lambda_0^{1/2} \Lambda v\|_{s+1} + \|\Lambda_0^{1/2} D_b \Lambda[\partial b] \phi\|_{s+1})\} \\ &\leq C\{E + \|\partial^2 \phi - \delta^2 Z \partial^2 \eta\|_{s+1} \\ &\quad + \delta^{3/2} (\|\Lambda \phi\|_{s+3/2} + \|\Lambda v\|_{s+3/2} + \|D_b \Lambda[\partial b] \phi\|_{s+3/2})\} \\ &\leq C(E + \delta (\|\Lambda_0^{1/2} \phi\|_{s+2} + \|\Lambda_0^{1/2} v\|_{s+2} + \|\partial b\|_{s+5/2} \|\Lambda_0^{1/2} \phi\|_{s+2})) \leq CE. \end{aligned}$$

The proof is complete. \square

In the following lemma we consider the case where η and ϕ depend also on the time t .

Lemma 8.4. *Under the hypothesis of Lemma 8.2, there exists a constant $C = C(M, c, s) > 0$ independent of δ such that for any $\delta \in (0, \delta_2]$ we have*

$$\begin{cases} \delta \|Z_t\|_{s+1} + \|v_t\|_{s+1} \leq C \|(\eta_t, \phi_t)\|_{s+2}, \\ \delta \|Z_{tt}\|_s + \|v_{tt}\|_s \leq C (\|\eta_{tt}\|_s + \delta \|\eta_{tt}\|_{s+1} + \|\phi_{tt}\|_{s+1} + \|(\eta_t, \phi_t)\|_{s+2}^2). \end{cases}$$

Proof. By the definition (6.1) of (Z, v) we have

$$\begin{aligned} Z_t &= (1 + \delta^2 |\nabla \eta|^2)^{-1} (\Lambda \phi_t + D_\eta \Lambda[\eta_t] \phi \\ &\quad + \nabla \eta_t \cdot \nabla \phi + \nabla \eta \cdot \nabla \phi_t - 2\delta^2 Z \nabla \eta \cdot \nabla \eta_t), \\ v_t &= \nabla \phi_t - \delta^2 (Z \nabla \eta_t + Z_t \nabla \eta). \end{aligned}$$

Therefore, by Lemmas 5.5, 6.3, and 8.2 we obtain the estimate for (Z_t, v_t) . Similarly, in view of

$$\begin{aligned} Z_{tt} &= (1 + \delta^2 |\nabla \eta|^2)^{-1} (\Lambda \phi_{tt} + D_\eta \Lambda[\eta_{tt}] \phi + 2D_\eta \Lambda[\eta_t] \phi_t + D_\eta^2 \Lambda[\eta_t, \eta_t] \phi \\ &\quad + \nabla \eta_{tt} \cdot \nabla \phi + \nabla \eta \cdot \nabla \phi_{tt} + 2\nabla \eta_t \cdot \nabla \phi_t \\ &\quad - 4\delta^2 Z_t \nabla \eta \cdot \nabla \eta_t - 2\delta^2 Z (\nabla \eta \cdot \nabla \eta_{tt} + |\nabla \eta_t|^2)), \\ v_{tt} &= \nabla \phi_{tt} - \delta^2 (Z_{tt} \nabla \eta + Z \nabla \eta_{tt} + 2Z_t \nabla \eta_t), \end{aligned}$$

we obtain the estimate for (Z_{tt}, v_{tt}) . \square

9. Proof of the main theorems

In this section we will give a proof of Theorems 2.1 and 3.1. The existence of the solution for fixed $\delta > 0$ can be proved by using approximate equations and taking the limit. See [26, 24, 25, 15] for details. We can also show the dependence of the solution on the initial data, that is, the well-posedness of the initial value problem. Therefore, to show Theorem 2.1 it is sufficient to derive a priori estimates of the smooth solution $(\eta^\delta, \phi^\delta)$ for a time interval $[0, T]$ independent of δ .

Suppose that $(\eta^\delta, \phi^\delta)$ is the solution of (2.14) and (2.15) and satisfies

$$(9.1) \quad \begin{cases} \mathcal{E}(t) \equiv \|\eta^\delta(t)\|_{s+3}^2 + \|\nabla\phi^\delta(t)\|_{s+2}^2 \\ \quad + \|\Lambda_0^{1/2}(\partial^3\phi^\delta(t) - \delta^2 Z^\delta \partial^3\eta^\delta(t))\|_s^2 \\ \leq N_1, \\ \|\eta^\delta(t)\|_{s+2}^2 + \|\nabla\phi^\delta(t)\|_{s+1}^2 \leq N_2, \\ 1 + \eta^\delta(x, t) - b(x) > c_0/2 \quad \text{for } x \in \mathbf{R}^n, 0 \leq t \leq T, 0 < \delta \leq \delta_0, \end{cases}$$

where Z^δ is determined by (6.1) from $(\eta^\delta, \phi^\delta)$ and positive constants N_1, N_2, T , and δ_0 will be determined later. In the following we simply write the constants depending only on (M_0, N_1, c_0, s) and (M_0, N_2, c_0, s) by C_1 and C_2 , respectively. By Lemmas 4.5 and 4.4, there exists a small $\delta_1 = \delta_1(M_0, N_2, c_0, s) > 0$ such that for any $\delta \in (0, \delta_1]$ and $\varphi \in H^1$ we have

$$(9.2) \quad C_2^{-1}(\Lambda\varphi, \varphi) \leq (\Lambda_0\varphi, \varphi) \leq C_2(\Lambda\varphi, \varphi).$$

By Lemmas 8.2 and 8.3, there exists a small $\delta_2 = \delta_2(M_0, N_1, c_0, s) \in (0, \delta_1]$ such that we have

$$(9.3) \quad \begin{cases} \|Z^\delta\|_{s+1} + \delta^{1/2}\|Z^\delta\|_{s+2} + \delta\|\Lambda_0^{1/2}Z^\delta\|_{s+2} \leq C_1 \\ \|v^\delta\|_{s+2} + \|\Lambda_0^{1/2}v^\delta\|_{s+2} \leq C_1, \\ \|f_1\|_s^2 + \|\Lambda_0^{1/2}f_2\|_s^2 \leq C_1\mathcal{E} \quad \text{for } 0 \leq t \leq T, 0 < \delta \leq \min\{\delta_0, \delta_2\}. \end{cases}$$

In view of

$$\begin{aligned} \eta_t^\delta &= \Lambda\phi^\delta, \quad \partial_j\eta_t^\delta = \Lambda(\partial_j\phi^\delta - \delta^2 Z^\delta \partial_j\eta^\delta) - \nabla \cdot (v^\delta \partial_j\eta^\delta) + D_b\Lambda[\partial_j b]\phi^\delta, \\ \eta_{tt}^\delta &= \Lambda(\phi_{tt} - \delta^2 Z^\delta \eta_{tt}) - \nabla \cdot (v^\delta \eta_{tt}), \\ \phi_t^\delta &= -\eta^\delta - \frac{1}{2}|\nabla\phi^\delta|^2 + \frac{1}{2}\delta^2(1 + \delta^2|\nabla\eta^\delta|^2)(Z^\delta)^2, \\ \phi_{tt}^\delta &= -\eta_{tt}^\delta - \nabla\phi^\delta \cdot \nabla\phi_{tt}^\delta + \delta^4(\nabla\eta^\delta \cdot \nabla\eta_{tt}^\delta)(Z^\delta)^2 + \delta^2(1 + \delta^2|\nabla\eta^\delta|^2)Z^\delta Z_{tt}^\delta, \end{aligned}$$

in the same way as the proof of Lemma 8.2 we obtain

$$(9.4) \quad \begin{cases} \|(\eta_t^\delta(t), \phi_t^\delta(t))\|_{s+2} + \|\eta_{tt}^\delta(t)\|_s + \delta\|\eta_{tt}^\delta(t)\|_{s+1} + \|\phi_{tt}^\delta(t)\|_{s+1} \leq C_1, \\ \|(\eta_t^\delta(t), \phi_t^\delta(t))\|_{s+2}^2 \leq C_1\mathcal{E}(t) \end{cases}$$

for $0 \leq t \leq T$ and $0 \leq \delta \leq \min\{\delta_2, \delta_0\}$. Therefore, by Lemma 8.4

$$\delta(\|Z_t^\delta\|_{s+1} + \|Z_{tt}^\delta\|_s) + \|v_t^\delta\|_{s+1} + \|v_{tt}^\delta\|_s \leq C_1.$$

Particularly, for $a^\delta = 1 + \delta^2(Z_t^\delta + v^\delta \cdot \nabla Z^\delta)$ we have $\|\nabla a^\delta\|_s + \|a_t^\delta\|_s \leq C_1$ for $0 \leq t \leq T$ and $0 \leq \delta \leq \min\{\delta_2, \delta_0\}$. Moreover, by the Sobolev inequality

$$\delta|Z_t^\delta + v^\delta \cdot \nabla Z^\delta|_\infty \leq C\delta(\|Z_t^\delta\|_s + \|v^\delta\|_s\|Z^\delta\|_{s+1}) \leq C_1.$$

Hence, setting $\delta_3 = \min\{\delta_2, (2C_1)^{-1}\}$ we obtain $1/2 \leq a^\delta(x, t) \leq 2$ for $x \in \mathbf{R}^n$, $0 \leq t \leq T$, and $0 < \delta \leq \min\{\delta_3, \delta_0\}$. Now, we can apply the energy estimate obtained in Lemma 7.6 to the quasi-linear system (8.3) and obtain

$$\begin{aligned} & \|\zeta^\delta(t)\|_s^2 + \|\Lambda_0^{1/2}\psi^\delta(t)\|_s^2 \\ & \leq C_2 e^{C_1 t} (\|\zeta^\delta(0)\|_s^2 + \|\Lambda_0^{1/2}\psi^\delta(0)\|_s^2) + C_1 \int_0^t e^{C_1(t-\tau)} \mathcal{E}(\tau) d\tau, \end{aligned}$$

where we used (9.2). By the second estimate in (9.4) we easily obtain

$$\begin{aligned} \|\eta^\delta(t)\|_{s+2}^2 + \|\nabla\phi^\delta(t)\|_{s+1}^2 & \leq \|\eta_0^\delta\|_{s+2}^2 + \|\nabla\phi_0^\delta\|_{s+1}^2 + C_1 \int_0^t \mathcal{E}(\tau) d\tau \\ & \leq \|\eta_0^\delta\|_{s+2}^2 + \|\nabla\phi_0^\delta\|_{s+1}^2 + C_1 t. \end{aligned}$$

Here, by Lemma 5.2 we see that

$$\begin{aligned} \|\nabla\partial^3\phi^\delta(t)\|_{s-1} & \leq 2\|\Lambda_0^{1/2}\psi^\delta(t)\|_{s-1/2} + 2\delta^2\|\Lambda_0^{1/2}(Z^\delta(t)\zeta^\delta(t))\|_{s-1/2} \\ & \leq 2\|\Lambda_0^{1/2}\psi^\delta(t)\|_s + C_1\delta^{3/2}\|\zeta^\delta(t)\|_s. \end{aligned}$$

Therefore, we obtain

$$\mathcal{E}(t) \leq (C_2 + C_1\delta)e^{C_1 t} (\|\eta_0^\delta\|_{s+3+1/2}^2 + \|\nabla\phi_0^\delta\|_{s+3}^2) + C_1 \int_0^t e^{C_1(t-\tau)} \mathcal{E}(\tau) d\tau,$$

which together with Gronwall's inequality yields that

$$\mathcal{E}(t) \leq (C_2 + C_1\delta)e^{C_1 T} (\|\eta_0^\delta\|_{s+3+1/2}^2 + \|\nabla\phi_0^\delta\|_{s+3}^2).$$

Moreover, we have $|\eta^\delta(t) - \eta_0^\delta|_\infty \leq C \int_0^t \|\eta_t^\delta(\tau)\|_s d\tau \leq c_0 C_1 t$. By setting $N_2 = \|\eta_0^\delta\|_{s+2}^2 + \|\nabla\phi_0^\delta\|_{s+1}^2 + 1$, $N_1 = 4C_2(\|\eta_0^\delta\|_{s+3+1/2}^2 + \|\nabla\phi_0^\delta\|_{s+3}^2)$, $\delta_0 = \min\{\delta_3, C_1^{-1}C_2\}$, and $T = (2C_1)^{-1} (< C_1^{-1} \log 2)$, we see that the estimates in (9.1) holds for $0 \leq t \leq T$ and $0 < \delta \leq \delta_0$. By (9.4) we also obtain a uniform bound for $\|(\eta_t^\delta(t), \phi_t^\delta(t))\|_{s+2}$. The proof of Theorem 2.1 is complete.

We proceed to prove Theorem 3.1. To this end, we first expand the Dirichlet-to-Neumann map $\Lambda = \Lambda(\eta, b, \delta)$ with respect to δ^2 . The next lemma is a mathematically rigorous version of the formal expansion (3.2).

Lemma 9.1. *Let $s > n/2$. Under Assumption 4.1 (A1) and*

$$\|J^{s+2}\nabla_X\theta\|_{L^2(\Omega_0)} + \sup_{0 \leq x_{n+1} \leq 1} \|\nabla_X\theta(\cdot, x_{n+1})\|_{s+3/2} \leq M,$$

there exists a constant $C = C(M, c, s) > 0$ independent of δ such that we have

$$\|\Lambda\phi + \nabla \cdot ((1 + \eta - b)\nabla\phi)\|_s \leq C\delta^2(\|\Lambda_0^{1/2}\phi\|_{s+2} + \|\nabla\phi\|_{s+1}).$$

Proof. Set $\Phi := \phi^{\hbar}$ and $\tilde{\Phi} := \Phi \circ \Theta$. Then, we have (4.6). Since $\partial_{n+1}\tilde{\Phi}(\cdot, 0) = 0$ and $\delta^{-2}\partial_{n+1}\tilde{\Phi}(\cdot, 1) = \Lambda\phi$, we see that

$$\begin{aligned} \Lambda\phi &= \int_0^1 \partial_{n+1}((\delta^{-2}(1 + \partial_{n+1}\theta_{n+1})^{-1} + p_{22})\partial_{n+1}\tilde{\Phi})dx_{n+1} \\ &= - \int_0^1 \nabla \cdot ((1 + \partial_{n+1}\theta_{n+1})E_n + \delta^2 P_{11})\nabla\tilde{\Phi} dx_{n+1} \\ &\quad - \int_0^1 \nabla \cdot (p_{12}\partial_{n+1}\tilde{\Phi})dx_{n+1}, \end{aligned}$$

where we used (5.9) and (5.10). By (4.2) we see that $\tilde{\Phi}(\cdot, x_{n+1}) = \phi - \int_{x_{n+1}}^1 \partial_{n+1}\tilde{\Phi}(\cdot, y)dy$ and $\int_0^1 (1 + \partial_{n+1}\theta_{n+1})dx_{n+1} = 1 + \eta - b$, so that

$$\begin{aligned} &\Lambda\phi + \nabla \cdot ((1 + \eta - b)\nabla\phi) \\ &= \int_0^1 \nabla \cdot \left((1 + \partial_{n+1}\theta_{n+1}) \int_{x_{n+1}}^1 \nabla\partial_{n+1}\tilde{\Phi}(\cdot, y)dy \right) dx_{n+1} \\ &\quad - \delta^2 \int_0^1 \nabla \cdot P_{11}\nabla\tilde{\Phi}dx_{n+1} - \int_0^1 \nabla \cdot (p_{12}\partial_{n+1}\tilde{\Phi})dx_{n+1}. \end{aligned}$$

Therefore, we obtain

$$\|\Lambda\phi + \nabla \cdot ((1 + \eta - b)\nabla\phi)\|_s \leq C(\delta^2\|J^{s+1}\nabla\tilde{\Phi}\|_{L^2(\Omega_0)} + \|J^{s+1}\partial_{n+1}\tilde{\Phi}\|_{L^2(\Omega_0)}).$$

On the other hand, it follows from (5.11) that

$$\|J^{s+1}\partial_{n+1}\tilde{\Phi}\|_{L^2(\Omega_0)} \leq C\delta^2\|J^{s+2}I_\delta\nabla_X\tilde{\Phi}\|_{L^2(\Omega_0)}.$$

These estimates together with Lemmas 5.6 and 5.7 imply the desired estimate. \square

By the uniform estimate obtained in Theorem 2.1, Lemma 9.1, and the standard compactness argument, we see that as $\delta \rightarrow +0$

$$(\eta^\delta, \nabla\phi^\delta) \rightarrow (\eta^0, u^0) \text{ weakly* in } L^\infty(0, T; H^{s+3} \times H^{s+2}),$$

where (η^0, u^0) is a unique solution of the shallow water equations (3.3) with initial conditions $(\eta^0, u^0)|_{t=0} = (\eta_0^0, u_0^0)$. Moreover, taking the limit $\delta \rightarrow +0$ of the identity $\text{rot}\nabla\phi^\delta = 0$ we see that u^0 satisfies the irrotational condition

(3.4). Next, we will show the strong convergence. It follows from (2.14), (3.3), and (3.4) that

$$(9.5) \quad \begin{cases} (\eta^\delta - \eta^0)_t + \nabla \cdot ((1 + \eta^\delta - b)(\nabla\phi^\delta - u^0) + (\eta^\delta - \eta^0)u^0) = \delta^2 f_3^\delta, \\ (\nabla\phi^\delta - u^0)_t + \nabla(\eta^\delta - \eta^0) + \frac{1}{2}\nabla((\nabla\phi^\delta + u^0) \cdot (\nabla\phi^\delta - u^0)) = \delta^2 f_4^\delta, \end{cases}$$

where

$$\begin{aligned} f_3^\delta &= \delta^{-2}(\Lambda\phi^\delta + \nabla \cdot ((1 + \eta^\delta - b)\nabla\phi^\delta)), \\ f_4^\delta &= \frac{1}{2}\nabla((1 + \delta^2|\nabla\eta^\delta|^2)^{-1}(\Lambda\phi^\delta + \nabla\eta^\delta \cdot \nabla\phi^\delta)) \\ &\quad \left(= \frac{1}{2}\nabla((1 + \delta^2|\nabla\eta^\delta|^2)(Z^\delta)^2) \right). \end{aligned}$$

By (9.3) and Lemma 9.1, we easily have $\|(f_3^\delta(t), f_4^\delta(t))\|_s \leq C$ for $0 \leq t \leq T$ and $0 < \delta \leq \delta_0$. Taking these equations into account, we will consider the following system of linear equations

$$(9.6) \quad \begin{cases} \zeta_t + \nabla \cdot (aw + b_1\zeta) = f_1, \\ w_t + \nabla\zeta + \nabla(b_2 \cdot w) = f_2, \end{cases}$$

where $a, b_1 = (b_{11}, \dots, b_{1n}), b_2 = (b_{21}, \dots, b_{2n}), f_1$, and f_2 are given function of x and t .

Lemma 9.2. *Let $s > n/2$ and suppose that*

$$M^{-1} \leq a(x, t) \leq M, \quad \|(a_t, \nabla a)\|_s + \|(b_1, b_2)\|_{s+1} \leq M.$$

Then, there exists a constant $C = C(M, s) > 0$ such that for any smooth solution (ζ, w) of (9.6) satisfying the irrotational condition $\text{rot} w = 0$ we have

$$\|(\zeta(t), w(t))\|_s^2 \leq C e^{Ct} \|(\zeta(0), w(0))\|_s^2 + C \int_0^t e^{C(t-\tau)} \|(f_1(\tau), f_2(\tau))\|_s^2 d\tau.$$

Proof. We define an energy function $E_s(t)$ by

$$E_s(t) := \|\zeta(t)\|_s^2 + (aJ^s w(t), J^s w(t)),$$

which is equivalent to $\|(\zeta(t), w(t))\|_s^2$. Then, we see that

$$\begin{aligned} \frac{d}{dt} E_s(t) &= 2(J^s \zeta_t, J^s \zeta) + 2(aJ^s w_t, J^s w) + (a_t J^s w, J^s w) \\ &= -2(\nabla \cdot ([J^s, a]w), J^s \zeta) - 2(\nabla \cdot ([J^s, b_1]\zeta), J^s \zeta) + ((\nabla \cdot b_1)J^s \zeta, J^s \zeta) \\ &\quad + 2(J^s f_1, J^s \zeta) - 2(a\nabla([J^s, b_2] \cdot w), J^s w) + ((\nabla \cdot (ab_2))J^s w, J^s w) \\ &\quad - 2(a(J^s w \cdot \nabla)b_2, J^s w) + 2(aJ^s f_2, J^s w) + (a_t J^s w, J^s w) \\ &\leq C E_s(t) + \|(f_1(t), f_2(t))\|_s^2. \end{aligned}$$

Therefore, the desired energy estimate comes from Gronwall's inequality. \square

Applying the energy estimate to (9.5) we obtain

$$\|\eta^\delta(t) - \eta^0(t)\|_s + \|\nabla\phi^\delta(t) - u^0(t)\|_s \leq C(\|\eta_0^\delta - \eta_0^0\|_s + \|\nabla\phi_0^\delta - u_0^0\|_s + \delta^2)$$

for $0 \leq t \leq T$ and $0 < \delta \leq \delta_0$ with a constant C independent of δ and t . This shows the strong convergence of the solution $(\eta^\delta, \nabla\phi^\delta)$ in $C([0, T]; H^s)$. Since we have a uniform bound of the solution $(\eta^\delta, \nabla\phi^\delta)$ in $C([0, T]; H^{s+3} \times H^{s+2})$, by the interpolation inequality we obtain the strong convergence of the solution in $C([0, T]; H^{s+3-\epsilon} \times H^{s+2-\epsilon})$ for each $\epsilon > 0$. The latter part of the theorem comes directly from the above estimate. The proof of Theorem 3.1 is complete.

DEPARTMENT OF MATHEMATICS
 FACULTY OF SCIENCE AND TECHNOLOGY
 KEIO UNIVERSITY
 3-14-1 HIYOSHI, KOHOKU-KU
 YOKOHAMA 223-8522, JAPAN
 e-mail: iguchi@math.keio.ac.jp

References

- [1] G. B. Airy, *Tides and waves*, Encyclopaedia Metropolitana, London **5** (1845), 241–396.
- [2] B. Alvarez-Samaniego and D. Lannes, *Large time existence for 3D water-waves and asymptotics*, Invent. Math. **171** (2008), 485–541.
- [3] W. Craig, *An existence theory for water waves and the Boussinesq and Korteweg-de Vries scaling limits*, Comm. Partial Differential Equations **10** (1985), 787–1003.
- [4] W. Craig, P. Guyenne, D. P. Nicholls and C. Sulem, *Hamiltonian long wave expansions for water waves over a rough bottom*, Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **461** (2005), 839–873.
- [5] L. C. Evans, *Partial differential equations*, Grad. Stud. Math. **19**, American Mathematical Society, Providence, RI, 1998.
- [6] K. O. Friedrichs, *On the derivation of the shallow water theory*, Appendix to: “The formulation of breakers and bores” by J. J. Stoker in Comm. Pure Appl. Math. **1** (1948), 1–87.
- [7] T. Iguchi, *A long wave approximation for capillary-gravity waves and an effect of the bottom*, Comm. Partial Differential Equations **32** (2007), 37–85.
- [8] ———, *A mathematical justification of the forced Korteweg-de Vries equation for capillary-gravity waves*, Kyushu J. Math. **60** (2006), 267–303.

- [9] T. Kano, *Une théorie trois-dimensionnelle des ondes de surface de l'eau et le développement de Friedrichs*, J. Math. Kyoto Univ. **26** (1986), 101–155 and 157–175 [French].
- [10] T. Kano and T. Nishida, *Sur les ondes de surface de l'eau avec une justification mathématique des équations des ondes en eau peu profonde*, J. Math. Kyoto Univ. **19** (1979), 335–370 [French].
- [11] ———, *Water waves and Friedrichs expansion*, Recent topics in nonlinear PDE, 39–57, North-Holland Math. Stud. **98**, North-Holland, Amsterdam, 1984.
- [12] ———, *A mathematical justification for Korteweg-de Vries equation and Boussinesq equation of water surface waves*, Osaka J. Math. **23** (1986), 389–413.
- [13] D. J. Korteweg and G. de Vries, *On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves*, Phil. Mag. **39** (1895), 422–443.
- [14] H. Lamb, *Hydrodynamics*, 6th edition, Cambridge University Press.
- [15] D. Lannes, *Well-posedness of the water-waves equations*, J. Amer. Math. Soc. **18** (2005), 605–654.
- [16] Y. A. Li, *A shallow-water approximation to the full water wave problem*, Comm. Pure Appl. Math. **59** (2006), 1225–1285.
- [17] V. I. Nalimov, *The Cauchy-Poisson problem*, Dinamika Splošn. Sredy **18** (1974), 104–210 [Russian].
- [18] L. V. Ovsjannikov, *To the shallow water theory foundation*, Arch. Mech. **26** (1974), 407–422.
- [19] ———, *Cauchy problem in a scale of Banach spaces and its application to the shallow water theory justification*, Applications of methods of functional analysis to problems in mechanics, 426–437, Lecture Notes in Math. **503**, Springer, Berlin, 1976.
- [20] G. Schneider and C. E. Wayne, *The long-wave limit for the water wave problem I, The case of zero surface tension*, Comm. Pure Appl. Math. **53** (2000), 1475–1535.
- [21] ———, *The rigorous approximation of long-wavelength capillary-gravity waves*, Arch. Rational Mech. Anal. **162** (2002), 247–285.
- [22] J. J. Stoker, *Water waves: the mathematical theory with application*, A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York.
- [23] F. Ursell, *The long-wave paradox in the theory of gravity waves*, Proc. Cambridge Philos. Soc. **49** (1953), 685–694.

- [24] S. Wu, *Well-posedness in Sobolev spaces of the full water wave problem in 2-D*, *Invent. Math.* **130** (1997), 39–72.
- [25] ———, *Well-posedness in Sobolev spaces of the full water wave problem in 3-D*, *J. Amer. Math. Soc.* **12** (1999), 445–495.
- [26] H. Yosihara, *Gravity waves on the free surface of an incompressible perfect fluid of finite depth*, *Publ. RIMS Kyoto Univ.* **18** (1982), 49–96.
- [27] V. E. Zakharov, *Stability of periodic waves of finite amplitude on the surface of a deep fluid*, *J. Appl. Mech. Tech. Phys.* **9** (1968), 190–194.