



Document details

< Back to results | 1 of 1

↗ Export ↴ Download 🖨️ Print ✉️ E-mail 📄 Save to PDF ☆ Add to List More... >

View at Publisher

Critical Care Medicine
2020, Pages E380-E390

A Single-Center Prospective Observational Study Comparing Resting Energy Expenditure in Different Phases of Critical Illness: Indirect Calorimetry Versus Predictive Equations

(📄 Article in press ?)

Tah, P.C.^{a,b}, Lee, Z.-Y.^a, Poh, B.K.^c, Abdul Majid, H.^{d,e,f}, Hakumat-Rai, V.-R.^{g,h}, Mat Nor, M.B.ⁱ, Kee, C.C.^j, Zaman, M.K.^k, Hasan, M.S.^a ✉️ 👤

^aDepartment of Anesthesiology, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia

^bDepartment of Dietetics, University of Malaya Medical Centre, Kuala Lumpur, Malaysia

^cNutritional Sciences Programme, Centre for Community Health Studies (ReaCH), Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

View additional affiliations ▾

Abstract

▾ View references (47)

Objectives: Several predictive equations have been developed for estimation of resting energy expenditure, but no study has been done to compare predictive equations against indirect calorimetry among critically ill patients at different phases of critical illness. This study aimed to determine the degree of agreement and accuracy of predictive equations among ICU patients during acute phase (≤ 5 d), late phase (6-10 d), and chronic phase (≥ 11 d). **Design:** This was a single-center prospective observational study that compared resting energy expenditure estimated by 15 commonly used predictive equations against resting energy expenditure measured by indirect calorimetry at different phases. Degree of agreement between resting energy expenditure calculated by predictive equations and resting energy expenditure measured by indirect calorimetry was analyzed using intraclass correlation coefficient and Bland-Altman analyses. Resting energy expenditure values calculated from predictive equations differing by $\pm 10\%$ from resting energy expenditure measured by indirect calorimetry was used to assess accuracy. A score ranking method was developed to determine the best predictive equations. **Setting:** General Intensive Care Unit, University of Malaya Medical Centre. **Patients:** Mechanically ventilated critically ill patients. **Interventions:** None. **Measurements and Main Results:** Indirect calorimetry was measured thrice during acute, late, and chronic phases among 305, 180, and 91 ICU patients, respectively. There were significant differences ($F= 3.447$; $p = 0.034$) in mean resting energy expenditure measured by indirect calorimetry among the three phases. Pairwise comparison showed mean resting energy expenditure measured by indirect calorimetry in late phase ($1,878 \pm 517$ kcal) was significantly higher than during acute phase ($1,765 \pm 456$ kcal) ($p = 0.037$). The predictive equations with the best agreement and accuracy for acute phase was Swinamer (1990), for late phase was Brandi (1999) and Swinamer (1990), and for chronic phase was Swinamer (1990). None of the resting energy expenditure calculated from predictive equations showed very good agreement or accuracy. **Conclusions:** Predictive equations tend to either over- or underestimate resting energy expenditure at different phases. Predictive equations with "dynamic" variables and respiratory data had better agreement with resting energy expenditure measured by indirect calorimetry compared with predictive equations developed for healthy adults or predictive equations based on "static" variables. Although none of the resting energy expenditure calculated from predictive equations had very good agreement, Swinamer (1990) appears to provide relatively good agreement across three phases and could be used to predict resting energy expenditure when indirect calorimetry is not available. © 2020 by the Society of Critical Care Medicine and Wolters Kluwer Health, Inc. All Rights Reserved.

Author keywords

basal metabolism calorimetry critical care critical illness nutritional requirements predictive equations

Metrics ⓘ View all metrics >



PlumX Metrics ▾

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

Set citation alert >

Set citation feed >

Related documents

Predictive equations versus measured energy expenditure by indirect calorimetry: A retrospective validation

Zusman, O. , Kagan, I. , Bendavid, I. (2019) *Clinical Nutrition*

Should we calculate or measure energy expenditure? Practical aspects in the ICU

Rattanachaiwong, S. , Singer, P. (2018) *Nutrition*

Evaluating the TARGET and EAT-ICU trials: How important are accurate caloric goals? Point-counterpoint: The pro position

Singer, P. , Pichard, C. , Rattanachaiwong, S. (2020) *Current Opinion in Clinical Nutrition and Metabolic Care*




View all related documents based on references

Find more related documents in Scopus based on:

Authors > Keywords >

References (47)

[View in search results format >](#)

All [Export](#)  [Print](#)  [E-mail](#)  [Save to PDF](#) [Create bibliography](#)

-
- 1 Singer, P., Anbar, R., Cohen, J., Shapiro, H., Shalita-Chesner, M., Lev, S., Grozovski, E., (...), Madar, Z.
The tight calorie control study (TICACOS): A prospective, randomized, controlled pilot study of nutritional support in critically ill patients

(2011) *Intensive Care Medicine*, 37 (4), pp. 601-609. Cited 326 times.
doi: 10.1007/s00134-011-2146-z

[View at Publisher](#)
-
- 2 McClave, S.A., Taylor, B.E., Martindale, R.G., Warren, M.M., Johnson, D.R., Braunschweig, C., McCarthy, M.S., (...), Compher, C.
Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.)

(2016) *Journal of Parenteral and Enteral Nutrition*, 40 (2), pp. 159-211. Cited 805 times.
doi: 10.1177/0148607115621863

[View at Publisher](#)
-
- 3 Singer, P., Blaser, A.R., Berger, M.M., Alhazzani, W., Calder, P.C., Casaer, M.P., Hiesmayr, M., (...), Bischoff, S.C.
ESPEN guideline on clinical nutrition in the intensive care unit ([Open Access](#))

(2019) *Clinical Nutrition*, 38 (1), pp. 48-79. Cited 168 times.
<http://www.elsevier-international.com/journals/clnu/>
doi: 10.1016/j.clnu.2018.08.037

[View at Publisher](#)
-
- 4 Heyland, D.K., Cahill, N., Day, A.G.
Optimal amount of calories for critically ill patients: Depends on how you slice the cake!

(2011) *Critical Care Medicine*, 39 (12), pp. 2619-2626. Cited 161 times.
doi: 10.1097/CCM.0b013e318226641d

[View at Publisher](#)
-
- 5 Singer, P., Singer, J.
Clinical guide for the use of metabolic carts: Indirect calorimetry - No longer the orphan of energy estimation

(2016) *Nutrition in Clinical Practice*, 31 (1), pp. 30-38. Cited 21 times.
<http://www.sagepub.com.ezproxy.um.edu.my/journalsProdDesc.nav?prodId=Journal201896>
doi: 10.1177/0884533615622536

[View at Publisher](#)
-