

A SINGULAR NONLINEAR SECOND-ORDER PERIODIC BOUNDARY VALUE PROBLEM

JUNYU WANG AND DAQING JIANG

(Received August 28, 1996, revised November 19, 1997)

Abstract. A singular nonlinear second-order periodic boundary value problem is studied and the uniqueness and existence of solutions are obtained by employing a priori estimates, perturbation techniques and comparison principles.

1. Introduction and main results. During the last two decades, singular nonlinear two-point boundary value problems, not including periodic ones, have been studied extensively. For details, see, for instance, papers [1]–[20] and the references therein. However, the works on singular nonlinear periodic boundary value problems are quite rarely seen. It is well known that periodic boundary value problems have always been attended to. So we study in this paper a singular nonlinear periodic boundary value problem of the form

$$(1) \quad \begin{cases} -u''(t) + \rho^2 u(t) = Q(t, u(t)), & 0 \leq t \leq 2\pi, \\ u(0) = u(2\pi), \quad u'(0) = u'(2\pi), \end{cases}$$

where ρ is a positive constant and the nonlinear function $Q(t, u) = f(t, u) + h(t, u)$ is assumed to be defined on $[0, 2\pi] \times (0, +\infty)$ and satisfy the following assumptions:

(A₁) For each fixed $u \in (0, +\infty)$, $f(t, u)$ is nonnegative integrable on $[0, 2\pi]$.

(A₂) For almost all $t \in [0, 2\pi]$, $f(t, u)$ is nonincreasing in $u > 0$ and

$$\int_0^{2\pi} f(s, 0+) ds > 0, \quad f(t, 0+) = \lim_{u \rightarrow 0+} f(t, u).$$

(A₃) $h(t, u)$ is a Caratheodory function defined on $[0, 2\pi] \times (0, \infty)$, i.e., for each fixed $u \in (0, \infty)$, the function $h(t, u)$ is nonnegative integrable on $[0, 2\pi]$ and for almost all $t \in [0, 2\pi]$, the function $h(t, u)$ is continuous in $u > 0$.

(A₄) There exists a nonnegative integrable function $k(t)$ defined on $[0, 2\pi]$ and a nonnegative continuous increasing function $H(u)$ defined on $(0, +\infty)$ such that

$$h(t, u) \leq k(t)H(u) \quad \text{for almost all } (t, u) \in [0, 2\pi] \times (0, \infty),$$

1991 *Mathematics Subject Classification.* Primary 34B15.

Key words and phrases: singular periodic boundary value problem, positive solution, uniqueness, existence, perturbation technique, comparison principle.

The work was partially supported by the NNSF of China.

where $H(u)$ and $k(t)$ respectively satisfy

$$(2) \quad \overline{\lim}_{u \rightarrow +\infty} \frac{H(u)}{u} = B$$

and

$$(3) \quad G(0)B \int_0^{2\pi} k(s)ds < 1.$$

Hence $G(0)$ is the positive maximum of the Green function

$$G(|t-s|) := G(t,s) := \begin{cases} \frac{e^{\rho(t-s)} + e^{\rho(2\pi-t+s)}}{2\rho(e^{2\rho\pi} - 1)}, & 0 \leq s \leq t \leq 2\pi, \\ \frac{e^{\rho(s-t)} + e^{\rho(2\pi-s+t)}}{2\rho(e^{2\rho\pi} - 1)}, & 0 \leq t \leq s \leq 2\pi. \end{cases}$$

A direct calculation shows that

$$\frac{e^{\rho\pi}}{\rho(e^{2\rho\pi} - 1)} = G(\pi) \leq G(t,s) \leq G(0) = \frac{e^{2\rho\pi} + 1}{2\rho(e^{2\rho\pi} - 1)} \quad \text{on } [0, 2\pi] \times [0, 2\pi].$$

The assumptions (A_1) and (A_2) allow but not require $f(t, u)$ to have singularity at $u=0$ and to be discontinuous with respect to u . For example, the function

$$f(t, u) = |\sin 8\pi t|^{-\alpha} \left(\sum_{j=1}^{\infty} \eta\left(\frac{1}{j} - u\right) + u^{-\beta} \right)$$

satisfies (A_1) and (A_2) . Here $\alpha \in (0, 1)$, $\beta > 0$, and $\eta(s)$ is the Heaviside function

$$\eta(s) := \begin{cases} 1, & s \geq 0, \\ 0, & s < 0. \end{cases}$$

A function $u(t)$ is said to be a solution to the problem (1), if

- (i) $u \in C^1[0, 2\pi]$, $u(0) = u(2\pi)$, $u'(0) = u'(2\pi)$,
- (ii) $u''(t)$ exists almost everywhere and integrable on $[0, 2\pi]$, and

$$-u''(t) + \rho^2 u(t) = Q(t, u(t)) \quad \text{a.e. on } [0, 2\pi].$$

Furthermore, if $u(t) > 0$ on $[0, 2\pi]$, then it is called a positive solution.

The main results of this paper are as follows.

THEOREM 1. *Let (A_1) – (A_4) hold. Then there exist two positive numbers δ and N such that for all solutions $u(t)$ to the problem (1)*

$$\delta \leq u(t) \leq N \quad \text{on } [0, 2\pi].$$

THEOREM 2. *Assume that (A_1) – (A_4) are satisfied and $f(t, u)$ is continuous in $u > 0$. Then the problem (1) has at least one positive solution.*

THEOREM 3. *Assume that $h(t, u) \equiv 0$ and $f(t, u)$ satisfies (A_1) – (A_2) . Then the problem (1) has a unique positive solution.*

Our arguments for establishing the existence and uniqueness of solutions to the problem (1) involve only the positivity of the Green function, a priori estimates, perturbation techniques, and comparison principles.

2. Proof of Theorem 1. It is readily verified that the problem (1) is equivalent to the integral equation

$$(4) \quad u(t) = \int_0^{2\pi} G(t, s)Q(s, u(s))ds, \quad 0 \leq t \leq 2\pi.$$

Let $u(t)$ be a solution to the problem (1) and let

$$m := \min\{u(t) : 0 \leq t \leq 2\pi\}, \quad M := \max\{u(t) : 0 \leq t \leq 2\pi\}.$$

Then, by (4), we obtain

$$G(\pi) \int_0^{2\pi} Q(s, u(s))ds \leq m \leq M \leq G(0) \int_0^{2\pi} Q(s, u(s))ds,$$

where $G(\pi)$ is the positive minimum of $G(t, s)$. Consequently, we have

$$(5) \quad M \leq mG(0)/G(\pi).$$

We now prove that there exists a $\delta > 0$ such that for all solutions $u(t)$ to the problem (1)

$$(6) \quad u(t) \geq \delta \quad \text{on} \quad [0, 2\pi].$$

If this is not the case, then there exists a sequence of solutions to the problem (1), $\{u_j(t)\}$, such that

$$m_j := \min\{u_j(t) : 0 \leq t \leq 2\pi\} \rightarrow 0 \quad \text{as} \quad j \rightarrow \infty.$$

From (A_1) , (A_2) , (A_3) , (5) and (6), we have

$$\begin{aligned} m_j &\geq G(\pi) \int_0^{2\pi} f(s, u(s))ds \\ &\geq G(\pi) \int_0^{2\pi} f(s, M_j)ds \\ &\geq G(\pi) \int_0^{2\pi} f(s, m_j G(0)/G(\pi))ds, \end{aligned}$$

where $M_j := \max\{u_j(t) : 0 \leq t \leq 2\pi\}$. Letting $j \rightarrow \infty$ in the above, we lead to

$$0 \geq G(\pi) \int_0^{2\pi} f(s, 0+) ds > 0,$$

a contradiction. This shows that (6) holds. Therefore, the condition (A_2) excludes the case when $u(t) \equiv 0$ is a solution to the problem (1).

Next, we prove that there exists an $N > 0$ such that for all solution $u(t)$ to the problem (1)

$$(7) \quad u(t) \leq N \quad \text{on} \quad [0, 2\pi].$$

From (3), we choose an $\varepsilon > 0$ such that

$$G(0)(B + \varepsilon) \int_0^{2\pi} k(s) ds < 1.$$

It follows from (2) that there exists an $N^* > \delta$ such that

$$H(u) \leq (B + \varepsilon)u \quad \text{for all} \quad u \geq N^*.$$

Let

$$N := \frac{N^* + G(0) \int_0^{2\pi} f(s, \delta) ds + G(0)H(N^*) \int_0^{2\pi} k(s) ds}{1 - G(0)(B + \varepsilon) \int_0^{2\pi} k(s) ds}$$

and let

$$M := \max\{u(t) : 0 \leq t \leq 2\pi\},$$

where $u(t)$ is an arbitrary solution to the problem (1). Without loss of generality, we may assume that $M \geq N^*$. From (4), we have

$$\begin{aligned} M &\leq G(0) \int_0^{2\pi} [f(s, u(s)) + h(s, u(s))] ds \\ &\leq G(0) \int_0^{2\pi} [f(s, \delta) + k(s)H(u(s))] ds \\ &< N^* + G(0) \int_0^{2\pi} f(s, \delta) ds + G(0) \int_0^{2\pi} k(s) ds H(N^*) \\ &\quad + G(0)(B + \varepsilon)M \int_0^{2\pi} k(s) ds \end{aligned}$$

i.e., $M \leq N$. This shows that (7) holds. Theorem 1 is thus proved.

3. Proof of Theorem 2. Let δ and N be two positive numbers determined by Theorem 1. Consider the modified problem

$$(8) \quad \begin{cases} -u''(t) + \rho^2 u(t) = Q^*(t, u(t)), & 0 \leq t \leq 2\pi, \\ u(0) = u(2\pi), \quad u'(0) = u'(2\pi), \end{cases}$$

where $Q^*(t, u) := f^*(t, u) + h^*(t, u)$ and

$$f^*(t, u) := \begin{cases} f(t, \delta) & \text{if } u < \delta, \\ f(t, u) & \text{if } u \geq \delta, \end{cases}$$

$$h^*(t, u) := \begin{cases} h(t, u) & \text{if } u \leq N, \\ h(t, N) & \text{if } u > N. \end{cases}$$

It is clear that both $f^*(t, u)$ and $h^*(t, u)$ satisfy (A_1) – (A_4) and hence Theorem 1 holds for the problem (8).

We define a mapping $\Phi: D \rightarrow D$ by

$$(\Phi u)(t) := \int_0^{2\pi} G(t, s) Q^*(s, u(s)) ds, \quad \forall u \in D,$$

where $D := \{u \in C[0, 2\pi]; u(t) \geq 0 \text{ on } [0, 2\pi]\}$.

By the definition of Φ , it is readily verified that Φ is a compactly continuous mapping from D into D . The Schauder fixed point theorem tells us that Φ has at least one fixed point in D . Let $u(t)$ be a fixed point in D . Then it is easy to check that $u(t)$ is a solution to the problem (8). Since $\delta \leq u(t) \leq N$ on $[0, 2\pi]$, $u(t)$ is also a positive solution to the problem (1).

4. Proof of Theorem 3. When $h(t, u) \equiv 0$, the following comparison principle holds.

LEMMA 1. *Let $f_j(t, u)$, $j=1, 2$, satisfy (A_1) and (A_2) and let $u_j(t)$, $j=1, 2$, be a solution to the problem (1) with $f = f_j$. If $f_1(t, u) \geq f_2(t, u)$ a.e. on $[0, 2\pi] \times (0, +\infty)$, then $u_1(t) \geq u_2(t)$ for all $t \in [0, 2\pi]$.*

PROOF OF LEMMA 1. Let $w(t) := u_1(t) - u_2(t)$. If the lemma is not true, then $w(t) < 0$ for some $t \in [0, 2\pi]$. Without loss of generality, we can consider only the following three cases.

Case I. $w(t) < 0$ on $[0, 2\pi]$. In this case, we have

$$(9) \quad -w''(t) + \rho^2 w(t) = f_1(t, u_1(t)) - f_2(t, u_2(t)) \geq 0$$

for almost all $t \in [0, 2\pi]$, i.e.,

$$w''(t) \leq \rho^2 w(t) < 0$$

for almost all $t \in [0, 2\pi]$, which implies that $w'(0) > w'(2\pi)$. This contradicts the fact that $w'(0) = w'(2\pi)$.

Case II. $w(0) = w(2\pi) \geq 0$ and $w(t) < 0$ for some $t \in (0, 2\pi)$. In this case, there exists a subinterval (a, b) of $[0, 2\pi]$ such that

$w(t) < 0$ in (a, b) , $w(a) = w(b) = 0$, and hence $w'(a) \leq 0 \leq w'(b)$.

By (9), we have

$$w''(t) \leq \rho^2 w(t) < 0 \quad \text{for almost all } t \in (a, b),$$

which implies that $w'(a) > w'(b)$. This is a contradiction.

Case III. $w(0) = w(2\pi) < 0$ and $w(t) \geq 0$ for some $t \in (0, 2\pi)$. In this case, there are two points $a, b \in (0, 2\pi)$, $a \leq b$, such that

$w(t) < 0$ in $[0, a) \cup (b, 2\pi]$, $w(a) = w(b) = 0$, and hence $w'(a) \geq 0 \geq w'(b)$.

By (9), we have

$$w''(t) \leq \rho^2 w(t) < 0 \quad \text{for almost all } t \in [0, a) \cup (b, 2\pi],$$

which implies that $w'(0) > w'(a) \geq w'(b) > w'(2\pi)$. This is also a contradiction.

The Lemma is thus proved.

In very much the same way, we can prove the following statement.

LEMMA 2. *If $h(t, u) \equiv 0$, then the problem (1) has at most one solution.*

When $f(t, u)$ is not continuous in $u > 0$ (for almost all $t \in [0, 2\pi]$), we employ a perturbation technique. We define

$$f(t, u, v) := \frac{1}{v} \int_u^{u+v} f^*(t, s) ds, \quad F(t, u, v) := \frac{1}{v} \int_{u-v}^u f^*(t, s) ds,$$

where $v > 0$, $f^*(t, u)$ is defined as in the proof of Theorem 2, and

$$f_j(t, u) := f\left(t, u, \frac{1}{j}\right), \quad F_j(t, u) := F\left(t, u, \frac{1}{j}\right), \quad j = 1, 2, \dots$$

Then $f_j(t, u)$ and $F_j(t, u)$ satisfy (A_1) , (A_2) ,

$$f_j(t, u) \leq f_{j+1}(t, u) \leq F_{j+1}(t, u) \leq F_j(t, u) \quad \text{on } [0, 2\pi] \times [0, +\infty),$$

and

$$\lim_{j \rightarrow \infty} f_j(t, u) = \lim_{j \rightarrow \infty} F_j(t, u) = f^*(t, u) \quad \text{pointwise on } [0, 2\pi] \times [0, +\infty),$$

since

$$\frac{\partial f(t, u, v)}{\partial u} \leq 0, \quad \frac{\partial F(t, u, v)}{\partial u} \geq 0, \quad \frac{\partial f(t, u, v)}{\partial v} \leq 0, \quad \frac{\partial F(t, u, v)}{\partial v} \geq 0,$$

$$\text{a.e. on } [0, 2\pi] \times [0, +\infty) \times (0, +\infty).$$

From Theorems 1 and 2, Lemmas 1 and 2, we conclude that the problem (1) with $f = f_j$ ($f = F_j$) has a unique solution $\delta \leq u_j(t) \leq N$ ($\delta \leq v_j(t) \leq N$) and

$$u_j(t) \leq u_{j+1}(t) \leq v_{j+1}(t) \leq v_j(t) \quad \text{on } [0, 2\pi].$$

Consequently, we have

$$(10) \quad u_j(t) \leq \lim_{j \rightarrow \infty} u_j(t) := u^*(t) \leq \lim_{j \rightarrow \infty} v_j(t) := v^*(t) \leq v_j(t) \quad \text{on } [0, 2\pi].$$

It follows by (4) that

$$\begin{aligned} u^*(t) \geq u_j(t) &= \int_0^{2\pi} G(t, s) f_j(s, u_j(s)) ds \\ &\geq \int_0^{2\pi} G(t, s) f_j(s, u^*(s)) ds \quad \text{on } [0, 2\pi]. \end{aligned}$$

Letting $j \rightarrow \infty$ in the above yields

$$u^*(t) \geq \int_0^{2\pi} G(t, s) f(s, u^*(s)) ds \quad \text{on } [0, 2\pi].$$

(Here we have used the fact that $\delta \leq u^*(t) \leq N$.) Similarly, we obtain

$$(11) \quad \begin{aligned} v^*(t) &\leq \int_0^{2\pi} G(t, s) f(s, v^*(s)) ds \\ &\leq \int_0^{2\pi} G(t, s) f(s, u^*(s)) ds \\ &\leq u^*(t) \quad \text{on } [0, 2\pi]. \end{aligned}$$

From (10) and (11), it follows that

$$u^*(t) = v^*(t) = \int_0^{2\pi} G(t, s) f(s, u^*(s)) ds \quad \text{on } [0, 2\pi].$$

This shows that $u^*(t)$ is a unique solution to problem (1).

This completes the proof of Theorem 3.

ACKNOWLEDGEMENT. The authors thank the referee for his valuable suggestion.

REFERENCES

- [1] A CALLEGARI AND A. NACHMAN, Some singular nonlinear differential equations arising in boundary layer theory, *J. Math. Anal. Appl.* 64 (1978), 96–105.
- [2] S. D. TALIAFERRO, A nonlinear singular boundary value problem, *Nonlinear Analysis* 3 (1979), 897–904.
- [3] A. NACHMAN AND A. CALLEGARI, A nonlinear boundary value problems in theory of pseudoplastic fluids, *SIAM J. Appl. Math.* 38 (1980), 275–281.
- [4] J. E. BOUILLET AND S. M. GOMES, An equation with a singular nonlinearity related to diffusion problems in one dimension, *Quart. Appl. Math.* 42 (1985), 395–402.
- [5] L. E. BOBISUD, D. O'REGAN AND W. D. ROYALTY, Singular boundary value problem, *App. Analysis* 23 (1986), 233–243.
- [6] JUNYU WANG, A two-point boundary value problem with singularity, *Northeast. Math. J.* 3 (1987), 281–291.

- [7] H. J. WEINISCHKE, On finite displacement of circular elastic membranes, *Math. Mech. in the Appl. Sci.* 9 (1987), 76–98.
- [8] C. J. VAN DUJIN, S. M. GOMES AND HONGFEI ZHANG, On a class of similarity solutions of the equation $u_t = (|u|^{m-1}u_x)_x$ with $m > -1$, *IMA J. Appl. Math.* 41 (1988), 147–163.
- [9] J. V. BAXLEY, A singular nonlinear boundary value problem: Membrane response of a spherical cap, *SIAM J. Appl. Math.* 48 (1988), 497–505.
- [10] J. A. GATICA, V. OLIKER AND P. WALTMAN, Singular nonlinear boundary value problems for second order differential equations, *J. Differential Equations* 79 (1989), 62–78.
- [11] D. O'REGAN, Positive solutions to singular and nonsingular second-order boundary value problems, *J. Math. Anal. Appl.* 142 (1989), 40–52.
- [12] JUNYU WANG, A free boundary problem for generalized diffusion equation, *Nonlinear Analysis* 14 (1990), 691–700.
- [13] J. A. CATICA, G. E. HERNANDEZ AND P. WALTMAN, Radially symmetric solutions of a class of singular elliptic equations, *Proc. Edinburgh Math. Soc.* 33 (1990), 168–180.
- [14] S. M. GOMES AND J. SPREKELS, Krasonskii's Theorem on operators compressing a cone: Application to some singular boundary value problems, *J. Math. Anal. Appl.* 153 (1990), 443–459.
- [15] JUNYU WANG AND JIE JIANG, The existence of positive solutions to a singular nonlinear boundary value problem, *J. Math. Anal. Appl.* 176 (1993), 322–329.
- [16] J. JANUS AND J. MYJAK, A generalized Emden-fowler equation with a negative exponent, *Nonlinear Analysis* 23 (1994), 953–970.
- [17] JUNYU WANG, On positive solutions of singular nonlinear two-point boundary problems, *J. Differential Equations* 107 (1994), 163–174.
- [18] JUNYU WANG, Solvability of singular nonlinear two-point boundary problems, *Nonlinear Analysis* 24 (1995), 555–561.
- [19] JUNYU WANG AND WEJIE GAO, A singular boundary value problem for the one-dimensional p -Laplacian, *J. Math. Anal. Appl.* 201 (1996), 851–866.
- [20] D. O'REGAN, Singular Dirichlet boundary value problem I. Superlinear and Nonresonant case, *Nonlinear Analysis* 29 (1997), 221–245.

DEPARTMENT OF MATHEMATICS
JILIN UNIVERSITY
CHANGCHUN 130023
PEOPLE'S REPUBLIC OF CHINA

DEPARTMENT OF MATHEMATICS
NORTHEAST NORMAL UNIVERSITY
CHANGCHUN 130024
PEOPLE'S REPUBLIC OF CHINA