

Swarthmore College

Works

Mathematics & Statistics Faculty Works

Mathematics & Statistics

2002

A Specht Module Analog For The Rook Monoid

Cheryl Grood

Swarthmore College, cgrood1@swarthmore.edu

Follow this and additional works at: <https://works.swarthmore.edu/fac-math-stat>



Part of the [Mathematics Commons](#)

[Let us know how access to these works benefits you](#)

Recommended Citation

Cheryl Grood. (2002). "A Specht Module Analog For The Rook Monoid". *Electronic Journal Of Combinatorics*. Volume 9, <https://works.swarthmore.edu/fac-math-stat/16>

This work is brought to you for free by Swarthmore College Libraries' Works. It has been accepted for inclusion in Mathematics & Statistics Faculty Works by an authorized administrator of Works. For more information, please contact myworks@swarthmore.edu.

A Specht Module Analog for the Rook Monoid

Cheryl Grood*

Department of Mathematics and Statistics
Swarthmore College, Swarthmore, PA, USA
cgrood1@swarthmore.edu

Submitted: July 20, 2001; Accepted: December 18, 2001.

MR Subject Classifications: 05E10, 20M30

Abstract

The wealth of beautiful combinatorics that arise in the representation theory of the symmetric group is well-known. In this paper, we analyze the representations of a related algebraic structure called the *rook monoid* from a combinatorial angle. In particular, we give a combinatorial construction of the irreducible representations of the rook monoid. Since the rook monoid contains the symmetric group, it is perhaps not surprising that the construction outlined in this paper is very similar to the classic combinatorial construction of the irreducible S_n -representations: namely, the Specht modules.

1 Introduction

Let R_n be the set of all $n \times n$ matrices that contain at most one entry of one in each column and row and zeroes elsewhere. Under matrix multiplication, R_n has the structure of a *monoid*, a set with an associative binary operation and an identity element. The monoid R_n is known both as the *symmetric inverse semigroup* and the *rook monoid*, the latter name stemming from the correspondence between the matrices in R_n and the placement of non-attacking rooks on an $n \times n$ chessboard. The number of rank r matrices in R_n is $\binom{n}{r}^2 r!$ and hence the rook monoid has a total of $\sum_{r=0}^n \binom{n}{r}^2 r!$ elements. Note that the set of rank n matrices in the rook monoid is isomorphic to S_n , the symmetric group on n letters.

It is often useful to associate each $n \times n$ matrix $(a_{ij}) \in R_n$ with a function σ , given by

$$\sigma(j) = \begin{cases} i & \text{if there exists an } i \text{ such that } a_{ij} = 1 \\ 0 & \text{otherwise.} \end{cases}$$

*The author wishes to thank Stephen Maurer for reading an earlier draft of this paper and giving many useful suggestions. The author is also grateful to Lou Solomon for encouraging her to write up these results and to the referee for several helpful comments.

This function is well-defined because at most one element in each column is nonzero. For example, with $n = 4$ the matrix

$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

corresponds to a map σ that sends 1 to 3, 2 to 0, 3 to 2, and 4 to 4. Note that when (a_{ij}) is a permutation matrix, the function σ is in fact the permutation associated with that matrix. Munn [2] introduced a concise way of representing elements of R_n using what he termed *cycle-link notation*. In this notation, a cycle (a_1, a_2, \dots, a_k) means, as in the case of permutations, that

$$a_1 \mapsto a_2 \mapsto \dots \mapsto a_k \mapsto a_1.$$

In contrast, a link $[b_1, b_2, \dots, b_l]$ means that $b_1 \mapsto b_2 \mapsto \dots \mapsto b_l$ and b_l maps to 0. The 4×4 matrix above would translate to $[1, 3, 2](4)$ in cycle-link notation.


Munn's results ([2], [3]) on the representations of rook monoids stemmed from his work on the more general theory of representations of finite semigroups. The representation theory of the rook monoid is particularly interesting because of its similarity to the representation theory of the symmetric group. Munn showed that the complex monoid algebra $\mathbb{C}R_n$ is semisimple and subsequently found the irreducible representations of R_n . He demonstrated that these irreducibles are indexed by partitions of nonnegative integers less than or equal to n , whereas the irreducible representations of S_n are indexed by partitions of n . In addition, Munn computed the characters of the R_n -irreducibles using irreducible characters of S_r , $1 \leq r \leq n$.

More recently, Solomon ([5], [6]) has investigated q -generalizations of the rook monoid as well as an analog of Schur-Weyl duality for R_n , and Halverson et al.[1] have found an R_n analog of the Murnaghan-Nakayama rule, a combinatorial construction of the irreducible characters of S_n . In this paper, we provide a combinatorial construction for the irreducible representations of R_n that is very similar to the Specht module construction of the irreducible representations of S_n . Our construction and notation closely follow the exposition in [4] on Specht modules.

2 Preliminaries

A *partition* is a finite sequence of nonnegative integers $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_m)$ arranged in a weakly decreasing order; that is, $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m \geq 0$. The λ_i are called the *parts* of the partition. The *weight* of λ , denoted $|\lambda|$, is the sum of its parts: $|\lambda| = \sum_{i=1}^m \lambda_i$. If $|\lambda| = r$, we write: $\lambda \vdash r$. The partition with all parts equal to 0 is the *empty partition*.

The *length* of λ , denoted $\ell(\lambda)$, is the maximum subscript j such that $\lambda_j > 0$. Associated to each partition λ is a Ferrers diagram of shape λ . The Ferrers diagram has λ_i boxes in its i -th row, and the boxes are left-justified. For example, the Ferrers diagram associated with $\lambda = (6, 4, 3, 2, 2, 1)$ is



The Ferrers diagram for the empty partition is denoted by \emptyset . If $\lambda \vdash r$, we say that a *tableau of shape λ* , or a λ -*tableau*, is a Ferrers diagram of shape λ with the boxes filled with distinct entries from the set $\{1, 2, \dots, r\}$. The following is an example of a tableau of shape $(4, 2, 1, 1)$.

1	3	4	8
2	6		
5			
7			

Tableaux are at the core of the Specht module construction for the irreducibles of S_n . We now introduce a new object which will serve the same function for R_n as the tableaux did for S_n .

Definition 2.1 *Let $\lambda \vdash r$, where $0 \leq r \leq n$. An n -tableau of shape λ , also called a λ_r^n -tableau, is a Ferrers diagram of shape λ filled with r distinct entries from the set $\{1, 2, \dots, n\}$.*

For instance,

1	5
3	7
4	

is an n -tableau of shape $(2, 2, 1)$ for any $n \geq 7$. Note that in the case where $r = n$, an n -tableau of shape λ is in fact a tableau of shape λ . The *content* of an λ_r^n -tableau t is the set of entries in t . An n -tableau t of shape λ is *standard* if the entries in its rows and columns increase left-to-right and top-to-bottom, respectively.

Notation 2.1 *Let t be a λ_r^n -tableau. Then $t_{i,j}$ will denote the entry contained in the box of the i th row and j th column of t .*

Two n -tableaux t_1 and t_2 of shape λ are *row-equivalent* if the corresponding rows of the two tableaux contain the same entries; in this case, we write $t_1 \sim t_2$.

Definition 2.2 An n -tabloid $\{t\}$ of shape λ , or an λ_r^n -tabloid $\{t\}$, is the set of all λ_r^n -tableaux that are row-equivalent to t ; i.e.,

$$\{t\} = \{s \mid s \sim t\}.$$

Let N^λ be the vector space over \mathbb{C} generated by the λ_r^n -tableaux; that is, N^λ is the set of all formal \mathbb{C} -linear combinations of λ_r^n -tableaux. We can define an action of R_n on N^λ by first determining how R_n acts on the basis of λ_r^n -tableaux and then linearly extending this action to the whole vector space. If t is an λ_r^n -tableau, we define σt to equal the zero vector if t contains an entry $t_{i,j}$ such that $\sigma(t_{i,j}) = 0$; when t contains no such entry, we say that $(\sigma t)_{i,j} = \sigma(t_{i,j})$. Similarly, if we let M^λ represent the vector space over \mathbb{C} generated by the λ_r^n -tabloids, we have an induced action of R_n on M^λ given by:

$$\sigma\{t\} = \begin{cases} \mathbf{0} & \text{if } \sigma t = \mathbf{0} \\ \{\sigma t\} & \text{otherwise.} \end{cases}$$

Since $s \sim t$ implies that $\sigma s \sim \sigma t$, this induced action is well-defined.

Suppose that t is an λ_r^n -tableau, and let C_i be the entries in the i th column of t . We now form the group

$$C_t = S_{C_1} \times S_{C_2} \times \cdots \times S_{C_i},$$

which we can think of as being contained in R_n . Each element in C_t stabilizes the columns of t , but note that C_t does not consist of *all* the elements in R_n that fix the columns of t . For each λ_r^n -tableau t , we will generate the following element of M^λ :

$$e_t = \sum_{\sigma \in C_t} \text{sgn}(\sigma) \sigma\{t\},$$

where *sgn* stands for the sign of the permutation σ . We will call e_t the *n -polytabloid associated with t* . Let R^λ be the vector space over \mathbb{C} generated by the set of n -polytabloids of shape λ ; let S^λ be the subspace of R^λ generated by $\{e_t \mid t, \lambda_r^n\text{-tableau; content } t = \{1, 2, \dots, r\}\}$. The subspaces S^λ , $\lambda \vdash r$, are exactly the so-called *Specht modules*, a complete set of distinct irreducible S_r -modules. We will show that the vector spaces R^λ form a complete set of distinct irreducible R_n -modules, and hence are analogous to the Specht modules for the symmetric group.

3 R_n -modules

The goal of this section is to show that for each λ such that $|\lambda| \leq n$, R^λ is an R_n -module. For that, we will need some new notation and several lemmas.

Definition 3.1 Let $\pi \in R_n$. Then $\hat{\pi}$ will denote the element in S_n that comes from changing all the links in π to cycles.

For example, if $\pi = [1, 4, 6](2, 3)[5]$, then $\widehat{\pi} = (1, 4, 6)(2, 3)(5)$. If $\pi \in S_n$, then clearly $\pi = \widehat{\pi}$.

Lemma 3.1 *Suppose $\pi \in R_n$ and t is a λ_r^n -tableau. If $\pi t \neq \mathbf{0}$, then $\pi t = \widehat{\pi} t$.*

Proof. Express π in cycle-link notation. Since $\pi t \neq \mathbf{0}$, none of the entries $t_{i,j}$ occur as a right-most element of a link of π . Thus $\pi(t_{i,j}) = \widehat{\pi}(t_{i,j})$ for all i, j and so $\pi t = \widehat{\pi} t$. \square

Lemma 3.2 *If $\sigma \in S_n$ and t is an λ_r^n -tableau with column entries C_1, \dots, C_l , then $C_{\sigma t} = \sigma C_t \sigma^{-1}$.*

Proof. Let $\pi = (i_1, i_2, \dots, i_j)(i_{j+1}, \dots, i_k) \cdots (i_m, \dots, i_n)$ be an element in S_n expressed in cycle notation. Now consider $\sigma C_t \sigma^{-1}$. If we view C_t as a subgroup of S_n , it follows that $\sigma C_t \sigma^{-1} = \sigma S_{C_1} \sigma^{-1} \times \cdots \times \sigma S_{C_l} \sigma^{-1}$. Because

$$\sigma \pi \sigma^{-1} = (\sigma(i_1), \sigma(i_2), \dots, \sigma(i_j))(\sigma(i_{j+1}), \dots, \sigma(i_k)) \cdots (\sigma(i_m), \dots, \sigma(i_n)),$$

it follows that $\sigma S_{C_i} \sigma^{-1} = S_{\sigma(C_i)}$, where $\sigma(C_i)$ represents the set of elements $\{\sigma(j) \mid j \in C_i\}$. So,

$$\sigma C_t \sigma^{-1} = S_{\sigma(C_1)} \times \cdots \times S_{\sigma(C_l)},$$

which is exactly $C_{\sigma t}$. \square

These lemmas allow us to prove the following proposition.

Proposition 3.3 *Suppose $\pi \in R_n$, t is a λ_r^n -tableau. If $\pi t = \mathbf{0}$, then $\pi e_t = \mathbf{0}$. Otherwise, $\pi e_t = e_{\widehat{\pi} t}$.*

Proof. First consider the case where $\pi t = \mathbf{0}$. Recall that $\pi t = \mathbf{0}$ if and only if $\pi\{t\} = \mathbf{0}$ as well. Each term in the linear combination of tabloids that form e_t contains the exact same entries as t . Therefore, if $\pi\{t\} = \mathbf{0}$, then $\pi\{s\} = \mathbf{0}$ for all $\{s\}$ in the linear combination e_t ; it follows that in this case, $\pi e_t = \mathbf{0}$.

If $\pi t \neq \mathbf{0}$, we have $\pi t = \widehat{\pi} t$ by Lemma 3.1. Thus, $\pi\{s\} = \widehat{\pi}\{s\}$ for every tabloid $\{s\}$ that appears in the linear combination e_t , and hence $\pi e_t = \widehat{\pi} e_t$. By manipulating the terms of $\widehat{\pi} e_t$, we can conclude that

$$\begin{aligned} \pi e_t &= \widehat{\pi} e_t \\ &= \widehat{\pi} \sum_{\sigma \in C_t} \text{sgn}(\sigma) \sigma \{t\} \\ &= \sum_{\sigma \in C_t} \text{sgn}(\sigma) \widehat{\pi} \sigma \{t\} \\ &= \sum_{\sigma \in C_t} \text{sgn}(\widehat{\pi} \sigma \widehat{\pi}^{-1}) \widehat{\pi} \sigma \widehat{\pi}^{-1} \widehat{\pi} \{t\} \\ &= \sum_{\sigma \in C_t} \text{sgn}(\widehat{\pi} \sigma \widehat{\pi}^{-1}) \widehat{\pi} \sigma \widehat{\pi}^{-1} \{\widehat{\pi} t\} \end{aligned}$$

$$\begin{aligned}
&= \sum_{\gamma \in \widehat{\pi} C_t \widehat{\pi}^{-1}} \text{sgn}(\gamma) \gamma \{\widehat{\pi} t\} \\
&= \sum_{\gamma \in C_{\widehat{\pi} t}} \text{sgn}(\gamma) \gamma \{\widehat{\pi} t\} \quad (\text{by Lemma 3.2}) \\
&= e_{\widehat{\pi} t}.
\end{aligned}$$

□

Thus, by Proposition 3.3, it follows that for any $\pi \in R_n$ and e_t in R^λ , πe_t equals either $\mathbf{0}$ or $e_{\widehat{\pi} t}$, both of which are again elements of R^λ . This immediately gives us the following corollary.

Corollary 3.4 R^λ is an R_n -module.

Furthermore, we are able to show easily that

Corollary 3.5 R^λ is a cyclic R_n -module.

Proof. Note that if s and t are both λ_r^n -tableaux, there exists an element σ in S_n such that $s = \sigma t$. By Proposition 3.3, we have that $\sigma e_t = e_{\widehat{\sigma} t}$. Since $\sigma \in S_n, \sigma = \widehat{\sigma}$, and so $\sigma e_t = e_{\sigma t}$, which in turn equals e_s . Since we can generate all the basis elements of R^λ using one element e_t , R^λ is cyclic. □

4 Irreducibility of R^λ

Let $|\lambda| = r$, with $r \leq n$. Recall that the subspace S^λ of R^λ is an irreducible S_r -module. We will use this fact to help show that R^λ is an irreducible R_n -module.

Theorem 4.1 If $|\lambda| \leq n$, R^λ is an irreducible R_n -module.

Proof. Let U be a nonzero submodule of R^λ . We need to show that U must in fact be equal to R^λ . Since R^λ is a cyclic module, it suffices to show that we can find a λ_r^n -tableau t such that $e_t \in U$.

Let \mathbf{u} be a nonzero element of U , and write \mathbf{u} as a linear combination of n -polytabloids:

$$\mathbf{u} = \sum_{t, \lambda_r^n\text{-tableau}} a_t e_t.$$

We now group the tabloids based on their content; *i.e.*,

$$\mathbf{u} = \sum_{\substack{X \subset \{1, \dots, n\} \\ |X| = r}} \left(\sum_{t, \text{tableau of content } X} a_t e_t \right).$$

Since $\mathbf{u} \neq \mathbf{0}$, there must be at least one set X_0 such that

$$\sum_{t, \text{ content } t = X_0} a_t e_t \neq \mathbf{0}.$$

If $X_0 = \{i_1, i_2, \dots, i_r\}$, where $i_1 < i_2 < \dots < i_r$, consider the element π in R_n defined by

$$\pi(k) = \begin{cases} 0 & \text{if } k \notin X_0 \\ j & \text{if } k = i_j. \end{cases}$$

Note that we have defined π in such a way so that $\pi t = \mathbf{0}$ if and only if content $t \neq X_0$. Multiplying \mathbf{u} by π , we obtain

$$\pi \mathbf{u} = \sum_{t, \lambda_r^n\text{-tableau}} a_t (\pi e_t).$$

By Proposition 3.3, if $\pi t = \mathbf{0}$, then $\pi e_t = \mathbf{0}$. We have defined π in such a way that all the terms in $\pi \mathbf{u}$ with content not equal to X_0 vanish. Thus, $\pi \mathbf{u}$ simplifies to

$$\pi \mathbf{u} = \sum_{t, \text{ content } t = X_0} a_t (\pi e_t),$$

which in turn equals $\sum_{t, \text{ content } t = X_0} a_t (\widehat{\pi} e_t)$, since both π and $\widehat{\pi}$ act the same on tabloids of content X_0 . Multiplying on the left by $\widehat{\pi}^{-1}$, we see that

$$\begin{aligned} \mathbf{u} &= \widehat{\pi}^{-1}(\pi \mathbf{u}) \\ &= \sum_{t, \text{ content } t = X_0} a_t \widehat{\pi}^{-1}(\widehat{\pi} e_t) \\ &= \sum_{t, \text{ content } t = X_0} a_t e_t, \end{aligned} \tag{1}$$

which is nonzero, so $\pi \mathbf{u}$ cannot be $\mathbf{0}$.

Now we apply Proposition 3.3 to (1), so that

$$\pi \mathbf{u} = \sum_{t, \text{ content } t = X_0} a_t e_{\widehat{\pi} t}.$$

Now observe that when the content of t is X_0 , the content of $\widehat{\pi} t$ equals $\{1, 2, \dots, r\}$, which in turn implies that $e_{\widehat{\pi} t} \in S^\lambda$. Since $\pi \mathbf{u}$ is a linear combination of elements in S^λ , $\pi \mathbf{u}$ itself is an element of S^λ . Hence $U \cap S^\lambda \neq \mathbf{0}$, since the intersection contains the nonzero element $\pi \mathbf{u}$. Viewing U as an S_r -module (by realizing $S_r \subset R_n$), we have that $U \cap S^\lambda$ is a nonzero S_r -submodule of S^λ . By the irreducibility of S^λ , we can conclude that $U \cap S^\lambda = S^\lambda$ and thus $S^\lambda \subseteq U$. Now pick any $e_t \in S^\lambda$; e_t must also be contained in U , and since R^λ is cyclic, we have that $U = R^\lambda$ as desired. \square

Now that we know that these modules are irreducible, we need to show that they are all distinct.

Theorem 4.2 *If R^λ and R^μ are isomorphic as R_n -modules, then $\lambda = \mu$.*

Proof. Let $\Theta : R^\lambda \longrightarrow R^\mu$ be an R_n -isomorphism. Let t be a λ_r^n -tableau. The image of e_t under Θ is a linear combination of $\{e_s \mid s, \text{ standard } \mu_r^n\text{-tableaux}\}$. If the entry i is in t , then $\pi = (1)(2) \cdots (\widehat{i}) \cdots (n)[i]$ annihilates t , and hence e_t as well. Thus, $\Theta(\pi e_t) = \mathbf{0}$, which implies that

$$\begin{aligned} \mathbf{0} &= \pi\Theta(e_t) \\ &= \pi \left(\sum_{s, \text{ standard } \mu_r^n\text{-tableau}} a_s e_s \right) \\ &= \sum_{s, \text{ standard } \mu_r^n\text{-tableau}} a_s \pi e_s \\ &= \sum_{\substack{s, \text{ standard } \mu_r^n\text{-tableau} \\ \pi s \neq \mathbf{0}}} a_s e_{\widehat{\pi}s} && \text{(by Proposition 3.3)} \\ &= \sum_{\substack{s, \text{ standard } \mu_r^n\text{-tableau} \\ \pi s \neq \mathbf{0}}} a_s e_s, \end{aligned}$$

since $\widehat{\pi}$ is the identity element. The only linear combination of basis elements that equals $\mathbf{0}$ is the trivial one; thus, $\Theta(e_t)$ is a linear combination of $\{e_s \mid s, \text{ standard } \mu_r^n\text{-tableau, } \pi s = \mathbf{0}\}$. However, the only tableaux that π annihilates are those that contain an entry of i . Since this argument holds for every i in the content of t , $\Theta(e_t)$ is a linear combination of μ_r^n -polytabloids e_s such that the content of t is a subset of the content of s . Since Θ is invertible, we in fact have that

$$\Theta(e_t) = \sum_{\substack{s, \text{ standard } \mu_r^n\text{-tableau} \\ \text{content } s = \text{content } t}} a_s e_s. \tag{2}$$

Recall that S^λ is the subspace of R^λ generated by the polytabloids with content $1, 2, \dots, |\lambda|$. It follows from Equation 2 that the image of S^λ under Θ is contained in S^μ ; that is, S^λ is isomorphic to a submodule of S^μ . The only way this can occur is if $\lambda = \mu$, which is what we wanted to show. \square

5 A Basis for R^λ

In closing, we give a combinatorial basis for our irreducible R_n -modules R^λ . Again, the story closely resembles what happens in the case of the irreducible S_r -modules S^λ . In that case, a basis for S^λ consists of $\{e_t \mid t \text{ is a standard tableau of shape } \lambda\}$. We will establish that a similar statement holds for R^λ :

Theorem 5.1 *Let $\lambda \vdash r$, with $0 \leq r \leq n$. Then the set*

$$\{e_t \mid t \text{ is a standard } \lambda_r^n\text{-tableau}\}$$

is a basis for R^λ .

Proof. First, we will show that the e_t are linearly independent. If the content of two λ_r^n -tableaux s and t differ, it is not hard to see that e_t and e_s are independent. Therefore, it suffices to show that

$$\{e_t \mid t, \text{ standard } \lambda_r^n\text{-tableau with content } X\}$$

is independent. Without loss of generality, assume that the content X is the set $\{1, 2, \dots, r\}$. Since these polytabloids form a basis for S^λ , they are independent, which is what we wanted to show.

Next, we compute the order of $\{e_t \mid t \text{ is a standard } \lambda_r^n\text{-tableau}\}$. There are $\binom{n}{r}$ ways to choose a fixed content X . Let f_λ denote the number of standard tableaux of shape λ . Then in fact f_λ counts the number of λ_r^n -tableaux with a given content X . Hence, the set $\{e_t \mid t \text{ is a standard } \lambda_r^n\text{-tableau}\}$ has $\binom{n}{r} f_\lambda$ elements. Since we have shown that these vectors are independent, we can conclude that the dimension of R^λ is at least $\binom{n}{r} f_\lambda$.

For semisimple algebras, the sum of the squares of the dimensions of its nonisomorphic irreducible modules equals the dimension of the algebra itself; *i.e.*,

$$\sum_{I, \text{ irreducible } A\text{-module}} (\dim I)^2 = \dim A.$$

Suppose there is some λ such that the dimension of R^λ is strictly larger than $\binom{n}{r} f_\lambda$. Then we would have

$$\begin{aligned} \sum_{r=0}^n \binom{n}{r}^2 r! &= \dim \mathbb{C}R_n \\ &< \sum_{r=0}^n \sum_{\lambda \vdash r} (\dim R^\lambda)^2 \\ &= \sum_{r=0}^n \sum_{\lambda \vdash r} \binom{n}{r}^2 (f_\lambda)^2 \\ &= \sum_{r=0}^n \binom{n}{r}^2 \left(\sum_{\lambda \vdash r} (f_\lambda)^2 \right). \end{aligned}$$

But, it is a well-known combinatorial identity that

$$\sum_{\lambda \vdash r} (f_\lambda)^2 = r!$$

(*cf.* [4]), and so every irreducible module R^λ must have dimension equal to $\binom{n}{r} f_\lambda$. Hence, the set of independent vectors $\{e_t \mid t \text{ is a standard } \lambda_r^n\text{-tableau}\}$ also spans R^λ . Thus we have shown that $\{e_t \mid t \text{ is a standard } \lambda_r^n\text{-tableau}\}$ is a basis for R^λ . \square

References

- [1] M. Dieng, T. Halverson, and V. Poladian, *Characters of the q -rook monoid*, preprint, 2001.
- [2] W.D. Munn, *Matrix representations of semigroups*, Proc. Camb. Phil. Soc., **53** (1957), 5-12.
- [3] W.D. Munn, *The characters of the symmetric inverse semigroup*, Proc. Camb. Phil. Soc., **53** (1957), 13-18.
- [4] B. Sagan, *The Symmetric Group*, Wadsworth and Brooks Cole, Pacific Grove, CA, 1991.
- [5] L. Solomon, *Representations of the Rook Monoid*, to appear in J. Algebra.
- [6] L. Solomon, *Representations of the Symmetric Inverse Semigroup*, Address given at C.R.M. Montréal, 1997.