

# A spectral sequence for the homology of a finite algebraic delooping

Birgit Richter

joint work in progress with Stephanie Ziegenhagen

Fourth Arolla Conference  
on Algebraic Topology  
2012

$E_n$ -homology

A resolution spectral sequence


A Blanc-Stover spectral sequence

Examples

## Little $n$ -cubes

Let  $C_n$  denote the operad of little  $n$ -cubes.  $C_n(r)$ ,  $r \geq 0$ .

$n = 2, r = 3$ :



$C_n$  acts on and detects  $n$ -fold based loop spaces.

## $E_n$ -homology

$(C_* C_n(r))_r, r \geq 1$  is an operad in the category of chain complexes. Let  $E_n$  be a cofibrant replacement of  $C_* C_n$ . For an augmented  $E_n$ -algebra  $A_*$  let  $\bar{A}_*$  denote the augmentation ideal.

The sth  $E_n$ -homology group of  $\bar{A}_*$ ,  $H_s^{E_n}(\bar{A}_*)$  is then the sth derived functor of indecomposables of  $\bar{A}_*$ .

**Theorem [Fresse 2011]**

There is an  $n$ -fold bar construction for  $E_n$ -algebras,  $B^n$ , such that

$$H_s^{E_n}(\bar{A}_*) \cong H_s(\Sigma^{-n} B^n(\bar{A}_*)).$$

I.e.,  $E_n$ -homology is the homology of an  $n$ -fold algebraic delooping.

## Some results

Cartan (50s):  $H_*^{E_n}$  of polynomial algebras, exterior algebras and some more.

Fresse (2011):  $X$  a nice space:  $B^n(C^*(X))$  determines the cohomology of  $\Omega^n X$ .

Livernet-Richter (2011): Functor homology interpretation for  $H_*^{E_n}$  for augmented commutative algebras.

$H_*^{E_n}(\bar{A}_*) \cong HH_{*+n}^{[n]}(A_*)$ , Hochschild homology of order  $n$  in the sense of Pirashvili.

What is  $H_*^{E_n}(\bar{A}_*)$  in other interesting cases such as Hochschild cochains,  $A_* = C^*(B, B)$ , or  $A_* = C_*(\Omega^n X)$ ?

## Setting

In the following  $k$  is a field, most of the times  $k = \mathbb{F}_2$  or  $k = \mathbb{Q}$ .  
The underlying chain complex of  $A_*$  is non-negatively or non-positively graded.  
Over  $\mathbb{F}_2$ :  $n = 2$ ; for  $\mathbb{Q}$ : arbitrary  $n$ .

# 1-restricted Lie algebras

## Definition

A 1-restricted Lie algebra over  $\mathbb{F}_2$  is a non-negatively graded  $\mathbb{F}_2$ -vector space,  $\mathfrak{g}_*$ , together with two operations, a Lie bracket of degree one,  $[-, -]$  and a restriction,  $\xi$ :

$$\begin{aligned}[-, -]: \quad \mathfrak{g}_i \times \mathfrak{g}_j &\rightarrow \mathfrak{g}_{i+j+1}, & i, j \geq 0, \\ \xi: \quad \mathfrak{g}_i &\rightarrow \mathfrak{g}_{2i+1} & i \geq 0.\end{aligned}$$

These satisfy the relations

1. The bracket is bilinear, symmetric and satisfies the Jacobi relation

$$[a, [b, c]] + [b, [c, a]] + [c, [a, b]] = 0 \text{ for all homogeneous } a, b, c \in \mathfrak{g}_*.$$

2. The restriction interacts with the bracket as follows:  
 $[\xi(a), b] = [a, [a, b]]$  and  $\xi(a + b) = \xi(a) + \xi(b) + [a, b]$  for all homogeneous  $a, b \in \mathfrak{g}_*$ .

1-rL: The category of 1-restricted Lie algebras.

# 1-restricted Gerstenhaber algebras

## Definition

A 1-restricted Gerstenhaber algebra over  $\mathbb{F}_2$  is a 1-restricted Lie algebra  $G_*$  together with an augmented commutative  $\mathbb{F}_2$ -algebra structure on  $G_*$  such that the multiplication in  $G_*$  interacts with the restricted Lie-structure as follows:

- ▶ (Poisson relation)

$$[a, bc] = b[a, c] + [a, b]c, \text{ for all homogeneous } a, b, c \in G_*.$$

- ▶ (multiplicativity of the restriction)

$$\xi(ab) = a^2\xi(b) + \xi(a)b^2 + ab[a, b] \text{ for all homogeneous } a, b \in G_*.$$

1-rG: the category of 1-restricted Gerstenhaber algebras.

In particular, the bracket and the restriction annihilate squares:

$[a, b^2] = 2b[a, b] = 0$  and  $\xi(a^2) = 2a^2\xi(a) + a^2[a, a] = 0$ . Thus if 1 denotes the unit of the algebra structure in  $G_*$ , then  $[a, 1] = 0$  for all  $a$  and  $\xi(1) = 0$ .



## Free objects and indecomposables

For a graded vector space  $V_*$  let  $1rL(V_*)$  be the free 1-restricted Lie algebra on  $V_*$ .

The free commutative algebra  $S(1rL(V_*))$  has a well-defined 1-rG structure and is the free 1-restricted Gerstenhaber algebra generated by  $V_*$ :

$$1rG(V_*) = S(1rL(V_*)).$$

For  $G_* \in 1rG$  let  $Q_{1rG}(G_*)$  be the graded vector space of indecomposables.

Note:  $Q_{1rG}(G_*) = Q_{1rL}(Q_a(G_*))$ .

# Homology of free objects

## Lemma

$$H_*(E_2(\bar{A}_*)) \cong 1rG(H_*(\bar{A}_*)).$$

Proof: Let  $X$  be a space. F. Cohen describes  $H_*(C_2(X); \mathbb{F}_2)$ .

Observation by Haynes Miller:  $H_*(C_2(X); \mathbb{F}_2) \cong 1rG(\bar{H}_*(X; \mathbb{F}_2))$ .

(Dyer-Lashof operations only give algebraic operations. )

Take  $X$  with  $\bar{H}_*(X; \mathbb{F}_2) \cong H_*(\bar{A}_*)$ , then

$$H_*(E_2(\bar{A}_*)) \cong H_*(C_2(X); \mathbb{F}_2).$$

# Resolution spectral sequence

## Theorem

There is a spectral sequence

$$E_{p,q}^2 \cong (\mathbb{L}_p Q_{1rG}(H_*(\bar{A}_*)))_q \Rightarrow H_{p+q}^{E_2}(\bar{A}_*)$$

Proof: Standard resolution  $E_2^{\bullet+1}(\bar{A}_*)$ .

$$E_{p,q}^1 : H_q^{E_2}(E_2^{p+1}(\bar{A}_*)) \cong H_q(E_2^p(\bar{A}_*))$$

$$H_q(E_2^p(\bar{A}_*)) \cong 1rG^p(H_*\bar{A}_*)_q \cong Q_{1rG}(1rG^{p+1}(H_*\bar{A}_*))_q.$$

$d^1$  takes homology wrt resolution degree.

## Example

For  $X$  connected:

$$(\mathbb{L}_p Q_{1rG}(H_*(C_*(\Omega^2 \Sigma^2 X; \mathbb{F}_2))))_* = (\mathbb{L}_p Q_{1rG}(1rG(\bar{H}_*(X; \mathbb{F}_2))))_*.$$

This reduces to  $\bar{H}_q(X; \mathbb{F}_2)$  in the  $(p = 0)$ -line and

$$H_q^{E_2}(C_*(\Omega^2 \Sigma^2 X; \mathbb{F}_2)) \cong \bar{H}_q(X; \mathbb{F}_2).$$

## Rational case

The rational case is much easier:

$$H_*(E_{n+1}\bar{A}_*) \cong nG(H_*(\bar{A}_*)),$$

the free  $n$ -Gerstenhaber algebra generated by the homology of  $\bar{A}_*$ .

We get:

$$E_{p,q}^2 \cong (\mathbb{L}_p Q_{nG}(H_*(\bar{A}_*)))_q \Rightarrow H_{p+q}^{E_{n+1}}(\bar{A}_*)$$

for every  $E_{n+1}$ -algebra  $\bar{A}_*$  over the rationals.

## General Blanc-Stover setting

Let  $\mathcal{C}$  and  $\mathcal{B}$  be some categories of graded algebras (e.g., Lie, Com,  $n$ -Gerstenhaber etc.) and let  $\mathcal{A}$  be a concrete category (such as graded vector spaces) and  $T: \mathcal{C} \rightarrow \mathcal{B}$ ,  $S: \mathcal{B} \rightarrow \mathcal{A}$ .

If  $TF$  is  $S$ -acyclic for every free  $F$  in  $\mathcal{C}$ , then there is a Grothendieck composite functor spectral sequence for all  $C$  in  $\mathcal{C}$

$$E_{s,t}^2 = (\mathbb{L}_s \bar{S}_t)(\mathbb{L}_* T)C \Rightarrow (\mathbb{L}_{s+t}(S \circ T))C.$$

- ▶ Note:  $T, S$  non-additive.
- ▶  $\bar{S}_t(\pi_* B) = \pi_t(SB)$  if  $B$  is free simplicial; otherwise it is defined as a coequaliser.
- ▶  $\bar{S}$  takes the homotopy operations on  $\pi_* B$  into account ( $B$  a simplicial object in  $\mathcal{B}$ ):  $\pi_* B$  is a  $\Pi$ - $\mathcal{B}$ -algebra.
- ▶  $\mathcal{B} = Com$ :  $\pi_*(B)$  has divided power operations.  $\mathcal{B} = rLie$ :  $\pi_* B$  inherits a Lie bracket and has some extra operations.

## In our case

### Theorem

- ▶  $k = \mathbb{F}_2$ : For any  $C \in 1rG$ :

$$E_{s,t}^2 = \mathbb{L}_s((\bar{Q}_{1rL})_t)(AQ_*(C|\mathbb{F}_2, \mathbb{F}_2)) \Rightarrow \mathbb{L}_{s+t}(Q_{1rG}).$$

- ▶ For  $k = \mathbb{Q}$  we get for all  $n$ -Gerstenhaber algebras  $C$ :

$$\mathbb{L}_s((\bar{Q}_{nL})_t)(AQ_*(C|\mathbb{Q}, \mathbb{Q})) \Rightarrow \mathbb{L}_{s+t}(Q_{nG}).$$

## Hochschild cochains, rational case

Let  $V$  be a vector space. Then  $C^*(TV, TV)$  is an  $E_2$ -algebra. How close is  $H_*^{E_2}(\bar{C}^*(TV, TV))$  to  $V$ , i.e., how free is  $C^*(TV, TV)$  as an  $E_2$ -algebra?

### Proposition

For  $V = \mathbb{Q}$ , i.e.  $TV = \mathbb{Q}[x]$  the  $E_2$ -homology of the reduced Hochschild cochain complex is non-trivial in all degrees  $r \geq -1$ , more precisely

$$H_r^{E_2}(\bar{C}^*(\mathbb{Q}[x], \mathbb{Q}[x])) \cong \mathbb{Q}$$

for all  $r \geq -1$ .

Thus in this case  $E_2$ -homology of the Hochschild cochains on  $T\mathbb{Q}$  is much larger than the vector space  $\mathbb{Q}$  we started with.



The calculations uses the equivalence of categories of  $n$ -Lie algebras and graded Lie-algebras given by  $n$ -fold (de)suspension. Thus (up to suspension) we have to calculate ordinary Lie-homology of  $AQ_*(HH^*(\mathbb{Q}[x], \mathbb{Q}[x]))$  and this is concentrated in homological degree zero and there it is  $\mathbb{Q}\langle x_0, y_{-1} \rangle$  with trivial 1-Lie structure.

$$\infty > \dim(V) \geq 2$$

In these cases we can determine the input for the Blanc-Stover spectral sequence:

$$AQ_*(HH^*(TV, TV)|\mathbb{Q}; \mathbb{Q}) \cong HH_{*+1}^{(1)}(\mathbb{Q} \rtimes M(-1); \mathbb{Q})$$

and the first Hodge summand  $HH_*^{(1)}(\mathbb{Q} \rtimes M(-1); \mathbb{Q})$  is additively isomorphic to the free graded Lie-algebra generated by the graded vector space  $M(-1)$ .

However,  $M(-1) = HH^1(TV, TV)$  is *not* free as a Lie-algebra.

# Chains on iterated loop spaces

Let  $k$  be  $\mathbb{Q}$  and let  $X$  be an  $(n + 1)$ -connected nice topological space.

Then  $H_*(\Omega^{n+1}X; \mathbb{Q}) \cong S(\Sigma^{-n}\pi_*(\Omega X) \otimes \mathbb{Q})$  as  $n$ -Gerstenhaber algebras.

## Proposition

$$\mathbb{L}_s(\mathbb{Q}_{nG})(H_*(\Omega^{n+1}X; \mathbb{Q}))_q \cong \mathrm{Tor}_{s+1, q+n}^{H_*(\Omega X; \mathbb{Q})}(\mathbb{Q}, \mathbb{Q}).$$

## Conjecture

This does not just look like a shifted version of the Rothenberg-Steenrod  $E^2$ -term, but there is an underlying isomorphism of spectral sequences. The RS-spsq converges to  $H_*(X; \mathbb{Q})$ .