




# REINWARDTIA

A JOURNAL ON TAXONOMIC BOTANY, PLANT SOCIOLOGY AND ECOLOGY

ISSN 0034 – 365 X | E-ISSN 2337 – 8824 | Accredited 792/AU3/P2MI-LIPI/04/2016



# REINWARDTIA

*A JOURNAL ON TAXONOMIC BOTANY, PLANT SOCIOLOGY AND ECOLOGY*

Vol. 15 (1): 1 – 65, June 27, 2016

## Chief Editor

Kartini Kramadibrata (Mycologist, Herbarium Bogoriense, Indonesia)

## Editors

Dedy Darnaedi (Taxonomist, Herbarium Bogoriense, Indonesia)

Tukirin Partomihardjo (Ecologist, Herbarium Bogoriense, Indonesia)

Joeni Setijo Rahajoe (Ecologist, Herbarium Bogoriense, Indonesia)

Marlina Ardiyani (Taxonomist, Herbarium Bogoriense, Indonesia)

Himmah Rustiami (Taxonomist, Herbarium Bogoriense, Indonesia)

Lulut Dwi Sulistyarningsih (Taxonomist, Herbarium Bogoriense, Indonesia)

Topik Hidayat (Taxonomist, Indonesia University of Education, Indonesia)

Eizi Suzuki (Ecologist, Kagoshima University, Japan)

Jun Wen (Taxonomist, Smithsonian Natural History Museum, USA)

Barry J Conn (Taxonomist, School of Life and Environmental Sciences, The University of Sydney, Australia)

David G. Frodin (Taxonomist, Royal Botanic Gardens, Kew, United Kingdom)

## Secretary

Endang Tri Utami

Rina Munazar

## Layout

Dede Aryanto

## Illustrators

Subari

Wahyudi Santoso

Anne Kusumawaty

Correspondence on editorial matters and subscriptions for Reinwardtia should be addressed to:

HERBARIUM BOGORIENSE, BOTANY DIVISION,  
RESEARCH CENTER FOR BIOLOGY–INDONESIAN INSTITUTE OF SCIENCES  
CIBINONG SCIENCE CENTER, JLN. RAYA JAKARTA – BOGOR KM 46,  
CIBINONG 16911, P.O. Box 25 CIBINONG  
INDONESIA  
PHONE (+62) 21 8765066; Fax (+62) 21 8765062  
E-MAIL: reinwardtia@mail.lipi.go.id  
<http://e-journal.biologi.lipi.go.id/index.php/reinwardtia>

A	B	
C	D	E
F	G	H
	I	J

Cover images: *Begonia manuselaensis* Ardaka & Ardi. A. Plant habit in situ; B. Plant habit ex situ; C. Lamina abaxial surface with the small red scales on the veins; D. Stipules. E. Male inflorescence; F. Solitary female inflorescence; G. Male flower; H. Female flowers; I. Ovary cross section; J. Fruit. Source of materials: Wisnu Ardi, WI 104 (BO, KRB, SING). Photo credits: (B-J) by Wisnu Ardi; A: I.G. Tirta

**The Editors would like to thank all reviewers of volume 15(1):**

Andrew Henderson - New York Botanical Garden, Bronx, New York, USA  
Andrew Powling - School of Biological Sciences, University of Portsmouth, Portsmouth, UK  
Axel D. Poulsen - Royal Botanic Garden Edinburgh, Edinburgh, Scotland, UK  
Jeffrey Lee Walck - Tennessee State University, Tennessee, USA  
Mark Hughes - Royal Botanic Garden Edinburgh, Edinburgh, Scotland, UK  
Mien A. Rifai - Akademi Ilmu Pengetahuan Indonesia (AIPI)  
Wong Khoon Meng - Singapore Botanic Garden, Singapore



## A STUDY OF THE UNDERGROWTH VEGETATION OF SEMPU ISLAND, EAST JAVA, INDONESIA

Received 03 September 2015; accepted 26 May 2016

**ASEP SADILI**

*Herbarium Bogoriense, Research Center for Biology, Indonesian Institute of Sciences, Jln. Raya Jakarta-Bogor Km 46, Cibinong 16911, Indonesia. Email: asep.sadili@gmail.com*

**KUSWATA KARTAWINATA**

*Integrative Research Center, The Field Museum, 1400 Lake Shore Drive, Chicago, IL 60605, USA. Email: kkjak@indo.net.id*

### ABSTRACT

SADILI, A. & KARTAWINATA, K. 2016. A study of the undergrowth vegetation of Sempu Island, East Java, Indonesia. *Reinwardtia* 15(1): 1 – 9. — A study of forest floor vegetation in coastal forest (site 1) and inland lowland forest (site 2) was carried out at the Sempu Island Nature Reserve, Malang Regency, East Java. The objective of the study is to obtain data on plant species diversity, species composition and structure of the forest floor vegetation. In each site the vegetation was sampled with a plot of 1 m × 50 m, which was divided into 50 subplots of 1 m × 1 m each. Overall from the two plots we recorded 59 species, 57 genera and 39 families. Shannon-Wiener diversity indices (H') were relatively high. *i.e.*, 4.47 in Plot 1 and 3.2 in Plot 2, with a mean of 3.84. The number of the seedlings of shrubs and trees was greater than that of the herbaceous species. The families having the highest number of species were Euphorbiaceae (6 species) and Fabaceae (5 species). Based on dominant species the vegetation in Plot 1 was designated as *Scleria lithosperma-Asystasia nemorum* community type, while in Plot 2 *Pterospermum javanicum-Knema* sp. community type. The similarity index between these community types was only 18%. Only *P. javanicum* showed a good regeneration and the regeneration of other species in the two community types was poor, indicating unclear floristic changes in the forest of the Sempu Island.

**Key words:** Seedling and herbaceous composition, Sempu Island Nature Reserve, species diversity.

### ABSTRAK

SADILI, A. & KARTAWINATA, K. 2016. Kajian vegetasi lantai hutan di Pulau Sempu, Jawa Timur, Indonesia. *Reinwardtia* 15(1): 1 – 9. — Penelitian vegetasi lantai di hutan pantai (lokasi 1) dan hutan pamah (lokasi 2) dilakukan di Cagar Alam Pulau Sempu, Kabupaten Malang, Jawa Timur. Tujuan penelitian untuk memperoleh data keanekaragaman dan komposisi jenis serta struktur vegetasi lantai hutan. Di setiap lokasi vegetasi dicuplik dengan menggunakan petak memanjang berukuran 1 m × 50 m yang dibagi menjadi 50 subpetak masing-masing berukuran 1 m × 1 m. Dua petak pencuplikan tersebut merekam 59 jenis 57 marga dari 39 suku. Indeks keanekaragaman jenis Shannon-Wiener (H') relatif tinggi, yaitu 4,47 di Petak 1 dan 3,2 di Petak 2, dengan rerata 3,84. Jenis-jenis semai pohon dan perdu lebih banyak dibandingkan dengan jenis-jenis terna. Suku yang memiliki jumlah jenis terbanyak adalah Euphorbiaceae (6 jenis) dan Fabaceae (5 jenis). Berdasarkan jenis dominan vegetasi di Petak 1 dinamakan tipe komunitas *Scleria lithosperma-Asystasia nemorum* sedangkan di Petak 2 disebut tipe komunitas *Pterospermum javanicum-Knema* sp. Indeks kesamaan antara dua tipe komunitas tersebut hanya 18%. Hanya *P. javanicum* yang menunjukkan regenerasi yang baik sedangkan regenerasi pohon jenis lainnya sebagian besar di kedua tipe komunitas tersebut tidak baik, sehingga arah perubahan komposisi komunitas hutan di Pulau Sempu tidak cukup jelas.

**Kata kunci:** Cagar Alam Pulau Sempu, keanekaragaman jenis, komposisi jenis terna dan semai.

### INTRODUCTION

Indonesia has a high number of vegetation types representing natural ecosystems with a high degree of plant species diversity, comprising the littoral vegetation at the sea level up to the nival vegetation type on permanent snow at the highest mountain top of Mt. Jaya in Papua (Kartawinata, 2005). Much of the lowland forests have suffered destruction by timber logging, coal and mineral mining, conversion into timber, rubber and oil palm plantations, human settlements and other uses. The remaining undisturbed primary lowland forests today can be found mainly in various protected areas, such as national parks and nature reserves. The Sempu Island Nature Reserve (Cagar

Alam Pulau Sempu), or Sempu Island NR for short, in East Java represents one of such areas.

To date the vegetation studies in Indonesia have been concerned mainly with the lowland forests, focusing primarily on tree life form (Kartawinata, 2005) and very few investigations have devoted to undergrowth communities (*e.g.* Ismail, 1975; Sumardja & Kartawinata, 1977; Sadili *et al.*, 2009; Sadili, 2010). In rain forests, trees are the dominant life form and the undergrowth consists of tree seedlings, shrubs and herbs, while the so called “herb layer” is a synusia of tree seedlings (Whitmore, 1986). Studies in tropical forest elsewhere revealed that herb diversity contributes a great deal to the total species diversity (Cicuzza *et al.*, 2013; Gentry & Dodson, 1987; Kiew, 1978;

Meijer, 1959; Nath *et al.*, 2005; Poulsen, 1996; Ting & Poulsen, 2009; Tran *et al.*, 2005). The understory vegetation in South American tropical forests is said to be rich in species, with herbaceous plants amounting to 21 to 47% of the total plant species (Croat, 1978; Gentry & Dodson, 1987). It has been reported that herbaceous plants represent 8 to 29% of the total plant species richness in moist to wet forests, 53% in dry forests and 6% in forests on white sandy soils (Hall & Swaine, 1976; Gentry & Dodson, 1987; Gentry & Emmons, 1987). Herbs and seedlings constitute an integral part of and have important functions in a forest ecosystem, hence their conservation is necessary (Irwanto, 2006). In lowland forests, dense herbs often form patches of closed vegetation in natural and man-made gaps and along foot paths and rivers that are well illuminated, but in the interior of mature rain forest, herbs occur mainly as scattered individuals and are even absent (Richards, 1996).

In view of these phenomena we undertook a study on the undergrowth communities in the coastal and lowland forests on the Sempu Island NR. Previous studies on the island have been concerned with the structure and composition of the tree layers in the mangrove forest (Prawiroatmodjo, 2013) and the dryland forests (Polosakan 2011, 2014) as well as with the conservation status of species, such as *Myristica teysmannii* (Risna, 2009). To date no studies on undergrowth communities have been previously undertaken on this island. In this study we took into a consideration the observation of Richards (1996) that the synusia of ground herbs is not synonymous with the "ground flora" of lowest (E) stratum because the majority of the plants in third layer are young trees, shrubs and woody climbers.

In our study the undergrowth community covered the E layer comprising herbs, seedlings of trees and shrubs but excluded epiphytes. We expect that the study will be useful to complement existing forest tree data needed for the management of the Sempu Island NR. Research is necessary to understand the relationship between various components of the ecosystem preserved, to facilitate and support the management. The objective of the present study is to obtain basic ecological data in the form of the structure and composition of the undergrowth communities in the coastal and the inland lowland forests that will be of use in the management of the reserve.

## MATERIALS AND METHODS

The Sempu Island NR covers the entire island with the total area of  $\pm 877$  ha and is managed by the Sub-Seksi Konservasi Sumber Daya Alam Malang (Malang Natural Resources Conservation

Subsection) of the Balai Besar Konservasi Sumber Daya Alam Jawa Timur (East Java Office of Natural Resource Conservation) (Kehutanan, 2009; MOF, 2009). Administratively it belongs to the Sendang Biru Subdistrict, Sumbermanjing Wetan District, Malang Regency and East Java Province. Geographically it is located at  $112^{\circ}40'45'' - 112^{\circ}42'45''$  East and  $8^{\circ}27'24'' - 8^{\circ}24'54''$  South. On the north side it borders with the Sempu Bay while on the east, south and north sides it is surrounded by the Indian Ocean (Fig. 1).

In the rainfall scheme of Schmidt & Ferguson (1951) the Sempu Island falls under the rainfall type C, indicating the slightly wet climate, where the prevalent natural vegetation is evergreen forest with an indication of slight seasonality by the presence of few deciduous tree species (Kartawinata, 2013). Fig. 2 is the climate diagram for Sumbermanjing weather station, about 60 km north of Sempu Island, based on Walter's (1973) method, using the 30-year rainfall data of LMG (1967) and Climate-Data.Org (2015) for temperature. It shows the drought period during the lowest rainfall in August (29 mm) and September (23 mm) and the wet period between October (115 mm) and July (119 mm), with the highest rainfall (322 mm) in December. In the climate diagram the drought period is indicated by the chart area enclosed by the rainfall curve and the temperature curve during the months of August and September, where during this period the evapotranspiration is assumed to exceed the precipitation (Walter, 1973). This period is critical for the life of plants, especially herbaceous plants that may be wilting and even dead. The soils on the island is rocky covered with thin layer of litter and having low moisture, low fertility, pH of 6-7 and temperature of  $30-32^{\circ}\text{C}$  (Kramadibrata *et al.*, 2010).

Mangrove forest, coastal forest, lowland forest and lake-side forest can be found on the Sempu Island. Some ecosystems have been disturbed by touristic activities and other human uses by people living in the surrounding areas. The field study was conducted from 29 September to 10 October 2010 within the research plots of Polosakan (2011 & 2014) located in a coastal forest (Site 1) at Waruwaru and a lowland forest (Site 2) at Telonpring (Fig. 1). Site 1 was located at  $08^{\circ}25'50.76''$  S and  $112^{\circ}41'40.76''$  E'. The forest at Site 1 was slightly disturbed by touristic activities. The soil was calcareous with a thin litter layer. The study by Polosakan (2011) revealed that the coastal forest was structurally dominated by small trees (diameter  $< 20$  cm) with a density of 705 trees/ha. The prevalent species, with the importance value  $> 10$ , were *Mallotus floribundus*, *Ficus* sp., *Drypetes ovalis*, *Artocarpus elasticus*, *Vitex pinnata*, *Aglaia odoratissima*, *Pterospermum diversifolium* and *Terminalia* sp. The families with the highest number of species




Fig. 1. The geographic position of the Sempu Island showing the study sites on the northern side of the island near Waruwaru.

were Euphorbiaceae, Sterculiaceae and Anacardiaceae. Site 2 at  $08^{\circ}26'13.13''$  S and  $112^{\circ}41'56.56''$  E was selected in the lowland forest located further inland which was denser than forest in Site 1 and was still in very good condition with no human disturbance. The soil was moister and thicker and physiography was more undulating than that in Site 1. Polosakan (2014) recorded that the tree density was 1005 trees/ha dominated by trees with diameters  $< 20$  cm and canopy height of 15-20 m. The prevalent tree species were *Streblus asper*, *Aglaia argentea*, *Drypetes longifolia* and *Pseuduvaria reticulata*. The emergent trees with height  $> 25$  m were *Pterospermum javanicum*, *Terminalia subspathulata*, *Bischofia javanica* and *Cratoxylum sumatranum*. The families with the highest number of species were Euphorbiaceae, Annonaceae, Clusiaceae and Sterculiaceae.

The undergrowth communities in each site was sampled with a plot of  $1 \text{ m} \times 50 \text{ m}$ , which was further divided into 50 subplots of  $1 \text{ m} \times 1 \text{ m}$  each. Within each subplot all species of herbs and seedlings (woody plants with height of  $< 50$  cm) were identified; the number of individuals and the percentage of cover of each species were measured. Voucher specimens of each species were collected and identified at the Herbarium Bogoriense, Research Center of Biology-LIPI, Cibinong. The nomenclature of plant names follows Backer & Bakhuizen van den Brink, Jr. (1963, 1965, 1968). Validation name for the new version was obtained through the Plant List. In each subplot the following habitat factors were measured: slope, soil properties (pH, moisture and fertility) with a portable direct soil meter and the

light reaching ground was estimated by the percentage of canopy opening. Data analysis follows the method of Mueller-Dombois & Ellenberg (1974) covering Relative Density (RD), Relative Dominance (RDo), Relative Frequency (RF) and Importance Value ( $IV = RD + RDo + RF$ ). Sorensen's similarity index based on density between the two sites was calculated using the Biodiversity Pro version 2 and diversity index ( $H'$ ) was calculated using the formula of Shannon-Wiener (Barbour *et al.*, 1987).

## RESULTS AND DISCUSSION

### Floristic composition

In the two plots studied we recorded 59 species, 57 genera and 39 families, with density of 315,000 individuals/ha, comprising 13 (22.03%) species of herbs and 46 (77.97%), species of woody seedlings (Table 1). It agrees with Whitmore (1986), who mentioned that in tropical forests the ground herb synusia is not so rich in species as the woody part. The low occurrence of herbs in our study was likely due to the fact that the study was carried out during the drought period in September-October (Fig. 2), where most herbs must have wilted and died away, leaving only the hardy species that could withstand drought, which in the present study were represented by *Apostasia* sp., *Asystasia nemorum*, *Cayratia japonica*, *Hedychium* sp., *Panicum* sp. *Pleomele* sp. and *Scleria lithosperma*.

Of the total 59 species, 53 species occurred in Plot 1 and 32 species in Plot 2, respectively. The two plots differed greatly with the index of similarity of only 18%. Similarity between the two




Fig. 2. Climate diagram for Sumbermanjing weather station, about 60 km north of Sempu Island, based on Walter's (1973) method, using data from LMG (1967) for rainfall and Climate-Data.Org (2015) for temperature. It shows a drought period in August and September and wet period between October and July.

plots were attributed to 26 species in Group 1 (Table 1), which were present in both plots at different values of density, dominance and frequency thus importance values. *Pterospermum javanicum*, *Knema* sp. *Microcos argentata*, *Ficus* sp., *Antidesma bunius*, *Ilex cymosa*, *Terminalia catappa* and *Aglaia argentea* were dominating (IV = 8.69–76.73%). The other species in Group 1 (Table 1) shared together by Plot 1 and Plot 2 were less significant compared to the above dominant species as they had much lower density, cover, frequency and importance values. The habitats vary from one place to another within an island, resulting in the variation in species diversity. Such a phenomenon occurred in the forest of the Sempu Island, as shown by the Shannon-Wiener Diversity Index of 4.47 for Plot 1 and 3.2 for Plot 2, with the mean of 3.84. Applying the criteria of Barbour *et al.* (1987) the plant species diversity in the forests of the Sempu Island was considered high ( $H' = 3-4$ ) to very high ( $H' > 4$ ).

Plot 1 was dominated by a sedge species, *Scleria lithosperma* (IV = 56.56), which was absent in Plot 2, indicating the community was a secondary growth, as it commonly grows in communities that developed abundantly following the clearance of primary forests as well as on open rocks and sandy habitats (Kern, 1974). The species was fairly well distributed throughout the subplots as indicated by its high frequency (RF = 5.63%). The second dominant species was *Asystasia nemorum* (IV = 31.24). The two species had the widest distribution in the subplots as indicated by highest RF (5.63% and 7.81%, respectively).

Based on these vegetation parameters the two species can be used to name the undergrowth community in Plot 1 as the "*Scleria lithosperma-Asystasia nemorum* community type". The community was further complemented by Species Group 2 (Table 1), consisting of 28 species, having much lower RF and IV and occurring only in Plot 1, hence characterizing the undergrowth vegetation in the coastal forest of the Sempu Island. It is interesting to note that there were only *Guettarda speciosa*, *Hibiscus tiliaceus* and *Terminalia catappa* representing the typical coastal forest species, especially those characteristics of the "*Barringtonia* formation". Similar situation was also observed in the composition of the tree layer reported by Polosakan (2011).

Plot 2 was jointly dominated by *Pterospermum javanicum*, *Knema* sp. and *Microcos argentata*, which were the most prevalent in having high values of dominance, density and frequency (Table 1). Other prevalent species with lower IV were *Ficus* sp., *Antidesma bunius*, *Ilex cymosa*, *Terminalia catappa* and *Aglaia argentea*. It should be noted that *Pterospermum javanicum* is a typical species found in forests on small islands and inland lowland forests in West Java (Ismail, 1975; Kartawinata, 1977; Kartawinata *et al.*, 1985; Sumardja & Kartawinata, 1977). Of interest was *Terminalia catappa*, which is generally the main component of any coastal forest (Kartawinata, 2013; Steenis & Schippers-Lammerstse, 1965), was more prevalent in Plot 2 representing the inland lowland forest, indicating the sign of



invasion. Species in Group 3 characterized the undergrowth community in the lowland inland primary forest, as the occurrence of these species was restricted to Plot 2. Except for *Garcinia* sp. and *Myristica teysmannii*, they are secondary growth species as they are commonly growing in the open forests and dry scrubs on open places especially in teak forest and seasonal forests (Backer & Bakhuizen van den Brink, Jr., 1963, 1965).

The microclimatic condition under the shade of tall trees and short trees in a forest with closed canopy is different from that under a forest with canopy gaps. In a closed-canopy forest, the light reaching the ground is low, humidity is high, temperature is low and evapotranspiration is low (Richard, 1996). Under such conditions the herbaceous species and seedlings do not always grow well and normal, as have been observed in Plot 1. The floristic differences between Plot 1 and Plot 2 were apparently attributed to the differences in the habitat conditions in the two plots, in particular the canopy gaps and slope (Table 1). The percentage of canopy gaps in Plot 1 was higher than in Plot 2, which was related to the higher density of trees and the more closed canopy in Plot 2. In a lowland forest, dense herbaceous vegetation frequently occur in natural and man-made openings, along paths and rivers, where the solar radiation is good, but in the interior of mature rain forest, herbs are found chiefly as scattered individuals and are sometimes absent (Richards, 1996). Herbs are common along streams, other wet places and canopy gaps, allowing greater penetration of light (Richards, 1996; Whitmore, 1986). Our field observations elsewhere show that undergrowth species may form patches of closed vegetation and are found mainly beneath canopy gaps.

The species-area curves (Fig. 3) initially showed the increase of the number of species with the

increase in areas in the two plots, but beginning to level off at the areas of 4000 m<sup>2</sup>– 4900 m<sup>2</sup> (40<sup>th</sup> – 49<sup>th</sup> subplots), indicating that the area of 50 m<sup>2</sup> may be considered a minimum area to sufficiently register the representative number of species within the community. It was further confirmed also by the regression index ( $R^2$ ) of  $\pm 0.927$  for Plot I and  $\pm 0.970$  for Plot II, which were approaching 1, implying that further increase of the number of species was stable or constant, *i.e.* one or two species only.

The species composition of the undergrowth communities in the Sempu Island NR was considered high compared to those in other communities along the south coast of Java *e.g.* Ismail (1975) at Sukawayana, Sumardja & Kartawinata (1977) at Pangandaran and Sadili (2010) at Cikepuh. The differences could be attributed to different natural habitat conditions as well as to human disturbance factors.

#### Distribution and regeneration

The distributional patterns of herbs and seedlings in the Sempu Island NR, as expressed by the frequency values, were the product of interactions among ecological processes that have been progressing thus far, combined with the reproductive capacity and adaptability of the species to the environment (Barbour *et al.*, 1987; Sadili, 2010). Table 1 shows that *Asystasia nemorum*, *Syzygium* sp., *Scleria lithosperma* and *Ficus* sp. in Plot 1 had a relatively better distribution compared to the others (RF > 5%), but in Plot 2 the RF values for *Asystasia nemorum* and *Syzygium* sp. were lower, except for *Ficus* sp. which was higher (FR=8.46%), while *Scleria lithosperma* was not present at all. Species with high frequency (FR > 5%) were *Knema* sp., *Microcos argentata*, *Pterospermum javanicum*, *Ficus* sp., *Ilex cymosa*, *Antidesma bunius* and *Terminalia catappa*. They were recorded in Plot 2




Fig. 3. Species-area curve for undergrowth communities in a coastal forest (Plot 1) and a lowland forest (Plot 2) at the Sempu Island NR, Malang, East Java.

Table 1. Species of tree (T), shrub (S), herb (H), and liana (L) in Plot 1 and Plot 2 expressed in terms of Relative Density (RD), Relative Dominance (RDo), Relative Frequency (RF) and Importance Value (IV= RD+RDo+RF), vegetation characteristics and hábitat of the undergrowth communities in the coastal and inland lowland forests of the Sempu Island Nature Reserve, Malang, East Java

No	Habitat	Site								
		Plot 1				Plot 2				
	Soil pH	6.78 ± 0.27				6.65 ± 0.41				
	Humidity (%)	30.71 ± 0.68				30.73 ± 1.08				
	Soil fertility scale	1.16 ± 0.41				1.48 ± 0.50				
	Canopy gap (%)	29.53 ± 11.91				9.68 ± 11.73				
	Slope (%)	5.00 ± 9.92				12.10 ± 13.52				
<b>Vegetation characteristics</b>										
	Density (Individuals/1 ha)	317.600				312.400				
	Number of species	53				32				
	Number of genus	47				30				
	Number of family	35				22				
	Shannon Diversity Index (H')	4.47				3.2				
	Species	Life-form	RDo	RD	RF	IV	RDo	RD	RF	IV
<b>Species Group 1</b>										
1	<i>Pterospermum javanicum</i> Jungh.	T	1.00	1.25	3.13	5.38	33.81	30.54	12.39	76.73
2	<i>Knema</i> sp.	T	0.88	0.13	0.63	1.64	13.93	17.67	15.71	47.31
3	<i>Microcos argentata</i> Burret	T	1.04	0.66	1.88	3.58	15.09	16.84	13.90	45.82
4	<i>Ficus</i> sp.	T	2.49	6.27	5.00	13.76	10.24	8.19	8.46	26.89
5	<i>Antidesma bunius</i> (L.) Spreng.	T	1.24	1.32	1.88	4.43	7.02	6.02	6.65	19.69
6	<i>Ilex cymosa</i> Blume	T	0.92	0.26	0.94	2.13	3.61	4.55	7.55	15.71
7	<i>Terminalia catappa</i> L.	T	1.28	1.45	3.44	6.17	2.38	2.88	6.34	11.61
8	<i>Aglaia argentea</i> Blume	T	1.02	0.59	1.25	2.87	2.11	2.05	4.53	8.69
9	<i>Lithocarpus sundaicus</i> (Blume) Rehder	T	1.04	0.66	1.25	2.95	0.78	0.90	1.21	2.88
10	<i>Desmodium</i> sp.	S	1.35	1.72	2.50	5.57	0.62	0.58	1.51	2.70
11	<i>Ardisia</i> sp.	T	1.06	0.73	0.63	2.41	0.64	0.64	1.21	2.49
12	<i>Mitrephora polypyrena</i> Miq.	T	0.92	0.26	0.31	1.50	0.25	0.38	1.81	2.45
13	<i>Crotalaria</i> sp.	S	0.88	0.13	0.31	1.33	0.30	0.32	1.21	1.83
14	<i>Polyalthia</i> sp.	T	1.10	0.86	1.88	3.83	0.56	0.38	0.60	1.55
15	<i>Dalbergia latifolia</i> Roxb.	T	0.87	0.07	0.31	1.24	0.34	0.38	0.60	1.32
16	<i>Lasianthus</i> sp.	T	1.24	1.32	2.19	4.74	0.12	0.19	0.91	1.22
17	<i>Streblus asper</i> Lour.	T	1.14	0.99	1.25	3.38	0.21	0.32	0.60	1.13
18	<i>Diospyros cauliflora</i> Blume	T	1.55	2.37	3.13	7.05	0.24	0.19	0.60	1.03
19	<i>Psychotria sarmentosa</i> Blume	T	1.65	2.70	3.75	8.10	0.19	0.13	0.60	0.93
20	<i>Pterospermum diversifolium</i> Blume	T	1.06	1.25	3.75	6.06	0.12	0.13	0.60	0.85

Table 1. Species of tree (T), shrub (S), herb (H), and liana (L) in Plot 1 and Plot 2 expressed in terms of Relative Density (RD), Relative Dominance (RDo), Relative Frequency (RF) and Importance Value (IV= RD+RDo+RF), vegetation characteristics and hábitat of the undergrowth communities in the coastal and inland lowland forests of the Sempu Island Nature Reserve, Malang, East Java (continued)

21	<i>Barringtonia racemosa</i> (L.) Spreng.	T	1.35	1.72	1.88	4.94	0.08	0.06	0.30	0.44
22	<i>Engelhardia</i> sp.	T	1.08	0.79	2.50	4.37	0.18	0.13	0.60	0.92
23	<i>Apostasia</i> sp.	H	0.96	0.40	0.31	1.67	0.09	0.13	0.30	0.52
24	<i>Garcinia celebica</i> L.	T	0.88	0.13	0.31	1.33	0.06	0.06	0.30	0.43
25	<i>Syzygium</i> sp.	T	1.61	3.30	5.63	10.54	0.40	0.26	0.60	1.26
26	<i>Asystasia nemorum</i> Nees	H	5.56	17.88	7.81	31.24	3.73	3.07	3.63	10.43
<b>Species Group 2</b>										
27	<i>Scleria lithosperma</i> (L.) Sw.	H	34.18	16.75	5.63	56.56				
28	<i>Cleistanthus glaber</i> Airy Shaw	T	3.29	10.36	4.38	18.02				
29	<i>Knema laurina</i> Warb.	T	2.23	6.79	4.69	13.71				
30	<i>Cayratia japonica</i> (Thunb.) Gagnep.	L	1.35	1.72	4.06	7.13				
31	<i>Bambusa</i> sp.	S	1.55	3.10	2.19	6.84				
32	<i>Guettarda speciosa</i> L.	T	1.10	0.86	2.19	4.15				
33	<i>Lophopetalum javanum</i> Turcz.	T	1.37	1.78	2.19	5.34				
34	<i>Panicum</i> sp.	H	1.34	1.65	1.56	4.55				
35	<i>Syzygium claviflorum</i> (Roxb.) Wall. ex A.M.Cowan & Cowan	T	1.14	0.99	1.88	4.00				
36	<i>Hibiscus tiliaceus</i> L.	T	1.20	1.19	1.56	3.95				
37	<i>Beilschmiedia madang</i> Blume	T	1.08	0.79	1.56	3.43				
38	<i>Peltophorum inerme</i> (Roxb.) Naves	T	1.06	0.73	1.56	3.35				
39	<i>Ficus punctata</i> Thunb.	T	1.02	0.59	1.25	2.87				
40	<i>Glochidion</i> sp.	T	1.02	0.59	0.94	2.55				
41	<i>Santiria</i> sp.	T	1.02	0.59	0.94	2.55				
42	<i>Dioscorea</i> sp.	L	0.94	0.33	1.25	2.52				
43	<i>Diospyros sumatrana</i> Miq.	T	1.02	0.59	0.63	2.24				
44	<i>Diospyros</i> sp.	T	0.98	0.46	0.63	2.07				
45	<i>Hedychium</i> sp.	H	0.88	0.13	0.63	1.64				
46	<i>Clematis nepalensis</i> Royle	L	0.90	0.20	0.31	1.41				
47	<i>Pleomele</i> sp.	S	0.90	0.20	0.31	1.41				
48	<i>Allophylus cobbe</i> (L.) Raeusch.	T	0.87	0.07	0.31	1.24				
49	<i>Artocarpus elasticus</i> Reinw. ex Blume	T	0.87	0.07	0.31	1.24				
50	<i>Diospyros javanica</i> Bakh.	T	0.87	0.07	0.31	1.24				
51	<i>Mallotus floribundus</i> (Blume) Müll.Arg.	T	0.87	0.07	0.31	1.24				
52	<i>Memecylon</i> sp.	T	0.87	0.07	0.31	1.24				
53	<i>Polyscias fruticosa</i> (L.) Harms	T	0.87	0.07	0.31	1.24				
<b>Species Group 3</b>										
54	<i>Croton argyratus</i> Blume	T					1.49	1.28	2.11	4.89
55	<i>Garcinia</i> sp.	T					0.45	0.58	1.51	2.54
56	<i>Myristica teysmannii</i> Miq.	T					0.28	0.26	1.21	1.75
57	<i>Dillenia</i> sp.	T					0.15	0.19	0.6	0.95
58	<i>Aporosa aurita</i> (Tul.) Miq.	T					0.09	0.13	0.3	0.52
59	<i>Leea aequata</i> L.	S					0.06	0.06	0.3	0.43

and also in Plot 1 but with lower values (Table 1). Species which were present with relatively high frequencies in Plot 1 and Plot 2 signified that they had better distribution and regeneration than the other species, implying that the habitat conditions at the Sempu Island NR were favorable for their survival and good growth. Polosakan (2011, 2014) recorded 49 species of trees in the coastal forest in the Sempu Island, of which, in the present study, only 9 species (18.37%) occurred in the seedling stage and of 63 species of trees in the lowland forest only 14 species (22.22%) were seedlings. It indicated that the regeneration in these forests were poor. The standing trees in the plots of Polosakan (2011, 2014) were represented only by *Aglaia argentea*, *Artocarpus elasticus*, *Diospyros cauliflora*, *Garcinia celebica*, *Guettarda speciosa*, *Mallotus floribundus*, *Mytrephora polyneura*, *Peltophorum inerme*, *Pterospermum diversifolium*, and *P. javanicum*. Among these species only *P. javanicum* had the highest IV = 76.73 and RF = 30.54% in Plot 2 and IV = 5.38 and RD = 1.25% in Plot 1. It implies that only *P. javanicum* will maintain itself and perhaps become more dominant in the inland forest in the future. This is, however, only weak indication and the situation will be different if the study is further enlarged by establishing more samples of tree and undergrowth layers over larger areas. *Myristica teysmannii* which was classified as endangered species (Risna, 2009) was present only in Plot 1 with relatively high density (RD = 17.67%) and frequency (RF = 15.71 hence it had a good chance to survive in the area.

## CONCLUSIONS

The undergrowth flora in the coastal forest formed the *Scleria lithosperma*—*Asystasia nemorum* community type and in the inland lowland forest the *Pterospermum javanicum*—*Knema* sp. community type and each community type had a high species diversity. The floristic similarity between the two communities was very low. The floristic composition in the two communities did not reflect a good regeneration of the forests and the direction of floristic change and the future tree composition of the forest was not clear. Only *Pterospermum javanicum* will maintain itself and perhaps become more dominant in the inland forest in the future. This is, however, only an indication and the situation will be more affirmative if the study is further enlarged to cover larger areas. Further investigation of undergrowth and tree layers in the coastal and inland forest over a wider area with higher sampling intensity is needed to provide good data for better management of the Sempu Island NR. The pressure of human activities including economic development upon the reserve is greatly

increasing which should be checked to ensure the future existence and viability of the reserve. More studies on floristic composition and structure of various plant communities in protected and non-protected areas should be carried out to clarify scientific explanation concerning the cause of patterns of species richness and also to provide data for utilization, management and conservation of biodiversity (Kartawinata, 2005).

## ACKNOWLEDGEMENTS

We are grateful to the Head of Herbarium Bogoriense the Botany Division, the Head of the Biology Research Center-LIPI and the Directorate of Higher Education of the Ministry of National Education (through the Ministry of Research and Technology) for providing financial support. We also thank the Head of the East Java Office of Natural Resource Conservation (Balai Besar Konservasi Sumber Daya Alam Jawa Timur), for providing a research permit and to Pak Ipin, the Head of the Sempu Island NR and Pak Saihu for providing field facilities and assistance. We should acknowledge also the members of the research team (*i.e.* Kartini Kramadibrata, Suhardjono, Ruddy Polosakan, Florentina I. Windadri, Agus Sujadi, Idang Sumanta and Dewi Rosalina) for their cooperation during the field work.

## REFERENCES

- BACKER, C. A. & BAKHUIZEN VAN DEN BRINK, Jr. R. C. 1963 & 1965. *Flora of Java*, Vol. 1 & 2. N. V. P. Noordhoff, Groningen.
- BACKER, C. A. & BAKHUIZEN VAN DEN BRINK, Jr. R. C. 1968. *Flora of Java*, Vol. 3. Wolters-Noordhoff N. V., Groningen.
- BARBOUR, G. M., BURCK, J. K. & PITTS, W. D. 1987. *Terrestrial Plant Ecology*. The Benjamin/Cummings Publishing Inc., New York.
- CICUZZA, D., KRÖMER, T., POULSEN, A. D., ABRAHAMCZYK, S., DELHOTAL, T., PIEDRA, H. M. & KESSLER, M. 2013. A transcontinental comparison of the diversity and composition of tropical forest understory herb assemblages. *Biodiversity and Conservation* 22: 755–772.
- CLIMATE-DATA.ORG. 2015. <http://id.climate-data.org/location/622273>.
- CROAT, T. B. 1978. *Flora of Baro Colorado Island*. Stanford University Press, Stanford.
- GENTRY, A. & DODSON, C. 1987. Contribution of non-trees to the species richness of a tropical rain forest. *Biotropica* 19: 149–156.
- GENTRY, A. H. & EMMONS, L. H. 1987. Geographical variation in fertility, phenology and composition of understorey Neotropical forests. *Biotropica* 19: 216–227.
- HALL, I. N. & SWAINE, M. D. 1976. Classification and ecology of closed-canopy forest in Ghana. *Journal of Ecology* 64: 913–951.
- IRWANTO. 2006. Penerapan AMDAL pada Pembangunan di Bidang Kehutanan. [http://saveforest.webs.com/amdal\\_kehutanan.html](http://saveforest.webs.com/amdal_kehutanan.html). Accessed 14 Maret 2014 (In Indonesian).
- ISMAL, A. R. 1975. *Analisa vegetasi tumbuhan*

- bawah di hutan Cagar Alam Sukawayana, Pelabuhan Ratu, Jawa Barat. [Skripsi], Departemen Biologi, Institut Teknologi Bandung, Bandung (In Indonesian).
- KARTAWINATA, K. 1977. Structure and composition of forests in some nature reserves in West Java, Indonesia. In: *Papers presented at the 13th Pacific Science Congress*. Vancouver, Canada. 1975. LIPI, Jakarta. Pp. 59–68.
- KARTAWINATA, K. 2005. Six decades of natural vegetation studies in Indonesia. In: SOEMODIHARDJO, S. & SASTRAPRADJA, S. D. (Eds.). *Six Decades of Science and Scientists in Indonesia*. Naturindo, Bogor. Pp. 95–140.
- KARTAWINATA, K. 2013. *Diversitas ekosistem alami Indonesia*. Yayasan Pustaka Obor Indonesia & LIPI Press, Jakarta (In Indonesian).
- KARTAWINATA, K., APANDI, A. & SUSELO, T. B. 1985. The forest of Peucang Island, Ujung Kulon National Park, West Java. In: SASTRAPRADJA, D. et al. (Eds.). *Proceedings of the Symposium on 100 Years Development of Krakatau and its Surrounding*. LIPI. 1: 448–452.
- KEHUTANAN. 2009. *Rencana Pengelolaan Tahunan Kawasan Cagar Alam Pulau Sempu*. Balai Besar Konservasi Sumber Daya Alam Jawa Timur. Surabaya. Unpublished (In Indonesian).
- KERN, J. H. 1974. Cyperaceae. *Flora Malesiana* I. 7: 435–753.
- KIEW, R. 1978. Floristic components of the ground flora of a tropical lowland rainforest at Gunung Mulu National Park, Sarawak. *Pertanika* 1: 112–119.
- KRAMADIBRATA, K., PRAWIROATMODJO, S., POLOSAKAN, R., WINDADRI, F. I., SADILI, A., SUJADI, A., ROSALINA, D. & SUMANTA, I. 2010. Kajian Ekosistem hutan dataran rendah Cagar Alam Pulau Sempu, Malang–Jawa Timur. *Laporan Perjalanan*. Pusat Penelitian Biologi LIPI. Bogor. Unpublished (In Indonesian).
- LMG (LEMBAGA METEOROLOGI DAN GEOFISIKA, 1967. Tjurah hujan rata2 di Djawa dan Madura (Mean rainfall in Java and Madura). *Meteorological Notes No. 8 Part I*. Lembaga Meteorologi dan Geofisika, Direktorat Djenderal Perhubungan Udara. Departemen Perhubungan. Jakarta.
- MEIJER, W. 1959. Plant sociological analysis of montane rain forest near Tjibodas, West Java. *Acta Bot. Neerl.* 8: 277–291.
- MOF (MINISTRY OF FORESTRY). 2009. *Annual Management Plan Sempu Island Nature Reserve area (Rencana Pengelolaan Tahunan Kawasan Cagar Alam Pulau Sempu)*. Center for Natural Resources Conservation in East Java. Surabaya.
- MUELLER-DOMBOIS, D. & ELLENBERG, H. 1974. *Aims and Methods of Vegetation Ecology*. John Wiley & Sons, Inc. New York. 547 p.
- NATH, P. C., ARUNACHALAM, A., KHAN, M. L., ARUNACHALAM, K. & BARBHUIYA, A. R. 2005. Vegetation analysis and tree population structure of tropical wet evergreen forests in and around Namdapha National Park, northeast India. *Biodiversity and Conservation* 14: 2109–2136.
- POLOSAKAN, R. 2011. Komposisi jenis dan struktur vegetasi di hutan pantai pada kawasan Cagar Alam Pulau Sempu Kabupaten Malang. In: WIDYATMOKO, D. et al. (Eds.). *Prosiding Seminar Nasional HUT Kebun Raya Cibodas Ke-159*. LIPI: 336–340 (In Indonesian).
- POLOSAKAN, R. 2014. Analisa vegetasi hutan pamah di kawasan Cagar Alam Pulau Sempu Kabupaten Malang. In: SUGIONO et al. (Eds.). *Prosiding Seminar Nasional Biodiversitas*. Universitas Negeri Solo: 212–216 (In Indonesian).
- POULSEN, A. D. 1996. Species richness and density of ground herbs within a plot of lowland rainforest in north-west Borneo. *Journal of Tropical Ecology* 12 (2): 177–190.
- PRAWIROATMODJO, S. 2013. Hutan mangrove Cagar Alam Pulau Sempu, Jawa Timur. *Jurnal Biologi Indonesia* 9: 121–139.
- RICHARDS, P. W. 1996. *The Tropical Rainforest*, Second Edition, Cambridge University Press, Cambridge.
- RISNA, R. A. 2009. *Autekologi dan studi populasi Myristica teijsmannii Miq. (Myristicaceae) di Cagar Alam Pulau Sempu Jawa Timur*. Institut Pertanian Bogor, Bogor. [MSc. Thesis].
- SADILI, A. 2010. Struktur dan komposisi jenis tumbuhan herba dan semai pada habitat satwa herbivor di Suaka Marga Satwa Cikepuh Sukabumi, Jawa Barat. *Berita Biologi* 10 (1): 51–58.
- SADILI, A., KARTAWINATA, K., KARTONEGORO, A., SOEDJITO, H. & SUMADIJAYA, A. 2009. Floristic composition and structure of subalpine summit habitats on Mt. Gede Pangrango Complex, Cibodas Biosphere Reserve, West Java. Indonesia. *Reinwardtia* 12(5): 391–404.
- SCHMIDT, F. H. & FERGUSON, J. H. 1951. *Rainfall types based on wet and dry period rations for Indonesia with Western New Guinea*. Verhandelingen Djawatan Meteorologi dan Geofisika, Djakarta.
- STEENIS, C. G. G. J. VAN & SCHIPPERS-LAMMERTSE, A. F. 1965. General part: Concise plant geography of Java. In: BACKER, C. A. & BAKHUIZEN VAN DEN BRINK, Jr., R. C. (Eds.). *Flora of Java*, vol. 2, N. V. P. Noordhoff, Groningen. Pp. 3–72.
- SUMARDJA, E. A. & KARTAWINATA, K. 1977. Vegetation analysis of the habitats of banteng (*Bos javanicus*) at the Pananjung-Pangandaran Nature Reserve, West Java. *BIOTROP Bulletin* 18: 3–43.
- TING, T. M. & POULSEN, A. D. 2009. Understorey vegetation at two mud volcanoes in North-East Borneo. *Journal of Tropical Forest Science* 21 (3): 198–209.
- TRAN, H., IIDA, S. & INOUE, S. 2005. Species composition, diversity and structure of secondary tropical forests following selective logging in Huong Son, Ha Tinh Province, Vietnam. *Jour. Fac. Agr.* 50 (2): 551–571.
- WALTER, H. 1973. *Vegetation of the earth*, (English edition). The English Universities Press, London.
- WHITMORE, T. C. 1986. *Tropical Rain Forest of the Far East*. Second Edition. ELBS/Oxford University Press, Oxford.



## ERRATUM

### REINWARDTIA Vol. 14 (2), 2015

1. Please change the existing epithet name in p. 307, LINE 2 on ABSTRACT after *Zingiber: engganoense*
2. Please change the existing epithet name in p. 307, LINE 2 on ABSTRAK after *Zingiber: engganoense*
3. Please change the existing epithet name in p. 307, COLUMN 2, LINE 4 on SPECIES DESCRIPTION after *Zingiber: engganoense*
4. Please change the existing epithet name in p. 307, COLUMN 2, LINE 9 on SPECIES DESCRIPTION after *Zingiber: engganoense*
5. Please change the existing epithet name in p. 308, COLUMN 2, LINE 3 on Phenology after *Zingiber: engganoense*
6. Please change the existing epithet name in p. 308, on Table 1 after *Zingiber: engganoense*
7. Please change the existing epithet name in p. 309, on Table 2 after *Zingiber: engganoense*
8. Please change the existing epithet name in p. 309, on Fig. 1. after *Zingiber: engganoense*
9. Please change the existing epithet name in p. 310, on Plate 1. after *Zingiber: engganoense*
10. Please change the existing word in p. 312, LINE 17 on *Alpinia macrocrista* with the following: KRB

## INSTRUCTION TO AUTHORS

**Scope.** *Reinwardtia* is a scientific irregular journal on plant taxonomy, plant ecology and ethnobotany published in December. Manuscript intended for a publication should be written in English.

**Titles.** Titles should be brief, informative and followed by author's name and mailing address in one-paragraphed.

**Abstract.** English abstract followed by Indonesian abstract of not more than 250 words. Keywords should be given below each abstract.

**Manuscript.** Manuscript is original paper and represent an article which has not been published in any other journal or proceedings. The manuscript of no more than 36 pages by using Times New Roman 11, MS Word for Windows of A4 with double spacing, submitted to the editor through <reinwardtia@mail.lipi.go.id>. New paragraph should be indented in by 5 characters. For the style of presentation, authors should follow the latest issue of *Reinwardtia* very closely. Author(s) should send the preferred running title of the article submitted. Every manuscript will be sent to two blind reviewers.

**Identification key.** Taxonomic identification key should be prepared using the aligned couplet type.

**Nomenclature.** Strict adherence to the International Code of Botanical Nomenclature is observed, so that taxonomic and nomenclatural novelties should be clearly shown. English description for new taxon proposed should be provided and the herbaria where the type specimens area deposited should be presented. Name of taxon in taxonomic treatment should be presented in the long form that is name of taxon, author's name, year of publication, abbreviated journal or book title, volume, number and page.

**Map/line drawing illustration/photograph.** Map, line drawing illustration, or photograph preferably should be prepared in landscape presentation to occupy two columns. Illustration must be submitted as original art accompanying, but separated from the manuscript. The illustration should be saved in JPG or GIF format at least 350 pixels. Legends or illustration must be submitted separately at the end of the manuscript.

**References.** Bibliography, list of literature cited or references follow the Harvard system as the following examples.

- Journal : KRAENZLIN, F. 1913. *Cyrtandraceae* novae Philippinenses I. *Philipp. J. Sci.* 8: 163–179.  
MAYER, V., MOLLER, M., PERRET, M. & WEBER, A. 2003. Phylogenetic position and generic differentiation of *Epithemateae* (*Gesneriaceae*) inferred from plastid DNA sequence data. *American J. Bot.* 90: 321–329.
- Proceedings : TEMU, S. T. 1995. Peranan tumbuhan dan ternak dalam upacara adat “Djoka Dju” pada suku Lio, Ende, Flores, Nusa Tenggara Timur. In: NASUTION, E. (Ed.). *Prosiding Seminar dan Lokakarya Nasional Etnobotani II*. LIPI & Perpustakaan Nasional: 263–268. (In Indonesian).  
SIMBOLON, H. & MIRMANTO, E. 2000. Checklist of plant species in the peat swamp forests of Central Kalimantan, Indonesia. In: IWAKUMA, T. *et al.* (Eds.) *Proceedings of the International Symposium on: Tropical Peatlands*. Pp.179-190.
- Book : RIDLEY, H. N. 1923. *Flora of the Malay Peninsula 2*. L. Reeve & Co. Ltd, London.
- Part of Book : BENTHAM, G. 1876. *Gesneriaceae*. In: BENTHAM, G. & HOOKER, J. D. *Genera plantarum 2*. Lovell Reeve & Co., London. Pp. 990–1025.
- Thesis : BAIRD, L. 2002. *A Grammar of Kéo: An Austronesian language of East Nusantara*. Australian National University, Canberra. [PhD. Thesis].
- Website : <http://www.nationaalherbarium.nl/fmcollectors/k/KostermansAJGH.htm>. Accessed 15 February 2012.





## **Reinwardtia**

Published by Herbarium Bogoriense, Botany Division, Research Center for Biology,  
Indonesian Institute of Sciences  
Address: Jln. Raya Jakarta-Bogor Km. 46 Cibinong 16911, P.O. Box 25 Cibinong  
Telp. (+ 62) 21 8765066; Fax (+62) 21 8765062  
E-mail: reinwardtia@mail.lipi.go.id

### ***REINWARDTIA* Author Agreement Form**

Title of article :

Name of Author(s) :

I/We hereby declare that:

- My/Our manuscript was based on my/our original work.
- It was not published or submitted to other journal for publication.
- I/we agree to publish my/our manuscript and the copyright of this article is owned by Reinwardtia.
- We have obtained written permission from copyright owners for any excerpts from copyrighted works that are included and have credited the sources in our article.

Author signature (s)

Date

---

Name



**REINWARDTIA**  
Vol. 15. No. 1. 2016  
CONTENTS  
Page

ASEP SADILI & KUSWATA KARTAWINATA. A study of the undergrowth vegetation of Sempu Island, East Java, Indonesia ...	1
RAHMAH, KUSWATA KARTAWINATA, NISYAWATI, WISNU WARDHANA & ERWIN NURDIN. Tree species diversity in the lowland forest of the core zone of the Bukit Duabelas National Park, Jambi, Indonesia .....	11
NASRIANTI SYAM, TATIK CHIKMAWATI & HIMMAH RUSTIAMI. A phenetic study of the <i>Calamus flabellatus</i> complex (Palmae) in West Malesia .....	27
DIAN LATIFAH, ROBERT A. CONGDON & JOSEPH A. HOLTUM. Regeneration strategies of Palms (Arecaceae) in response to cyclonic disturbances .....	43
I MADE ARDHAKA, WISNU HANDOYO ARDI, NI KADEK EROSI UNDAHARTA & I. G. TIRTA. A new species of <i>Begonia</i> (Begoniaceae) from Manusela National Park, Seram .....	61
ERRATUM REINWARDTIA Vol. 14(2), 2015 .....	65

Reinwardtia is a LIPI accredited Journal (792/AU3/P2MI-LIPI/04/2016)  
<http://e-journal.biologi.lipi.go.id/index.php/reinwardtia>

Herbarium Bogoriense  
Botany Division  
Research Center for Biology – Indonesian Institute of Sciences  
Cibinong Science Center  
Jln. Raya Jakarta – Bogor, Km 46  
Cibinong 16911, P.O. Box 25 Cibinong  
Indonesia