

# Pacific Journal of Mathematics

## **A SUB-ELLIPTIC ESTIMATE FOR A CLASS OF INVARIANTLY DEFINED ELLIPTIC SYSTEMS**

LESLEY MILLMAN SIBNER AND ROBERT JULES SIBNER

## A SUB-ELLIPTIC ESTIMATE FOR A CLASS OF INVARIANTLY DEFINED ELLIPTIC SYSTEMS

L. M. SIBNER AND R. J. SIBNER

We consider a certain invariantly defined nonlinear system of partial differential equations on a Riemannian manifold. Since a special case describes a steady, irrotational, compressible flow on the manifold, it is natural to refer to the (square of) the pointwise norm of the solution as the speed of the flow and to the density of the flow. Under appropriate restrictions on the density, the system is elliptic and we obtain a sub-elliptic estimate and a maximum principle for the speed of the flow in terms of the curvature of the manifold.

**Introduction.** Let  $M$  be an  $n$ -dimensional Riemannian manifold, and  $A^p(M)$  the space of smooth  $p$ -forms on  $M$ . For  $\omega \in A^p(M)$ ,  $x \in M$ , let  $Q(\omega) = (\omega, \omega)(x) = *(\omega \wedge *\omega)(x)$  denote the pointwise norm of the form  $\omega$ . Let  $\rho: C^\infty(M) \rightarrow R$  be a given bounded smooth strictly positive function which we call the density function.

In the following, we consider the invariantly defined nonlinear system of equations for  $\omega \in A^p(M)$ :

$$(1) \quad \begin{aligned} d\omega &= 0 \\ \delta(\rho(Q(\omega))\omega) &= 0. \end{aligned}$$

If  $p=1$ , this system describes the motion of a compressible fluid on  $M$  and reduces to a single second order equation for the potential function. If the metric is Euclidean and  $\rho(Q) = (1 - (\gamma - 1)/2Q)^{1/\gamma-1}$ , it becomes the gas dynamics equation for polytropic flow in  $R^n$ . If  $\rho \equiv 1$ , one obtains the Laplace-Beltrami equation.

To be more explicit, if  $\omega$  is a solution of (1), then it is also a solution of a homogeneous second order quasi-linear system,  $A\omega = 0$ . In local coordinates, let

$$\omega = \omega_{i_1 \dots i_p} dx^{i_1} \wedge \dots \wedge dx^{i_p}, \quad \omega^{j_1 \dots j_p} = g^{j_1 i_1} \dots g^{j_p i_p} \omega_{i_1 \dots i_p},$$

and assume  $\rho = \rho(Q(\omega))$ . Then,  $A: A^p(M) \rightarrow A^p(M)$  is given by

$$\begin{aligned} (A\omega)_{i_1 \dots i_p} = & \sum_{i,j} g^{ji} \left\{ \rho \frac{\partial^2 \omega_{i_1 \dots i_p}}{\partial x^i \partial x^j} \right. \\ & \left. + 2\rho' \omega^{j_1 \dots j_p} \sum_{k=1}^p \omega_{i_1 \dots i_{k-1} i i_{k+1} \dots i_p} \frac{\partial^2 \omega_{j_1 \dots j_p}}{\partial x^{i_k} \partial s^j} \right\} \\ & + \text{lower order terms.} \end{aligned}$$

(We will observe the usual summation convention wherever possible.)

A computation shows that the system (1) is *elliptic* for  $\omega \in A^p(M)$  if and only if

$$(2) \quad \frac{d}{dQ}(\rho^2 Q) > 0 \quad \text{for all } x \in M.$$

If  $M$  is compact and (1) is elliptic, there is a unique *weak* solution in each cohomology class (Sibner [7]). The question of smooth solutions is unresolved except in the case  $p = 1$  ([7]) or if the metric is Euclidean (Uhlenbeck [10]).

Assuming that the system (1) is elliptic, we shall derive an inequality for the function  $Q$ , of the following kind:

$$(3) \quad LQ + B(\omega) \geq 0$$

where  $\omega$  is a solution of (1),  $L$  is a single second order elliptic operator with no zero order term, and  $B$  is a quadratic form whose sign depends upon the curvature tensor. Such an inequality leads to a maximum principle and is perhaps a step in the direction of elliptic regularity for the system (1) (see [10]). An inequality of the form (3) and a maximum principle were previously proved by the authors for 1-forms on surfaces ( $n = 2$ ) (see [9]).

1. The inequality satisfied by  $Q$ . Let  $I = i_1 \cdots i_p, J = j_1 \cdots j_p$  be multi-indices and set  $g^{IJ} = g^{i_1 j_1} \cdots g^{i_p j_p}$ . Then,  $Q(\omega) = g^{JJ} \omega_J \omega_I$  in this notation. In terms of the Riemannian metric, let  $|\nabla \omega|^2 = g^{kl} g^{JJ} \nabla_k \omega_J \nabla_l \omega_I$ . Define the curvature form (Lichnerowicz [5, 6])

$$K(\omega) = g^{JJ} \omega_J \left\{ \sum_{s=1}^p K_{i_s}^t \omega_{i_1 \cdots t \cdots i_p} + \sum_{\substack{1 \cdots p \\ t < s}} K_{i_s i_t}^{vu} \omega_{i_1 \cdots v \cdots u \cdots i_p} \right\}.$$

Computing the Laplacian of  $Q$ , one obtains

$$-\frac{1}{2} \Delta Q = \frac{1}{2} g^{jj} \nabla_j \nabla_i Q = |\nabla \omega|^2 + (g^{JJ} \omega_J)(g^{ji} \nabla_j \nabla_i \omega_I)$$

and using the formula for  $\Delta \omega$  ([11]) one obtains the fundamental identity for  $Q$ :

$$(1.1) \quad -\frac{1}{2} \Delta Q = |\nabla \omega|^2 + K(\omega) - p!(\omega, \omega).$$

Next, let  $\omega$  be a solution of (1) and replace  $\omega$  in (1.1) by  $\rho \omega$ . Using the fact that  $\delta \rho \omega = 0$ , one obtains

$$(1.2) \quad -\frac{1}{2} \Delta(\rho^2 Q) + p!(\delta d \rho \omega, \rho \omega) = \rho^2 K(\omega) + |\nabla(\rho \omega)|^2.$$

We shall show that the left hand side is a second order differ-

ential operator on  $Q$ ,  $LQ = L_1Q + L_2Q$ , and will compute its principal part.

$$L_1Q = -\frac{1}{2} \Delta(\rho^2Q) = \frac{1}{2}(\rho^2 + 2\rho\rho'Q)g^{ji} \frac{\partial^2 Q}{\partial x^i \partial x^j} + \text{first order terms}.$$

Using the fact that  $d\omega = 0$ ,

$$p!(\delta d\rho\omega)_{i_1 \dots i_p} = -g^{ji} \nabla_j \left( \rho' \omega_{i_1 \dots i_p} \frac{\partial Q}{\partial x^i} - \rho' \sum_{k=1}^p \omega_{i_1 \dots i_{k-1} i_{k+1} \dots i_p} \frac{\partial Q}{\partial x^{ik}} \right).$$

Therefore,

$$\begin{aligned} L_2Q &= p!(\delta d\rho\omega, \rho\omega) \\ &= -g^{ji} \rho\rho' \omega_{i_1 \dots i_p} \omega^{i_1 \dots i_p} \frac{\partial^2 Q}{\partial x^i \partial x^j} + g^{ji} \rho\rho' \omega^{i_1 \dots i_p} \sum_{k=1}^p \omega_{i_1 \dots i_{k-1} i_{k+1} \dots i_p} \frac{\partial^2 Q}{\partial x^{ik} \partial x^j} \\ &\quad + \text{first order terms}. \end{aligned}$$

Combining coefficients of  $\partial^2 Q / \partial x^i \partial x^j$ ,

$$L_2Q = \left\{ -g^{ji} \rho\rho'Q + \rho\rho' \omega_{i_1 \dots i_p} \sum_{k=1}^p g^{ji_k i_{k+1} \dots i_p} \right\} \frac{\partial^2 Q}{\partial x^i \partial x^j} + \dots$$

and

$$LQ = \left\{ \frac{1}{2} g^{ji} \rho^2 + \rho\rho' \omega_{i_1 \dots i_p} \sum_{k=1}^p g^{ji_k i_{k+1} \dots i_p} \right\} \frac{\partial^2 Q}{\partial x^i \partial x^j} + \dots.$$

**PROPOSITION 1.1.** *If the system (1) is elliptic at a solution  $\omega$ , then  $L$  is an elliptic operator.*

*Proof.* The principal symbol of  $L_1$  on a cotangent vector  $\pi = (\pi_1, \dots, \pi_n) \neq 0$  is

$$\frac{1}{2}(\rho^2 + 2\rho\rho'Q)g^{ji} \pi_i \pi_j > 0$$

using the ellipticity condition (2). Therefore,  $L_1$  is elliptic.

Choose geodesic normal coordinates at a point, in which case  $g^{ij} = \delta^{ij}$ ,  $\omega^i = \omega_i$ , and  $Q(\omega) = \sum_I (\omega_i)^2$ . The principal symbol of  $L_2$  in these coordinates becomes:

$$\begin{aligned} & -\rho\rho' \left( Q\delta^{ij} - \sum_{k=1}^p \omega_{i_1 \dots i_{k-1} j i_{k+1} \dots i_p} \omega_{i_1 \dots i_{k-1} i i_{k+1} \dots i_p} \right) \pi_i \pi_j \\ & = -\rho\rho' \left\{ (\sum (\omega_{i_1 \dots i_p})^2) (\sum \pi_i^2) - \left( \sum_{k=1}^p \omega_{i_1 \dots i_{k-1} i i_{k+1} \dots i_p} \pi_i \right)^2 \right\}. \end{aligned}$$

If  $\rho' \leq 0$ , one sees immediately from the Schwartz inequality that this expression is nonnegative. Therefore, if  $\rho' \leq 0$ , the principal symbol of  $L = L_1 + L_2$  is positive definite and  $L$  is elliptic.

If  $\rho' \geq 0$ , the principal symbol of  $L$  in geodesic normal coordinates is

$$\frac{1}{2}\rho^2 \sum \pi_i^2 + \rho\rho'(\sum \omega_{i_1 \dots i_{k-1} i_{k+1} \dots i_p} \pi_i)^2 > 0$$

using the fact that  $\rho$  is positive. Hence,  $L$  is elliptic.

Summarizing the results of this section, we have shown

**PROPOSITION 1.2.** *Let  $\omega \in A^p(M)$  be a solution of the homogeneous elliptic system (1). Then  $Q(\omega)$  is a solution of the single scalar equation*

$$(1.3) \quad LQ = \rho^2 K(\omega) + |\nabla(\rho\omega)|^2$$

where  $L$  is a second order elliptic operator having no zero order terms.

**2. The maximum principle.** Our main result is the following.

**THEOREM.** *Let  $\omega$  be a solution of the elliptic system (1). Then,*

(a)  $Q = (\omega, \omega)$  cannot have a relative maximum at a point  $x_0$  where  $K(\omega)|_{x_0} > 0$ .

(b) If  $Q$  has a relative maximum at a point  $x_0$ , in a neighborhood  $N$  of which,  $K(\omega) \geq 0$  then

(i)  $Q$  is constant on  $N$ .

(ii)  $K(\omega) \equiv 0$  in  $N$ .

(iii)  $\nabla\omega \equiv 0$  on  $N$ .

*Proof.* As in [9], statements (a) and (i) of (b) follow from Proposition 1.2 and the Hopf Maximum Principle. But if  $Q$  is constant in  $N$ , then  $LQ \equiv 0$  in  $N$  which gives, again by Proposition 1.2, statements (ii) and (iii).

If  $p = 1$ , the curvature expression  $K(\omega)$  reduces to  $R^{ij}\omega_i\omega_j$  where  $R^{ij}$  is the Ricci curvature tensor. One speaks of  $R^{ij}\omega_i\omega_j$  as the Ricci curvature in the direction  $\omega$ . In the language of gas dynamics we have the

**COROLLARY.** *A subsonic compressible flow on  $M$  cannot assume its maximum speed at a point where the Ricci curvature is positive in the direction of the flow. If the maximum speed is attained at a point of a region  $N$  in which the Ricci curvature in the flow direction is nonnegative, then the curvature must in fact be zero, the speed  $Q$  must be constant, and the flow parallel in  $N$  (i.e., the covariant derivatives,  $\nabla_i\omega_j \equiv 0$ ). If it is further known that*

the Ricci curvature is positive definite, then  $Q \equiv 0$  and hence  $\omega \equiv 0$  in  $N$ .

*Added in proof.* The authors, with P. D. Smith, have obtained a regularity theorem for the system (1) using the estimate (3). It will appear in a forthcoming paper.

#### REFERENCES

1. L. Bers, *Mathematical Aspects of Subsonic and Transonic Gas Dynamics*, Wiley, New York, 1958.
2. S. Bochner, *Vector fields and Ricci curvature*, Bull. Amer. Math. Soc., **52** (1946), 776-797.
3. ———, *Curvature and Betti numbers*, Annals of Math., **49** (1948), 379-390.
4. E. Hopf, *Elementare Bemerkungen über die Lösungen partieller Differentialgleichungen zweiter Ordnung vom elliptischen Typus*, Sitzber. Preuss. Akad. Wiss. Physik-math. Kl, **19** (1927), 147-152.
5. A. Lichnerowicz, *Courbure et nombres de Betti d'une variété riemannienne compacte*, Compt. Rend. Acad. Sci. Paris, **226** (1948), 1678-1680.
6. ———, *Courbure, nombres de Betti espaces symmetriques*, Proc. Intern. Congr. of Math., **2** (1952), 216-223.
7. L. M. Sibner and R. J. Sibner, *A nonlinear Hodge-de Rham theorem*, Acta Math., **125** (1970), 57-73.
8. ———, *Nonlinear Hodge theory: Applications*, Advances in Math., **31** (1979), 1-16.
9. ———, *A maximum principle for compressible flow*, Proc. Amer. Math. Soc., **71** (1978), 103-108.
10. K. Uhlenbeck, *Regularity for a class of nonlinear elliptic systems*, Acta Math., **138** (1977), 219-240.
11. K. Yano, *Integral formulas in Riemannian Geometry*, Marcel Dekker, New York, 1970.

Received May 8, 1980. Research of the first author was partially supported by NSF Grant MCS78-03276 and research of the second author was partially supported by NSF Grant MCS78-03268.

POLYTECHNIC INSTITUTE OF NEW YORK

BROOKLYN, NY 11201

AND

CITY UNIVERSITY OF NEW YORK

BROOKLIN COLLEGE

BROOKLIN, NY 11210



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California  
Los Angeles, CA 90024

HUGO ROSSI

University of Utah  
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University  
Stanford, CA 94305

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$102.00 a year (6 Vols., 12 issues). Special rate: \$51.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1981 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan



Thomas E. Armstrong and William David Sudderth, Nearly strategic measures .....	251
John J. Buoni, Artatrana Dash and Bhushan L. Wadhwa, Joint Browder spectrum .....	259
Jack Paul Diamond, Hypergeometric series with a $p$ -adic variable .....	265
Raymond Frank Dickman, Jack Ray Porter and Leonard Rubin, Completely regular absolutes and projective objects .....	277
James Kenneth Finch, On the local spectrum and the adjoint .....	297
Benno Fuchssteiner, An abstract disintegration theorem .....	303
Leon Gerber, The volume cut off a simplex by a half-space .....	311
Irving Leonard Glicksberg, An application of Wermer's subharmonicity theorem .....	315
William Goldman, Two examples of affine manifolds .....	327
Yukio Hirashita, On the Weierstrass points on open Riemann surfaces .....	331
Darrell Conley Kent, A note on regular Cauchy spaces .....	333
Abel Klein and Lawrence J. Landau, Periodic Gaussian Osterwalder-Schrader positive processes and the two-sided Markov property on the circle .....	341
Brenda MacGibbon, $\mathcal{H}$ -Borelian embeddings and images of Hausdorff spaces .....	369
John R. Myers, Homology 3-spheres which admit no PL involutions .....	379
Boon-Hua Ong, Invariant subspace lattices for a class of operators .....	385
Chull Park, Representations of Gaussian processes by Wiener processes .....	407
Lesley Millman Sibner and Robert Jules Sibner, A sub-elliptic estimate for a class of invariantly defined elliptic systems .....	417
Justin R. Smith, Complements of codimension-two submanifolds. III. Cobordism theory .....	423
William Albert Roderick Weiss, Small Dowker spaces .....	485
David J. Winter, Cartan subalgebras of a Lie algebra and its ideals. II .....	493