

 Open access • Journal Article • DOI:10.1088/0004-637X/728/2/120

A SUB-SATURN MASS PLANET, MOA-2009-BLG-319Lb — [Source link](#)

[N. Miyake](#), [Takahiro Sumi](#), [Subo Dong](#), [Subo Dong](#) ...+127 more authors

Institutions: [Nagoya University](#), [Ohio State University](#), [Princeton University](#), [Las Cumbres Observatory Global Telescope Network](#) ...+39 more institutions

Published on: 20 Feb 2011 - [The Astrophysical Journal](#) (American Astronomical Society)

Topics: [Gravitational microlensing](#), [Einstein radius](#), [Planetary system](#), [Light curve](#) and [Planetary mass](#)

Related papers:

- [A COLD NEPTUNE-MASS PLANET OGLE-2007-BLG-368Lb: COLD NEPTUNES ARE COMMON](#)
- [Discovery of a Jupiter/Saturn Analog with Gravitational Microlensing](#)
- [Microlensing event moa-2007-blg-400: exhuming the buried signature of a cool, jovian-mass planet](#)
- [Sub-Saturn Planet MOA-2008-BLG-310Lb: Likely To Be In The Galactic Bulge](#)
- [A Low-Mass Planet with a Possible Sub-Stellar-Mass Host in Microlensing Event MOA-2007-BLG-192](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/a-sub-saturn-mass-planet-moa-2009-blg-319lb-hhtr7kzp0x>

A SUB-SATURN MASS PLANET, MOA-2009-BLG-319Lb

N. MIYAKE^{1,69}, T. SUMI^{1,69}, SUBO DONG^{2,70,71}, R. STREET^{3,4,72}, L. MANCINI^{5,6,7,73}, A. GOULD^{8,71}, D. P. BENNETT^{9,69,74},
Y. TSAPRAS^{3,10,72}, J. C. YEE^{8,71}, M. D. ALBROW^{11,74}, I. A. BOND^{12,69}, P. FOUQUE^{13,74}, P. BROWNE^{14,72,73}, C. HAN^{15,71},
C. SNODGRASS^{16,17,72,73}, F. FINET^{18,73}, K. FURUSAWA^{1,69}, K. HARPSØE^{19,73}, W. ALLEN^{20,71}, M. HUNDERTMARK^{21,73},
M. FREEMAN^{22,69}, D. SUZUKI^{1,69},

AND

F. ABE¹, C. S. BOTZLER²², D. DOUCHIN²², A. FUKUI¹, F. HAYASHI¹, J. B. HEARNshaw²³, S. HOSAKA¹, Y. ITOW¹, K. KAMIYA¹,
P. M. KILMARTIN²⁴, A. KORPELA²⁵, W. LIN¹², C. H. LING¹², S. MAKITA¹, K. MASUDA¹, Y. MATSUBARA¹, Y. MURAKI²⁶,
T. NAGAYAMA²⁷, K. NISHIMOTO¹, K. OHNISHI²⁸, Y. C. PERROTT²², N. RATTENBURY²², TO. SAITO²⁹, L. SKULJAN¹²,
D. J. SULLIVAN²⁵, W. L. SWEATMAN¹², P. J. TRISTRAM²⁴, K. WADA²⁶, P. C. M. YOCK²²

(THE MOA COLLABORATION)⁶⁹

G. BOLT³⁰, M. BOS³¹, G. W. CHRISTIE³², D. L. DEPOY³³, J. DRUMMOND³⁴, A. GAL-YAM³⁵, B. S. GAUDI⁸, E. GORBIKOV³⁶,
D. HIGGINS³⁷, K.-H. HWANG¹⁵, J. JANCZAK⁸, S. KASPI^{36,38}, C.-U. LEE³⁹, J.-R. KOO⁴⁰, S. KOZŁOWSKI⁸, Y. LEE⁴¹, F. MALLIA⁴²,
A. MAURY⁴², D. MAOZ³⁶, J. MCCORMICK⁴³, L. A. G. MONARD⁴⁴, D. MOORHOUSE⁴⁵, J. A. MUÑOZ⁴⁶, T. NATUSCH⁴⁷,
E. O. OFEK^{48,75}, R. W. POGGE⁸, D. POLISHOOK³⁶, R. SANTALLO⁴⁹, A. SHPORER³⁶, O. SPECTOR³⁶, AND G. THORNLEY⁴⁵

(THE μ FUN COLLABORATION)⁷¹

A. ALLAN⁵⁰, D. M. BRAMICH^{51,74}, K. HORNE^{14,74}, N. KAINS¹⁴, I. STEELE⁵²

(THE ROBO NET COLLABORATION)⁷²

V. BOZZA^{5,6,53}, M. J. BURGDORF^{54,55}, S. CALCHI NOVATI^{5,6}, M. DOMINIK^{14,72,74,76}, S. DREIZLER²¹, M. GLITRUP⁵⁶,
F. V. HESSMAN²¹, T. C. HINSE^{19,57}, U. G. JØRGENSEN^{19,58}, C. LIEBIG^{14,59}, G. MAIER⁵⁹, M. MATHIASSEN¹⁹, S. RAHVAR^{60,61},
D. RICCI¹⁸, G. SCARPETTA^{5,6,53}, J. SKOTTFELT¹⁹, J. SOUTHWORTH⁶², J. SURDEJ¹⁸, J. WAMBSGANSS⁵⁹, F. ZIMMER⁵⁹

(THE MiNDSTEP CONSORTIUM)⁷³

V. BATISTA^{63,64}, J. P. BEAULIEU^{63,64,65}, S. BRILLANT⁶⁶, A. CASSAN^{63,64}, A. COLE⁶⁷, E. CORRALES^{63,64}, CH. COUTURES^{63,64},
S. DIETERS^{63,64,67}, J. GREENHILL⁶⁷, D. KUBAS^{63,64,66}, J. MENZIES⁶⁸

(THE PLANET COLLABORATION)⁷⁴

¹ Solar-Terrestrial Environment Laboratory, Nagoya University, Nagoya 464-8601, Japan; nmiyake@stelab.nagoya-u.ac.jp, sumi@stelab.nagoya-u.ac.jp, furusawa@stelab.nagoya-u.ac.jp, dsuke@stelab.nagoya-u.ac.jp, abe@stelab.nagoya-u.ac.jp, afukui@stelab.nagoya-u.ac.jp, fhayashi@stelab.nagoya-u.ac.jp, hosaka@stelab.nagoya-u.ac.jp, itow@stelab.nagoya-u.ac.jp, kkamiya@stelab.nagoya-u.ac.jp, makita@stelab.nagoya-u.ac.jp, kmasuda@stelab.nagoya-u.ac.jp, ymatsu@stelab.nagoya-u.ac.jp, nishimo@stelab.nagoya-u.ac.jp

² Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA; dong@ias.edu

³ Las Cumbres Observatory Global Telescope Network, 6740 Cortona Dr., Suite 102, Goleta, CA 93117, USA

⁴ Department of Physics, Broida Hall, University of California, Santa Barbara, CA 93106-9530, USA

⁵ Dipartimento di Fisica "E. R. Caianiello," Università degli Studi di Salerno, Via Ponte Don Melillo, 84085 Fisciano (SA), Italy

⁶ Istituto Internazionale per gli Alti Studi Scientifici (IIASS), Via G. Pellegrino 19, 84019 Vietri sul Mare (SA), Italy

⁷ Dipartimento di Ingegneria, Università del Sannio, Corso Garibaldi 107, 82100 Benevento, Italy

⁸ Department of Astronomy, Ohio State University, 140 W. 18th Ave., Columbus, OH 43210, USA; gould@astronomy.ohio-state.edu, gaudi@astronomy.ohio-state.edu, jyee@astronomy.ohio-state.edu, pogge@astronomy.ohio-state.edu, simkoz@astronomy.ohio-state.edu

⁹ Department of Physics, University of Notre Dame, Notre Dame, IN 46556, USA; bennett@nd.edu

¹⁰ Astronomy Unit, School of Mathematical Sciences, Queen Mary, University of London, London E1 4NS, UK

¹¹ Department of Physics and Astronomy, University of Canterbury, Private Bag 4800, Christchurch, New Zealand

¹² Institute of Information and Mathematical Sciences, Massey University, Private Bag 102-904, North Shore Mail Centre, Auckland, New Zealand; i.a.bond@massey.ac.nz

¹³ LATT, Université de Toulouse, CNRS, 14 Avenue Edouard Belin, 31400 Toulouse, France

¹⁴ SUPA, University of St. Andrews, School of Physics & Astronomy, North Haugh, St. Andrews, KY16 9SS, UK

¹⁵ Department of Physics, Institute for Basic Science Research, Chungbuk National University, Chongju 361-763, Republic of Korea; cheongho@astroph.chungbuk.ac.kr

¹⁶ European Southern Observatory, Alonso de Cordova 3107, Casilla 19001, Santiago 19, Chile

¹⁷ Max Planck Institute for Solar System Research, Max-Planck-Str. 2, 37191 Katlenburg-Lindau, Germany

¹⁸ Institut d'Astrophysique et de Géophysique, Allée du 6 Août 17, Sart Tilman, Bât. B5c, 4000 Liège, Belgium

¹⁹ Niels Bohr Institutet, Københavns Universitet, Juliane Maries Vej 30, 2100 København Ø, Denmark

²⁰ Vintage Lane Observatory, Blenheim, New Zealand; whallen@xtra.co.nz

²¹ Institut für Astrophysik, Georg-August-Universität, Friedrich-Hund-Platz 1, 37077 Göttingen, Germany

²² Department of Physics, University of Auckland, Auckland, New Zealand

²³ Department of Physics and Astronomy, University of Canterbury, Private Bag 4800, Christchurch, New Zealand

²⁴ Mt. John University Observatory, University of Canterbury, P.O. Box 56, Lake Tekapo 8770, New Zealand

²⁵ School of Chemical and Physical Sciences, Victoria University, Wellington, New Zealand

²⁶ Department of Physics, Konan University, Kobe, Nishiokamoto 8-9-1 Japan

²⁷ Department of Physics and Astrophysics, Faculty of Science, Nagoya University, Nagoya 464-8602, Japan

²⁸ Nagano National College of Technology, Nagano 381-8550, Japan

²⁹ Tokyo Metropolitan College of Aeronautics, Tokyo 116-8523, Japan

³⁰ Craigie Observatory, Perth, Australia; gbolt@iinet.net.au

³¹ Molehill Astronomical Observatory, Auckland, New Zealand; molehill@ihug.co.nz

³² Auckland Observatory, Auckland, New Zealand; gwchristie@christie.org.nz

³³ Department of Physics and Astronomy, Texas A&M University, College Station, TX, USA; depay@physics.tamu.edu

³⁴ Possum Observatory, Patutahi, New Zealand; john_drummond@xtra.co.nz

- ³⁵ Department of Particle Physics and Astrophysics, Weizmann Institute of Science, 76100 Rehovot, Israel; avishay.gal-yam@weizmann.ac.il
- ³⁶ School of Physics and Astronomy and Wise Observatory, Tel-Aviv University, Tel-Aviv 69978, Israel; shai@wise.tau.ac.il, dani@wise.tau.ac.il, david@wise.tau.ac.il, shporer@wise.tau.ac.il, odedspec@wise.tau.ac.il, evgenyg@wise.tau.ac.il
- ³⁷ Hunters Hill Observatory, Canberra, Australia; dhi67540@bigpond.net.au
- ³⁸ Department of Physics, Technion, Haifa 32000, Israel
- ³⁹ Korea Astronomy and Space Science Institute, Daejeon, Republic of Korea
- ⁴⁰ Department of Astronomy and Space Science, Chungnam National University, Daejeon, and Korea Astronomy and Space Science Institute, Daejeon, Republic of Korea
- ⁴¹ Department of Astronomy and Space Science, Chungnam National University, Republic of Korea
- ⁴² Campo Catino Austral Observatory, San Pedro de Atacama, Chile; francomallia@campocatinobservatory.org, alain@spaceobs.com
- ⁴³ Farm Cove Observatory, Centre for Backyard Astrophysics, Pakuranga, Auckland, New Zealand; farmcoveobs@xtra.co.nz
- ⁴⁴ Bronberg Observatory, Centre for Backyard Astrophysics, Pretoria, South Africa; lagmonar@nmisa.org
- ⁴⁵ Kumeu Observatory, Kumeu, New Zealand; acrux@orcon.net.nz, guy.thornley@gmail.com
- ⁴⁶ Departamento de Astronomía y Astrofísica, Universidad de Valencia, E-46100 Burjassot, Valencia, Spain
- ⁴⁷ AUT University, Auckland, New Zealand; tim.natusch@aut.ac.nz
- ⁴⁸ Division of Physics, Mathematics and Astronomy, California Institute of Technology, Pasadena, CA 91125, USA
- ⁴⁹ Southern Stars Observatory, Faaa, Tahiti, French Polynesia; obs930@southernstars-observatory.org
- ⁵⁰ School of Physics, University of Exeter, Stocker Road, Exeter EX4 4QL, UK
- ⁵¹ European Southern Observatory, Karl-Schwarzschild-Straße 2, 85748 Garching bei München, Germany
- ⁵² Astrophysics Research Institute, Liverpool John Moores University, Liverpool CH41 1LD, UK
- ⁵³ INFN, Gruppo Collegato di Salerno, Sezione di Napoli, Italy
- ⁵⁴ Deutsches SOFIA Institut, Universität Stuttgart, Pfaffenwaldring 31, 70569 Stuttgart, Germany
- ⁵⁵ SOFIA Science Center, NASA Ames Research Center, Mail Stop N211-3, Moffett Field, CA 94035, USA
- ⁵⁶ Department of Physics & Astronomy, Aarhus University, Ny Munkegade 120, 8000 Århus C, Denmark
- ⁵⁷ Armagh Observatory, College Hill, Armagh, BT61 9DG, UK
- ⁵⁸ Centre for Star and Planet Formation, Københavns Universitet, Øster Voldgade 5-7, 1350 København Ø, Denmark
- ⁵⁹ Astronomisches Rechen-Institut, Zentrum für Astronomie der Universität Heidelberg (ZAH), Mönchhofstr. 12-14, 69120 Heidelberg, Germany
- ⁶⁰ Department of Physics, Sharif University of Technology, P.O. Box 11365-9161, Tehran, Iran
- ⁶¹ School of Astronomy, IPM (Institute for Studies in Theoretical Physics and Mathematics), P.O. Box 19395-5531, Tehran, Iran
- ⁶² Astrophysics Group, Keele University, Staffordshire, ST5 5BG, UK
- ⁶³ Institut d'Astrophysique de Paris, UPMC Univ Paris 06, UMR7095, F-75014, Paris, France
- ⁶⁴ CNRS, UMR7095, Institut d'Astrophysique de Paris, F-75014 Paris, France
- ⁶⁵ Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK
- ⁶⁶ European Southern Observatory, Casilla 19001, Vitacura 19, Santiago, Chile
- ⁶⁷ School of Math and Physics, University of Tasmania, Private Bag 37, GPO Hobart, Tasmania 7001, Australia
- ⁶⁸ South African Astronomical Observatory, P.O. Box 9 Observatory 7925, South Africa

Received 2010 October 8; accepted 2010 December 8; published 2011 January 28

ABSTRACT

We report the gravitational microlensing discovery of a sub-Saturn mass planet, MOA-2009-BLG-319Lb, orbiting a K- or M-dwarf star in the inner Galactic disk or Galactic bulge. The high-cadence observations of the MOA-II survey discovered this microlensing event and enabled its identification as a high-magnification event approximately 24 hr prior to peak magnification. As a result, the planetary signal at the peak of this light curve was observed by 20 different telescopes, which is the largest number of telescopes to contribute to a planetary discovery to date. The microlensing model for this event indicates a planet–star mass ratio of $q = (3.95 \pm 0.02) \times 10^{-4}$ and a separation of $d = 0.97537 \pm 0.00007$ in units of the Einstein radius. A Bayesian analysis based on the measured Einstein radius crossing time, t_E , and angular Einstein radius, θ_E , along with a standard Galactic model indicates a host star mass of $M_L = 0.38^{+0.34}_{-0.18} M_\odot$ and a planet mass of $M_p = 50^{+44}_{-24} M_\oplus$, which is half the mass of Saturn. This analysis also yields a planet–star three-dimensional separation of $a = 2.4^{+1.2}_{-0.6}$ AU and a distance to the planetary system of $D_L = 6.1^{+1.1}_{-1.2}$ kpc. This separation is ~ 2 times the distance of the snow line, a separation similar to most of the other planets discovered by microlensing.

Key words: gravitational lensing: micro – planetary systems

Online-only material: color figure

1. INTRODUCTION

We present the eleventh microlensing planet, following 10 previous discoveries (Bond et al. 2004; Udalski et al. 2005; Beaulieu et al. 2006; Gould et al. 2006; Gaudi et al. 2008; Bennett et al. 2008; Dong et al. 2009; Sumi et al. 2010; Janczak et al. 2010). Microlensing is unique among exoplanet detection methods in that it is sensitive to planets with masses down to $1 M_\oplus$ (Bennett & Rhie 1996) at relatively large separations, typically between 1 AU and 6 AU, depending on the mass of the host star. These separations are generally beyond the “snow line” at ~ 2.7 AU M/M_\odot (Ida & Lin 2004; Lecar et al.

⁶⁹ Microlensing Observations in Astrophysics (MOA) Collaboration, <http://www.phys.canterbury.ac.nz/moa>

⁷⁰ Sagan Fellow.

⁷¹ Microlensing Follow Up Network (μ FUN), <http://www.astronomy.ohio-state.edu/~microfun>

⁷² RoboNet, <http://robonet.lco.gt.net>

⁷³ Microlensing Network for the Detection of Small Terrestrial Exoplanets (MiNDSTeP), <http://www.mindstep-science.org>

⁷⁴ Probing Lensing Anomalies Network (PLANET), <http://planet.iap.fr>

⁷⁵ Einstein Fellow.

⁷⁶ Royal Society University Research Fellow.

2006; Kennedy & Kenyon 2008), the region where planets can form most quickly, according to the core accretion theory. Microlensing confirms this expectation, as a statistical analysis of the prevalence of planets found by microlensing shows that Saturn mass planets beyond the snow line are more common than the higher mass gas giants found by radial velocities in shorter period orbits (Gould et al. 2010), although the microlensing results are consistent with an extrapolation of the radial velocity results for solar-mass stars to larger orbital distances (Cumming et al. 2008). Furthermore, Sumi et al. (2010) have shown that the number of planets (per logarithmic interval) increases with decreasing mass ratio, q , as $q^{-0.7 \pm 0.2}$, down to $\sim 10 M_{\oplus}$. So, cold Neptunes seem to be even more common than cold Saturns. While the number of planets found by microlensing is relatively small, it is the cold Neptunes and Saturns discovered by microlensing that represent the most common types of exoplanet yet to be discovered. Microlensing has also found the first Jupiter/Saturn analog planetary system (Gaudi et al. 2008; Bennett et al. 2010), and it should soon be possible to use the microlensing results to determine how the properties of exoplanet systems vary with distance from the Galactic center.

Searches for exoplanets via the microlensing method are currently conducted by two survey groups, the Microlensing Observations in Astrophysics (MOA; Bond et al. 2001; Sumi et al. 2003) and the Optical Gravitational Lensing Experiment (OGLE; Udalski 2003), which monitor $\sim 40 \text{ deg}^2$ of the Galactic bulge to identify stellar microlensing events that can be searched for planetary signals. The planetary signals have durations that range from a few hours to a few days, so a global network of telescopes is needed to search for and characterize planetary signals. The follow-up groups that complete this telescope network are the Microlensing Follow-Up Network (μFUN), RoboNet, Microlensing Network for the Detection of Small Terrestrial Exoplanets (MiNDSTeP), and the Probing Lensing Anomalies NETwork (PLANET). These narrow field-of-view (FOV) follow-up telescopes can provide very high cadence observations of a small number of events that are known to be interesting, due to known or suspected planetary deviations in progress (Sumi et al. 2010) or high-magnification events, which have very high planet detection efficiency (Griest & Safizadeh 1998; Rhie et al. 2000; Rattenbury et al. 2002). The very wide (2.2 deg^2) FOV of the MOA-II 1.8 m telescope with 80M pixel CCD camera MOA-cam3 (Sako et al. 2008) provides high-cadence survey observations of the entire Galactic bulge, and this allows MOA to identify suspected planetary deviations in progress and to predict high magnification ($A_{\text{max}} \gtrsim 100$) for events with relatively short timescales (Einstein radius crossing time $t_E < 20$ days). MOA-2009-BLG-319 is one such short timescale high-magnification event that was identified as a high-magnification event based on MOA data taken ~ 24 hr prior to peak magnification.

In this paper, we report the discovery of a sub-Saturn mass planet in the microlensing event, MOA-2009-BLG-319. We describe the observations and data sets in Section 2. The light curve modeling is presented in Section 3. We discuss the measurement of the source magnitude and color in Section 4 and derive the angular Einstein radius in Section 5. In Section 6, we search for a microlensing parallax signal. In Section 7, we use a Bayesian analysis to estimate the masses and distances of the host star and the planet, based on the angular Einstein radius and microlens parallax. We present our conclusion in Section 8.

2. OBSERVATIONS

For the bulk of the 2009 observing season, the MOA group was the only microlensing survey group in operation because the OGLE group completed the OGLE-III survey on 2009 May 3, in order to upgrade to the OGLE-IV camera with a much wider FOV. Prompted in part by this fact, MOA adopted a new observing strategy for the 2009 observing season in order to increase the planet detection efficiency. The top six fields (a total of 13.2 deg^2) yielded 54% of the microlensing events found by MOA in previous seasons, and these were observed every 15 minutes. The next six fields (with 25% of the previous years' events) were observed every 47 minutes, and most of the remaining 10 fields were observed every 95 minutes. This new observing strategy yielded 563 microlensing alert events in 2009, an increase of about 100 over the 2008 total. MOA-2009-BLG-319 was the first of four of these events to yield an apparent planetary signal.

The event MOA-2009-BLG-319 [(R.A., decl.)_{J2000.0} = ($18^{\text{h}}06^{\text{m}}58^{\text{s}}.13$, $-26^{\circ}49'10''.89$), (l, b) = (4.202, -3.014)] was detected and announced as a normal microlensing alert event by the MOA collaboration on 2009 June 20 (HJD' \equiv HJD $- 2450000 = 5003.056$). The discovery announcement provided a model for this event, which indicated that this was a high-magnification event, and so MOA immediately began follow-up observations in the I and V bands with the University of Canterbury's 0.6 m Boller & Chivens (B&C) telescope at Mt. John Observatory. Public access to the MOA photometry over the next two nights led the μFUN , RoboNet, and MiNDSTeP collaborations to begin observations of this event ~ 2.5 days after its discovery. Three days after the discovery, the MOA data indicated that this event was quite likely to reach high magnification, and the μFUN group issued a high-magnification alert by email to all interested observers, estimating a peak magnification of $A_{\text{max}} > 100$ (at 1σ) 18 hr later at HJD' = 5006.875. This alert message noted "low-level systematics" in the MOA data, which were, in fact, not systematic errors at all. Instead, this light curve feature was the first (weak) planetary caustic crossing. Then, 14 hr later at June 24 UT 01:12 HJD' \simeq 5006.55, data from the μFUN SMARTS CTIO telescope in Chile provided clear evidence for a second, much stronger, caustic crossing feature, which prompted μFUN to issue an anomaly alert. This feature was readily confirmed by the MiNDSTeP observer at La Silla from data already in hand (see Figure 1). A large number of telescopes responded to this anomaly alert, resulting in continuous photometric monitoring of the remainder of the planetary signal with no gaps larger than 5 minutes until after the planetary signal finished, some ~ 20 hr later.

The complete data set for MOA-2009-BLG-319 consists of observations from 20 different observatories representing the MOA, μFUN , RoboNet, MiNDSTeP, PLANET groups, as well as the InfraRed Survey Facility (IRSF) telescope in South Africa. Specifically, the data set includes data from the following telescopes and passbands: MOA-II (New Zealand) 1.8 m wide R band, the Mt. John Observatory B&C (New Zealand) 0.61 m I and V bands, μFUN Auckland Observatory (New Zealand) 0.4 m R -band, μFUN Bronberg (South Africa) 0.35 m unfiltered, μFUN SMARTS CTIO (Chile) 1.3 m V , I , and H bands, μFUN Campo Catino Austral (CAO, Chile) 0.5 m unfiltered, μFUN Farm Cove (New Zealand) 0.35 m unfiltered, μFUN IAC80 (Tenerife, Spain) 0.8 m I band, μFUN Mt. Lemmon (Arizona, USA) 1.0 m I band, μFUN Southern Stars Observatory (SSO, Tahiti) 0.28 m unfiltered, μFUN




Figure 1. Light curve of planetary microlensing event MOA-2009-BLG-319. The top panel shows the data points and the best-fit model light curve with finite source and limb-darkening effects. The three lower panels show close-up views of the four caustic crossing light curve regions and the residuals from the best-fit light curve. The photometric measurements from MOA, B&C, Auckland, Bronberg, CAO, CTIO, Farm Cove, and LOAO are plotted as filled dots with colors indicated by the legend in the top panel. The other data sets are plotted with open circles. The data sets of μ FUN Bronberg and SSO have been averaged into 0.01 day bins, and the RoboNet FTN and FTS data sets are shown in 0.005 day bins, for clarity.

Vintage Lane Observatory (New Zealand) 0.41 m unfiltered, μ FUN Wise (Israel) 0.46 m unfiltered, μ FUN Palomar (USA) 1.5 m *I* band, RoboNet Faulkes Telescope North (FTN, Hawaii) 2.0 m SDSS-*I* band, RoboNet Faulkes Telescope South (FTS, Australia) 2.0 m SDSS-*I* band, RoboNet Liverpool Telescope (La Palma) 2.0 m SDSS-*I* band, MiNDSTeP Danish (La Silla) 1.54 m *I* band, PLANET Canopus (Australia) 1.0 m *I* band, PLANET SAAO (South Africa) 1.0 m *I* band, and IRSF (South Africa) 1.4 m *J*, *H*, and *K_S* bands. This is more follow-up telescopes than have been used for previous planetary microlensing discoveries.

The light curve for this event had four distinct caustic crossing features, which were all observed with good-to-excellent sampling. The first is a weak caustic entry at $\text{HJD}' \sim 5006.05$, which is observed by MOA. The second is a caustic exit at magnification $A \sim 60$ at $\text{HJD}' \sim 5006.6$. This region of the light curve is covered by the CTIO, Danish, Liverpool, and Wise telescopes. The next light curve feature is a strong caustic entry, which produced the light curve peak at $A_{\text{max}} \sim 205$, at $\text{HJD}' \sim 5006.96$. The final caustic exit occurs shortly thereafter at $\text{HJD}' \sim 5007.0$ at a magnification of $A \sim 180$. This main peak covering the third and fourth caustic crossing has excellent coverage, observed by 16 telescopes.

The images were reduced using several different photometry methods. The MOA data sets were reduced by the MOA Difference Image Analysis (DIA) pipeline (Bond et al. 2001). The μ FUN data sets except the CTIO *H* band and Bronberg were reduced by the MOA DIA pipeline and pySIS version

3.0 (Albrow et al. 2009), which is based on the numerical kernel method invented by Bramich (2008). The CTIO *H* band and Bronberg data sets were reduced using the OSU DoPHOT pipeline. The Danish data were reduced by the DIAPL image subtraction software (Wozniak 2000). The RoboNet and PLANET data sets were reduced by pySIS version 3.0. The IRSF data set was reduced by the DoPHOT pipeline. The error bars for the data points are re-normalized so that χ^2 per degree of freedom for the best-fit model is nearly one.

All of these data sets are used for modeling except for the CTIO *V* and *H* bands, the Canopus and SAAO *I* band, and the IRSF *J*, *H*, and *K_S* bands. The CTIO *V* band, the Canopus and SAAO *I* band, and IRSF *J*-, *H*-, and *K_S*-band data sets do not have many observations and do not cover the planetary deviation region of the light curve. The CTIO *H*-band data were not used in the modeling because of a cyclic pattern caused by intrapixel sensitivity variations and image dithering. For our modeling of microlensing parallax effects, we have used a binned data set in order to speed up the modeling calculations. Note that we checked that an analysis with unbinned data gives the same results.

3. MODELING

Inspection of Figure 1 indicates that the event exhibits a number of caustic crossings, so we expect that this event, like most planetary microlensing events, will exhibit significant finite source effects. The first step in modeling is

Table 1

Limb-darkening Coefficients for the Source Star with Effective Temperature $T_{\text{eff}} = 5500$ K, Surface Gravity $\log g = 4.5$, and Metallicity $\log[M/H] = 0.0$ (Claret 2000)

Filter Color	V	R	I	J	H	K
c	0.3866	0.2556	0.1517	-0.0234	-0.2154	-0.1606
d	0.4303	0.5027	0.5281	0.6021	0.7695	0.6324

therefore to measure the source color, which then enables us to determine the limb-darkening parameters for the various light curves.

3.1. Source Color

Once a microlensing model is found, the dereddened source color and magnitude $[I, (V - I)]_0$ can be determined by comparing the instrumental values of these quantities to those of the red clump (Yoo et al. 2004). This is described in Section 4. However, before a good model can be found, the limb-darkening coefficients must be determined, which requires an estimate of the source color. This seemingly endless loop can be broken by making a model-independent measurement of the instrumental source color from a regression of V -band flux on I -band flux (and then comparing this value to the instrumental clump color). We find $(V - I)_0 = 0.82$, as reported in more detail in Section 4.

3.2. Limb Darkening

We adopt a two-parameter square-root law (Claret 2000) for the surface brightness of the source,

$$S_\lambda(\vartheta) = S_\lambda(0)[1 - c(1 - \cos \vartheta) - d(1 - \sqrt{\cos \vartheta})]. \quad (1)$$

Here, c and d are the limb-darkening coefficients, $S_\lambda(0)$ is the central surface brightness of the source, and ϑ is the angle between the normal to the stellar surface and the line of sight, i.e., $\sin \vartheta = \theta/\theta_*$, where θ is the angular distance from the center of the source.

Based on the source color estimate of $(V - I) = 0.82$, the source is likely to have a G8 spectral type and an effective temperature of $T_{\text{eff}} = 5475$ K according to Bessell & Brett (1988). We use limb-darkening parameters from Claret (2000) for a source star with effective temperature $T_{\text{eff}} = 5500$ K, surface gravity $\log g = 4.5$, and metallicity $\log[M/H] = 0.0$ as presented in Table 1.

3.3. Best-fit Model

We search for the best-fit binary lens model using a variation of the Markov Chain Monte Carlo (MCMC) algorithm (Verde et al. 2003) due to Doran & Mueller (2004) and Bennett (2010) that frequently changes the ‘‘jump function’’ in order to find the χ^2 minimum more quickly. There are three lensing parameters in common with single lens events, the time of the closest approach to the center of mass t_0 , the Einstein crossing time t_E , and the minimum impact parameter u_0 . Binary lens models require four additional parameters: the planet–star mass ratio q , the binary lens separation d , which is projected onto the plane of the sky and normalized by the angular Einstein radius θ_E , the angle of the source trajectory relative to the binary lens axis α , and source radius relative to the Einstein radius $\rho = \theta_*/\theta_E$. In addition, for each data set and passband, there are two parameters to describe the unmagnified source and background fluxes in that band.

We begin by conducting a very broad parameter search. The parameter search has been conducted by two independent




Figure 2. Caustic is plotted by the solid curve for the MOA-2009-BLG-319 best-fit model, and the dashed line indicates the source star trajectory. The circle represents the source star size. The source star crosses the caustic curve four times, with peak magnification of $A_{\text{max}} \sim 205$ during the third caustic crossing at $\text{HJD}' \sim 5006.96$.

codes. We perform 300 separate χ^2 minimizations with initial parameters distributed over the ranges $-5 < \log q < -1$, $-3 < \log d < 0.3$, in order to identify the parameter regimes of models that could explain the light curve. The initial parameters with $\log d > 0.3$ were not prepared because of the $d \leftrightarrow d^{-1}$ symmetry. We find that the only models consistent with the observed light curve have $q \sim 10^{-4}$ and $d \sim 1$ and that the best-fit model has $q = (3.95 \pm 0.02) \times 10^{-4}$, $d = 0.97537 \pm 0.00007$, and other parameters as listed in Table 2. The projected position of the planet is pretty close to the Einstein ring, and therefore d was well constrained. The light curves and caustic of this event are shown in Figure 1 and Figure 2, respectively, which resemble Figure 8 in Wambsganss (1997). Here we assumed no orbital motion of the planet around the host star in our model. So, the d and α are the average separation and angle during half a day when the source is crossing the caustics. The changes of these parameters due to the orbital motion during this period could be of the same order or slightly larger than the nominal MCMC error of the average values given above. These changes do not affect the results in this analysis because they are much smaller than the uncertainty given in Section 7.

4. SOURCE MAGNITUDE AND COLOR

The dereddened source magnitude and color can be estimated as follows. We locate the clump in the color–magnitude diagram (CMD) of stars within $2'$ of the source star, shown in Figure 3, with the following method. The stars, which are $I < 17$ mag and $(V - I) > 1.5$ mag, were used for the clump estimate. Among them, the stars within 0.3 mag from the clump centroid were picked up. Note that the clump in the first turn was assumed. Then, the mean magnitude of I and mean color $(V - I)$ were calculated using the stars within 0.3 mag and replaced as the new clump centroid. This was iterated until the clump centroid position is converged. Therefore, we find the clump as $[I, (V - I)]_{\text{clump}} = (15.31, 1.91)$. The best model source brightness and color are obtained as $[I, (V - I)]_s = (19.82, 1.69)$ from the fits. With a 0.05 mag correction due to blending by fainter stars in this crowded field (Bennett et al. 2010), this yields

$$[I, (V - I)]_s - [I, (V - I)]_{\text{clump}} = (4.51, -0.22). \quad (2)$$

We adopt the dereddened red clump giant (RCG) magnitude $M_{I,0,\text{clump}} = -0.25$ and color $(V - I)_{0,\text{clump}} = 1.04$ from

Table 2
Best-fit Model Parameters with Various Effects, Finite Source, Orbital and Terrestrial Parallax, and u_0

Orbital Parallax	Terrestrial Parallax	$u_0 > 0$	$u_0 < 0$	t_0 HJD'	t_E (days)	u_0 (10^{-3})	q (10^{-4})	d	α (rad)	ρ (10^{-3})	$\pi_{E,E}$	$\pi_{E,N}$	χ^2
		○		5006.99482	16.57	6.22	3.95	0.97537	5.7677	1.929	7023.8
			σ	0.00006	0.08	0.03	0.02	0.00007	0.0005	0.010	
			○	5006.99485	16.56	-6.23	3.95	0.97540	0.5156	1.931	7023.8
			σ	0.00005	0.08	0.03	0.02	0.00006	0.0005	0.009	
○		○		5006.99480	16.59	6.22	3.95	0.97540	5.7673	1.929	0.40	0.30	7023.2
			σ	0.00007	0.09	0.03	0.02	0.00007	0.0005	0.011	
○			○	5006.99482	16.56	-6.23	3.95	0.97534	0.5155	1.931	0.40	-0.30	7023.4
			σ	0.00006	0.08	0.03	0.02	0.00007	0.0005	0.010	
	○	○		5006.99477	16.61	6.21	3.94	0.97540	5.7671	1.926	-0.23	0.12	7017.6
			σ	0.00006	0.08	0.03	0.02	0.00007	0.0005	0.010	0.07	0.04	
	○		○	5006.99483	16.57	-6.23	3.95	0.97542	0.5161	1.931	-0.02	0.26	7019.2
			σ	0.00006	0.07	0.03	0.02	0.00007	0.0005	0.010	0.04	0.07	
○	○	○		5006.99478	16.60	6.21	3.94	0.97540	5.7673	1.926	-0.15	0.15	7017.7
			σ	0.00006	0.08	0.03	0.02	0.00007	0.0004	0.010	0.07	0.05	
○	○		○	5006.99481	16.56	-6.23	3.95	0.97538	0.5162	1.932	-0.04	0.23	7019.4
			σ	0.00006	0.07	0.03	0.02	0.00007	0.0005	0.009	0.04	0.07	

Notes. The lines with “σ” list the 1σ error of parameters given by MCMC. HJD' \equiv HJD-2450000. Note that the u_0 conventions are the same as Figure 2 of Gould (2004). χ^2 value is the result of the fitting with 18 data sets, which have 7210 data points. The model search with finite source and orbital parallax effects were done by a grid search.




Figure 3. $(V - I, I)$ CMD of the stars within $2'$ of the MOA-2009-BLG-319 source using μ FUN CTIO data calibrated to OGLE-II. The filled triangle and square indicate the source and blend stars, respectively, assuming that the blended light comes from a single star. The filled circle indicates the center of the red clump giant distribution.

Bennett et al. (2008), which is based on Girardi & Salaris (2001) and Salaris & Girardi (2002). Rattenbury et al. (2007) find that the clump in this field lies 0.12 mag in the foreground of the Galactic center, which we take to be at $R_0 = 8.0 \pm 0.3$ kpc (Yelda et al. 2010). Hence, the distance modulus of the clump is DM = 14.40. This yields a dereddened RCG centroid in this field of

$$[I, (V - I)]_{\text{clump},0} = (14.15, 1.04). \quad (3)$$

Assuming that the source suffers the same extinction as the clump, we use the best-fit source magnitude and color to obtain the dereddened values for the source,

$$\begin{aligned} [I, (V - I)]_{S,0} &= (14.15, 1.04) + (4.51, -0.22) \\ &= (18.66, 0.82). \end{aligned} \quad (4)$$

A comparison of $(V - I)_{S,0}$ estimated by this method to

14 spectra of microlensed source stars at high magnification (Bensby et al. 2010) suggests that $(V - I)_{S,0}$ is determined with an uncertainty of 0.06 mag. For the uncertainty in $I_{S,0}$, we estimate uncertainties of 0.08 from R_0 , 0.05 from the Galactic bulge RCG centroid, and 0.05 from the Rattenbury et al. (2007) offset from the Galactic center, which when added in quadrature yields a total uncertainty of 0.11 mag.

Equation (3) implies extinction of $A_I = 1.16 \pm 0.11$ and reddening $E(V - I) = 0.87 \pm 0.08$, which is consistent within the error with $E(V - I) = 0.97 \pm 0.03$ from the OGLE-II extinction map (Sumi 2004).

5. MEASUREMENT OF THE ANGULAR EINSTEIN RADIUS, θ_E

The sharp caustic crossing features in the MOA-2009-BLG-319 light curve resolve the finite angular size of the source star, and these finite source effects allow us to determine the angular Einstein radius θ_E and the lens-source relative proper motion $\mu_{\text{rel}} = \theta_E/t_E$. Following Yoo et al. (2004), we use the dereddened color and magnitude of the source $[I, (V - I)]_{S,0}$ from Equation (4). Next, we obtain the source angular radius using the source V and K magnitude. We estimate $(V - K)_0$ from $(V - I)_0$ and the Bessell & Brett (1988) color-color relations for dwarf stars,

$$[K, (V - K)]_{S,0} = (17.67, 1.81) \pm (0.14, 0.15). \quad (5)$$

We also estimate the K magnitude using IRSF data, $K_{S,0} = 18.09 \pm 0.42$. This is consistent with but less accurate than the K magnitude estimated from $(V - I)_0$. So, we use the K magnitude estimated from $(V - I)_0$. For main-sequence stars, the relationship between color, brightness, and a star angular radius θ_* was determined by Kervella et al. (2004) to be

$$\log(2\theta_*) = 0.0755(V - K) + 0.5170 - 0.2K, \quad (6)$$

which with K and $(V - K)$ from Equation (5) implies

$$\theta_* = 0.66 \pm 0.06 \mu\text{as}. \quad (7)$$




Figure 4. Contours of $\Delta\chi^2 = 1, 4, 9, 16$ with orbital and terrestrial parallax parameters. The left panel is the result with $u_0 > 0$ and the right panel is with $u_0 < 0$. The best-fit result with $u_0 > 0$ is better than $u_0 < 0$ about $\Delta\chi^2 = 1.7$. Furthermore, the best-fit model with and without parallax is different with $\Delta\chi^2 = 6.1$.

(A color version of this figure is available in the online journal.)

The fit parameter $\rho \equiv \theta_*/\theta_E$ is the source star angular radius in units of the angular Einstein radius. Thus, the angular Einstein radius θ_E is

$$\theta_E = \frac{\theta_*}{\rho} = 0.34 \pm 0.03 \text{ mas}. \quad (8)$$

Therefore, the source-lens relative proper motion μ is

$$\mu = \frac{\theta_E}{t_E} = 7.52 \pm 0.65 \text{ mas yr}^{-1}. \quad (9)$$

6. MICROLENSING PARALLAX EFFECT

The event timescale is not long, $t_E = 16.6$ days, so one does not expect to detect the orbital microlensing parallax effect (Refsdal 1966; Gould 1992; Alcock et al. 1995). However, the very sharp third peak was observed simultaneously from Australia, New Zealand, and Hawaii, i.e., along two nearly perpendicular base lines of length, $0.36 R_\oplus$ and $1.25 R_\oplus$, respectively. Therefore, there is some chance that these data will reveal a signal due to terrestrial microlensing parallax (Hardy & Walker 1995; Holz & Wald 1996; Gould et al. 2009).

Microlensing parallax is usually described by the parallax parameter, π_E , which is the amplitude of the two-dimensional microlens parallax vector, and the two components of this vector are denoted by $\pi_{E,E}$ and $\pi_{E,N}$, which are the east and north components of the vector on the sky. The microlens parallax vector has the same direction as the lens-source proper motion, perpendicular to the line of sight. It is related to the lens-source relative parallax π_{rel} and the angular Einstein radius θ_E (Gould 2000) by

$$\pi_E = \frac{\pi_{\text{rel}}}{\theta_E}, \quad \pi_{\text{rel}} = \pi_L - \pi_S, \quad (10)$$

where π_L and π_S are the lens and the source parallaxes, respectively.

Our initial search for microlensing parallax included both the orbital and terrestrial effect, as is necessary for a physically correct model. Our initial fits indicated a weak microlensing parallax signal, so we searched for orbital parallax and terrestrial parallax signals separately, in order to determine which type of parallax signal is being seen and to test for possible systematic errors. We must also consider alternative model solutions due to the $u_0 > 0 \leftrightarrow u_0 < 0$ degeneracy first noted by Smith et al. (2003). As the model results listed in Table 2 indicate, orbital parallax can improve the fit χ^2 by only $\Delta\chi^2 =$

0.6, with two additional parameters, which is not at all significant. The best terrestrial parallax model, however, does give a formally significant χ^2 improvement of $\Delta\chi^2 = 6.2$, but this improvement decreases to $\Delta\chi^2 = 6.1$ for the best physical (terrestrial plus orbital) parallax model. With two additional parameters, this is formally significant at almost the 95% confidence level. Figure 4 shows the $\Delta\chi^2$ contours for microlensing parallax fits to the MOA-2009-BLG-319 light curve.

The best-fit parallax model has $u_0 > 0$ and $(\pi_{E,E}, \pi_{E,N}) = (-0.15, 0.15) \pm (0.07, 0.05)$, while the best-fit $u_0 < 0$ model has a χ^2 value that is larger than the best-fit $u_0 > 0$ solution by 1.7 and only an improvement of $\Delta\chi^2 = 4.4$ over the best-fit non-parallax solution. Thus, the best $u_0 < 0$ model is neither a significant improvement over the best non-parallax model nor significantly worse than the best parallax model. We find that χ^2 improvement for the best-fit parallax model comes from the Mt. John observatory (MOA-II 1.8 m and Canterbury 0.6m) telescopes alone, with a total χ^2 improvement $\Delta\chi^2 = 7.3$, while the contribution of all the other data sets is $\Delta\chi^2 = -1.2$ (i.e., the parallax model is disfavored). One would expect that χ^2 should improve for the many other data sets, and the fact that it does not suggest that the parallax signal may not be real.

If we assume that the scalar parallax measurement of π_E is correct, then it implies that the lens system is located in the inner Galactic disk. Due to the flat rotation curve of the Galaxy, the stars at this location are rotating much faster than the typical line of sight to a Galactic bulge star. As a result, the direction of the parallax vector (which is parallel to the lens-source relative velocity) is most likely to be in the direction of Galactic rotation, which is $\sim 30^\circ$ east of north. This is similar to the direction of the parallax vector for the best $u_0 < 0$ model, but it is roughly perpendicular to that for the $u_0 > 0$ model. So, the $u_0 > 0$ solution appears to be disfavored on a priori grounds.

Because of the low significance of the microlensing parallax signal and the indications of possible systematic problems with the measurement of the parallax parameters, we will use only an upper limit on the microlensing parallax effect in our analysis.

7. THE LENS PROPERTIES

We can place lower limits on the lens mass and distance with our measured angular Einstein radius, θ_E , and our upper limit on the amplitude of the microlens parallax vector, π_E . The lens mass is given by

$$M = \frac{\theta_E}{\kappa\pi_E}, \quad (11)$$

where $\kappa = 4G/(c^2 \text{ AU}) = 8.1439 \text{ mas } M_\odot^{-1}$. With our upper limit from the previous section, $\pi_E < 0.5$, gives a lower limit on the total mass of the lens system, $M > 0.08 M_\odot$. This implies that the lens primary is more massive than a brown dwarf and must be a star or stellar remnant. From Equation (10), this implies that the source-lens relative parallax is $\pi_{\text{rel}} < 0.17 \text{ mas}$.

The vast majority of source stars for microlensing events seen toward the bulge are stars in the bulge, and the MOA-2009-BLG-319 source magnitude and colors are consistent with a bulge G-dwarf source. So, it is reasonable to assume that the source is a bulge star with a distance of $D_S \approx 8.0 \text{ kpc}$. This implies that the lens parallax is $\pi_L = \pi_{\text{rel}} + \pi_S < 0.30 \text{ mas}$, from Equation (10). The lens parallax is related to the distance by $\pi_L = 1 \text{ AU}/D_L$, so a lower limit on the lens distance is $D_L > 3.33 \text{ kpc}$.

An upper limit on the lens mass may be obtained if we assume that the planetary host star is a main-sequence star and not a stellar remnant. We can consider the blended flux seen at the same location of the source beyond the measured source flux from the microlensing models. If we attribute this blended flux to a single star, we can follow the reasoning of Section 4 in order to estimate the dereddened magnitude of the blend star:

$$(I, V - I)_{b,0} = (17.78, 0.75) \pm (0.12, 0.14), \quad (12)$$

under the (conservative) assumption that the blend star lies behind all the foreground dust. We can now use this as an upper limit on the brightness of a main-sequence lens star. From Schmidt-Kaler (1982) and Bessell & Brett (1988), we find an upper limit on the host star mass of $M < 1.14 M_\odot$.

As we found finite source effects in the light curve, we can break out one degeneracy of the lens star mass M , distance D_L , and velocity v . We calculated the probability distribution from Bayesian analysis by combining this equation and the measured values of θ_E and t_E with the Galactic model (Han & Gould 2003) assuming that the distance to the Galactic center is 8 kpc. We included the upper limit of microlens parallax amplitude. A constraint of the upper limit for blending light was also included for the lens mass upper limit. The probability distribution from a Bayesian analysis is shown in Figure 5. The host star is a K- or M-dwarf star with a mass of $M_L = 0.38^{+0.34}_{-0.18} M_\odot$ and distance $D_L = 6.1^{+1.1}_{-1.2} \text{ kpc}$, planetary mass $M_p = 50^{+44}_{-24} M_\oplus$, and projected separation $r_\perp = 2.0^{+0.4}_{-0.4} \text{ AU}$. The physical three-dimensional separation, $a = 2.4^{+1.2}_{-0.6} \text{ AU}$, was estimated by putting a planetary orbit at random inclination, eccentricity, and phase (Gould & Loeb 1992).

8. DISCUSSION AND CONCLUSION

We have reported the discovery of a sub-Saturn mass planet in the light curve of the microlensing event MOA-2009-BLG-319. This event was observed by 20 telescopes, the largest number of telescopes to participate in a microlensing planet discovery to date. The lens system has a mass ratio $q = (3.95 \pm 0.02) \times 10^{-4}$ and a separation $d = 0.97537 \pm 0.00007$ Einstein radii. The lens-source relative proper motion was determined to be $\mu_{\text{rel}} = 7.52 \pm 0.65 \text{ mas yr}^{-1}$ from the measurement of finite source effects. A slightly better light curve fit can be obtained when the (terrestrial) microlensing parallax effect is included in the model, yielding an improvement of $\Delta\chi^2 = 6.1$. This is very marginal statistical significance, and there are indications that systematic errors may influence the result. So, we use our microlensing parallax analysis to set an upper limit of $\pi_E < 0.5$.




Figure 5. Probability distribution from a Bayesian analysis for the distance, D_L , mass, M_L , and the physical three-dimensional separation a . The vertical solid lines indicate the median values. The dark and light shaded regions indicate the 68% and 95% limits. The solid curve in the top panel indicates the mass–distance relation of the lens from the measurement of θ_E assuming $D_S = 8 \text{ kpc}$. Note that D_S is not fixed in the actual Bayesian analysis.

The probability distribution estimated from a Bayesian analysis indicates that the lens host star mass is $M_L = 0.38^{+0.34}_{-0.18} M_\odot$ with a sub-Saturn mass planet $M_p = 50^{+44}_{-24} M_\oplus$ and the physical three-dimensional separation $a = 2.4^{+1.2}_{-0.6} \text{ AU}$. The distance of the lens star is $D_L = 6.1^{+1.1}_{-1.2} \text{ kpc}$. The known microlensing exoplanets are summarized in Figure 6 and Table 3. MOA-2009-BLG-319Lb lies at ~ 2.3 times the distance of the snow line, which is estimated to be at $a_{\text{snow}} = 2.7 \text{ AU } M/M_\odot$. This is similar to the separation of other planets found by microlensing (Sumi et al. 2010).

There are some indications of low-level systematic deviations from the best-fit model remaining in the light curve, near the third and fourth caustic crossing features (see the bottom panel residuals in Figure 1), which do not affect the results in this analysis. These systematic light curve deviations might be caused by orbital motion of the lens, a second planet, or systematic photometry errors. A more detailed analysis will be performed in the future when the adaptive optics images from the Keck telescope are reduced, and this analysis may shed more light on the mass and distance of the host star.

The next few years are expected to see an increase in the rate of microlensing planet discoveries. The OGLE group has started the OGLE-IV survey with their new 1.4 deg^2 CCD camera in

Table 3
Parameters of Exoplanets Discovered by Microlensing

Name	Host Star Mass $M_L (M_\odot)$	Distance D_L (kpc)	Planet Mass M_p	Separation a (AU)	Mass Estimated by
OGLE-2003-BLG-235Lb	$0.63^{+0.07}_{-0.09}$	$5.8^{+0.6}_{-0.7}$	$2.6^{+0.8}_{-0.6} M_J$	$4.3^{+2.5}_{-0.8}$	θ_E , lens brightness
OGLE-2005-BLG-071Lb	0.46 ± 0.04	3.2 ± 0.4	$3.8 \pm 0.4 M_J$	3.6 ± 0.2	θ_E, π_E , detection of the lens
OGLE-2005-BLG-169Lb	$0.49^{+0.23}_{-0.29}$	$2.7^{+1.6}_{-1.3}$	$13^{+6}_{-8} M_\oplus$	$2.7^{+1.7}_{-1.4}$	θ_E , Bayesian
OGLE-2005-BLG-390Lb	$0.22^{+0.21}_{-0.11}$	$6.6^{+1.0}_{-1.0}$	$5.5^{+5.5}_{-2.7} M_\oplus$	$2.6^{+1.5}_{-0.6}$	θ_E , Bayesian
OGLE-2006-BLG-109Lb	$0.51^{+0.05}_{-0.04}$	1.49 ± 0.19	$231 \pm 19 M_\oplus$	2.3 ± 0.5	θ_E, π_E
c			$86 \pm 7 M_\oplus$	$4.5^{+2.1}_{-1.0}$	θ_E, π_E
OGLE-2007-BLG-368Lb	$0.64^{+0.21}_{-0.26}$	$5.9^{+0.9}_{-1.4}$	$20^{+7}_{-8} M_\oplus$	$3.3^{+1.4}_{-0.8}$	θ_E , Bayesian
MOA-2007-BLG-192Lb	$0.084^{+0.015}_{-0.012}$	$0.70^{+0.21}_{-0.12}$	$3.2^{+5.2}_{-1.8} M_\oplus$	$0.66^{+0.19}_{-0.14}$	θ_E, π_E
MOA-2007-BLG-400Lb	$0.30^{+0.19}_{-0.12}$	$5.8^{+0.6}_{-0.8}$	$0.83^{+0.49}_{-0.31} M_J$	$0.72^{+0.38}_{-0.16} / 6.5^{+3.2}_{-1.2}$	θ_E , Bayesian
MOA-2008-BLG-310Lb	0.67 ± 0.14	> 6.0	$28^{+58}_{-23} M_\oplus$	$1.4^{+0.7}_{-0.3}$	θ_E , Bayesian
MOA-2009-BLG-319Lb	$0.38^{+0.34}_{-0.18}$	$6.1^{+1.1}_{-1.2}$	$50^{+44}_{-24} M_\oplus$	$2.4^{+1.2}_{-0.6}$	θ_E , Bayesian

Notes. MOA-2007-BLG-400Lb has two solutions due to a strong close/wide model degeneracy, and details of the MOA-2008-BLG-310Lb parameters are discussed by Janczak et al. (2010) and Sumi et al. (2010).




Figure 6. Exoplanets as a function of mass vs. semimajor axis. The red circles with error bars indicate planets found by microlensing. The filled circles indicate planets with mass measurements, while open circles indicate Bayesian mass estimates. MOA-2009-BLG-319Lb is indicated by the gold-filled open circle. The black dots and blue squares indicate the planets discovered by radial velocities and transits, respectively. The magenta and green triangles indicate the planets detected via direct imaging and timing, respectively. The non-microlensing exoplanet data were taken from The Extrasolar Planets Encyclopaedia (<http://exoplanet.eu/>). The planets in our solar system are indicated with initial letters.

2010 March. This will allow OGLE to survey the bulge at a cadence almost as high as that of MOA-II, but with better seeing that should yield a substantial increase in the rate of microlensing planet discoveries. MOA also plans an upgrade to a $\sim 10 \text{ deg}^2$ MOA-III CCD camera in a few years, which will allow an even higher cadence Galactic bulge survey. The Korean Microlensing Telescope Network (KMTNet) is funded to dramatically increase the longitude coverage of microlensing survey telescopes. They plan three wide FOV telescopes to go in South Africa, Australia, and South America. When these telescopes come online, we anticipate another dramatic increase in the microlensing planet discovery rate.

The MOA project and a part of authors were supported by the Grant-in-Aid for Scientific Research, the grant JSPS20340052, JSPS18253002, JSPS Research fellowships, the Global COE Program of Nagoya University “Quest for Fundamental Principles in the Universe” from JSPS and MEXT of Japan, the

Marsden Fund of New Zealand, the New Zealand Foundation for Research and Technology, and grants-in-aid from Massey University and the University of Auckland. N.M. was supported by JSPS Research Fellowships for Young Scientists. T.S. was supported by the grant JSPS20740104. D.P.B. was supported by grants NNX07AL71G and NNX10AI81G from NASA and AST-0708890 from the NSF. A.G. and S.D. were supported in part by NSF AST-0757888. A.G., S.D., S.G., and R.P. were supported in part by NASA NNG04GL51G. Work by S.D. was performed in part under contract with the California Institute of Technology (Caltech) funded by NASA through the Sagan Fellowship Program. J.C.Y. was supported by an NSF Graduate Research Fellowship. Work by C.H. was supported by the Creative Research Initiative program (2009-0081561) of the National Research Foundation of Korea. Astronomical research at Armagh Observatory was supported by the Department of Culture, Arts, and Leisure (DCAL), Northern Ireland, UK. F.F., D.R., and J.S. acknowledge support from the Communauté française de Belgique—Actions de recherche concertées—Académie universitaire Wallonie-Europe.

REFERENCES

- Albrow, M. D., et al. 2009, *MNRAS*, **397**, 2099
 Alcock, C., et al. 1995, *ApJ*, **454**, L125
 Beaulieu, J.-P., et al. 2006, *Nature*, **439**, 437
 Bennett, D. P. 2010, *ApJ*, **716**, 1408
 Bennett, D. P., & Rhie, S. H. 1996, *ApJ*, **472**, 660
 Bennett, D. P., et al. 2008, *ApJ*, **684**, 663
 Bennett, D. P., et al. 2010, *ApJ*, **713**, 837
 Bensby, T., et al. 2010, *A&A*, **512**, A41
 Bessell, M. S., & Brett, J. M. 1988, *PASP*, **100**, 1134
 Bond, I. A., et al. 2001, *MNRAS*, **327**, 868
 Bond, I. A., et al. 2004, *ApJ*, **606**, L155
 Bramich, D. M. 2008, *MNRAS*, **386**, L77
 Claret, A. 2000, *A&A*, **363**, 1081
 Cumming, A., Butler, R. P., Marcy, G. W., Vogt, S. S., Wright, J. T., & Fischer, D. A. 2008, *PASP*, **120**, 531
 Dong, S., et al. 2009, *ApJ*, **698**, 1826
 Doran, M., & Mueller, C. M. 2004, *J. Cosmol. Astropart. Phys.*, **JCAP09(2004)003**
 Gaudi, B. S., et al. 2008, *Science*, **319**, 927
 Girardi, L., & Salaris, M. 2001, *MNRAS*, **323**, 109
 Gould, A. 1992, *ApJ*, **392**, 442
 Gould, A. 2000, *ApJ*, **542**, 785
 Gould, A. 2004, *ApJ*, **606**, 319
 Gould, A., & Loeb, A. 1992, *ApJ*, **396**, 104

- Gould, A., et al. 2006, *ApJ*, **644**, L37
Gould, A., et al. 2009, *ApJ*, **698**, L147
Gould, A., et al. 2010, *ApJ*, **720**, 1073
Griest, K., & Safizadeh, N. 1998, *ApJ*, **500**, 37
Han, C., & Gould, A. 2003, *ApJ*, **592**, 172
Hardy, S. J., & Walker, M. A. 1995, *MNRAS*, **276**, L79
Holz, D. E., & Wald, R. M. 1996, *ApJ*, **471**, 64
Ida, S., & Lin, D. N. C. 2004, *ApJ*, **616**, 567
Janczak, J., et al. 2010, *ApJ*, **711**, 731
Kennedy, G. M., & Kenyon, S. J. 2008, *ApJ*, **673**, 502
Kervella, P., Thévenin, F., Di Folco, E., & Ségransan, D. 2004, *A&A*, **426**, 297
Lecar, M., Podolak, M., Sasselov, D., & Chiang, E. 2006, *ApJ*, **640**, 1115
Rattenbury, N. J., Mao, S., Sumi, T., & Smith, M. C. 2007, *MNRAS*, **378**, 1064
Rattenbury, N. J., Bond, I. A., Skuljan, J., & Yock, P. C. M. 2002, *MNRAS*, **335**, 159
Refsdal, S. 1966, *MNRAS*, **134**, 315
Rhie, S. H., et al. 2000, *ApJ*, **533**, 378
Sako, T., et al. 2008, *Exp. Astron.*, **22**, 51
Salaris, M., & Girardi, L. 2002, *MNRAS*, **337**, 332
Schmidt-Kaler, Th. 1982, in Landolt-Börnstein: Numerical Data and Functional Relationships in Science and Technology, Vol. 2b, ed. K. Schaifers & H. H. Voigt (Berlin: Springer)
Smith, M. C., Mao, S., & Paczyński, B. 2003, *MNRAS*, **339**, 925
Sumi, T. 2004, *MNRAS*, **349**, 193
Sumi, T., et al. 2003, *ApJ*, **591**, 204
Sumi, T., et al. 2010, *ApJ*, **710**, 1641
Udalski, A. 2003, *Acta Astron.*, **53**, 291
Udalski, A., et al. 2005, *ApJ*, **628**, L109
Verde, L., et al. 2003, *ApJS*, **148**, 195
Wambsganss, J. 1997, *MNRAS*, **284**, 172
Woźniak, P. R. 2000, *Acta Astron.*, **50**, 421
Yelda, S., Ghez, A. M., Lu, J. R., Do, T., Clarkson, W., & Matthews, K. 2010, arXiv:1002.1729
Yoo, J., et al. 2004, *ApJ*, **603**, 139