

## ORIGINAL ARTICLE

# A Survey of Genome Editing Activity for 16 Cas12a Orthologs

Bernd Zetsche,<sup>1-5,\*</sup> Jonathan Strecker,<sup>1-4,\*</sup> Omar O. Abudayyeh,<sup>1-4,6</sup> Jonathan S. Gootenberg,<sup>1-4,7</sup>  
David A. Scott<sup>1-4</sup> and Feng Zhang<sup>1-4</sup>

<sup>1</sup> Broad Institute of MIT and Harvard, Cambridge, USA

<sup>2</sup> McGovern Institute for Brain Research at MIT, Cambridge, USA

<sup>3</sup> Department of Brain and Cognitive Science, MIT, Cambridge, USA

<sup>4</sup> Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, USA

<sup>5</sup> Department of Developmental Pathology, Institute of Pathology, Bonn Medical School, Bonn, Germany

<sup>6</sup> Harvard-MIT Division of Health Science and Technology, Cambridge, USA

<sup>7</sup> Department of Systems Biology, Harvard Medical School, Boston, USA

\*These authors contributed equally to this work.

(Received for publication on August 6, 2019)

(Revised for publication on October 18, 2019)

(Accepted for publication on October 18, 2019)

(Published online in advance on November 14, 2019)

The class 2 CRISPR-Cas endonuclease Cas12a (previously known as Cpf1) offers several advantages over Cas9, including the ability to process its own array and the requirement for just a single RNA guide. These attributes make Cas12a promising for many genome engineering applications. To further expand the suite of Cas12a tools available, we tested 16 Cas12a orthologs for activity in eukaryotic cells. Four of these new enzymes demonstrated targeted activity, one of which, from *Moraxella bovoculi* AAX11\_00205 (Mb3Cas12a), exhibited robust indel formation. We also showed that Mb3Cas12a displays some tolerance for a shortened PAM (TTN versus the canonical Cas12a PAM TTTV). The addition of these enzymes to the genome editing toolbox will further expand the utility of this powerful technology. (DOI: 10.2302/kjm.2019-0009-OA)

**Keywords:** genome editing; CRISPR-Cas; gene therapy

### Introduction

The ability to edit the genome of living cells enables a broad range of downstream genetic analyses and has the potential for therapeutic use to resolve pathogenic mutations. Over the past several years, enzymes from CRISPR-Cas systems, which provide bacteria and archaea with adaptive immunity, have emerged as powerful tools for eukaryotic gene editing. In nature, CRISPR-Cas systems acquire DNA snippets that match invading viruses or foreign nucleic acids, creating a memory bank of infection. These snippets are then transcribed into short RNA guides, which are used by Cas proteins to detect invading

nucleic acids. Once a sequence match is found, Cas nucleases destroy the foreign nucleic acid. In particular, Class 2 CRISPR-Cas systems are well-suited for development as molecular technologies because they contain single effector enzymes. These effector enzymes, such as Cas9, are RNA-guided DNA endonucleases that have been harnessed for a range of genome engineering applications.<sup>1</sup>

Although Cas9 was the first such enzyme to be developed as a genome editing tool,<sup>2,3</sup> three orthologs of Cas12a (a single RNA-guided class 2 effector previously known as Cpf1) from *Francisella novicida* U112 (Fn-Cas12a), *Acidaminococcus* sp. BV3L6 (AsCas12a), and *Lachnospiraceae* bacterium ND2006 (LbCas12a), have

Reprint requests to: Feng Zhang, PhD, Broad Institute of MIT and Harvard, 75 Ames Street, Cambridge, MA 02142, USA, E-Mail: zhang@broadinstitute.org

The supplemental file is available in the on-line version only.

Copyright © 2019 by The Keio Journal of Medicine

also been used for genome editing in eukaryotic cells.<sup>4–8</sup> Endonucleases of the Cas12a family differ from the Cas9 family in several ways: (i) Cas12a utilizes T-rich proto-spacer adjacent motifs (PAMs) located 5' of the targeted DNA sequence, (ii) target cleavage occurs distally from the PAM and results in sticky-end overhangs, (iii) Cas12a is guided by a single CRISPR RNA (crRNA) and does not require trans-activating CRISPR RNA; and (iv) Cas12a possesses both RNase and DNase activity, which allows it to process its own CRISPR array.<sup>7,9</sup> These features make Cas12a particularly useful in certain situations, such as targeting AT-rich genomic regions and multiplexed gene targeting.<sup>8,10</sup> Additionally, Cas12a has been shown to possess non-specific single-stranded DNA cleavage activity after it has been activated by target binding, which has been leveraged for nucleic acid detection.<sup>11–14</sup> Finally, Cas12a is more specific than Cas9 in certain contexts, making it well-suited to applications in which high specificity is critical.<sup>15,16</sup>

Given previous work showing that different Cas enzyme orthologs exhibit a range of activity in eukaryotic cells<sup>2,16,17</sup> and indicating the potential advantages of Cas12a, we sought to identify additional Cas12a orthologs with high activity in eukaryotic cells. Here we examine 16 new Cas12a-family proteins for nuclease activity in human cells. We identify four orthologs that can induce insertion/deletion (indel) events at targeted genomic loci. One ortholog, from *Moraxella bovoculi* AAX11\_00205 (Mb3Cas12a), exhibited comparable activity to AsCas12a and LbCas12a when targeting sites containing TTTV (V=A, C, or G) PAMs. We also show that Mb3Cas12a can recognize a TTN PAM, but with lower efficiency than the conserved TTTV PAM. Together, these new orthologs expand the genome editing toolbox, providing new enzymes that can be used for tailored applications.

## Materials and Methods

*Computational search for Cas12a orthologs.* Cas12a orthologs were selected as previously described.<sup>7</sup>

*Cell culture and transfection.* HEK293T cells were maintained at 37°C with 5% CO<sub>2</sub> in Dulbecco's Modified Eagle Medium (Gibco) supplemented with 10% fetal bovine serum (HyClone) and 2 mM GlutaMAX (Life Technology). For indel analysis, 22,000 cells were seeded per well of a 96-well plate (Corning) 1 day before transfection. Each well was transfected with 100 ng Cas12a-encoding plasmid (see the supplemental file) and 50 ng guide-encoding plasmid or PCR fragment, or 150 ng Cas12a and guide-encoding plasmid, using Lipofectamine 2000 (Thermo Fisher Scientific). Cells were harvested 3 days after transfection using QuickExtract DNA extraction solution according to the manufacturer's protocol and analyzed by surveyor assay or deep sequencing. To generate Cas12a-containing whole-cell lysate, 120,000 cells were seeded per well of a 24-well plate (Corning) 1

day before transfection. Each well was transfected with 500 ng Cas12a-encoding plasmid, and cell lysate was harvested 2 days after transfection.


*In vitro PAM identification assay.* The *in vitro* PAM identification assay was performed as described previously.<sup>18</sup> Briefly, whole-cell lysate from HEK293T cells overexpressing one of the Cas12a orthologs was prepared with lysis buffer (20 mM HEPES, 100 mM KCl, 5 mM MgCl<sub>2</sub>, 1 mM DTT, 5% glycerol, 0.1% Triton X-100) supplemented with EDTA-free Complete Protease Inhibitor Cocktail (Roche). CrRNA with corresponding direct repeat sequences were transcribed *in vitro* using custom oligonucleotides and a HiScribe T7 *in vitro* Transcription Kit (NEB) according to the manufacturer's recommended protocol for small RNA transcripts. The PAM library consisted of a pUC19 plasmid carrying a degenerate 8-bp sequence 5' of a 33-bp target site.<sup>7</sup> The library was pre-cleaved with XmnI and column purified prior to use (Qiagen). Each *in vitro* cleavage reaction consisted of 1 μl 10× CutSmart buffer (NEB), 200 ng PAM library, 500 ng *in vitro* transcribed crRNA, 10 μl cell lysate, and water for a total volume of 20 μl. Reactions were incubated at 37°C for 1 h and stopped by adding 500 μl buffer PB (Qiagen) followed by column purification. Purified DNA was amplified and sequenced using a MiSeq (Illumina) with a single-end 150-cycle kit. Sequencing results were entered into the PAM discovery pipeline.<sup>7</sup>

*Surveyor assay.* The surveyor assay was performed as previously described.<sup>19</sup> Briefly, genomic regions flanking a target site for each gene were amplified by PCR, and the products were purified using a QiaQuick Spin Column (Qiagen). Total purified PCR products (400 ng) were mixed with 2 μL 10 Taq DNA Polymerase buffer (Enzymatics) and ultrapure water to a final volume of 20 μL. Re-annealing was achieved by heating to 95°C for 2 min followed by a slow cool down to 10°C (~2.5°C per min). Re-annealed products were treated with surveyor nuclease (IDT) according to the manufacturer's protocol. Cleavage products were then visualized on 10% Novex TBE polyacrylamide gels (Life Technologies). Gels were stained with SYBR Gold DNA stain (Life Technologies) for 10 min and imaged with a Gel Doc imaging system (Bio-Rad).

*Deep Sequencing.* Targeted regions were amplified using a previously described two-step PCR protocol.<sup>19</sup> Indels were counted computationally as previously described.<sup>18</sup> Briefly, each amplicon was searched for exact matches within a 70-bp window around the cut site. For each sample, the indel rate was determined as (number of reads with indel) / (number of total reads). Samples with fewer than 1000 total reads were not included in subsequent analyses.

## Results

We selected 16 uncharacterized Cas12a-family proteins



**Fig. 1** Analysis of Cas12a ortholog diversity.


(A) Phylogenetic tree of 16 new Cas12a orthologs and 3 Cas12a orthologs with confirmed activity in eukaryotic cells (1-FnCas12a, 7-AsCas12a, and 13-LbCas12a). The approximate location of the RuvC subdomains and the nuclease (Nuc) domain are shaded in blue and pink respectively. (B) Alignment of direct repeat sequences of Cas12a orthologs. Sequences that are removed post crRNA maturation are shown in gray letters. Non-conserved bases are colored red. The stem duplex is shaded gray. (C) RNA secondary structures for mature crRNAs from 1-FnCas12a, 7-AsCas12a, 13-LbCas12a, and 23-Mb3Cas12a, predicted using Geneious 2 software.

with varying degrees of homology to three Cas12a orthologs (FnCas12a, AsCas12a, and LbCas12a)<sup>4,7</sup> with confirmed activity in eukaryotic cells (**Fig. 1A**). The direct repeat (DR) sequences of crRNAs associated with Cas12a orthologs show high levels of homology (**Fig. 1B**) and are predicted to fold into almost identical secondary structures (**Fig. 1C**). The homology is particularly strong at the stem structure and the AAUU motif (**Fig. 1C**), which is required for efficient crRNA maturation,<sup>9</sup> suggesting that the mechanism of crRNA maturation may be conserved within the Cas12a-family.

We performed a previously described *in vitro* assay<sup>18</sup> to determine the sequence of the PAM for each Cas12a ortholog (**Fig. 2A**). Of the 16 new Cas12a proteins, ten were active *in vitro* and recognized a T-rich PAM located 5' of the targeted sequence (**Fig. 2B**), just as previously characterized Cas12a proteins do.<sup>7</sup>

Next, we tested the 16 Cas12a orthologs for activity in human cells. We chose a previously validated target

within *VEGFA*, located next to a TTTG PAM that is permissive to all Cas12a orthologs. HEK293T cells were transfected with plasmids encoding humanized Cas12a orthologs together with PCR amplified fragments comprising a U6 promoter fused to the corresponding crRNA sequence (**Fig. 3A**). Four of the new Cas12a orthologs [*Thiomicrospira* sp. Xs5 (TsCas12a), *Moraxella bovoculi* AAX08\_00205 (Mb2Cas12a), *Moraxella bovoculi* AAX11\_00205 (Mb3Cas12a), and *Butyrivibrio* sp. NC3005 (BsCas12a)] were able to induce detectable indel events, as measured by surveyor nuclease assay (**Fig. 3B**). We tested these orthologs with six additional guides targeting either *DNMT1* or *EMX1* next to TTTV PAMs (**Fig. 3C**) and compared them to the activity of AsCas12a and LbCas12a. For all four new Cas12a enzymes, indel frequencies of >20% could be detected for at least two guides, but only Mb3Cas12a was able to induce robust indel levels with all six guides comparable to those of AsCas12a and LbCas12a. The apparent difference in activi-



**Fig. 2** PAM identification for Cas12a orthologs.


(A) Schematic for in vitro PAM screen. A library of plasmids bearing randomized 5' PAM sequences was cleaved by individual Cas12a nucleases and their corresponding crRNAs. Uncleaved plasmid DNA was PCR amplified and sequenced to identify depleted PAM sequences. (B) PAM sequences for ten Cas12a orthologs identified by in vitro PAM screens.

ties between Mb2Cas12a and Mb3Cas12a was somewhat surprising given that these orthologs share a predicted homology of 94.7%.

Because Mb3Cas12a was predicted to recognize a less restrictive PAM than the TTTV consensus PAM of AsCas12a and LbCas12a (**Fig. 2B**), we tested the ability of Mb3Cas12a to cleave endogenous DNA at TTN PAMs. To this end, we designed 64 guides: 16 guides for *DNMT1*, *EMX1*, *GRIN2b*, or *VEGFA*, targeting next to any combination of NTTN PAMs. To compare the activity of Mb3Cas12a, AsCas12a, and LbCas12a at NTTN PAMs, we transfected HEK293T cells with two plasmids, one expressing Cas12a and one expressing the crRNA, and assessed indel frequencies at each target site by deep sequencing. The average activity at TTTV PAMs was approximately 18% for Mb3Cas12a, 28% for AsCas12a, and 13% for LbCas12a (**Fig. 4A**). A few guides targeting next to NTTN PAMs (three for MbCas12a and one for AsCas12a) resulted in activity between 25–45% indels. However, whereas Mb3Cas12a performed better than AsCas12a and LbCas12a at NTTN PAMs, the average activity was relatively low with approximately 5.3% for Mb3Cas12a, approximately 2.7% for AsCas12a, and approximately 1.4% for LbCas12a. Comparing activity across all VTTN PAMs revealed statistically significant differences in indel activity between Mb3Cpf1 and LbCpf1 (mean 5.26% vs 1.38%,  $P = 0.0117$ ), but not between Mb3Cpf1 and AsCpf1 (mean 5.26% vs 2.69%,  $P = 0.1343$ ).

Based on the in vitro PAM screen, Mb3Cas12a tolerates Cs or Ts within its PAM. To assess the tolerance for Cs at position 2 and 3 of the Mb3Cas12a PAM, we used 18 guides targeting *DNMT1* or *EMX1* next to RTTN and NYYN PAMs (R=A or G, Y=C or T). HEK293T cells were transfected with a single plasmid expressing Mb3Cas12a and crRNA. The activity of each guide was determined using the surveyor nuclease assay. Guides targeting next to RTTN, RCTN, and RTCN PAMs had an average activity of approximately 15%, approximately 9%, and approximately 4%, respectively; however, guides targeting next to RCCN PAMs were mostly inactive (**Fig. 4B**). Taken together, our data show that Mb3Cas12a is active in human cells and shows robust activity at TTTV PAMs at levels comparable to those of AsCas12a and LbCas12a. Furthermore, Mb3Cas12a can reliably target sites with RTTV PAMs, albeit with lower overall activity.





**Fig. 3** Activity of Cas12a orthologs in human cells.

(A) Sixteen human codon-optimized Cas12a orthologs were expressed in HEK293T cells using CMV-driven expression vectors. The corresponding crRNA was expressed from PCR amplified fragments containing a U6 promoter fused to the crRNA sequence. NLS, nuclear localization signal. (B) Comparison of in vitro activity using a pre-validated guide targeting *VEGFA* next to a TTTV (V=A, C, or G) PAM. Indel frequencies were detected by surveyor assay. Red triangles indicate cleaved fragments. The percent indel frequencies are the averages of three bioreplicates. (C) The activities of four new Cas12a orthologs compared to AsCas12a and LbCas12a using six guides targeting either *EMX1* or *DNMT1*. Each data point represents one guide; indel frequencies were determined by surveyor assay and are shown as the means of all guides with SEMs.

## Discussion

Reaching the full potential of CRISPR-based genome editing will require a suite of tools to ensure that there are optimal enzymes for a range of genomic contexts. This will be particularly important for the therapeutic deployment of CRISPR, where the target site will be constrained by the genetic variations found in individual patients. Moreover, to tackle the full landscape of pathogenic mutations, a number of different gene editing strategies will be needed beyond simple gene knockout. Consequently, having an array of Cas enzymes that can be used in human cells is essential for the continued development of this technology.

Here we examined 16 new Cas12a family proteins for potential use in genome editing. Four of these, *Thiomicrospira* sp. Xs5 (TsCas12a), *Moraxella bovoculi* AAX08\_00205 (Mb2Cas12a), *Moraxella bovoculi* AAX11\_00205 (Mb3Cas12a), and *Butyrivibrio* sp. NC3005 (BsCas12a), exhibited activity in human cells. We chose HEK293 cells as a model for exploring these new Cas12a orthologs because there is a wealth of published data available on the efficiencies of other Cas enzymes in these cells. Previously, we observed only weak activity of FnCas12a in mammalian cells.<sup>7</sup> However, a recent study found that FnCas12a exhibits robust activity in plant cells,<sup>4</sup> indicating that Cas12a orthologs might have different activities depending on the organism. Therefore,


it may be informative in future studies to test the activities of these new Cas12a orthologs in different cell types and organisms.

Further analysis of the PAM requirements of the most active new ortholog, Mb3Cas12a, showed that it has a less restricted PAM (TTV) than AsCas12a and LbCas12a, which are active only at the canonical TTTV PAM. Alignment of Mb3Cas12a to other Cas12a orthologs did not suggest any immediate reason for the more relaxed PAM (data not shown), and further work will be required to investigate the structural basis for this altered PAM requirement.

Given the advantageous properties of Cas12a, such as its inherent high specificity and distinct PAM preference, this family of enzymes represents a powerful addition to the gene editing toolbox. Here, we further expanded the utility of Cas12a by identifying new orthologs that are active in human cells.

## Author Contributions

B.Z., J.S., and F.Z. conceived this study. B.Z. and J.S. performed the experiments with help from all authors. O.A. and J.G. analyzed PAM detection data. D.S. contributed to computational analysis of Cas12 orthologs. F.Z. supervised the research. B.Z. and F.Z. wrote the manuscript with input from all authors.



**Fig. 4** Evaluation of activity with relaxed PAM sequences.

(A) Mb3Cas12a, AsCas12a, and LbCas12a were tested for recognition of NTTN PAMs using four guides per PAM, targeting four different genes (*DNMT1*, *EMX1*, *GRIN2b*, or *VEGFA*). Indel frequencies were determined by deep sequencing. Each data point represents the average of three bioreplicates for one guide. Data are shown as means with SEMs. (B) Mb3Cas12a was tested with 18 guides targeting either *DNMT1* or *EMX1* next to a RTTN and NYYN PAM (R=A or G, Y=C or T). Each data point represents one guide; data are shown as means with SEMs.

### Acknowledgments

We thank R. Macrae, R. Belliveau, G. Faure, and L. Gao for discussions and support. J.S. is supported by the Human Frontier Science Program. F.Z. is a New York Stem Cell Foundation–Robertson Investigator. F.Z. is supported by National Institutes of Health grants (1R01-HG009761, 1R01-MH110049, and 1DP1-HL141201); the Howard Hughes Medical Institute; the New York Stem Cell, Edward Mallinckrodt, Jr., and G. Harold and Leila Mathers Foundations; the Poitras Center for Psychiatric Disorders Research at MIT; the Hock E. Tan and K. Lisa Yang Center for Autism Research at MIT; J. and P. Poitras; and the Phillips Family. F.Z. is a co-founder and advisor of Beam Therapeutics, Editas Medicine, Arbor Biotechnologies, Sherlock Biosciences, and Pairwise Plants. The authors plan to make the reagents widely available to the academic community through Addgene and to provide software tools via the Zhang lab website (zlab.bio). A patent has been filed relating to the presented data.

### Conflicts of Interest

The authors have declared that no conflict of interest exists.

### References

- Zhang, F. Development of CRISPR-Cas systems for genome editing and beyond. DOI:10.1017/S0033583519000052 52, 653 (2019).
- Cong L, Ran FA, Cox D, Lin S, Barretto R, Habib N, Hsu PD, Wu X, Jiang W, Marraffini LA, Zhang F: Multiplex genome engineering using CRISPR/Cas systems. *Science* 2013; **339**: 819–823. PMID:23287718, DOI:10.1126/science.1231143
- Mali P, Yang L, Esvelt KM, Aach J, Guell M, DiCarlo JE, Norville JE, Church GM: RNA-guided human genome engineering via Cas9. *Science* 2013; **339**: 823–826. PMID:23287722, DOI:10.1126/science.1232033
- Endo A, Masafumi M, Kaya H, Toki S: Efficient targeted mutagenesis of rice and tobacco genomes using Cpf1 from *Francisella novicida*. *Sci Rep* 2016; **6**: 38169. PMID:27905529, DOI:10.1038/srep38169
- Kim Y, Cheong SA, Lee JG, Lee SW, Lee MS, Baek IJ, Sung YH: Generation of knockout mice by Cpf1-mediated gene targeting. *Nat Biotechnol* 2016; **34**: 808–810. PMID:27272387, DOI:10.1038/nbt.3614
- Ma S, Liu Y, Liu Y, Chang J, Zhang T, Wang X, Shi R, Lu W, Xia X, Zhao P, Xia Q: An integrated CRISPR *Bombyx mori* genome editing system with improved efficiency and expanded target sites. *Insect Biochem Mol Biol* 2017; **83**: 13–20. PMID:28189747, DOI:10.1016/j.ibmb.2017.02.003
- Zetsche B, Gootenberg JS, Abudayyeh OO, Slaymaker IM, Makarova KS, Essletzbichler P, Volz SE, Joung J, van der Oost J, Regev A, Koonin EV, Zhang F: Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. *Cell* 2015; **163**: 759–771. PMID:26422227, DOI:10.1016/j.cell.2015.09.038
- Zetsche B, *et al*: Multiplex gene editing by CRISPR–Cpf1 using a single crRNA array. *Nat Biotechnol* 2016. PMID:27918548
- Fonfara I, Richter H, Bratović M, Le Rhun A, Charpentier E: The CRISPR-associated DNA-cleaving enzyme Cpf1 also processes precursor CRISPR RNA. *Nature* 2016; **532**: 517–521. PMID:27096362, DOI:10.1038/nature17945
- Wang M, Mao Y, Lu Y, Tao X, Zhu J: Multiplex gene editing in rice using the CRISPR-Cpf1 system. *Mol Plant* 2017; **10**: 1011–1013. PMID:28315752, DOI:10.1016/j.molp.2017.03.001
- Chen JS, Ma E, Harrington LB, Da Costa M, Tian X, Palefsky JM, Doudna JA: CRISPR-Cas12a target binding unleashes indiscriminate single-stranded DNase activity. *Science* 2018; **360**: 436–439. PMID:29449511, DOI:10.1126/science.aar6245
- Li SY, Cheng QX, Liu JK, Nie XQ, Zhao GP, Wang J: CRISPR-Cas12a has both cis- and trans-cleavage activities on single-stranded DNA. *Cell Res* 2018; **28**: 491–493. PMID:29531313, DOI:10.1038/s41422-018-0022-x
- Gootenberg JS, Abudayyeh OO, Kellner MJ, Joung J, Collins JJ, Zhang F: Multiplexed and portable nucleic acid detection platform with Cas13, Cas12a, and Csm6. *Science* 2018; **360**: 439–444. PMID:29449508, DOI:10.1126/science.aaq0179

14. Li SY, Cheng QX, Wang JM, Li XY, Zhang ZL, Gao S, Cao RB, Zhao GP, Wang J: CRISPR-Cas12a-assisted nucleic acid detection. *Cell Discov* 2018; **4**: 20. PMID:29707234, DOI:10.1038/s41421-018-0028-z
15. Kleinstiver BP, Tsai SQ, Prew MS, Nguyen NT, Welch MM, Lopez JM, McCaw ZR, Aryee MJ, Joung JK: Genome-wide specificities of CRISPR-Cas Cpf1 nucleases in human cells. *Nat Biotechnol* 2016; **34**: 869–874. PMID:27347757, DOI:10.1038/nbt.3620
16. Strecker J, Jones S, Koopal B, Schmid-Burgk J, Zetsche B, Gao L, Makarova KS, Koonin EV, Zhang F: Engineering of CRISPR-Cas12b for human genome editing. *Nat Commun* 2019; **10**: 212. PMID:30670702, DOI:10.1038/s41467-018-08224-4
17. Ran FA, Cong L, Yan WX, Scott DA, Gootenberg JS, Kriz AJ, Zetsche B, Shalem O, Wu X, Makarova KS, Koonin EV, Sharp PA, Zhang F: In vivo genome editing using *Staphylococcus aureus* Cas9. *Nature* 2015; **520**: 186–191. PMID:25830891, DOI:10.1038/nature14299
18. Gao L, Cox DB, Yan WX, Manteiga JC, Schneider MW, Yamano T, Nishimasu H, Nureki O, Crosetto N, Zhang F: Engineered Cpf1 variants with altered PAM specificities. *Nat Biotechnol* 2017; **35**: 789–792. PMID:28581492, DOI:10.1038/nbt.3900
19. Hsu PD, Scott DA, Weinstein JA, Ran FA, Konermann S, Agarwala V, Li Y, Fine EJ, Wu X, Shalem O, Cradick TJ, Marraffini LA, Bao G, Zhang F: DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat Biotechnol* 2013; **31**: 827–832. PMID:23873081, DOI:10.1038/nbt.2647