



United States  
Department of  
Agriculture

Forest Service

Southern Forest  
Experiment Station


New Orleans,  
Louisiana

General Technical Report  
SO-86



# A Synopsis of the Taxonomic Revisions in the Genus *Ceratocystis* Including a Review of Blue-Staining Species Associated with *Dendroctonus* Bark Beetles

Thelma J. Perry



## SUMMARY

Taxonomic revisions in both the teleomorphic (sexual) and anamorphic (asexual) forms of the genus *Cerutocystis* Ellis & Halstead are chronicled in this review. Recognized species associated with *Dendroctonus* Erichson bark beetles are summarized, and several species that have been published as recombinations, species that were previously described as *Ophiostoma*, and species that have not yet been recombined are listed.

# A Synopsis of the Taxonomic Revisions in the Genus *Ceratocystis* Including a Review of Blue-Staining Species Associated with *Dendroctonus* Bark Beetles

Thelma J. Perry

## INTRODUCTION

Because the fungi of the genera *Ceratocystis* Ellis & Halstead (1890) and *Ophiostoma* H. & P. Sydow (1919) are wood stainers and plant pathogens, they are economically important. Among the best known are the blue stain fungus *Ophiostoma minus* (Hedgc.) H. & P. Sydow ( $\equiv$  *Ceratocystis minor* [Hedgc.] Hunt), the oak wilt fungus *Ceratocystis fagacearum* (Bretz) Hunt, the Dutch elm disease fungus *Ophiostoma ulmi* (Buism.) Nannfeldt ( $\equiv$  *Ceratocystis ulmi* [Buism.] C. Moreau), the sweet potato rot fungus *Ceratocystis fimbriata* Ellis & Halstead, and the root pathogen *Leptographium wageneri* (Kendr.) Wingfield ( $\equiv$  *Verticicladiella Wageneri*: Kendrick).

Insects disperse these fungi's spores that are produced in a sticky substance by both the teleomorphs (sexual form) and the anamorphs (asexual form) (table 1) and readily adhere to beetle body parts. Most Scolytidae (bark beetles) transport one or more *Ceratocystis* species. These fungi are found in or near bark beetle galleries. Some species may assist the beetles as tree drying agents (Bramble and Holst 1940; Nelson 1934) or may elicit tree defensive reactions (Mattson and others, 1988). Some *Ceratocystis* species are ambrosia fungi, which serve as food for developing beetle larvae (Whitney 1971, 1982). Other species compete with immature beetles for space and nutrients in the phloem (Barras 1970; Bridges and Perry 1985; Franklin 1970).

The fungal-bark-beetle-host associations have been listed in publications such as Barras and Perry (1975), Upadhyay (1981), Whitney (1982), and Beaver (1989). Because of recent taxonomic revisions of the genus *Ceratocystis* in both the anamorphic and teleomorphic forms, there is a need for an updated bibliography of fungal-insect-host interactions. Harrington (1988) lists 34 *Leptographium* Lagerberg & Melin species, 20 of which have a known *Ophiostoma* teleomorph, and 30 species of bark beetles that vector these *Ophiostoma* I *Leptographium* fungi. Literature searches for bark beetle-fungal associations should

not be limited to the keyword *Ceratocystis* but should also include the fungi *Ophiostoma* and *Ceratocystiopsis* Upadhyay & Kendrick.

Because of the disagreements about the taxonomic status of the genus *Ceratocystis*, various reclassifications have contributed to the use of incorrect names in the fungal-bark beetle literature. My objectives are (1) to chronicle the events that led to revisions in both the teleomorphic genera and the anamorphic form-genera that make up the genus *Ceratocystis* and (2) to provide a quick and easy guide to the accepted taxonomic names of these fungi.

Most of the recent taxonomic disagreements occur at the level or order, family, and genus (fig. 1). In 1980, Benny and Kimbrough described the accepted order as Ophiostomatales with four genera: *Ophiostoma*, *Ceratocystis*, *Ceratocystiopsis*, and *Sphaeronaemella* Karsten. In 1932, Nannfeldt had erected the family Ophiostomataceae for the genus *Ophiostoma*. This is the traditionally accepted family for the genus *Ceratocystis* and the synonymized genera *Europhium* Parker and *Ophiostoma*. Von Arx and van der Walt (1988) accepted the order Ophiostomatales with the family Ophiostomataceae representing the genera *Ceratocystiopsis*, *Europhium*, and *Ophiostoma*. They accepted the anamorphic form-genus *Raffaelea* v. Arx & Hennebert and 16 other form-genera that are described in Upadhyay (1981) (table 2), with the exception of *Verticicladiella* S. Hughes, which is synonymized as *Leptographium* (Wingfield 1985). In 1988, Wingfield and others described *Knoxdaviesia*, a new anamorph of the genus *Ceratocystiopsis*. Von Arx and van der Walt (1988) established the family Pyxidiophoraceae for species with the anamorph *Chalara* (Corda) Rabenhorst, representing the genera *Ceratocystis* Ellis & Halstead, *Cryptendoxyla* Malloch & Cain, *Mycorhynchidium* Malloch & Cain, and *Pyxidiophora* Brefeld & Tavel.

At the generic level, three genera are currently accepted: (1) *Ceratocystis* Ellis & Halstead, containing only those species with the form-genus *Chalara* (Corda), Rabenhorst; (2) *Ophiostoma* H. & P. Sydow;

Thelma J. Perry is a microbiologist at Forest Insect and Disease Research, U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station, Pineville, LA 71360.

Table 1.—*Dendroctonus* species (Wood 1982) and associated blue-staining fungi


<i>Dendroctonus</i> sp.	Fungal associate	Publication
<i>Dendroctonus adjunctus</i> Blandford = <i>D. convexifrons</i> Hopkins	<i>Ophiostoma stenoceras</i> (Robak) Melin & Nannfeldt ≡ <i>Ceratocystis stenoceras</i> (Robak) C. Moreau = <i>Ceratocystis gossypina</i> var. <i>robusta</i> Davidson ≡ <i>Ceratostomella stenoceras</i> Robak = <i>Ceratocystis gossypina</i> Davidson <i>Ophiostoma adjuncti</i> (Davids.) Harrington ≡ <i>Ceratocystis adjuncti</i> Davidson <i>Leptographium pyrinum</i> Davidson <i>Leptographium</i> sp. Lagerberg & Melin	Davidson 1971  Davidson 1978 Davidson 1978 Harrington 1988
<i>Dendroctonus brevicomis</i> Le Conte	<i>Ophiostoma minus</i> H. & P. Sydow ≡ <i>Ceratocystis minor</i> (Hedgc.) Hunt ≡ <i>Ceratostomella minor</i> Hedgcock = <i>Ceratostomella pini</i> Munch = <i>Ceratostomella exigua</i> Hedgcock = <i>Ceratostomella pseudotsugae</i> Rumbold <i>Ophiostoma nigrocarpum</i> (Davids.) de Hoog ≡ <i>Ceratocystis nigrocarpa</i> Davidson	Mathre 1964  Rumbold 1931 Rumbold 1936 Davidson 1966
<i>Dendroctonus frontalis</i> Zimmermann	<i>Ophiostoma minus</i> H. & P. Sydow ≡ <i>Ceratocystis minor</i> (Hedgc.) Hunt ≡ <i>Ceratostomella minor</i> Hedgcock = <i>Ceratostomella pini</i> Munch = <i>Ceratostomella exigua</i> Hedgcock = <i>Ceratostomella pseudotsugae</i> Rumbold <i>Ophiostoma nigrocarpum</i> (Davids.) de Hoog ≡ <i>Ceratocystis nigrocarpa</i> Davidson <i>Ceratocystiopsis ranaculosus</i> Perry & Bridges	Rumbold 1931 Rumbold 1936 Davidson 1966 Bridges & Perry 1987
<i>Dendroctonus jeffreyi</i> Hopkins	<i>Ophiostoma ips</i> (Rumb.) Nannfeldt ≡ <i>Ceratocystis ips</i> (Rumb.) C. Moreau = <i>Ceratostomella montia</i> Rumbold	Mathre 1964 Rumbold 1941
<i>Dendroctonus ponderosae</i> Hopkins = <i>D. monticolae</i> Hopkins	<i>Ophiostoma huntii</i> (Robins.-Jeff.) de Hoog & Scheffer ≡ <i>Ceratocystis huntii</i> Robinson-Jeffrey & Grinchenko <i>Ophiostoma minus</i> H. & P. Sydow ≡ <i>Ceratocystis minor</i> (Hedgc.) Hunt ≡ <i>Ceratostomella minor</i> Hedgcock = <i>Ceratostomella pini</i> Munch = <i>Ceratostomella exigua</i> Hedgcock = <i>Ceratostomella pseudotsugae</i> Rumbold <i>Ophiostoma ips</i> (Rumb.) Nannfeldt ≡ <i>Ceratocystis ips</i> (Rumb.) C. Moreau = <i>Ceratostomella montia</i> Rumbold <i>Ophiostoma piliferum</i> (Fries) H. & P. Sydow ≡ <i>Ceratocystis pilifera</i> (Fries) C. Moreau = <i>Sphaeria pilifera</i> Fries = <i>Ceratocystis shrenkiana</i> (Hedgc.) C. Moreau <i>Ophiostoma clavigerum</i> (Robins.-Jeff. & Davids.) Harrington = <i>Ceratocystis clavigem</i> (Robins.-Jeff. & Davids.) Upadhyay ≡ <i>Europhium clavigerum</i> Robinson-Jeffrey & Davidson <i>Ceratocystiopsis minuta</i> (Siem.) Upadhyay & Kendrick ≡ <i>Ceratocystis minuta</i> (Siem.) Hunt ≡ <i>Ophiostoma minutum</i> Siemaszko <i>Leptographium</i> Lagerberg & Melin	Robinson-Jeffrey & Grinchenko 1964  Rumbold 1931 Rumbold 1936 Mathre 1964 Rumbold 1941  Mathre 1964 Robinson-Jeffrey & Davidson 1964 Mathre 1964 Harrington 1988, Robinson 1962

Table 1.—*Dendroctonus* species (Wood 1982) and associated blue-staining fungi—Continued

<i>Dendroctonus</i> sp.	Fungal associate	Publication
<i>Dendroctonus</i> <i>pseudotsugae</i> Hopkins	<i>Ophiostoma minus</i> H. & P. Sydow	Mathre 1964
	≡ <i>Ceratocystis minor</i> (Hedgc.) Hunt	
	≡ <i>Ceratostomella minor</i> Hedgcock	Rumbold 1936
	= <i>Ceratostomella exigua</i> Hedgcock	
	= <i>Ceratostomella pini</i> Munch	Rumbold 1936
	= <i>Ceratostomella pseudotsugae</i> Rumbold	Wingfield 1983
	<i>Leptographium abietinum</i> (Peck) Wingfield	Harrington 1988
	≡ <i>Verticicladiella abietina</i> (Peck) S. Hughes	
	= <i>Sporocybe abietina</i> Peck	
	<i>Leptographium</i> sp. A	
<i>Dendroctonus</i> <i>rufipennis</i> (Kirby) = <i>D. engelmanni</i> Hopkins = <i>D. piceaperda</i> Hopkins	<i>Ophiostoma bicolor</i> Davidson & Wells	Davidson 1955
	≡ <i>Ceratocystis bicolor</i> (Davids. & Wells) Davidson	Davidson 1958
	<i>Ophiostoma piceaperda</i> (Rumb.) von Arx	Mathre 1964
	≡ <i>Ceratocystis piceaperda</i> (Rumb.) C. Moreau	
	≡ <i>Ceratostomella piceaperda</i> Rumbold	Rumbold 1936
	= <i>Ceratocystis europioides</i> Wright & Cain	Wright & Cain 1961
	<i>Ophiostoma penicillatum</i> (Gros.) Siemaszko	Davidson 1955
	≡ <i>Ceratocystis penicillata</i> (Gros.) C. Moreau	
	≡ <i>Ceratostomella penicillata</i> Grossmann	
	= <i>Ophiostoma truncicola</i> Davidson	
	<i>Ophiostoma coerulescens</i> (Munch) Nannfeldt	Davidson, 1955
	≡ <i>Ceratocystis coerulescens</i> (Munch) Bakshi	
	≡ <i>Endoconidiophora coerulescens</i> Munch	
<i>Leptographium abietinum</i> (Peck) Wingfield	Wingfield 1983, Harrington 1988	
≡ <i>Verticicladiella abietina</i> (Peck) S. Hughes		
= <i>Sporocybe abietina</i> Peck		
<i>Leptographium engelmanni</i> Davidson	Davidson 1955	
<i>Dendroctonus</i> <i>terebrans</i> (Olivier)	<i>Ophiostoma ips</i> (Rumb.) Nannfeldt	Rane & Tatter 1987
	≡ <i>Ceratocystis ips</i> (Rumb.) C. Moreau	
	= <i>Ceratostomella montia</i> Rumbold	
	<i>Leptographium procerum</i> (Kendr.) Wingfield	Wingfield 1983, Harrington 1988
	≡ <i>Verticicladiella procerum</i> Kendrick	
<i>Leptographium terebrantis</i> Barras & Perry	Barras & Perry 1971	
<i>Dendroctonus</i> <i>valens</i> Le Conte	<i>Ophiostoma piliferum</i> (Fries) H. & P. Sydow	Goheen & Cobb 1978
	≡ <i>Ceratocystis pilifera</i> (Fries) C. Moreau	
	= <i>Sphaeria pilifera</i> Fries	
	<i>Ophiostoma wagneri</i> (Goheen & Cobb) Harrington	Goheen & Cobb 1978
	≡ <i>Ceratocystis wagneri</i> Goheen & Cobb	
	<i>Ophiostoma ips</i> (Rumb.) Nannfeldt	Rumbold 1931
	≡ <i>Ceratocystis ips</i> (Rumb.) C. Moreau	
	= <i>Ceratostomella montia</i> Rumbold	Wingfield 1983
	<i>Leptographium terebrantis</i> Barras & Perry	Rumbold 1931
	<i>Ophiostoma piceaperdum</i> (Rumb.) von Am	
	≡ <i>Ceratocystis piceaperda</i> (Rumb.) C. Moreau	
≡ <i>Ceratostomella piceaperda</i> Rumbold		
= <i>Ceratocystis europioides</i> Wright & Cain	Wright & Cain 1961	
<i>Dendroctonus</i> Erichson	<i>Ceratocystis leucocarpa</i> Davidson*	Davidson 1966
	<i>Ophiostoma aureum</i> (Robins-Jeff. & Davids.) Harrington	Robinson-Jeffrey & Davidson 1968
	≡ <i>Ceratocystis aurea</i> (Robins-Jeff. & Davids.) Upadhyay	
	≡ <i>Europhium aureum</i> Robinson-Jeffrey & Davidson	
	<i>Ophiostoma robustum</i> (Robins-Jeff. & Davids.) Harrington	Robinson & Davidson 1968
	≡ <i>Ceratocystis robusta</i> (Robins-Jeff. & Davids.) Upadhyay	
≡ <i>Europhium robustum</i> Robinson-Jeffrey & Davidson		

\*This species has not been examined or published as *Ophiostoma*.

### Benny & Kimbrough (1980)



### Von Arx & Van Der Walt (1988)




Figure 1 — Synopsis of the most recent classification schemes of these genera.

Table 2.—*Form-genera with asexual conidiation, Nag Raj and Kendrick (1975), Upadhyay (1981), Upadhyay and Kendrick (1975), Wingfield (1985)*

Exogenous, Holoblastic	
<i>Acremonium</i> Link ex Fries	Simple, reduced conidiophores
<i>Allescheriella</i> Henn.	Nonspecific, hyaline
<i>Hyalodendron</i> Diddens	Acropetal chains, simple, hyaline
<i>Hyalopesotum</i> Upadhyay & Kendrick	Anellidic, synnematos, hyaline
<i>Hyalorhinocladia</i> Upadhyay & Kendrick	Mononematous, sympodial, hyaline
<i>Gabarnaudia</i> Samson & W. Gams	Mononematous, phialidic, hyaline
<i>Graphilbum</i> Upadhyay & Kendrick	Synnematous, sympodial, hyaline
<i>Graphium</i> Corda	Synnematous, annellidic, pigmented
<i>Graphiocladiella</i> Upadhyay	Mononematous, annellidic, hyaline
<i>Knoxdaviesia</i> Wingfield, Van Wyk & Marassas	Mononematous, phialidic, hyaline
<i>Leptographium</i> Lagerb. & Melin	Mononematous, annellidic, pigmented
<i>Pachnodium</i> Upadhyay & Kendrick	Nonspecific
<i>Pesotum</i> Crane & Schoknecht	<b>Synnematous</b> , sympodial, pigmented
<i>Sporothrix</i> Hekt. & Perkins ex. Nicot & Mariat	Simple, sympodial, hyaline
<i>Verticillidiella</i> S. Hughes = <i>Leptographium</i> , Wingfield	Mononematous, sympodial and annellidic, pigmented
Endogenous, Enteroblastic	
<i>Chalara</i> (Corda) Rabenh.	Phialidic, simple, hyaline
<i>Chalaropsis</i> Peyronel = <i>Chalara</i>	
<i>Phialocephala</i> Kendrick	Phialidic, mononematous, hyaline
<i>Phialoglyphium</i> Upadhyay & Kendrick	Phialidic, synnematos, pigmented
<i>Thielaoiopsis</i> Went = <i>Chalara</i>	

and (3) *Cerutocystiopsis* Upadhyay & Kendrick for the other form-genera (table 2).

## DISCUSSION

When both the teleomorph and the anamorph are known, the fungus is called a holomorph (the whole fungus). The teleomorph, which reproduces with ascospores, is the sexual form of the fungus. It may have one (mono-anamorphic) or many (pleo-anamorphic) asexual stages. The anamorph, or asexual form, is placed into a form-genus. Form-genera, formerly known as fungi imperfecti, are the asexual, conidial-producing forms of a fungus. Form-genera are classified according to the method of conidia formation and conidiophore morphology (conidiogenesis) and may or may not be associated with a teleomorph. In the Ophiostomataceae, conidiogenesis may be **holoblastic** (when both inner and outer walls contribute to the formation of conidia) or enteroblastic (when the inner wall or neither wall contributes to the formation of conidia). Older terms for holoblastic conidiogenesis are exoconidial and exogenous; for enteroblastic, endoconidial and endogenous. These terms are defined in Kendrick (1971), Hennebert and Weresub (1977), and Ainsworth and Bisby's *Dictionary of the Fungi*

(Hawksworth and others 1983). In *The Whole Fungus* (Kendrick 1979), noteworthy scientists have written stimulating discussions of the history, conidial formation, terms, and relative importance of anamorphs as taxonomic features.

Scientists have made taxonomic revisions in both the teleomorphs and the anamorphs of the genus *Cerutocystis*. Since cultural and microscopic analyses alone are not always sufficient to delineate genus and species characteristics, new methods of **ultrastructural**, chemical, biochemical, immunological, and genetic analyses are being used to realign the genus *Cerutocystis* and its anamorphs.

More complete historical data on this genus can be found in Bakshi (1951), Hunt (1956), and Upadhyay (1981). Over the years, several genera have been synonymized and placed in the genus *Cerutocystis*. Bakshi (1951) synonymized six genera and the species *Endoconidiophora coerulescens* Munch and *Ophiostoma coerulescens* (Munch) Nannfeldt as *Cerutocystis coerulescens* (Munch) Bakshi. In this enteroblastic species, spores (conidia) are produced within the spore-bearing structure (conidiophore). He also described *Cerutocystis galeiformis*, a holoblastic species where spores were formed at the "end of hypha" or conidiophore (*Cephalosporium* Corda) or at the "ends of the collection of hyphae" or conidiophores

(*Graphium* Corda and *Leptographium*). Ellis and Halstead's (1890) generic concept of *Ceratocystis* was then changed to accommodate species that produced both enteroblastic and holoblastic spores. Hunt (1956) accepted Bakshi's (1951) changes, synonymized two more genera, redescribed the genus *Ceratocystis*, completed the transfer of synonymized species, made new combinations, and described new species.

In 1974, de Hoog proposed the resurrection of the genus *Ophiostoma* with the type species *O. piliferum* for species producing holoblastic conidia. Von Hohnel (1918) originally described the genus as *Linostoma*, an invalid name, which Sydow and Sydow (1919) renamed *Ophiostoma*, and Bakshi (1951) synonymized with the genus *Ceratocystis*.

Melin and Nannfeldt (1934) divided the genus *Ophiostoma* into sections to accommodate species that produced their spores inside the conidial bearing structure (enteroblastic) and to accommodate the holoblastic species. De Hoog used Nannfeldt's sections in Melin and Nannfeldt (1934) to divide the two genera based on the type of conidia formed, either enteroblastic or holoblastic. In 1975, Weijman and de Hoog divided the genus *Ceratocystis* as *Ceratocystis sensu stricto* (in the strict sense) to contain all species with enteroblastic conidiogenesis and *Ceratocystis sensu lato* (in the wide sense) as the *Ophiostoma* group for holoblastic species. In 1975, Upadhyay and Kendrick proposed to split the genus *Ceratocystis* and to establish the genus *Ceratocystiopsis* for those species with falcate ascospores and ascoma (perithecia) with short necks.

In 1981, Upadhyay published a monograph of the *Ceratocystis* species, synonymized *Ophiostoma* and *Europhium*, and transferred 13 of 15 *Ceratocystis* species to *Ceratocystiopsis*. In 1984, de Hoog and Scheffer established *Ophiostoma* as a valid genus characterized by having cellulose, lignin, and rhamnose in their cell walls; growing on cycloheximide media; and producing holoblastic conidia. The genus *Ceratocystis* was limited to species with enteroblastic *Chalara* anamorphs and those that were cellulose negative and sensitive to cycloheximide. De Hoog and Scheffer accepted Upadhyay and Kendrick's genus *Ceratocystiopsis*.

The genus *Ceratocystis* now contains only those species having the enteroblastic form-genus *Chalara*, with the type species *C. fimbriata* Ellis & Halstead (table 3). *Ceratocystiopsis*, with the type species *C. minuta* (Siem.) Upadhyay & Kendrick, is an exception in that it does have an enteroblastic *Chalara* anamorph as well as holoblastic anamorphs. The holoblastic *Ophiostoma*, type species *O. piliferum*, has several form-genera of which two or more anamorphs may develop in culture. *Ophiostoma clavigerum* (Robins.-Jeff. & Davids.) Harrington (1987), when continually subcultured, has produced as many as six

Table 3.—*Ceratocystis* species with *Chalara* anamorphs, de Hoog (1974) and von Arx (1987)

de Hoog	
<i>Ceratocystis adiposa</i>	(Butler) C. Moreau
<i>C. autogmpa</i>	Bakshi
<i>C.!</i> <i>coerulescens</i>	(Munch) Bakshi
<i>C. fagacearum</i>	(Bretz) Hunt
<i>C. fimbriata</i>	Ellis & Halstead
<i>C. major</i>	(van Beyma) C. Moreau
<i>C. moniliformis</i>	(Hedgcock) C. Moreau
<i>C. musarum</i>	Riedl
<i>C. paradoxa</i>	(Dade) C. Moreau
<i>C. radicola</i>	(Bliss) C. Moreau
<i>C. uariospora</i>	(Davidson) C. Moreau
<i>C. virescens</i>	(Davidson) C. Moreau
von Arx	
<i>C. adiposum</i>	(Butler) C. Moreau
<i>C. coerulescens</i>	(Munch) Bakshi
<i>C. fagacearum</i>	(Bretz) Hunt
<i>C. fimbriata</i>	Ellis & Halstead
<i>C. paradoxa</i>	(Dade) C. Moreau

forms: *Graphiocladiella* Upadhyay, *Verticicladiella*, *Hyalorhinocladiella* Upadhyay & Kendrick, *Leptographium*, an annellidic yeast, and a holoblastic yeast (Tsuneda and Hiratsuka 1984).

DeHoog in Malloch (1979) said of the *Verticicladella-Phialographium* Upadhyay & Kendrick-*Leptographium* complex that older cultures degenerate and tend to form less well-structured, less complex, and less energy-consuming anamorphs.

Recognized form-genera now total 17 (table 2). Nag Raj and Kendrick (1975) synonymized *Chalaropsis* Peyronel and *Thielaviopsis* Went as *Chalara*. Upadhyay (1981) described 17 form-genera, but Wingfield (1985) reclassified *Verticicladiella* as *Leptographium*, and Wingfield and others (1988) described a new species, *Knoxdaviesia*. Harrington (1988) felt that most *Ophiostoma* anamorphs could probably be accommodated in the form-genera *Sporothrix* Hekt. & Perkins ex Nicot & Mariat, *Leptographium*, and *Graphium*. Form-genera of *Ceratocystis*, *Ceratocystiopsis*, and *Ophiostoma* need taxonomic study as recommended by the 1986 International Council of Botanical Names and by Kananaskis II, published as *The Whole Fungus* (Kendrick 1979).

Mycologists, plant pathologists, and botanists do not agree on the higher classifications of fungi. For instance, Ainsworth and Bisby's *Dictionary of the Fungi* (Hawksworth and others 1983) recognizes the Kingdom Fungi but does not recommend any one scheme for classes in the Subdivision Ascomycotina. Von Arx (197) advocated a limited number of orders and the omission of higher ranks. At an August 1990 meeting in Bad Windsheim, West Germany, specialists in the Ophiostomatales presented papers with startling opinions of this group of fungi (pers.



comm. Dr. M. Blackwell). Upadhyay would retain the family Ophiostomataceae in the order Microascales; Samuels would place Ophiostoma and Ceratocystis in the Order Sordariales and the Family Chaetosphaeriaceae; Wingfield would synonymize *Ceratocystiopsis* with *Ophiostoma*, except *Ceratocystiopsis falcata* (Wright & Cain) Upadhyay, which would reside in *Ceratocystis sensu stricto*.

Several authors discuss anamorphs; others discuss molecular methods, pathogenicity, vectors and dispersal, sapstain, tree response and host defense reactions, and chemistry. Kendrick<sup>1</sup> (Wingfield and others 1990) states that "since we do not yet have all this information (ultrastructure, development patterns, anamorph-teleomorph relationships, wall-chemistry, chemistry of slime, vectors and dispersal mechanisms, pathogenicity, and genetics) for any of the more than one-hundred members of the Ophiostomatales, it is clear that plenty of work remains to be done before we will properly understand this fascinating and important group of fungi."

Article 59 of the International Code of Botanical Nomenclature (Greuter and others 1983) states that the name of the holomorphic Ascomycete (the whole fungus, with both anamorph and teleomorph) takes precedence over the name of the anamorph. For example, one mycangial fungus of *Dendroctonus frontalis* Zimmermann, the southern pine beetle, is heterothallic (sexual reproduction requiring the union of two compatible mating types) and has been described as the anamorph *Sporothrix*<sup>2</sup>, but since the anamorphic name was not validly described, the fungus must be cited by the name of the teleomorph, *Ceratocystiopsis ranaculosus* Perry & Bridges (Bridges and Perry 1987, Harrington and Zambino 1990). Mycologists have agreed that anamorphs that have been validly described and published may be cited

separately, such as the anamorph *Sporothrix subannulata* Livingston & Davidson and its teleomorph *Ophiostoma subannulatum* Livingston & Davidson (Livingston and Davidson 1987).

A list of the *Ceratocystis* species that have been transferred (published as *Ophiostoma* species by new authors, table 4) can be found in this publication. If a described species of *Ceratocystis* having an anamorph other than *Chalara* is not listed as recombined in de Hoog and Scheffer (1984) and Harrington (1987, 1988) (examples 1, 2), then consult Upadhyay (1981) (example 3) for the original author of a *Ceratocystis* species that had been originally described as or transferred to an *Ophiostoma* species (table 5). More than a hundred described species of *Ceratocystis* exist. Some species were originally described as *Ophiostoma*;

Table 4.—*New and recombined Ophiostoma species by other authors*

---

<i>Ophiostoma ainoue</i> Solheim (1986)
<i>O. cucullulatum</i> Solheim (1986)
<i>O. davidsonii</i> (Olchowecki & Reid) Solheim (1986)
<i>O. europhioides</i> (Wright & Cain) Solheim (1986)
<i>O. flexuosum</i> Solheim (1986)
<i>O. longirostellatum</i> (Bakshi) v. Arx & Muller (1954)
<i>O. roboris</i> Georgescu & Teodoru (1948)
<i>O. sagmatospora</i> (Wright & Cain) Solheim (1986)
<i>O. subannulatum</i> Livingston & Davidson (1987)

---

Table 5.—*Ceratocystis species that were originally described as, or transferred to Ophiostoma, de Hoog (1974) and Upadhyay (1981)*

---

<i>Ophiostoma bicolor</i> Davidson & Wells
<i>O. brunneo-ciliatum</i> Mathieson-Kaarik
<i>O. ips</i> (Rumb.) Nannfeldt
<i>O. zeptographioides</i> (Davidson) v. Arx
<i>O. microsporium</i> (Davidson) de Hoog
<i>O. minus</i> (Hedgc.) H. & P. Sydow
<i>O. multiannulatum</i> (Hedgcock & Davidson) Hendrix
<i>O. narcissi</i> Limber
<i>O. nigrocurpum</i> (Davidson) de Hoog
<i>O. olivaceum</i> Mathieson
<i>O. penicillitum</i> Mathieson
<i>O. perfectum</i> (Davidson) de Hoog
<i>O. piceae</i> (Munch) H. & P. Sydow
<i>O. piceuperdum</i> (Rumbold) v. Arx
<i>O. piliferum</i> (Fries) H. & P. Sydow
<i>O. polonicum</i> Siemaszko
<i>O. rostrocyllindricum</i> (Davidson) v. Arx
<i>O. stenocerus</i> (Robek) Melin & Nannfeldt
<i>O. tetropii</i> Matheison
<i>O. triungulosporem</i> Butin
<i>O. ulmi</i> (Buisman) Nannfeldt

---

<sup>1</sup>Blackwell, Meredith. 1990. Personal communication with Dr. Blackwell, Department of Botany, Louisiana State University, Baton Rouge, LA 70803, in Sept. 1990, regarding "The Biology and Taxonomy of the Ophiostomatales", an international symposium, held in Bad Windsheim, West Germany, on August 21-24, 1990. Program and Abstracts organized by Wingfield, M.J., Department of Microbiology, University of the Orange Free State, P.O. Box 339, Bloemfontein 9300, Republic of South Africa; Seifert, K., Biosystematics Research Institute, Central Experiment Farm, William Saunders Building, Ottawa, Ontario, KIA OC6, Canada; Webber, J.F., Forest Research Station, Alice Holt Lodge, Wrecclesham, Farnham, Surrey, GU10 4LH, England. The proceedings will be published at a later date.

<sup>2</sup>Harrington, T.C. 1991. Personal communication with Dr. Harrington, Department of Plant Pathology, Iowa State University, Ames, IA 50011, who reviewed the manuscript in January, 1991, and disagreed with the taxonomic identification of this holomorph and stated that it needs further examination.

some species have been transferred to the genus *Ophiostoma*, and some have not yet been recombined (tables 6 & 7). Recombinations formed in the transfer of species must follow the rules of the International Code of Botanical Nomenclature, Article 33. The rules have been restated in the International Commission on the Taxonomy of Fungi (ICTF) Code of Practice for Systematic Mycologists (Sigler and Hawksworth 1987). Care must be taken to note the Latin or Greek endings of genera and species. For example, the genus *Ophiostoma* has a neuter ending, so the species ending must also be neuter.

Table 6.—*Ceratocystis* species not recombined, Upadhyay (1981)

---

<i>Ceratocystis acericola</i> Griffin
<i>C. aequiuaginata</i> Olchowecki & Reid
<i>C. allantospora</i> Griffin
<i>C. angusticollis</i> Wright & Cain
<i>C. arborea</i> Olchowecki & Reid
<i>C. brunneo-crinita</i> Wright & Cain
<i>C. California</i> Devay, Davidson and Moller
<i>C. columnaris</i> Olchowecki & Reid
<i>C. deltoideospora</i> Olchowecki & Reid
* <i>C. fimicola</i> (Marchal) Upadhyay
* <i>C. helvellae</i> (Karsten) Upadhyay
<i>C. hyalothecium</i> Davidson
<i>C. introcitrina</i> Olchowecki & Reid
<i>C. leucocarpa</i> Davidson
<i>C. magnifica</i> Griffin
<i>C. olivaceapini</i> Davidson
<i>C. populicola</i> Olchowecki & Reid
<i>C. pseudominor</i> Olchowecki & Reid
<i>C. stenospora</i> Griffin
<i>C. tenella</i> Davidson
<i>C. torticiliata</i> Olchowecki & Reid
<i>C. tubicollis</i> Olchowecki & Reid

---

\*Upadhyay synonymized these species, but Benny and Kimbrough (1980) and Cannon and Hawksworth (1982) recognize the genus *Sphaeronaemella* Karsten as valid.

Table 7.—*Ceratocystis* species not recombined by other authors

---

<i>Ceratocystis grandicarpa</i> Kowalski & Butin (1989)
<i>C. kubanicum</i> Scerbin-Parfenenko (1953)
<i>C. nothofagi</i> Butin (1984)
<i>C. novae-zelandiae</i> Hutchinson & Reid (1988)
<i>C. prolifera</i> Kowalski & Butin (1989)
<i>C. ualachium</i> Georgescu, Teodoru & Badea (1948)

---

## EXAMPLES

1. *Ophiostoma araucariae* (Butin) de Hoog & Scheffer  $\equiv$  *Ceratocystis araucariae* Butin.
2. *Ophiostoma abiocarpum* (Davids.) Harrington  $\equiv$  *Ceratocystis abiocarpa* Davidson
3. *Ophiostoma minus* (Hedgc.) H. & P. Sydow  $\equiv$  *Ceratocystis minor* (Hedgc.) Hunt.

The symbol " $\equiv$ " designates an obligate synonym. Both names are based on the same type specimen. They are nomenclatural synonyms. The symbol " $\equiv$ " designates a facultative synonym--names based on different type specimens (Hawksworth and others 1983); See table 1.

## SYNOPSIS OF RELEVANT PUBLICATIONS

- 1890 Ellis and Halstead, in Halstead (1890), established the genus *Ceratocystis* without a formal generic description.
- 1891 Halstead and Fairchild established the generic concept of the genus and provided the type species *C. fimbriata*, with an endogenous conidial form.
- 1919 Sydow and Sydow established the genus *Ophiostoma* for *Linostoma* von Hohnel, an invalid name. Von Hohnel (1918) characterized the genus *Linostoma* as having hyaline ascospores formed in evanescent asci and having fringes of ostiolar "cilia."
- 1932 Nannfeldt erected the family Ophiostomataceae for the genus *Ophiostoma*.
- 1934 Melin and Nannfeldt divided *Ophiostoma* into two sections. The species of section *Brevirostrata* Nannfeldt have short, conical perithecial necks, and the species assigned to section *Longirostrata* were characterized by longer filiform necks; Longirostrata was further divided to contain species with both endogenous conidia, *Chalara*-type (synonymized *Endoconidiophora* Munch), and exogenous conidia.
- 1951 Bakshi revived the generic name *Ceratocystis* for species with both endogenous and exogenous conidia. He regarded *Ophiostoma*, *Endoconidiophora*, *Rostrella* Zimmermann, *Linostoma* von Hohnel, and *Grosmania* G. Goidanich as synonyms. *Ceratostomella* Saccardo was retained as a separate genus. He described three new species and transferred the species *Endoconidiophora coeruleascens* Munch (Munch 1907) and *Ophiostoma coeruleascens* (Munch) Nannfeldt (Melin and

- Nannfeldt 1934) to the genus *Cerutocystis*, as *Cerutocystis coeruleascens* (Munch) Bakshi.
- 1956 Hunt published the first comprehensive monograph of *Cerutocystis* species, placed the genus in the order Plectascales, accepted Bakshi's emendations (that the genus contains species with both endogenous and exogenous conidia), and completed the transfer of species. He synonymized *Sphaeria* Haller ex Fries and *Sphaeronaemella* Karsten ex Seeler under *Cerutocystis*, and *Fugascus* Falck was listed as *nomen nudum* (a genus without adequate description). *Cerutostomella* was also excluded from synonymy. He presented a key to the species and sectioned them according to conidial development: section (1) species with an endoconidial imperfect stage, the Endoconidial Group; section (2) species with a *Leptographium* or *Graphium* imperfect state consisting of the *Grossmannia* Group and the *Ophiostoma* Group in part; and section (3) species with mycelial conidia only, the *Ophiostoma* Group in part.
- 1957 Parker established the genus *Europhium* with the type species *E. trinacriforme*. The species was distinguished from *Ceratocystis* by its neckless, closed ascocarps. Benny and Kimbrough (1980) and Upadhyay (1981) synonymized this genus with *Cerutocystis*; von Arx and van der Walt (1988) resurrected the genus and placed it in the family Ophiostomataceae.
- 1962 Kendrick redescribed the genus *Verticicladiella* S. Hughes, differentiated the genus *Leptographium* Lagerberg & Melin, placed three species in *Verticicladiella*, and described four new species, *V. procera*, *V. brachiata*, *V. antibiotica*, and *V. wagnerii*.
- 1964 Rosinski and Campana demonstrated both chitin and cellulose in the hyphal walls of *Ceratocystis ulmi* (Buisman) C. Moreau.
- 1965 Kendrick and Molnar described a new *Ceratocystis* with the first-described *Verticicladiella* anamorph.
- 1965 Rosinski confirmed the occurrence of cellulose in *C. ulmi*.
- 1967 Smith and others differentiated four species by their cell wall chemistry and divided them into two categories. The endoconidial (enteroblastic or endogenous) forms *Cerutocystis fagacearum* (Bretz) Hunt and *C. fimbriata* Ellis & Halstead have no cellulose in their cell walls, and the exoconidial (holoblastic or exogenous) forms of *C. olivacea* (Mathiesen) Hunt have both chitin and cellulose in their cell walls.
- 1968 Griffin published a monograph of 32 *Ceratocystis* species found in Ontario with descriptions, figures, distribution data, and a key to 60 species based on ascospore morphology. Eleven species were described as new.
- 1968 Robinson-Jeffrey and Davidson described three new species of *Europhium* (*clavigerum*, *aureum*, and *robustum*), which Upadhyay, in 1981, synonymized as *Ceratocystis* species including the type species *E. trinacriforme*. In 1987, Harrington recombined the four species as *Ophiostoma*.
- 1970 Von Arx urged' revision of the form-genus *Graphium* because conidia of described species were formed exogenously as well as endogenously. He cited the genus as *Ceratocystis* Ellis & Halstead with 50 species and accepted Bakshi's synonyms but did not accept Bakshi's (1951) or Hunt's (1956) emendations.
- 1971 Spencer and Gorin, to aid classification, grouped species of *Ceratocystis* and *Graphium* according to their polysaccharide components. The endoconidial (endogenous) group formed glucomannans or galactoglucomannans rather than rhamnose- and mannose-containing polysaccharides as in the exoconidial (exogenous) group.
- 1972 Cain suggested a relationship to filamentous yeastlike taxa as *Cephaloascus* Hanawa and said that *Verticicladiella* may be more closely related to hyphal yeast genera with hat-shaped ascospores than to phialidic *Chalara*-like anamorphs.
- 1973 Muller and von Arx placed the genus *Ceratocystis* in the order Sphaeriales of the Pyrenomycetes and the family Ophiostomataceae Nannfeldt, removing it from the orders Plectascales (Nannfeldt 1932, Hunt 1956), the Microascales of the Plectomycetes (Luttrell 1951, 1955; Kendrick and Molnar 1965) and the Ophiostomatales (Rosinski 1961). The order Sphaeriales contains four genera: *Europhium* Parker, *Sphaeronaemella* Karsten ex Seeler, *Chadefuudia* Feldman, and *Ceratocystis* (Ellis & Halstead) Bakshi.
- 1974 De Hoog concluded that the genus *Ceratocystis* should be separated into 2 separate genera based on the morphology of the conidial states: *Cerutocystis* Ellis and Halstead with the anamorphs *Chalara*, *Chalaropsis* Peyronel, and *Thielaviopsis* Went accepting 11 species (see table 2), and the genus *Ophiostoma* H. & P. Sydow containing the anamorphs *Sporothrix* Hektoen & Perkins ex Nicot & Mariat, *Verticicladiella* S. Hughes, *Leptographium* Lagerberg & Melin, and *Graphium* Corda. He proposed four new *Ophiostoma* combinations, enlarged the concept of the form-genus *Sporothrix*, and described new species.

- 1974 Jewell studies cellulose distribution in cell walls of 47 species of *Ceratocystis* and 4 species of *Europhium* cytochemically and using x-ray diffraction analysis. Thirty-one species of *Ceratocystis* and all of the *Europhium* species contained cellulose.
- 1974 Olchowecki and Reid published a monograph of 50 *Ceratocystis* species found in Manitoba with a key to 70 species of which 25 were described as new. They provided a first-time conidial description for three species. They subdivided the genus into four groups based on ascospore morphology; (1) the Minuta Group, (2) the Ips Group, (3) the Fimbriata Group, and (4) the Pilifera Group.
- 1974 Von Arx urged revision of the genus *Ceratocystis* and suggested that species with *Chalara*-like anamorphs be retained within the genus, but those species having holoblastic conidia be transferred to *Ophiostoma*. He maintained *Europhium* as a distinct genus.
- 1975 Nag Raj and Kendrick placed the form-genera *Chalaropsis* and *Thielaviopsis* in synonymy with *Chalara*.
- 1975 Upadhyay and Kendrick proposed dividing the genus *Ceratocystis* and establishing a new genus *Ceratocystiopsis* based on Olchowecki and Reid's Minuta group that included 19 species with falcate ascospores and **ascoma** with short necks. The new genus was classed as Ascomycotina, Plectomycetes, Microascales, Ophiostomataceae. They listed 13 anamorphs of *Ceratocystis* and *Ceratocystiopsis*, 4 of which were described as new: *Hyalorhinocla diella*, *Graphilbum*, *Hyalopesotum*, and *Pachnodium*. *Europhium* was considered a synonym of *Ceratocystis*, and *Ophiostoma* was not accepted as a valid genus. Olchowecki and Reid's three other groups-Ips, Fimbriata, and Pilifera-were accepted.
- 1975 Weijman and de Hoog discussed the results of cell wall cellulose (Smith and others 1967) and the presence of rhamnose (Spencer & Gorin 1971) as an aid in the classification of teleomorphs and anamorphs. They divided the genus *Ceratocystis* into two groups: (1) the phialidic endogenous conidial forms- *Chalara* and allied genera-and (2) the exogenous conidial forms of the *Ophiostoma* group (*Graphium*-like states). *Ceratocystis sensu stricto* for the *Chalara* type was used where rhamnose and cellulose were absent.
- 1977 Redhead and Malloch placed the genus *Ceratocystis* in the order Endomycetales and the family Endomycetaceae as a yeast-related genus. The family Ophiostomataceae was synonymized with the family Endomycetaceae.
- 1978 Upadhyay dropped emendation of the genus *Ceratocystis* by Bakshi (1951) and Hunt (1956) to conserve the original description by Ellis and Halstead.
- 1979 The Proceedings of the Second International Mycological Congress (Kananaskas II) was published in two volumes as *The Whole Fungus* (Kendrick 1979). Included were discussions of fungal morphology, classification, ecology, evolution, and techniques. New terms such as holomorph, teleomorph, and anamorph were defined. Malloch, von Arx, de Hoog, and Luttrell presented divergent views of *Ceratocystis* classification. Kendrick and DiCosmo's paper of teleomorph-anamorph connections listed 70 species of *Ceratocystis* and 2 species of *Ceratocystiopsis*.
- 1980 Benny and Kimbrough proposed a new order, Ophiostomatales, having only one family Ophiostomataceae (Nannfeldt 1932) that included the genera *Ophiostoma*, *Ceratocystis*, *Ceratocystiopsis*, and *Sphaeronaemella*. *Europhium* was synonymized, and *Chadefaudia* Feldman was rejected from the family. They resurrected the genus *Sphaeronaemella*, which Hunt (1956) had placed in synonymy with the genus *Ceratocystis*.
- 1981 Harrington published on cycloheximide as an aid to classification of the species. *Ceratocystis sensu stricto* (*Chalara* anamorphs) is sensitive to cycloheximide, but *Ceratocystis sensu lato* (*Ophiostoma* spp.) is not sensitive.
- 1981 Upadhyay published a monograph on the genera *Ceratocystis* Ellis & Halstead and *Ceratocystiopsis* Upadhyay & Kendrick. He placed the two genera in the order Microascales of the Plectomycetes. *Ophiostoma*, *Rostrella*, *Endoconidiophora*, *Grosmunnia*, *Sphaeronaemella*, and *Europhium* were rejected as valid genera and were reduced to synonymy with the genus *Ceratocystis*. He divided the genus *Ceratocystis* into four sections: (1) *Ophiostoma*, ascospores lacking a hyaline gelatinous sheath; (2) *Ips*, ascospores always surrounded by a hyaline gelatinous sheath, rectangular, never curved; (3) *Ceratocystis*, ascospores sheathed, partly curved; and, (4) *Endoconidiophora*, sheath elongated, inequilateral.
- 1983 Ainsworth and Bisby's *Dictionary of the Fungi* (Hawksworth and others 1983) listed the order Ophiostomatales and the family Ophiostomataceae as having 2 or 3 genera, 13 synonyms, and 85 species.
- 1983 Urosevic redescribed five *Ceratocystis* species that cause tracheomycoses of oak as *Ophiostoma* species.

- 1984 De Hoog and Scheffer published a historical reappraisal of the genera *Ceratocystis* and *Ophiostoma*. They concluded that all species with *Chalara* anamorphs are *Ceratocystis sensu strictu* and distinct from *Ceratocystiopsis*. All others having conidial anamorphs are *Ophiostoma* (other than *Chalara*, which has cell walls with rhamnose and is resistant to cycloheximide). The latter included *Europhiurn*. They recombined 14 *Ceratocystis* species as *Ophiostoma* species (table 8).
- 1985 Kendrick recognized the order Ophiostomatales having 15 genera and 130 species.
- 1985 Wingfield reclassified *Verticicladiella* as *Leptographium*. He noted that the annellidic as well as the sympodial type of growth occurs in the *Leptographium* species.
- 1986 Solheim presented a key to genera of the family Ophiostomataceae and a key to ten Norwegian *Ophiostoma* species and one *Ceratocystiopsis* species; *O. ainoae*, *O. cucullatum*, and *O. flexuosum* were described as new.
- 1987 Bridges and Perry described a new species: *Ceratocystiopsis runuculosus* Perry & Bridges, having a *Sporothrix* anamorph.

Table 8.-Recombined *Ceratocystis* species by de Hoog and Scheffer (1974)

---

<i>Ophiostoma araucariae</i> (Butin) de Hoog & Scheffer. ≡ <i>Ceratocystis araucariae</i> Butin
<i>Ophiostoma bacillisporum</i> (Butin & Zimmermann) de Hoog & Scheffer ≡ <i>Ceratocystis bacillispora</i> Butin & Zimmermann
<i>Ophiostoma brevicolla</i> (Davidson) de Hoog & Scheffer ≡ <i>Ceratocystis brevicollis</i> Davidson
<i>Ophiostoma distortum</i> (Davidson) de Hoog & Scheffer ≡ <i>Ceratocystis distorta</i> Davidson
<i>Ophiostoma dryocoetidis</i> (Kendrick & Molnar) de Hoog & Scheffer ≡ <i>Ceratocystis dryocoetidis</i> Kendrick & Molnar
<i>Ophiostoma francke-grosmaniae</i> (Davidson) de Hoog & Scheffer ≡ <i>Ceratocystis francke-grosmaniae</i> Davidson
<i>Ophiostoma huntii</i> (Robinson-Jeffrey) de Hoog & Scheffer ≡ <i>Ceratocystis huntii</i> Robinson-Jeffery
<i>Ophiostoma megalobrunneum</i> (Davidson & Toole) de Hoog & Scheffer ≡ <i>Ceratocystis megalobrunnea</i> Davidson & Toole
<i>Ophiostoma nigrum</i> (Davidson) de Hoog & Scheffer ≡ <i>Ceratocystis nigra</i> Davidson
<i>Ophiostoma populinum</i> (Hinds & Davidson) de Hoog & Scheffer ≡ <i>Ceratocystis populina</i> Hinds & Davidson
<i>Ophiostoma rostrocoronatum</i> (Davidson & Eslyn) de Hoog & Scheffer ≡ <i>Ceratocystis rostrocoronata</i> Davidson & Eslyn
<i>Ophiostoma seticolle</i> (Davidson) de Hoog & Scheffer ≡ <i>Ceratocystis seticollis</i> Davidson
<i>Ophiostoma sparsum</i> (Davidson) de Hoog & Scheffer ≡ <i>Ceratocystis sparsa</i> Davidson
<i>Ophiostoma tremulo-aureum</i> (Davidson & Hinds) de Hoog & Scheffer ≡ <i>Ceratocystis tremulo-aurea</i> Davidson & Hinds

---

- 1987 Von Arx accepted five species in the genus *Ceratocystis*. All have *Chalara* anamorphs, but only *C. fimbriata* has hat-shaped ascospores.
- 1987 Harrington recombined 11 *Ceratocystis* species as *Ophiostoma* (table 9).
- 1988 Hutchinson and Reid presented a key to seven potential wood-staining *Ceratocystis*. *Ceratocystis novae-zelandiae* was described as new. They listed one species of *Ceratocystiopsis* and a *Sphaeronaemella* species.
- 1988 Von Arx and van der Walt accepted the order Ophiostomatales by Benny and Kimbrough for the three genera *Ceratocystis*, *Ceratocystiopsis*, and *Ophiostoma* and the family Ophiostomataceae for *Ceratocystiopsis* and *Ophiostoma*. However, they proposed a new family, Pyxidiphoraceae, for *Ceratocystis* because of their *Chalara* anamorphs and the lack of cellulose in their cell walls (fig. 1).
- 1988 Wingfield and others described a new species *Ceratocystiopsis proteae* with a new anamorph genus *Knoxdaviesia*.
- 1989 Kowalski and Butin redescribed four species of *Ceratocystis* from oak in Poland and described two new species. *Ceratostomella quercus* Geor-gev. and *Ophiostoma roboris* were synonymized as *Ceratocystis piceae* (Munch) Bakshi.
- 1990 Harrington and Zambino rejected *Ceratocystis* minor var. *barrasi* J. Taylor as the mycangial fungus of *Dendroctonus frontalis* and established *Ceratocystiopsis ranaculosus* as the correct fungus based on isozyme analysis and mating studies.

Table 9.-Recombined *Ceratocystis* species by Harrington (1987)

---

<i>Ophiostoma abiocarpum</i> (Davidson)	Harrington
≡ <i>Ceratocystis abiocarpa</i> Davidson	
<i>Ophiostoma adjuncti</i> (Davidson)	Harrington
≡ <i>Ceratocystis adjuncti</i> Davidson	
<i>Ophiostoma aureum</i> (Robinson-Jeffrey & Davidson)	Harrington
≡ <i>Europhium aureum</i> Robinson-Jeffrey & Davidson	
≡ <i>Ceratocystis aurea</i> (Robinson-Jeffrey & Davidson)	Upadhyay
<i>Ophiostoma cainii</i> (Olchowecki & Reid)	Harrington
≡ <i>Ceratocystis cainii</i> Olchowecki & Reid	
<i>Ophiostoma clavigerum</i> (Robinson-Jeffrey & Davidson)	Harrington
≡ <i>Europhium clavigerum</i> Robinson-Jeffrey & Davidson	
≡ <i>Ceratocystis clavigem</i> (Robinson-Jeffrey & Davidson)	Upadhyay
<i>Ophiostoma crassivaginatum</i> (Griffin)	Harrington
≡ <i>Ceratocystis crassivaginata</i> Griffin	
≡ <i>Ceratocystiopsis crassivaginata</i> (Griffin)	Upadhyay
<i>Ophiostoma grandifoliae</i> (Davidson)	Harrington
≡ <i>Ceratocystis grandifoliae</i> Davidson	
<i>Ophiostoma robustum</i> (Robinson-Jeffrey & Davidson)	Harrington
≡ <i>Europhium robustum</i> Robinson-Jeffrey & Davidson	
≡ <i>Ceratocystis robusta</i> (Robinson-Jeffrey & Davidson)	Upadhyay
<i>Ophiostoma trinacriforme</i> (Parker)	Harrington
≡ <i>Europhium trinacriforme</i> Parker	
≡ <i>Ceratocystis trinacriforme</i> (Parker)	Upadhyay
<i>Ophiostoma valdivianum</i> (Butin)	Harrington
≡ <i>Ceratocystis valdiviana</i> Butin	
<i>Ophiostoma wagneri</i> (Goheen & Cobb)	Harrington
≡ <i>Ceratocystis wagneri</i> Goheen & Cobb	

---

- 1990 Malloch and Blackwell proposed the genus *Kathistes* and presented a key to the genera likely to be confused with *Ceratocystiopsis* — *Gabarnaudia*, *Sphaeronaemella*, *Ceratocystis*, *Ophiostoma*, *Pyxidiophora*, *Rhynchonectria*, *Treleasia*, *Subbaromyces*, *Spumatoria*, and *Klasterskya*.
- 1990 Van Wyk and Wingfield reviewed the controversial taxonomic status of *Ceratocystis*, *Ceratocystiopsis*, and *Ophiostoma* regarding the development of asci, ascospores, and **centrum** structure. They suggested that additional ultrastructural studies are required to clarify these relationships.

### LITERATURE CITED

- Bakshi, B.K. 1951. Studies on four species of *Ceratocystis*, with a discussion on fungi causing sap-stain in Britain. Commonwealth Mycological Institute Mycology Research Paper. **35:1–16**.
- Barr-as, S.J. 1970. Antagonism between *Dendroctonus frontalis* and the fungus *Ceratocystis minor*. Annals of the Entomological Society of America. **63:1187–1190**.
- Barras, S.J.; Perry, T.J. 1971. *Leptographium terebrantis* sp. nov. associated with *Dendroctonus terebrans* in loblolly pine. Mycopathologia et Mycologia Applicata. **43:1–10**.
- Barras, S.J.; Perry, T.J. 1975. Interrelationships among microorganisms, bark or ambrosia beetles and woody host tissue. An annotated bibliography 1965-1974. Gen. Tech. Rep. **SO-10**. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 34 p.
- Beaver, R.A. 1989. Insect-fungus relationships in the bark and ambrosia beetles. Symposia of the Royal Entomological Society of London, 1989. London: Academic Press. 121-143.
- Benny, G.L.; Kimbrough, J.W. 1980. A synopsis of the orders and families of Plectomycetes with keys to genera. Mycotaxon. **12:1–91**.
- Bramble, W.C.; Holst, E.C. 1940. Fungi associated with *Dendroctonus frontalis* in killing shortleaf pines and their effect on conduction. *Phytopathology*. **30:881–899**.
- Bridges, J.R.; Perry, T.J. 1985. Effects of mycangial fungi on gallery construction and distribution of bluestain in southern pine beetle-infested pine bolts. *Journal of Entomological Science*. **20:271–275**.
- Bridges, J.R.; Perry, T.J. 1987. *Ceratocystiopsis ranaculosus* sp. nov. associated with the southern pine beetle. *Mycologia*. **79:630–633**.
- Butin, H.; Aquilar, A.M. 1984. Blue-stain fungi on *Nothofagus* from Chile-including two new species of *Ceratocystis* Ellis & Halst. *Phytopathologische Zeitschrift*. **109:80–89**.
- Cain, R.F. 1972. Evolution of fungi. *Mycologia*. **64:1–14**.
- Cannon, P.F.; Hawksworth, D.L. 1982. A re-evaluation of *Melanospora* Corda and similar *Pyrenomyces* with a revision of the British species. *Botanical Journal of the Linnean Society*. **84:115–160**.
- Davidson, R.W. 1955. Wood-staining fungi associated with bark beetles in Engelmann spruce in Colorado. *Mycologia*. **47:58–67**.
- Davidson, R.W. 1958. Additional species of *Ophiostomataceae* from Colorado. *Mycologia*. **50:661–670**.
- Davidson, R.W. 1966. New species of *Ceratocystis* from conifers. *Mycopathologia et Mycologia Applicata*. **28:273–286**.
- Davidson, R.W. 1971. New species of *Ceratocystis*. *Mycologia*. **63:5–15**.
- Davidson, R.W. 1978. Staining fungi associated with *Dendroctonus adjuncti* in pines. *Mycologia*. **70:35–40**.
- de Hoog, G.S. 1974. The genera *Blastobotrys*, *Sporothrix*, *Calcarisporium* and *Calcarisporiella* gen. nov. *Studies in Mycology*. **7:1–84**.
- de Hoog, G.S.; Scheffer, R.J. 1984. *Ceratocystis* versus *Ophiostoma*: A reappraisal. *Mycologia*. **76:292–299**.
- Franklin, R.T. 1970. Observations on the blue stain-southern pine beetle relationship. *Journal of the Georgia Entomological Society*. **5:53–57**.
- Georgescu, C.; Teodoru, I.; Badea, M. 1948. Le Dessechement en masse du **chene**. Institutul de Cercetari si Experimentatie Foriestiera. Bucharest: Analele; **11: 185–223**.
- Goheen, D.J.; Cobb, F.W., Jr. 1978. Occurrence of *Verticicladiella wagnerii* and its perfect state, *Ceratocystis wagneri* sp. nov., in insect galleries. *Phytopathology*. **68:1192–1195**.
- Greuter, W. and others, eds. 1983. International code of botanical nomenclature adopted by the Thirteenth International Botanical Congress. Sydney, August 1981. [Regnum Vegetabile no. 1111. Utrecht: Bohn, Scheltema & Holkema. 472 p.
- Griffin, H.D. 1968. The genus *Ceratocystis* in Ontario. *Canadian Journal of Botany*. **46:689–718**.
- Halstead, B.D. 1890. Some fungous diseases of the sweet potato. The black rot. *New Jersey Agricultural Experiment Station Bulletin*. **76:7–14**.
- Halstead, B.D.; Fairchild, D.G. 1891. Sweet-potato black rot. *Journal of Mycology*. **7: 1–1 1**.
- Harrington, T.C. 1981. Cycloheximide sensitivity as a taxonomic character in *Ceratocystis*. *Mycologia*. **73:1123–1129**.

- Harrington, T.C. 1987. New combinations in *Ophiostoma* of *Ceratocystis* species with *Leptographium* anamorphs. *Mycotaxon*. **28:39–43**.
- Harrington, T.C. 1988. *Leptographium* species, their distribution, hosts and insect vectors. In: Harrington, T.C.; Cobb, F.W., Jr., eds. *Leptographium* root diseases on conifers. St. Paul, MN: American Phytopathology Society Press: 1-39.
- Harrington, T.C.; Zambino, P.J. 1990. *Ceratocystiopsis ranaculosus*, not *Ceratocystis minor* var. *barasii*, is the mycangial fungus of the southern pine beetle. *Mycotaxon*. **38: 103-115**.
- Hawksworth, D.L.; Sutton, B.C.; Ainsworth, G.C. 1983. Ainsworth and Bisby's dictionary of the fungi. 7th ed. Kew, Surrey, England: Commonwealth Mycological Institute. 445 p.
- Hennebert, D.L.; Weresub, L.K. 1977. Terms for states and forms of fungi, their names and types. *Mycotaxon*. **6:207–211**.
- Hunt, J. 1956. Taxonomy of the genus *Ceratocystis*. *Lloydia*. **19: 1-58**.
- Hutchinson, L.J.; Reid, J. 1988. Taxonomy of some potential wood-staining fungi from New Zealand 1. *Ophiostomataceae*. *New Zealand Journal of Botany*. **26:63–81**.
- Jewell, T.R. 1974. A qualitative study of cellulose distribution in *Ceratocystis* and *Europhium*. *Mycologia*. **66: 139-146**.
- Kendrick, W.B. 1962. The *Leptographium* complex. *Verticicladiella* Hughes. *Canadian Journal of Botany*. **40:771–797**.
- Kendrick, W.B., ed. 1971. Taxonomy of the fungi imperfecti. Toronto: University of Toronto Press. 309 p.
- Kendrick, W.B., ed. 1979. The whole fungus. In: Proceedings of the 2nd International Mycological Conference held at the Environmental Sciences Centre of the University of Calgary; Kananaskis, Alberta, Canada. Ottawa: National Museums of Canada and the Kananaskis Foundation. 793 p.
- Kendrick, W.B. 1985. The fifth kingdom. Ottawa: Mycologue Publications. 363 p.
- Kendrick, W.B.; Molnar, A.C. 1965. A new *Ceratocystis* and its *Verticicladiella* imperfect state associated with the bark beetle *Dryocoetes confusus* on *Abies Zasiocarpa*. *Canadian Journal of Botany*. **43:39–43**.
- Kowalski, T.; Butin, H. 1989. Taxonomy of known and new species of *Ceratocystis* from oak (*Quercus robur* L.). *Journal of Phytopathology*. **24:236–248**.
- Livingston, W.H.; Davidson, R.W. 1987. *Ophiostoma subannulatum*, a new fungal species pathogenic to grand fir roots. *Mycologia*. **79:144–147**.
- Luttrell, E.S. 1951. Taxonomy of the Pyrenomycetes. University of Missouri Studies. **24:1–121**
- Luttrell, E.S. 1955. The ascostromatic ascomycetes. *Mycologia*. **47:511–532**.
- Malloch, D. 1979. Plectomycetes and their anamorphs. In: Kendrick, B., ed. The whole fungus. Ottawa: National Museums of Canada: 153-165. Vol. 1.
- Malloch, D.; Blackwell, M. 1990. *Kathistes*, a new genus of pleomorphic ascomycetes. *Canadian Journal of Botany*. **68:1712–1721**.
- Mathre, D.E. 1964. Survey of *Ceratocystis* spp. associated with bark beetles in California. *Contributions of the Boyce Thompson Institute*. **22:353–362**.
- Mattson, W.J.; Levieux, J.; Bernard-Dagan, C., eds. 1988. Mechanisms of woody plant defenses against insects: search for a pattern. London: Springer-Verlag. 416 p.
- Melin, E.; Nannfeldt, J.A. 1934. Researches into the blueing of ground wood-pulp. *Svenska Skogsvårdsfoereningens Tidskrift*. **32:397–616**.
- Muller, E.; von Arx, J.A. 1973. Pyrenomycetes: Meliolales, Coronophorales, Sphaeriales. In: Ainsworth, G.C., Sparrow, F.K.; Sussman, A.S., eds. The fungi. New York: Academic Press: 87-132. Vol. 4A.
- Munch, E. 1907. Die blaufaule des nadelhoizes. *Naturwissenschaftliche Zeitschrift fur Forst-und Landwirtschaft*. **5:531–573**.
- Nag Raj, T.R.; Kendrick, W.B. 1975. A monograph of *Chalara* and allied genera. Waterloo: Wilfred Laurier University Press. 200 p.
- Nannfeldt, J.A. 1932. Studien uber die morphologie und systematik der nichtlichenisierten inoperculaten Discomyceten. *Nova Acta Regiae Societatis Scientiarum Upsaliensis*. Ser. 4. **8(2):1–368**.
- Nelson, R.M. 1934. Effect of bluestain on southern pines attacked by bark beetles. *Phytopathologische Zeitschrift*. **7:327–353**.
- Olchowecki, A.; Reid, J. 1974. Taxonomy of the genus *Ceratocystis* in Manitoba. *Canadian Journal of Botany*. **52:1675–1711**.
- Parker, A.K. 1957. *Europhium*, a new genus of the Ascomycetes with a *Leptographium* imperfect state. *Canadian Journal of Botany*. **35:173–179**.
- Rane, K.K.; Tattar, T.A. Pathogenicity of blue-stain fungi associated with *Dendroctonus terebrans*. *Plant Disease*. **71:879–883**.
- Redhead, S.A.; Malloch, D. 1977. The *Endomycetaceae*: new concepts, new taxa. *Canadian Journal of Botany*. **55:1701–1711**.
- Robinson, R.C. 1962. Blue stain fungi in lodgepole pine (*Pinus contorta* Dougl. var. *latifolia* Engelm.) infested by the mountain pine beetle (*Dendroctonus monticolae* Hopk.). *Canadian Journal of Botany*. **40:609–614**.
- Robinson-Jeffrey, R.C.; Davidson, R.W. 1968. Three new *Europhium* species with *Verticicladiella* imperfect states on blue-stained pine. *Canadian Journal of Botany*. **46:1523–1527**.
- Robinson-Jeffrey, R.C.; Grinchenko, A.H.H. 1964. A new fungus in the genus *Ceratocystis* occurring on



- blue-stained lodgepole pine attacked by bark beetles. *Canadian Journal of Botany*. **42**:527–532.
- Rosinski, M.A. 1961. Development of the ascocarp of *Cerutocystis ulmi*. *American Journal of Botany*. **48**:285–293.
- Rosinski, M.A. 1965. Further confirmation of the occurrence of cellulose in *Cerutocystis ulmi*. *Mycologia*. **57**:668.
- Rosinski, M.A.; Campana, R.J. 1964. Chemical analysis of the cell wall of *Cerutocystis ulmi*. *Mycologia*. **56**:738–744.
- Rumbold, C.T. 1931. Two blue-staining fungi associated with bark-beetle infestation of pines. *Journal of Agricultural Research*. **43**:847–873.
- Rumbold, C.T. 1936. Three blue-staining fungi, including two new species, associated with bark beetles. *Journal of Agricultural Research*. **52**:419–437.
- Rumbold, C.T. 1941. A blue-stain fungus, *Cerutocystis montium* n. sp. and some yeasts associated with two species of *Dendroctonus*. *Journal of Agricultural Research*. **62**:589–601.
- Scerbin-Parfenenko, A.L. 1953. Rakovye a sosudistye boleznj listvennych porod. Goslesbimizdat. [Vol. no. unknown] 1-89.
- Sigler, L.; Hawksworth, D.L. 1987. International commission on the taxonomy of fungi (ITCF) code of practice for systemic mycologists. *Mycopathologia*. **99**:3–7.
- Smith, M.J.; Patik, C.M.; Rosinski, M.A. 1967. A comparison of cellulose production in the genus *Cerutocystis*. *Mycologia*. **59**:965–969.
- Solheim, H. 1986. Species of Ophiostomataceae isolated from *Picea ubies* infested by the bark beetle *Ips typographus*. *Nordic Journal of Botany*. **6**:199–207.
- Spencer, J.F.T.; Gorin, P.A.J. 1971. Systematics of the genera *Cerutocystis* and *Gruphium*. Proton magnetic resonance spectra of the mannose-containing polysaccharides as an aid in classification. *Mycologia*. **63**:387–402.
- Sydow, Von H.; P. Sydow. 1919. Mykologische mitteilungen. *Annales Mycologici*. **17**(1):3–47.
- Tsuneda, A.; Hiratsuka, Y. 1984. Sympodial and anellidic conidiation in *Cerutocystis clavigera*. *Canadian Journal of Botany*. **62**:2618–2624.
- Upadhyay, H.P. 1978. Proposal for the conservation of the generic name *Cerutocystis* Ell. & Halst. (1890) against *Sphueronuemellu* Karsten (1884). *Taxon*. **27**:553.
- Upadhyay, H.P. 1981. A monograph of the genus *Cerutocystis* and *Cerutocystiopsis*. Athens: University of Georgia Press. 176 p.
- Upadhyay, H.P.; Kendrick, W.B. 1975. Prodromus for a revision of *Cerutocystis* (Microascales, Ascomycetes) and its conidial states. *Mycologia*. **67**:798–805.
- Urosevic, B. 1983. Tracheomycotic diseases in oak. *Communicationes Instituti Forestalis Cechosloveniae*. **13**:85–100.
- van Wyk, P.W.J.; Wingfield, M.J. 1990. Ascospore development in *Cerutocystis sensu lato* (Fungi): a review. *Bothalia*. **20**(2):141–145.
- von Arx, J.A. 1970. The genera of fungi sporulating in pure culture. Lehre: Cramer. 288 p.
- von Arx, J.A. 1974. The genera of fungi sporulating in pure culture. 2nd ed. Vaduz: Cramer. 315 p.
- von Arx, J.A. 1979. Ascomycetes as Fungi Imperfecti. In: Kendrick, B., ed. The whole fungus. Ottawa: National Museums of Canada: 201-214. Vol. 1.
- von Arx, J.A. 1987. A re-evaluation of the Eurotiales. *Persoonia*. **13**:273–300.
- von Arx, J.A.; Muller, E. 1954. Die Gattungen der amerosporen Pyrenomyceten. *Beitrage zur Kryptogamenflora der Schweiz*. **11**: 1-434.
- von Arx, J.A.; van der Walt, J.P. 1988. Ophiostomatales and Endomycetales. In: de Hoog, G.S.; Smith, M.Th.; Weijman, A.C.M., eds. The expanding realm of yeast-like fungi: Proceedings of an international symposium on the perspectives of taxonomy, ecology and phylogeny of yeast and yeast-like fungi; 1987 Aug. 3-7; Amersfort, The Netherlands. Amsterdam: Elsevier Science: 167-176.
- von Hohnel, F. 1918. Mykologische fragmente CXCIV. Uber die gattung *Purodiopsis* Maublanc. *Annales Mycologici*. **16**:40–41.
- Weijman, A.C.M.; de Hoog, G.S. 1975. On the subdivision of the genus *Cerutocystis*. *Antonie van Leeuwenhoek. Journal of Microbiology and Serology*. **41**:353–360.
- Whitney, H.S. 1971. Association of *Dendroctonus ponderosae* (Coleoptera:Scolytidae) with blue stain fungi and yeasts during brood development in lodgepole pine. *Canadian Entomologist*. **103**:1495–1503.
- Whitney, H.S. 1982. Relationships between bark beetles and symbiotic organisms. In: Mitton, J.B.; Sturgeon, K., eds. A system for the study of evolutionary biology. Austin: University of Texas Press: 183–211.
- Whitney, H.S.; Cobb, F.W., Jr. 1972. Non-staining fungi associated with the bark beetle *Dendroctonus brevicomis* (Coleoptera:Scolytidae) on *Pinus ponderosae*. *Canadian Journal of Botany*. **50**:1943–1945.
- Wingfield, M.J. 1983. Association of *Verticicladiella procera* and *Leptographium terebrantis* with insects in the Lake States. *Canadian Journal of Forest Research*. **13**:1238–1245.
- Wingfield, M.J. 1985. Reclassification of *Verticicladiella* based on conidial development. *Transactions of the British Mycological Society*. **85**:81–93.

- Wingfield, M.J.; van Wyk, P.S.; Marasas, W.F.O.  
1988. *Ceratocystiopsis proteae* sp. nov., with a new anamorph genus. *Mycologia*. **80:23–30**.
- Wood, S.L. 1982. The bark and ambrosia beetles of North and Central America (Coleoptera: Scolytidae), a taxonomic monograph. Great Basin Naturalist Memoirs # 6. Provo, UT: Brigham Young University. 1359 p.
- Wright, E.F.; Cain, R.F. 1961. New species of *Ceratocystis*. Canadian Journal of Botany. **39:1215–1230**.

Perry, Thelma J. 1991. A synopsis of the taxonomic revisions in the genus *Ceratocystis* including a review of blue-staining species associated with *Dendroctonus* bark beetles. Gen. Tech. Rep. SO-86. New Orleans, LA: US. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 16 p.

Lists *Ophiostoma* species previously described as *Ceratocystis* and summarizes events that led to revisions in the genera.

**Keywords:** *Ceratocystiopsis*, *Ophiostoma*

Persons of any race, color, national origin, sex, age, religion, or with any handicapping condition are welcome to use and enjoy all facilities, programs, and services of the USDA. Discrimination in any form is strictly against agency policy, and should be reported to the Secretary of Agriculture, Washington, DC 20250.



