

 Open access • Journal Article • DOI:10.1017/S096012951100003X

## A system of interaction and structure v: The exponentials and splitting

— [Source link](#) 

Alessio Guglielmi, Lutz Straßburger

**Institutions:** University of Bath, École Polytechnique

**Published on:** 01 Jun 2011 - Mathematical Structures in Computer Science (Cambridge University Press)

**Topics:** Splitting theorem, Commutative property, Deep inference and Linear logic

Related papers:

- [A system of interaction and structure](#)
- [Linear Logic and Noncommutativity in the Calculus of Structures](#)
- [A system of interaction and structure ii: the need for deep inference](#)
- [A Purely Logical Account of Sequentiality in Proof Search](#)
- [A Local System for Classical Logic](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/a-system-of-interaction-and-structure-v-the-exponentials-and-ai2f2kbt1v>



*Citation for published version:*

Guglielmi, A & Straburger, L 2011, 'A system of interaction and structure V: the exponentials and splitting', *Mathematical Structures in Computer Science*, vol. 21, no. 3, pp. 563-584.  
<https://doi.org/10.1017/S096012951100003X>

*DOI:*

[10.1017/S096012951100003X](https://doi.org/10.1017/S096012951100003X)

*Publication date:*

2011

[Link to publication](#)

© Cambridge University Press 2011

## University of Bath

### Alternative formats

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# A system of interaction and structure V: the exponentials and splitting

ALESSIO GUGLIELMI<sup>†</sup> and LUTZ STRABBURGER<sup>‡</sup>

<sup>†</sup>*Department of Computer Science, University of Bath, Bath BA2 7AY, United Kingdom*  
Email: [alessio@uglielmi.name](mailto:alessio@uglielmi.name)

<sup>‡</sup>*INRIA Saclay–Île-de-France and École Polytechnique, Laboratoire d'Informatique (LIX),  
Rue de Saclay, 91128 Palaiseau Cedex, France*  
Email: [lutz@lix.polytechnique.fr](mailto:lutz@lix.polytechnique.fr)

Received 22 January 2004; revised 3 December 2010

System NEL is the mixed commutative/non-commutative linear logic BV augmented with linear logic's exponentials, or, equivalently, it is MELL augmented with the non-commutative self-dual connective *seq*. NEL is presented in deep inference, because no Gentzen formalism can express it in such a way that the cut rule is admissible. Other recent work shows that system NEL is Turing-complete, and is able to express process algebra sequential composition directly and model causal quantum evolution faithfully. In this paper, we show cut elimination for NEL, based on a technique that we call *splitting*. The splitting theorem shows how and to what extent we can recover a sequent-like structure in NEL proofs. When combined with a 'decomposition' theorem, proved in the previous paper of this series, splitting yields a cut-elimination procedure for NEL.

## 1. Introduction

This is the fifth in a series of papers dedicated to the proof theory of a self-dual, non-commutative, linear connective called *seq*, in the context of linear logic. The addition of *seq* to multiplicative linear logic yields a logic that we call BV. This logic is the main subject of study of this series of papers. BV is conjectured to be the same as pomset logic, which was studied by Retoré in Retoré (1997) and other papers.

BV was defined in Guglielmi (2007) (the first paper of this series), where a sound, complete and cut-free system for BV was given, together with a cut-elimination procedure. The proof system of BV departs radically from the traditional sequent calculus methodology, and instead adopts *deep inference* as the design principle. Briefly, this means that proofs can be freely composed by the same connectives used for formulae, or, equivalently, inference rules can be applied arbitrarily deeply inside formulae. Guglielmi (2007) provides an introduction to deep inference, which was, in fact, originally conceived precisely for the purposes of capturing BV.

<sup>†</sup> Alessio Guglielmi is supported by an ANR Senior Chaire d'Excellence entitled 'Identity and Geometric Essence of Proofs' and by the INRIA ARC 'REDO: Redesigning Logical Syntax'.

<sup>‡</sup> Lutz Straßburger is supported by the ANR project 'INFER: Theory and Application of Deep Inference' and by the INRIA ARC 'REDO: Redesigning Logical Syntax'.

The formalism used so far in this series of papers is called the *calculus of structures*, or *CoS*, and we will use it again in this paper. This is, conceptually, the simplest formalism in deep inference, being just a special form of term rewriting. More sophisticated formalisms in deep inference are emerging, in particular, *open deduction* (Guglielmi *et al.* 2010a). This formalism improves on CoS because it has an increased algebraic flavour and allows for a speed-up in the size of proofs. However, these differences do not affect cut-elimination procedures, in the sense that from a cut-elimination procedure in CoS, one can trivially obtain a cut-elimination procedure in open deduction. For this reason, the results in this paper are broadly valid within the deep-inference paradigm.

The use of deep inference is necessary: Alwen Tiu showed in the second paper of this series (Tiu 2006) that it is impossible for the sequent calculus to provide a sound, complete and cut-free proof system for BV. This is proved by exhibiting an infinite set of BV tautologies with a cleverly designed structure such that any bounded-depth, cut-free inference system (in particular, any sequent calculus system) is either unsound or incomplete on the set of tautologies. The design of these tautologies exploits the ability of the seq connective, together with the usual ‘par’ disjunction of linear logic, to bury at arbitrary depths inside formulae certain key structures that have to be ‘unlocked’, by inference, before any other parts of the formula are touched. We stress the fact that this behaviour is independent of the logical formalism employed to describe it. So, deep inference appears to be the most natural choice of proof-system design methodology because of its ability to apply inference at arbitrary depths inside formulae.

BV might be considered exotic as a logic, but it has a very natural algebraic character, that already found applications in diverse fields. We mention here:

- (1) its ability to capture very precisely the sequential connective of Milner’s CCS (and hence of other process algebras) (Bruscoli 2002);
- (2) a better axiomatisation of causal quantum computation than linear logic provides (Blute *et al.* 2008; Blute *et al.* 2010); and
- (3) a new class of categorical models (Blute *et al.* 2009).

We know that the proof system BV is NP-complete (Kahramanoğulları 2008a), and its feasibility for proof search has also been studied (Kahramanoğulları 2004).

This fifth paper, together with the fourth paper (Straßburger and Guglielmi 2009) in the series, is devoted to the proof theory of system BV when it is enriched with linear logic’s exponentials. We call the resulting system NEL (non-commutative exponential linear logic). We can also think of NEL as MELL (multiplicative exponential linear logic (Girard 1987)) augmented with seq. NEL, which was first presented in Guglielmi and Straßburger (2002), is conservative over both BV and over MELL augmented by the mix and nullary mix rules (Fleury and Retoré 1994; Retoré 1993; Abramsky and Jagadeesan 1994). Like BV, NEL cannot be expressed with a cut-free proof system outside of deep inference because of the counterexample in Tiu (2006) (mentioned above), and because NEL is conservative over BV.

An important feature of NEL is that it is Turing-complete – see Straßburger (2003c) for a proof. This makes for an interesting comparison with MELL, whose complexity is currently unknown. MELL is expected to be decidable, but several years of research have

not settled the question. If MELL turns out to be decidable, then seq would be the decisive factor that allows us to cross the border between decidability and Turing-completeness. This would have an intuitive explanation in the fact that seq can be employed to simulate the structure of a Turing machine tape, which is precisely what MELL, which is fully commutative, apparently cannot do.

Further interest in NEL comes from the possibility of enhancing, in a natural way, the range of applications of BV. Similarly to what happens when exponentials are added to multiplicative linear logic, the augmented expressivity of NEL over BV can be employed to better capture process algebras, for example, and this is indeed an active area of research. We might equally expect that NEL will further improve our ability to describe quantum evolution phenomena, and we expect to find enriched categorical models over those for BV.

Each of the two NEL papers in the series is devoted to a theorem: *splitting* in this paper and *decomposition* in the previous paper (Straßburger and Guglielmi 2009). Together, the two theorems immediately yield a cut-elimination procedure, and the cut-elimination result is claimed in this paper.

Splitting (which was introduced in Guglielmi (2007)) is, in a sense, a way of rebuilding into deep inference the structure of Gentzen sequent-calculus proofs to the extent possible in the presence of par and seq. The technique consists of first blocking the access of inference rules to a part of the formula to be proved, however deep, and then removing from the context of this blocked part as much ‘logical material’ as possible. In other words, we prove as much as we can of a given formula in the presence of a part that has been blocked. The splitting theorem states properties of what is left of the context of the blocked part, in relation to the shape of the blocked part. It turns out that the splitting property is just a generalisation of the shape of Gentzen calculi rules, and coincides precisely with them when we stipulate that the blocked part of a formula is at the shallowest possible level.

Splitting is, *a priori*, a hard theorem to prove, but, thanks to the decomposition theorem proved in Straßburger and Guglielmi (2009), we only need to prove it for a fragment of NEL, and this is what we do in this paper. Once splitting is available, cut elimination follows immediately.

The main results of this paper have already been presented, without proof, in Guglielmi and Straßburger (2002) – the proofs of the statements have been available for several years in a manuscript on the web.

## 2. The system

In this section we give a brief summary of the system NEL as defined in Straßburger and Guglielmi (2009) – see that paper for further details and introductory comments.

**Definition 2.1.** There are countably many *positive* and *negative atoms*. They, positive or negative, are denoted by  $a, b, \dots$ . *Structures* are denoted by  $S, P, Q, R, T, U, V, W, X$

Associativity

$$\begin{aligned}
 [\vec{R} \wp [\vec{T} \wp \vec{U}]] &= [\vec{R} \wp \vec{T} \wp \vec{U}] \\
 (\vec{R} \otimes (\vec{T} \otimes \vec{U})) &= (\vec{R} \otimes \vec{T} \otimes \vec{U}) \\
 \langle \vec{R} \triangleleft \langle \vec{T} \triangleleft \vec{U} \rangle \rangle &= \langle \vec{R} \triangleleft \vec{T} \triangleleft \vec{U} \rangle
 \end{aligned}$$

Commutativity

$$\begin{aligned}
 [\vec{R} \wp \vec{T}] &= [\vec{T} \wp \vec{R}] \\
 (\vec{R} \otimes \vec{T}) &= (\vec{T} \otimes \vec{R})
 \end{aligned}$$

Unit

$$\begin{aligned}
 [\circ \wp \vec{R}] &= [\vec{R}] \\
 (\circ \otimes \vec{R}) &= (\vec{R}) \\
 \langle \circ \triangleleft \vec{R} \rangle &= \langle \vec{R} \rangle \\
 \langle \vec{R} \triangleleft \circ \rangle &= \langle \vec{R} \rangle
 \end{aligned}$$

Singleton

$$[R] = (R) = \langle R \rangle = R$$

Negation

$$\begin{aligned}
 \bar{\circ} &= \circ \\
 \overline{[R_1 \wp \dots \wp R_h]} &= (\bar{R}_1 \otimes \dots \otimes \bar{R}_h) \\
 \overline{(R_1 \otimes \dots \otimes R_h)} &= [\bar{R}_1 \wp \dots \wp \bar{R}_h] \\
 \overline{\langle R_1 \triangleleft \dots \triangleleft R_h \rangle} &= \langle \bar{R}_1 \triangleleft \dots \triangleleft \bar{R}_h \rangle \\
 \overline{?R} &= !\bar{R} \\
 \overline{!R} &= ?\bar{R} \\
 \bar{\bar{R}} &= R
 \end{aligned}$$

Contextual Closure

$$\text{if } R = T \text{ then } S\{R\} = S\{T\}$$

Fig. 1. Basic equations for the syntactic equivalence =.

and Z. The structures of the language NEL are generated by

$$S ::= a \mid \circ \mid \underbrace{[S \wp \dots \wp S]}_{>0} \mid \underbrace{(S \otimes \dots \otimes S)}_{>0} \mid \underbrace{\langle S \triangleleft \dots \triangleleft S \rangle}_{>0} \mid ?S \mid !S \mid \bar{S}$$

where  $\circ$ , the *unit*, is not an atom and  $\bar{S}$  is the *negation* of the structure  $S$ . Structures with a hole that does not appear in the scope of a negation are denoted by  $S\{ \}$ . The structure  $R$  is a *substructure* of  $S\{R\}$ , and  $S\{ \}$  is its *context*. We will simplify the notation for context in cases where structural parentheses fill the hole exactly: for example,  $S[R \wp T]$  stands for  $S\{[R \wp T]\}$ . The structures of the language NEL are equivalent modulo the relation  $=$ , defined in Figure 1. There,  $\vec{R}$ ,  $\vec{T}$  and  $\vec{U}$  stand for finite, non-empty sequences of structures (elements of the sequences are separated by  $\wp$ ,  $\triangleleft$  or  $\otimes$ , as appropriate in the context).

**Definition 2.2.** Figure 2 shows the system SNEL (*symmetric non-commutative exponential linear logic*). The rules  $\text{ai}\downarrow$ ,  $\text{ai}\uparrow$ ,  $\text{s}$ ,  $\text{q}\downarrow$ ,  $\text{q}\uparrow$ ,  $\text{p}\downarrow$ ,  $\text{p}\uparrow$ ,  $\text{e}\downarrow$ ,  $\text{e}\uparrow$ ,  $\text{w}\downarrow$ ,  $\text{w}\uparrow$ ,  $\text{b}\downarrow$ ,  $\text{b}\uparrow$ ,  $\text{g}\downarrow$  and  $\text{g}\uparrow$  are called *atomic interaction*, *atomic cut*, *switch*, *seq*, *coseq*, *promotion*, *copromotion*, *empty*, *coempty*, *weakening*, *coweakening*, *absorption*, *coabsorption*, *digging* and *codigging*, respectively. The *down fragment* of SNEL is  $\{\text{ai}\downarrow, \text{s}, \text{q}\downarrow, \text{p}\downarrow, \text{e}\downarrow, \text{w}\downarrow, \text{b}\downarrow, \text{g}\downarrow\}$ , the *up fragment* is  $\{\text{ai}\uparrow, \text{s}, \text{q}\uparrow, \text{p}\uparrow, \text{e}\uparrow, \text{w}\uparrow, \text{b}\uparrow, \text{g}\uparrow\}$ . Figure 3 then shows *system NEL*, where the rule  $\circ\downarrow$  is called *unit*.

All inference rules in SNEL have the form

$$\rho \frac{S\{T\}}{S\{R\}}$$

$$\begin{array}{ccc}
 \text{ai} \downarrow \frac{S\{\circ\}}{S[a \otimes \bar{a}]} & & \text{ai} \uparrow \frac{S(a \otimes \bar{a})}{S\{\circ\}} \\
 \\
 \text{s} \frac{S([R \otimes U] \otimes T)}{S[(R \otimes T) \otimes U]} & & \\
 \\
 \text{q} \downarrow \frac{S\langle [R \otimes U] \triangleleft [T \otimes V] \rangle}{S\langle (R \triangleleft T) \otimes (U \triangleleft V) \rangle} & & \text{q} \uparrow \frac{S\langle (R \triangleleft U) \otimes (T \triangleleft V) \rangle}{S\langle (R \otimes T) \triangleleft (U \otimes V) \rangle} \\
 \\
 \text{p} \downarrow \frac{S\{![R \otimes T]\}}{S[!R \otimes ?T]} & & \text{p} \uparrow \frac{S(?R \otimes !T)}{S\{?(R \otimes T)\}} \\
 \\
 \text{e} \downarrow \frac{S\{\circ\}}{S\{!\circ\}} & & \text{e} \uparrow \frac{S\{?\circ\}}{S\{\circ\}} \\
 \\
 \text{w} \downarrow \frac{S\{\circ\}}{S\{?R\}} & & \text{w} \uparrow \frac{S\{!R\}}{S\{\circ\}} \\
 \\
 \text{b} \downarrow \frac{S[?R \otimes R]}{S\{?R\}} & & \text{b} \uparrow \frac{S\{!R\}}{S(!R \otimes R)} \\
 \\
 \text{g} \downarrow \frac{S\{??R\}}{S\{?R\}} & & \text{g} \uparrow \frac{S\{!R\}}{S\{!!R\}}
 \end{array}$$

Fig. 2. System SNEL.

$$\begin{array}{ccc}
 \circ \downarrow \frac{}{\circ} & \text{ai} \downarrow \frac{S\{\circ\}}{S[a \otimes \bar{a}]} & \text{e} \downarrow \frac{S\{\circ\}}{S\{!\circ\}} \\
 \\
 \text{s} \frac{S([R \otimes U] \otimes T)}{S[(R \otimes T) \otimes U]} & \text{q} \downarrow \frac{S\langle [R \otimes U] \triangleleft [T \otimes V] \rangle}{S\langle (R \triangleleft T) \otimes (U \triangleleft V) \rangle} & \text{p} \downarrow \frac{S\{![R \otimes T]\}}{S[!R \otimes ?T]} \\
 \\
 \text{w} \downarrow \frac{S\{\circ\}}{S\{?R\}} & \text{b} \downarrow \frac{S[?R \otimes R]}{S\{?R\}} & \text{g} \downarrow \frac{S\{??R\}}{S\{?R\}}
 \end{array}$$

Fig. 3. System NEL.

saying that if a structure matches  $R$  in a context  $S\{ \}$ , it can be rewritten as specified by  $T$  in the same context  $S\{ \}$  (or *vice versa* when reasoning top-down). The unit rule of NEL is special in this respect, as it has no context; however, we only use it for convenience, and we could easily do without it by slightly adapting the notion of proof.

A derivation  $\Delta$  is a chain of consecutive applications of instances of inference rules. A derivation with no premise is called a *proof*, denoted by  $\Pi$ . A system  $S$  *proves*  $R$  if there

is in the system  $\mathcal{S}$  a proof  $\Pi$  whose conclusion is  $R$ . We use

$$\begin{array}{c} T \\ \mathcal{S} \parallel \Delta \\ R \end{array}$$

to denote a derivation  $\Delta$  with premise  $T$  and conclusion  $R$  whose rules are in  $\mathcal{S}$ , and

$$\begin{array}{c} \mathcal{S} \parallel \Pi \\ R \end{array}$$

to denote a proof  $\Pi$  in  $\mathcal{S}$  whose conclusion is  $R$ .

**Definition 2.3.** A rule  $\rho$  is *derivable* in the system  $\mathcal{S}$  if  $\rho \notin \mathcal{S}$  and for every instance

$$\rho \frac{T}{R}$$

there exists a derivation from  $T$  to  $R$  in  $\mathcal{S}$ . We say that a rule  $\rho$  is *admissible* for the system  $\mathcal{S}$  if  $\rho \notin \mathcal{S}$  and for every proof in  $\mathcal{S} \cup \{\rho\}$  there is a proof in  $\mathcal{S}$  with the same conclusion. Two systems are *equivalent* if they prove the same structures. Two systems  $\mathcal{S}$  and  $\mathcal{S}'$  are *strongly equivalent* if for every derivation from  $T$  to  $R$  in  $\mathcal{S}$  there is a derivation from  $T$  to  $R$  in  $\mathcal{S}'$ , and *vice versa*.

Notice that interaction and cut are atomic in SNEL; we can define their general versions as follows.

**Definition 2.4.** The following rules are called *interaction* and *cut*, respectively:

$$i\downarrow \frac{S\{\circ\}}{S[R \wp \bar{R}]} \quad \text{and} \quad i\uparrow \frac{S(R \otimes \bar{R})}{S\{\circ\}}$$

where  $R$  and  $\bar{R}$  are called *principal structures*.

The following two propositions were shown in Straßburger and Guglielmi (2009).

**Proposition 2.5.** The rule  $i\downarrow$  is derivable in  $\{ai\downarrow, s, q\downarrow, p\downarrow, e\downarrow\}$ , and, dually, the rule  $i\uparrow$  is derivable in the system  $\{ai\uparrow, s, q\uparrow, p\uparrow, e\uparrow\}$ .

**Proposition 2.6.** Each rule  $\rho\uparrow$  in SNEL is derivable in  $\{i\downarrow, i\uparrow, s, \rho\downarrow\}$ , and, dually, each rule  $\rho\downarrow$  in SNEL is derivable in the system  $\{i\downarrow, i\uparrow, s, \rho\uparrow\}$ .

As an immediate consequence of Propositions 2.5 and 2.6, we get the following proposition.

**Proposition 2.7.** The systems  $NEL \cup \{i\uparrow\}$  and  $SNEL \cup \{\circ\downarrow\}$  are strongly equivalent.

In the remainder of this paper we will give the proof of the cut-elimination theorem, which, as usual in deep inference, means proving that the up-fragment is admissible.






Fig. 4. Decomposition of derivations (Theorem 2.11).

**Theorem 2.8 (Cut admissibility).** System NEL is equivalent to  $\text{SNEL} \cup \{\circ\downarrow\}$ .

The following corollaries are immediate consequences of cut admissibility.

**Corollary 2.9.** The rule  $i\uparrow$  is admissible for system NEL.

**Corollary 2.10.** For any two structures *T* and *R*, we have

$$\text{SNEL} \begin{array}{c} T \\ \parallel \\ R \end{array} \text{ if and only if } \text{NEL} \begin{array}{c} \parallel \\ [\bar{T} \otimes R] \end{array}$$

Our cut elimination proof relies on the following theorem, which is a special case of a more general one, whose proof can be found in Straßburger and Guglielmi (2009).

**Theorem 2.11 (Decomposition).** Every derivation  $\Delta$  in SNEL can be rewritten as shown in Figure 4.

### 3. Splitting

Three approaches to cut elimination in deep inference have been explored in previous papers, but none of them can be applied in our case.

First, we could hope to rely on semantics, as Brünnler and Tiu did in Brünnler and Tiu (2001) for classical logic. However, to date, there is no provability semantics for NEL (in the sense, for example, of phase spaces (Okada 1999) or other model theoretic semantics) that we could use for a completeness argument.

Second, we could hope to use the approach taken in Brünnler (2003), where Brünnler presents a simple syntactic method that employs the atomicity of cut together with certain proof theoretical properties of classical logic. More recent papers that refine his method by using *atomic flows* are Guglielmi and Gundersen (2008) and Guglielmi *et al.* (2010b). However, this approach cannot be used for NEL because contraction cannot be applied to arbitrary formulas as in classical logic.

Third, we could hope to do as in (Guglielmi and Straßburger 2001; Straßburger 2003b), where we relied on permutations of rules. However, traditional techniques based on simple notions of rule-instance permutation cannot work. To use permutation arguments for NEL requires a more general notion of permutation than the usual one. The following remark shows an example that illustrates the point and gives some hints to anybody who would like to proceed in that direction.

**Remark 3.1.** Let  $P$  and  $Q$  be arbitrary provable structures with  $P \neq \circ \neq Q$ . Consider then the following proof, where we first stack two proofs of  $P$  and  $Q$  one on top of the other and then interweave two structures over the atoms  $a, b$  and  $c$  and their duals, and where the bottom rule instance is a  $q\uparrow$ :

$$\begin{array}{c}
 \text{NEL} \parallel \\
 P \\
 \text{NEL} \parallel \\
 (P \otimes Q) \\
 2 \cdot \text{ai}\downarrow \frac{}{(P \otimes \langle [b \wp \bar{b}] \triangleleft [c \wp \bar{c}] \rangle \otimes Q)} \\
 q\downarrow \frac{}{(P \otimes \langle [b \triangleleft c] \wp \langle \bar{b} \triangleleft \bar{c} \rangle \rangle \otimes Q)} \\
 s \frac{}{(P \otimes \langle [b \triangleleft c] \rangle \wp \langle \langle \bar{b} \triangleleft \bar{c} \rangle \rangle \otimes Q)} \\
 \text{ai}\downarrow \frac{}{(P \otimes \langle [a \wp \bar{a}] \triangleleft \langle [b \triangleleft c] \rangle \wp \langle \langle \bar{b} \triangleleft \bar{c} \rangle \rangle \otimes Q)} \\
 q\downarrow \frac{}{(P \otimes \langle [a \triangleleft b \triangleleft c] \rangle \wp \langle \bar{a} \triangleleft \langle \langle \bar{b} \triangleleft \bar{c} \rangle \rangle \otimes Q)} \\
 s \frac{}{[(P \otimes \langle a \triangleleft b \triangleleft c \rangle) \wp \langle \bar{a} \triangleleft \langle \langle \bar{b} \triangleleft \bar{c} \rangle \rangle \otimes Q]} \\
 q\uparrow \frac{}{[(P \otimes \langle a \triangleleft b \rangle) \triangleleft c] \wp \langle \bar{a} \triangleleft \langle \langle \bar{b} \triangleleft \bar{c} \rangle \rangle \otimes Q]}
 \end{array} \tag{1}$$

We can further suppose that the atoms  $a, b, c$  and their duals do not appear in  $P$  and  $Q$ . If so, any permutation of the bottom  $q\uparrow$  instance over the rule instances immediately above it leads to an unprovable structure, because some two dual atoms  $a/\bar{a}$  or  $b/\bar{b}$  or  $c/\bar{c}$  would become connected by a connective different from a  $\wp$ . In order for a proof to exist, any rule instance changing the mutual logical relations of the  $a/\bar{a}, b/\bar{b}$  and  $c/\bar{c}$  atoms in

the conclusion must be an instance of  $q\uparrow$ . As an example of the contrary, consider

$$q\downarrow \frac{\langle [(P \otimes \langle a \triangleleft b \rangle) \wp \langle \bar{a} \triangleleft (\langle \bar{b} \triangleleft \bar{c} \rangle \otimes Q)] \triangleleft c \rangle]}{\langle [(P \otimes \langle a \triangleleft b \rangle) \triangleleft c \rangle \wp \langle \bar{a} \triangleleft (\langle \bar{b} \triangleleft \bar{c} \rangle \otimes Q)] \rangle]} \quad (2)$$

where  $c$  and  $\bar{c}$  are not able to be joined in any  $ai\downarrow$  going upwards. This represents a difficulty for any argument based on permuting up the  $q\uparrow$  rule.

However, we can observe that the entire proofs of  $P$  and  $Q$  could be permuted below the  $q\uparrow$  instance, step by step. The two structures  $P$  and  $Q$  act as sort of ‘locks’ for the substructure made of  $a, b$  and  $c$  atoms (and their duals). Once the locks are open, by independently proving them and so reducing them to the unit, the permuting up of the  $q\uparrow$  instance can take place.

So, an approach to proof normalisation based on permutations is not necessarily ruled out completely, but appears to be complicated and is not pursued in this paper.

By using the idea in this example, it is not difficult to construct another example (different from the one used in Tiu (2006)) that shows the necessity of deep inference for designing cut-free systems for logics incorporating a self-dual non-commutative connective, like NEL.

Luckily, to prove cut admissibility for NEL we can use the technique called *splitting* in Guglielmi (2007). As that paper explains, this technique establishes a clear connection with the sequent calculus, at least for the fragments of proof systems that allow for a sequent calculus presentation (in our case, the commutative fragment).

See Guglielmi (2007) for an intuitive explanation of splitting – we will only mention here that the technique relies on two separate phases:

(1) *Context reduction*:

If a structure  $S\{R\}$  is provable, then  $S\{ \}$  can be reduced (by performing inference steps going upwards in the derivation) to the structure  $[\{ \} \wp U]$ , for some  $U$ , such that  $[R \wp U]$  is provable (the hole can be filled by any structure that does not play an active part in inference steps).

(2) *Splitting*:

If  $[(R \otimes T) \wp P]$  is provable, then  $P$  can be reduced to  $[P_1 \wp P_2]$  such that  $[R \wp P_1]$  and  $[T \wp P_2]$  are provable.

Context reduction is in turn proved by splitting, which is then at the core of the matter. In this section we will state and prove splitting proper; we will then tackle context reduction in Section 4.

For notational convenience, we define *system* NELc to be the system obtained from NEL by removing the rules for weakening, absorption and digging (which was called *non-core* in Straßburger and Guglielmi (2009)):

$$NELc = NEL \setminus \{w\downarrow, b\downarrow, g\downarrow\} = \{\circ\downarrow, ai\downarrow, s, q\downarrow, p\downarrow, e\downarrow\}.$$

The following lemma was called ‘shallow splitting’ in Guglielmi (2007). The proof is very similar, so we will not give it in full here. But note that we organise the case analysis here in a different way from in Guglielmi (2007) so that we can reuse it for Lemmas 3.3 and 3.4.

**Lemma 3.2 (Splitting).** Let  $R, T, P$  be any NEL structures.

(i) If  $[(R \otimes T) \wp P]$  is provable in NELc, then there are structures  $P_R$  and  $P_T$  such that

$$\begin{array}{c} [P_R \wp P_T] \\ \text{NELc} \parallel \\ P \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \\ [R \wp P_R] \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \\ [T \wp P_T] \end{array}$$

(ii) If  $[\langle R \triangleleft T \rangle \wp P]$  is provable in NELc, then there are structures  $P_R$  and  $P_T$  such that

$$\begin{array}{c} \langle P_R \triangleleft P_T \rangle \\ \text{NELc} \parallel \\ P \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \\ [R \wp P_R] \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \\ [T \wp P_T] \end{array}$$

*Proof.* We prove both statements simultaneously by structural induction on the number of atoms in the conclusion and the length (number of rule instances) of the proof, ordered lexicographically. Without loss of generality, we can assume  $R \neq \circ \neq T$  (otherwise both statements are trivially true).

(i) Consider the bottommost rule instance  $\rho$  in the proof of  $[(R \otimes T) \wp P]$ . We can distinguish three kinds of case:

(a) The first kind appears when the redex of  $\rho$  is inside  $R, T$  or  $P$ . Then we have the following situation:

$$\begin{array}{c} \text{NELc} \parallel \Pi \\ \rho \frac{[(R' \otimes T) \wp P]}{[(R \otimes T) \wp P]} \end{array}$$

where we can apply the induction hypothesis to  $\Pi$  because it is one rule shorter (if  $\rho = \text{ai}\downarrow$  also the conclusion is smaller). We get

$$\begin{array}{c} [P_R \wp P_T] \\ \text{NELc} \parallel \Delta_P \\ P \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \Pi_R \\ [R' \wp P_R] \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \Pi_T \\ [T \wp P_T] \end{array}$$

From  $\Pi_R$ , we can get

$$\begin{array}{c} \text{NELc} \parallel \Pi'_R \\ \rho \frac{[R' \wp P_R]}{[R \wp P_R]} \end{array}$$

and we are done. The situation is similar if the redex of  $\rho$  is inside  $T$  or  $P$ .

(b) The second kind of case is where the substructure  $(R \otimes T)$  is inside the redex of  $\rho$ , but is not modified by  $\rho$ . These cases can be compared with the ‘commutative cases’ in the usual sequent calculus cut-elimination argument. We will only show one representative example – a complete case analysis can be found in Guglielmi (2007)

and Straßburger (2003a). Suppose we have

$$\text{NELc} \Vdash \Pi$$

$$\text{q}\downarrow \frac{[\langle (R \otimes T) \wp P_1 \wp P_3 \rangle \triangleleft P_2 \rangle \wp P_4]}{[(R \otimes T) \wp \langle P_1 \triangleleft P_2 \rangle \wp P_3 \wp P_4]}$$

We can apply the induction hypothesis to  $\Pi$  because it is one rule shorter (the size of the conclusion does not change). This gives us

$$\begin{array}{c} \langle Q_1 \triangleleft Q_2 \rangle \\ \text{NELc} \Vdash \Delta_1 \\ P_4 \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \Vdash \Pi_1 \\ [(R \otimes T) \wp P_1 \wp P_3 \wp Q_1] \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \Vdash \Pi_2 \\ [P_2 \wp Q_2] \end{array}$$

We can apply the induction hypothesis again to  $\Pi_1$ , because the number of atoms in the conclusion is now strictly smaller (because we can assume that the instance of  $\text{q}\downarrow$  is not trivial). We get

$$\begin{array}{c} [P_R \wp P_T] \\ \text{NELc} \Vdash \Delta_2 \\ [P_1 \wp P_3 \wp Q_1] \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \Vdash \Pi_R \\ [R \wp P_R] \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \Vdash \Pi_T \\ [T \wp P_T] \end{array}$$

We can now build the following derivation from  $\Delta_1$ ,  $\Delta_2$  and  $\Pi_2$ :

$$\begin{array}{c} [P_R \wp P_T] \\ \text{NELc} \Vdash \Delta_2 \\ [P_1 \wp P_3 \wp Q_1] \\ \text{NELc} \Vdash \Pi_2 \\ \text{q}\downarrow \frac{[\langle [P_1 \wp Q_1] \triangleleft [P_2 \wp Q_2] \rangle \wp P_3]}{[\langle P_1 \triangleleft P_2 \rangle \wp P_3 \wp \langle Q_1 \triangleleft Q_2 \rangle]} \\ \text{NELc} \Vdash \Delta_1 \\ [\langle P_1 \triangleleft P_2 \rangle \wp P_3 \wp P_4] \end{array}$$

and we are done. All other cases in this group are similar.

- (c) The final type of case is where the substructure  $(R \otimes T)$  is destroyed by  $\rho$ . These cases can be compared to the ‘key cases’ in a standard sequent calculus cut-elimination argument. We have only one possibility. The most general situation is as follows:

$$\text{NELc} \Vdash \Pi$$

$$\text{s} \frac{[\langle [(R_1 \otimes T_1) \wp P_1] \otimes R_2 \otimes T_2 \rangle \wp P_2]}{[(R_1 \otimes R_2 \otimes T_1 \otimes T_2) \wp P_1 \wp P_2]}$$

where one of  $R_1$  and  $R_2$  might be  $\circ$ , but not both of them (and similarly for  $T_1$  and  $T_2$ ). As before, we can apply the induction hypothesis to  $\Pi$  and get

$$\begin{array}{c} [Q_1 \wp Q_2] \\ \text{NELc} \parallel \Delta_1 \\ P_2 \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \Pi_1 \\ [(R_1 \otimes T_1) \wp P_1 \wp Q_1] \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \Pi_2 \\ [(R_2 \otimes T_2) \wp Q_2] \end{array}$$

We can apply the induction hypothesis again to  $\Pi_1$  and  $\Pi_2$ . (Because we assume that the instance of  $\mathfrak{s}$  is non-trivial, the conclusions are strictly smaller than the one of the original proof.) We get

$$\begin{array}{c} [P_{R_1} \wp P_{T_1}] \\ \text{NELc} \parallel \Delta_3 \\ [P_1 \wp Q_1] \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \Pi_{R_1} \\ [R_1 \wp P_{R_1}] \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \Pi_{T_1} \\ [T_1 \wp P_{T_1}] \end{array}$$

and

$$\begin{array}{c} [P_{R_2} \wp P_{T_2}] \\ \text{NELc} \parallel \Delta_4 \\ Q_2 \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \Pi_{R_2} \\ [R_2 \wp P_{R_2}] \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \Pi_{T_2} \\ [T_2 \wp P_{T_2}] \end{array}$$

Now let  $P_R = [P_{R_1} \wp P_{R_2}]$  and  $P_T = [P_{T_1} \wp P_{T_2}]$ . We can build

$$\begin{array}{c} [P_{R_1} \wp P_{R_2} \wp P_{T_1} \wp P_{T_2}] \\ \text{NELc} \parallel \Delta_4 \\ [P_{R_1} \wp P_{T_1} \wp Q_2] \\ \text{NELc} \parallel \Delta_3 \\ [P_1 \wp Q_1 \wp Q_2] \\ \text{NELc} \parallel \Delta_1 \\ [P_1 \wp P_2] \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \Pi_{R_1} \\ [R_1 \wp P_{R_1}] \\ \text{NELc} \parallel \Pi_{R_2} \\ \frac{[(R_1 \otimes [R_2 \wp P_{R_2}]) \wp P_{R_1}]}{\mathfrak{s} [(R_1 \otimes R_2) \wp P_{R_1} \wp P_{R_2}]} \end{array}$$

and a similar proof of  $[(T_1 \otimes T_2) \wp P_{T_1} \wp P_{T_2}]$ , and we are done.

- (ii) The case for  $[(R \triangleleft T) \wp P]$  is similar to the one for  $[(R \otimes T) \wp P]$ , and we leave it as an exercise. □

We can now tackle modalities, for which we can also exhibit a splitting lemma.

**Lemma 3.3 (Splitting for modalities).** Let  $R$  and  $P$  be any NEL structures.

- (i) If  $[!R \wp P]$  is provable in NELc, then there are structures  $P_1, \dots, P_h$  for some  $h \geq 0$  such that

$$\begin{array}{c} [?P_1 \wp \dots \wp ?P_h] \\ \text{NELc} \parallel \\ P \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \\ [R \wp P_1 \wp \dots \wp P_h] \end{array}$$

(ii) If  $[?R \wp P]$  is provable in NELc, then there is a structure  $P_R$  such that

$$\text{NELc} \parallel \begin{array}{c} !P_R \\ P \end{array} \quad \text{and} \quad \text{NELc} \parallel \begin{array}{c} [R \wp P_R] \end{array}$$

*Proof.* The proof is similar to the previous one. We use the same induction measure and the same pattern in the case analysis as before:

(i) We again consider the bottommost rule instance  $\rho$  in the proof of  $[!R \wp P]$ , and we have the same three classes of cases as in the proof of Lemma 3.2:

(a) The redex of  $\rho$  is inside  $R$  or  $P$ .

This case is the same as in the proof of Lemma 3.2.

(b) The substructure  $!R$  is inside the redex of  $\rho$ , but is not changed by  $\rho$ .

This case is almost literally the same as for Lemma 3.2. We only have to replace  $(R \otimes T)$  by  $!R$ , and

$$\text{NELc} \parallel \begin{array}{c} [P_R \wp P_T] \\ \Delta_2 \\ [P_1 \wp P_3 \wp Q_1] \end{array} \quad \text{by} \quad \text{NELc} \parallel \begin{array}{c} [?P_1 \wp \dots \wp ?P_h] \\ \Delta_2 \\ [P_1 \wp P_3 \wp Q_1] \end{array}$$

(As for the previous lemma, the full details can be found in Straßburger (2003a).)

(c) The substructure  $!R$  is destroyed by  $\rho$ .

There are two possibilities ( $\rho = e\downarrow$  and  $\rho = p\downarrow$ ):

$$e\downarrow \frac{\text{NELc} \parallel \begin{array}{c} [o \wp P] \\ [!o \wp P] \end{array}}{\text{NELc} \parallel \begin{array}{c} [o \wp P] \\ [!o \wp P] \end{array}} \quad \text{and} \quad p\downarrow \frac{\text{NELc} \parallel \Pi \quad [![R \wp P_1] \wp Q_2]}{[!R \wp ?P_1 \wp Q_2]}$$

For  $\rho = e\downarrow$ , we are done immediately by letting  $h = 0$ . For  $\rho = p\downarrow$ , we can apply the induction hypothesis to  $\Pi$  and get structures  $P_2, \dots, P_h$  such that

$$\text{NELc} \parallel \begin{array}{c} [?P_2 \wp \dots \wp ?P_h] \\ Q_2 \end{array} \quad \text{and} \quad \text{NELc} \parallel \begin{array}{c} [R \wp P_1 \wp P_2 \wp \dots \wp P_h] \end{array}$$

We then immediately get

$$\text{NELc} \parallel \begin{array}{c} [?P_1 \wp ?P_2 \wp \dots \wp ?P_h] \\ [?P_1 \wp Q_2] \end{array}$$

(ii) As before, we consider the bottommost rule instance  $\rho$  in the proof of  $[?R \wp P]$ :

(a) The redex of  $\rho$  is inside  $R$  or  $P$ .

This case is the same as before.

- (b) The substructure  $?R$  is inside the redex of  $\rho$ , but is not changed by  $\rho$ .  
 As before, this case is almost literally the same as in the proof of Lemma 3.2. This time we have to replace  $(R \otimes T)$  by  $?R$ , and

$$\begin{array}{ccc} [P_R \wp P_T] & & !P_R \\ \text{NELc} \parallel \Delta_2 & \text{by} & \text{NELc} \parallel \Delta_2 \\ [P_1 \wp P_3 \wp Q_1] & & [P_1 \wp P_3 \wp Q_1] \end{array}$$

- (c) The substructure  $?R$  is destroyed by  $\rho$ .  
 For this case there is only one possibility:

$$\begin{array}{c} \text{NELc} \parallel \Pi \\ \rho \downarrow \frac{[![R \wp P_1] \wp P_2]}{[?R \wp !P_1 \wp P_2]} \end{array}$$

We can apply part (i) of the lemma and get

$$\begin{array}{ccc} [?Q_1 \wp \dots \wp ?Q_h] & & \text{NELc} \parallel \Pi_R \\ \text{NELc} \parallel \Delta & \text{and} & [R \wp P_1 \wp Q_1 \wp \dots \wp Q_h] \\ P_2 & & \end{array}$$

Now let  $P_R = [P_1 \wp Q_1 \wp \dots \wp Q_h]$ . We can then build

$$\begin{array}{c} ![P_1 \wp Q_1 \wp \dots \wp Q_h] \\ \{ \rho \downarrow \} \parallel \\ [!P_1 \wp ?Q_1 \wp \dots \wp ?Q_h] \\ \text{NELc} \parallel \Delta \\ [!P_1 \wp P_2] \end{array}$$

as desired. □

**Lemma 3.4 (Splitting for atoms).** Let  $a$  be any atom and  $P$  be any NEL structure. If there is a proof

$$\begin{array}{c} \text{NELc} \parallel \\ [a \wp P] \end{array}$$

then there is a derivation

$$\begin{array}{c} \bar{a} \\ \text{NELc} \parallel \\ P \end{array}$$



*Proof.* After the previous two proofs, this is an almost trivial exercise. Case (a) is as before, and for (b), we have to replace  $(R \otimes T)$  by  $a$ , and

$$\begin{array}{ccc} [P_R \wp P_T] & & \bar{a} \\ \text{NELc} \parallel \Delta_2 & \text{by} & \text{NELc} \parallel \Delta_2 \\ [P_1 \wp P_3 \wp Q_1] & & [P_1 \wp P_3 \wp Q_1] \end{array}$$

For case (c), the only possibility is

$$\begin{array}{c} \text{NELc} \parallel \Pi' \\ \text{ai} \downarrow \frac{P_1}{[a, \bar{a}, P_1]} \end{array}$$

from which we immediately get

$$\begin{array}{c} \bar{a} \\ \text{NELc} \parallel \\ [\bar{a}, P_1] \end{array}$$

as desired. □

#### 4. Context reduction

The idea of context reduction is to reduce a problem that concerns an arbitrary (deep) context  $S\{ \}$  to a problem that only concerns a shallow context  $[\{ \} \wp P]$ . In the case of cut elimination, for example, we will then be able to apply splitting.

Before giving the statement, we need to define the *modality depth* of a context  $S\{ \}$  to be the number of ! and ? in whose scope the  $\{ \}$  occurs. In the following lemma, the  $\{ \}$  is treated as an ordinary atom.

**Lemma 4.1 (Context reduction).** Let  $R$  be a NEL structure and  $S\{ \}$  be a context. If  $S\{R\}$  is provable in NELc, then there is a structure  $P_R$  such that

$$\begin{array}{ccc} ! \cdots ![\{ \} \wp P_R] & & \\ \text{NELc} \parallel \Delta & \text{and} & \text{NELc} \parallel \Pi \\ S\{ \} & & [R \wp P_R] \end{array}$$

where the number of ! in front of  $[\{ \} \wp P_R]$  is the modality depth of  $S\{ \}$ .

*Proof.* We proceed by structural induction on the context  $S\{ \}$ . The base case, when  $S\{ \} = \{ \}$ , is trivial.

We can now distinguish four cases:

- (a)  $S\{ \} = [(S'\{ \} \otimes T) \wp P]$  where, without loss of generality,  $T \neq \circ$ . Note that we do allow  $P = \circ$ . We can apply splitting (Lemma 3.2) to the proof of  $[(S'\{R\} \otimes T) \wp P]$  and get

$$\text{NELc} \parallel \begin{array}{c} [P_S \wp P_T] \\ \Delta_P \\ P \end{array} \quad \text{and} \quad \text{NELc} \parallel \begin{array}{c} \Pi_S \\ [S'\{R\} \wp P_S] \end{array} \quad \text{and} \quad \text{NELc} \parallel \begin{array}{c} \Pi_T \\ [T \wp P_T] \end{array}$$

Because  $T \neq \circ$ , we can now apply the induction hypothesis to  $\Pi_S$  and get

$$\text{NELc} \parallel \begin{array}{c} ! \cdots ![\{ \} \wp P_R] \\ \Delta' \\ [S'\{ \} \wp P_S] \end{array} \quad \text{and} \quad \text{NELc} \parallel \begin{array}{c} \Pi \\ [R \wp P_R] \end{array}$$

From this we can build

$$\begin{array}{c} ! \cdots ![\{ \} \wp P_R] \\ \text{NELc} \parallel \Delta' \\ [S'\{ \} \wp P_S] \\ \text{NELc} \parallel \Pi_T \\ \text{S} \frac{[(S'\{ \} \otimes [T \wp P_T]) \wp P_S]}{[(S'\{ \} \otimes T) \wp P_S \wp P_T]} \\ \text{NELc} \parallel \Delta_P \\ [(S'\{ \} \otimes T) \wp P] \end{array}$$

as desired.

- (b) The cases  $S\{ \} = [(S'\{ \} \triangleleft T) \wp P]$  and  $S\{ \} = [(T \triangleleft S'\{ \}) \wp P]$  are handled similarly to (a).
- (c) For the case  $S\{ \} = [!S'\{ \} \wp P]$ , we can apply splitting (Lemma 3.3) to the proof of  $[!S'\{R\} \wp P]$  and get

$$\text{NELc} \parallel \begin{array}{c} [?P_1 \wp \cdots \wp ?P_h] \\ \Delta_P \\ P \end{array} \quad \text{and} \quad \text{NELc} \parallel \begin{array}{c} \Pi_S \\ [S'\{R\} \wp P_1 \wp \cdots \wp P_h] \end{array}$$

Applying the induction hypothesis to  $\Pi_S$ , we get  $P_R$  such that

$$\text{NELc} \parallel \begin{array}{c} ! \cdots ![\{ \} \wp P_R] \\ \Delta' \\ [S'\{ \} \wp P_1 \wp \cdots \wp P_h] \end{array} \quad \text{and} \quad \text{NELc} \parallel \begin{array}{c} \Pi \\ [R \wp P_R] \end{array}$$

From this we can build

$$\begin{array}{c}
 !! \cdots ![\{ \} \wp P_R] \\
 \text{NELc} \parallel \Delta' \\
 ![S'\{ \} \wp P_1 \wp \cdots \wp P_h] \\
 \{p\downarrow\} \parallel \\
 [!S'\{ \} \wp ?P_1 \wp \cdots \wp ?P_h] \\
 \text{NELc} \parallel \Delta_P \\
 [!S'\{ \} \wp P]
 \end{array}$$

Note that in this case the number of ! in front of  $[\{ \} \wp P_R]$  increases.

(d) The case where  $S\{ \} = [?S'\{ \} \wp P]$  is similar to (c). □

### 5. Elimination of the up fragment

In this section, we will first show four lemmas, which are all easy consequences of splitting and which say that the core up rules of system SNEl are admissible if they are applied in a shallow context  $[\{ \} \wp P]$ . Then we will show how context reduction is used to extend these lemmas to any context. As a result, we get a proof of cut elimination that can be considered modular, in the sense that the four core up rules  $ai\uparrow$ ,  $q\uparrow$ ,  $p\uparrow$  and  $e\uparrow$  are shown to be admissible, each independently of the others.

**Lemma 5.1.** Let  $P$  be a structure and  $a$  be an atom. If  $[(a \otimes \bar{a}) \wp P]$  is provable in NELc, then  $P$  is also provable in NELc.

*Proof.* We apply splitting to the proof of  $[(a \otimes \bar{a}) \wp P]$ . This yields

$$\begin{array}{ccc}
 \begin{array}{c} [P_a \wp P_{\bar{a}}] \\ \text{NELc} \parallel \\ P \end{array} & \text{and} & \begin{array}{c} \text{NELc} \parallel \\ [a \wp P_a] \end{array} & \text{and} & \begin{array}{c} \text{NELc} \parallel \\ [\bar{a} \wp P_{\bar{a}}] \end{array}
 \end{array}$$

By applying Lemma 3.4, we get derivations from  $\bar{a}$  to  $P_a$  and from  $a$  to  $P_{\bar{a}}$ . From these we can build our proof

$$\begin{array}{c}
 \circ\downarrow \frac{\quad}{\circ} \\
 ai\downarrow \frac{\quad}{[\bar{a} \wp a]} \\
 \text{NELc} \parallel \\
 [P_a \wp P_{\bar{a}}] \\
 \text{NELc} \parallel \\
 P
 \end{array}$$

as desired. □

**Lemma 5.2.** Let  $R, T, U, V$  and  $P$  be any NEL structures. If  $[(\langle R \triangleleft U \rangle \otimes \langle T \triangleleft V \rangle) \wp P]$  is provable in NELc, then  $[(R \otimes T) \triangleleft (U \otimes V)] \wp P$  is also provable in NELc.

*Proof.* By applying splitting several times to the proof of  $[(\langle R \triangleleft U \rangle \otimes \langle T \triangleleft V \rangle) \wp P]$ , we get structures  $P_R, P_T, P_U$  and  $P_V$  such that

$$\begin{array}{c} \langle P_R \triangleleft P_U \rangle \wp \langle P_T \triangleleft P_V \rangle \\ \text{NELc} \parallel \\ P \end{array} \quad \text{NELc} \parallel \quad \text{NELc} \parallel \quad \text{NELc} \parallel \quad \text{NELc} \parallel \quad \begin{array}{c} [R \wp P_R] \\ [U \wp P_U] \\ [T \wp P_T] \\ [V \wp P_V] \end{array}$$

Combining these, we can build the proof

$$\begin{array}{c} \text{NELc} \parallel \\ \text{s, s, s, s} \frac{\langle [R \wp P_R] \otimes [T \wp P_T] \triangleleft ([U \wp P_U] \otimes [V \wp P_V]) \rangle}{\langle [(R \otimes T) \wp P_R \wp P_T] \triangleleft [(U \otimes V) \wp P_U \wp P_V] \rangle} \\ \text{q}\downarrow, \text{q}\downarrow \frac{\langle [(R \otimes T) \triangleleft (U \otimes V)] \wp \langle P_R \triangleleft P_U \rangle \wp \langle P_T \triangleleft P_V \rangle \rangle}{\text{NELc} \parallel} \\ [(R \otimes T) \triangleleft (U \otimes V)] \wp P \end{array}$$

as desired. □

**Lemma 5.3.** Let  $R, T$  and  $P$  be any NEL structures. If  $[(?R \otimes !T) \wp P]$  is provable in NELc, then  $[?(R \otimes T) \wp P]$  is also provable in NELc.

*Proof.* As above, we apply splitting several times to the proof of  $[(?R \otimes !T) \wp P]$  and get structures  $P_R, P_1, \dots, P_h$  such that

$$\begin{array}{c} [!P_R \wp ?P_1 \wp \dots \wp ?P_h] \\ \text{NELc} \parallel \\ P \end{array} \quad \text{and} \quad \text{NELc} \parallel \quad \text{and} \quad \text{NELc} \parallel \quad \begin{array}{c} [R \wp P_R] \\ [T \wp P_1 \wp \dots \wp P_h] \end{array}$$

Combining these, we can build the proof

$$\begin{array}{c} \text{NELc} \parallel \\ \text{s, s} \frac{![R \wp P_R] \otimes [T \wp P_1 \wp \dots \wp P_h]}{![R \otimes T] \wp P_R \wp P_1 \wp \dots \wp P_h} \\ \{p\downarrow\} \parallel \\ [?(R \otimes T) \wp !P_R \wp ?P_1 \wp \dots \wp ?P_h] \\ \text{NELc} \parallel \\ [?(R \otimes T) \wp P] \end{array}$$

as desired. □

**Lemma 5.4.** Let  $P$  be any NEL structure. If  $[?\circ \wp P]$  is provable in NELc, then  $[\circ \wp P]$  is also provable in NELc.

*Proof.* We leave this as a trivial exercise. □

Using context reduction (Lemma 4.1), we can extend the statements of Lemmas 5.1–5.4 from shallow contexts  $[\{ \} \otimes P]$  to arbitrary contexts  $S\{ \}$ , which we do in the following lemma.

**Lemma 5.5.** Let  $R, T, U$  and  $V$  be any structures,  $a$  be an atom and  $S\{ \}$  be any context. Then we have the following:

- (i) If  $S(a \otimes \bar{a})$  is provable in NELc, then so is  $S\{\circ\}$ .
- (ii) If  $S(\langle R \triangleleft U \rangle \otimes \langle T \triangleleft V \rangle)$  is provable in NELc, then so is  $S(\langle (R \otimes T) \triangleleft (U \otimes V) \rangle)$ .
- (iii) If  $S(?R \otimes !T)$  is provable in NELc, then so is  $S\{?(R \otimes T)\}$ .
- (iv) If  $S\{?\circ\}$  is provable in NELc, then so is  $S\{\circ\}$ .

*Proof.* All four statements are proved similarly – we will only show the third one here. Let a proof of  $S(?R \otimes !T)$  be given. We apply context reduction to get a structure  $P$  such that

$$\begin{array}{c} ! \cdots ![\{ \} \otimes P] \\ \text{NELc} \parallel \Delta \\ S\{ \} \end{array} \quad \text{and} \quad \begin{array}{c} \text{NELc} \parallel \Pi \\ [?(R \otimes !T) \otimes P] \end{array}$$

By Lemma 5.3, there is a proof  $\Pi'$  of  $[?(R \otimes T) \otimes P]$ , and plugging  $?(R \otimes T)$  into the hole of  $\Delta$ , we can build

$$\begin{array}{c} \{\circ\downarrow, e\downarrow\} \parallel \\ ! \cdots !\circ \\ \text{NELc} \parallel \Pi' \\ ! \cdots ! [?(R \otimes T) \otimes P] \\ \text{NELc} \parallel \Delta \\ S\{?(R \otimes T)\} \end{array}$$

It is obvious that the other statements can be proved in the same way. □

**Lemma 5.6.** If a structure  $R$  is provable in  $\text{NELc} \cup \{\text{ai}\uparrow, \text{q}\uparrow, \text{p}\uparrow, \text{e}\uparrow\}$ , it is also provable in NELc.

*Proof.* The instances of the rules  $\text{ai}\uparrow, \text{q}\uparrow, \text{p}\uparrow$  and  $\text{e}\uparrow$  are removed one after the other (starting with the topmost one) using Lemma 5.5. □

We can now very easily give a proof of the cut-elimination theorem for the system NEL.

*Proof of Theorem 2.8.* Cut elimination is obtained in two steps:

$$\begin{array}{ccc} \begin{array}{c} \circ\downarrow \text{---} \\ \circ \\ \text{SNEL} \parallel \\ R \end{array} & \xrightarrow{1} & \begin{array}{c} \text{NELc} \cup \{\text{ai}\uparrow, \text{q}\uparrow, \text{p}\uparrow, \text{e}\uparrow\} \parallel \\ R' \\ \{\text{w}\downarrow, \text{b}\downarrow, \text{g}\downarrow\} \parallel \\ R \end{array} & \xrightarrow{2} & \begin{array}{c} \text{NELc} \parallel \\ R' \\ \{\text{w}\downarrow, \text{b}\downarrow, \text{g}\downarrow\} \parallel \\ R \end{array} \end{array}$$

Step 1 is an application of the decomposition (Theorem 2.11). The instances of  $g\uparrow$ ,  $b\uparrow$  and  $w\uparrow$  disappear because their premise must be the unit  $\circ$ , which is impossible. Step 2 is just Lemma 5.6.  $\square$

This technique shows how admissibility can be proved uniformly for the cut rule (the atomic one) and the other up rules, which are actually very different rules from the cut. Hence, our technique is more general than cut elimination in the sequent calculus for two reasons:

- (1) It applies to connectives that admit no sequent calculus definition, such as  $\text{seq}$ .
- (2) It can be used to show admissibility of non-infinitary rules that involve no negation, like  $q\uparrow$  and  $p\uparrow$ .

## 6. Perspectives

We think that the techniques developed here for splitting can be exported to the many modal logics already available in deep inference (some of which have no known cut-free presentation in Gentzen formalisms). The reason is that linear logic modalities have a similar behaviour to those of modal logic. This is particularly obvious if we observe that the promotion rule of NEL is the same as the K rule of all modal logics in deep inference (corresponding to the K axiom of basic modal logic). Of course, contraction in linear logic, and in NEL, is restricted, but the splitting theorems, crucially, do not make any use of it.

Apart from its use in proving cut elimination, splitting is a powerful tool for reducing proof search non-determinism in deep inference proof systems. This is explored in Kahramanođulları (2006; 2008b).

We are currently investigating, in the context of the INRIA ARC project REDO, the relations between splitting and the focusing technique in linear logic (Andreoli 1992; Miller 1996), which is at the basis of ludics (Girard 2001). It appears that focusing can be justified and greatly generalised by splitting in deep inference. It seems that splitting is a way to explore the duality between any subformula and its context, so revealing a new logical symmetry.

## Acknowledgments

We thank Dale Miller, as well as David Baelde, Nicolas Guenot and Vivek Nigam for fruitful discussions on focusing and splitting.

## References

- Abramsky, S. and Jagadeesan, R. (1994) Games and full completeness for multiplicative linear logic. *Journal of Symbolic Logic* **59** (2) 543–574.
- Andreoli, J.-M. (1992) Logic programming with focusing proofs in linear logic. *Journal of Logic and Computation* **2** (3) 297–347.
- Blute, R., Panangaden, P. and Straßburger, L. (2008) The logic BV and quantum causality. In: *Trends in Logic VI: Logic and the foundations of physics: space, time and quanta*.

- Blute, R., Panangaden, P. and Slavnov, S. (2009) Deep inference and probabilistic coherence spaces.
- Blute, R. F., Guglielmi, A., Ivanov, I. T., Panangaden, P. and Straßburger, L. (2010) A logical basis for quantum evolution and entanglement.
- Bruscoli, P. (2002) A purely logical account of sequentiality in proof search. In: Stuckey, P.J. (ed.) Logic Programming, 18th International Conference. *Springer-Verlag Lecture Notes in Artificial Intelligence* **2401** 302–316.
- Brünnler, K. (2003) Atomic cut elimination for classical logic. In: Baaz, M. and Makowsky, J.A. (eds.) CSL 2003. *Mathematical Structures in Computer Science* **2803** 86–97.
- Brünnler, K. and Tiu, A. (2001) A local system for classical logic. In: Nieuwenhuis, R. and Voronkov, A. (eds.) LPAR 2001. *Springer-Verlag Lecture Notes in Artificial Intelligence* **2250** 347–361.
- Fleury, A. and Retoré, C. (1994) The mix rule. *Mathematical Structures in Computer Science* **4** (2) 273–285.
- Girard, J.-Y. (1987) Linear logic. *Theoretical Computer Science* **50** 1–102.
- Girard, J.-Y. (2001) Locus solum: From the rules of logic to the logic of rules. *Mathematical Structures in Computer Science* **11** (3) 301–506.
- Guglielmi, A. (2007) A system of interaction and structure. *ACM Transactions on Computational Logic* **8** (1).
- Guglielmi, A. and Gundersen, T. (2008) Normalisation control in deep inference via atomic flows. *Logical Methods in Computer Science* **4** (1:9) 1–36.
- Guglielmi, A. and Straßburger, L. (2002) A non-commutative extension of MELL. In: Baaz, M. and Voronkov, A. (eds.) Logic for Programming, Artificial Intelligence, and Reasoning, LPAR 2002. *Springer-Verlag Lecture Notes in Artificial Intelligence* **2514** 231–246.
- Guglielmi, A., Gundersen, T. and Parigot, M. (2010a) A proof calculus which reduces syntactic bureaucracy. In: Lynch, C. (ed.) 21st International Conference on Rewriting Techniques and Applications. *Leibniz International Proceedings in Informatics (LIPIcs)* **6**, Schloss Dagstuhl–Leibniz-Zentrum für Informatik 135–150.
- Guglielmi, A., Gundersen, T. and Straßburger, L. (2010b) Breaking paths in atomic flows for classical logic. In: *Proceedings of the 25th Annual IEEE Symposium on Logic in Computer Science (LICS 2010)*, IEEE Computer Society 284–293.
- Guglielmi, A. and Straßburger, L. (2001) Non-commutativity and MELL in the calculus of structures. In: Fribourg, L. (ed.) Computer Science Logic, CSL 2001. *Springer-Verlag Lecture Notes in Computer Science* **2142** 54–68.
- Kahramanoğulları, O. (2004) System BV without the equalities for unit. In: Aykanat, C., Dayar, T. and Körpeoğlu, I. (eds.) 19th International Symposium on Computer and Information Sciences, ISCIS 2004. *Springer-Verlag Lecture Notes in Computer Science* **3280** 986–995.
- Kahramanoğulları, O. (2006) Reducing nondeterminism in the calculus of structures. In: Hermann, M. and Voronkov, A. (eds.) LPAR 2006. *Springer-Verlag Lecture Notes in Artificial Intelligence* **4246** 272–286.
- Kahramanoğulları, O. (2008a) System BV is NP-complete. In: de Queiroz, R. and Macintyre, A. (eds.) 12th Workshop on Logic, Language, Information and Computation. *Annals of Pure and Applied Logic* **152** (1-3) 107–121.
- Kahramanoğulları, O. (2008b) Interaction and depth against nondeterminism in proof search.
- Miller, D. (1996) Forum: A multiple-conclusion specification logic. *Theoretical Computer Science* **165** 201–232.
- Okada, M. (1999) Phase semantic cut-elimination and normalization proofs of first- and higher-order linear logic. *Theoretical Computer Science* **227** (1-2) 333–396.

- Retoré, C. (1993) *Réseaux et Séquents Ordonnés*, Ph.D. thesis, Université Paris VII.
- Retoré, C. (1997) Pomset logic: A non-commutative extension of classical linear logic. In: de Groote, Ph. and Hindley, J. R. (eds.) *Typed Lambda Calculus and Applications, TLCA'97. Springer-Verlag Lecture Notes in Computer Science* **1210** 300–318.
- Straßburger, L. (2003a) *Linear Logic and Noncommutativity in the Calculus of Structures*, Ph.D. thesis, Technische Universität Dresden.
- Straßburger, L. (2003b) MELL in the Calculus of Structures. *Theoretical Computer Science* **309** (1-3) 213–285.
- Straßburger, L. (2003c) System NEL is undecidable. In: De Queiroz, R. Pimentel, E. and Figueiredo, L. (eds.) 10th Workshop on Logic, Language, Information and Computation (WoLLIC). *Electronic Notes in Theoretical Computer Science* **84**.
- Straßburger, L. and Guglielmi, A. (2009) A system of interaction and structure IV: The exponentials and decomposition.
- Tiu, A. (2006) A system of interaction and structure II: The need for deep inference. *Logical Methods in Computer Science* **2** (2) 1–24.