



REVIEW

Open Access

# A systematic review of the evidence for Canada's Physical Activity Guidelines for Adults

Daren ER Warburton<sup>1,2\*</sup>, Sarah Charlesworth<sup>1,2</sup>, Adam Ivey<sup>1,2</sup>, Lindsay Nettlefold<sup>1,2</sup>, Shannon SD Bredin<sup>3</sup>

## Abstract

This systematic review examines critically the scientific basis for *Canada's Physical Activity Guide for Healthy Active Living* for adults. Particular reference is given to the dose-response relationship between physical activity and premature all-cause mortality and seven chronic diseases (cardiovascular disease, stroke, hypertension, colon cancer, breast cancer, type 2 diabetes (diabetes mellitus) and osteoporosis). The strength of the relationship between physical activity and specific health outcomes is evaluated critically. Literature was obtained through searching electronic databases (e.g., MEDLINE, EMBASE), cross-referencing, and through the authors' knowledge of the area. For inclusion in our systematic review articles must have at least 3 levels of physical activity and the concomitant risk for each chronic disease. The quality of included studies was appraised using a modified Downs and Black tool. Through this search we identified a total of 254 articles that met the eligibility criteria related to premature all-cause mortality (N = 70), cardiovascular disease (N = 49), stroke (N = 25), hypertension (N = 12), colon cancer (N = 33), breast cancer (N = 43), type 2 diabetes (N = 20), and osteoporosis (N = 2). Overall, the current literature supports clearly the dose-response relationship between physical activity and the seven chronic conditions identified. Moreover, higher levels of physical activity reduce the risk for premature all-cause mortality. The current Canadian guidelines appear to be appropriate to reduce the risk for the seven chronic conditions identified above and all-cause mortality.

## Introduction

There is considerable literature supporting the importance of habitual physical activity in the primary and secondary prevention of varied chronic conditions [1-16]. Routine physical activity is thought to be of benefit for over 25 chronic conditions [17]. Seven chronic diseases in particular have been associated with a physically inactive lifestyle including coronary artery disease, stroke, hypertension, colon cancer, breast cancer, type 2 diabetes (diabetes mellitus) and osteoporosis [18-20].


Canada has played a leading role in the development of physical activity guidelines for individuals across the lifespan. This includes the development (in 1998) of "Canada's Physical Activity Guide to Healthy Active Living" for adults between the ages of 20 and 55 yr [21], which was followed by "Canada's Physical Activity Guide to Healthy Active Living for Older Adults" [22], and "Canada's Physical Activity Guide to Healthy Active Living for Children and Youth" [23]. The adult

guidelines (which are now approximately 10 years old) state generally that 20-55 yr adults should accumulate 60 min of daily physical activity or 30 min of moderate to vigorous exercise on at least 4 days a week [18,19].

We reported recently that Canada's adult guidelines were consistent with other international guidelines and were supported by a compelling body of literature [18,19]. We revealed strong evidence that routine physical activity was effective in the primary prevention of cardiovascular disease, stroke, hypertension, breast cancer, colon cancer, type 2 diabetes and osteoporosis. Moreover, physical activity appears to play an important role in the prevention of obesity and obesity-related co-morbidities. However, implicit in the adult guidelines is the belief that there is a dose-response relationship between physical activity and the associated health benefits. Moreover, a central belief in these guidelines and most international physical activity guidelines is that the dose-response relationship is curvilinear with the greatest health benefits seen in physically inactive individuals who become "more physically active." In fact, a consistent pattern (shown in Figure 1) has been hypothesized, wherein

\* Correspondence: [darenwb@interchange.ubc.ca](mailto:darenwb@interchange.ubc.ca)

<sup>1</sup>Cardiovascular Physiology and Rehabilitation Laboratory, University of British Columbia, Vancouver, Canada



**Figure 1** Theoretical relationship between the risk for chronic disease and physical activity/fitness.

there are marked changes in health status with relatively minor increments in physical activity/fitness in individuals that are the least active/fit. Generally, the health benefits have been thought to level off at the upper end of the physical activity/fitness continuum (Figure 1). However, recent work (such as that provided by Gledhill and Jamnik in the Canadian Physical Activity and Life-style Approach) has speculated that there are likely multiple dose-response curves for various endpoints [24].

The primary purpose of this systematic review was to examine critically the current literature to determine whether or not a dose-response relationship exists between habitual physical activity and chronic disease. In particular, we sought to determine whether the key messaging "Every little bit counts, but more is even better - everyone can do it!" of the adult physical activity guidelines is supported by a strong body of evidence.

Due to the breadth of literature, we have chosen to focus on the relationship between physical activity and all-cause mortality, and the seven chronic conditions that are thought to be reduced greatly with habitual physical activity (i.e., cardiovascular disease (excluding stroke), stroke, hypertension, colon cancer, breast cancer, type 2 diabetes and osteoporosis) (see Table 1). Owing to the nature of the physical activity guidelines, the emphasis of this paper was on primary prevention, despite the clear evidence that routine physical activity is also an effective secondary preventative strategy against many chronic conditions [16,18,19]. Accordingly, our primary objectives were to examine the evidence for a dose-response relationship between: 1) physical activity and all-cause mortality, and 2) physical activity and incidence of the following chronic conditions (cardiovascular disease (except stroke), stroke, hypertension, type 2 diabetes, colon cancer, breast cancer, and osteoporosis.

**Table 1** Relative risks (RR) and population attributable risks (PAR%) for physical inactivity in Canada, Australia, and the USA.

Disease	Canada		Australia		USA	
	RR	PAR%	RR	PAR%	RR	PAR%
CHD	1.45	19.4	1.5	18	2.0	22
Stroke	1.60	24.3	2.0	16	na	Na
Hypertension	1.30	13.8	na	na	1.5	12
Colon Cancer	1.41	18.0	1.5	19	2.0	22
Breast Cancer	1.31	14.2	1.1	9	1.2	5
Type 2 Diabetes	1.50	21.1	1.3	13	1.5	12
Osteoporosis	1.59	24.0	1.4*	18*	2.0	18*

Source: Canadian Data [20]; Australian Data [161]; US Data: [162]. \*Evaluated the incidence of falls/fractures.

## Methods

### Criteria for considering studies for this review

Our research team utilized a rigorous, systematic, and evidence-based approach to examine critically the levels of evidence on physical activity and the risk for premature mortality and chronic disease. Any studies that evaluated the relationship between at least three different levels of physical activity and mortality or incidence of chronic disease were eligible for inclusion. Therefore, excluded studies included those that examined only the most active versus least active populations (e.g., sedentary/inactive vs. physically active). Any form of physical activity/exercise measurement (e.g., self-report, pedometer, accelerometer, maximal aerobic power ( $\text{VO}_2 \text{max}$ ) was eligible for inclusion. The key outcomes were mortality and incidence of chronic disease. Only published, English language studies examining adults (e.g., 19-65 yr) were included. Participants must have previously been healthy (asymptomatic) adults without established chronic disease. There was no restriction according to study design.

To examine the relative risk reductions associated with physical activity, we calculated the mean and median risk reductions across studies focusing on the highest level versus the lowest level of physical activity/fitness. For each study we also determined whether or not a dose-response relationship was present (i.e., reflecting a progressive decrease in the risk with increasing physical activity/fitness levels).

### Search strategy

Literature searches were conducted in the following electronic bibliographical databases:

- MEDLINE (1950-March 2008, OVID Interface);
- EMBASE (1980- March 2008, OVID Interface),
- CINAHL (1982- March 2008, OVID Interface);

- PsycINFO (1840- March 2008, Scholars Portal Interface);
- Cochrane Library (-March 2008),
- SPORTDiscus (-March 2008).

The Medical Subject Headings (MeSH) were kept broad. See tables 2, 3, 4, 5, 6, 7, 8 and 9 for the complete search strategy and keywords used. The electronic search strategies were created and carried out by researchers experienced with systematic reviews of the literature (DW and LN). The citations and applicable electronic versions of the article (where available) were downloaded to an online research management system (RefWorks, Bethesda, Maryland, USA).

**Screening**

Two reviewers (LN and SC) screened independently the title and abstract of the citations to identify potential

**Table 2 Results of the MEDLINE literature search regarding all-cause mortality.**

#	Searches (28 Feb 2008)	Results
1	exp Physical Fitness/	15236
2	Motor Activity/	49721
3	exp Physical Endurance/	15383
4	exp Exercise/	57742
5	exp Exertion/	88903
6	exp Sports/	71887
7	exp exercise therapy/	17231
8	exp exercise tolerance/	4192
9	exp health behaviour/	59409
10	leisure time physical activity	996
11	occupational physical activity.mp	190
12	exp Pilability/	2279
13	exp Muscle Strength/	5717
14	muscs power.mp	965
15	exp Back/	12821
16	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15	291635
17	dose-response.mp	321066
18	intensity.mp	142881
19	volume.mp	298471
20	exp Energy Metabolism/	206808
21	exp oxygen consumption/	83352
22	exp time factors/	763712
23	17 or 18 or 19 or 20 or 21 or 22	1651633
24	16 and 23	67698
25	exp Mortality/	190058
26	all cause mortality.mp	4618
27	25 or 26	192720
28	24 and 27	421
29	limit 28 to (english and humans and "all adult (19 plus years)")	279

**Table 3 Results of the MEDLINE literature search regarding cardiovascular disease.**

Search #	Searches (3 Mar 2008)	Results
1	exp Physical Fitness/	15244
2	Motor Activity/	49751
3	exp Physical Endurance/	15408
4	exp Exercise/	57806
5	exp Exertion/	88967
6	exp Sports/	71931
7	exp exercise therapy/	17243
8	exp exercise tolerance/	4205
9	exp health behaviour/	59467
10	leisure time physical activity.mp	998
11	occupational physical activity.mp	191
12	exp Pilability/	2289
13	exp Muscle Strength/	5731
14	muscs power.mp	965
15	exp Back/	12822
16	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15	291817
17	dose-response.mp	321198
18	intensity.mp	142955
19	volume.mp	298620
20	exp Energy Metabolism/	206886
21	exp oxygen consumption/	83387
22	exp time factors/	764091
23	17 or 18 or 19 or 20 or 21 or 22	1652372
24	16 and 23	67760
25	exp Cardiovascular Diseases/	1411730
26	exp Heart diseases/	675083
27	exp Myocardial Infarction/	116070
28	exp Death, Sudden Cardiac/	6772
29	exp Coronary Artery Disease/	18137
30	exp Coronary Disease/	144236
31	exp Vascular Diseases	1018275
32	25 or 26 or 27 or 28 or 29 or 30 or 31	1411730
33	24 and 32	9603
34	limit 33 to (english language and humans and "all adult (19 plus years)")	5544

articles for inclusion. Duplicate citations were removed. The reviewers were not blinded to the authors or journals. Biographies of key studies and reviews in the field were also cross-referenced for further articles. For those articles that appeared relevant, the full text was obtained and data was extracted using a common template. In cases of disagreement, discussion with a third reviewer (DW) was used to achieve consensus. Full (100%) consensus was achieved. All studies that were excluded during the citation and full-article screening processes were recorded along with the reasons for exclusion.

**Table 4 Results of the MEDLINE literature search regarding stroke.**

Search #	Searches (29 Feb 2008)	Results
1	exp Physical Fitness/	15241
2	Motor Activity/	49744
3	exp Physical Endurance/	15387
4	exp Exercise/	57764
5	exp Exertion/	88921
6	exp Sports/	71907
7	exp exercise therapy/	17237
8	exp exercise tolerance/	4196
9	exp health behaviour/	59430
10	leisure time physical activity.mp	996
11	occupational physical activity.mp	190
12	exp Piability/	2288
13	exp Muscle Strength/	5720
14	muscs power.mp	965
15	exp Back/	12821
16	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15	291718
17	dose-response.mp	321133
18	intensity.mp	142919
19	volume.mp	298526
20	exp Energy Metabolism/	206837
21	exp oxygen consumption/	83359
22	exp time factors/	763871
23	17 or 18 or 19 or 20 or 21 or 22	1651958
24	16 and 23	67720
25	exp Stroke/	45243
26	exp Cerebrovascular Disorders/	196243
27	exp Brain Ischemia/	58943
28	exp Brain Infarction/ or exp Cerebral Infarction	21357
29	exp Infarction, Middle Cerebral Artery/ or exp Intracranial Aneurysm/ or exp Subarachnoid	46725
30	Hemorrhage/ or exp Cerebral Hemorrhage/exp Ischemic Attack, Transient/	14753
31	25 or 26 or 27 or 28 or 29 or 30	196243
32	24 and 31	692
33	limit 32 to (english language and humans and "all adult (19 plus years)")	291

### Data Extraction

Two reviewers (LN and SC) completed standardized data extraction forms, which were verified by two other reviewers (DW and SB). We extracted information regarding the study design, the country where the study was conducted, the participant characteristics, the sample size, the objectives of the study, the methodologies employed, the major outcomes (i.e., mortality, incidence of chronic disease, physical activity levels/classifications), and the comments and conclusions made based on the findings of the study. The reviewers were not blinded to the journal or the author names when extracting information from the articles.

**Table 5 Results of the MEDLINE literature search regarding hypertension.**

Search #	Searches (3 Mar 2008)	Results
1	exp Physical Fitness/	15244
2	Motor Activity/	49751
3	exp Physical Endurance/	15408
4	exp Exercise/	57806
5	exp Exertion/	88967
6	exp Sports/	71931
7	exp exercise therapy/	17243
8	exp exercise tolerance/	4205
9	exp health behaviour/	59467
10	leisure time physical activity.mp	998
11	occupational physical activity.mp	191
12	exp Piability/	2289
13	exp Muscle Strength/	5731
14	muscs power.mp	965
15	exp Back/	12822
16	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15	291817
17	dose-response.mp	3211987
18	intensity.mp	142955
19	volume.mp	298620
20	exp Energy Metabolism/	206886
21	exp oxygen consumption/	83387
22	exp time factors/	764091
23	17 or 18 or 19 or 20 or 21 or 22	1652372
24	exp Hypertension/	168466
25	exp Blood Pressure/	205571
26	exp Blood Pressure Determination/ or exp Blood Pressure Monitoring, Ambulatory/ or exp Blood	18244
27	Pressure Monitors/24 or 25 or 26	336025
28	16 and 23 and 27	5647
29	limit 28 to (english language and humans and "all adult (19 plus years)")	3642

### Level of Evidence

The approach used to establish the level and grade of evidence was consistent with that used during creation of the "Canadian clinical practice guidelines on the management and prevention of obesity in adults and children" [25]. The level of evidence provides information regarding the strength of the evidence in favour of physical activity/exercise in the primary prevention of premature mortality and the seven chronic diseases of primary interest. This evaluation process is based on a pre-defined and objective criteria (see Table 10).

The grade for each article provides information regarding whether physical activity is effective in the primary prevention of the varied conditions evaluated (Table 10). Where applicable this grade informs the reader about the potential risk of the physical activity. A study that receives the highest grading would indicate

**Table 6 Results of the MEDLINE literature search regarding colon cancer.**

Search #	Searches (3 Mar 2008)	Results
1	exp Physical Fitness/	15244
2	Motor Activity/	49751
3	exp Physical Endurance/	15408
4	exp Exercise/	57806
5	exp Exertion/	88967
6	exp Sports/	71931
7	exp exercise therapy/	17243
8	exp exercise tolerance/	4205
9	exp health behaviour/	59467
10	leisure time physical activity.mp	998
11	occupational physical activity.mp	191
12	exp Piability/	2289
13	exp Muscle Strength/	5731
14	muscs power.mp	965
15	exp Back/	12822
16	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15	291817
17	dose-response.mp	321198
18	intensity.mp	142955
19	volume.mp	298620
20	exp Energy Metabolism/	206886
21	exp oxygen consumption/	83387
22	exp time factors/	764091
23	17 or 18 or 19 or 20 or 21 or 22	1652372
24	exp Colonic Neoplasms/	51780
25	exp Rectal Neoplasms/	28011
26	exp Colorectal Neoplasms/	99982
27	exp Colorectal Neoplasms/, Hereditary Nonpolyposis/ or exp Intestinal Neoplasms.	117563
28	24 or 25 or 26 or 27	117563
29	16 and 23 and 28	108
30	limit 29 to (53nenglish language and humans and "all adult (19 plus years)")	77

that the benefits clearly outweigh the risks and receive a strong recommendation.

#### Quality Assessment

The quality of each study was also established using the procedures of Gorber et al. [26]. Owing to the fact that only observational study designs were included in our systematic review, we used the Downs and Black [27] scale to assess the quality of non-randomized investigations. Similar to the work of Prince et al. [28] we chose to include the most relevant components of the scoring tool. Therefore, a modified version of the Downs and Black checklist was used with the final checklist consisting of 15 items with a maximum score of 15 points. Higher points reflected a superior quality of investigation.

**Table 7 Results of the MEDLINE literature search regarding breast cancer.**

Search #	Searches (28 Feb 2008)	Results
1	exp Physical Fitness/	15236
2	Motor Activity/	49721
3	exp Physical Endurance/	15383
4	exp Exercise/	57742
5	exp Exertion/	88903
6	exp Sports/	71887
7	exp exercise therapy/	17231
8	exp exercise tolerance/	4192
9	exp health behaviour/	59409
10	leisure time physical activity.mp	996
11	occupational physical activity.mp	190
12	exp Piability/	2279
13	exp Muscle Strength/	5717
14	muscs power.mp	965
15	exp Back/	12821
16	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15	291635
17	dose-response.mp	321066
18	intensity.mp	142881
19	volume.mp	298471
20	exp Energy Metabolism/	206808
21	exp oxygen consumption/	83352
22	exp time factors/	763712
23	17 or 18 or 19 or 20 or 21 or 22	1651633
24	exp Breast Neoplasms/	149817
25	16 and 23 and 24	296
26	limit 25 to (54 nenglish language and humans and "all adult (19 plus years)"	216

#### Results

##### Physical Inactivity and All-Cause Mortality

A total of 2040 citations were identified during the electronic database search (Figure 2). Of these citations, 288 were identified in MEDLINE, 222 in EMBASE, 496 in Cochrane, and 1034 in the CINAHL/SportDiscus/PsychInfo search. A total of 167 duplicates were found, leaving a total of 1873 unique citations. A total of 1696 articles were excluded after scanning, leaving a total of 177 articles for full review. From these articles 130 were excluded after full review leaving 47 articles for inclusion in the systematic review. An additional 23 articles were added to the review based on the authors' knowledge of the area. The reasons for exclusion included review articles (n = 26), commentary (n = 10), did not report 3 levels of physical activity (n = 24), no objective measure of physical activity (n = 2), report (n = 15), not a formal study (n = 11), not related to all-cause mortality (n = 27), the participants were too young (n = 1), not able to retrieve articles (n = 7), and other (n = 7). Therefore, a total of 70 articles were included in the

**Table 8 Results of the MEDLINE literature search regarding type 2 diabetes.**

Search #	Searches (29 Feb 2008)	Results
1	exp Physical Fitness/	15241
2	Motor Activity/	49744
3	exp Physical Endurance/	15387
4	exp Exercise/	57764
5	exp Exertion/	88921
6	exp Sports/	71907
7	exp exercise therapy/	17237
8	exp exercise tolerance/	4196
9	exp health behaviour/	59430
10	leisure time physical activity.mp	996
11	occupational physical activity.mp	190
12	exp Piability/	2288
13	exp Muscle Strength/	5720
14	muscs power.mp	965
15	exp Back/	12821
16	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15	291718
17	dose-response.mp	321133
18	intensity.mp	142919
19	volume.mp	298526
20	exp Energy Metabolism/	206837
21	exp oxygen consumption/	83359
22	exp time factors/	763871
23	17 or 18 or 19 or 20 or 21 or 22	1651958
24	16 and 23	67720
25	exp Blood Glucose/or exp Diabetes Mellitus, Type 2/	132583
26	exp Hyperglycemia/	16214
27	exp Glucose Intolerance/ or exp Glucose Tolerance Test/	24986
28	exp Hyperinsulinism/	30490
29	25 or 26 or 27 or 28	165157
30	29 and 24	3006
31	Limit 30 to (english language and humans and "all adult (19 plus years)")	1985

systematic review of the literature regarding the relationship between physical activity and premature mortality.

The majority of the studies included in our systematic review were prospective cohort investigations (Table 11). These studies involved a total of 1,525,377 participants; averaging 21,791 participants per study (range 302-252,925). There were a total of 111,125 reported cases of premature all-cause mortality (ranging per study from 43-10,952). The total length of study follow-up for the prospective cohort studies averaged 11.1 yr (ranging from 0.5-28 yr). The articles were published over a 22 yr period ranging from 1985 to 2007. These studies involved large samples of men and women from regions throughout the world.

**Table 9 Results of the MEDLINE literature search regarding osteoporosis.**

Search #	Searches (29 feb 2008)	Results
1	exp Physical Fitness/	15241
2	Motor Activity/	49744
3	exp Physical Endurance/	15387
4	exp Exercise/	57764
5	exp Exertion/	88921
6	exp Sports/	71907
7	exp exercise therapy/	17237
8	exp exercise tolerance/	4196
9	exp health behaviour/	59430
10	leisure time physical activity.mp	996
11	occupational physical activity.mp	190
12	exp Piability/	2288
13	exp Muscle Strength/	5720
14	muscs power.mp	965
15	exp Back/	12821
16	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15	291718
17	dose-response.mp	321133
18	intensity.mp	142919
19	volume.mp	298526
20	exp Energy Metabolism/	206837
21	exp oxygen consumption/	83359
22	exp time factors/	763871
23	17 or 18 or 19 or 20 or 21 or 22	1651958
24	exp Osteoporosis, Postmenopausal/ or exp Osteoporosis/	31532
25	exp Fractures, Bone/ or exp Bone Density/	125269
26	exp Bone Diseases/ or exp Bone Diseases, Metabolic/	308084
27	exp "Bone and bones"/	369634
28	exp Tensile Strength/	12050
29	exp Compressive Strength	2838
30	24 or 25 or 26 or 27 or 28 or 29	642158
31	16 and 23 and 30	2138
32	Limit 31 to (english language and humans and "all adult (19 plus years)")	1193

We observed a mean 31% lower risk for all-cause mortality in the most active individuals. The median risk reduction was 32%. It is important to highlight that many of these studies included women, with sub-analyses that revealed similar risk reductions between sexes. Our findings are consistent with previous reports [15,16,29-31]. The majority (90%) of the studies supported the health benefits of physical activity demonstrating a significant risk reduction in physically active individuals. *The level of evidence would be considered to be a Level 2A based on the presence of overwhelming evidence from observational trials.* The studies examined were generally of a good quality with a mean (and median) score of 12 out of 15 (range 10-14).

**Table 10 The levels and grade of evidence scaling criteria applied to the articles.**

Level of Evidence	Criteria
Level 1	Randomized control trials without important limitations
Level 2	<ul style="list-style-type: none"><li>• Randomized control trials with important limitations</li><li>• Observational studies (non-randomized clinical trials or cohort studies) with overwhelming evidence</li></ul>
Level 3	Other observational studies (prospective cohort studies, case-control studies, case series)
Level 4	Inadequate or no data in population of interest Anecdotal evidence or clinical experience
Grade of Evidence	Criteria
Grade A	Strong recommendation (action can apply to most individuals in most circumstances) <ul style="list-style-type: none"><li>• Benefits clearly outweigh risks (or vice-versa)</li><li>• Evidence is at Level 1, 2, or 3</li></ul>
Grade B	Weak recommendation (action may differ depending on individual's characteristics or other circumstances) <ul style="list-style-type: none"><li>• Unclear if benefits outweigh risks</li><li>• Evidence is at Level 1, 2, or 3</li></ul>
Grade C	Consensus recommendation (alternative actions may be equally reasonable) <ul style="list-style-type: none"><li>• Unclear if benefits outweigh risks</li><li>• Evidence is at Level 3 or 4</li></ul>

A clear dose-response relationship was also observed with marked reductions in the risk for all-cause mortality occurring with relatively small increments in physical activity (Figure 3). To examine more closely the temporal relationship between physical activity and all-cause mortality we calculated the (unadjusted) relative risks associated with incremental levels of physical activity/fitness using the reported cases of all-cause mortality and the number of participants (per group) in each investigation. In some instances, we were required to calculate the number of participants based on the reported incidence rates and person years, or based on data obtained directly from the authors (2 investigations). We were not able to obtain this information in 18 investigations, and as such this analysis was restricted to the remaining 52 investigations. There was considerable variability in the methods of classifying the physical activity/fitness levels of the participants. Accordingly, Figure 3 illustrates the mean relative risk reduction according to three separate study types including those that subdivided participants into tertiles, quartiles and quintiles, respectively. This figure demonstrates clearly the dose-response relationship between physical activity and all-cause mortality. Collectively, the literature is consistent indicating that the current Canadian guidelines (approximately 4.2 MJ/wk, 1000 kcal/wk) are associated with a 20-30% lower risk for premature all-cause mortality, with greater health benefits with high volumes and/or intensities of activity. In our analyses it was apparent that the greatest differences in risk occurred between the lowest adjacent activity/fitness categories, suggesting that sedentary individuals can markedly


reduce their risk for all-cause mortality with relatively minor increments in physical activity. This is consistent with the current messaging of Canada's physical activity guidelines.

The strength of the relationship between physical fitness and premature mortality has been well-established [6,32,33]. In our analyses there were greater risk reductions in studies that took objective measures of physical fitness. We observed an average risk reduction of approximately 45%, which was consistent between men and women. A risk reduction of greater than 50% was not uncommon in these studies. For instance, Myers et al. (2004) reported that being fit or physically active was associated with greater than 50% lower mortality risk in men. They also noted that a 4.2 MJ/wk (1000 kcal/wk) increase in physical activity, or a 1 metabolic equivalent (MET) higher physical fitness level was associated with a mortality benefit of around 20%. It is also important to highlight that longitudinal studies evaluating changes in physical activity or fitness have revealed a lower premature mortality risk [16,34-41]. As we previously reported, routine physical activity or elevated physical fitness also appears to reduce the risk for premature mortality in individuals with risk factors for chronic disease [42,43].

**Implications**

Since the seminal work of Morris and colleagues (in the 1950s [44,45]) and the early work of Paffenbarger (in the 1970s [46,47]) there has been considerable research (especially epidemiological evidence) documenting the health benefits of engaging in routine physical activity and/or being physically fit [17,48]. Both physical activity





fitness and all-cause mortality. As demonstrated below and in Table 11 and Figure 1, this literature is extensive.

The assessment of the relationship between all-cause mortality is complicated by the inclusion of deaths related to suicides, homicide, and accidents [18,19,52]. Nonetheless, the available evidence is incontrovertible; individuals who are habitually physically active and/or physically fit are at a markedly reduced risk for premature all-cause mortality [15,16,18,19]. In Canada, physical inactivity is a major cause of premature mortality from diseases of the cardiovascular system (33.3%), cancers (29.1%), and type 2 diabetes (3.5%) [53]. Globally, physical inactivity has been linked with 2 million premature deaths per year, including 22% of cases of coronary heart disease, and 10-16% of cases of breast cancer, colon cancer, rectal cancer and type 2 diabetes [54]. As such, the promotion of the health benefits of physical activity is of paramount importance for the effective prevention of chronic disease and premature mortality on a national and international scale.

In summary, there is a clear dose-response relationship between physical activity and premature all-cause mortality. Physically active individuals have an approximate risk reduction of 31% in comparison to physically inactive individuals. When objective measures of aerobic fitness are taken the risk reductions are even greater approximating 45%.

#### **Recommendation #1**

*For a reduced risk for premature mortality, it is recommended that individuals should participate in 30 min or more of moderate to vigorous exercise on most days of the week. Greater health benefits appear to occur with higher volumes and/or intensities of activity. [Level 2, Grade A]*

#### **Primary Prevention of Cardiovascular Disease**

In our systematic search of the literature, a total of 9408 citations were identified during the electronic database search (Figure 4). Of these citations, 5973 were identified in MEDLINE, 2561 in EMBASE, 193 in Cochrane, and 681 in the CINAHL/SportDiscus/PsychInfo search. A total of 923 duplicates were found, leaving a total of 8485 unique citations. A total of 8138 articles were excluded after scanning, leaving a total of 347 articles for full review. An additional 20 articles were added through cross-referencing. From these articles 319 were excluded after full review leaving 33 articles for inclusion in the systematic review. The reasons for exclusion included non-experimental studies (n = 45), only effect on cardiovascular disease risk factors (n = 115), did not report 3 levels of physical activity (n = 12), subjects less than 18 yr of age (n = 4), reviews, summaries, dissertations, thesis, and abstracts (n = 30), clinical population (n = 14), not on cardiovascular disease or did not fit

(a behaviour) and physical fitness (an attained state) appear to be related to health status in a dose-dependent fashion, with physical fitness demonstrating the strongest relationship [18,19]. Numerous reports indicate that physical inactivity and/or low physical fitness are associated with an increased risk for chronic disease and premature all-cause and disease-specific mortality [2,43,49-51]. Some of the most compelling research includes the relationship between physical activity/



**Table 11 Studies examining the relationship between physical activity and all-cause mortality.**

Publication Country Study Design Quality Score	Objective	Population	Methods	Outcome	Comments and Conclusions
Blair et al 1989 [7]	To study physical fitness (PF) and risk of all-cause mortality in men and women.	<ul style="list-style-type: none"> <li>• n = 13,344 (10,224 men; 3,120 women)</li> <li>• Sex: Men and women</li> </ul>	Baseline and 8 year follow-up	<ul style="list-style-type: none"> <li>• 283 deaths</li> </ul>	Low levels of PF increase the risk for premature mortality.
USA		<ul style="list-style-type: none"> <li>• Age: 20-&gt;60 years (yr)</li> </ul>	PF assessment: Maximal treadmill exercise test.	Adjusted risk ratio (RR), 95% confidence interval (CI)	
Prospective cohort		<ul style="list-style-type: none"> <li>• Characteristics: Participants were given a preventative Medicine examination including maximal treadmill exercise test</li> </ul>	Fitness categorized into quintiles:	Men	
D & B score = 12			Q1 = least fit Q2 Q3 Q4 Q5 = most fit	<ul style="list-style-type: none"> <li>• Q1 = 3.44 (2.05-5.77)</li> <li>• Q2 = 1.37 (0.76-2.50)</li> <li>• Q3 = 1.46 (0.81-2.63)</li> <li>• Q4 = 1.17 (0.63-2.17)</li> <li>• Q5 = 1.00 (referent)</li> </ul> Women <ul style="list-style-type: none"> <li>• Q1 = 4.65 (2.22-9.75)</li> <li>• Q2 = 2.42 (1.09-5.37)</li> <li>• Q3 = 1.43 (0.60-3.44)</li> <li>• Q4 = 0.76 (0.27-2.11)</li> <li>• Q5 = 1.00 (referent)</li> </ul>	
Myers et al 2004 [32]	To determine the effects of PF and physical activity (PA) on all-cause mortality.	<ul style="list-style-type: none"> <li>• n = 6,213</li> <li>• Sex: Men</li> <li>• Age: Mean 59.0 ± 11.2 yr</li> <li>• Characteristics: Men referred for exercise testing</li> </ul>	Baseline and mean 5.5 ± 2.0 year follow-Up	<ul style="list-style-type: none"> <li>• 1,256 deaths</li> </ul>	Being fit or active is associated with >50% reductions in mortality risk.
USA			PF assessment: Treadmill test to measure VO <sub>2</sub> peak	PF Level hazard ratio (HR) (95% CI) <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	PF predicted mortality more strongly than PA.
Prospective cohort				<ul style="list-style-type: none"> <li>• G2 = 0.59 (0.52-0.68)</li> <li>• G3 = 0.46 (0.39-0.55)</li> <li>• G4 = 0.28 (0.23-0.34)</li> </ul>	Increasing PA (by 1000 kcal/wk or 1 MET) confers a mortality benefit of 20%.
D & B score = 12			PA assessment: Self reported PA divided into 4 groups	PA Level HR (95% CI) <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

			G2	• G2 = 0.63 (0.36-1.10)	
			G3	• G3 = 0.42 (0.23-0.78)	
			G4 = Highest level	• G4 = 0.38 (0.19-0.73)	
Blair et al 1995 [36]	To evaluate the relationship between changes in PF and risk of mortality in men.	• n = 9,777	4.9 year mean follow-up	• 223 deaths	Men who maintained or increased adequate PF had a reduced risk for all-cause mortality than individuals who were consistently unfit.
USA		• Sex: Men • Age: 20-82 yr • Characteristics: Participants were given a preventative medicine examination including maximal treadmill exercise test	PF assessment: Maximal exercise test at baseline and follow-up	RR (95% CI) • G1 = 1.00 (referent)	
Prospective cohort				• G2 = 0.56 (0.41-0.75) • G3 = 0.52 (0.38-0.70) • G4 = 0.33 (0.23-0.47)	
D & B score = 13			Groups based on changes in PF G1 = unfit to unfit G2 = unfit to fit G3 = fit to unfit G4 = fit to fit		
Bijnen et al 1999 [37]	To examine the association of PA at baseline and 5 years	• n = 472	1985 and 1990	• 118 deaths	Recent levels of PA were more important for mortality risk than PA 5 years previously.
Netherlands	previously with all- cause mortality risk in a cohort of elderly Dutch men.	• Sex: Men • Age: >65 yr • Characteristics: Mostly independently living elders (~95%) • Zutphen Elderly Study	PA assessment: Questionnaire, divided into tertiles: Lowest Middle Highest	Multivariate adjusted RR (95% CI)  PA in 1985: Lowest tertile = 1.00 (referent) Middle tertile  • Total activity = 1.25 (0.79- 1.99) • Walking = 0.97 (0.60-1.57) • Bike = 0.97 (0.59-1.57) • Gardening = 0.66 (0.39-1.10) • Other = 1.08 (0.66-1.78) • Heavy activity = 0.73 (0.45-1.17) • Non heavy activity = 0.89 (0.57-1.40)	Becoming or remaining sedentary increased the mortality risk.
Retrospective cohort					
D & B score = 12					

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

	<p>Highest tertile</p> <ul style="list-style-type: none"> <li>• Total activity = 1.25 (0.73-2.12)</li> <li>• Walking = 0.94 (0.58-1.55)</li> <li>• Bike = 1.07 (0.61-1.88)</li> <li>• Gardening = 0.77 (0.42-1.39)</li> <li>• Other = 1.24 (0.74-2.07)</li> <li>• Heavy activity = 0.76 (0.44-1.32)</li> <li>• Non heavy activity = 0.94 (0.58-1.53)</li> </ul> <p>PA in 1990:  Lowest tertile = 1.00 (referent)</p> <p>Middle tertile</p> <ul style="list-style-type: none"> <li>• Total activity = 0.56 (0.35-0.89)</li> <li>• Walking = 0.82 (0.51-1.32)</li> <li>• Bike = 0.49 (0.29-0.82)</li> <li>• Gardening = 1.67 (1.00-2.79)</li> <li>• Other = 0.93 (0.53-1.65)</li> <li>• Heavy activity = 1.19 (0.73-1.92)</li> <li>• Non heavy activity = 0.61 (0.38-0.99)</li> </ul> <p>Highest tertile</p> <ul style="list-style-type: none"> <li>• Total activity = 0.44 (0.25-0.80)</li> <li>• Walking = 1.17 (0.70-1.96)</li> <li>• Bike = 0.43 (0.23-0.80)</li> <li>• Gardening = 1.03 (0.55-1.94)</li> <li>• Other = 0.74 (0.44-1.23)</li> <li>• Heavy activity = 0.72 (0.40-1.31)</li> <li>• Non heavy activity = 0.65 (0.40-1.05)</li> </ul>
--	---

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Gregg et al 2003 [39]	To examine the relationship of changes in PA and mortality among older women.	• n = 9,518	Baseline (1986-1988) and median 10.6 year follow-up (1992-1994)	• 2,218 deaths	Increasing and maintaining PA levels could lengthen life for older women but appears to provide less benefit for women aged at least 75 years and those with poor health status.
USA		• Sex: Women	PA Assessment: Questionnaire, divided into quintiles of PA (kcal/wk)		
		• Age: ≥ 65 yr		Multivariate adjusted HRR (95% CI): Quintiles of total	
		• Characteristics: White community dwelling participants from 4 US research centres			
Prospective cohort			Q1= <163 Q2 = 163-503 Q3 = 504-1045 Q4 = 1046-1906 Q5 = ≥ 1907	PA • Q1 = 1.00 (referent) • Q2 = 0.73 (0.64-0.82) • Q3 = 0.77 (0.68-0.87) • Q4 = 0.62 (0.54-0.71) • Q5 = 0.68 (0.59-0.78)	
D & B score = 13			Quintiles of walking(kcal/wk) Q1 = <70 Q2 = 70-186 Q3 = 187-419 Q4 = 420-897 Q5 = 898	Walking HRR (95% CI) • Q1 = 1.00 (referent) • Q2 = 0.91 (0.81-1.02) • Q3 = 0.78 (0.68-0.88) • Q4 = 0.71 (0.63-0.82) • Q5 = 0.71 (0.62-0.82)	
				Multivariate adjusted HRR (95% CI) Change in activity level: Sedentary at baseline • Staying sedentary = 1.00 (referent) • Became active = 0.52 (0.40-0.69) Mod / high active at baseline • Became sedentary = 0.92 (0.77-1.09) • Stayed active = 0.68 (0.56-0.82)	
Wannamethee et al 1998 [40]	To study the relationship between heart rate, PA and all-cause mortality.	• n = 5,934	Baseline (1978-1980) and 12-14 year follow-up	• 219 deaths	Maintaining or taking up light or moderate PA reduces mortality in older men.

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

UK	Prospective cohort	<ul style="list-style-type: none"> <li>Sex: Men</li> <li>Age: Mean 63 yr</li> <li>Characteristics: Healthy, sedentary (4,311 were considered "healthy" in 1992)</li> <li>The British Regional Heart Study</li> </ul>	PA assessment: Questionnaire, split into groups  PA score G1 = Inactive/occasional G2 = Light G3 = Moderate G4 = Moderately vigorous/Vigorous Regular walking (min/d) G1 = 0 G2 = <20 G3 = 21-40 G4 = 41-60 G5 = ≥ 60 Recreational activity, 4 groups G1 = Inactive/fairly Inactive G2 = Average 4 hr/weekend G3 = Fairly active >4 h/weekend G4 = Very active Sporting activity, 3 Groups G1 = None G2 = Occasional G3 = >1 time/month	Multivariate adjusted RR (95% CI), PA  • G1 = 1.00 (referent) • G2 = 0.61 (0.43-0.86) • G3 = 0.50 (0.31-0.79) • G4 = 0.65 (0.45-0.94)  Regular walking • G1 = 1.00 (referent) • G2 = 1.15 (0.73-1.79) • G3 = 1.06 (0.75-1.50) • G4 = 0.97 (0.65-1.46) • G5 = 0.62 (0.37-1.05)  Recreational activity • G1 = 1.00 (referent) • G2 = 0.95 (0.43-1.07) • G3 = 0.68 (0.43-1.07) • G4 = 0.34 (0.35-1.00)  Sporting activity • G1 = 1.00 (referent) • G2 = 0.50 (0.25-1.03) • G3 = 0.88 (0.64-1.23)	
Paffenbarger et al 1986 [63]	To examine the PA and life-style characteristics of Harvard alumni for the relationship with all-cause mortality.	<ul style="list-style-type: none"> <li>n = 16,936</li> <li>Sex: Men</li> <li>Age: 35-74</li> <li>Characteristics: Harvard alumni</li> </ul>	12-16 year follow-up (1962 to 1978)  Records of freshman year physical examinations and records of intercollegiate sport	• 1,413 deaths  Age adjusted RR (95% CI):  Those who walked  • G1 = 1.00 (referent)	The findings suggest a protective effect of exercise against all-cause mortality.

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

D & B score = 14			PA assessment: Mailed questionnaires surveying post college PA	• G2 = 0.85 • G3 = 0.79 Trend $p = 0.0009$	Physical Activity Index (95% CI):
				• G1 = 1.00 (referent) • G2 = 0.78 • G3 = 0.73 • G4 = 0.63 • G5 = 0.62	
			Exercise reported: Walking (miles/wk) 3 groups G1 = <3 G2 = 3-8 G3 = ≥ 9	• G6 = 0.52 • G7 = 0.46 • G8 = 0.62 Trend $p = <0.0001$	
				PA index (kcal/wk) 3 groups: G1 = <500 G2 = 500-999 G3 = 1000-1499 G4 = 1500-1999 G5 = 2000-2499 G6 = 2500-2999 G7 = 3000-3499 G8 = >3500 Cox proportional hazard models	
Schnohr et al 2007 [64]	To determine the impact of walking duration and intensity on all-cause mortality.	• n = 7,308 (3,204 male; 4,104 female)	Baseline and an average of 12 year follow-up	• 1,391 deaths	The findings indicate that the relative intensity and not duration of walking is the most important in relation to all-cause mortality.
Denmark		• Sex: Male and female • Age: 20-93 yr	PA assessment: Questionnaire, 4 durations and 3 intensities	Multivariate adjusted HR (95% CI):	
Prospective cohort		• Characteristics: Participants with no history of CHD, stroke or cancer and who had no difficulty in walking		Men	
D & B score = 12		• The Copenhagen City Heart Study		• G1 = 1.00 (referent) • G2 = 0.38 (0.25-0.58)	





**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

			<p>G1 = Rarely/never G2 = 1 time/week to a few times/month</p> <p>G3 = 2-4 times/week G4 = &gt;4 times/week</p> <p>Activity index G1 = Low G2 = Medium G3 = High</p>	<p>Trend p = &lt;0.001</p> <p>Frequency of vigorous PA per week</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.83 (0.69-0.99)</li> <li>• G3 = 0.74 (0.59-0.93)</li> <li>• G4 = 0.62 (0.42-0.90)</li> </ul> <p>Trend p = 0.009</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.77 (0.69-0.86)</li> <li>• G3 = 0.68 (0.60-0.77)</li> </ul> <p>Trend p = &lt;0.001</p>	
Paffenbarger et al 1993 [67]	To analyze changes in the lifestyles of Harvard College alumni and the association of these changes with mortality.	<ul style="list-style-type: none"> <li>• n = 10,269</li> <li>• Sex: Men</li> <li>• Age: 45-84 yr (in 1977)</li> </ul>	Baseline (1977) and 8 year follow-up (1985)	<ul style="list-style-type: none"> <li>• 476 deaths</li> </ul> <p>Beginning moderate sports activity was associated with 23% lower risk of death (95% CI 4%-42%, <math>p = 0.015</math>) than those not taking up moderate activity</p>	Beginning moderately vigorous sports activity was associated with lower rates of death from all causes among middle aged and older men.
USA		<ul style="list-style-type: none"> <li>• Characteristics: Participants with no reported life- threatening disease</li> </ul>	PA Assessment: Questionnaire – blocks walked daily, stairs climbed daily and type, frequency and duration of weekly sports and recreational activities		
Prospective cohort					
D & B score = 13			Physical activity index (kcal/wk) Sports and recreational activities Light <4.5 METs Moderate >4.5 METs Weekly lists of deaths were obtained from the Harvard college alumni office		

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

			Proportional hazard models with Poisson regression methods		
Katzmarzyk and Craig 2002 [154]	To quantify the relationship between musculoskeletal fitness and all-cause mortality.	<ul style="list-style-type: none"> <li>• n = 8,116 (3,933 male; 4,183 female)</li> </ul>	Baseline (1981) and	<ul style="list-style-type: none"> <li>• 238 deaths</li> </ul>	Some components of musculoskeletal fitness are predictive of mortality.
		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>	13 year follow-up	RR (95% CI) adjusted for age, smoking status, body mass and VO <sub>2max</sub>	
Canada			Musculoskeletal fitness (sit ups, push ups, grip strength, sit and reach) measures divided into quartiles		
Prospective cohort		<ul style="list-style-type: none"> <li>• Age: 20-69 yr</li> <li>• Characteristics: Participants who had musculoskeletal fitness measurements taken</li> </ul>	Q1 = lowest	Sit ups	
			Q2	Men	
			Q3	<ul style="list-style-type: none"> <li>• Q1 = 2.72 (1.56-4.64)</li> </ul>	
			Q4 = highest	<ul style="list-style-type: none"> <li>• Q2 = 1.32 (0.73-2.41)</li> <li>• Q3 = 1.61 (0.90-2.87)</li> <li>• Q4 = 1.00 (referent)</li> </ul>	
D & B score = 11		<ul style="list-style-type: none"> <li>• Canadian Fitness Survey</li> </ul>	Cox proportional hazard ratio model	Women	
				<ul style="list-style-type: none"> <li>• Q1 = 2.26 (1.15-4.43)</li> <li>• Q2 = 2.24 (1.07-4.67)</li> <li>• Q3 = 1.27 (0.59-2.72)</li> <li>• Q4 = 1.00 (referent)</li> </ul>	
				Push-ups	
				Men	
				<ul style="list-style-type: none"> <li>• Q1 = 1.25 (0.77-2.05)</li> <li>• Q2 = 1.17 (0.71-1.90)</li> <li>• Q3 = 0.94 (0.55-1.62)</li> <li>• Q4 = 1.00 (referent)</li> </ul>	
				Women	
				<ul style="list-style-type: none"> <li>• Q1 = 0.61 (0.32-1.17)</li> <li>• Q2 = 0.81 (0.45-1.47)</li> <li>• Q3 = 0.87 (0.48-1.58)</li> <li>• Q4 = 1.00 (referent)</li> </ul>	
				Grip strength (kg)	
				Men	
				<ul style="list-style-type: none"> <li>• Q1 = 1.49 (0.86-2.59)</li> </ul>	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				<ul style="list-style-type: none"> <li>• Q2 = 1.42 (0.82-2.45)</li> <li>• Q3 = 1.59 (0.95-2.68)</li> <li>• Q4 = 1.00 (referent)</li> </ul>	
				Women	
				<ul style="list-style-type: none"> <li>• Q1 = 1.08 (0.58-1.99)</li> <li>• Q2 = 0.62 (0.44-1.56)</li> <li>• Q3 = 1.25 (0.70-2.23)</li> <li>• Q4 = 1.00 (referent)</li> </ul>	
				Sit and reach (cm)	
				Men	
				<ul style="list-style-type: none"> <li>• Q1 = 1.06 (0.64-1.74)</li> <li>• Q2 = 1.01 (0.61-1.66)</li> <li>• Q3 = 1.20 (0.74-1.95)</li> <li>• Q4 = 1.00 (referent)</li> </ul>	
				Women	
				<ul style="list-style-type: none"> <li>• Q1 = 1.18 (0.66-2.10)</li> <li>• Q2 = 1.07 (0.60-1.91)</li> <li>• Q3 = 0.77 (0.44-1.46)</li> <li>• Q4 = 1.00 (referent)</li> </ul>	
Andersen et al 2000 [163]	To evaluate the relationship between levels of OPA, LTPA, cycling to work and sports participation and all-cause mortality.	<ul style="list-style-type: none"> <li>• n = 30,640 (17,265 men; 13,375 women)</li> </ul>	14.5 year follow-up	<ul style="list-style-type: none"> <li>• 8,549 deaths</li> </ul>	LTPA was inversely associated with all-cause mortality in both men and women in all age groups.
Denmark			PA assessment: Questionnaire for LTPA, divided into:	Incidence of all-cause mortality and PA	
Prospective cohort		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: 20-93 years (yr)</li> </ul>		Multivariate adjusted RR (95% CI)	
			G1 = Low		
		<ul style="list-style-type: none"> <li>• Characteristics: Participants of the Copenhagen City Heart Study, Glostrup Population Study and Copenhagen Male Study</li> </ul>	G2 = Moderate		
D & B score = 13			G3 = High	Age 20-44 yr	
				Men	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.73 (0.56-0.96)</li> <li>• G3 = 0.74 (0.55-1.01)</li> </ul>	
				Women	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				<ul style="list-style-type: none"> <li>• G2 = 0.75 (0.54-1.04)</li> <li>• G3 = 0.66 (0.42-1.05)</li> </ul> <p>Age 45-64 yr</p> <p>Men</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.75 (0.67-0.84)</li> <li>• G3 = 0.75 (0.67-0.85)</li> </ul> <p>Women</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.73 (0.65-0.83)</li> <li>• G3 = 0.66 (0.56-0.77)</li> </ul> <p>Age &gt;65 yr</p> <p>Men</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.62 (0.53-0.73)</li> <li>• G3 = 0.60 (0.50-0.72)</li> </ul> <p>Women</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.52 (0.45-0.61)</li> <li>• G3 = 0.49 (0.39-0.61)</li> </ul> <p>All age groups</p> <p>Men</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.72 (0.66-0.78)</li> <li>• G3 = 0.71 (0.65-0.78)</li> </ul> <p>Women</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.65 (0.60-0.71)</li> <li>• G3 = 0.59 (0.52-0.67)</li> </ul>	
Barengo et al 2004 [164]	To investigate whether moderate or high LTPA are associated with reduced CVD and all-cause mortality, independent of CVD risk factors and other forms of PA in men and women.	<ul style="list-style-type: none"> <li>• n = 31,677 (15,853 men; 16,824 women)</li> <li>• Sex: Men and women</li> </ul>	20 year follow-up	HRR (95% CI)	Moderate and high levels of LTPA and OPA are associated with reduced premature all-cause mortality.
Finland		<ul style="list-style-type: none"> <li>• Age: 30-59 yr</li> <li>• Characteristics: Participants from eastern and south-western Finland</li> </ul>	PA assessment: Questionnaire self administered to measure OPA, LTPA and commuting activity	LTPA	<ul style="list-style-type: none"> <li>• 1.00 (referent) = low</li> <li>• 0.91 (0.84-0.98) = mod, Men</li> </ul>

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Prospective cohort					
D & B score = 14					
			<ul style="list-style-type: none"> <li>• 0.79 (0.70-0.90) = high, Men</li> <li>• 0.89 (0.81-0.98) = mod, women</li> <li>• 0.98 (0.83-1.16) = high, women</li> </ul>		
			OPA		
			<ul style="list-style-type: none"> <li>• 1.00 (referent) = low</li> <li>• 0.75 (0.68-0.83) = mod, men</li> <li>• 0.77 (0.71-0.84) = active, men</li> <li>• 0.79 (0.70-0.89) = mod, women</li> <li>• 0.78 (0.70-0.87) = active, women</li> </ul>		
Bath 2003 [165]	To examine differences between older men and women on the self-rated health mortality relationship.	• n = 1,042 (406 men; 636 women at baseline)	Baseline, 4 and 12 years post	Number of deaths: At 4 years 242 (106 men; 136 women)	The self-rated health-mortality relationship can be explained by health and related factors among older men and women.
UK		• Sex: Men and women		• At 12 years 665 (287 men; 378 women)	
Prospective cohort		• Age: >65 yr • Characteristics: Community-dwelling Elderly	General physical health 14-item health index (Ebrahin et al 1987) scoring from 0-14 (no health problems – multiple health problems)	Multivariate adjusted HR (95% CI)	
D & B score = 11					
			<ul style="list-style-type: none"> <li>• The Nottingham Longitudinal Study of Activity and Ageing</li> </ul>		
			Men after 4 years <ul style="list-style-type: none"> <li>• High = 1.00 (referent)</li> <li>• Med = 1.19 (0.61-2.33)</li> <li>• Low = 1.51 (0.75-3.03)</li> </ul>		
			PA assessment: Self-rated health surveys, divided into 3 levels of PA: High Medium Low		
			Women after 4 years <ul style="list-style-type: none"> <li>• High = 1.00 (referent)</li> <li>• Med = 1.03 (0.58-1.82)</li> <li>• Low = 1.51 (0.86-2.67)</li> </ul>		
			Men after 12 years		



**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

			Cox proportional hazards regression Models	<ul style="list-style-type: none"> <li>• High = 1.00 (referent)</li> <li>• Med = 1.28 (0.94-1.74)</li> <li>• Low = 1.13 (0.82-1.55)</li> </ul> <p>Women after 12 years</p> <ul style="list-style-type: none"> <li>• High = 1.00 (referent)</li> <li>• Med = 1.20 (0.90-1.61)</li> <li>• Low = 1.23 (0.93-1.62)</li> </ul>	
Bijnen et al 1998 [166]	To describe the association between PA and mortality (CVD, stroke, all-cause) in elderly men.	<ul style="list-style-type: none"> <li>• n = 802</li> <li>• Sex: Men</li> <li>• Age: 64-84 yr</li> </ul>	10 year follow-up	<ul style="list-style-type: none"> <li>• 373 deaths</li> </ul>	PA may protect against all- cause mortality in elderly men
Netherlands		<ul style="list-style-type: none"> <li>• Characteristics: Retired Dutch men</li> </ul>	PA assessment: Questionnaire, divided into groups:	Multivariate adjusted RR (95% CI)	
Prospective cohort			<p>G1 = Lowest</p> <p>G2 = Middle</p> <p>G3 = Highest</p>	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.80 (0.63-1.02)</li> <li>• G3 = 0.77 (0.59-1.00)</li> </ul> <p>p = 0.04</p>	
D & B score = 12					
Blair et al 1993 [167]	To evaluate the relationship of sedentary living habits to all-cause mortality in women.	<ul style="list-style-type: none"> <li>• n = 3,120</li> <li>• Sex: Women</li> <li>• Age: Not available</li> </ul>	Baseline and 8 year follow-up	<ul style="list-style-type: none"> <li>• 43 deaths</li> </ul>	There is a graded inverse relationship between PF and all-cause mortality in women.
USA		<ul style="list-style-type: none"> <li>• Characteristics: Participants were given a preventative medicine examination</li> </ul>	PF assessment: PF measured via maximal treadmill exercise test;	Age adjusted death rates (per 10,000 person years) by fitness	
Prospective				<ul style="list-style-type: none"> <li>• Low Fitness = 40</li> <li>• Mod Fitness = 16</li> <li>• High Fitness = 7</li> </ul>	The lack of relationship between PA and death rate was believed to be due to an inadequate assessment of PA.
D & B score = 14			PA assessment: Questionnaire	No difference between levels of PA	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Blair et al 1996 [168]	To review the association of PF to all-cause and CVD mortality.	<ul style="list-style-type: none"> <li>• n = 32,421 (25,341 men; 7,080 women)</li> </ul>	Baseline and average 8 year follow-up (range 0.1-19.1 years)	<ul style="list-style-type: none"> <li>• 601 deaths in men</li> <li>• 89 deaths in women</li> </ul>	The study observed a steep inverse gradient of death rates across low, moderate and high PF levels. The association was strong and remained after adjustment for potential confounding factors.
USA		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: 20-80 yr (mean 43 yr)</li> </ul>	PF assessment: Treadmill test; duration was used to assign participants to sex specific groups:	RR (95% CI) in low PF vs. high PF	
Prospective cohort		<ul style="list-style-type: none"> <li>• Characteristics: Participants were excluded if they did not reach 85% of their age predicted maximal heart rate on the maximal exercise treadmill test</li> </ul>		Men	
D & B score = 14			Low (least fit 20%) Moderate (next 40%) High (most fit 40%)	<ul style="list-style-type: none"> <li>• 1.52 (1.28-1.82)</li> <li>Women</li> <li>• 2.10 (1.36-3.26)</li> </ul>	Adjusted deaths per 10,000 person years according to PF
		<ul style="list-style-type: none"> <li>• Aerobics Center Longitudinal Study</li> </ul>	Proportional hazard modeling	<ul style="list-style-type: none"> <li>Men</li> <li>• Low = 49</li> <li>• Med = 27</li> <li>• High = 23</li> <li>Women</li> <li>• Low = 29</li> <li>• Med = 13</li> <li>• High = 14</li> </ul>	
Boyle et al 2007 [169]	To examine the association between PA and the risk of incident disability, including impairment in activities of daily living and instrumental activities of daily living in community based older persons free from dementia.	<ul style="list-style-type: none"> <li>• n = 1,020</li> <li>• Sex: Men and women</li> <li>• Age: 54-100 yr</li> </ul>	2.6 year follow-up	<ul style="list-style-type: none"> <li>• 156 deaths</li> </ul>	The risk of death decreased 11% with each hour of PA/wk.
			PA assessment: Questionnaire, hr/wk of PA Incidence of all-cause mortality	HR for all-cause mortality	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

USA		<ul style="list-style-type: none"><li>• Characteristics: Participants from 40 retirement communities across Chicago</li></ul>		The risk of death was 11% lower for each hr/wk of PA	
Prospective cohort					
D & B score = 13		<ul style="list-style-type: none"><li>• Rush Memory and Aging Project</li></ul>			
Bucksch et al 2005 [170]	To examine the effect of moderately intense PA on all-cause mortality.	<ul style="list-style-type: none"><li>• n = 7,187 (3,742 men; 3,445 women)</li><li>• Sex: Men and women</li></ul>	Baseline (1984-1986) and 12-14 yr follow-up (1998)	<ul style="list-style-type: none"><li>• 943 deaths</li></ul> RR (95% CI) for achieving recommended PA vs. not achieving recommendation	Participants who achieved recommended amounts of MPA or VPA were at a significantly lower risk of death than their sedentary counterparts.
Germany		<ul style="list-style-type: none"><li>• Age: 30-69 yr</li><li>• Characteristics: Participants were healthy and physically active during leisure time</li></ul>	PA assessment: Questionnaire (Minnesota Leisure Time Physical Activity questionnaire) divided into groups based on: Achieving recommended amount of MPA (30 min, 5 d/wk (≥2.5 h/wk))		
Prospective cohort				Women <ul style="list-style-type: none"><li>• MPA = 0.65 (0.51-0.82)</li><li>• VPA = 0.78 (0.57-1.08)</li><li>• MPA or VPA = 0.60 (0.47-0.75)</li></ul> Men <ul style="list-style-type: none"><li>• MPA = 0.90 (0.77-1.01)</li><li>• VPA = 0.74 (0.61-0.90)</li><li>• MPA or VPA = 0.80 (0.68-0.94)</li></ul>	
D & B score = 13			Achieving recommended amount of VPA (20 min, 3 d/wk (≥ 1 h/wk))	RR (95% CI) for volume of lifestyle activities (kcal/kg/wk)	
			Volume of lifestyle activities (kcal/kg/wk)	Women	
			G1 = 0	<ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li></ul>	
			G2 = <14	<ul style="list-style-type: none"><li>• G2 = 0.79 (0.57-1.08)</li></ul>	
			G3 = 14-33.5	<ul style="list-style-type: none"><li>• G3 = 0.68 (0.50-0.94)</li></ul>	
			G4 = ≥ 33.5	<ul style="list-style-type: none"><li>• G4 = 0.57 (0.41-0.79)</li></ul>	
				p < 0.001	
				Men	
				<ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li></ul>	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				<ul style="list-style-type: none"> <li>• G2 = 0.98 (0.76-1.17)</li> <li>• G3 = 0.80 (0.63-1.00)</li> <li>• G4 = 0.91 (0.74-1.13)</li> </ul> <p><math>p = 0.20</math></p> <p>Adjusted for age, other recommendation, social class, smoking, BMI, cardio risk factor index, alcohol intake, chronic disease index and dietary factors</p>	
Bucksch and Helmert 2004 [171]	To examine LTPA and premature death in the general population of former West Germany.	<ul style="list-style-type: none"> <li>• n = 7,187 (3,742 men; 3,445 women)</li> <li>• Sex: Men and women</li> <li>• Age: 30-69 yr</li> <li>• Characteristics: Participants were selected on the basis of the German Cardiovascular Prevention Study</li> </ul>	Baseline (1984-1986) and 12-14 year follow-up (1998)	<ul style="list-style-type: none"> <li>• 943 deaths</li> </ul> <p>RR (95% CI)</p> <p>Men, LTPA</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	LTPA is inversely associated with all-cause mortality in men and women.
Germany			PA assessment: Questionnaire (Minnesota Leisure Time Physical Activity questionnaire) divided into groups based on: LTSA (h/wk)		
Prospective cohort				<ul style="list-style-type: none"> <li>• G2 = 0.85 (0.78-0.93)</li> <li>• G3 = 0.64 (0.50-0.82)</li> <li>• G4 = 0.70 (0.54-0.91)</li> </ul> <p><math>p &lt; 0.001</math></p> <p>Men, LTPA index</p>	
D & B score = 14		<ul style="list-style-type: none"> <li>• The National Health Survey of the German Federal Institute of Population Research (1984-1998)</li> </ul>	<p>G1 = 0</p> <p>G2 = &lt;1</p> <p>G3 = 1-2</p> <p>G4 = &gt;2</p> <p>The LTSA-index (kcal/kg/wk)</p> <p>G1 = 0</p> <p>G2 = 1-10</p> <p>G3 = 10-25</p> <p>G4 = &gt;25</p> <p>Mortality – Records from the mandatory population registries</p>	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.92 (0.70-1.23)</li> <li>• G3 = 0.89 (0.69-1.17)</li> <li>• G4 = 0.61 (0.44-0.84)</li> </ul> <p><math>p &lt; 0.01</math></p> <p>Women, LTPA</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.93 (0.82-1.04)</li> <li>• G3 = 0.69 (0.48-0.98)</li> <li>• G4 = 0.57 (0.35-0.94)</li> </ul> <p><math>p &lt; 0.01</math></p> <p>Women, LTPA index</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

			Cox proportional hazard regression model	<ul style="list-style-type: none"> <li>• G2 = 0.68 (0.45-1.01)</li> <li>• G3 = 0.79 (0.51-1.21)</li> <li>• G4 = 0.46 (0.25-0.85)</li> <li><math>p &lt; 0.01</math></li> <li>Adjusted for age, social class, smoking, BMI, cardio risk factor index, alcohol intake, chronic disease index and dietary factors</li> </ul>	
Carlsson et al 2006 [172]	To investigate the association between PA and mortality in post-menopausal women.	<ul style="list-style-type: none"> <li>• n = 27,734</li> <li>• Sex: Women</li> <li>• Age: 51-83 yr</li> </ul>	Baseline (1997) and 2-7 year follow-up (1999-2004)	<ul style="list-style-type: none"> <li>• 1,232 deaths</li> </ul>	The study indicates that even fairly small amounts of activity will reduce mortality in older women.
Sweden		<ul style="list-style-type: none"> <li>• Characteristics: Women who participated in a population based Screening programme in 1987</li> </ul>		RR (95% CI) adjusted for lifestyle and medical problems	
Prospective cohort			PA assessment: Questionnaires for: METs/day, different PA (walking/biking), LTPA, OPA, household PA, TV watching and reading	PA (METs/day)	
D & B score = 12		<ul style="list-style-type: none"> <li>• The Swedish Mammography Cohort</li> </ul>		<ul style="list-style-type: none"> <li>• &gt;50 = 1.00 (referent)</li> <li>• 45-50 = 1.05 (0.77-1.42)</li> <li>• 40-45 s = 1.09 (0.81-1.46)</li> <li>• 45-40 = 1.26 (0.94-1.70)</li> <li>• &lt;35 = 2.56 (1.85-3.53)</li> </ul>	
			Mortality – Records from the National Population Register	Different PA Walking/biking (min/d) <ul style="list-style-type: none"> <li>• &gt; 90 = 1.00 (referent)</li> <li>• 60-90 = 1.01 (0.76-1.34)</li> <li>• 40-60 = 0.92 (0.70-1.20)</li> <li>• 20-40 = 0.96 (0.75-1.23)</li> <li>• &lt;20 = 1.16 (0.90-1.50)</li> <li>• Almost never = 1.94 (1.51-2.50)</li> </ul>	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

			LTPA (hr/wk) • >5 = 1.00 (referent) • 4-5 = 0.95 (0.74-1.22) • 2-3 = 1.02 (0.83-1.26) • 1 = 1.09 (0.88-1.36) • <1 = 1.91 (1.56-2.35) OPA • Heavy manual labour = 1.00 (referent) • Walking/lifting/ a lot carrying = 0.96 (0.55-1.70) • Walking/lifting/ not a lot carrying = 1.00 (0.60-1.68) • Mostly standing = 0.91 (0.52-1.61) • Seated 50% of time = 0.97 (0.58-1.62) • Mostly sedentary = 1.93 (1.15-3.25) Household work (hr/d) • >8 h/d = 1.00 (referent) • 7-8 = 0.68 (0.49-0.93) • 5-6 = 0.66 (0.51-0.87) • 3-4 = 0.83 (0.64-1.06) • 1-2 = 0.89 (0.69-1.15) • <1 = 1.73 (1.30-2.32) Adjusted for age		
Crespo et al 2002 [173]	To study the relationship between PA and obesity with all-cause mortality in Puerto Rican men.	• n = 9,136 (1962-1965)	Baseline and 12 year follow-up	• 1,445 deaths	Some PA is better than none in protecting against all-cause mortality. The benefits are independent of body weight.
Puerto Rico		• Sex: Men	PA assessment: Questionnaire, divided into 4 groups based on METs G1 = low G2 G3 G4 = high	Multivariate OR (95% CI) adjusted for age	
Prospective cohort		• Age: 35-79 yr	Multivariate logistic function model	• C1 = 1.00 (referent)	
D & B score = 12		• Characteristics: Participants with no known coronary heart disease		• C2 = 0.67 (0.57-0.78)	



**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

		<ul style="list-style-type: none"> <li>• The Puerto Rico Heart Health Program</li> </ul>		<ul style="list-style-type: none"> <li>• C3 = 0.63 (0.54-0.74)</li> <li>• C4 = 0.54 (0.46-0.64)</li> <li><math>p &lt; 0.0001</math></li> <li>Multivariate adjusted OR (95% CI)</li> <li>• C1 = 1.00 (referent)</li> <li>• C2 = 0.68 (0.58-0.79)</li> <li>• C3 = 0.63 (0.54-0.75)</li> <li>• C4 = 0.55 (0.46-0.65)</li> <li><math>p &lt; 0.0001</math></li> </ul>	
Davey Smith et al 2000 [174]	To examine the relationship of PA and various causes of death.	<ul style="list-style-type: none"> <li>• n = 6,702 (at baseline)</li> </ul>	Baseline (1969-1970) and 25 year follow-up	<ul style="list-style-type: none"> <li>• 926 deaths</li> </ul>	In the study, an inverse association of both LTPA and walking pace with mortality from all-causes was seen.
UK		<ul style="list-style-type: none"> <li>• Sex: Men</li> </ul>	PA assessment: Questionnaire with 3 groups for walking pace (Slower, same, faster) and 3 groups for LTPA (inactive, moderately active, active)	Age adjusted RR (95% CI) for walking pace	
Prospective cohort D & B score = 13		<ul style="list-style-type: none"> <li>• Age: 40-64 yr</li> <li>• Characteristics: Participants from rural northern Japan</li> <li>• Whitehall study</li> </ul>		<ul style="list-style-type: none"> <li>• Slower = 2.47 (2.2-2.8)</li> <li>• Same = 1.35 (1.2-1.5)</li> <li>• Faster = 1.00 (referent)</li> <li><math>p &lt; 0.001</math></li> <li>Fully adjusted RR (95% CI) for walking pace</li> <li>• Slower = 1.87 (1.6-2.1)</li> <li>• Same = 1.21 (1.1-1.3)</li> <li>• Faster = 1.00 (referent)</li> <li><math>p &lt; 0.001</math></li> <li>Age adjusted RR (95% CI) for LTPA</li> <li>• Inactive = 1.44 (1.3-1.6)</li> <li>• Mod = 1.13 (1.0-1.2)</li> <li>• Active = 1.00 (referent)</li> <li><math>p &lt; 0.001</math></li> <li>Fully adjusted RR (95% CI) for LTPA</li> <li>• Inactive = 1.20 (1.1-1.3)</li> <li>• Mod = 1.07 (1.0-1.2)</li> <li>• Active = 1.00 (referent)</li> <li><math>p &lt; 0.001</math></li> </ul>	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Eaton et al 1995 [175]	To determine whether self-reported PA predicts a decreased rate of CHD and all-cause mortality in middle aged men.	<ul style="list-style-type: none"> <li>• n = 8,463</li> </ul>	21 year follow-up	<ul style="list-style-type: none"> <li>• 2,593 deaths</li> </ul>	Baseline levels of self-reported LTPA predicted a decreased rate of CHD and all-cause mortality.
Europe, Israel, mid eastern Asia, Northern Africa		<ul style="list-style-type: none"> <li>• Sex: Men</li> </ul>	PA assessment: Questionnaire for LTPA	Age adjusted RR (95% CI) LTPA	
Prospective cohort		<ul style="list-style-type: none"> <li>• Age: ≥40 yr</li> </ul>	G1 = Sedentary	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
D & B score = 12		<ul style="list-style-type: none"> <li>• Characteristics: Government employees without known CVD</li> </ul>	G2 = Light	<ul style="list-style-type: none"> <li>• G2 = 0.84 (0.74-0.94)</li> </ul>	
			G3 = Light daily	<ul style="list-style-type: none"> <li>• G3 = 0.81 (0.73-0.90)</li> </ul>	
			G4 = Heavy	<ul style="list-style-type: none"> <li>• G4 = 0.84 (0.72-0.98)</li> </ul>	
			Questionnaire for OPA	OPA	
			G1 = Sitting	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
			G2 = Standing	<ul style="list-style-type: none"> <li>• G2 = 0.99 (0.88-1.12)</li> </ul>	
			G3 = Walking	<ul style="list-style-type: none"> <li>• G3 = 1.09 (0.99-1.20)</li> </ul>	
			G4 = Physical labour	<ul style="list-style-type: none"> <li>• G4 = 1.16 (1.03-1.30)</li> </ul>	
Fang et al 2005 [176]	To assess the association of exercise and CVD outcome among persons with different blood pressure status.	<ul style="list-style-type: none"> <li>• n = 9,791 (3,819 men; 5,972 women)</li> </ul>	17 year follow-up	Incidence of all-cause mortality and PA	A significant effect of exercise on mortality in normotensive subjects was not found.
USA		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>	PA assessment: Questionnaire with 3 groups	Multivariate adjusted HR (95% CI)	
Prospective cohort		<ul style="list-style-type: none"> <li>• Age: 25-74 yr</li> </ul>	G1 = Least exercise	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
D & B score = 12		<ul style="list-style-type: none"> <li>• Characteristics: Non-institutionalized participants</li> </ul>	G2 = Moderate exercise	<ul style="list-style-type: none"> <li>• G2 = 0.75 (0.53-1.05)</li> </ul>	
			G3 = Most exercise	<ul style="list-style-type: none"> <li>• G3 = 0.71 (0.45-1.12)</li> </ul>	
Fried et al 1998 [177]	To determine the disease, functional and personal characteristics that jointly predict mortality.	<ul style="list-style-type: none"> <li>• n = 5,886</li> </ul>	5 year follow-up	<ul style="list-style-type: none"> <li>• 646 deaths</li> </ul>	PA was a predictor of 5-year mortality.
USA		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>	PA assessment: Self reported exercise (5 groups)	Incidence of all-cause mortality and PA	
Prospective cohort		<ul style="list-style-type: none"> <li>• Age: ≥65 yr</li> </ul>	MPA or VPA (kJ/wk)	Multivariate adjusted RR (95% CI)	
D & B score = 11		<ul style="list-style-type: none"> <li>• Characteristics: Community dwelling elders</li> </ul>	G1 = ≤282	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
			G2 = 283-1789	<ul style="list-style-type: none"> <li>• G2 = 0.78 (0.60-1.00)</li> </ul>	
			G3 = 1790-4100	<ul style="list-style-type: none"> <li>• G3 = 0.81 (0.63-1.05)</li> </ul>	
			G4 = 4101-7908	<ul style="list-style-type: none"> <li>• G4 = 0.72 (0.55-0.93)</li> </ul>	
			G5 = >7908	<ul style="list-style-type: none"> <li>• G5 = 0.56 (0.43-0.74)</li> </ul>	
				p < 0.005	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Fujita et al 2004 [178]	To examine the relationship between walking duration and all-cause mortality in a Japanese cohort.	• n = 41,163 (20,004 men; 21,159 women)	Baseline (1990) and 11 year follow-up (2001)	• 1,879 deaths	Time spent walking was associated with a reduced risk for all-cause mortality.
Japan		• Sex: Men and women	PA assessment: Questionnaire Walking, 3 levels:  G1 = ≤30 min G2 = 30 min to 1 hr G3 = ≥1 hr	Age and sex adjusted RR (95% CI) for time spent walking (hr/d)	
Prospective cohort D & B score = 13		• Age: 40-64 yr • Characteristics: Healthy, sedentary	Cox proportional hazard model	Whole group • G1 = 1.22 (1.09-1.35) • G2 = 1.09 (0.95-1.22) • G3 = 1.00 (referent) p < 0.001 Men only • G1 = 1.14 (1.00-1.30) • G2 = 1.03 (0.90-1.19) • G3 = 1.00 (referent) p = 0.061 Women only • G1 = 1.40 (1.16-1.68) • G2 = 1.23 (1.01-1.49) • G3 = 1.00 (referent) p < 0.001 RR (95% CI) for time spent walking (hr/d) (adjusted for age, education, marital status, past history of diseases, smoking, drinking, BMI and dietary variables) Whole group • G1 = 1.17 (1.04-1.31) • G2 = 1.06 (0.93-1.20) • G3 = 1.00 (referent) p = 0.011 Men • G1 = 1.08 (0.94-1.25) • G2 = 0.98 (0.84-1.14) • G3 = 1.00 (referent) p = 0.318 Women • G1 = 1.38 (1.12-1.70)	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				<ul style="list-style-type: none"> <li>• G2 = 1.24 (1.00-1.54)</li> <li>• G3 = 1.00 (referent)</li> <li><math>p &lt; 0.001</math></li> </ul>	
Glass et al 1999 [179]	To examine any association between social activity, productive activity and PA and mortality in older people.	<ul style="list-style-type: none"> <li>• n = 2,761 (1,169 men; 1,143 women)</li> </ul>	13 year follow-up	Incidence of all-cause mortality by fitness activity quartile	More active elderly people were less likely to die than those who were less active.
USA		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>	PA assessment: Interview, Amount of activity	13 yr mortality by amount of activity	
Prospective cohort		<ul style="list-style-type: none"> <li>• Age: <math>\geq 65</math> yr</li> </ul>	G1 = Low	<ul style="list-style-type: none"> <li>• G1 = 74.0</li> </ul>	
D & B score = 12		<ul style="list-style-type: none"> <li>• Characteristics: Healthy elders</li> </ul>	G2 = Low-medium	<ul style="list-style-type: none"> <li>• G2 = 69.8</li> </ul>	
			G3 = Medium-high	<ul style="list-style-type: none"> <li>• G3 = 62.4</li> </ul>	
			G4 = High	<ul style="list-style-type: none"> <li>• G4 = 55.2</li> </ul>	
Gulati et al 2003 [180]	To determine whether exercise capacity is a predictor for all-cause mortality in asymptomatic women.	<ul style="list-style-type: none"> <li>• n = 5,721</li> </ul>	Baseline (1992) and 8 year follow-up (2000)	<ul style="list-style-type: none"> <li>• 180 deaths</li> </ul>	This study confirmed that exercise capacity is an independent predictor of death in asymptomatic women, greater than what has been previously established among men.
USA		<ul style="list-style-type: none"> <li>• Sex: Women</li> </ul>	PF Assessment: Treadmill stress test Exercise capacity (METs)	For every 1 MET increase there was a reduced death risk of 17% ( $p < 0.001$ )	
			G1 = $<5$		
			G2 = 5-8		
			G3 = $>8$		
Prospective cohort		<ul style="list-style-type: none"> <li>• Age: Mean <math>52 \pm 11</math> yr</li> </ul>		Age-adjusted RR	
D & B score = 11		<ul style="list-style-type: none"> <li>• Characteristics: Asymptomatic women</li> </ul>		<ul style="list-style-type: none"> <li>• G1 = 2.0 (1.3-3.2)</li> </ul>	
		<ul style="list-style-type: none"> <li>• St James Women Take Heart Project</li> </ul>		<ul style="list-style-type: none"> <li>• G2 = 1.6 (1.1-2.4)</li> </ul>	
				<ul style="list-style-type: none"> <li>• G3 = 1.00 (referent)</li> </ul>	
				Adjusted for Framingham Risk Score	
				<ul style="list-style-type: none"> <li>• G1 = 3.1 (2.1-4.8)</li> </ul>	
				<ul style="list-style-type: none"> <li>• G2 = 1.9 (1.3-2.9)</li> </ul>	
				<ul style="list-style-type: none"> <li>• G3 = 1.00 (referent)</li> </ul>	
Haapanen et al 1996 [181]	To examine the association between LTPA and all-cause mortality.	<ul style="list-style-type: none"> <li>• n = 1,072</li> </ul>	Baseline and a 10 yr 10 month follow-up	<ul style="list-style-type: none"> <li>• 168 deaths</li> </ul>	Low PA is a risk factor for all-cause mortality.
Finland		<ul style="list-style-type: none"> <li>• Sex: Men</li> </ul>	PA assessment: Self-reported LTPA, divided into 4 groups by EE (kJ/wk)	RR (95% CI) according to EE group	
			G1 = 0-3349		
			G2 = 3350-6279		
			G3 = 6280-8791		
			G4 = $>8791$		

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Prospective cohort		• Age: 35-63 yr	Mortality–National Death Index search	• G1 = 2.74 (1.46-5.14)	
D & B score = 14		• Characteristics: Healthy, sedentary	Cox proportional HR	• G2 = 1.10 (0.55-2.21) • G3 = 1.74 (0.87-3.50) • G4 = 1.00 (referent)	
Hakim et al 1998 [182]	To examine the association between walking and mortality in retired men.	• n = 707	Baseline and 12 yr follow-up	• 208 deaths	The findings in older physically capable men indicate that regular walking is associated with a lower overall mortality rate.
USA		• Sex: Men		RR (95% CI) according to distance walked Adjusted for age	
Prospective cohort		• Age: 61-81 yr	PA assessment: Questionnaire	• G1 vs. G3 = 1.9 (1.3-2.9)	
D & B score = 12		• Characteristics: Retired non-smoking men who were physically capable of participating in low intensity activities on a daily basis	Distance walked (miles/day)		
			G1 = 0.0-0.9 G2 = 1.0-2.0 G3 = 2.1-8.0	• G1 vs. G3 = 1.6 (1.2-2.2) • G2 vs. G3 = 1.2 (0.8-1.7) Trend $p = 0.002$	
		• Honolulu Heart Program		Adjusted for risk factors • G1 vs. G3 = 1.8 (1.2-2.7) • G1 vs. G2 = 1.5 (1.1-2.1) • G2 vs. G3 = 1.1 (0.8-1.7) Trend $p = 0.01$	
Hillsdon et al 2004 [183]	To examine whether VPA is associated with all-cause mortality.	• n = 10,522 (4,929 men; 5,593 women)	>10 year follow-up	• 825 deaths	Questionnaire respondents who reported engaging in VPA less than twice a week experienced a 37% reduced risk of all-cause mortality compared with respondents who reported a lower frequency of VPA.
UK		• Sex: Men and women	PA assessment: Questionnaire for frequency of VPA	Age and sex adjusted RR (95% CI)	
Prospective Cohort		• Age: 35-64 yr	G1 = Never, <1 time/month	• G1 = 1.00 (referent)	
		• Characteristics: Healthy, sedentary	G2 = <2 times/wk	• G2 = 0.57 (0.42-0.79)	
		• OXCHECK study	G3 = >2 times/wk	• G3 = 0.72 (0.54-0.95)	
D & B score = 11				Fully adjusted RR (95% CI) • G1 = 1.00 (referent) • G2 = 0.63 (0.45-0.89) • G3 = 0.81 (0.60-1.09)	
			Mortality – Recorded from the Office of National Statistics		

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

			Cox proportional HR		
Hu et al 2005 [184]	To examine the association of PA and BMI and their combined effect with the risk of total, CVD and cancer mortality.	<ul style="list-style-type: none"> <li>• n = 47,212 (22,528 men; 24,684 women)</li> <li>• Sex: Men and women</li> <li>• Age: 25-64 yr</li> <li>• Characteristics: Participants from eastern Finland</li> </ul>	17.7 year follow-up	<ul style="list-style-type: none"> <li>• 7,394 deaths</li> </ul>	Regular PA is an important indicator for decreased risk of all-cause mortality. PA has a strong independent effect on mortality.
Finland			PA assessment: Questionnaire for PA level, divided into 3 groups	Adjusted HR (95% CI)	
Prospective cohort				Men	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.74 (0.68-0.81)</li> <li>• G3 = 0.63 (0.58-0.70)</li> </ul>	
D & B score = 12			G1 = Low G2 = Moderate G3 = High	Trend $p = <0.001$	
				Women	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.64 (0.58-0.70)</li> <li>• G3 = 0.58 (0.52-0.64)</li> </ul>	
				Trend $p = <0.001$	
Hu et al 2004 [185]	To examine the association of BMI and PA with death.	<ul style="list-style-type: none"> <li>• n = 116,564</li> <li>• Sex: Women</li> <li>• Age: 30-55 yr</li> <li>• Characteristics: Females free of known CVD and cancer</li> </ul>	Baseline (1976) and 24 year follow-up	<ul style="list-style-type: none"> <li>• 10,282 deaths</li> </ul>	Reduced PA is a strong and independent predictor of death.
USA			PA assessment: Questionnaire for PA level, divided into 3 groups (hr/week)	Multivariate RR (95% CI) by PA (hr/wk)	
Prospective cohort			G1 = $\geq 3.5$ G2 = 1.0-3.4 G3 = $<1.0$	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.18 (1.10-1.26)</li> <li>• G3 = 1.52 (1.41-1.63)</li> </ul>	
D & B score = 11			BMI ( $\text{kg/m}^2$ )	Multivariate RR (95% CI) by PA adjusted for BMI	
			G1 = $<25$ G2 = 25-29 G3 = 30	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.14 (1.06-1.22)</li> <li>• G3 = 1.44 (1.34-1.55)</li> </ul>	
			Cox proportional HR		
Kampert et al 1996 [186]	To examine PF and PA in relation to all-cause and cancer mortality.	<ul style="list-style-type: none"> <li>• n = 32,421 (25,341 men; 7,080 women)</li> </ul>	Baseline (1970) and ~8 year follow-up (1989)	<ul style="list-style-type: none"> <li>• 690 deaths</li> </ul>	The data support the hypothesis that an active and fit way of life delays death.

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

USA Prospective cohort  D & B score = 13		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: 20-88 yr (mean ~43)</li> <li>• Characteristics: Predominantly white and from the middle and upper socioeconomic strata</li> </ul>	PA assessment: Questionnaire, divided into quintiles of activity (min/wk)	Adjusted RR (95% CI) by quintiles of activity	
			Men	<ul style="list-style-type: none"> <li>• Sedentary = 1.00 (referent)</li> <li>• C1-2 = 0.71 (0.58-0.97)</li> <li>• C3 = 0.83 (0.59-1.16)</li> <li>• C4 = 0.57 (0.30-1.08)</li> <li>• C5 = 0.92 (0.29-2.88)</li> </ul>	
			Male activity categories	Trend $p = 0.011$	
			Sedentary = 855 C1-2 = 1,072 C3 = 1,292 C4 = 1,453 C5 = 1,601	Women <ul style="list-style-type: none"> <li>• Sedentary = 1.00 (referent)</li> <li>• C1-2 = 0.68 (0.39-1.17)</li> <li>• C3 = 0.39 (0.09-1.65)</li> <li>• C4-5 = 1.14 (0.27-4.80)</li> </ul> Trend $p = 0.217$	
Kaplan et al 1996 [187]	To assess LTPA and its association with all cause mortality.	• n = 6,131 (3298 men; 2833 women)	28 year follow-up	• 1,226 deaths	The data provide further support for the importance of PA and indicate that the protective effect of PA is a robust one.
USA  Prospective cohort  D & B score = 13		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: 16-94 yr</li> <li>• Characteristics: Northern Californian adults</li> </ul>	PA assessment: Three questions about PA, with scores 0 (never), 2 (sometimes) or 4 (often).	Incidence of all-cause mortality and PA	
				Death rates/1000 person years	
				Men	
			Tertiles of PA score	<ul style="list-style-type: none"> <li>• T1 = 24.68</li> <li>• T2 = 11.37</li> <li>• T3 = 7.59</li> </ul>	
			T1 = 0-2	Women	
			T2 = 4-6		
			T3 = 8-12	<ul style="list-style-type: none"> <li>• T1 = 18.03</li> </ul>	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				<ul style="list-style-type: none"> <li>• T2 = 7.66</li> <li>• T3 = 3.88</li> </ul>	
Khaw et al 2006 [188]	To examine the relationship between PA patterns over 1 year and total mortality.	<ul style="list-style-type: none"> <li>• n = 22,191 (9,984 men; 12,207 women)</li> <li>• Sex: Men and women</li> <li>• Age: 45-79 yr</li> <li>• Characteristics: Community living participants</li> </ul>	8 year follow-up	<ul style="list-style-type: none"> <li>• 1,553 deaths</li> </ul>	Even very moderate levels of usual PA are associated with reductions in mortality.
UK			PA assessment: Questionnaire, divided into 4 groups of PA	Incidence of all-cause mortality and PA Adjusted RR (95% CI) All	
Prospective cohort			G1 = Inactive	• G1 = 1.00 (referent)	
			G2 = Moderately inactive	• G2 = 0.83 (0.73-0.95)	
				• G3 = 0.68 (0.58-0.80)	
			G3 = Moderately active	• G4 = 0.68 (0.57-0.81)	
			G4 = Active	Age <65	
				• G1 = 1.00 (referent)	
				• G2 = 1.01 (0.78-1.31)	
				• G3 = 0.81 (0.62-1.07)	
				• G4 = 0.82 (0.62-1.09)	
				Age >65	
				• G1 = 1.00 (referent)	
				• G2 = 0.77 (0.66-0.91)	
				• G3 = 0.65 (0.53-0.79)	
				• G4 = 0.64 (0.50-0.80)	
Kohl et al 1996 [189]	To determine the association of maximal exercise hemodynamic responses with risk of all-cause mortality.	<ul style="list-style-type: none"> <li>• n = 26,621 (20,387 men; 6,234 women)</li> <li>• Sex: Men and women</li> <li>• Age: Male mean 42.2 yr; female mean 41.9 Yr</li> <li>• Characteristics: Apparently healthy patients of a preventive medicine centre</li> </ul>	Average 8.1 year follow-up	<ul style="list-style-type: none"> <li>• 348 deaths in men and 66 in women</li> </ul>	The results suggest an exaggerated SBP or an attenuated heart rate response to maximal exercise may indicate an elevated risk for mortality.
USA				Adjusted RH (95% CI) by maximal exercise test HR Men	
Prospective cohort			PF assessment: Maximal exercise test HR (bpm), divided into 4 Groups:	• Q1 = 1.00 (referent)	
			G1 = <171	• Q2 = 0.61 (0.44-0.85)	
			G2 = 171-178	• Q3 = 0.69 (0.51-0.93)	
			G3 = 179-188	• Q4 = 0.60 (0.41-0.87)	
			G4 = >188	Trend $p < 0.05$	
D & B score = 12					



**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				Women	
				• Q1 = 1.00 (referent)	
				• Q2 = 1.23 (0.65-2.32)	
				• Q3 = 0.69 (0.30-1.63)	
				• Q4 = 0.71 (0.22-2.24)	
				Trend $p > 0.05$	
Kujala et al 1998 [190]	To investigate LTPA and mortality in a cohort of twins.	• n = 15,902 (7,925 men; 7,977 women)	Baseline 1975 and death outcome from 1977-1994	• 1,253 deaths	LTPA is associated with reduced mortality, even after genetic and other familial factors are taken into account.
Finland		• Sex: Men and women		HR (95% CI)	
		• Age: 25-64 yr			
		• Characteristics: Healthy, Finnish same sex twins	PA assessment: Questionnaire, quintiles of fitness in MET hours/day	Adjusted for age and sex	
Prospective cohort				• Sedentary = 1.00 (referent)	
				• OE = 0.71 (0.62-0.81)	
				• CE = 0.57 (0.45-0.74)	
		• The Finnish Twin Cohort		Trend $p = 0.001$	
D & B score = 13			Q1 = <58		
			Q2 = 59-1.29		
			Q3 = 1.30-2.49	Adjusted for age, sex, smoking	
			Q4 = 2.50-4.49		
			Q5 = >4.50	• Sedentary = 1.00 (referent)	
				• OE = 0.76 (0.67-0.87)	
				• CE = 0.68 (0.53-0.88)	
			Categorized into:		
			-Sedentary		
			-Occasional exerciser (OE)	Trend $p = 0.001$	
			-Conditioning exerciser (CE)	Adjusted for age, sex, smoking, occupational group, alcohol	
				• Sedentary = 1.00 (referent)	
				• OE = 0.80 (0.69-0.91)	
				• CE = 0.76 (0.59-0.98)	
				Trend $p = 0.002$	
				HR (95% CI) among 434 same sex twin pairs compared with sedentary category in 1975	
				• Sedentary = 1.00 (referent)	
				• OE = 0.66 (0.46-0.94)	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				<ul style="list-style-type: none"> <li>• CE = 0.44 (0.23-0.83)</li> </ul> <p>Trend <math>p = 0.005</math> Adjusted for smoking</p> <ul style="list-style-type: none"> <li>• Sedentary = 1.00 (referent)</li> <li>• OE = 0.70 (0.48-1.01)</li> <li>• CE = 0.56 (0.29-1.09)</li> </ul> <p>Trend <math>p = 0.04</math> Adjusted for smoking, occupational group, alcohol</p> <ul style="list-style-type: none"> <li>• Sedentary = 1.00 (referent)</li> <li>• OE = 0.73 (0.50-1.07)</li> <li>• CE = 0.56 (0.29-1.11)</li> </ul> <p>Trend <math>p = 0.06</math> OR (95% CI) in quintiles among 434 same sex twin pairs compared with sedentary category in 1975</p> <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.85</li> <li>• Q3 = 0.72</li> <li>• Q4 = 0.68</li> <li>• Q5 = 0.60</li> </ul>	
LaCroix et al 1996 [191]	To determine whether walking is associated with a reduced risk of CVD hospitalization and death in older adults.	<ul style="list-style-type: none"> <li>• n = 1,645 (615 men; 1030 women)</li> </ul>	4.2 year follow-up	RR (95% CI) by category of walking	Walking more than 4 hr/wk was associated with a reduced risk of mortality from all-causes.
		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>	PA assessment: Questionnaire for walking h/wk, divided into 3 groups		
USA		<ul style="list-style-type: none"> <li>• Age: <math>\geq 65</math> yr</li> </ul> <p>Characteristics: Participants from a group health co-operative</p>	G1 = <1 hr/week G2 = 1-4 hr/week G3 = >4 hr/week	Men	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>
Prospective cohort				<ul style="list-style-type: none"> <li>• G2 = 0.78 (0.43-1.45)</li> <li>• G3 = 0.89 (0.49-1.62)</li> </ul>	
D & B score = 12				Women	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.50 (0.28-0.90)</li> <li>• G3 = 0.48 (0.25-0.83)</li> </ul> <p>Age 65-74 yr</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.81 (0.40-1.61)</li> </ul>

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				<ul style="list-style-type: none"> <li>• G3 = 1.13 (0.60-2.15)</li> </ul> <p>Age ≥75 yr</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.63 (0.37-1.08)</li> <li>• G3 = 0.46 (0.25-0.84)</li> </ul> <p>High functioning</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.73 (0.38-1.41)</li> <li>• G3 = 0.89 (0.48-1.65)</li> </ul> <p>Limited functioning</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.60 (0.34-1.05)</li> <li>• G3 = 0.51 (0.28-0.92)</li> </ul>	
Lam et al 2004 [192]	To investigate the relationship LTPA and mortality in Hong Kong.	<ul style="list-style-type: none"> <li>• n = 24,079 cases (13,778 men; 10,301 women);</li> </ul>	10 years prior	Multivariate adjusted OR (95% CI) by LTPA	The data confirm and extend previous findings in Caucasian populations on the association between LTPA and longevity.
Hong Kong		<ul style="list-style-type: none"> <li>• n = 13,054 controls (3,918 men; 9,136 women)</li> </ul>	PA assessment: Questionnaire for LTPA, divided into 3 groups	Men	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.60 (0.54-0.67)</li> <li>• G3 = 0.66 (0.60-0.73)</li> </ul>	
Case-Control		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: ≥35 yr</li> <li>• Characteristics: All ethnic Chinese</li> </ul>	G1 = <1 times per month	Women	
D & B score = 12			G2 = 1-3 times per month	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.81 (0.74-0.88)</li> <li>• G3 = 0.71 (0.66-0.77)</li> </ul>	
			G3 = ≥4 times per month		
Lan et al 2006 [193]	To investigate the relationship between exercise and all-cause mortality.	<ul style="list-style-type: none"> <li>• n = 2,113 (1,081 men; 1,032 women)</li> <li>• Sex: Men and women</li> <li>• Age: ≥65 yr</li> <li>• Characteristics: Non-institutionalized elders</li> <li>• Taiwan National Health Interview Survey</li> </ul>	Baseline and 2 year follow-up	<ul style="list-style-type: none"> <li>• 197 deaths</li> </ul> <p>HR (95% CI) by LTPA frequency</p> <p>Adjusted for age and sex</p>	Older persons are recommended to expend at least 1000 kcal/wk through regular exercise for mortality reduction.
Taiwan			PA assessment: Questionnaire for LTPA (frequency/wk)		
Prospective cohort			G1 = Sedentary	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.49 (0.36-0.67)</li> <li>• G3 = 0.20 (0.09-0.46)</li> </ul>	Protection of exercise against death also increases with the number of activities.
D & B score = 13			G2 = 1 time/wk		
			G3 = ≥2 times/wk		

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

			Questionnaire for EE (kcal/wk), divided into 5 groups:	<p>Trend <math>p = &lt;0.001</math></p> <p>Multivariate adjusted</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.70 (0.50-0.98)</li> <li>• G3 = 0.35 (0.15-0.82)</li> </ul> <p>Trend <math>p = 0.014</math></p>	
			<p>G1 = Sedentary</p> <p>G2 = &lt;500</p> <p>G3 = 500-999</p> <p>G4 = 1000-1999</p> <p>G5 = <math>\geq 2000</math></p>	<p>HR (95% CI) by EE</p> <p>Adjusted for age and sex</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.64 (0.41-1.01)</li> <li>• G3 = 0.55 (0.35-0.85)</li> <li>• G4 = 0.30 (0.17-0.53)</li> <li>• G5 = 0.24 (0.12-0.48)</li> </ul> <p>Trend <math>p &lt; 0.001</math></p> <p>Multivariate adjusted</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.80 (0.49-1.30)</li> <li>• G3 = 0.74 (0.46-1.17)</li> <li>• G4 = 0.50 (0.27-0.90)</li> <li>• G5 = 0.43 (0.21-0.87)</li> </ul> <p>Trend <math>p = 0.043</math></p>	
Laukkanen et al 2001 [194]	To examine the relationship between maximal oxygen uptake and overall mortality.	<ul style="list-style-type: none"> <li>• n = 1,294</li> </ul>	Baseline and 10.7 year follow-up	<ul style="list-style-type: none"> <li>• 124 deaths</li> </ul>	PF has a strong, graded, inverse association with overall mortality.
Finland		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: 42.0-61.3 yr (mean 52.1)</li> <li>• Characteristics: Men free from CVD, COPD, and cancer at baseline</li> </ul>	PF assessment: Exercise tolerance test, 4 groups by maximal oxygen uptake (ml/kg/min)	<p>Adjusted RR (95% CI) by quartile</p>	
Prospective cohort				<p>Maximal oxygen uptake</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.47 (0.71-3.01)</li> <li>• G3 = 2.79 (1.44-5.39)</li> <li>• G4 = 3.85 (2.02-7.32)</li> </ul> <p>Linear trend <math>p = &lt;0.001</math></p>	
D & B score = 14			<p>G1 = &gt;37.1</p> <p>G2 = 32.3-37.1</p> <p>G3 = 27.6-32.2</p>		

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

			G4 = <27.6	Test duration	
			Test duration (min)	• G1 = 1.00 (referent)	
			G1 = >11.2	• G2 = 2.22 (1.08-4.55)	
			G2 = 9.6-11.2	• G3 = 2.23 (1.11-4.49)	
			G3 = 8.2-9.5	• G4 = 3.94 (2.01-7.74)	
			G4 = <8.2	Linear trend $p < 0.001$	
Lee and Paffenbarger 2000 [195]	To compare various levels of PA with mortality.	• n = 13,485	Baseline and 15 year follow-up	• 2,539 deaths	The study provides some support for recommendations that emphasize MPA. A benefit of VPA is also evident.
		• Sex: Men		RR (95% CI)	
		• Age: Mean 57.5 yr	PA assessment:	• G1 = 1.00 (referent)	
		• Characteristics: Men who matriculated as undergraduates in 1916-1950			
USA			Questionnaires for LTPA index (including walking, stair climbing, sports and recreational activity),	• G2 = 0.80 (0.72-0.88)	
Prospective cohort				• G3 = 0.74 (0.65-0.83)	
		• The Harvard Alumni Health Study		• G4 = 0.80 (0.69-0.93)	
				• G5 = 0.73 (0.64-0.84)	
D & B score = 12				Trend $p = < 0.001$	
			5 groups (kJ/wk)		
			G1 = <4200		
			G2 = 4200-8399		
			G3 = 8400-12599		
			G4 = 12600-16799		
			G5 = ≥ 16800		
Lee et al 1995 [196]	To examine the independent association of vigorous and non-vigorous PA with longevity.	• n = 17,321	Follow-up 22-26 years	• 3,728 deaths	There is a graded inverse relationship between PA and mortality. Vigorous, but not non-vigorous activities are associated with longevity.
		• Sex: Men			
		• Age: Mean 46 yr	PA assessment: Questionnaires for EE (kJ/wk), quintiles	RR (95% CI) by EE (kJ/wk)	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

USA		• Characteristics: Harvard University alumni, without self-reported physician diagnosed cardiovascular disease, cancer or chronic obstructive pulmonary disease		Q1= 1.00 (referent)	
Prospective cohort			Q1 = ≤ 630 Q2 = 630-1680 Q3 = 1680-3150 Q4 = 3150-6300 Q5 = >6300	• Q2 = 0.94 (0.86–1.04) • Q3 = 0.95 (0.86–1.05) • Q4 = 0.91 (0.83 - 1.01) • Q5 = 0.91 (0.82-1.00)	
D & B score = 12				RR (95% CI) by EE (Vigorous activity, kJ/wk) • Q1 = 1.00 (referent) • Q2 = 0.88 (0.82-0.96)  • Q3 = 0.92 (0.82-1.02) • Q4 = 0.87 (0.77-0.99) • Q5 = 0.87 (0.78-0.97)	
Lee et al 2004 [197]	To investigate the effect of various PA patterns on all-cause mortality.	• n = 8,421	Baseline 1988 and follow-up 1993	• 1,234 deaths	The results suggest that regular PA generating 1000 kcal/wk or more should be recommended for lowering mortality rates. Among those with no major risk factors, even 1-2 episodes per week generating 1000 kcal or more can postpone mortality.
USA		• Sex: Men • Age: Mean 66 yr		Age adjusted RR (95% CI) by PA pattern	
Prospective cohort		• Characteristics: Participants free of major chronic disease	PA assessment: Questionnaire for PA (kcal/wk), 4 groups	• G1 = 1.00 (referent) • G2 = 0.75 (0.63-0.90) • G3 = 0.82 (0.63-1.07) • G4 = 0.61 (0.53-0.69)	
D & B score = 11		• The Harvard Alumni Health Study	G1 = <500 (Sedentary)  G2 = 500-999 (Insufficiently active) G3 = ≥ 1000 (Weekend warrior) G4 = Regularly active	Multivariate adjusted  • G1 = 1.00 (referent) • G2 = 0.75 (0.62-0.91) • G3 = 0.85 (0.65-1.11)	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				• G4 = 0.64 (0.55-0.73)	
Leitzmann et al 2007 [198]	To examine PA guidelines in relation to mortality.	• n = 252,925 (142,828 male; 110,097 women)	Baseline and 6 month follow-up	• 7,900 deaths	Following PA guidelines is associated with lower risk of death. Mortality benefit may also be achieved by engaging in less than recommended activity levels.
USA		• Sex: Men and women	PA assessment: Questionnaire for MPA and VPA, 5 groups each MPA (h/wk)	Multivariate adjusted RR (95% CI) according to activity	
Prospective cohort		• Age: 50-71 yr		MPA	
		• Characteristics: Participants free of CVD, cancer or emphysema		• G1 = 1.00 (referent)	
		• The National Institute of Health-American Association of Retired Persons		• G2 = 0.85 (0.79-0.93)	
D & B score = 13				• G3 = 0.79 (0.74-0.85)	
			G1 = sedentary	• G4 = 0.76 (0.71-0.82)	
			G2 = <1	• G5 = 0.68 (0.63-0.74)	
			G3 = 1-3	Trend $p = <0.001$	
			G4 = 4-7	VPA	
			G5 = >7		
			VPA (frequency/wk)	• G1 = 1.00 (referent)	
			G1 = inactive	• G2 = 0.77(0.71-0.83)	
			G2 = <1	• G3 = 0.77 (0.72-0.82)	
			G3 = 1-2	• G4 = 0.68 (0.63-0.73)	
			G4 = 3-4	• G5 = 0.71 (0.66-0.77)	
			G5 = $\geq 5$	Trend $p = <0.001$	
			Cox proportional HR		
Leon et al 1997 [199]	To examine the long-term association of LTPA and risk of death from coronary heart disease and all-causes.	• n = 12,138	16 year follow-up	• 1,904 deaths	The data suggest that a relatively small amount of daily moderate intensity LTPA can reduce premature mortality in middle-aged and older men at high risk for CHD.
		• Sex: Men			
		• Age: 35-57 yr	PA assessment: Minnesota LTPA questionnaire, categorized by frequency/month and average duration, deciles (min/d)	Multivariate adjusted RR (95% CI) by deciles of LTPA	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

USA		<ul style="list-style-type: none"> <li>• Characteristics: Men who at entry to the study were free of clinical evidence of CHD or other serious medical problems but were at the upper 10%-15% of a CHD probability score distribution derived from the FHS data</li> </ul>			
Prospective cohort				<ul style="list-style-type: none"> <li>• D1 = 1.00 (referent)</li> <li>• D2-4 = 0.85 (0.73-0.99)</li> <li>• D5-7 = 0.87 (0.75-1.02)</li> <li>• D8-10 = 0.83 (0.71-0.97)</li> </ul>	
D & B score = 12			D1 = 4.9 D2-4 = 22.7 D5-7 = 53.9 D8-10 = 140.4		
		<ul style="list-style-type: none"> <li>• Multiple Risk Factor Intervention Trial</li> </ul>	Cox proportional HR		
Lissner et al 1996 [200]	To examine the relationship of OPA and LTPA on all-cause mortality in women.	<ul style="list-style-type: none"> <li>• n = 1,405</li> <li>• Sex: Women</li> <li>• Age: 38-60 yr</li> <li>• Characteristics: Free from major disease at baseline</li> </ul>	Baseline and 20 year follow-up	<ul style="list-style-type: none"> <li>• 277 deaths</li> </ul>	Decreases in PA as well as low initial levels are strong risk factors for mortality.
Sweden			PA assessment: Questionnaire for OPA and LTPA, 3 groups	RR (95% CI) by LTPA	
Prospective cohort		<ul style="list-style-type: none"> <li>• The Gothenburg Prospective Study of Women</li> </ul>		20 year follow-up LTPA during age 20-38 years <ul style="list-style-type: none"> <li>• Low = 1.00 (referent)</li> <li>• Med = 0.66 (0.34-1.26)</li> <li>• High = 0.46 (0.21-1.01)</li> </ul>	
D & B score = 10			G1 = Low G2 = Medium G3 = High	LTPA during age 39-60 years <ul style="list-style-type: none"> <li>• Low = 1.00 (referent)</li> <li>• Med = 0.56 (0.35-0.90)</li> <li>• High = 0.44 (0.22-0.91)</li> </ul> LTPA during the past 12 months <ul style="list-style-type: none"> <li>• Low = 1.00 (referent)</li> <li>• Med = 0.56 (0.39-0.82)</li> <li>• High = 0.45 (0.24-0.86)</li> </ul> 20 year follow-up OPA during age 20-38 years	



**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				<ul style="list-style-type: none"> <li>• Low = 1.00 (referent)</li> <li>• Med = 0.59 (0.18-1.87)</li> <li>• High = 0.50 (0.16-1.58)</li> </ul> <p>OPA during age 39-60 years</p> <ul style="list-style-type: none"> <li>• Low = 1.00 (referent)</li> <li>• Med = 0.66 (0.21-2.08)</li> <li>• High = 0.47 (0.14-1.52)</li> </ul> <p>OPA during the past 12 months</p> <ul style="list-style-type: none"> <li>• Low = 1.00 (referent)</li> <li>• Med = 0.28 (0.17-0.46)</li> <li>• High = 0.24 (0.14-0.43)</li> </ul>	
Manini et al 2006 [201]	To determine whether energy expenditure is associated with all-cause mortality in older adults.	<ul style="list-style-type: none"> <li>• n = 302 (150 men; 152 women)</li> <li>• Sex: Men and women</li> </ul>	Mean follow-up of 6.15 years	<ul style="list-style-type: none"> <li>• 55 deaths</li> </ul> <p>HR (95% CI) by tertiles of PA EE</p>	Free-living activity EE was strongly associated with lower risk of mortality.
USA		<ul style="list-style-type: none"> <li>• Age: 70-82 yr</li> </ul>	PA assessment: Questionnaire, divided into tertiles of PA EE (kcal/d)		
Prospective cohort		<ul style="list-style-type: none"> <li>• Characteristics: High-functioning community dwelling elders</li> </ul>		Adjusted for age, sex, race and study site	
D & B score = 13			<p>T1 = &lt;521</p> <p>T2 = 521-770</p> <p>T3 = &gt;770</p>	<ul style="list-style-type: none"> <li>• T1 = 1.00 (referent)</li> <li>• T2 = 0.63 (0.29-1.18)</li> <li>• T3 = 0.37 (0.15-0.76)</li> </ul> <p>Trend <math>p = 0.009</math></p> <p>Adjusted for age, sex, race, study site, weight, height, percent body fat and sleep duration</p> <ul style="list-style-type: none"> <li>• T1 = 1.00 (referent)</li> <li>• T2 = 0.57 (0.30-1.09)</li> <li>• T3 = 0.31 (0.14-0.69)</li> </ul> <p>Trend <math>p = 0.004</math></p> <p>Adjusted for age, sex, race, study site, self rated health, education, smoking, CVD, lung disease, diabetes, hip or knee osteoarthritis, osteoporosis, cancer and depression</p> <ul style="list-style-type: none"> <li>• T1 = 1.00 (referent)</li> <li>• T2 = 0.65 (0.33-1.28)</li> </ul>	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				• T3 = 0.33 (0.15-0.74) Trend $p = 0.007$	
Matthews et al 2007 [202]	To determine the effects of exercise and non-exercise PA on mortality.	• n = 67,143  • Sex: Women • Age: 40-70 yr • Characteristics: Women without heart disease, stroke or cancer	Baseline and an average of 5.7 year follow-up	• 1,091 deaths  RR (95% CI)	Overall PA levels are an important determinant of longevity.
China			PA assessment: Interview to report (MET h/d), 4 groups Overall activity	Multivariate adjustment	
Prospective cohort				Overall activity (MET hr/d) • G1 = 1.00 (referent) • G2 = 0.81 (0.69-0.96)	
D & B score = 12		• The Shanghai Women's Health Study	G1 = ≤ 9.9 G2 = 10.0-13.6 G3 = 13.7-18.0 G4 = ≥ 18.1  Adult exercise G1 = 0 G2 = 0.1-3.4 G3 = 3.5-7.0 G4 = ≥ 7.1 Cox proportional hazard models	• G3 = 0.67 (0.57-0.80) • G4 = 0.61 (0.51-0.73) Trend $p = 0.000$  Adult exercise (MET hr/d) • G1 = 1.00 (referent) • G2 = 0.84 (0.74-0.96) • G3 = 0.77 (0.59-0.99) • G4 = 0.64 (0.36-1.14) Trend $p = 0.008$	
Menotti and Seccareccia 1985 [203]	To investigate the relationship between OPA and all-cause mortality.	• n = 99,029  • Sex: Men • Age: 40-59 yr • Characteristics: Men employed on the Italian railway system	Baseline and 5 year follow-up	• 2,661 deaths  Age adjusted death rates per 1000 over 5 years classified by PA only	The results suggest that PA may play a role in the prediction of fatal events.
Italy				• Sedentary = 26.20 • Moderate = 27.05 • Heavy = 27.35	
Prospective cohort					

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

D & B score = 12				Age adjusted death rates per 1,000 over 5 years classified by PA and job responsibility	
			G1 = sedentary – low	• G1 = 30.00	
			G2 = sedentary – med	• G2 = 25.20	
			G3 = sedentary – high	• G3 = 25.80	
			G4 = moderate – low	• G4 = 26.30	
			G5 = moderate – med	• G5 = 28.50	
			G6 = moderate – high	• G6 = 25.80	
			G7 = heavy – low	• G7 = 26.90	
			G8 = heavy – med	• G8 = 30.80	
Mensink et al 1996 [204]	To compare various indices for PA and their association with cardiovascular risk factors as well as total and CVD mortality.	• n = 15,436 (7,689 men; 7797 women)	5-8 year follow-up	Incidence of all-cause mortality and PA	An inverse relation of PA and total mortality.
Germany		• Sex: Men and women	PA assessment: Questionnaire Total activity, 3 groups	Adjusted RR (95% CI)	
		• Age: 25-69 yr			
Prospective cohort		• Characteristics: Participants from communities in Western Germany		Total activity, men	
			G1 = Low	• G1 = 1.00 (referent)	
			G2 = Moderate	• G2 = 0.56 (0.30-1.04)	
			G3 = High	• G3 = 0.78 (0.42-1.44)	
				Total activity, women	
			LTPA, 3 groups	• G1 = 1.00 (referent)	
			G1 = Low	• G2 = 1.24 (0.60-2.58)	
			G2 = Moderate	• G3 = 1.29 (0.58-2.85)	
			G3 = High		
			Conditioning activity, 3 groups	LTPA, men	
				• G1 = 1.00 (referent)	
			G1 = No activity	• G2 = 0.61 (0.35-1.05)	
			G2 = Moderate	• G3 = 0.79 (0.48-1.31)	
			G3 = High	LTPA, women	
				• G1 = 1.00 (referent)	
			Sports activity, 4 groups	• G2 = 0.94 (0.51-1.75)	
				• G3 = 0.81 (0.44-1.49)	
			G1 = no sports		
			G2 = <1 hour	Conditioning activity, men	
			G3 = 1-2 hours	• G1 = 1.00 (referent)	
			G4 = >2 hours	• G2 = 0.76 (0.44-1.34)	
				• G3 = 0.67 (0.36-1.25)	
D & B score = 12					

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				Conditioning activity, women • G1 = 1.00 (referent) • G2 = 0.38 (0.13-1.06) • G3 = 0.80 (0.42-1.54) Sports Activity, men • G1 = 1.00 (referent) • G2 = 0.49 (0.26-0.95) • G3 = 0.57 (0.30-1.09) • G4 = 0.36 (0.16-0.79) Sports activity, women • G1 = 1.00 (referent) • G2 = 0.38 (0.12-1.23) • G3 = 0.52 (0.23-1.17) • G4 = 0.28 (0.07-1.17)	
Morgan and Clarke 1997 [205]	To assess the value of broadly based customary PA scores in predicting 10-year mortality in elderly people.	• n = 1,042 (407 men; 635 women)  • Sex: Men and women  • Age: ≥65 yr • Characteristics: British elders  • Nottingham Longitudinal Study of Activity and Aging	10 year follow-up  PA assessment: Questionnaire for PA, 3 groups  G1 = Low G2 = Intermediate  G3 = High	Incidence of all-cause mortality and PA  HR (95% CI) Men • G1 = 1.59 (1.12-2.25) • G2 = 1.35 (0.96-1.89)  • G3 = 1.00 (referent) Women • G1 = 2.07 (1.53-2.79) • G2 = 1.53 (1.12-2.09) • G3 = 1.00 (referent)	A wide range of customary or habitual PA, can provide indices showing both cross sectional and predictive validity for 10 year mortality.
UK  Prospective cohort  D & B score = 12					
Myers et al 2002 [206]	To compare PF and PA levels with all-cause mortality.	• n = 6,213  • Sex: Men • Age: Mean 59 ± 11 yr	Baseline and mean 6.2 ± 3.7 year follow-up	• 1,256 deaths  Age adjusted RR (95% CI) by quintile	Exercise capacity is a more powerful predictor of mortality among men than other established risk factors for CVD.

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

USA		<ul style="list-style-type: none"> <li>• Characteristics: Participants with a normal exercise test result (n = 2,534) and participants with an abnormal exercise test or CVD or both (n = 3,679)</li> </ul>	PF assessment: Treadmill test for VO <sub>2</sub> peak, divided into quintiles (METs)	<ul style="list-style-type: none"> <li>• Q1 = 4.5 (3.0-6.8)</li> </ul>	
Prospective cohort				<ul style="list-style-type: none"> <li>• Q2 = 2.4 (1.5-3.8)</li> <li>• Q3 = 1.7 (1.1-2.8)</li> <li>• Q4 = 1.3 (0.7-2.2)</li> </ul>	
D & B score = 12			Q1 = Lowest level 1.0-5.9 Q2 Q3 Q4 Q5 = Highest level ≥13.0	<ul style="list-style-type: none"> <li>• Q5 = 1.00 (referent)</li> </ul>	
Ostbye et al 2002 [207]	To analyze the effect of smoking and other modifiable risk factors on ill health, defined in a multidimensional fashion.	<ul style="list-style-type: none"> <li>• n = 12,956</li> </ul>	6 year follow-up	<ul style="list-style-type: none"> <li>• 782 deaths</li> </ul>	Quitting smoking and increasing exercise levels are the lifestyle interventions most likely to improve overall health.
USA		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: 50-60 yr</li> </ul>	PA assessment: Questionnaire for PA, 4 groups	Incidence of all-cause mortality and PA	
Prospective cohort		<ul style="list-style-type: none"> <li>• Characteristics: Participants from the Health and Retirement Study (HRS) only</li> </ul>	G1 = Sedentary	Death rates (95% CI) per 1000 population/yr	
D & B score = 13			G2 = Light G3 = Moderate G4 = Heavy	<ul style="list-style-type: none"> <li>• G1 = 20.6 (17.8-24.0)</li> <li>• G2 = 9.1 (8.1-9.5)</li> <li>• G3 = 8.3 (7.5-9.2)</li> <li>• G4 = 4.4 (3.5-5.6)</li> </ul>	
Paffenbarger et al 1994 [208]	To study the adoption or maintenance of PA and other optional lifestyle patterns for their influence on mortality rates of Harvard College alumni.	<ul style="list-style-type: none"> <li>• n = 14,786</li> </ul>	Follow-up between	<ul style="list-style-type: none"> <li>• 2,343 deaths</li> </ul>	Adopting a physically active lifeway delays mortality and extends longevity.
		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: 45-84 yr (in 1977)</li> </ul>	1977 and 1988	RR (95% CI) of mortality according to PA	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

USA			PA assessment: Questionnaire for blocks walked daily, stairs climbed daily and type, frequency and duration of weekly sports and recreational activities		
		Characteristics: Harvard College alumni			
Prospective cohort				Physical activity index (kcal/wk)	
				• G1 = 1.00 (referent)	
				• G2 = 1.13 (1.01-1.26)	
				• G3 = 0.72 (0.64-0.82)	
				• G4 = 0.77 (0.69-0.85)	
D & B score = 14			Physical activity index (kcal/wk) Sports and recreational activities were scored according to intensity and duration	Walking (km/wk)	
				• G1 = 1.00 (referent)	
				• G2 = 1.21 (1.08-1.35)	
				• G3 = 0.94 (0.83-1.07)	
				• G4 = 0.89 (0.78-1.01)	
				Moderately vigorous sports play (METs)	
			Light < 4.5 METs Moderate ≥ 4.5 METs		
				• G1 = 1.00 (referent)	
				• G2 = 1.11 (0.93-1.33)	
				• G3 = 0.73 (0.65-0.81)	
				• G4 = 0.72 (0.64-0.80)	
				Adjusted for potential confounding influences	
Richardson et al 2004 [209]	To investigate the impact of a sedentary lifestyle on all-cause mortality.	• n = 9,611 (4,642 men; 4,969 women)  • Sex: Men and women • Age: 51-61 yr	Baseline (1992) and 8 year follow-up	• 810 deaths	A sedentary lifestyle is associated with a higher risk of death in pre-retirement aged adults.
USA				OR (95% CI)	
Prospective cohort		• Characteristics: Participants born between 1931-1941 and who not institutionalized in 1992	PA assessment: Questionnaire for PA, 3 groups:	• G1 = 1.00 (referent)	
				• G2 = 0.64 (0.52-0.81)	
			G1 = Sedentary G2 = occasional or light G3 = Regular MVPA	• G3 = 0.62 (0.44-0.85) p = 0.01	
D & B score = 13					

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

		• Health and Retirement Study			
Rockhill et al 2001 [210]	To determine the association between recreational PA and mortality in women.	• n = 80,348	Baseline (1980) and follow-up between 1982-1996	• 4,871 deaths	People who are more physically active are at reduced mortality risk relative to those who are less active.
USA		• Sex: Women • Age: 30-55 yr		Multivariate adjusted RR (95% CI) by (hr/wk)	
Prospective cohort		• Characteristics: Free from CVD or cancer at baseline • Nurses Health Study	PA assessment: Questionnaire in 1980 and up-dated every 2- 4 years, 5 groups of PA (hr/wk)	• G1 = 1.00 (referent)	
D & B score = 11				• G2 = 0.82 (0.76-0.89) • G3 = 0.75 (0.69-0.81) • G4 = 0.74 (0.68-0.81) • G5 = 0.71 (0.61-0.82) p<0.001	
			G1 = <1 G2 = 1-1.9 G3 = 2-3.9 G4 = 4-6.9 G5 = ≥7		
Rosengren and Wilhelmsen 1997 [211]	To investigate the effect of OPA and LTPA on risk of death.	• n = 7,142	Baseline (1970-1973) and 20 year follow-up	• 2,182 deaths	The study demonstrates the protective effect of LTPA on mortality.
Sweden		• Sex: Men • Age: 47-55 yr • Characteristics: Without symptomatic CHD	PA assessment: Postal questionnaires, 3 groups:	Unadjusted RR (95% CI) • G1 = 1.00 (referent) • G2 = 0.74 (0.68-0.82) • G3 = 0.73 (0.68-0.79)	
Prospective cohort			G1 = Sedentary G2 = Moderately active G3 = Regular exercise	Multivariate adjustment • G1 = 1.00 (referent) • G2 = 0.84 (0.77-0.93) • G3 = 0.83 (0.77-0.90)	
D & B score = 13					

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Schnohr et al 2003 [212]	To assess the associations of regular LTPA and changes in LTPA with risk of death.	<ul style="list-style-type: none"> <li>• n = 7,023 (4,471 men; 5,676 women)</li> <li>• Sex: Men and women</li> <li>• Age: 20-79 yr</li> <li>• Characteristics: Participants from the Copenhagen City Heart Registered Population</li> </ul>	18 year follow-up	<ul style="list-style-type: none"> <li>• 2,725 deaths</li> <li>Incidence of all-cause mortality and PA and changes in PA</li> </ul>	Maintaining or adopting a moderate or high degree of PA was associated with lower risk of death.
Denmark					
Prospective cohort			G1 = Low-low G2 = Low-moderate G3 = Low-high G4 = Moderate- low G5 = Moderate-Moderate G6 = Moderate-high G7 = High-low G8 = High-moderate G9 = High-high	Adjusted RR (95% CI) Men <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.64 (0.49-0.83)</li> <li>• G3 = 0.64 (0.47-0.87)</li> <li>• G4 = 0.73 (0.56-0.96)</li> <li>• G5 = 0.71 (0.57-0.88)</li> <li>• G6 = 0.64 (0.51-0.81)</li> <li>• G7 = 1.11 (0.76-1.62)</li> <li>• G8 = 0.66 (0.51-0.85)</li> <li>• G9 = 0.61 (0.48-0.76)</li> </ul> Women <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.75 (0.57-0.97)</li> <li>• G3 = 0.72 (0.50-1.05)</li> <li>• G4 = 0.70 (0.54-0.91)</li> <li>• G5 = 0.64 (0.52-0.79)</li> <li>• G6 = 0.58 (0.45-0.73)</li> <li>• G7 = 0.72 (0.48-1.07)</li> <li>• G8 = 0.61 (0.47-0.80)</li> <li>• G9 = 0.66 (0.51-0.85)</li> </ul>	
D & B score = 12					
Schnohr et al 2004 [213]	To examine whether the relationship between established risk factors and mortality differs with socioeconomic status as measured by level of education.	<ul style="list-style-type: none"> <li>• n = 30,635 (16,236 men; 14,399 women)</li> </ul>	16 year follow-up	<ul style="list-style-type: none"> <li>• 10,952 deaths</li> </ul>	The study shows the strong predictive effect of PA on mortality is independent of education level.



**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

	• Sex: Men and women	Socioeconomic status assessment: level of education	Incidence of all-cause mortality and PA stratified by years of education
Denmark	• Age: 20-93 yr • Characteristics: Participants from the Copenhagen City Heart Registered Population		
Prospective cohort D & B score = 12		PA assessment: Questionnaire	Deaths <8 years of education
			Men
		4 groups of PA	G1 = 916
		G1 = none or very little	G2 = 1693
		G2 = 2-4 h/wk of LPA	G3 = 1012
		G3 = >4 h/wk of LPA or 2-4 h/wk of high level activity	G4 = 67
		G4 = Competition level or >4 h/wk of hard level activity	Women
			• G1 = 872
			• G2 = 1298
			• G3 = 346
			• G4 = 10
			8-11 years of education
			Men
			• G1 = 432
			• G2 = 1040
			• G3 = 616
			• G4 = 33
			Women
			• G1 = 363
			• G2 = 852
			• G3 = 268
			• G4 = 10
			>11 years of education
			Men
			• G1 = 104
			• G2 = 302
			• G3 = 182
			• G4 = 11
			Women

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

				<ul style="list-style-type: none"> <li>• G1 = 48</li> <li>• G2 = 129</li> <li>• G3 = 61</li> <li>• G4 = 3</li> </ul>	
Schnohr et al 2006 [214]	To investigate the association between LTPA and mortality.	<ul style="list-style-type: none"> <li>• n = 4,894 (2,136 men; 2,758 women)</li> </ul>	Baseline (1976) and start of follow-up in 1981-1983 (to 2000)	<ul style="list-style-type: none"> <li>• 1,787 deaths</li> </ul>	Long-term moderate or high PA was associated with significantly lower mortality in men and women.
Denmark		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: 20-79 yr</li> <li>• Characteristics: Healthy males and women</li> </ul>		RR (95% CI)	
Prospective cohort			PA assessment: Survey for LTPA, 3 groups:	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.64 (0.56-0.73)</li> <li>• G3 = 0.56 (0.48-0.65)</li> </ul>	
D & B score = 13		<ul style="list-style-type: none"> <li>• The Copenhagen City Heart Study</li> </ul>	G1 = Low G2 = Mod G3 = High	Trend $p < 0.001$  Multivariate adjustment <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.78 (0.68-0.89)</li> <li>• G3 = 0.75 (0.64-0.87)</li> </ul> Trend $p = 0.001$	
Schooling et al 2006 [215]	To examine how a Comprehensive assessment of baseline health status affects the relationship between obesity or PA and mortality.	<ul style="list-style-type: none"> <li>• n = 54,088 (17,849 men; 36,239 women)</li> </ul>	4.1 year follow-up	<ul style="list-style-type: none"> <li>• 3,819 deaths</li> </ul>	PA, which normally has a negative relationship with adiposity, had the largest impact on survival for the health states, with the strongest inverse relationship between BMI and mortality.
Hong Kong		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>	PA assessment: Interview for PA min/d, 3 groups	Incidence of all-cause mortality and PA	
Prospective cohort		<ul style="list-style-type: none"> <li>• Age: <math>\geq 65</math> yr</li> <li>• Characteristics: Chinese elders</li> </ul>	G1 = None G2 = $\leq 30$ min/d G3 = $\geq 30$ min/d	Adjusted HR (95% CI) <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.83 (0.76-0.91)</li> <li>• G3 = 0.73 (0.67-0.80)</li> </ul> Trend $p < 0.001$	
D & B score = 13					

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Sundquist et al 2004 [216]	To study the association between varying levels of PA and all-cause mortality in the elderly.	<ul style="list-style-type: none"> <li>• n = 3,206 (1,414 men; 1,792 women)</li> <li>• Sex: Men and women</li> <li>• Age: ≥65 yr</li> <li>• Characteristics: Non-institutionalized elders</li> </ul>	Baseline (1988-1989) and follow-up in 2000	<ul style="list-style-type: none"> <li>• 1,806 deaths</li> </ul>	Even occasional PA decreases the risk of mortality among elderly people.
Sweden			PA assessment: Questionnaire for PA, 5 groups	Age-adjusted HR (95% CI)	
Prospective cohort				Men	
				• G1 = 1.00 (referent)	
				• G2 = 0.74 (0.62-0.87)	
				• G3 = 0.57 (0.44-0.73)	
				• G4 = 0.51 (0.41-0.64)	
		The Swedish Annual Level-of-Living Survey (Statistics Sweden)	G1 = none G2 = occasionally	• G5 = 0.60 (0.44-0.82)	
D & B score = 12			G3 = once per week G4 = twice per week G5 = vigorously at least twice per week	Women	
				• G1 = 1.00 (referent)	
				• G2 = 0.70 (0.59-0.82)	
				• G3 = 0.59 (0.46-0.77)	
				• G4 = 0.47 (0.35-0.62)	
				• G5 = 0.54 (0.31-0.94)	
			Cox proportional HR	Men and women	
				Multivariate adjustment	
				• G1 = 1.00 (referent)	
				• G2 = 0.72 (0.64-0.81)	
				• G3 = 0.60 (0.50-0.71)	
				• G4 = 0.50 (0.42-0.59)	
				• G5 = 0.60 (0.46-0.79)	
Talbot et al 2007 [217]	To investigate how changes in LTPA affect all-cause mortality.	<ul style="list-style-type: none"> <li>• n = 2,092 (1,316 men; 776 women)</li> <li>• Sex: Men and women</li> <li>• Age: 19-&lt;90 yr</li> </ul>	Baseline in 1958 for males and in 1978 for females and an average follow-up of 21.2 ± 9.4 years for men and 10.2 ± 5.6 years for women	<ul style="list-style-type: none"> <li>• 628 deaths (538 male; 90 female)</li> </ul>	Greater declines in total and high-intensity LTPA are independent predictors of all-cause mortality.
USA				RR (95% CI) for standard deviation of rate of change in LTPA	

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Prospective cohort	<ul style="list-style-type: none"> <li>• Characteristics: Community residents, generally with above average income, high education and with good or excellent self related health</li> </ul>	<p>(If RR is &lt;1 then a SD increase is associated with decrease mortality. If RR is &gt;1, then a SD increase is associated with increase in mortality)</p>	
D & B score = 13	The Baltimore Longitudinal Study of Aging	PA assessment: Questionnaire for LTPA (METs min/24 h), 3 groups G1 = low G2 = medium G3 = high Rate of change (ROC)	Multivariate adjustment Men <70 years • G1 = 0.96 (0.84-1.08) • G2 = 0.91 (0.79-1.04) • G3 = 0.42 (0.33-0.53) • ROC low = 0.90 (0.80-1.01) • ROC med = 1.01 (0.90-1.14) • ROC high = 0.78 (0.65-0.94) Men >70 years • G1 = 0.95 (0.82-1.10) • G2 = 0.89 (0.76-1.05) • G3 = 0.78 (0.62-0.97) • ROC low = 1.07 (0.93-1.24) • ROC med = 1.13 (1.00-1.27) • ROC high = 0.91 (0.75-1.12) Women <70 years • G1 = 0.75 (0.53-1.07) • G2 = 0.61 (0.36-1.03) • G3 = 0.80 (0.50-1.30) • ROC low = 1.02 (0.74-1.40) • ROC med = 1.38 (0.86-2.28) • ROC high = 0.90 (0.63-1.27) Women >70 years • G1 = 0.85 (0.63-1.15) • G2 = 0.78 (0.39-1.59) • G3 = 0.62 (0.32-1.22) • ROC low = 1.10 (0.85-1.42) • ROC med = 0.96 (0.46-2.03) • ROC high = 0.70 (0.40-1.22)

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Trolle-Lagerros et al 2005 [218]	To quantify the effect of PA on overall mortality in younger women and to assess the effect of past versus current activity.	<ul style="list-style-type: none"> <li>• n = 99,099</li> </ul>	11.4 year follow-up	<ul style="list-style-type: none"> <li>• 1,313 deaths</li> </ul>	Current PA substantially reduces mortality among women. The association is observed even with low levels of PA and is accentuated with increased PA.
		<ul style="list-style-type: none"> <li>• Sex: Women</li> <li>• Age: 30-49 yr</li> </ul>	PA assessment: Questionnaire using a 5 point scale, 5 groups	Incidence of all-cause mortality and PA past and current	
Sweden and Norway		<ul style="list-style-type: none"> <li>• Characteristics: Participants from Norway and one region of Sweden</li> </ul>			
Retrospective cohort			G1 = Sedentary	Adjusted HR (95% CI)	
			G2 = Low	PA at enrolment	
			G3 = Moderate	• G1 = 1.00 (referent)	
			G4 = High	• G2 = 0.78 (0.61-1.00)	
			G5 = Vigorous	• G3 = 0.62 (0.49-0.78)	
				• G4 = 0.58 (0.44-0.75)	
				• G5 = 0.46 (0.33-0.65)	
				Trend $p < 0.0001$	
				PA at age 30 yr	
				• G1 = 1.00 (referent)	
				• G2 = 0.79 (0.55-1.15)	
				• G3 = 0.90 (0.64-1.28)	
				• G4 = 0.98 (0.68-1.42)	
				• G5 = 0.96 (0.65-1.44)	
				Trend $p = 0.22$	
				PA at age 14 yr	
				• G1 = 1.00 (referent)	
				• G2 = 0.95 (0.66-1.38)	
				• G3 = 0.96 (0.69-1.34)	
				• G4 = 0.88 (0.62-1.25)	
				• G5 = 1.06 (0.75-1.51)	
				Trend $p = 0.62$	
Villeneuve et al 1998 [219]	To examine the relationship between PF, PA and all-cause mortality.	<ul style="list-style-type: none"> <li>• n = 14,442 (6,246 men; 8,196 women)</li> </ul>	Baseline (1981) and 7 year follow-up	RR (95% CI) by EE, multivariate adjustment	There was a reduction in mortality risk associated with even modest participation in activities of low intensity.
		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: 20-69 yr</li> </ul>			
Canada			PA assessment: Questionnaire for EE (kcal/kg/day), 5 groups	LTPA, men	


**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Prospective cohort	D & B score = 11	Canadian Fitness Survey	• Characteristics: Asymptomatic for CVD	• G1 = 1.00 (referent)	
			G1 = 0-<0.5 G2 = 0.5-<1.5 G3 = 1.5-<3.0 G4 = ≥ 3.0 G5 = ≥ 0.5 PF levels: Recommended Minimum  Undesirable Refusal  Multivariate Poisson regression analysis	• G2 = 0.81 (0.59-1.11) • G3 = 0.79 (0.54-1.13) • G4 = 0.86 (0.61-1.22) • G5 = 0.82 (0.65-1.04)*  Non vigorous LTPA, men • G1 = 1.00 (referent) • G2 = 0.81 (0.56-1.17) • G3 = 0.70 (0.44-1.13) • G4 = 0.82 (0.53-1.27) • G5 = 0.78 (0.59-1.04)*  LTPA, women • G1 = 1.00 (referent)  • G2 = 0.94 (0.69-1.30) • G3 = 0.92 (0.64-1.34) • G4 = 0.71 (0.45-1.11) • G5 = 0.88 (0.68-1.04)* Non vigorous LTPA, women • G1 = 1.00 (referent) • G2 = 0.97 (0.69-1.36) • G3 = 0.87 (0.57-1.33) • G4 = 0.72 (0.43-1.21) • G5 = 0.89 (0.67-1.17)* RR (95% CI) by fitness levels, adjusted for age, sex and smoking Recommended = 1.00 (referent) • Minimum = 1.02 (0.69-1.51) • Undesirable = 1.52 (0.72-3.18) • Refusal = 1.04 (0.45-2.39)	
Weller and Corey 1998 [220]	To study the relationship between PA and mortality in women.	• n = 6,620  • Sex: Women • Age: ≥;30 yr	Baseline and 7 year follow-up	• 449 deaths  OR (95% CI)	PA is inversely associated with risk of death in women.

**Table 11: Studies examining the relationship between physical activity and all-cause mortality. (Continued)**

Canada		<ul style="list-style-type: none"> <li>• Characteristics: Without known heart disease</li> <li>• Canadian Fitness Survey</li> </ul>	PA assessment: Questionnaires for: EE (kcal/kg/d), quartiles	EE (kcal/kg/d)	
Prospective cohort				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.91 (0.66-1.25)</li> <li>• Q3 = 0.94 (0.72-1.23)</li> <li>• Q4 = 0.89 (0.67-1.17)</li> </ul>	
D & B score = 11			Q1 = lowest Q2 = Q3 = Q4 = highest LTPA, 3 groups G1 = Sedentary G2 = Mod G3 = High	LTPA levels <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.63 (0.46-0.86)</li> <li>• G3 = 0.76 (0.59-0.98)</li> </ul>	
			Walking, 3 groups G1 = < half the time G2 = half the time G3 = > half the time	Walking <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.64 (0.49-0.82)</li> <li>• G3 = 0.64 (0.47-0.86)</li> </ul>	
Yu et al 2003 [221]	To examine the relationship between LTPA and all-cause mortality.	• n = 1,975	Baseline and 10 year follow-up	• 252 deaths	The study found a strong inverse association between heavy LTPA and all-cause mortality.
UK		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: 49-64 yr</li> <li>• Characteristics: Without a history of CHD at baseline</li> </ul>	PA assessment: Questionnaire (Minnesota LTPA index, kcal/d), 3 group	Age adjusted HR (95% CI)	
Prospective cohort				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.73 (0.54-0.99)</li> <li>• G3 = 0.74 (0.55-1.04)</li> </ul> Trend $p = 0.046$	
D & B score = 11			G1 = Light to no activity G2 = Moderate activity G3 = Heavy activity	Multivariate adjusted <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.79 (0.58-1.08)</li> <li>• G3 = 0.76 (0.56-1.04)</li> </ul> Trend $p = 0.083$	

D & B score, Downs and Black quality score; PF, physical fitness; YR, years; RR, risk ratio; 95% CI, 95% confidence interval; PA, physical activity; VO2 peak, peak oxygen consumption; HR, hazard ratio; min/d, minutes per day; kcal/wk, kilocalories per week; LTPA, leisure-time physical activity; MET, metabolic equivalent; VO2 max, maximal oxygen consumption; OPA, occupational physical activity; CVD, cardiovascular disease; hr/wk, hours per week; MPA, moderate physical activity; kcal/kg/wk, kilocalories per kilogram per week; kJ/wk, kilojoules per week; EE, energy expenditure; G, groups; EE, energy expenditure; BMI, body mass index; C, class; kg/m<sup>2</sup>, kilogram by meters squared; HR, heart rate; BPM, beats per minute; MVPA, moderate to vigorous physical activity; OR, odds ratio; Q, quartile or quintile; RCT, randomized clinical trial; T, tertiles; TPA, total physical activity; VPA, vigorous physical activity; mL/kg/min, milliliters per kilogram per minute.



**Figure 3** Mean relative risk reduction in all-cause mortality across physical activity/fitness categories.


definition of cardiovascular disease ( $n = 78$ ), and other ( $n = 19$ ). Therefore, a total of 49 articles were included in the systematic review of the literature regarding the relationship between physical activity and the incidence of cardiovascular disease.

The majority of the studies included in our systematic review were prospective cohort investigations (Table 12). These studies involved a total of 726,474 participants; averaging 12,313 participants per study (range 680-88,393). There were a total of 34,815 reported cases of cardiovascular disease (ranging per study from 42-2,596). The total length of study follow-up for the prospective cohort studies averaged 14.1 yr (ranging from 2-29 yr). The articles were published over a 32 yr period ranging from 1975 to 2007. These studies involved large samples of men and women from regions throughout the world.

Similar to the all-cause mortality data, the risk for cardiovascular disease demonstrates a graded inverse dose-response relationship to physical activity and fitness. The relative reduction in the incidence of cardiovascular disease averages 33% (median risk reduction of 36%),

with greater risk reductions in studies that employed objective measures of aerobic fitness. It is not uncommon for studies to demonstrate a 50% or higher risk reduction when an objective measure of physical fitness was taken (Table 12). The importance of physical activity may actually be underestimated owing to multivariate control for many confounding factors (as discussed previously) and the fact that effects of within-person variation in physical activity are often not considered [55]. The relative risk reduction appears to be similar for men and women, and also appear to extend to non-Caucasian populations [56]. Some evidence also exists indicating that small amounts of physical activity are associated with lower cardiovascular-disease related mortality [57,58]. Similar to all-cause mortality, physical activity confers health benefits independent of other known risk factors [42,59]. *Collectively, the level of evidence would be considered to be Level 2A based on the presence of overwhelming evidence from observational trials.* The quality of the investigations was generally high with a mean (and median) Downs and Black score of 12 (range 9-14).





**Figure 4** Results of the Literature Search for Cardiovascular Disease.

## Implications

Research in the field began with the landmark work of Morris and colleagues, which demonstrated that men in physically demanding occupations (bus conductors and postmen) had a significantly lower risk of heart disease than individuals who worked in less demanding jobs (bus drivers and office workers) [45]. Since then

considerable research has examined the relationship between physical activity and the risk for cardiovascular disease. In fact, several systematic reviews of the literature have been developed regarding the role of habitual physical activity in the primary and secondary prevention of cardiovascular disease [33,60-62]. The research to date has been consistent and compelling, habitual physical activity reduces markedly the risk for cardiovascular disease.

Based on the available literature, there is compelling evidence that the recommendation of 30 min of moderate intensity exercise on most days of the week (equivalent to 4.2 MJ/wk or 1000 kcal/wk) reaches a threshold associated with significant reductions in cardiovascular-related mortality [32,63]. Brisk walking has also been shown to be preferable to a slower pace [64]. However, weekly exercise volumes of less than 4.2 MJ (1000 kcal) may be cardio-protective [14,59,65-67]. For instance, Lee et al. (2001) found that as little as 1 hr/wk of walking was associated with a 50% lower cardiovascular disease mortality in one sample of women. Wisløff et al. [58] reported that a single weekly bout of self-reported high intensity exercise was associated with a lower risk of cardiovascular death relative to those reporting no activity in both men (RR = 0.61, 95% CI = 0.49-0.75), and women (RR = 0.49, 95% CI = 0.27-0.89). Moreover, no additional benefit was seen with higher durations or frequency of exercise sessions [58]. The authors stated that this evidence challenges "current recommendations that require at least 1000 kcal of caloric expenditure per week to achieve exercise-induced protection against premature cardiovascular death." However, this research is in fact supportive of the Canadian guidelines which recognize the potential health benefits of low volumes of physical activity as reflected by the statement "Every little bit counts, but more is even better - everyone can do it!" It however should be noted that the statement "more is even better" is supported by a strong evidence base.

### Recommendation #2

*For a reduced risk for cardiovascular disease-related events and mortality, it is recommended that individuals participate in 30 min or more of moderate to vigorous exercise on most days of the week. Greater health benefits appear to occur with high volume and/or intensities of activity. Health benefits may also occur with as little as one hr of brisk walking per week. [Level 2, Grade A]*

### The Primary Prevention of Stroke

Stroke affects a significant proportion of Canadian society with approximately 50,000 new cases each year [68]. The relationship between physical activity and the risk for stroke is compelling, supporting at least a 25-

**Table 12 Studies examining the relationship between physical activity and cardiovascular disease.**

Publication Country Study Design Quality Score	Objective	Population	Methods	Outcome	Comments and Conclusions
Paffenbarger and Hale 1975 [47] USA  Prospective cohort  D & B score = 12	To evaluate the role of PA in reducing coronary mortality among longshoremen	<ul style="list-style-type: none"> <li>• n = 6,351</li> <li>• Sex: Men</li> <li>• Age: 35-74 yr</li> <li>• Characteristics: Longshoreman</li> </ul>	22 years of follow up, or until reached the age of 75 yr   PA assessment: Energy and oxygen cost requirements of longshoring jobs	RR (95% CI) Sudden death  <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 3.5</li> <li>• G3 = 2.8</li> </ul> Delayed death  <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.4</li> </ul> Activity level G1 = Heavy (5.2-7.5 kcal/min) G2 = Moderate (2.4-5.0 kcal/min) G3 = Light (1.5-2.0 kcal/min)  Outcome measure: Death from CHD	VPA is associated with reduced risk of coronary mortality, particularly sudden cardiac death.
Manson et al 2002 [56]  USA  Prospective cohort	To compare the roles of walking and vigorous exercise in the prevention of CV events in a large, ethnically diverse cohort of postmenopausal women.	<ul style="list-style-type: none"> <li>• n = 73,743</li> <li>• Sex: Women</li> <li>• Age: 50-79 yr</li> <li>• Characteristics: Healthy, Post Menopausal</li> <li>• Women's Health Initiative Observational Study</li> </ul>	Enrolment from 1994-98 Clinic visit for baseline screening,   PA assessment: Questionnaire for: Total exercise (MET- hr/wk) G1 = 0-2.4  G2 = 2.5-7.2	<ul style="list-style-type: none"> <li>• Number of New Cases: 345</li> <li>• Total Number of CVD events: 1551</li> </ul> Age adjusted RR (95% CI) Total exercise (MET-hr/wk)  <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.73 (0.53-0.99)</li> </ul>	Both walking and VPA are associated with substantial reductions in the incidence of CHD events.

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

D & B score = 12			G3 = 7.3-13.4	• G3 = 0.69 (0.51-0.95)	
			G4 = 13.5-23.3	• G4 = 0.68 (0.50-0.93)	
			G5 = ≥ 23.4	• G5 = 0.47 (0.33-0.67)	
				p = <0.001	
				Walking (MET-hr/wk)	
			Walking (MET-hr/wk)	• G1 = 1.00 (referent)	
			G1 = None	• G2 = 0.71 (0.53-0.96)	
			G2 = 0.1-2.5	• G3 = 0.60 (0.44-0.83)	
			G3 = 2.6-5.0	• G4 = 0.54 (0.39-0.76)	
			G4 = 5.1-10.0	• G5 = 0.61 (0.44-0.84)	
			G5 > 10	p = 0.004	
			Time for VPA (min)	Vigorous exercise	
			G1 = None	• G1 = 1.00 (referent)	
			G2 = 1-60	• G2 = 1.12 (0.79-1.60)	
			G3 = 61-100	• G3 = 0.56 (0.32-0.98)	
			G4 = 101-150	• G4 = 0.73 (0.43-1.25)	
			G5 = >150	• G5 = 0.58 (0.34-0.99)	
				p = 0.008	
			Outcome Measure: Incidence of CVD and CHD		
Wisloff et al 2006 [58]	To study the association between the amount and intensity of exercise and CVD mortality.	• n = 56,072 (27,143 men; 28,929 women)	Length of follow-up: 16 ± 4 yr	• Number of Cases: 1,603 male, 993 female	Men and women who exercise to a moderate degree and spend less than the recommended energy (< 1000 kcal/wk) are at lower risk of dying from heart disease than those who never exercise.
Norway		• Sex: Men and women	PA assessment: Questionnaire for LTPA, 4 groups	Multivariate RR (95% CI)	
Prospective cohort		• Age: ≥ 20 yr		Men	
		• Characteristics: Free form CVD	Men	• Q1 = 1.00 (referent)	
			Q1 = None	• Q2 = 0.66 (0.50-0.87)	
			Q2 = 1/wk >30 min high	• Q3 = 0.83 (0.65-1.06)	
			Q3 = 2-3/wk > 30 min high	• Q4 = 0.77 (0.59-1.01)	
			Q4 = ≥ 4/wk > 30 min high	Women	
				• Q1 = 1.00 (referent)	
D & B score = 12		• HUNT study			

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

			Women	• Q2 = 0.63 (0.31-1.29)	
			Q1 = None	• Q3 = 0.66 (0.32-1.34)	
			Q2 = 1/wk ≤ 30 min low	• Q4 = 0.86 (0.45-1.62)	
			Q3 = 1/wk ≤ 30 min high		
			Q4 = 2-3/wk ≤ 30 min low		
			Outcome Measure: Ischaemic heart disease mortality		
			Cox proportional HR		
Lee et al 2001 [59]	To examine the relationship between PA (specifically walking) and CHD among women, including those at high risk for CHD.	• n = 39,372	Recruitment of Participants: Sept 1992-May 1995	• Number of Cases: 244	Even light to moderate activity is associated with lower CHD rates in women.
USA and Puerto Rico		• Sex: Women			
		• Age: ≥ 45 yr		Multivariate RR (95% CI) Time spent walking	
		• Characteristics: Healthy	PA assessment: Questionnaires Divided into 4 or 5 groups:	• G1 = 1.00 (referent)	
		• Women's Health Study		• G2 = 0.86 (0.57-1.29)	As little as 1 hour of walking per week predicted lower risk.
Prospective cohort				• G3 = 0.49 (0.28-0.86)	
				• G4 = 0.48 (0.29-0.78)	
				p = <0.001	
D & B score = 12			Time spent walking		
			G1 = No regular walking	Walking pace	
			G2 = 1-59 min/wk	• G1 = 1.00 (referent)	
			G3 = 1.0-1.5 h/wk	• G2 = 0.56 (0.32-0.97)	
			G4 = ≥ 2.0 h/wk	• G3 = 0.71 (0.47-1.05)	
			Walking pace (km/h)	• G4 = 0.52 (0.30-0.90)	
			G1 = No regular walking	p = 0.02	
			G2 = 3.2		
			G3 = 3.2-4.7		
			G4 = ≥ 4.8	EE (kcal/wk)	
				• Q1 = 1.00 (referent)	
			EE (kcal/wk)	• Q2 = 0.79 (0.56-1.12)	
			G1 = 200	• Q3 = 0.55 (0.37-0.82)	
			G2 = 200-599	• Q4 = 0.75 (0.50-1.12)	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

			<p>G3 = 600-1499 and G4 = 1500 or more</p> <p>Energy expenditure for VPA (kcal/wk)</p> <p>G1 = No vigorous, &lt;200 kcal/wk</p> <p>G2 = No vigorous, ≥ 200 kcal/wk</p> <p>G3 = Vigorous, 1-199 kcal/wk</p> <p>G4 = Vigorous, 200-499 kcal/wk</p> <p>G5 = Vigorous, ≥ 500 kcal/wk</p>	<p><math>p = 0.03</math></p> <p>Energy expended VPA (kcal/wk)</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.65 (0.46-0.91)</li> <li>• G3 = 1.18 (0.79-1.78)</li> <li>• G4 = 0.96 (0.60-1.55)</li> <li>• G5 = 0.63 (0.38-1.04)</li> </ul>	
Paffenbarger et al 1993 [67]	To analyze changes in the lifestyle of Harvard Alumni and the associations of these changes to mortality.	<ul style="list-style-type: none"> <li>• n = 10,269</li> <li>• Sex: Men</li> <li>• Age: 45-84 yr</li> <li>• Characteristics: Health, Harvard College Alumni</li> </ul>	<p>Baseline measure in 1962 or 1967 with a follow up in 1977</p> <p>PA assessment: Mailed questionnaires included questions on type, duration, intensity, frequency of PA.</p> <p>Outcome Measure: CHD deaths between 1977 and 1985</p> <p>Cox proportional hazards model</p> <p>Poisson regression methods</p> <p>The Mantel extension of the Mantel-Haenszel test</p>	<p>Alumni who increased their PA index to 2000 kcal or more per week had a 17% lower risk of death from CHD than those who were sedentary (<math>p = 0.507</math>)</p> <p>Men who took up moderate took up moderately vigorous activity had a 41% lower risk than those who continued not to engage in such activity (<math>p = 0.044</math>)</p>	<p>Moderately vigorous sports activity was associated with lower rates of death from CHD among middle aged and older men</p>
USA					
Prospective cohort					
D & B score = 13					
Haapanen et al 1997 [77]	To examine the association between duration and intensity of LTPA and the risk of CHD.	<ul style="list-style-type: none"> <li>• n = 2,840 (1,500 men; 1,340 women)</li> </ul>	<p>Length of Follow-up: 10 yrs</p>	<ul style="list-style-type: none"> <li>• Incident Rates (per 1000 person-years) for CHD = 108 for men and 75 for women.</li> </ul>	<p>Total EE had an inverse and independent association with risk of CHD in middle aged Finnish men but not among women.</p>

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Finland		• Sex: Men and women	PA assessment: Questionnaire for LTPA EE (kcal/wk)	Multivariate RR (95% CI) LTPA and CHD mortality	
Prospective cohort		• Age: 35-63 yr		Men	
		• Characteristics: Healthy	Men	• G1 = 1.98	
			G1 = 0-1100	• G2 = 1.33	
			G2 = 1101-1900	• G3 = 1.00 (referent)	
			G3 = >1900		
			Women	• G1 = 1.25	
			G1 = 0-900	• G2 = 0.73	
			G2 = 901-1500	• G3 = 1.00 (referent)	
			G3 = >1500		
			Outcome Measure: CHD mortality		
			Cox proportional HR		
Barengo et al 2004 [164]	To investigate whether moderate or high LTPA are associated with a reduced CVD and all-cause mortality, independent of CVD risk factors and other forms of PA in men and women.	• n = 31,677 (15,853 men; 16,824 women)	20 year follow-up	• Number of Cases (Men): 1,661	Moderate and high levels of LTPA and OPA are associated with reduced CVD mortality.
			PA assessment: Questionnaire for LTPA and OPA, 3 groups	• Number of Cases (Women): 778	
Finland		• Sex: Men and women		HR (95% CI) LTPA, men	
Prospective cohort		• Age: 30-59	G1 = Low activity	• G1 = 1.00 (referent)	
		• Characteristics: Participant from eastern and south-western Finland	G2 = Moderate activity	• G2 = 0.91 (0.82-1.00)	
			G3 = High activity	• G3 = 0.83 (0.69-0.99)	
				LTPA, women (referent)	
				• G1 = 1.00	
				• G2 = 0.83 (0.71-0.96)	
				• G3 = 0.89 (0.68-1.18)	
				OPA, men	
				• G1 = 1.00 (referent)	
				• G2 = 0.75 (0.64-0.87)	
				• G3 = 0.77 (0.69-0.87)	
				OPA, women	
D & B score = 13					
D & B score = 14					

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.73 (0.60-0.88)</li> <li>• G3 = 0.77 (0.65-0.91)</li> </ul>	
Bijnen et al 1998 [166]	To describe the association between the PA pattern of elderly men and CHD mortality.	<ul style="list-style-type: none"> <li>• n = 802</li> <li>• Sex: Men</li> <li>• Age: 64-84 yr</li> </ul>	Length of Follow-up: 10	<ul style="list-style-type: none"> <li>• Number of Cases: 90</li> </ul>	PA did not show a protective effect on death from CHD.
Netherlands		<ul style="list-style-type: none"> <li>• Characteristics: Free from Serious Illness</li> </ul>	PA assessment: Questionnaire, divided into 3 groups	RR (95% CI)	
Prospective cohort			G1 = Lowest	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.63 (0.38-1.05)</li> <li>• G3 = 0.85 (0.51-1.44)</li> </ul>	
D & B score = 13		<ul style="list-style-type: none"> <li>• Ethnicity: Dutch</li> <li>• Zutphen Elderly Study</li> </ul>	G2 = Middle G3 = Highest Outcome Measure: CHD Mortality Cox Proportional HR		
Davey-Smith et al 2000 [174]	To examine the association between two measures of physical activity (LTPA and usual walking pace) with cause specific mortality (CHD).	<ul style="list-style-type: none"> <li>• n = 6,702</li> <li>• Sex: Men</li> <li>• Age: 40-64 yr</li> <li>• Whitehall Study</li> </ul>	Length of Follow-up: 25 yrs	<ul style="list-style-type: none"> <li>• Number of Cases: 955</li> </ul>	Inverse associations of both LTPA and walking pace with mortality from CHD were seen.
England			PA assessment: Questionnaire during examination for walking pace and LTPA	RR (95% CI) by walking pace	
Prospective cohort				<ul style="list-style-type: none"> <li>• G1 = 1.45 (0.9-2.2)</li> <li>• G2 = 1.30 (1.1-1.6)</li> <li>• G3 = 1.00 (referent)</li> </ul>	
D & B score = 11			Walking pace	$p < 0.01$ Multivariate RR (95% CI) by LTPA level	
			G1 = Slower	• G1 = 1.24 (1.0-1.5)	
			G2 = Same	• G2 = 0.94 (0.8-1.2)	
			G3 = Faster	• G3 = 1.00	
			LTPA	$p < 0.05$	
			G1 = Inactive		

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

			G2 = Moderate G3 = Active Outcome Measure: CHD Mortality Cox Proportional HR		
Eaton et al 1995 [175]	To determine whether self reported PA predicts a decreased risk of CHD.	• n = 8,463 (LTPA), 8,418 (OPA)	Length of Follow-up: 21 yrs	• Number of Cases: 709  Age adjusted RR (95% CI) by LTPA level	Baseline levels of self reported LTPA predicted a decreased rate of CHD.
USA		• Sex: Men • Age: 40 yr • Characteristics: Healthy, free of CHD	PA assessment: Interview	• G1 = 1.00 (referent) • G2 = 0.79 (0.63-0.99) • G3 = 0.73 (0.59-0.89)	
Prospective cohort			LTPA	• G4 = 0.71 (0.52-0.98)	
D & B score = 11		Ethnicity: Israeli	G1 = Sedentary G2 = Light G3 = Light Daily G4 = Heavy  OPA G1 = Sitting G3 = Walking G4 = Physical Labour Outcome Measure: CHD Death Cox Proportional HR	Age adjusted RR (95% CI) by OPA level • G1 = 1.00 (referent) • G2 = 0.99 (0.75-1.18) • G3 = 0.94 (0.78-1.12) • G4 = 0.87 (0.67-1.10)	
Hillsdon et al 2004 [183]	To examine whether a short, easily administered measure of PA is associated with the risk of death from all causes and specific causes.	• n = 10,522 (4,929 men; 5,593 women)	Length of Follow-up: > 10 yrs	• Number of Cases: 155  Multivariate RR (95% CI) by PA level	Self reported VPA is associated with the risk of future mortality.
UK		• Sex: Men and women	PA assessment: Questionnaire, 3 groups:	• G1 = 1.00 (referent) • G2 = 0.46 (0.19-1.12) • G3 = 0.96 (0.53-1.75)	
Prospective cohort		• Age: 35-64 yr  • Characteristics: no history of chest pain	G1 = Never / <1 time/month  G2 = <2 times/wk G3 = ≥ 2 times/wk		
D & B score = 11					



**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

			Outcome Measure: IHD mortality Cox proportional HR		
Leon et al 1997 [199]	To study the relationship of PA to CHD in a well defined population at above average risk for CHD over a 16 yr observation period.	<ul style="list-style-type: none"><li>• n = 12,138</li></ul>	Follow up for 16 years	Age Adjusted RR (95% CI)	A relatively small amount (10-36 min/d) of daily moderate intensity LTPA can significantly reduce premature mortality from CHD in middle aged men at high risk for CHD.
USA		<ul style="list-style-type: none"><li>• Sex: Men</li><li>• Age: 35-57 yr</li></ul>	PA assessment: Questionnaire at baseline (Minnesota LTPA questionnaire), divided/ grouped into deciles of LTPA (min/d)	<ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.71 (0.56-0.91)</li></ul>	
		<ul style="list-style-type: none"><li>• Characteristics: Free of CHD but in the upper 10-15% of a CHD probability risk score</li></ul>		<ul style="list-style-type: none"><li>• G3 = 0.75 (0.59-0.96)</li></ul>	
Prospective cohort				<ul style="list-style-type: none"><li>• G4 = 0.69 (0.54-0.96)</li></ul>	
D & B score = 11			G1 = D1: (0-9 min/d)	Multivariate adjusted RR (95% CI) <ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.75 (0.54-0.96)</li></ul>	
		Multiple risk factor intervention trial	G2 = D2-4: (10-36 min/d) G3 = D5-7: (37-75 min/d) G4 = D8-10: (76-359 min/d) Outcome Measure: CHD Mortality	<ul style="list-style-type: none"><li>• G3 = 0.81 (0.64-1.04)</li><li>• G4 = 0.75 (0.59-0.96)</li></ul>	
Rosengren et al 1997 [211]	To examine the long term effect of OPA and LTPA on the risk of death from CHD.	<ul style="list-style-type: none"><li>• n = 7,142</li></ul>	Length of Follow-up: 20 yrs	Number of Cases: 684	There appears to be a protective effect of LTPA on CHD-related death.
		<ul style="list-style-type: none"><li>• Sex: Men</li><li>• Age: 47-55 yr</li></ul>		Multivariate RR (95% CI) for LTPA	
Sweden		<ul style="list-style-type: none"><li>• Characteristics: Swedish men</li></ul>	PA assessment: Questionnaire for LTPA, 3 groups	<ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li></ul>	
				<ul style="list-style-type: none"><li>• G2 = 0.84 (0.71-1.00)</li><li>• G3 = 0.84 (0.73-0.96)</li></ul>	
Prospective cohort			G1 = Sedentary G2 = Moderately active		

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

D & B score = 13			G3 = Regular exercise		
			Outcome Measure: CHD death		
			Proportional HR		
Schnohr et al 2006 [214]	To describe the associations between different levels of LTPA and subsequent causes of death.	• n = 4,894 (2,136 men; 2,758 women)	Participants included in the study were only those whose PA levels did not change over 5 years	• Number of Cases: 292	There was an inverse and significant dose- response association between LTPA and CHD-related mortality.
Denmark			Adjusted RR (95% CI) Whole group		
			• G1 = 1.00 (referent)		
			• G2 = 0.71 (0.51-0.99)		
Prospective cohort			• G3 = 0.56 (0.38-0.82)		
D & B score = 12			PA assessment:		
			Questionnaire LTPA		
			G1 = <4 METS	Men	
			G2 = 4-6 METS	• G1 = referent	
			G3 = >6 METS	• G2 = survived 4.9 yrs longer	
				• G3 = survived 6.8 yrs longer	
			Cox proportional HR		
			Women		
			• G1 = referent		
			• G2 = survived 5.5 yrs longer		
			• G3 = survived 6.4 yrs longer		
Weller et al 1998 [220]	To examine the relationship between PA and mortality.	• n = 6,620	Length of Follow-up: 7 yrs	• Number of Cases: 109	LTPA is inversely associated with risk of fatal MI.
Canada			PA assessment:		
			Questionnaire, 4 groups for LTPA (kcal/kg/day) and non-LTPA (kcal/kg/day)		
			OR (95% CI) by LTPA		
			• Q1 = 1.00 (referent)		
			• Q2 = 0.61 (0.07-1.19)		
			• Q3 = 0.84 (0.52-1.37)		
Prospective cohort			• Q4 = 0.63 (0.36-1.09)		
D & B score = 9			LTPA (kcal/kg/day)	OR (95% CI) by non-LTPA	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

			Q1 = $\geq 0$ Q2 = $\geq 0.1$ Q3 = $\geq 0.5$ Q4 = $\geq 1.6$ Non-LTPA (kcal/kg/day) Q1 = $\geq 0$ Q2 = $\geq 2.8$ Q3 = $\geq 5.9$ Q4 = $\geq 9.9$ Outcome Measure: Fatal MI Logistic regression analysis	• Q1 = 1.00 (referent) • Q2 = 0.71 (0.44-1.16) • Q3 = 0.57 (0.33-0.97) • Q4 = 0.49 (0.26-0.92)	
Yu et al 2003 [221]	To examine the optimal intensity of LTPA to decrease the risk of CHD mortality in middle aged British men.	• n = 1,975  • Sex: Men  • Age: 49-64 yr • Characteristics: Healthy, no previous history of CHD	10 year follow-up  PA assessment: Questionnaire (Minnesota LTPA questionnaire), 3 groups	• Number of Cases: 82  Multivariate adjusted HR (95% CI)  • G1 = 1.00 (referent) • G2 = 0.74 (0.44-1.25)  • G3 = 0.55 (0.31-0.98)  $p = 0.039$	Strong significant inverse relationship between heavy LTPA and CHD mortality.        Relationship was not significant for low- moderate intensity LTPA and OPA.
UK  Prospective cohort  D & B score = 11		• Caerphilly collaborative heart study	Total activity level (kcal/day) G1 = 0.0 - 161.6  G2 = 161.8 - 395.3 G3 = 395.5 - 2747.2 Cox proportional HR		
Altieri et al 2004 [222]	To assess the possible protective role of PA on CHD.	• n = 985 (507 men; 478 women)	PA assessment: Questionnaire for OPA, divided into quartiles	Number of Cases: 507  OR (95% CI) for CHD and OPA	LTPA from 15-19 yrs as well as OPA from 30 - 39 yrs both have a significant inverse relationship with risk of non fatal acute MI.
Italy  Case Control		• Sex: Men and women  • Age: < 79 yr	Q1 = lowest Q2 Q3	• Q1 = 1.00 (referent) • Q2 = 0.63 (0.39-1.03) • Q3 = 0.56 (0.35-0.90)	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

D & B score = 11		<ul style="list-style-type: none"> <li>• Characteristics: Case: Patients admitted to Hospital with non-fatal Acute MI. Controls: Patients admitted to hospital for acute condition unrelated to known or potential risk factors for acute MI</li> </ul>	Q4 = highest	<ul style="list-style-type: none"> <li>• Q4 = 0.57 (0.34-0.95)</li> </ul>	
				$p = 0.045$	
			Outcome Measure: Non Fatal acute MI Unconditional logistic regression		
Batty et al 2003 [223]	To examine the relationship between physical activity and three mortality endpoints in healthy persons.	<ul style="list-style-type: none"> <li>• n = 6,474</li> </ul>	Length of Follow-up: 25 yr	<ul style="list-style-type: none"> <li>• Number of Cases: 837</li> </ul>	A suggestion that the symptomatic nature of ischemia appeared to modify the affects of
		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: 40-64 yr</li> </ul>		<ul style="list-style-type: none"> <li>• Number of Dropouts: 158</li> </ul>	
UK		<ul style="list-style-type: none"> <li>• Characteristics: British civil servants who underwent a resting ECG</li> </ul>	PA assessment: Questionnaire for LTPA, divided into 3 groups:	HR (95% CI) for CHD and LTPA	
Prospective cohort			G1 = Inactive	<ul style="list-style-type: none"> <li>• G1 = 1.14 (0.9-1.4)</li> <li>• G2 = 0.94 (0.8-1.1)</li> </ul>	PA on total and CHD mortality.
			G2 = Moderate		
			G3 = Active	<ul style="list-style-type: none"> <li>• G3 = 1.00 (referent)</li> </ul>	
D & B score = 13			Outcome Measure: CHD mortality Cox Proportional HR		
Chen and Millar [224]	To examine the potential protective effect of LTPA on the incidence of heart disease and depression.	<ul style="list-style-type: none"> <li>• n = 15,670</li> </ul>	Length of Follow-up: 2 yrs	<ul style="list-style-type: none"> <li>• 100 cases</li> </ul>	Regular and at least MPA can be beneficial to heart health.
		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: <math>\geq 20</math> yr</li> </ul>		Adjusted OR (95% CI)	
			PA assessment: EE from self administered questionnaire, 4 groups (kcal/kg/day)		
Canada		<ul style="list-style-type: none"> <li>• Characteristics: Healthy and free from heart disease</li> </ul>		<ul style="list-style-type: none"> <li>• G1 = 5.0 (1.84-13.59)</li> </ul>	
				<ul style="list-style-type: none"> <li>• G2 = 3.7 (1.26-10.67)</li> </ul>	
Prospective cohort				<ul style="list-style-type: none"> <li>• G3 = 1.00 (referent)</li> </ul>	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

D & B score = 11	National Population Health Survey		G1 = Sedentary G2 = Light (<1.5) G3 = Moderate (1.5-2.9)	• G4 = 1.3 (0.41-3.89)	
			G4 = Active (≥ 3) Outcome Measure: CHD incidence Multiple logistic regression		
Conroy et al 2005 [225]	To examine the relationship between 1) PA during young adulthood and middle age, and 2) PA during each time period and CHD during middle age and older women.	• n = 37,169  • Sex: Women • Age: ≥ 45 yr	Length of Follow-up: 9 yrs	• Number of Cases: 477	PA during middle age predicts lower risk of CHD
US		• Characteristics: Healthy women health professionals  • Women's Health Study	PA assessment: Questionnaire for EE (kcal/wk) and months/yr	Multivariate RR (95% CI) Baseline PA and incidence of CHD  • G1 = 1.00 (referent) • G2 = 0.62 (0.48-0.80) • G3 = 0.61 (0.48-0.79) • G4 = 0.61 (0.46-0.81)	
Cohort study			Baseline PA (kcal/wk)		
D & B score = 11			G1 = <200 G2 = 200-599 G3 = 600-1499 G4 = ≥ 1500	p = <0.001  Past PA and incidence of CHD • G1 = 1.00 (referent) • G2 = 0.76 (0.57-1.02) • G3 = 0.95 (0.72-1.24) • G4 = 1.04 (0.78-1.39) • G5 = 0.81 (0.58-1.14)	
			Past PA Months per year G1 = 0 G2 = 1-3 G3 = 4-6 G4 = 7-9 G5 = 10-12		
			Outcome Measure: Incidence of CHD Cox proportional hazard regression		

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Dorn et al 1999 [226]	To examine the long-term relationships between total PA and mortality from all causes and CHD in the general population.	• n = 1,461 (698 men; 763 women)	Length of Follow-up: 29 years	• Number of Cases: 109 men, 81 women	PA favorably influences mortality risks in non- obese men and younger women.
USA		• Sex: Men and women	PA assessment: Questionnaire	Multivariate RR (95% CI) for PAI in non- obese men	
Prospective cohort		• Age: 15-96 yr		• 0.40 (0.19-0.88) for 1 kcal/kg/h	
		• Characteristics: Healthy, free from CHD, diabetes, and Stroke.	Outcome Measure: CHD Mortality	Multivariate RR (95% CI) for PAI in obese men	
				• 1.86 (0.86-4.03) for 1 kcal/kg/h	
D & B score = 11			Cox Proportional Hazard Ratio	Multivariate RR (95% CI) for PAI in women < 60 yrs	
		• Ethnicity: White.		• 0.42 (0.11-1.52) for 1 kcal/kg/h	
				Multivariate RR (95% CI) for PAI in women > 60 yrs	
				• 1.78 (0.77-4.09) for 1 kcal/kg/h	
Folsom et al 1997 [227]	To examine the association of PA at baseline with CHD incidence.	• n = 13,999 (6,166 men; 7833 women)	Length of Follow-up: 4-7 yrs	• Number of Cases: 223 men, 97 women, Multivariate RR (95% CI) LTPA, men	No significant relationships.
USA		• Sex: Men and women	PA assessment: Questionnaire during home interview, divided into quartiles of LTPA and sports activity		
Prospective cohort		• Age: 45-64 yr		• Q1 = 1.00 (referent)	
		• Characteristics: no CHD at baseline		• Q2 = 1.08 (0.75-1.55)	
				• Q3 = 0.83 (0.51-1.36)	
				• Q4 = 0.89 (0.59-1.35)	
D & B score = 9			Q1 = Low		
		• Ethnicity: Black and non Black	Q2	LTPA, women	
			Q3	• Q1 = 1.00 (referent)	
		• Atherosclerosis Risk in Communities Study	Q4 = High	• Q2 = 0.74 (0.42-1.31)	
				• Q3 = 1.07 (0.55-2.09)	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

			Outcome Measure: CHD incidence Poisson Regression	<ul style="list-style-type: none"> <li>• Q4 = 0.64 (0.34-1.24)</li> </ul>	
				Multivariate RR (95% CI) Sports, men <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 1.15 (0.79-1.68)</li> <li>• Q3 = 1.03 (0.68-1.54)</li> <li>• Q4 = 0.83 (0.56-1.23)</li> </ul> Sports, women <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.99 (0.58-1.67)</li> <li>• Q3 = 0.64 (0.32-1.27)</li> <li>• Q4 = 0.72 (0.37-1.38)</li> </ul>	
Fransson et al 2004 [228]	To estimate the influence of LTPA and OPA on acute MI.	<ul style="list-style-type: none"> <li>• n = 4069 (2,742 men; 1,327 women)</li> </ul>	PA assessment: Questionnaire for LTPA, 5 groups	<ul style="list-style-type: none"> <li>• Number of Cases: 1,204 men, 550 women</li> </ul>	Exercise seems to reduce the risk of MI.
Sweden		<ul style="list-style-type: none"> <li>• Sex: Men and Women</li> </ul>	G1 = Seldom G2 = Sometimes	OR (95% CI)	
Case Control		<ul style="list-style-type: none"> <li>• Age: 45-70 yr</li> <li>• Characteristics: Cases: Diagnosed with acute MI</li> </ul>	G3 = 1x/wk G4 = 2-3x/wk	LTPA, men <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
D & B score = 12			G5 = >3x/wk	<ul style="list-style-type: none"> <li>• G2 = 0.76 (0.61-0.95)</li> <li>• G3 = 0.67 (0.51-0.88)</li> <li>• G4 = 0.63 (0.49-0.83)</li> <li>• G5 = 0.53 (0.38-0.73)</li> </ul>	
		<ul style="list-style-type: none"> <li>• Stockholm Heart Epidemiology</li> </ul>	Questionnaire for total physical activity, 3 groups G1 = Passive G2 = Somewhat active G3 = Active	LTPA, women <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.69 (0.49-0.98)</li> <li>• G3 = 0.38 (0.25-0.58)</li> <li>• G4 = 0.62 (0.38-1.01)</li> <li>• G5 = 0.31 (0.15-0.66)</li> </ul>	
			Questionnaire for sitting at work, 3 groups G1 = Less than half the time G2 = About half the time G3 = More than half the time	Total physical activity, men <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.66 (0.47-0.94)</li> </ul>	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

			Outcome Measure: Acute MI	<ul style="list-style-type: none"> <li>• G3 = 0.46 (0.31-0.69)</li> </ul>	
			Conditional and unconditional logistics regression	Total physical activity, women <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.34 (0.22-0.53)</li> <li>• G3 = 0.16 (0.07-0.37)</li> </ul> Sitting at work, men <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.91 (0.73-1.15)</li> <li>• G3 = 0.90 (0.72-1.12)</li> </ul> Sitting at work, women <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.77 (0.51-1.17)</li> <li>• G3 = 0.47 (0.31-0.69)</li> </ul>	
Fransson et al 2006 [229]	To evaluate whether LTPA compensates for the increased risk of acute MI associated with overweight and obesity.	• n = 4069 (2,742 men; 1,327 women)	PA Assessment: Questionnaire for LTPA, 3 groups	Number of Cases: 1204 men, 550 women	Regular LTPA seems to provide protection against MI and non- fatal MI.
Sweden		• Sex: Men and women	G1 = Very little /occasional walks	Multivariate OR (95% CI) for acute MI LTPA, men	
Case Control		• Age: 45-70 yr	G2 = Occasional / once per week	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.70 (0.58-0.84)</li> <li>• G3 = 0.57 (0.46-0.71)</li> </ul>	
D & B score = 12		• Characteristics: Cases: had acute MI	G3 = Twice per week or more	LTPA, women	
			Outcome measure: Acute MI	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.52 (0.40-0.68)</li> <li>• G3 = 0.44 (0.30-0.65)</li> </ul>	
			Conditional and unconditional logistics regression	Multivariate OR (95% CI) for non-fatal MI LTPA, men <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.79 (0.65-0.96)</li> <li>• G3 = 0.63 (0.50-0.79)</li> </ul>	



**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

				LTPA, women	
				• G1 = 1.00 (referent)	
				• G2 = 0.64 (0.48-0.86)	
				• G3 = 0.58 (0.39-0.87)	
Haapanen-Niemi 2000 [230]	To investigate the independent associations and the possible interaction of BMI LTPA and perceived physical performance and functional capacity with the risk of mortality.	• n = 2,212 (1,090 men; 1,122 women)	Length of Follow-up: 16 yrs	• Number of Cases: 208 all cause deaths, 54% of those CVD. 73% of CVD deaths due to CHD	Increase perceived PF is associated with a reduced risk of CHD mortality in men.
Finland		• Sex: Men and women	PA assessment: Postal Survey		
Prospective cohort		• Age: 35-63 yr	Total LTPA energy expenditure (kcal/wk)	Multivariate RR (95% CI) Total LTPA EE index and CHD mortality, men	
		• Characteristics: Healthy		• G1 = 1.00 (referent)	
			G1 = High	• G2 = 0.88 (0.44-1.76)	
			G2 = Moderate	• G3 = 1.70 (0.90-3.21)	
D & B score = 13		• Ethnicity: Finnish	G3 = Low	p = 0.056	
			Perceived physical fitness compared to age-mates	Multivariate RR (95% CI) Perceived physical fitness, men	
			G1 = Better	• G1 = 1.00 (referent)	
			G2 = Similar	• G2 = 2.82 (1.06-7.46)	
			G3 = Worse	• G3 = 4.64 (1.56-13.84)	
			Outcome Measure: CHD mortality	p = 0.011	
				Total LTPA EE index and CHD mortality, women	
			Cox proportional HR	• G1 = 1.00 (referent)	
				• G2 = 0.43 (0.16-1.16)	
				• G3 = 1.17 (0.51-2.68)	
				p = 0.046	
				Multivariate RR (95% CI) Perceived physical fitness, women	
				• G1 = 1.00 (referent)	
				• G2 = 0.82 (0.32-2.16)	
				• G3 = 1.89 (0.57-6.27)	
				p = 0.154	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Kannel et al 1986 [231]	To examine the role of low levels of OPA and LTPA in the development of CV morbidity and mortality over the short and long term.	<ul style="list-style-type: none"> <li>• n = 1,166</li> <li>• Sex: Men</li> <li>• Age: 45-65 yr</li> <li>• Characteristics:</li> </ul>	Length of Follow-up: 24 yrs	<ul style="list-style-type: none"> <li>• Number of Cases: 220 mortality, 371 morbidity</li> </ul>	Rate of CHD Mortality and Morbidity decreases with increased level of PA but no association was found with physical demand of work
USA			PA assessment: Questionnaire during examination	Cumulative 24 year age adjusted rate per 1000 people	
Prospective cohort				24 hr PA index for LTPA CHD mortality	
D & B score = 11			PA index: G1 = <29	<ul style="list-style-type: none"> <li>• G1 = 255</li> <li>• G2 = 184</li> </ul>	
			G2 = 30-34	• G3 = 152	
			G3 = >34	p < 0.01	
			Physical demand of work	24 hr PA index for LTPA CHD incidence	
			G1 = Sedentary	• G1 = 414	
			G2 = Light	• G2 = 353	
			G3 = Medium	• G3 = 311	
			G4 = Heavy		
			Outcome Measure: CHD mortality and Morbidity	Physical demand of work and CHD mortality	
				• G1 = 216	
			Cox proportional HR	• G2 = 209	
				• G3 = 169	
				• G4 = 170	
				Physical demand of work and CHD incidence:	
				• G1 = 355	
				• G2 = 405	
				• G3 = 307	
				• G4 = 325	
Kaprio et al 2000 [232]	To examine the contribution of genetic and other familial factors to the relationship between LTPA and CHD.	<ul style="list-style-type: none"> <li>• n = 8,205</li> <li>• Sex: Men</li> <li>• Age: 25-69 yr</li> <li>• Characteristics: Same sex twin pairs, free of CVD</li> </ul>	Length of Follow-up: 18 yrs	<ul style="list-style-type: none"> <li>• Number of Cases: 723</li> </ul>	LTPA compared to being sedentary helps prevent CHD in men.
Finland			PA assessment: Questionnaire for LTPA, 3 groups:	Multivariate RR (95% CI)	
				• G1 = 1.00 (referent)	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Prospective cohort	D & B score = 12		G1 = Sedentary G2 = Occasional Exercisers  G3 = Conditioning Exercisers Outcome Measure: Hospitalization or death from CHD Poisson regression	$p = 0.010$	<ul style="list-style-type: none"> <li>• G2 = 0.84 (0.70-1.01)</li> <li>• G3 = 0.68 (0.50-0.92)</li> </ul>
Lakka et al 1994 [233]	To investigate the independent associations of LTPA and maximal oxygen uptake with the risk of acute MI.	<ul style="list-style-type: none"> <li>• n = 1,166</li> <li>• Sex: Men</li> <li>• Age: 42-61 yr</li> </ul>	Baseline examination: 1984-1989		Conditioning LTPA and VO <sub>2</sub> max had an inverse, graded and independent association with the risk
Finland		<ul style="list-style-type: none"> <li>• Characteristics: Healthy with normal ECG</li> </ul>	PA assessment: Questionnaire for conditioning PA (h/wk), 3 groups (h/wk)		
Prospective cohort			G1 = <0.7 G2 = 0.7 G3 = >2.2	Adjusted RH (95% CI) by conditioning PA level  <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.11 (0.58-2.12)</li> <li>• G3 = 0.31(0.12-0.85)</li> </ul>	
D & B score = 13		<ul style="list-style-type: none"> <li>• Kuopio Ischaemic Heart Disease Risk Factor Study</li> </ul>	PF assessment: VO <sub>2</sub> max (ml/kg/min)  G1 = <28.0 G2 = 28.0-33.6 G3 = >33.6 Outcome event: acute MI Cox proportional HR	Adjusted RG (95% CI) by VO <sub>2</sub> max  <ul style="list-style-type: none"> <li>• G1 = 1.00</li> <li>• G2 = 0.76 (0.38-1.50)</li> <li>• G3 = 0.26 (0.10-0.68)</li> </ul>	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Laukkanen et al 2004 [234]	To determine whether $VO_{2peak}$ predicts CVD morbidity and mortality in a sample of men as related to conventional risk factors, medications or underlying chronic disease.	<ul style="list-style-type: none"> <li>• 1,294 healthy; 1,057 unhealthy</li> </ul>	PF Assessment: $VO_2$ peak (ml/kg/min) measured by exercise test with an electrically braked cycle ergometer, divided into quartiles	<ul style="list-style-type: none"> <li>• Number of Cases: 204 CV deaths, 323 non-fatal coronary events</li> </ul>	Dose-response relationship between directly measured PF and CVD death among healthy men at baseline.
Finland		<ul style="list-style-type: none"> <li>• Sex: Men</li> </ul>		Healthy men with low $VO_2$ peak (lowest quartile) had an increased risk	
Prospective cohort		<ul style="list-style-type: none"> <li>• Age: 42-60 yr</li> <li>• Characteristics: Healthy and not healthy participants</li> </ul>	Q1 = <27.6 Q2 = 27.6-32.2	Adjusted RR (95% CI) by PF quartile Fatal MI	Unfit men with unfavorable risk profiles are the risk group that would benefit the most from preventative measures.
D & B score = 11			Q3 = 32.3-37.1	• 3.29 (0.86-12.90)	
		<ul style="list-style-type: none"> <li>• Kuopio Ischaemic Heart Disease Risk Factor Study</li> </ul>	Q4 = >37.2		
			Outcome Measure: Incidence of fatal and non fatal CVD during 13 year follow-up Cox proportional HR	Non-Fatal MI • 2.16 (1.12-4.18)	
Lee et al 2000 [235]	To investigate whether different durations of exercise episode are associated with different risk of CHD.	<ul style="list-style-type: none"> <li>• n = 7,307</li> </ul>	Baseline survey in 1988	<ul style="list-style-type: none"> <li>• Number of Cases: 482</li> </ul>	Longer durations of PA bouts are not associated with decreased CHD risk compared with shorter bouts, once total EE is taken into account.
USA		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: Mean 66.1 ± 7.5</li> </ul>	PA assessment: Survey for EE (kJ/wk), divided into 5 groups and episodes of PA (min), divided into 6 groups	Multivariate adjusted RR (95% CI) by EE	
		<ul style="list-style-type: none"> <li>• Characteristics: Healthy</li> </ul>		<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.80 (0.57-1.12)</li> <li>• G3 = 0.80 (0.55-1.16)</li> <li>• G4 = 0.74 (0.47-1.17)</li> </ul>	
Prospective cohort		<ul style="list-style-type: none"> <li>• Harvard Alumni Study</li> </ul>		<ul style="list-style-type: none"> <li>• G5 = 0.62 (0.41-0.94)</li> </ul>	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

D & B score = 12			Energy expenditure (kJ/wk)		As long as the total EE is similar, more frequent shorter bouts or longer less frequent bouts have an equivalent reduction in CHD risk.
			G1 = <4,200	Multivariate adjusted RR (95% CI) by duration of PA episode	
			G2 = 4,200-8,399		
			G3 = 8,400-12,599		
			G4 = 12,600-16,799	• G1 = 1.00 (referent)	
			G5 = ≥ 16,800	• G2 = 1.15 (0.70-1.87)	
				• G3 = 1.01 (0.68-1.51)	
				• G4 = 1.11 (0.67-1.84)	
			Duration of PA episode (min)	• G5 = 1.18 (0.77-1.80)	
			G1 = None	• G6 = 1.25 (0.83-1.87)	
			G2 = 1-15		
			G3 = 16-30		
			G4 = 31-45		
			G5 = 46-60		
			G6 = >60		
			Outcome Measure: Fatal and Non Fatal CHD		
			Proportional hazards regression		
Lee et al 2003 [236]	To investigate whether moderate-intensity exercise is associated with reduced CHD.	• n = 7,337	PA assessment: Survey rating usual level of exertion when exercising, divided into tertiles	• Number of Cases: 551	Inverse association between relative intensity of PA and the risk of CHD.
USA		• Sex: Male		Multivariate adjustment RR (95% CI)	
		• Age: Mean 66.1 yr		• T1 = 1.00 (referent)	
		• Characteristics: Healthy		• T2 = 0.87 (0.70-1.09)	
				• T3 = 0.92 (0.75-1.14)	
Prospective cohort		Harvard Alumni Study	Energy expenditure (kcal/wk)		
			T1 = <1000		
			T2 = 1000-2499		
			T3 = ≥ 2500		
			Cox proportional HR		
D & B score = 13					

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Lemaitre et al 1999 [237]	To investigate whether regular participation in moderate intensity activity confers overall protection from sudden primary cardiac arrest.	<ul style="list-style-type: none"> <li>• n = 355 cases, 503 controls</li> </ul>	PA assessment: Interview (with spouses) for LTPA, 7 groups	<ul style="list-style-type: none"> <li>• 355 cases</li> </ul>	Participation in moderate intensity LTPA was associated with a decreased risk of primary cardiac arrest.
USA		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: 25-74 yr</li> </ul>	G1 = No activity G2 = Gardening only ≤ 60 min/wk G3 = Gardening only > 60 min/wk G4 = Walking ≤ 60 min/wk G5 = Walking > 60 min/wk G6 = Moderate intensity LTPA (not walking or gardening) G7 = High intensity LTPA Logistic regression analysis	RR (95% CI) <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.52 (0.21-1.28)</li> <li>• G3 = 0.34 (0.13-0.89)</li> <li>• G4 = 0.45 (0.17-1.19)</li> <li>• G5 = 0.27 (0.11-0.67)</li> <li>• G6 = 0.31 (0.13-0.74)</li> <li>• G7 = 0.34 (0.16-0.75)</li> </ul>	
Case control		<ul style="list-style-type: none"> <li>• Characteristics: Previously healthy prior to primary cardiac arrest. Control Subjects: Individually matched to case patients on age (within 7 years) and sex at a ratio of about 2:1 were randomly selected from community by random-digit dialing</li> </ul>			
D & B score = 11					
Lemaitre et al 1995 [238]	To examine whether LTPA decreases the risk of MI in postmenopausal women.	<ul style="list-style-type: none"> <li>• n = 1,193</li> </ul>	PA assessment: Phone interview for LTPA, divided into quartiles of EE (mean kcal/wk)	<ul style="list-style-type: none"> <li>• Number of Cases: 268</li> </ul>	Risk of MI among postmenopausal women is decreased by 50% with modest LT energy expenditures, equivalent to 30-45 min of walking for exercise three times per week
USA		<ul style="list-style-type: none"> <li>• Sex: Women</li> <li>• Age: Mean 67 yr</li> <li>• Characteristics: Postmenopausal Cases: Diagnosed with non-fatal MI Controls: free from MI</li> </ul>	Q1 = 71 Q2 = 472 Q3 = 1183 Q4 = 3576 Outcome Measure: Diagnosed with non-fatal MI Logistic regression analysis	Multivariate RR (95% CI) <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.52 (0.34-0.80)</li> <li>• Q3 = 0.40 (0.26-0.63)</li> <li>• Q4 = 0.40 (0.25-0.63)</li> </ul> <p>p = &lt;0.001</p>	
Case control					
D & B score = 11					

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Li et al 2006 [239]	To examine independent and joint associations of PA and adiposity with CHD incidence.	<ul style="list-style-type: none"> <li>• n = 88,393</li> <li>• Sex: Women</li> </ul>	Length of Follow-up: 20 yrs	<ul style="list-style-type: none"> <li>• Number of Cases: 2,358</li> <li>• Number of Dropouts: &lt;2% lost to follow-up contributes to the development of CHD in women.</li> </ul>	Physical inactivity independently contributes to the development of CHD in women.
USA		<ul style="list-style-type: none"> <li>• Age: 34-59 yr</li> <li>• Characteristics: Nurses</li> </ul>	PA assessment: Questionnaire for LTPA (hr/wk), 3 groups	up	
Prospective cohort		<ul style="list-style-type: none"> <li>• Nurses' Health Study</li> </ul>		Multivariate HR (95% CI)	
D & B score = 12			G1 = ≥3.5 G2 = 1-3.49 G3 = <1 Outcome Measure: CHD incidence Cox proportional HR	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.34 (1.18-1.51)</li> <li>• G3 = 1.43 (1.26-1.63)</li> </ul>	
Lemaitre et al 1995 [240]	To evaluate the effect of PA on MI occurrence.	<ul style="list-style-type: none"> <li>• n = 1,107 (726 controls, 381 cases)</li> </ul>	PA assessment: Questionnaire, 3-5 groups depending on variable	OR (95% CI),	PA level was inversely associated with occurrence of MI in both sexes, although the association presented a significant linear trend only for women; in men it suggested a u-shaped relation.
Portugal		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>		Total PA, men	
Case control		<ul style="list-style-type: none"> <li>• Age: ≥ 40 yr</li> <li>• Characteristics: Case: Admitted to Hospital and diagnosed with first episode of MI Control: Healthy, no history of CHD</li> </ul>	Total PA (MET hr/day), men	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.54 (0.33-0.88)</li> <li>• G3 = 0.34 (0.20-0.59)</li> <li>• G4 = 0.59 (0.36-0.98)</li> </ul>	
D & B score = 12			G1 = 28.3-32.1 G2 = 32.2-33.3 G3 = 33.4-36.5 G4 = 36.6-40.3 G5 = 40.4-83.1 Total PA (MET hr/day), women	<ul style="list-style-type: none"> <li>• G5 = 0.90 (0.56-1.45)</li> </ul> Trend $p = 0.827$ Total PA, women	
				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.39 (0.21-0.73)</li> </ul>	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

			Q1 = 28.9-32.7 Q2 = 32.8-34.1 Q3 = 34.2-37.8 Q4 = 37.8-70.6	• Q3 = 0.33 (0.17-0.64) • Q4 = 0.22 (0.11-0.47) $p = <0.001$	
			Sport participation (MET hr/day), men G1 = 0.0 G2 = 0.1-1.0 G3 = 1.1-2.0 G4 = 2.1-3.6 G5 = 3.7-15.4	Sport participation, men • G1 = 1.00 (referent) • G2 = 0.36 (0.19-0.69), • G3 = 0.72 (0.41-1.26), • G4 = 0.42 (0.23-0.76), • G5 = 0.31 (0.16-0.62) $p = <0.001$	
Lovasi et al 2007 [241]	To investigate the shape of the relationship between LTPA and MI risk.	• n = 4,094	PA assessment: Telephone interview (Minnesota LTPA Questionnaire)	• Number of Cases: 697	Time engaged in LTPA, even non strenuous LTPA was associated with a lower risk of MI, and the shape of this relationship was non- linear
USA		• Sex: Men and women • Age: 64 ± 9 yr • Characteristics: Group Health Cooperative Members		Adjusted OR (95% CI) LTPA and non fatal CHD • G1 = 1.00 (referent)	
Case control			LTPA G1 = None G2 = <2 G3 = 2-5 G4 = 5-9 G5 = >9 h/wk	• G2 = 0.88 (0.66-1.17) • G3 = 0.62 (0.46-0.83) • G4 = 0.61 (0.45-0.82) • G5 = 0.59 (0.44-0.80)	
D & B score = 11			Strenuous LTPA G1 = None G2 = non strenuous LTPA G3 = Any Strenuous LTPA Outcome measure: non fatal CHD Logistic regression	Adjusted RR (95% CI) Strenuous LTPA and non Fatal CHD • G1 = 1.00 (referent) • G2 = 0.76 (0.59-0.99) • G3 = 0.53 (0.40-0.70)	



**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Manson et al 1999 [242]	To assess the comparative roles of walking and vigorous exercise in the prevention of coronary events in women.	<ul style="list-style-type: none"> <li>• n = 72,488</li> </ul>	PA assessment:	<ul style="list-style-type: none"> <li>• Number of Cases: 645 coronary events</li> </ul>	Both walking and VPA are associated with a substantial reductions in incidence of CHD. Risk reductions for each were similar hen total PAy was similar. Walking 3 or more hours per week could reduce the risk of CHD by 30-40%.
		<ul style="list-style-type: none"> <li>• Sex: Women</li> </ul>	Questionnaire with detailed information on PA.		
		<ul style="list-style-type: none"> <li>• Age: 40-65 yr</li> </ul>		Multivariate RR (95% CI) by total PA score	
USA		<ul style="list-style-type: none"> <li>• Characteristics: Healthy, no Previous history of CHD</li> </ul>		<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
Prospective cohort			Total PA score	<ul style="list-style-type: none"> <li>• G2 = 0.88 (0.71-1.10)</li> <li>• G3 = 0.81(0.64-1.02)</li> </ul>	
			G1 = 1-2.0	<ul style="list-style-type: none"> <li>• G4 = 0.74 (0.58-0.95)</li> </ul>	
		Nurses' Health Study	G2 = 2.1-4.6	<ul style="list-style-type: none"> <li>• G5 = 0.66 (0.51-0.86)</li> </ul>	
D & B score = 12			G3 = 4.7-10.4	p = 0.002	
			G4 = 10.5-21.7		
			G5 = >21.7		
				Multivariate RR (95% CI) by walking activity	
			Walking, in those who did not participate in VPA: (MET hr/wk)	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.78 (0.57-1.06)</li> </ul>	
			G1 = 0.5	<ul style="list-style-type: none"> <li>• G3 = 0.88 (0.65-1.21)</li> </ul>	
			G2 = 0.6-2.0	<ul style="list-style-type: none"> <li>• G4 = 0.70 (0.51-0.95)</li> </ul>	
			G3 = 2.1-3.8	<ul style="list-style-type: none"> <li>• G5 = 0.65 (0.47-0.91)</li> </ul>	
			G4 = 3.9-9.9	p = 0.02	
			G5 = ≥ 10		
				Multivariate RR (95% CI) by walking pace	
			Walking pace (mph)	<ul style="list-style-type: none"> <li>• 1.00 (referent)</li> <li>• 0.75 (0.59-0.96)</li> <li>• 0.64 (0.47-0.88)</li> </ul>	
			G1 = <2.0		
			G2 = 2.0-2.9		
			G3 = ≥ 3.0		

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Mora et al 2007 [243]	To investigate whether differences in several CV risk factors mediate the effect of PA on reduced risk of CVD.	<ul style="list-style-type: none"> <li>• n = 27,055</li> <li>• Sex: Women</li> <li>• Age: ≥ 45 yr</li> </ul>	10.9 ± 1.6 yr of follow up	<ul style="list-style-type: none"> <li>• Number of Cases: 640</li> </ul>	There remained a borderline significant inverse association between PA and risk of CHD after adjustment for all sets of risk factors.
USA		<ul style="list-style-type: none"> <li>• Characteristics: Healthy</li> </ul>	PA assessment: Questionnaires at study entry for categories of EE from PA (kcal/wk), 4 groups	HR (95% CI), basic model	
Prospective cohort		<ul style="list-style-type: none"> <li>• Women's health study</li> </ul>		<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.84 (0.67-1.06)</li> <li>• G3 = 0.76 (0.61-0.96)</li> <li>• G4 = 0.62 (0.48-0.82)</li> </ul>	
D & B score = 13			G1 = <200 G2 = 200-599	p = 0.001	While all sets of risk factors should some mediation on the effect of PA on CHD none made the relationship insignificant
			G3 = 600-1499 G4 = ≥ 1500	Multivariate adjusted HR (95% CI)	
			Outcome measure: Incidence of CVD and	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.71 (0.58-0.87)</li> <li>• G3 = 0.64 (0.52-0.78)</li> <li>• G4 = 0.48 (0.38-0.62)</li> </ul>	
			Cox proportional HR	p = <0.001	
O'Connor et al 1995 [244]	To examine the association between intensity of exercise and CHD risk.	<ul style="list-style-type: none"> <li>• n = 680 (532 men and 148 women)</li> </ul>	PA assessment: Home interview for PA, divided into quartiles	<ul style="list-style-type: none"> <li>• Number of Cases: 340</li> </ul>	Significant inverse association between PA level and the risk of non fatal MI in men, which persisted after adjustment for other risk factors.
USA		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>	Q1 = Lowest Q2 Q3 Q4 = Highest	Adjusted OR (95% CI) by PA level, men	
Case control		<ul style="list-style-type: none"> <li>• Age: &lt; 76 yr</li> <li>• Characteristics: Cases: Diagnosed MI (non-fatal), no previous history of CHD. Controls: no history of CHD.</li> </ul>		<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.60 (0.32-1.13)</li> <li>• Q3 = 0.41 (0.21-0.78)</li> <li>• Q4 = 0.41 (0.22-0.77)</li> </ul>	
D & B score = 12			Outcome Measure: non-fatal MI	Adjusted OR (95% CI) by PA level, women	
				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> </ul>	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

		Moderate- vigorous sports men	Cut-points kcal/wk	<ul style="list-style-type: none"> <li>• Q2 = 1.07 (0.27-4.17)</li> </ul>	
		Q1 = Lowest		<ul style="list-style-type: none"> <li>• Q3 = 2.02 (0.56-7.38)</li> </ul>	
		Q2		<ul style="list-style-type: none"> <li>• Q4 = 1.29 (0.31-5.35)</li> </ul>	
		Q3		$p = 0.51$	
		Q4 = Highest			
				Adjusted OR (95% CI) by moderate-vigorous sports, men	
				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> </ul>	
		Moderate- vigorous sports Women		<ul style="list-style-type: none"> <li>• Q2 = 1.12 (0.60-2.10)</li> </ul>	
		Cut-points kcal/wk		<ul style="list-style-type: none"> <li>• Q3 = 0.61 (0.30-1.24)</li> </ul>	
		Q1 = Lowest		<ul style="list-style-type: none"> <li>• Q4 = 0.43 (0.20-0.92)</li> </ul>	
		Q2		$p = 0.02$	
		Q3			
		Q4 = Highest		Adjusted OR (95% CI) by moderate-vigorous sports, women	
		Logistic regression analysis		<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> </ul>	
				<ul style="list-style-type: none"> <li>• Q2 = 1.31 (0.37-4.66)</li> </ul>	
				<ul style="list-style-type: none"> <li>• Q3 = 1.90 (0.44-8.28)</li> </ul>	
				<ul style="list-style-type: none"> <li>• Q4 = 0.35 (0.07-1.84)</li> </ul>	
				$p = 0.62$	
Rastogi et al 2004 [245]	To examine the relation between PA and CHD risk in India.	• n = 1,050	PA assessment: Questionnaire	Number of Cases: 350	Observed a strong and dose dependent inverse association between LTPA and non fatal CHD.
		• Sex: Men and women		Multivariate OR (95% CI) by LTPA	
USA		• Age: 21-74 yr	LTPA (MET min/d)	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
		• Characteristics: Cases: Diagnosed with MI (non fatal) Controls: non- cardiac patients	G1 = 0	<ul style="list-style-type: none"> <li>• G2 = 0.96 (0.59-1.55)</li> </ul>	
Case control			G2 = 0-145	<ul style="list-style-type: none"> <li>• G3 = 0.44 (0.27-0.71)</li> </ul>	
D & B score = 12			G3 = ≥145	$p = 0.001$	
			Sedentary time (min/d)	Multivariate OR (95% CI) by sedentary time	
			G1 = <70	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
			G2 = 70-130	<ul style="list-style-type: none"> <li>• G2 = 1.15 (0.68-1.95)</li> </ul>	
			G3 = 130-215	<ul style="list-style-type: none"> <li>• G3 = 1.04 (0.61-1.76)</li> </ul>	
			G4 = ≥215	<ul style="list-style-type: none"> <li>• G4 = 1.88 (1.09-3.21)</li> </ul>	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

			<i>p</i> = 0.02		
			Outcome Measure: Non-fatal MI		
			Conditional logistic regression		
Rodriguez et al 1994 [246]	To examine the relationship between PA and 23 yr incidence of CHD morbidity and mortality.	<ul style="list-style-type: none"> <li>• n = 7,074</li> <li>• Sex: Men</li> <li>• Age: 45-64 yr</li> </ul>	23 year follow-up	<ul style="list-style-type: none"> <li>• Number of Cases: 789</li> </ul>	PA was associated with a significant reduction in the risk of CHD morbidity and mortality.
USA		<ul style="list-style-type: none"> <li>• Characteristics: Japanese- American living in Oahu, Hawaii in 1965, &lt; 65 years to reduce effect of retirement on PA levels</li> </ul>	PA assessment: Questionnaire for PA index, divided into tertiles	Age adjusted RR (95% CI), CHD incidence	
Prospective cohort			T1 = Low	<ul style="list-style-type: none"> <li>• T1 = 1.00 (referent)</li> <li>• T2 = 1.01 (.86-1.19)</li> <li>• T3 = 0.83 (0.86-1.19)</li> </ul>	These data support the hypothesis that PA is associated with a favorable profile of CVD risk factors.
			T2 = Moderate		
			T3 = High		
D & B score = 11			Cox proportional regression model	Multivariate adjusted RR (95% CI), CHD incidence	
				<ul style="list-style-type: none"> <li>• T1 = 1.00 (referent)</li> <li>• T2 = 1.07 (0.90-1.26)</li> </ul>	This study did not show a dose-response relationship since the medium tertile of PA showed increased rates of CHD compared to the inactive group.
		<ul style="list-style-type: none"> <li>• The Honolulu Heart Program</li> </ul>		<ul style="list-style-type: none"> <li>• T3 = 0.95 (0.80-1.14)</li> </ul>	
				Age adjusted RR (95% CI), CHD mortality	
				<ul style="list-style-type: none"> <li>• T1 = 1.00 (referent)</li> <li>• T2 = 1.12 (0.88-1.44)</li> <li>• T3 = 0.74 (0.56-0.97)</li> </ul>	
				Multivariate adjusted RR (95% CI)	
				<ul style="list-style-type: none"> <li>• T1 = 1.00 (referent)</li> <li>• T2 = 1.19 (0.93-1.53)</li> <li>• T3 = 0.85 (0.65-1.13)</li> </ul>	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Rothenbacher et al 2003 [247]	To estimate the risk for CHD associated with LTPA.	• n = 791 (312 cases; 479 controls)	PA assessment: Interview	Number of Cases: 312	LTPA showed a clear inverse association with risk of CHD.
Germany		• Sex: Men and Women	LTPA (h/wk) G1 = 0 G2 = <1 G3 = 1-2 G4 = >2	Multivariate OR (95% CI), LTPA Winter • G1 = 1.00 (referent) • G2 = 0.48 (0.27-0.84) • G3 = 0.54 (0.369-0.82)  • G4 = 0.27 (0.19-0.47)	
Case control	Age: 40-68 yr Characteristics: Cases: stable CHD diagnosed within 2 years, no recent MI, Controls: no history of CHD.				
D & B score = 12			Workday activity by bike/foot, (min/workday) G1 = <15 G2 = 15-30 G3 = 30-60 G4 = >60 Outcome Measure: non fatal CHD Unconditional logistic regression, linear regression model	Summer • G1 = 1.00 (referent) • G2 = 0.85 (0.47-1.53) • G3 = 0.60 (0.38-0.95) • G4 = 0.39 (0.26-0.59) Multivariate OR (95% CI), workday activity by bike/foot • G1 = 1.00 (referent)  • G2 = 0.53 (0.30-0.93) • G3 = 0.36 (0.21-0.62) • G4 = 0.58 (0.36-0.94)	
Seccareccia and Menotti 1992 [248]	To examine the relationship between OPA and the risk of CHD death.	• n = 1,621  • Sex: Men • Age: 40-59 yr  • Characteristics: Healthy	25 year of follow-up  PA assessment: Questionnaire for OPA (kcal/d), 3 groups  G1 = Sedentary, < 2400 G2 = Moderate, 2400-3199 G3 = Heavy ≥ 3200	• 189 cases  Age Standardized CHD and deaths rates:  • G1 = 18.9 ± 3.1 • G2 = 13.1 ± 1.7 • G3 = 11.0 ± 0.9	Increase in OPA is inversely related to risk of CHD death.
Italy					
Prospective cohort D & B score = 11					

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

			Indicators of PF including HR, vital capacity, FEV in 3/4 of sec, and corrected arm circumference (minus contribution of fat). End Point: Fatal CHD		
Sesso et al 2000 [249]	To examine the association of the quantity and intensity of PA with CHD risk and the impact of other coronary risk factors.	• n = 12,516	PA assessment: Questionnaire	Number of Cases: 2,135	L-Shaped association between PA and the risk of CHD, with a reduction in CHD risk of approximately 20% for total PA levels >4200 kJ/wk
USA		• Sex: Men • Age: 39-88 yr • Characteristics: Healthy	PA Index (kJ/wk) G1 = <2100 G2 = 2100-4199	Multivariate HR (95% CI) • G1 = 1.00 (referent) • G2 = 0.90 (0.79-1.03) • G3 = 0.81 (0.71-0.92)	
Prospective cohort		• Harvard Alumni	G3 = 4200-8399 G4 = 8400-12599 G5 = >12600	• G4 = 0.80 (0.69-0.93) • G5 = 0.81 (0.71-0.94) p = 0.003	
D & B score = 12		Study			Suggests that vigorous activities are associated with a reduced risk of CHD, whereas moderate or light PA has no clear association with risk of CHD.
			Cox proportional HR		
Sundquist et al 2005 [250]	To examine the long term effect of LTPA on incident cases of CHD.	• n = 5,196 (2,645 men, 2,551 women)	PA assessment: Questionnaire Levels of PA	Age and sex adjusted RR (95% CI)	Positive long term effect of LTPA on CHD risk among men and women.
Sweden		• Sex: Men and women	Q1 = None Q2 = Occasionally Q3 = 1-2 times per week	• Q1 = 1.00 (referent) • Q2 = 0.72 (0.51-1.00) • Q3 = 0.64 (0.46-0.89) • Q4 = 0.46 (0.29-0.74)	
Prospective cohort		Age: 35-74 yr	Q4 = Vigorous ≥2 times per week	Multivariate adjusted RR (95% CI)	
D & B score = 11		• Characteristics: Those not hospitalized for CHD in the last 2 years and those who rate their general health as poor were excluded	Outcome Measure: Fatal or non fatal CHD	• Q1 = 1.00 (referent) • Q2 = 0.76 (0.55-1.07) • Q3 = 0.74 (0.53-1.04) • Q4 = 0.59 (0.37-0.95)	
			Cox regression model		

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Talbot et al 2002 [251]	To examine the contributions of LTPA and aerobic fitness to the risk of coronary events in healthy younger and older adults.	<ul style="list-style-type: none"> <li>• n = 689</li> <li>• Sex: Men</li> <li>• Age:</li> </ul>	Surveys began in 1960 and were completed on every visit	<ul style="list-style-type: none"> <li>• Number of Cases: 63</li> </ul>	In younger men PF predicts a reduced risk of CHD but not LTPA.
USA		51.6 ± 16.8 yr		After adjusting for coronary risk factors there was:	
Prospective cohort		<ul style="list-style-type: none"> <li>• Characteristics: Community dwelling</li> </ul>	PA assessment: Survey for LTPA (97 activities) at every visit.	RR: 0.53 (p < 0.001) and	In older men, high intensity LTPA and PF appear to be of similar importance in reducing CHD risk.
D & B score = 12		<ul style="list-style-type: none"> <li>• Baltimore Longitudinal Study of Aging</li> </ul>	PF assessment: Treadmill VO <sub>2</sub> max test on alternate visits	RR: 0.61 (p = 0.024) in older men. Total LTPA was unrelated to coronary risk in either age group. With 3 levels of LTPA intensity substituted for total LTPA: RR = 0.39 for tertile 3 vs. tertile 1	
Tanasescu et al 2002 [252]	To assess the amount, type and intensity of PA in relation to risk of CHD in men.	<ul style="list-style-type: none"> <li>• n = 44,452</li> <li>• Sex: Men</li> <li>• Age: 40-75 yr</li> </ul>	PA assessment: Questionnaire	<ul style="list-style-type: none"> <li>• Number of Cases: 1,700</li> </ul>	Total PA, running, weight training, and walking were associated with a reduced risk for CVD.
USA		<ul style="list-style-type: none"> <li>• Characteristics: Health professionals, no history of CHD and in good health</li> </ul>	Total PA (MET hr/wk)	Age adjusted HR (95% CI) by total PA	
Prospective cohort			Q1 = 0-6.32	• Q1 = 1.00 (referent)	
			Q2 = 6.33-14.49	• Q2 = 0.85 (0.74-0.98)	
			Q3 = 14.50-25.08	• Q3 = 0.78 (0.67-0.92)	
			Q4 = 25.09-41.98	• Q4 = 0.72 (0.62-0.83)	The average exercise intensity was associated with a reduced risk (independent of total PA).
D & B score = 11			Q5 = > 41.99	• Q5 = 0.58 (0.49-0.68) p = .001	
		<ul style="list-style-type: none"> <li>• Health Professionals follow-up study</li> </ul>	Exercise intensity (METs)	Age adjusted HR (95% CI) by exercise intensity	
			G1 = Low-1-4	• G1 = .00 (referent)	

**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

			<p>G2 = Mod.-4-6 G3 = High 6-12 Walking pace independent of total volume of PA (mph)</p> <p>Q1 = &lt;2 Q2 = 2-3 Q3 = 3-4 Q4 = &gt; 4</p> <p>Outcome Measure: Nonfatal MI or Fatal CHD occurring during follow-up Cox proportional HR</p>	<p>• G2 = 0.94 (0.83-1.04) • G3 = 0.83 (0.72-0.97)</p> <p><math>p = 0.02</math> Age adjusted HR (95% CI) by walking pace</p> <p>• Q1 = 1.00 (referent) • Q2 = 0.72 (0.54-0.94) • Q3 = 0.61 (0.45-0.81) • Q4 = 0.51 (0.31-0.84)</p> <p><math>p &lt; 0.001</math></p>	
Vatten et al 2006 [253]	To investigate whether obesity-related CV mortality could be modified by PA.	• n = 54,284 (27,769 men; 26,515 women)	Length of Follow-up: 16 years	• Number of Cases: 2,462	Increased PA reduces the risk of death in women, but not in men.
Norway		• Sex: Men and women	PA assessment: Questionnaire	Multivariate HR (95% CI), men	
Prospective cohort		Age: ≥ 20 yr	Divided into 4 groups	• Q1 = 1.00 (referent) • Q2 = 1.01 (0.89-1.16) • Q3 = 0.98 (0.84-1.14)	
		• Characteristics: Free from CVD at baseline	Q1 = High	• Q4 = 1.18 (1.00-1.38)	
			Q2 = Medium Q3 = Low	$p = 0.11$	
D & B score = 12		• HUNT study	Q4 = Never	Multivariate HR (95% CI), women	
			Outcome Measure: Ischemic heart disease mortality	• Q1 = 1.00 (referent) • Q2 = 1.23 (1.01-1.51) • Q3 = 1.54 (1.24-1.91) • Q4 = 1.52 (1.23-1.88)	
			Cox proportional HR	$p < 0.001$	




**Table 12: Studies examining the relationship between physical activity and cardiovascular disease. (Continued)**

Wagner et al 2002 [254]	To investigate if the association between PA patterns and incidence of coronary events could explain the gradient in CHD observed between 2 countries.	<ul style="list-style-type: none"> <li>• n = 9,758</li> </ul>	Length of Follow-up: 5 yrs	Number of Cases: 167 hard CHD, 154 angina events	Beneficial effect of LTPA EE on hard CHD incidence in middle aged men.
		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>	PA assessment: Questionnaire for LTPA, 3 groups:	Number of Dropouts: < 2%	
Ireland/France		<ul style="list-style-type: none"> <li>• Age: 50-59 yr</li> <li>• Characteristics: Healthy at Baseline</li> </ul>		HR (95% CI), hard events	
Prospective cohort			G1 = Lowest	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
			G2 = Middle	<ul style="list-style-type: none"> <li>• G2 = 0.73 (0.51-1.05)</li> </ul>	
			G3 = Highest	<ul style="list-style-type: none"> <li>• G3 = 0.66 (0.46-0.96)</li> </ul>	
D & B score = 12			Outcome Measure: CHD hard events and Angina	p = 0.04	
				HR (95% CI), angina	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
			Cox proportional HR	<ul style="list-style-type: none"> <li>• G2 = 0.83 (0.55-1.25)</li> <li>• G3 = 1.28 (0.88-1.86)</li> </ul>	
				p = 0.10	

D & B score, Downs and Black quality score; YR, years; G, groups; CHD, coronary heart disease; RR, risk ratio; 95% CI, 95% confidence interval; PA, physical activity; VPA, vigorous physical activity; CV, cardio vascular; MET, metabolic equivalent; kcal/wk, kilocalories per week; Q, quartile or quintile; km/h, kilometers per hour; LTPA, leisure-time physical activity; HR, hazard ratio; OPA, occupational physical activity; kcal/kg/day kilocalories per kilogram per day; MI, myocardial infarction; ECG, electrocardiogram; kcal/kg/h kilocalories per kilogram per hour; mph, miles per hour; CVD, cardiovascular disease.

30% risk reduction in the most active individuals [31]. In fact, in a review of the literature Katzmarzyk and Janssen [20] reported that lack of physical activity carried a relative risk of 1.60 (95% CI = 1.42-1.80) for stroke, similar to or higher than that for coronary heart disease (1.45), hypertension (1.30), colon cancer (1.41), breast cancer (1.31), type 2 diabetes (1.50), and osteoporosis (1.59).

In our systematic review of the literature, a total of 1104 citations were identified during the electronic database search (Figure 5). Of these citations, 405 were identified in MEDLINE, 183 in EMBASE, 227 in Cochrane, and 289 in the CINAHL/SportDiscus/PsychInfo search.



**Figure 5** Results of the Literature Search for Stroke.

A total of 13 duplicates were found, leaving a total of 1091 unique citations. A total of 1011 articles were excluded after scanning, leaving a total of 80 articles for full review. An additional 9 articles were retrieved through cross-referencing and the authors' knowledge of the field. From these articles 64 were excluded after full review leaving 25 articles for inclusion in the systematic review. The reasons for exclusion included non-experimental/weak design (poor execution introducing bias) ( $n = 16$ ), did not contain three levels of physical activity or not possible to determine dose-response relationship ( $n = 14$ ), reviews, summaries, meta-analyses ( $n = 17$ ), dissertations, thesis, abstracts ( $n = 8$ ), and other ( $n = 9$ ). Therefore, a total of 25 articles were included in the systematic review of the literature regarding the relationship between physical activity and the primary prevention of stroke (Table 13).

The data providing dose-response information is all observational in nature, involving both case control and cohort investigations. These studies (predominantly prospective cohort designs) included a total of 479,336 participants; averaging 17,753 subjects per study (range 428-73,265). There were a total of 12,361 reported cases of stroke (ranging per study from 32-2,863). The total length of study follow-up for the prospective cohort studies averaged 13.2 yr (ranging from 6-26 yr). The articles were published over a 14 yr period ranging from 1993 to 2007. These studies involved large samples of men and women from regions throughout the world including studies from the USA (11), UK (2), Iceland (1), Denmark (2), Norway (4), Netherlands (1), Finland (2), Japan (1), Australia (1) and Greece (1). Very few studies [69,70] examined non-Caucasian participants.

*We found strong evidence that physical activity was associated with a reduced risk for stroke. The level of evidence was consistent with a Level 3A classification.* We observed an average risk reduction of 31% across all studies (median = 29%). In comparison to cardiovascular disease, there was more variability in the risk reductions in stroke in the highest activity/fitness group. The quality of the investigations was also generally quite good with a mean (and median) Downs and Black score of 13 (range 11-15).

The risk reductions appear to be even greater in studies that assessed physical fitness directly. For instance, in data from the Aerobics Center Longitudinal Study [71] the high fitness group (estimated peak METs = 13.1) and the moderate fitness group (estimated peak METs 10.5) had significantly lower risks of stroke mortality (68 and 63%, respectively) than the least fit men (estimated peak METs 8.5).

A dose-response relationship did emerge when examining the literature. However, as illustrated by others this was extremely variable amongst studies and varied

**Table 13 Studies examining the relationship between physical activity and stroke.**

Publication Country Study Design Quality Score	Objective	Population	Methods	Outcome	Comments and Conclusions
Wisloff et al 2006 [58]	To assess exercise amount and intensity in relation to subsequent CVD mortality (including stroke).	• n = 27,143 men, 28,929 women	16 year follow up	Multivariate adjusted RR (95% CI) Men	Both high and low- intensity exercise may be associated with a reduced risk of stroke in both men and women.
Norway		• Sex: Men and women • Age: ≥ 20 yr • Characteristics: free from CVD • HUNT Study	PA Assessment: Questionnaire  PA  G1 = None G2 = <1/wk G3a = 1/wk ≤ 30 min low G3b = 1/wk ≤ 30 min high G3c = 1/wk > 30 min low G3d = 1/wk > 30 min high  G4a = 2-3/wk ≤ 30 min low G4b = 2-3/wk ≤ 30 min high G4c = 2-3/wk > 30 min low G4d = 2-3/wk > 30 min high G5a = ≥ 4/wk ≤ 30 min low G5b = ≥ 4/wk ≤ 30 min high G5c = ≥ 4wk > 30 min low G5d = ≥ 4/wk > 30 min high Outcome Measure: IHD mortality Cox proportional HR	G1 = 1.00 (referent) G2 = 0.90 (0.70-1.17) G3a = 0.90 (0.64-1.26) G3b = 0.59 (0.27-1.27) G3c = 0.62 (0.40-0.95) G3d = 0.51 (0.31-0.86) G4a = 0.72 (0.49-1.05) G4b = 0.63 (0.31-1.30) G4c = 1.02 (0.72-1.44)  G4d = 0.59 (0.37-0.92) G5a = 0.97 (0.70-1.36) G5b = 0.68 (0.27-1.66) G5c = 0.81 (0.65-1.20) G5d = 0.67 (0.49-1.11)  RR (95% CI) Women G1 = 1.00 (referent) G2 = 1.01 (0.81-1.25) G3a = 0.88 (0.68-1.15) G3b = 0.98 (0.46-2.10) G3c = 0.63 (0.42-0.94) G3d = 1.00 (0.50-1.98) G4a = 0.91 (0.70-1.17) G4b = 1.44 (0.78-2.65) G4c = 0.62 (0.44-0.88) G4d = 0.77 (0.36-1.66) G5a = 0.74 (0.56-0.99) G5b = 0.40 (0.10-1.62) G5c = 0.63 (0.45-0.89) G5d = 0.51 (0.21-1.26)	
Prospective cohort					
D & B score = 12					

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

Abbott et al 2003 [69]	To examine the way in which risk factor effects on the incidence of thromboembolic and hemorrhagic stroke can change over a broad range of ages.	• n = 7,589	6, 15 and 26 year follow up	Incidence rates per 1000 of stroke:	The protective effect of PA on reducing risk of stroke increased with age.
USA		• Sex: Men • Age: 45-93 yr	PA assessment: Using PA index over a 24 hour period PA information collected at study enrolment 1965-1968 and updated at physical examinations that occurred at 6, 15 and 26 years into follow-up.	• G1 = 9.0 (49) • G2 = 17.8 (124)	
Prospective cohort		• Characteristics: Free from CHD and stroke at enrolment; Japanese ancestry living on the island of Oahu, Hawaii.	Grouped into 4 age groups, yr:	• G3 = 33.4 (112)	
D & B score = 14		• Honolulu Heart Program	G1 = 45-54 G2 = 55-64  G3 = 65-74  G4 = 75-93 Outcome Measure: diagnosis of fatal and non fatal stroke during 26 years of follow-up Cox proportional HR	• G4 = 48.1 (111) Incidence of stroke event increased with advancing age p <0.001 There appeared to be a small protective effect within each age group. Inverse relations increased with age (p = 0.046). The protective effect of PA became significant in men >77 years (p = 0.032)	
Gillium et al 1996 [70]	To examine the relationship between recreational and non-recreational PA and risk of stroke.	• n = 2,368 men, 2,713 women	11.6 year follow up	Number of Cases: 249 white women, 270 white men, 104 black	Sedentary behaviour was found to be associated with increased risk of stroke.
USA		• Sex: Men and women • Age: 45-74 yr	PA assessment: Questionnaire divided into tertiles: T1 = Low T2 = Medium T3 = High	RR (95% CI) Black men and women Recreational PA • T1 = 1.33 (0.67-2.63)  • T2 = 1.33 (0.63-2.79) • T3 = 1.00 (referent)	
Prospective cohort		• Ethnicity: Black and white • NHANES I	Outcome Measure: Total Stroke Cox proportional HR	Non-recreational PA • T1 = 1.40 (0.90-2.16)	
D & B score = 12					

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

	<ul style="list-style-type: none"> <li>• T2 = 1.41 (0.74-2.70)</li> <li>• T3 = 1.00 (referent)</li> </ul>
RR (95% CI) White men age 45-64	
Recreational PA	<ul style="list-style-type: none"> <li>• T1 = 1.24 (0.63-2.41)</li> <li>• T2 = 1.17 (0.61-2.27)</li> <li>• T3 = 1.00 (referent)</li> </ul>
Non-recreational PA	<ul style="list-style-type: none"> <li>• T1 = 1.07 (0.40-2.86)</li> <li>• T2 = 1.75 (1.04-2.96)</li> <li>• T3 = 1.00 (referent)</li> </ul>
RR (95% CI) White women age 45-64	
Recreational PA	<ul style="list-style-type: none"> <li>• T1 = 3.13 (0.95-10.32)</li> <li>• T2 = 1.80 (0.52-6.22)</li> <li>• T3 = 1.00 (referent)</li> </ul>
Non-recreational PA	<ul style="list-style-type: none"> <li>• T1 = 3.51 (1.66-7.46)</li> <li>• T2 = 1.07 (0.57-1.99)</li> <li>• T3 = 1.00 (referent)</li> </ul>
RR (95% CI) White men age 65-74	
Recreational PA	<ul style="list-style-type: none"> <li>• T1 = 1.29 (0.58-1.88)</li> <li>• T2 = 0.86 (0.58-1.28)</li> <li>• T3 = 1.00 (referent)</li> </ul>
Non-recreational	<ul style="list-style-type: none"> <li>• T1 = 1.82 (1.15-2.88)</li> <li>• T2 = 1.20 (0.88-1.64)</li> <li>• T3 = 1.00 (referent)</li> </ul>
RR (95% CI) White women age 65-75	
Recreational PA	<ul style="list-style-type: none"> <li>• T1 = 1.55 (0.95-2.53)</li> <li>• T2 = 1.27 (0.76-2.12)</li> <li>• T3 = 1.00 (referent)</li> </ul>
Non-recreational PA	<ul style="list-style-type: none"> <li>• T1 = 1.82 (1.10-3.02)</li> <li>• T2 = 1.42 (1.01-2.00)</li> <li>• T3 = 1.00 (referent)</li> </ul>

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

Lee and Blair 2002 [71]	To examine the association between PF and stroke mortality in men.	<ul style="list-style-type: none"> <li>• n = 16,878</li> <li>• Sex: Men</li> <li>• Age: 40-87 yrs</li> <li>• Aerobics Center Longitudinal Study</li> </ul>	Baseline medical evaluation between 1971 and 1994 with average follow up period of 10 years	Average estimated maximal METs <ul style="list-style-type: none"> <li>• T1 = 8.5 MET</li> <li>• T2 = 10.5 MET</li> <li>• T3 = 13.1 MET</li> </ul>	Moderate and high levels of PF were associated with lower risk of stroke mortality in men.
USA					
Prospective cohort			PF assessment: Maximal exercise tolerance test, divided into tertiles	RR (95% CI) adjusted for age and exam year <ul style="list-style-type: none"> <li>• T1 = 1.00 (referent)</li> <li>• T2 = 0.35 (0.16-0.77)</li> <li>• T3 = 0.28 (0.11-0.71)</li> </ul>	
D & B score = 13			T1 = Low T2 = Moderate T3 = High Cox proportional HR	Trend p = 0.005	
Hu et al 2000 [72]	To examine the association between PA and risk of total stroke and stroke sub- types in women.	<ul style="list-style-type: none"> <li>• n = 72,488</li> <li>• Sex: Women</li> <li>• Age:40-65 yr</li> <li>• Characteristics: Nurses</li> <li>• Nurses' Health Study</li> </ul>	Baseline measurement in 1986 with follow-up questionnaire in 1988 and 1992	<ul style="list-style-type: none"> <li>• 407 cases of stroke (258 ischemic strokes, 67 subarachnoid hemorrhages, 42 intracerebral hemorrhages, and 40 strokes of unknown type)</li> </ul>	PA, including moderate-intensity exercise such as walking, is associated with a substantial reduction in risk of total and ischemic stroke in a dose- response manner.
USA					
Prospective cohort			PA assessment: Questionnaire for total PA (MET h/wk), divided into quintiles, walking activity (MET h/wk), divided into quintiles and walking pace	Multivariate RR (95% CI) for total stroke by total PA level <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.98</li> <li>• Q3 = 0.82</li> <li>• Q4 = 0.74</li> <li>• Q5 = 0.66</li> </ul>	
D & B score = 13			Total PA (MET h/wk)  Q1 = 0 - 2.0 Q2 = 2.1 - 4.6  Q3 = 4.7 - 10.4 Q4 = 10.5-21.7	p = 0.005  Multivariate RR (95% CI) for ischemic Stroke by total PA level	

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

	• Q1 = 1.00 (referent)
Q5 = > 21.7	
	• Q2 = 0.87
Walking activity (MET h/wk)	• Q3 = 0.83
Q1 = 0.5	• Q4 = 0.76
Q2 = 0.6 - 2.0	• Q5 = 0.52
Q3 = 2.1 - 3.8	p = 0.003
Q4 = 3.9 - 10	
Q5 = 10	Multivariate RR (95% CI) for total stroke by walking activity
Walking pace (mph)	• Q1 = 1.00 (referent)
G1 < 2.0	• Q2 = 0.76
G2 = 2-2.9	• Q3 = 0.78
G3 3.0	• Q4 = 0.70
	• Q5 = 0.66
Outcome measure: Stroke incidence	p = 0.01
	Multivariate RR (95% CI) for ischemic stroke by walking activity
Pooled logistic regression	
Cox proportional HR	• Q1 = 1.00 (referent)
	• Q2 = 0.77
	• Q3 = 0.75
	• Q4 = 0.69
	• Q5 = 0.60
	p = 0.02
	Multivariate RR (95% CI) for total stroke by usual Walking Pace
	• G1 = 1.00 (referent)
	• G2 = 0.81
	• G3 = 0.49
	p < 0.001
	Multivariate RR (95% CI) for ischemic stroke by usual walking pace
	• G1 = 1.00 (referent)
	• G2 = 0.71
	• G3 = 0.47
	p < 0.001

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

Lee et al 1999 [74]	To examine the association between exercise and stroke risk.	<ul style="list-style-type: none"> <li>• n = 21,823</li> <li>• Sex: Men</li> <li>• Age: 40-84 yr</li> </ul>	11.1 year follow up	Number of Cases: 533	VPA is associated with a decreased risk of stroke in men.
USA			PA assessment: Questionnaire for frequency of VPA, divided into 4 groups	Multivariate RR1 (95% CI) for total stroke by VPA	
Prospective cohort				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.79 (0.61-1.03)</li> </ul>	Inverse association with PA seemed to be mediated through beneficial effects on body weight, BP, cholesterol and glucose tolerance.
D & B score = 13			G1 < 1 time/week G2 = 1 time/week G3 = 2-4 times/week G4 ≥ 5 times/week	• G3 = 0.80 (0.65-0.99) • G4 = 0.79 (0.61-1.03) p = 0.04 RR2 (95% CI) for total stroke by VPA	
			RR1 = adjusted for smoking, alcohol consumption, history of angina and parental history of MI at <60 years	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.81 (0.61-1.07)</li> </ul>	
				<ul style="list-style-type: none"> <li>• G3 = 0.88 (0.70-1.10)</li> <li>• G4 = 0.86 (0.65-1.13)</li> </ul>	
				p = 0.25	
			RR2 = adjusted for all of the above plus, BMI, history of, hypertension, high cholesterol and diabetes	RR2 (95% CI) for ischemic stroke by VPA	
				VPA	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.90 (0.66-1.22)</li> <li>• G3 = 0.95 (0.74-1.22)</li> <li>• G4 = 0.97 (0.71-1.32)</li> </ul>	
			Outcome Measure: Total Stroke (Ischemic and Hemorrhagic)	p = 0.81	
				RR2 (95% CI) for hemorrhagic stroke by VPA	
			Cox proportional HR	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.54 (0.25-1.13)</li> <li>• G3 = 0.71 (0.41-1.23)</li> <li>• G4 = 0.54 (0.26-1.15)</li> </ul>	
				p = 0.10	



**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

Bijnen et al 1998 [166]	To describe the association between the PA patterns of elderly men and stroke mortality.	<ul style="list-style-type: none"> <li>• n = 802</li> </ul>	10 year follow up	Number of Cases: 47	No significant finding
Denmark		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: 64-84 yr</li> <li>• Characteristics: Not all free from previous stroke</li> </ul>	PA assessment: Questionnaire for LTPA, divided into tertiles	Multivariate adjusted RR (95% CI) <ul style="list-style-type: none"> <li>• T1 = 1.00 (referent)</li> <li>• T2 = 0.65 (0.33-1.25)</li> <li>• T3 = 0.55 (0.24-1.26)</li> </ul>	
Prospective cohort			T1 = Lowest		
			T2	$p = 0.12$	
			T3 = Highest		
D & B score = 15			Outcome Measure: Stroke Mortality Cox proportional HR		
Schnohr et al 2006 [214]	To describe the association between different levels of LTPA and subsequent causes of death (stroke).	<ul style="list-style-type: none"> <li>• n = 2136 men, 2,758 women</li> </ul>	5 year follow up	RR (95% CI), univariate	Although RR for of death from stroke was below 1 for both moderate and high compared with low PA, this association did not reach the level of statistical significance.
Copenhagen		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: 20 – 79 yr</li> <li>• Characteristics: Healthy, PA level did not change between 2 examinations, 5 years apart</li> </ul>	PA assessment: Questionnaire for LTPA, divided into 3 groups	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.64 (0.39-1.05)</li> <li>• G3 = 0.70 (0.41-1.21)</li> </ul> Trend $p = 0.4$	
Prospective cohort			G1 = Low PA (<4 METS)		
			G2 = Moderate PA (4-6 METS)	RR (95% CI), multivariate:	
			G3 = High PA (>6 METS)	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.67 (0.40-1.12)</li> <li>• G3 = 0.76 (0.43-1.34)</li> </ul>	
D & B score = 13		<ul style="list-style-type: none"> <li>• Copenhagen City Heart Study</li> </ul>	Multivariate Analysis Kaplan-Meier Plots	Trend $p = 0.6$	
			Linear, Logistical and Cox Regression.		
Vatten et al 2006 [253]	To investigate whether obesity-related CV mortality could be modified by PA.	<ul style="list-style-type: none"> <li>• n = 26,515 men, 27,769 women</li> </ul>	16 year follow up	Number of Cases: 994 women, 771 men	Lower levels of TPA are associated with an increased risk of stroke.

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

		<ul style="list-style-type: none"><li>• Sex: Men and women</li></ul>	PA assessment: Questionnaire for total amount of PA, divided into 4 groups		
Norway		<ul style="list-style-type: none"><li>• Age: 20 yr</li><li>• Characteristics: Free from CVD at baseline</li></ul>		Multivariate HR (95% CI), men <ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li></ul>	
Prospective cohort				<ul style="list-style-type: none"><li>• Q2 = 1.05 (0.85-1.30)</li></ul>	
D & B score = 14		<ul style="list-style-type: none"><li>• HUNT study</li></ul>	G1 = High G2 = medium G3 = low G4 = never	<ul style="list-style-type: none"><li>• Q3 = 1.21 (0.95-1.54)</li><li>• Q4 = 1.35 (1.05-1.74)</li></ul> <p>p = 0.009</p>	
			Outcome Measure: Stroke mortality	Multivariate HR (95% CI), women <ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li><li>• Q2 = 1.16 (0.93-1.45)</li><li>• Q3 = 1.45 (1.14-1.86)</li></ul>	
			Cox proportional HR	<ul style="list-style-type: none"><li>• Q4 = 1.45 (1.14-1.83)</li></ul> <p>p &lt; 0.001</p>	
Agnarsson et al 1999 [255]	To examine the association of LTPA and pulmonary function with the risk of stroke.	<ul style="list-style-type: none"><li>• n = 4,484</li><li>• Sex: Men</li><li>• Age: 45-80</li></ul>	Length of Follow-up: 10.6 ± 3.6 years	Number of Cases: 249	Apparent protective effect of regular continued LTPA in middle age men on the risk of ischemic stroke.
Iceland		<ul style="list-style-type: none"><li>• Characteristics: no history of Stroke</li></ul>	PA assessment: Questionnaire for LTPA (h/wk) and type of activity (intensity), each divided into 3 groups	Adjusted for age and smoking RR (95% CI) for total stroke by LTPA level	
Prospective cohort		<ul style="list-style-type: none"><li>• Reykjavik Study</li></ul>		<ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.84 (0.63-1.13)</li><li>• G3 = 0.73 (0.40-1.35)</li></ul>	
D & B score = 13			LTPA summer/winter G1 = none  G2 = ≤ 5 h/wk G3 = ≥ 6 h/wk	Adjusted for age and smoking RR (95% CI) for ischemic stroke by LTPA level	
			Type of Activity	<ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.72 (0.51-1.01)</li></ul>	

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

			G1 = none		
			G2 = low intensity	• G3 = 0.78 (0.41-1.48)	
			G3 = high Intensity		
			Outcome Measure: Total and ischemic Stroke	RR (95% CI) for total stroke by type of activity	
				• G1 = 1.0,0 (referent)	
				• G2 = 0.75 (0.53-1.08)	
				• G3 = 1.10 (0.78-1.57)	
			Cox proportional HR	RR (95% CI) for ischemic stroke by type of activity	
				• G1 = 1.00 (referent)	
				• G2 = 0.72 (0.44-1.07)	
				• G3 = 0.96 (0.64-1.44)	
Ellekjaer et al 2000 [256]	To examine the association between different levels of LTPA and stroke mortality in middle-aged and elderly women.	• n = 14,101	Baseline 1984-1986: 2 self administered questionnaires and clinical measurements included in the screening program.	Number of cases: 457	This study demonstrates a consistent, negative association between PA and stroke mortality in women.
Norway		• Sex: Women • Age: 50 yr		Multivariate RR (95% CI), all age groups	
Prospective cohort		• Characteristics: free from stroke at baseline		• G1 = 1.00 (referent) • G2 = 0.77	
			PA assessment: Questionnaire for LTPA, divided into 3 groups	• G3 = 0.52	
D & B score = 14				Multivariate RR (95% CI), age 50–69 years	
			G1 = low		The most active women had approx. 50% lower risk of death from stroke compare to inactive women.
			G2 = medium	• G1 = 1.00 (referent)	
			G3 = high	• G2 = 0.57 • G3 = 0.42	
			Outcome Measure: Death from stroke	p = 0.0021	
				Multivariate RR (95% CI), age 70-79 years	

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

				Cox proportional HR	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.79</li> <li>• G3 = 0.56</li> </ul> <p><math>p = 0.0093</math></p> <p>Multivariate RR (95% CI), age 80-101 years</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.91</li> <li>• G3 = 0.57</li> </ul> <p><math>p = 0.1089</math></p>	
Evenson et al 1999 [257]	To examine the relationship between PA and ischemic stroke risk.	<ul style="list-style-type: none"> <li>• n = 14,575</li> <li>• Sex: Men and women</li> <li>• Age: 45-64 yr</li> <li>• Atherosclerosis Risk in Communities Study</li> </ul>	7.2 year follow up	Number of Cases: 189  Number of Dropouts: 0%  Sport, Incidence of Ischemic Stroke  Multivariate adjusted RR (95% CI) by sport  <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q3= 0.83 (0.52-1.32)</li> </ul>	PA was weakly associated with a reduced risk of ischemic stroke among middle aged adults.
USA					
Prospective cohort					
D & B score = 14				Outcome Measure: Ischemic Stroke  Multivariate Poisson and Cox proportional HR	
				Multivariate adjusted RR (95% CI) by LTPA <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 =</li> <li>• Q3 = 0.89 (0.57-1.37)</li> </ul> Multivariate adjusted RR (95% CI) by OPA <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 =</li> <li>• Q3 = 0.69 (0.47-1.00)</li> </ul>	
Haheim et al 1993 [258]	To determine the risk factors of stroke incidence and mortality.	<ul style="list-style-type: none"> <li>• n = 14,403</li> <li>• Sex: Men</li> <li>• Age: 40-49 yr</li> </ul>	Baseline Screening from May 1972-December 1973.	HR (95% CI) for stroke incidence	Increased LTPA is associated with a reduced risk of stroke incidence but not mortality.
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.64 (0.38-1.08)</li> </ul>	

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

Norway			PA assessment: Questionnaire for LTPA, divided into groups	• G3 = 0.36 (0.15-0.80)	
Prospective cohort				HR (95% CI) for stroke mortality	
			G1 = Sedentary	• G1 = 1.00, (referent)	
			G2 = Moderate	• G2 = 0.82 (0.33-2.35)	
			G3 = Intermediate or Great	• G3 = 0.29 (0.03-1.51)	
D & B score = 14			Outcome Measure: Incidence of stroke morbidity and mortality until study end date, December 31, 1984. Cox proportional HR		
Hu et al 2005 [259]	To assess the relationship of different types of PA with total and type-specific stroke risk.	• n = 47,721  • Sex: Men and women	PA assessment: Mailed questionnaire for LTPA, OPA and commuting PA, divided into groups as follows:	RR (95% CI) by LTPA, men  • G1 = 1.00 (referent) • G2 = 0.83 • G3 = 0.72 $p < 0.001$	A high level of LTPA reduces the risk of all subtypes of stroke. Daily active commuting also reduces the risk of ischemic stroke.
Finland		• Age: 25-64 • Characteristics: Healthy at baseline			
Prospective cohort			LTPA levels: G1 = Low G2 = Moderate G3 = High	RR (95% CI) by LTPA, women • G1 = 1.00 (referent) • G2 = 0.86 • G3 = 0.75 $p = 0.007$	
D & B score = 13			OPA: G1 = Light  G2 = Moderate G3 = Hard	RR (95% CI) by LTPA, men and women  • G1 = 1.00 (referent) • G2 = 0.85  • G3 = 0.73 $p < 0.001$	
			Commuting PA: G1 = Motorized or no work, G2 = walking or cycling 1-29 min G3 = walking or cycling $\geq 30$ min.	RR (95% CI) by OPA, men • Not significant	

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

			Outcome Measure: Incidence of fatal or non-fatal stroke occurring during follow-up until end of 2003. Mean follow-up of 19 years.		
			Cox proportional hazard	RR (95% CI) by OPA, women <ul style="list-style-type: none"><li>• Not significant</li></ul> RR (95% CI) by OPA, men and women <ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.90</li><li>• G3 = 0.87</li></ul> $p = 0.007$ RR (95% CI) by commuting PA, men <ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.91</li><li>• G3 = 0.85</li></ul> $p = 0.047$ RR (95% CI) by commuting PA, women <ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.86</li><li>• G3 = 0.85</li></ul> $p = 0.018$ RR (95% CI) by commuting PA, men and women <ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.89</li><li>• G3 = 0.85</li></ul> $p = 0.002$	
Kiely et al 1994 [260]	To examine the influence of increased PA on stroke risk in members of the Framingham study cohort.	<ul style="list-style-type: none"><li>• n = 1,897 men 2,299 women</li><li>• Sex: Men and women</li><li>• Age: 28-62 yr</li><li>• Characteristics: Free from stroke</li></ul>	Baseline measurement in 1954-1955 and follow up in either 1968-1969 or 1971- 1972	Multivariate adjusted RR (95% CI) at first examination, men (mean age 50 years)	Medium and high levels of PA among men are protective against stroke relative to low levels.
USA				<ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.90 (0.62-1.31) <math>p = 0.59</math></li><li>• G3 = 0.84 (0.59-1.18) <math>p = 0.31</math></li></ul>	
Prospective cohort			PA assessment: Questionnaire for metabolic work done during a typical 24 hr period, divided into 3 groups		

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

D & B score = 12			<p>Multivariate adjusted RR (95% CI) at first examination, women (mean age 50 years)</p> <p>Protective effect of PA was slightly less for high levels of PA compared to medium levels for older men.</p>		
			<p>G1 = Low</p> <p>G2 = Medium</p> <p>G3 = High</p> <p>Outcome Measure: Incidence of stroke, as defined by the first occurrence of atherothrombotic brain infarctions, cerebral embolism or other type of stroke, during 32 years of follow-up.</p>		
			<p>• G1 = 1.00 (referent)</p> <p>• G2 = 1.21 (0.89-1.63) <math>p = 0.23</math></p> <p>• G3 = 0.89 (0.60-1.31) <math>p = 0.54</math></p>		
			<p>Multivariate adjusted RR (95% CI) at second examination, men (mean age 63 years)</p> <p>• G1 = 1.00 (referent)</p> <p>• G2 = 0.41 (0.24-0.89) <math>p = 0.0007</math></p> <p>• G3 = 0.53 (0.34-0.84) <math>p = 0.007</math></p>		
			<p>Multivariate adjusted RR (95% CI) at second examination, women (mean age 64 years)</p>		
			<p>Cox proportional HR</p> <p>• G1 = 1.00 (referent)</p> <p>• G2 = 0.97 (0.64-1.47) <math>p = 0.67</math></p> <p>• G3 = 1.21 (0.75-1.96) <math>p = 0.43</math></p>		
Krurup et al 2007 [261]	To compare the reported level of PA performed during the week preceding an ischemic stroke with that of community controls.	<p>• n = 127 cases 301 controls</p> <p>• Sex: Men and women</p> <p>• Age: <math>\geq 40</math> yr</p>	<p>PA assessment:</p> <p>Questionnaire about PA 1 week prior to stroke (cases) and 1 week prior to questionnaire (controls), divided into PASE scores and quartiles</p>	<p>Univariate OR (95% CI)</p> <p>PASE Score</p> <p>• Q1 = 1.00 (referent)</p> <p>• Q2 = 0.51 (0.28-0.95)</p> <p>• Q3 = 0.27 (0.14-0.54)</p>	<p>Stroke patients are less physically active in the week preceding an ischemic stroke when compared to age and sex-matched controls. Increasing PASE score was inversely, log-linearly and significantly associated with OR for ischemic stroke.</p>

Denmark

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

Case control		<ul style="list-style-type: none"> <li>Characteristics: Case: Stroke Patients (20% had history of Stroke), Controls: 4% had history of stroke</li> </ul>		<ul style="list-style-type: none"> <li>Q4 = 0.08 (0.03-0.20)</li> </ul>	
D & B score = 14			Q1 = 0-49 Q2 = 50-99 Q3 = 100-149 Q4 = 150+	Multivariate OR (95% CI) PASE Score <ul style="list-style-type: none"> <li>Q1 = 1.00 (referent)</li> <li>Q2 = 0.53 (0.26-1.08)</li> <li>Q3 = 0.27 (0.12-0.59)</li> </ul>	
			Outcome measure: Ischemic stroke Chi squared Kruskal-Wallis Statistics Multivariate conditional logistic regression	<ul style="list-style-type: none"> <li>Q4 = 0.09 (0.03-0.25)</li> </ul>	
Kurl et al 2003 [262]	To examine the relationship of PF with subsequent incidence of stroke. Also to compare PF with conventional risk factors as a predictor for future stroke.	<ul style="list-style-type: none"> <li>n = 2,011</li> </ul>	Baseline examinations conducted between March 1984 and December 1989 with average follow up period of 11 years	Multivariate HR (95% CI), any stroke	Low PF was associated with an increased risk of any stroke and ischemic stroke.
Finland		<ul style="list-style-type: none"> <li>Sex: Men</li> <li>Age: 42, 48, 54 or 60 yrs</li> </ul>		<ul style="list-style-type: none"> <li>Q1 = 1.00 (referent)</li> <li>Q2 = 1.39 (0.70-2.77)</li> <li>Q3 = 1.32 (0.66-2.65)</li> <li>Q4 = 2.30 (1.18-4.06)</li> </ul>	
Prospective cohort		<ul style="list-style-type: none"> <li>Characteristics: Free from stroke or pulmonary disease</li> <li>Kuopio Ischaemic Heart Disease Risk Factor Study</li> </ul>		Trend $p = 0.01$	
			PF assessment: Maximal exercise test on cycle ergometer. $\text{VO}_2$ max (ml/kg/min) divided into quartiles	Multivariate HR (95% CI), ischemic stroke	
D & B score = 14				<ul style="list-style-type: none"> <li>Q1 = 1.00 (referent)</li> <li>Q2 = 1.28 (0.56-2.94)</li> <li>Q3 = 1.64 (0.74-3.65)</li> </ul>	
			Q1 = >35.3	<ul style="list-style-type: none"> <li>Q4 = 2.40 (1.09-5.25)</li> </ul>	
			Q2 = 30.3-35.3		
			Q3 = 25.2-30.2	Trend $p = 0.01$	



**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

			Q4 = <25.2		
			Outcome Measure: Stroke incidence		
			Cox proportional HR		
Myint et al 2006 [263]	To examine the association between a combination of OPA and LTPA with risk of subsequent stroke.	• n = 22,602	Baseline measurement in	Model A: Used all 4 categories of PA	Higher levels of PA assessed using a single simple pragmatic tool based on both OPA and LTPA is associated with reduced stroke risk.
UK		• Sex: Men • Age: 40-79 yr • Characteristics: Healthy at baseline	1993-1997	HR (95% CI), men and women • G1 = 1.00 (referent) • G2 = 0.78 (0.61-1.00)	
Prospective cohort		• European Prospective Investigation in Cancer-Norfolk	PA assessment: Questionnaire for PA (includes LTPA and OPA) divided into 4 groups	• G3 = 0.66 (0.49-0.91) • G4 = 0.70 (0.49-0.99)	
D & B score = 11			G1 = Inactive G2 = moderately inactive G3 = moderately active G4 = active	$p = 0.024$ HR (95% CI), men • G1 = 1.00 (referent) • G2 = 0.75 (0.52-1.09)  • G3 = 0.55 (0.35-0.86) • G4 = 0.67 (0.43-1.05)	
			Outcome Measure: Incidence of fatal and non fatal stroke.	$p = 0.41$ Women not significant $p = 0.50$	
			Cox proportional HR	Model B: Used 3 categories of PA (G3 and G4 combined combined) HR (95% CI), men and women • G1 = 1.00 (referent) • G2 = 0.78 (0.61-1.00) • G3 = 0.68 (0.52-0.88) $p = 0.009$ HR (95% CI), men • G1 = 1.00 (referent) • G2 = 0.75 (0.52-1.09), • G3 = 0.61 (0.43-0.86) $p = 0.019$	

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

				Women not significant p = 0.34	
Noda et al 2005 [264]	To examine the impact of exercise on CVD (stroke) mortality in Asian populations.	<ul style="list-style-type: none"> <li>• n = 31,023 men, 42,242 women</li> <li>• Sex: Men and women</li> <li>• Age: 40 -79 yr</li> <li>• Ethnicity: Asian</li> </ul>	9.7 year follow up	Number of Cases: 186 men, 141 women  Number of Dropouts: 3.4%	PA through walking and sports participation may reduce the risk of mortality from ischemic stroke
Japan			PA assessment: Questionnaire for PA (walking and sports participation (h/ day), divided into quartiles:		
Prospective cohort				Multivariate adjusted HR (95% CI) by duration of walking PA, men	
				<ul style="list-style-type: none"> <li>• Q1 = 1.03 (0.63-1.69)</li> <li>• Q2 = 1.00 (referent)</li> <li>• Q3 = 0.56 (0.35-0.91)</li> <li>• Q4 = 0.71 (0.49-1.02)</li> </ul>	
D & B score = 13			Q1 = <0.5 Q2 = 0.5 Q3 = 0.6-0.9 Q4 = >1.0		
			Outcome Measure: Death from ischemic stroke	Multivariate adjusted HR (95% CI) by duration of walking PA, women	
				<ul style="list-style-type: none"> <li>• Q1 = 1.38 (0.82-2.33)</li> <li>• Q2 = 1.00 (referent)</li> </ul>	
			Cox proportional HR	<ul style="list-style-type: none"> <li>• Q3 = 0.56 (0.32-0.97)</li> <li>• Q4 = 0.73 (0.48-1.13)</li> </ul>	
				Multivariate adjusted HR (95% CI) by sport PA, men	
				<ul style="list-style-type: none"> <li>• Q1 = 1.34 (0.86-2.08)</li> <li>• Q2 = 1.00 (referent)</li> <li>• Q3 = 1.22 (0.66-2.25)</li> <li>• Q4 = 0.84 (0.45-1.57)</li> </ul>	
				Multivariate adjusted HR (95% CI) by sport PA, women	
				<ul style="list-style-type: none"> <li>• Q1 = 1.07 (0.64-1.77)</li> <li>• Q2 = 1.00 (referent)</li> <li>• Q3 = 0.62 (0.25-1.58)</li> <li>• Q4 = 0.73 (0.31-1.70)</li> </ul>	

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

Paganini-Hill and Barreto 2001 [265]	To identify risk factors and preventative measures for stroke in elderly men and women.	<ul style="list-style-type: none"> <li>• n = 4,722 men, 8,532 women</li> <li>• Sex: Men and women</li> </ul>	Baseline survey in 1981- 1982.	Multivariate adjusted RR (95% CI) for total hemorrhagic occlusion by exercise, men	Emphasized role of lifestyle modification in the primary prevention of stroke.
USA		<ul style="list-style-type: none"> <li>Age: 44-101 yr</li> <li>• Characteristics: no previous history of stroke. Residence of a retirement community in Southern California</li> </ul>	PA assessment: Questionnaire on amount of hours per day of exercise	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.88</li> <li>Q3 = 0.83</li> </ul>	
Prospective cohort			G1 = <0.5		
			G2 = <0.1	Multivariate adjusted RR (95% CI) for total hemorrhagic occlusion by exercise, women	
			G3 = 1+		
D & B score = 13			Outcome Measure: Incidence of hemorrhagic occlusion strokes up until December 31, 1998.	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.91</li> <li>• Q3 = 0.85</li> </ul>	
			Poisson Regression 40 year follow up		
Pitsavos et al 2004 [266]	To investigate the interaction between PA in men with LVH on stroke mortality.	<ul style="list-style-type: none"> <li>• n = 489</li> <li>• Sex: Men</li> </ul>		Number of cases: 67	PA reduced the risk of stroke in men without LVH.
USA		<ul style="list-style-type: none"> <li>• Age: 40-59 yr</li> <li>• Characteristics: Those without LVH</li> </ul>	PA assessment: Questionnaire	RR (95% CI)	
			G1 = Sedentary	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.64 (0.45-0.91)</li> </ul>	
Prospective cohort			G2 = Moderate	<ul style="list-style-type: none"> <li>• G3 = 0.72 (0.51-1.02)</li> </ul>	
		<ul style="list-style-type: none"> <li>• Corfu Cohort (Greece) from Seven Countries Study</li> </ul>	G3 = Hard		
D & B score = 12			Outcome Measure: Stroke mortality Cox proportional HR		

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

Sacco et al 1998 [267]	To investigate the association between LTPA and ischemic stroke.	<ul style="list-style-type: none"> <li>• n = 369 case, 678 control</li> </ul>	Case Subjects were recruited during hospitalization, self referral or from monitoring non hospitalized stroke. Controls were eligible if they had never been diagnosed with stroke and were >39 years.		LTPA was related to a decreased occurrence of ischemic stroke in elderly, multiethnic, urban subjects.
USA		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>		O R (95% CI) for duration of LTPA and stroke	
Case control		<ul style="list-style-type: none"> <li>• Age: &gt; 39 yr</li> <li>• Characteristics: Case Subjects: Diagnosed with first cerebral infarction after July 1, 1993. Control Subjects: Never diagnosed with stroke</li> </ul>		<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.42</li> </ul>	
D & B score = 14				<ul style="list-style-type: none"> <li>• G3 = 0.35</li> <li>• G4 = 0.31</li> </ul>	
			PA assessment: Questionnaire Divided into duration of LTPA (h/wk)		
		<ul style="list-style-type: none"> <li>• Northern Manhattan Stroke Study</li> </ul>	G1 = 0 G2 = <2 G3 = 2-<5 G4 = ≥ 5 Multivariate conditional logistic regression Baseline data collection from 1982-1983 in East Boston (MA), New Haven (CT) and Iowa and Washington counties (IA).		
Simonsick et al 1993 [268]	To examine the association between recreational PA among physically capable older adults and incidence of selected chronic diseases and mortality over 3 and 6 years.	<ul style="list-style-type: none"> <li>• n = 1,815</li> </ul>		After 3 years Iowa	No consistent relationship between PA and stroke was found after 3 or 6 years across all 3 population cohorts.
		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: ≥ 65 yrs</li> </ul>		OR (95% CI) Stroke and activity level	

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

USA	<ul style="list-style-type: none"> <li>• Characteristics: Physically capable to do heavy work around the house, walk up and down a flight of stairs and walk a half mile without help.</li> </ul>	<ul style="list-style-type: none"> <li>• T1 = 0.22 (0.08-0.61)</li> </ul>
Prospective cohort		<ul style="list-style-type: none"> <li>• T2 = 1.05 (0.60-1.84)</li> <li>• T3 = 1.00 (Referent)</li> </ul>
D & B score = 12	<p>PA assessment: Questionnaire</p> <p>T1 = High T2 = Moderate and T3 = Inactive</p> <p>• Established Populations for Epidemiologic Studies of the Elderly</p> <p>Outcome Measure: Stroke incidence during 3 and 6 year follow-ups.</p> <p>Logistic Regression</p>	<p>New Haven</p> <p>OR (95% CI) Stroke and activity level</p> <ul style="list-style-type: none"> <li>• T1 = 1.06 (0.38-2.95)</li> <li>• T2 = 1.26 (0.54-2.92)</li> <li>• T3 = 1.00 (Referent)</li> </ul> <p>East Boston</p> <p>OR (95% CI) Stroke and activity level</p> <ul style="list-style-type: none"> <li>• T1 = 0.59 (0.17-1.95)</li> <li>• T2 = 1.08 (0.52-2.27)</li> <li>• T3 = 1.00 (Referent)</li> </ul> <p>After 6 years</p> <p>Iowa</p> <p>OR (95% CI) Stroke and activity level</p> <ul style="list-style-type: none"> <li>• T1 = 0.56 (0.31-1.00)</li> <li>• T2 = 0.97 (0.64-1.48)</li> <li>• T3 = 1.00 (Referent)</li> </ul> <p>New Haven</p> <p>OR (95% CI) Stroke and activity level</p> <ul style="list-style-type: none"> <li>• T1 = 1.05 (0.52-2.12)</li> <li>• T2 = 1.29 (0.72-2.32)</li> <li>• T3 = 1.00 (Referent)</li> </ul> <p>East Boston</p> <p>OR (95% CI) Stroke and activity level</p> <ul style="list-style-type: none"> <li>• T1 = 1.21 (0.56-2.61)</li> </ul>

**Table 13: Studies examining the relationship between physical activity and stroke. (Continued)**

				<ul style="list-style-type: none"> <li>• T2 = 1.73 (0.98-3.06)</li> <li>• T3 = 1.00 (Referent)</li> </ul>	
Thrift et al 2002 [269]	To examine whether intracerebral hemorrhage is associated with dynamic or static exercise.	<ul style="list-style-type: none"> <li>• n = 662</li> <li>• Sex: Men and women</li> <li>• Age: 18-80 yr</li> </ul>	PA assessment: Interview, divided into 3 groups: frequency of vigorous activity	Number of Cases: 331	Findings not significant after multivariate analysis.
Australia		<ul style="list-style-type: none"> <li>• Characteristics: Cases: first episode of intracerebral hemorrhage Controls: Neighbours of cases</li> </ul>		Multivariate OR (95% CI) by frequency of VPA	
Case control			G1 = Never G2 = Rarely G3 = Once or more per month	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.68 (0.36-1.27)</li> <li>• G3 = 0.66 (0.39-1.11)</li> </ul>	
D & B score = 14			OPA level	p = 0.094	
			G1 = Sedentary G2 = Light to moderate G3 = Heavy	Multivariate OR (95% CI) by OPA level	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.94 (0.59-1.48), p = 0.773</li> <li>• G3 = 1.18 (0.57-2.46), p = 0.650</li> </ul>	
			Outcome Measure: Intracerebral hemorrhage		
			Multiple logistic regression		

D & B score, Downs and Black quality score; YR, years; wk, week; CVD, cardiovascular disease; G, groups; PA, physical activity; CHD, coronary heart disease; RR, risk ratio; 95% CI, 95% confidence interval; T, tertile; PF, physical fitness; MET, metabolic equivalent; Q, quartile or quintile; OPA, occupational physical activity; LTPA, leisure-time physical activity; HR, hazard ratio; VPA, vigorous physical activity; LVH, left ventricular hypertrophy.

according to the type of stroke (ischemic or haemorrhagic) [52]. For instance, 12 studies (46%) revealed a dose-response relationship in one or more measures of occupational and/or leisure-time physical activity and the risk for stroke. It is difficult to determine the minimal and optimal physical activity dosage for the prevention of stroke. Brisk walking has been associated with a lower risk of total and ischemic stroke [72]. In the Harvard Alumni study, the risk of stroke was lower at a weekly energy expenditure of 4.2-8.4 MJ/wk (1000-1999 kcal/wk) (RR = 0.76 (95% CI, 0.59 to 0.98)). With expenditures of 8.4-12.6 MJ/wk (2000-2999 kcal/wk) the RR dropped to 0.54 (0.38 to 0.76) [73]. Thus, the recommended daily expenditure of Canada's physical activity guidelines is sufficient to reduce the risk for stroke. Further research is required to clearly determine the risk reductions at exercise volumes less than 4.2 MJ/wk (1000 kcal/wk).

In summary, the results of these studies (taken as a whole) indicate that occupation- and leisure time-related physical activity are inversely related to the risk for stroke. Both physically active men and women have a lower risk of stroke, and it appears that this benefit may be present for both ischemic and haemorrhagic stroke [74]. The relationship between physical activity and stroke appears to be consistent between men and women. Unfortunately, relatively limited data exists in non-Caucasian populations.

**Recommendation #3**


*For a reduced risk of stroke, it is recommended that individuals should participate in 30 min or more of moderate to vigorous exercise on most days of the week. Brisk walking appears to be protective against the development of stroke. It remains to be determined whether lower volumes of physical activity lead to a reduced risk for stroke. [Level 3, Grade A]*

**Primary Prevention of Hypertension**

A total of 6287 citations were identified during the electronic database search (Figure 6). Of these citations, 4054 were identified in MEDLINE, 1360 in EMBASE, 253 in Cochrane, and 620 in the CINAHL/SportDiscus/PsychInfo search. A total of 40 duplicates were found, leaving a total of 6247 unique citations. A total of 6167 articles were excluded after scanning, leaving a total of 80 articles for full review. An additional five articles were found through cross-referencing and the reviewers' personal files. From these articles 72 were excluded after full review for the following reasons: weak design (n = 4), did not contain three levels of physical activity or not possible to determine dose-response relationship (n = 19), reviews, summaries, meta-analyses (n = 8), not dealing with hypertension (n = 2), only reported on changes in blood pressure (n = 27), clinical population

(n = 7), and other (n = 6). Therefore, a total of 12 articles were included in the systematic review of the literature regarding the relationship between physical activity and the primary prevention of hypertension. The majority of the literature examining the dose-response (for at least three levels of physical activity/fitness) involved prospective cohort analyses (83%).

As shown in Table 14, 12 investigations examined the dose-response (i.e., three or more levels) relationship between physical activity and the incidence of hypertension. This involved a total of 112,636 participants, averaging 10,240 subjects per study (range 1,243-41,837).



**Table 14 Studies examining the relationship between physical activity and hypertension.**

Publication Country Study Design Quality Score	Objective	Population	Methods	Outcome	Comments and Conclusions
Rankinen et al 2007 [75]	To investigate the contributions of DNA sequence variation in candidate genes, PF and BMI, as well as their interactions to the incidence of hypertension.	• n = 629 cases; 605 controls	10 year follow up	PF showed the strongest association with HTN risk among all subjects as well as sex-specific models. Each 1-MET increment in PF was associated with 19% (12- 14%), 16% (9-22%), 32% (17- 45%) risk reduction in all subjects, men and women respectively.	PF is a significant predictor of the risk of hypertension.
USA		• Sex: Men and women	All subjects required to have 2 clinic visits at least 2 years apart.		
Case control		• Age: Case: 43.3 (9.2) yr Control: 42.7 (8.9) yr	PF assessment: treadmill test (Blake protocol)		
D & B score = 13		• Characteristics: Healthy with BP 134/86 mmHg or less at their first clinic visit. Cases: those who developed hypertension during the follow-up period. Controls were those who did not develop hypertension	Outcome measure: Incidence of hypertension during follow-up. Incident cases of hypertension were defined as physician diagnosed hypertension with medication or SBP $\geq$ 140 mmHg and/of DBP $\geq$ 90 mmHg t-tests and chi-square tests Logistic regression modelling	When divided into quartiles on the basis of sex specific MET cut-offs, the third and fourth quartiles had a 58% (41-71%) and 63% (47-75%) lower risk of hypertension compared to the 1 <sup>st</sup> quartile.	
Pereira et al 1999 [76]	To examine PA and incident hypertension in men and women.	• n = 7,459	PA Assessment: Questionnaire for leisure, sport and work index, divided into quartiles	White Men	There is an inverse association between PA and incident hypertension in White middle aged men. White men in the highest quartiles of sport and leisure activity had statistically significant reductions in the odds of developing hypertension of 23 and 34% respectively, compared to men in the lower quartiles.
USA		• Sex: Men and women	Q1 = Lowest	Leisure Index Model 1	
Prospective cohort		• Age: 45-65 yr	Q2	• Q1 = 1.00 (referent)	
D & B score = 12		• Characteristics: No history of angina, MI, evidence of MI, angioplasty or other CV surgery or hypertension	Q3	• Q2 = 0.95 (0.70-1.28)	
		• Atherosclerosis Risk in Communities Study	Q4 = Highest	• Q3 = 0.83 (0.63-1.09)	
			Model 1 adjusted for: Age, education, baseline BP and study centre	• Q4 = 0.64 (0.46-0.89)	



**Table 14: Studies examining the relationship between physical activity and hypertension. (Continued)**

				Model 2 adjusted for: Covariates in model 1 and smoking, alcohol consumption, parental history of hypertension, energy, sodium, potassium and caffeine intake, BMI, waist to hip ratio, menopausal status and hormone use	Trend $p = 0.01$	
				Outcome Measure: Incidence of hypertension as defined as a SBP 140 mmHg and/or a DBP 90 mmHg or use of antihypertensive medications.	Leisure Index Model 2	
				Unconditional logistic regression Orthogonal polynomial coefficients	• Q1 = 1.00 (referent)	
					• Q2 = 0.99 (0.72-1.35)	
					• Q3 = 0.86 (0.65-1.13)	
					• Q4 = 0.66 (0.47-0.94)	
					Trend $p = 0.01$	
					Women	
					Sport Index Model 1	
					• Q1 = 1.00 (referent)	
					• Q2 = 1.26 (0.78-2.05)	
					• Q3 = 1.06 (0.61-1.84)	
					• Q4 = 1.92 (1.12-3.29)	
					Trend $p = 0.04$	
					Men	
					Sport Index Model 1	
					• Q1 = 1.00 (referent)	
					• Q2 = 1.23 (0.91-1.66)	
					• Q3 = 0.92 (0.70-1.22)	
					• Q4 = 0.74 (0.54-1.02)	
					Trend $p = 0.02$	
					Sport Index Model 2	
					• Q1 = 1.00 (referent)	
					• Q2 = 1.26 (0.93-1.71)	
					• Q3 = 0.95 (0.71-1.26)	
					• Q4 = 0.77 (0.55-1.08)	
					Trend $p = 0.05$	
Haapanen et al 1997 [77]	To assess the association between PA and hypertension.	• n = 732 men; 796 women	10 year follow up (1980 baseline)	Age adjusted incidence rates of hypertension Total energy expenditure High as referent:	Increased EE during LTPA and increased intensity of these activities were associated with reduced risk for incident hypertension (age adjusted) in men but not women.	

**Table 14: Studies examining the relationship between physical activity and hypertension. (Continued)**

Finland		• Sex: Men and women	PA assesment: Questionnaire for EE (kcal/wk), divided into tertiles		
Prospective cohort		• Age: 35-65 years	Male	Male:	
D & B score = 11		• Characteristics: Free of hypertension at baseline. Excluded those unable to participate in regular PA due to poor health	T1 = Low = 0-1100	• T1 = 1.00 (referent)	
			T2 = Medium = 1101-1900	• T2 = 1.66	
			T3 = High >1900	• T3 = 1.73	
				Trend $p = 0.021$	
			Female	Female:	
			T1 = Low = 0-900	• T1 = 1.00 (referent)	
			T2 = Medium = 901-1500	• T2 = 0.94	
			T3 = High = >1500	• T3 = 1.16	
			Outcome measure: Incidence of hypertension through self reported diagnosis and death certificates	Trend $p = 0.648$	
			Cox proportional HR		
Paffenbarger et al 1983 [78]	To examine the relationship of student and alumnus PA patterns and other characteristics with incident hypertension.	• n = 14,998	PA Assessment: Questionnaire for PA based on number of stairs ascended, blocks walked and hours per week of light and vigorous sports play, yard work etc.	There was no significantly reduced risk for hypertension in men who climbed 50 plus stairs per day (compared to < 50 stairs); who walked 5 plus blocks per day (compared to < 5 blocks); or who played light sports (compared to those who did not).	Contemporary vigorous exercise was inversely related to hypertension risk.
USA		• Sex: Men			
Prospective cohort		• Age: 35-74 yr	Outcome measure: Diagnosis of hypertension by physicians using criteria of SBP > 160 mmHg and/or DBP > 95 mmHg	The 59% of men who did engage in vigorous sports were at 35% greater risk of hypertension than the 41% who did not.	
				RR = 1.35	
				Trend $p = <0.001$	
D & B score = 12		• Characteristics: free of hypertension Harvard Alumni Study	Multivariate estimates	Alumni on the low side of the physical activity index (< 2000 kcal/wk) had a 30% increased risk of hypertension then those $\geq$ 2000 kcal/wk.	
				RR = 1.30	
				Trend $p = 0.004$	

**Table 14: Studies examining the relationship between physical activity and hypertension. (Continued)**

Paffenbarger et al 1997 [79]	To investigate the quantity and intensity of energy expenditure required to delay hypertension and prevent premature death.	• n = 6,390	PA Assessment: Questionnaire for weekly sports play, divided into tertiles	RR (95% CI)	Lack of vigorous sports play independently increased the risk of developing hypertension.
USA		• Sex: Men	T1 = None	• T1 = 1.00 (referent)	
Prospective cohort		• Age: 45-84 yr	T2 = Light Only (< 4.5 METs)	• T2 = 1.04 (0.77-1.40)	
D & B score = 12		• Characteristics: Free of hypertension, CHD, diabetes, COPD and potentially malignant cancer in 1977	T3 = Moderately vigorous (≥ 4.5 METs)	• T3 = 0.77 (0.62-0.96)	
		• Harvard Alumni Study	Outcome measure: Incident hypertension	Trend $p = 0.004$	
Hu et al 2004 [81]	To discover whether regular PA can reduce the risk of hypertension in normal weight and overweight men and women.	• n = 8,302 men; 9,139 women	11 year follow up	Multivariate adjusted HR (95% CI), men	Regular PA can reduce the risk of hypertension. The protective effect of PA was observed in both sexes regardless of level of obesity.
Finland		• Sex: Men and women	PA assessment: Questionnaire for OPA, LTPA and commuting PA, divided into tertiles	• T1 = 1.00 (referent)	
Prospective cohort		Age: 25-64 yr	T1 = Low	• T2 = 0.63	
D & B score = 13		Characteristics: Healthy and free of hypertension at baseline	T2 = Medium	• T3 = 0.59	
			T3 = High	Trend $p = < 0.001$	
			Outcome Measure: Incidence of drug treated hypertension	Multivariate adjusted HR (95% CI), women	
			Cox proportional HR	• T1 = 1.00	
				• T2 = 0.82	
				• T3 = 0.71	
				Trend $p = 0.005$	
Gu et al 2007 [82]	To determine the 8-year incidence of HTN and its risk factors among Chinese adults.	• n = 10,525	Baseline Examination in 1991 with 8 year follow up	RR (95% CI), men	Increasing PA has the potential to reduce incidence of hypertension.
China		• Sex: Men and women	PA assessment: Questionnaire administered by trained staff, divided into groups	• G1 = 1.00 (referent)	
Prospective cohort		Age: ≥ 40 yr	G1 = Low	• G2 = 1.12 (0.86-1.46)	
D & B score = 13		Characteristics: Healthy and free from hypertension at baseline.	G2 = Medium	• G3 = 1.27 (1.10-1.47)	
			G3 = High	RR (95% CI), women	

**Table 14: Studies examining the relationship between physical activity and hypertension. (Continued)**

			Outcome measure: HTN as defined at SBP $\geq$ 140 mmHg and/or DBP $\geq$ 90 mmHg or current use of antihypertensive medication	• G1 = 1.00 (referent)	
			t-tests, chi squared tests, Cochran-Armitage modeling, Modified Poisson approach	• G2 = 1.14 (0.98-1.34)	
				• G3 = 1.22 (1.02-1.45)	
Hayashi et al 1999 [83]	To investigate the association of the duration of the walk to work and LTPA with the risk for hypertension.	• n = 6,017	PA assessment: Questionnaire on health related behaviours and exercise Walk time to work	RR (95% CI) Frequency walk time to work (minutes)	The duration of walk to work was associated with a decreased risk of hypertension even after adjustment.
Japan		• Sex: Men	T1 = 0-10 min	• T1 = 1.00 (referent)	Regular PA (at least once weekly) was inversely related to the risk of incident hypertension
Prospective cohort		• Age: 35-60 yr	T2 = 11-20 min	• T2 = 0.65 (0.47-0.90)	
D & B score = 12		• Characteristics: Free from HTN at baseline. All employees at gas company in Osaka Japan. All had sedentary jobs.	T3 = $\geq$ 21 min	• T3 = 0.72 (0.59-0.88)	
			Outcome measure: Diagnosed with hypertension (as defined by a SBP $\geq$ 160 mmHg, a DBP $\geq$ 95 mmHg, or use of antihypertensive medication)	Trend $p = < 0.001$	
			Cox proportional HR		
Nakanishi et al 2005 [84]	To examine the relationship of overall PA to the risk of developing hypertension in normotensive Japanese male office workers over a 7 year observation period.	• n = 2,548	7 year follow up	Multivariate adjusted RR (95% CI) by PA level only	The rate of rise in both SBP and DBP in each follow-up year decreased with higher EE and that the risk of developing hypertension decreased in a dose dependent manner with higher daily life activity level.
Japan		• Sex: Men		Q1 = 1.00 (referent)	Analysis stratified by the presence of or absence of a risk factor showed the negative association of daily life activity with the risk of developing hypertension for men at both low and high risk. This tendency was also observed among men in all 3 categories of normotension.
Prospective cohort		• Age: 35-59 yr	PA assessment: 1-day activity record and reported the type and frequency on a weekly basis of LTPA, divided into quartiles (kcal/kg/d)	Q2 = 0.84 (0.72-0.98)	

**Table 14: Studies examining the relationship between physical activity and hypertension. (Continued)**

D & B score = 12		<ul style="list-style-type: none"><li>• Characteristics Healthy at baseline. No hypertension or CHD. All office workers for a Japanese company</li></ul>	<ul style="list-style-type: none"><li>• Q1 = &lt;33.3</li><li>• Q2 = 33.3-36.9</li><li>• Q3 = 37.0-40.3</li><li>• Q4 = 40.4</li></ul> <p>3 categories of normotensive BP Low Normal: SBP &lt; 120, DBP &lt; 80 Normal: SBP 120-130, DBP 80- 85 High Normal: SBP 130-139 DBP 85-89</p> <p>3 categories of normotensive BP Low Normal: SBP &lt; 120, DBP &lt; 80 Normal: SBP 120-130, DBP 80- 85 High Normal: SBP 130-139 DBP 85-89</p> <p>Cox proportional hazard model</p>	<ul style="list-style-type: none"><li>• Q3 = 0.75 (0.63-0.88)</li><li>• Q4 = 0.54 (0.45-0.64)</li></ul> <p>Trend <i>p</i> = &lt; 0.001</p> <p>Multivariate adjusted RR (95% CI) by PA level, low normal BP</p> <ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li></ul> <ul style="list-style-type: none"><li>• Q2 = 0.70 (0.47-1.05)</li></ul> <ul style="list-style-type: none"><li>• Q3 = 0.55 (0.37-0.83)</li><li>• Q4 = 0.43 (0.28-0.65)</li></ul> <p>Trend <i>p</i> = &lt;0.001</p> <p>Multivariate adjusted RR (95% CI) by PA level, normal BP</p> <ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li><li>• Q2 = 0.89 (0.68-1.16)</li><li>• Q3 = 0.69 (0.52-0.91)</li><li>• Q4 = 0.50 (0.37-0.68)</li></ul> <p>Trend <i>p</i> = &lt;0.001</p> <p>Multivariate adjusted RR (95% CI) by PA level, high normal BP</p> <ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li><li>• Q2 = 0.86 (0.69-1.07)</li><li>• Q3 = 0.88 (0.69-1.11)</li><li>• Q4 = 0.60 (0.46-0.78)</li></ul> <p>Trend <i>p</i> = 0.001</p>	
Foy et al 2006 [85]	To examine whether insulin resistance is associated with the effect of vigorous or moderate PA on baseline BP.	<ul style="list-style-type: none"><li>• n = 1,599</li></ul>	Baseline examination in 1992-1993	Unadjusted OR (95% CI)	Participants who meet or exceed current caloric expenditure recommendations for VPA demonstrate significantly less hypertension than do sedentary or underactive individuals.

**Table 14: Studies examining the relationship between physical activity and hypertension. (Continued)**

USA		• Sex: Men and women	PA assessment: VPA over the past year was determined via a 1-year recall of physical activity (kcal/d), divided into 3 groups	• T1 = 1.00 (referent)	
Cross sectional		• Age: 40-69 yr	• T1 = 0	• T2 = 0.69 (0.53-0.88)	
D & B score = 12		• Characteristics: Community dwelling adults	• T2 = 1-149 kcal/day	• T3 = 0.57 (0.45-0.74)	
		• Insulin Resistance Atherosclerosis Study	• T3 = >150 kcal/day	• Trend $p = < 0.001$	
				Adjusted OR (95% CI)	
				• T1 = 1.00 (referent)	
				• T2 = 0.82 (0.62-1.09)	
				• T3 = 0.73 (0.55-0.98)	
				Trend $p = 0.004$	
Folsom et al 1990 [270]	To examine the relationship between fat distribution and the 2-yr incidence of hypertension and stroke.	• n = 41,837	Baseline mailed survey in 1986: Pa assessment: Questionnaire for LTPA	• 978 cases	High PA reduced the risk of hypertension only before adjusting for other factors.
USA		• Sex: Women	T1 = Low	Age Adjusted RR (95% CI)	
Prospective cohort		• Age: 55-69 years (yr)	T2 = Medium	• T1 = 1.00 (referent)	
D & B score = 12		• Characteristics: All free of HTN at baseline	T3 = High	• T2 = 0.9 (0.7-1.1)	
			Mantel-Haenszel method	• T3 = 0.7 (0.6-0.9)	
			Multiple logistic regression		
Levenstein et al 2001 [271]	To examine the effects of a variety of psychosocial factors on the development of HTN in men and women in the general population.	• n = 1,031 men, 1,326 women	Questionnaires in 1965 and 1974, cohort followed until 1994	LTPA predictor of hypertension OR (95% CI)	Risk of HTN was reduced with increases in LTPA in women.
USA		• Sex: Men and women	PA assessment: LTPA rated on a scale of 0-16 points and analysed as a continuous variable	• All Subjects: 0.94 (0.91-0.97)	
Prospective cohort		• Characteristics: Free of hypertension at baseline	Outcome measure: Incidence of hypertension (defined as those who are taking antihypertensive medications)	• Women: 0.90 (0.87-0.94)	
D & B score = 13		• Alameda cohort study	Logistic regression analysis	• Men: 0.98 (0.94-1.02)	

D & B score, Downs and Black quality score; YR, years; PF, physical fitness; BMI, body mass index; MET, metabolic equivalent; PA, physical activity; MI, myocardial infarction; G, groups; Q, quartile or quintile; 95% CI, confidence interval; SBP, systolic blood pressure; DBP, diastolic blood pressure; EE, energy expenditure; kcal/wk, kilocalories per week; T, tertile; RR, risk ratio; HR, hazard ratio; CHD, coronary heart disease; COPD, chronic obstructive pulmonary disease; OPA, occupational physical activity; LTPA, leisure-time physical activity; BP, blood pressure; kcal/day, kilocalories per day.

There were a total of 11,441 reported cases of hypertension (ranging per study from 118-2,936). The total length of study follow-up averaged 8.6 yr (ranging from 0-16 yr). The articles were published over a 24 yr period ranging from 1983 to 2007.

All studies reviewed demonstrated positive effects of physical activity on the risk for hypertension. Of these studies all (7; 58%) revealed an inverse and graded relationship between hypertension and at least one measure of physical activity or fitness. Across all studies, when comparing the most active/fit group versus the least active/fit group we found an average RR of 0.68 (median = 0.70, range 0.37 to 0.90). Therefore, we observed that physical activity/fitness was associated with an average risk reduction of 32% for hypertension. It should be noted that the study [75] demonstrating the largest risk reduction (63%) evaluated cardiorespiratory fitness directly during a maximal treadmill test. This supports research (as discussed previously) which indicates that physical fitness is a better predictor of chronic disease than physical activity [6,18,19,32,33]. *Taken as a whole, the level of evidence can be classified as Level 3A.* The quality of studies was generally good with a mean Downs and Black score of 11 (median = 11, range = 10-12).

Five studies showed variable results (i.e., no clearly defined dose-response) while generally supporting the inverse relationship between physical activity/fitness and hypertension [76-80]. The variability in the response appears to be the result of different activity/fitness classifications and/or differing subject populations. For instance, some studies revealed that the dose-response relationships differed between genders and/or ethnicities [76,77]. Pereira et al. [76] revealed a 30% reduction in the risk for hypertension in the most active white men. There were graded dose-response relationships between indices of both leisure and sport activities in the white men.

However, there was a lack of association between physical activity and hypertension in white women and African American men and women. Similarly, Haapenen et al. [77] revealed a stronger association in men than in women. However, it should be noted clearly that other studies included in this systematic review evaluated women demonstrating a graded response [81]. Moreover, several studies were conducted with non-Caucasian populations and demonstrated a dose-dependent benefit [82-85]. In fact, data was obtained from varied regions of the world including USA (7), Japan (2), China (1), and Finland (1). Therefore, there is evidence to suggest that the protective effects of physical activity with respect to hypertension are transferable to women and non-Caucasian populations. However, further research is clearly warranted that examines the relationship between physical activity and hypertension in persons of

different ethnicities. Moreover, further research is needed to determine the effects of impact of socio-economic status on the observed relationships.

Some studies have indicated that vigorous activity is required to reduce the risk for hypertension. For instance, Paffenbarger [78] revealed that Harvard Alumni who did not engage in vigorous sports play were at a 35% higher risk for developing hypertension. However, there was no difference in the risk for hypertension in men who climbed >50 stairs per day, walked more than 5 city blocks daily, or engaged in light sports only. Similarly, the Paffenbarger and Lee [79] study revealed that moderately vigorous sports play was associated with a lower risk for hypertension, but physical activity (kcal/wk), walking distance (km/wk) and the amount of stairs climbed (floors/wk) were not significant predictors of the risk for hypertension. Collectively, this research group concluded that these findings highlighted the importance of the intensity of effort.

However, it should be noted that many of the studies in our systematic review observed the protective effect with moderate intensity physical activities. Findings from randomized controlled trials have also provided strong evidence that moderate intensity aerobic exercise is sufficient to reduce blood pressure and the risk for hypertension, particularly in at risk individuals [86,87]. The American College of Sports Medicine [88] recently advocated that to prevent hypertension, individuals should exercise on most, and preferably all, days of the week at a moderate intensity, for 30 min or more per day (continuous or accumulated). They also recommended supplementing endurance type activities with resistance exercise. This is supported by research indicating that moderate intensity resistance training can reduce blood pressure [89]. Collectively, this research and our current summary of the dose-response literature indicates that physical activity levels that are of a moderate to vigorous intensity are sufficient to lead to marked reductions in the risk for hypertension.

#### Implications

The impact of hypertension on North American society is enormous. In the US, 31% of non-institutionalized adults over the ages of 20 are currently thought to have hypertension [90]. In Canada, approximately 20% of adults report a diagnosis of hypertension including over 4 million Canadians [91-93]. It has been estimated that a 55 yr old Canadian with normal blood pressure has a greater than 90% chance of developing hypertension before the age of 80 yr [92]. The primary prevention of hypertension is of paramount importance to the attenuation of the risks and costs associated with hypertension and related comorbidities.

There is clear evidence that routine physical activity and/or increased physical fitness reduce greatly the risk

for hypertension in both normotensive and hypertensive individuals [18,19]. Extensive research has been conducted in the area including numerous prospective trials and various randomized controlled trials. Numerous reviews of the literature (of epidemiological and randomized controlled trials) have supported an inverse relationship between physical activity/fitness and in the incidence of hypertension [20,87,89,94-102]. In a recent systematic review of the prospective literature, Katzmarzyk and Janssen (2004) calculated that physically inactive individuals were at a 30% higher risk for hypertension (RR = 1.30 (95% CI = 1.16-1.46)) with a population attributable risk of 13.8% in Canada [20]. Acute bouts of exercise have also been shown to lead to transient changes in blood pressure that are potentially of health benefit [98]. For instance, blood pressure is often reduced after a single exercise session for 12-22 hr [88,103].


It is clear that routine physical activity is effective in both the primary and secondary prevention of hypertension. However, the optimal dosage of physical activity/exercise remains somewhat unclear. Our review of the literature examined critically the relationship between multiple levels of physical activity/fitness and the incidence of hypertension (in individuals without diagnosed hypertension). As identified above this evidence was compelling supporting the protective effects of habitual physical activity in the primary prevention of hypertension.

**Recommendation #4**

*For a reduced risk for hypertension, it is recommended that individuals should participate in 30 min or more of moderate to vigorous exercise on most days of the week. [Level 3, Grade A]*

**Primary Prevention of Colon and Breast Cancer  
Colon Cancer**

In our systematic search of the colon cancer literature, a total of 252 citations were identified during the electronic database search (Figure 7). Of these citations, 83 were identified in MEDLINE, 44 in EMBASE, 25 in Cochrane, and 100 in the CINAHL/SportDiscus/PsychInfo search. A total of 15 duplicates were found, leaving a total of 237 unique citations. A total of 164 articles were excluded after screening, leaving a total of 73 articles for full review. From these articles 47 were excluded after full-text review leaving 26 articles for inclusion, and an additional 7 articles were added from the authors' personal files. The reasons for exclusion included non-experimental/weak design (n = 8), reviews, summaries, meta-analyses (n = 13), editorial/comment (n = 3), not dealing specifically with colon cancer (n = 4), did not contain three levels of physical activity or not possible to determine dose-response relationship



**Figure 7** Results of the Literature Search for Colon Cancer.

(n = 9), and other (n = 10). Therefore, a total of 33 articles were included in the systematic review of the literature regarding the relationship between physical activity and the primary prevention of colon cancer.

These studies involved a total of 1,433,103 participants; averaging 43,427 participants per study (range 142-413,044). There were a total of 17,959 reported cases of colon cancer (ranging per study from 93-1,993). The total length of study follow-up for the prospective cohort studies averaged 10.7 yr (ranging from 4-26 yr). The articles were published over a 23 yr period ranging from 1985 to 2008. These studies involved large samples of men and women from regions throughout the world.

A dose-dependency of this relationship was present in the majority of the studies. When comparing the most



active/fit group versus the least active/fit group we found a mean risk reduction of 30% (median = 32%) across all studies. The most compelling literature was that which evaluated the relationship between moderate-to-vigorous leisure time physical activity. Based on the literature reviewed and the volume of activity assessed it would appear that Canada's guidelines for physical activity are sufficient to lower the risk for the development of colon cancer in asymptomatic adults. *The level of evidence would be considered to be Level 2A.* The studies were generally of a higher quality with a mean Downs and Black score of 13 (median = 14, range = 11-15).

It should be noted that there was considerable variability in the findings and conclusions of the studies (Table 15). As discussed later, the literature was further confounded by the fact that the relative risks associated with physical activity were often controlled (through multivariate analyses) for various potential confounding factors, which may actually inappropriately decrease the level of risk reduction associated with physical activity [31]. Moreover, similar to other chronic conditions this literature was limited greatly by the lack of consistent physical activity assessment and description. In many instances, it was difficult to determine the actual absolute volume and/or intensity of activity for each category of comparison. However, despite these limitations the results of these studies (taken as a whole) indicate that both occupation- and leisure time-related physical activity are inversely related to the risk of colon cancer.

#### **Breast Cancer**

As reviewed eloquently by others, the epidemiological evidence relating physical activity to a decreased incidence of breast cancer is persuasive. A recent systematic review of the literature found that more than 60 observational trials have examined the relationship between physical activity and breast cancer [31]. Previous reviews of the literature have revealed compelling and consistent findings indicating that habitual physical activity is associated with a reduced risk for breast cancer ranging from 20-80% [31,104].

Various investigations have attempted to evaluate the dose-response relationship between physical activity and the incidence of breast cancer (Table 16). Despite the volume of evidence available questions still remain regarding the minimal and optimal volume of exercise required to reduce the risk for breast cancer. As discussed by others [31,104] the findings are as varied as the investigations.

In our systematic search of the literature, a total of 571 citations were identified during the electronic database search (Figure 8). Of these citations, 228 were identified in MEDLINE, 89 in EMBASE, 56 in Cochrane, and 198 in the CINAHL/SportDiscus/PsychInfo search.

A total of 46 duplicates were found, leaving a total of 571 unique citations. A total of 411 articles were excluded after scanning, leaving a total of 114 articles for full review. From these articles 77 were excluded after full review leaving 37 articles for inclusion in the systematic review. An additional 6 articles were found through the reviewers' personal files. The reasons for exclusion included not containing three levels of physical activity or not possible to determine dose-response relationship ( $n = 1$ ), reviews, summaries, meta-analyses ( $n = 20$ ), report ( $n = 5$ ), editorial/comment ( $n = 21$ ), not a research article ( $N = 11$ ), not dealing specifically with breast cancer ( $n = 4$ ), not relevant ( $n = 5$ ), not primary prevention ( $n = 3$ ), and other ( $n = 10$ ). Therefore, a total of 43 articles were included in the systematic review of the literature regarding the relationship between physical activity and the primary prevention of breast cancer.

The data providing dose-response information is all observational in nature, involving both case control and cohort investigations. These studies involved a total of 1,861,707 participants averaging 44,326 subjects per study (range 526-680,000). There were a total of 80,247 reported cases of breast cancer (ranging per study from 109-17,986). The total length of study follow-up for the prospective cohort studies averaged 10.5 yr (ranging from 4-31 yr). The articles were published over a 14 yr period ranging from 1993-2007. These studies involved large samples of men and women from regions throughout the world.

The literature with respect to the primary prevention of breast cancer is as compelling as that found with respect to colon cancer. There is strong evidence that routine physical activity is associated with a reduced risk for the development of breast cancer. However, this literature is also confounded by many shortcomings (similar to other cancer literature) including considerable variability in the statistical analyses employed, the physical activity measurement tools used, and the experimental designs.

The overall risk reduction for breast cancer for individuals that are habitually physically active (at or above Canada's guidelines for physical activity) is thought to approximate 20-40% [31,105]. In our analyses, we found very similar findings. When comparing the most active group versus the least active group we found a mean (and median) risk reduction of 20% across all studies. *The level of evidence would be considered to be Level 2A.* Generally, the articles were of high quality with a mean Downs and Black score of 13 (median = 13, range = 9-14).

A dose-dependency of this relationship is also generally present in the majority of the studies. For instance, greater than 50% studies revealed a dose-response

**Table 15 Studies examining the relationship between physical activity and colon cancer.**

Publication Country Study Design Quality Score	Objective	Population	Methods	Outcome	Comments and Conclusions
Hou et al 2004 [272]	To examine the effect of various forms of PA on colon cancer risk, with particular attention to commuting PA.	• n = 931 case, 1,552 control	PA assessment: Interview for the following variables	• Number of cases: 931	Regular frequent PA over a long period of time reduces risk of CC.
China		• Sex: Men and women		Multivariate OR (95% CI) by OPA, men	
Case control D & B score = 14		• Age: 30-74 yr • Characteristics: Case: diagnosed with CC. controls: selected randomly from residents of urban Shanghai.	OPA (kJ/min) G1 = <8	• G1 = 1.00 (referent) • G2 = 1.23 (0.93-1.64) • G3 = 0.81 (0.59-1.19)	
			G2 = 8-12 G3 = >12	p = 0.10	
			Commuting PA (MET hr/wk) G1 = <48.3 G2 = 48.3-94.3 G3 = >94.3	Multivariate OR (95% CI) by OPA, women • G1 = 1.00 (referent) • G2 = 0.96 (0.69-1.16) • G3 = 0.64 (0.39-1.02) p = 0.009	
			LTPA (MET hr/wk) G1 = < 9.2 G2 = 9.2-13.6 G3 = >13.6	Multivariate OR (95% CI) Commuting PA, men • G1 = 1.00 (referent) • G2 = 1.11 (0.31-1.23) • G3 = 0.52 (0.27-0.87)	
			Outcome Measure: incident CC	p<0.001	
			Multiple logistic regression	Multivariate OR (95% CI) Commuting PA, women • G1 = 1.00 (referent) • G2 = 0.87 (0.42-1.52) • G3 = 0.56 (0.21-0.91) p = 0.007	
				Multivariate OR (95% CI) LTPA, men • G1 = 1.00 (referent) • G2 = 1.17 (0.13-1.95) • G3 = 0.72 (0.41-1.07) p = 0.06	

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

				Multivariate OR (95% CI) LTPA, women	
				• G1 = 1.00 (referent)	
				• G2 = 1.03 (0.41-1.59)	
				• G3 = 0.84 (0.13-2.25)	
				$p = 0.15$	
Boutron-Ruault et al 2001 [273]	To determine which step of the adenoma-carcinoma pathway was influenced by OPA and recreational PA.	• n = 480	PA assessment: Questionnaire and classified into 3 groups	Number of cases: 171	A sedentary lifestyle was associated with a high risk of CC.
France		• Sex: Men and women	G1 = Low	Age and gender adjusted OR (95% CI), OPA	
Case control		• Age: 30-79 years	G2 = Medium	• G1 = 1.00 (referent)	
D & B score = 13		• Characteristics: Cases had 1 <sup>st</sup> diagnosis of colorectal adenoma, controls were polyp free.	G3 = High	• G2 = 1.3 (0.8-2.0)	
				• G3 = 0.5 (0.3-0.9)	
				$p = 0.005$	
			Outcome Measure: Incident CC		
			Multiple logistic regression	Age and gender adjusted OR (95% CI), LTPA	
				• G1 = 1.00 (referent)	
				• G2 = 0.7 (0.4-1.1)	
				• G3 = 0.3 (0.2-0.5)	
				$p = <0.0001$	
				Age and gender adjusted OR (95% CI), Global PA	
				• G1 = 1.00 (referent)	
				• G2 = 0.8 (0.5-1.2)	
				• G3 = 0.3 (0.2-0.6)	
				$p = 0.0003$	
Brownson et al 1991 [274]	To investigate the risks of 16 cancer types in relation to OPA.	• n = 17,147	PA assessment: Medical records and classified into 3 groups:	Number of cases: 1,838	OPA is inversely related to risk of CC.
		• Sex: Men		Multivariate OR (95% CI)	
USA		• Age: ≥ 20 yr	OPA	G1 = 1.00 (referent)	
		• Characteristics: White, working	G1 = Low - Activity required <20% of time	G2 = 1.2 (1.0-1.5)	
Case controll			G2 = Moderate - Activity required 20-80% of time	G3 = 1.1 (1.0-1.3)	

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

D & B score = 15			G3 = High - Activity required >80% of time	p = 0.05	
			Outcome Measure: CC		
			Maximum likelihood estimates		
Calton et al 2006 [275]	To examine the relationship between PA and colon cancer risk in women.	• n = 31,783	11 year follow up	Number of cases: 243	Results do not support the hypothesis that PA is related to a lower incidence of CC in women.
USA		• Sex: Women	PA Assessment: Questionnaire / Phone interviews for the following variables, divided into 4 or 5 groups		
Prospective cohort		• Age: 61.1 yr		Multivariate RR (95% CI), TPA	
D & B score = 12		• Characteristics: Free from cancer at baseline		• G1 = 1.00 (referent)	
				• G2 = 1.45 (0.98-2.15)	
				• G3 = 1.16 (0.77-1.75)	
				• G4 = 1.27 (0.84-1.91)	
				• G5 = 1.15 (0.76-1.75)	
				p = 0.77	
			TPA (MET h/d)		
			G1 = 34.0-48.5	Multivariate RR (95% CI), MPA	
			G2 = 48.51-54.3	• G1 = 1.00 (referent)	
			G3 = 54.31-59.0	• G2 = 1.23 (0.82-1.83)	
			G4 = 59.1-64.9	• G3 = 1.47 (0.99-2.21)	
			G5 = 65.0-98.1	• G4 = 0.94 (0.61-1.46)	
				• G5 = 1.07 (0.70-1.62)	
			MPA (h/d)	p = 0.80	
			G1 = 0-3.0		
			G2 = 3.01-5.0		
			G3 = 5.01-6.70	Multivariate RR (95% CI), VPA	
			G4 = 6.71-8.14	• Q1 = 1.00 (referent)	
			G5 = 8.15-18.0	• Q2 = 1.19 (0.85-1.66)	
				• Q3 = 0.87 (0.59-1.29)	
			VPA (h/d)	• Q4 = 1.10 (0.78-1.55)	
			Q1 = 0	p = 0.80	
			Q2 = 0.1-1.0		

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

			Q3 = 1.1-2.0 Q4 = 2.1-14.0 Outcome Measure: Incidence of CC Cox proportional HR		
Chao et al 2004 [276]	To examine how the characteristics of recreational PA affect its association with colon cancer incidence among older.	• n = 151,174 (70,403 men; 80,771 women)	7 year follow up	Number of cases: 940	Increased amounts of time spent in recreational PA is associated with substantially lower risk of CC.
USA		• Sex: Men and women	PA assessment: Questionnaire for the following variables	Multivariate RR (95% CI) by recreational PA, men	
Prospective cohort		• Age: mean 63 yr		• G1 = 1.00 (referent)	
D & B score = 12		• Cancer prevention study II Nutrition Cohort		• G2 = 0.91 (0.69-1.19)	
			Recreational PA (h/wk)	• G3 = 0.72 (0.52-1.01)	
			G1 = None	• G4 = 0.86 (0.64-1.15)	
			G2 = <2	• G5 = 0.77 (0.54-1.08)	
			G3 = 2-3	• G6 = 0.58 (0.39- 0.87)	
			G4 = 4-6	p = 0.007	
			G5 = 7		
			G6 = ≥ 8	Multivariate RR (95% CI) by recreational PA, women	
			Recreational (MET h/wk)	• G1 = 1.00 (referent)	
			G1 = None	• G2 = 1.01 (0.70-1.44)	
			G2 = <7, 7-13	• G3 = 1.01 (0.68-1.49)	
			G3 = 14-23	• G4 = 0.97 (0.66-1.43)	
			G4 = 24-29	• G5 = 1.03 (0.65-1.65)	
			G5 = ≥ 30	• G6 = 0.65 (0.39-1.11)	
				p = 0.14	
			Walking (h/wk)		
			Q1 = None		
			Q2 = <4	Multivariate RR (95% CI) by recreational PA, men and women	
			Q3 = 4-6	• G1 = 1.00 (referent)	
			Q4 = ≥ 7	• G2 = 0.94 (0.75-1.16)	
				• G3 = 0.83 (0.65-1.07)	
			Walking plus other	• G4 = 0.89 (0.71-1.12)	

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

	activities (h/wk)	• G5 = 0.85 (0.64-1.12)
	Q1 = None	• G6 = 0.60 (0.44-0.83)
	Q2 = <4	<i>p</i> = 0.002
	Q3 = 4-6	
	Q4 = ≥ 7	Multivariate RR (95% CI) by MET h/ wk men
Outcome Measure: Incidence of CC		• G1 = 1.00 (referent)
		• G2 = 0.90 (0.68-1.18)
Cox proportional HR		• G3 = 0.83 (0.59-1.16)
		• G4 = 0.75 (0.55-1.01)
		• G5 = 0.86 (0.63-1.19)
		• G6 = 0.60 (0.41-0.87)
		<i>p</i> = 0.005
		Multivariate RR (95% CI) by MET h/ wk women
		• G1 = 1.00 (referent)
		• G2 = 1.02 (0.71-1.46)
		• G3 = 0.98 (0.65-1.47)
		• G4 = 1.0 (0.68-1.47)
		• G5 = 0.94 (0.60-1.48)
		• G6 = 0.77 (0.48-1.24)
		<i>p</i> = 0.15
		Multivariate RR (95% CI) by MET h/ wk men and women
		• G1 = 1.00 (referent)
		• G2 = 0.93 (0.75-1.16)
		• G3 = 0.88 (0.68-1.13)
		• G4 = 0.84 (0.66-1.06)
		• G5 = 0.89 (0.68-1.15)
		• G6 = 0.65 (0.49-0.87)
		<i>p</i> = 0.002
		Multivariate RR (95% CI) by walking, Men
		• Q1 = 1.00 (referent)
		• Q2 = 0.87 (0.66-1.15)
		• Q3 = 0.83 (0.60-1.16)
		• Q4 = 0.88 (0.61-1.25)
		<i>p</i> = 0.34

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

			<p>Multivariate RR (95% CI) by walking, women</p> <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 1.00 (0.70-1.44)</li> <li>• Q3 = 1.08 (0.71-1.63)</li> <li>• Q4 = 1.18 (0.71-1.95)</li> </ul> <p><math>p = 0.41</math></p> <p>Multivariate RR (95% CI) by walking plus other activities, men</p> <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.73 (0.53-1.02)</li> <li>• Q3 = 0.85 (0.58-1.24)</li> <li>• Q4 = 0.53 (0.36-0.78)</li> </ul> <p><math>p = 0.02</math></p> <p>Multivariate RR (95% CI) by walking plus other activities, women</p> <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.99 (0.67-1.47)</li> <li>• Q3 = 0.72 (0.43-1.19)</li> <li>• Q4 = 0.59 (0.36-0.98)</li> </ul> <p><math>p = 0.07</math></p>		
Colbert et al 2001 [277]	To examine the association between OPA and LTPA and colon cancer in male smokers.	<ul style="list-style-type: none"> <li>• n = 29,133</li> </ul>	12 year follow-up	Number of cases: 152	OPA is protective against CC in a dose-response manner.
USA		<ul style="list-style-type: none"> <li>• Sex: Men</li> </ul>			
Prospective cohort		<ul style="list-style-type: none"> <li>• Age: 50-69 yr</li> </ul>	PA assessment: Interview for OPA and LTPA	Multivariate RR (95% CI) by OPA	
D & B score = 13		<ul style="list-style-type: none"> <li>• Characteristics: Smokers</li> <li>• Alpha- Tocopherol, Beta-Carotene Cancer Prevention Study</li> </ul>		<ul style="list-style-type: none"> <li>• G1 = 0.61 (0.39-0.98)</li> <li>• G2 = 1.00 (referent)</li> <li>• G3 = 0.60 (0.34-1.04)</li> <li>• G4 = 0.45 (0.26-0.78)</li> </ul> <p><math>p = 0.003</math></p>	
			OPA		
			G1 = Non-worker		
			G2 = Sedentary		
			G3 = Light	Multivariate RR (95% CI), by LTPA	
			G4 = Moderate	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.82 (0.59-1.13)</li> </ul>	
			LTPA		
			G1 = Sedentary		
			G2 = Active		

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

			Outcome Measure: incident CC Cox proportional HR		
Dosemeci et al 1993 [278]	To examine associations between PA and cancer sites among workers in Turkey.	• n = 6,236 (3,486 cases in men and 379 cases in women; 2,127 control men and 244 control women)	PA assessment: Stanford Occupational Classification code system.	Number of cases: 93	Occupational EE is inversely related to risk of CC.
Turkey		• 93 cases for CC		Multivariate OR (95% CI) by total occupational EE	
Case control		• Sex: Men and women • Age: not indicated	Total Occupational EE (kj/min)	• G1 = 1.6 (0.9-2.8) • G2 = 1.1 (0.6-2.0)	
D & B score = 13		• Characteristics: All hospitalized Cases: Diagnosed with CC. Controls: included subjects diagnosed as non- cancers and cancers which there is no suggestion of an association with PA.	G1 = <8	• G3 = 1.0 (referent)	
			G2 = 8-12 G3 = >12	p = 0.04 When adjusted for socioeconomic status p = 0.03	
			Sitting time at work (h/d) Levels:	Multivariate OR (95% CI) by sitting time at work	
			G1 = <2 G2 = 2-6 G3 = >6	• G1 = 1.00 (referent) • G2 = 1.5 (0.9-2.5) • G3 = 1.5 (0.8-3.0)	
				p = 0.03 When adjusted for socioeconomic status p = 0.03	
			Outcome Measure: Incident CC Maximum likelihood estimates		
Friedenreich et al 2006 [279]	To investigate the role of PA in the development of colon cancer.	• n = 413,044  • Sex: Men and women	4 year follow-up	Multivariate RR (95% CI), TPA	Inverse association between PA and risk of CC, particularly for right sided tumours.
UK		• Age: 35-70 yr • Characteristics: Free of cancer at baseline	PA assessment: modified Baecke Questionnaire	• Q1 = 1.00 (referent)  • Q2 = 0.92 (0.76-1.12) • Q3 = 0.86 (0.70-1.04)	
Prospective cohort		• European Prospective Investigation into Nutrition and Cancer. (EPIC)	TPA	• Q4 = 0.78 (0.59-1.03)	
			Q1 = Inactive	p = 0.04	



**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

D & B score = 14			Q2 = Moderately inactive Q3 = Moderately active Q4 = Active Household PA (MET-h/wk) Q1 = <19.5 Q2 = 19.5-39.6 Q3 = 39.6-73.9 Q4 = ≥ 73.9	Multivariate RR (95% CI), TPA and right sided CC • Q1 = 1.00 (referent) • Q2 = 1.79 (0.59-1.06) • Q3 = 0.64 (0.47-0.86) • Q4 = 0.65 (0.43-1.00) $p = 0.004$	
				Multivariate RR (95% CI), household PA and right sided CC • Q1 = 1.00 (referent) • Q2 = 0.97 (0.75-1.27) • Q3 = 0.84 (0.64-1.12) • Q4 = 0.74 (0.54-1.02) $p = 0.05$	
			Outcome Measure: Incident CC		
			Cox proportional HR		
Giovannucci et al 1995 [280]	To examine the association between PA and colon cancer.	• n = 47,723	6 year follow-up	Multivariate RR (95% CI)	A moderate level of PA was related to a substantially lower risk of CC in this cohort of middle age to elderly men.
		• Sex: Men • Age: 40-75 yr	PA assessment: Questionnaire	• G1 = 1.00 (referent) • G2 = 0.73 (0.48-1.10)	
USA		• Characteristics: Health professionals		• G3 = 0.94 (0.63-1.39) • G4 = 0.78 (0.51-1.20)	
Prospective cohort		• Health Professionals Follow-up Study	Outcome Measure: Incidence of colon cancer	• G5 = 0.53 (0.32-0.88) $p = 0.03$	
D & B score = 12			Mantel-Haenszel estimator and logistic regression		
Isomura et al 2006 [281]	To examine the relationship of OPA, LTPA, commuting, housework and shopping with colorectal cancer risk.	• n = 1545 (778 cases, 767 controls)	PA assessment: Questionnaire and interview for the following variables	• Number of cases: 778	Adds to the evidence that PA confers decreased risk of CC, especially of distal CC in both men and women.
Japan		• Sex: Men and women		Multivariate OR (95% CI) for all CC by OPA, men • G1 = 1.00 (referent) • G2 = 0.9 (0.6-1.4)	
Case control		• Age: 20-74 yr	OPA, men		

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

D & B score = 12	• Characteristics: Free from cancer at baseline	G1 = Sedentary	• G3 = 0.7 (0.4-1.0)
		G2 = Moderate G3 = Hard	$p = 0.06$
	• Fukuoka colorectal cancer study	OPA, women	Multivariate OR (95% CI) for proximal CC by OPA, men
		G1 = Sedentary	• G1 = 1.00 (referent)
		G2 = Active	• G2 = 1.2 (0.6-2.2)
			• G3 = 0.7 (0.4-1.4)
		Total non-OPA, men (MET-h/wk)	$p = 0.45$
		G1 = 0.0	Multivariate OR (95% CI) for distal CC by OPA, men
		G2 = 0.1-15.9	• G1 = 1.00 (referent)
		G3 = 16.0	• G2 = 0.8 (0.4-1.4)
			• G3 = 0.6 (0.4-1.0)
			$p = 0.047$
		Total non-OPA women (MET hr/wk)	
		G1 = 0.0	
		G2 = 0.1-15.9	
		G3 = 16.0	Multivariate OR (95% CI) for all CC by non-OPA, men
		Moderate or hard non-OPA, men (MET hr/wk)	• G1 = 1.00 (referent)
		G1 = 0.0	• G2 = 0.9 (0.6-1.4)
		G2 = 0.1-14.9	• G3 = 0.8 (0.5-1.2)
		G3 = $\geq 15.0$	$p = 0.22$
			Multivariate OR (95% CI) for proximal CC by non-OPA, men
		Moderate or hard non-OPA, women (MET hr/wk)	• G1 = 1.00 (referent)
		G1 = 0.0	• G2 = 1.2 (0.6-2.1)
		G2 = 0.1-14.9	• G3 = 0.9 (0.5-1.7)
		G3 = 15.0	$p = 0.69$
	Outcome Measure: Incident CC		Multivariate OR (95% CI) for distal CC by non-OPA, men
			• G1 = 1.00 (referent)
	Multiple logistic regression analysis		• G2 = 0.8 (0.5-1.3)

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

<ul style="list-style-type: none"> <li>• G3 = 0.7 (0.4-1.1)</li> </ul>
<i>p</i> = 0.19
Multivariate OR (95% CI) for all CC by non-OPA, women
<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.9 (0.5-1.5)</li> <li>• G3 = 0.8 (0.5-1.4)</li> </ul>
<i>p</i> = 0.45
Multivariate OR (95% CI) for proximal CC by non-OPA, women
<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.5 (0.7-3.3)</li> <li>• G3 = 1.6 (0.7-3.6)</li> </ul>
<i>p</i> = 0.41
Multivariate OR (95% CI) for distal CC by non-OPA, women
<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.7 (0.4-1.3)</li> <li>• G3 = 0.6 (0.3-1.1)</li> </ul>
<i>p</i> = 0.12
Multivariate OR (95% CI) for all CC by moderate or hard non-OPA, men
<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.8 (0.6-1.2)</li> <li>• G3 = 0.8 (0.5-1.1)</li> </ul>
<i>p</i> = 0.24
Multivariate OR (95% CI) for proximal CC by moderate or hard non-OPA, men
<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.1 (0.6-2.1)</li> <li>• G3 = 1.0 (0.6-1.8)</li> </ul>
<i>p</i> = 0.99
Multivariate OR (95% CI) for distal CC by moderate or hard non-OPA, men
<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.7 (0.4-1.1)</li> <li>• G3 = 0.7 (0.4-1.0)</li> </ul>
<i>p</i> = 0.12

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

				Multivariate OR (95% CI) for all CC by moderate or hard non-OPA, women <ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 1.0 (0.6-1.6),</li><li>• G3 = 0.8 (0.5-1.4)</li></ul> <p>p = 0.35</p> Multivariate OR (95% CI) for proximal CC by moderate or hard non-OPA, women <ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 1.3 (0.6-2.5)</li><li>• G3 = 1.3 (0.6-2.7)</li></ul> <p>p = 0.59</p> Multivariate OR (95% CI) for distal CC by moderate or hard non-OPA, women <ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.8 (0.5-1.5)</li><li>• G3 = 0.5 (0.3-1.1)</li></ul> <p>p = 0.41</p>	
Johnsen et al 2006 [282]	To investigate the effects of OPA on colon cancer incidence.	<ul style="list-style-type: none"><li>• n = 54,478 (28,356 men, 26,122 women)</li><li>• Sex: Men and women</li></ul>	7.6 year follow-up	<ul style="list-style-type: none"><li>• Number of cases: 140 women, 157 men</li><li>• Number of dropouts: &lt;0.8%</li></ul>	No support for the hypothesis that OPA measured by MET-score may be associated with a lower risk of CC.
Denmark Prospective cohort		<ul style="list-style-type: none"><li>• Age: 50-64 yr</li><li>• Characteristics: Free of Cancer at baseline</li><li>• Diet, Cancer and Health Study</li></ul>	PA assessment: Questionnaire for OPA by MET score, 4 groups	Multivariate RR (95% CI), men <ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li><li>• Q2 = 1.11 (0.69-1.77)</li><li>• Q3 = 1.17 (0.77-1.79)</li><li>• Q4 = 0.95 (0.58-1.55)</li></ul>	
D & B score = 13			Q1 = Sitting		
			Q2 = Standing		
			Q3 = Manual		
			Q4 = Not working		
			Outcome Measure: Incidence of colon cancer	Multivariate RR (95% CI), women <ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li><li>• Q2 = 1.15 (0.68-1.93)</li><li>• Q3 = 1.34 (0.83-2.16)</li><li>• Q4 = 0.96 (0.60-1.53)</li></ul>	
			Cox proportional HR		

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

Larsen et al 2006 [283]	To examine the relationship between PA and colorectal cancer.	<ul style="list-style-type: none"> <li>• n = 6,961</li> </ul>	PA assessment: Questionnaire (scored from 2-12), divided into quartiles:	Number of cases: 108	Inactivity was not a significant risk factor for advanced colonic neoplasia.
Norway		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> <li>• Age: 50-64</li> <li>• Characteristics: No history of colorectal surgery, radiotherapy, cardiopulmonary disease, anticoagulant therapy, coronary episode.</li> </ul>	Q1 = 2-4	RR (95% CI) <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.61 (0.32-1.16)</li> </ul>	
Cross-sectional evaluation within a randomized controlled trial			Q2 = 5	<ul style="list-style-type: none"> <li>• Q3 = 0.75 (0.45-1.26)</li> </ul>	
			Q3 = 6	<ul style="list-style-type: none"> <li>• Q4 = 0.56 (0.34-0.92)</li> </ul>	
			Q4 = 7-12	p = 0.04	
			Outcome Measure: Positive test for colonic neoplasia	Multivariate RR (95% CI)	
D & B score = 13				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.64 (0.33-1.25)</li> <li>• Q3 = 0.82 (0.47-1.43)</li> <li>• Q4 = 0.67 (0.39-1.16)</li> </ul>	
			Multivariate logistic regression analysis	p = 0.23	
Larsson et al 2006 [284]	To investigate the association between PA and colorectal cancer.	<ul style="list-style-type: none"> <li>• n = 45,906</li> <li>• Sex: Men</li> <li>• Age: 45-79 yr</li> </ul>	7.1 year follow-up	Number of cases: 309 (133 proximal, 138 distal)	Results support a role of PA in reducing the risk of CC.
Sweden		<ul style="list-style-type: none"> <li>• Characteristics: Free of cancer at baseline</li> </ul>	PA assessment: Questionnaire for the following variables	Multivariate HR (95% CI) by LTPA	
Prospective cohort				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.66 (0.43-1.02)</li> </ul>	
			LTPA (min/day)	<ul style="list-style-type: none"> <li>• Q3 = 0.68 (0.46-1.01)</li> <li>• Q4 = 0.56 (0.37-0.83)</li> </ul>	
D & B score = 14			Q1 = <10	p = 0.01	
			Q2 = 10-29		
			Q3 = 30-59		

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

			Q4 = $\geq 60$	Multivariate HR (95% CI) by home/housework PA	
			Home/housework PA (h/day)	• Q1 = 1.00 (referent)	
				• Q2 = 0.75 (0.58-0.97)	
			Q1 = none	• Q3 = 0.75 (0.58-0.97)	
			Q2 = $<1$	• Q4 = 0.68 (0.48-0.96)	
			Q3 = 1-2	$p = 0.01$	
			Q4 = $\geq 3$		
			Incidence of Proximal CC(h/day)	Multivariate HR (95% CI) for distal CC by LTPA	
			G1 = $<1$	• Q1 = 1.00 (referent)	
			G2 = 1-2		
			G3 = $\geq 3$	• Q2 = 0.51 (0.28-0.93)	
				• Q3 = 0.50 (0.29-0.87)	
				• Q4 = 0.40 (0.22-0.70)	
				$p = 0.01$	
			Outcome Measure: Incident CC		
				Multivariate HR (95% CI) for proximal CC by home/housework PA	
			Cox proportional HR	• G1 = 1.00 (referent)	
				• G2 = 0.78 (0.53-1.14)	
				• G3 = 0.50 (0.29-0.89)	
				$p = 0.02$	
Lee and Paffenbarger 1994 [285]	To predict cancer risk using prospective assessments of PA.	• n = 17,607	26 year follow-up	• Number of cases: 280	Found a trend, of borderline statistical significance toward decreasing CC risk with increasing PA.
		• Sex: Men		• Number of dropouts: 14%	
		• Age: 30-79 yr	PA assessment: Questionnaire for PA level (kcal/wk)		
USA		• Characteristics: Healthy at baseline		Multivariate RR (95% CI), Model A: PA in 1962/1966 and updated in 1977	
		• Harvard College Alumni		• G1 = 1.00 (referent)	
Prospective cohort			G1 = $<1000$	• G2 = 1.07 (0.81-1.42)	
			G2 = 1000-2499	• G3 = 1.08 (0.81-1.46)	
			G3 = $\geq 2500$	$p = 0.58$	

D & B score = 13

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

			Outcome Measure: Incidence of fatal and non fatal CC	Multivariate RR (95% CI), Model B: PA in both 1962/1966 and 1977	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.75 (0.42-1.35)</li> <li>• G3 = 0.94 (0.54-1.64)</li> </ul>	
			Cox proportional HR	p = 0.76	
Lee et al 1997 [286]	To investigate whether PA alters the risk of developing CC in men.	• n = 20,614	10.9 year follow-up	Number of cases: 217	Data does not support the hypothesis that PA is related inversely to risk of developing CC.
		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: 40-84 yrs</li> </ul>	PA assessment: Questionnaire for the following variables	Multivariate RR (95% CI), frequency of	
USA		• Characteristics: Physicians, free of cancer at baseline		PA at baseline	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.1 (0.7-1.7)</li> <li>• G3 = 1.2 (0.8-1.6)</li> <li>• G4 = 1.1 (0.7-1.6)</li> </ul>	
Prospective cohort			Frequency of PA at baseline (times/ week)	p = 0.6	
		Physicians Health Study	G1 = <1 G2 = 1 G3 = 2-4 G4 = 5+	RR (95% CI), frequency of PA at baseline and 36 months	
D & B score = 15			Frequency of PA at baseline and 36 months G1 = 1/<1 G2 = <1/1+ G3 = 1+< 1 G4 = 1+/1+	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.2 (0.5-2.7)</li> <li>• G3 = 1.4 (0.9-2.3)</li> <li>• G4 = 1.3 (0.9-2.0)</li> </ul>	
			Outcome Measure: Incidence of fatal and non-fatal CC Cox proportional HR		

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

Lee et al 2007 [287]	To examine the association between PA and the risk of developing CRC in Japanese men and women.	• n = 65,022	6 year follow-up	Number of cases: 154 proximal CC, 166 distal CC	PA may prevent CC among Japanese men.
Japan		• Sex: Men and women • Age: 40-69 yr • Characteristics	PA assessment: Questionnaire for PA level (median MET hr/d)	Multivariate RR (95% CI) for CC men • Q1 = 1.00 (referent)	
Prospective cohort		• Ethnicity: Japanese	Q1 = 28.25 Q2 = 33.25 Q3 = 35.25 Q4 = 43.75	• Q2 = 0.87 (0.61-1.26) • Q3 = 0.62 (0.41-0.95) • Q4 = 0.58 (0.39-0.87) p = 0.006	
D & B score = 13			Outcome Measure: Incidence of CC Cox proportional HR	Multivariate RR (95% CI) for proximal CC men • Q1 = 1.00 (referent) • Q2 = 0.89 (0.52-1.51) • Q3 = 0.44 (0.22-0.86) • Q4 = 0.29 (0.14-0.60) p < 0.001 Multivariate RR (95% CI) for distal CC Men • Q1 = 1.00 (referent) • Q2 = 0.92 (0.54-1.54) • Q3 = 0.75 (0.42-1.33) • Q4 = 0.89 (0.53-1.51) p = 0.685 PA level and incidence of CC women Total CC • Q1 = 1.00 (referent) • Q2 = 1.03 (0.65-1.64) • Q3 = 0.91 (0.57-1.47) • Q4 = 0.89 (0.54-1.49) p = 0.610 Proximal CC women • Q1 = 1.00 (referent) • Q2 = 1.14 (0.61-2.12) • Q3 = 1.01 (0.53-1.89)	



**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

				<ul style="list-style-type: none"> <li>• Q4 = 0.55 (0.24-1.26)</li> </ul> <p>p = 0.151</p> <p>Distal CC women</p> <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 1.09 (0.52-2.29)</li> <li>• Q3 = 0.77 (0.34-1.74)</li> <li>• Q4 = 1.37 (0.66-2.85)</li> </ul> <p>p = 0.401</p>	
Longnecker et al 1995 [288]	To examine the relationship between OPA and vigorous LTPA and the risk of cancer of the right colon and rectum.	<ul style="list-style-type: none"> <li>• n = 242 rectal cancer and 703 controls</li> </ul>	PA assessment: Interview for vigorous LTPA and OPA (coded and self-reported), divided into groups:	Number of cases: 163	The amount of time spent at vigorous LTPA was associated with a decreased risk of cancer of the right colon.
USA		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: ≥ 31 yr</li> <li>• Characteristics: Case: Diagnosed with adenocarcinoma of the right colon or rectum. Controls: Both community and hospital. No history of large bowel cancer.</li> </ul>		RR (95% CI) by vigorous LTPA	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.73 (0.23-2.29)</li> </ul>	
Case control				<ul style="list-style-type: none"> <li>• G3 = 0.47 (0.16-1.36)</li> <li>• G4 = 0.60 (0.35-1.00)</li> </ul>	
			Vigorous LTPA (h/wk)		
			G1 = 0	p = 0.03	
			G2 = ≤ 0.5		
			G3 = 1	Multivariate OR (95% CI) by vigorous LTPA	
			G4 = >1		
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.81 (0.26-2.54)</li> <li>• G3 = 0.36 (0.11-1.14)</li> <li>• G4 = 0.57 (0.33-0.97)</li> </ul>	
			Coded lifetime OPA		
			G1 = Sedentary		
			G2 = light work		
			G3 = moderate	p = 0.06	
			G4 = heavy		
			Self reported lifetime OPA	Multivariate OR (95% CI) by coded lifetime OPA	
			G1 = Sedentary	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.79 (0.39-1.61)</li> <li>• G3 = 0.79 (0.36-1.74)</li> <li>• G4 = 0.99 (0.30-3.22)</li> </ul>	
			G2 = light work		
			G3 = more than light work		

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

				<p><math>p = 0.42</math></p> <p>Outcome Measure: Diagnosed with CC</p> <p>Multivariate OR (95% CI) by self reported lifetime OPA</p> <p>Conditional Logistic Regression</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.85 (0.41-1.76)</li> <li>• G3 = 0.68 (0.31-1.52)</li> </ul> <p><math>p = 0.15</math></p>	
Mai et al 2007 [289]	To examine in detail the relationship between recreational PA and invasive CC among women.	<ul style="list-style-type: none"> <li>• n = 120,147</li> <li>• Sex: Women</li> <li>• Age: 22-84 yr</li> <li>• Characteristics: no prior history of CC</li> <li>• California Teachers Study</li> </ul>	7 year follow-up	<p>Number of cases: 395</p> <p>RR (95% CI) by MPA over past 3 years</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.95 (0.72-1.24)</li> <li>• G3 = 0.78 (0.62-0.97)</li> </ul> <p><math>p = 0.02</math></p> <p>RR (95% CI) by strenuous + moderate (lifetime) PA:</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.79 (0.56-1.11)</li> <li>• G3 = 0.64 (0.44-0.93)</li> </ul> <p><math>p = 0.04</math></p>	Modest inverse association between recreational PA and CC.
USA			PA assessment: Questionnaire		
Prospective cohort			MPA over past 3 yrs (h/wk/yr)		
D & B score = 15			<p>G1 = 0-0.50</p> <p>G2 = 0.51-1.99</p> <p>G3 = <math>\geq 2.00</math></p> <p>Strenuous + Moderate (lifetime) PA (h/wk/yr)</p> <p>G1 = 0.0-0.50</p> <p>G2 = 0.51-3.99</p> <p>G3 = <math>\geq 4.00</math></p> <p>Outcome Measure: Incidence of invasive adenocarcinoma of the colon</p> <p>Cox proportional HR</p>		
Martinez et al 1997 [290]	To examine whether LTPA could significantly influence the risk of CC in women.	<ul style="list-style-type: none"> <li>• n = 89,448</li> <li>• Sex: Women</li> </ul>	6 year follow-up	<p>Number of cases: 212</p>	Significant inverse association between LTPA and incidence of CC in women.

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

		• Age: 30-55 yr	PA assessment: Questionnaire for LTPA	Multivariate RR (95% CI) for all CC	
USA		• Characteristics: Nurses, free from cancer at baseline		• G1 = 1.00 (referent)	
Prospective cohort			G1 = <2 G2 = 2-4 G3 = 5-10 G4 = 11-21 G5 = >21	• G2 = 0.71 (0.44-1.15) • G3 = 0.78 (0.50-1.20) • G4 = 0.67 (0.42-1.07) • G5 = 0.54 (0.33-0.90)	
D & B score = 14			Outcome Measure: Incidence of CC	$p = 0.03$ Multivariate RR (95% CI) for distal CC	
			Mantel-Haenszel Estimator and logistic regression models	• G1 = 1.00 (referent) • G2 = 0.92 (0.48-1.79) • G3 = 0.81 (0.43-1.55)  • G4 = 0.71 (0.36-1.41) • G5 = 0.31 (0.12-0.77) $p = 0.01$ Multivariate RR (95% CI) for proximal CC	
				• G1 = 1.00 (referent) • G2 = 0.54 (0.23-1.22) • G3 = 0.79 (0.40-1.56) • G4 = 0.62 (0.30-1.32) • G5 = 0.77 (0.38-1.58) $p = 0.67$	
Nilsen et al 2008 [291]	To study the separate associations of recreational PA with the incidence of, and mortality from cancer in the ascending, transverse, descending and sigmoid segments of the colon.	• n = 59,369	17 year follow-up	Number of cases: 736	Strong inverse associations between recreational PA and risk of cancer morbidity and mortality of the transverse and sigmoid colon but no association for cancer in the ascending and descending colon.
		• Sex: Men and women	PA assessment: Questionnaire for frequency and duration of recreational PA	HR (95% CI) by frequency of recreational PA, men	
Norway		• Age: not indicated		• G1 = 1.00 (referent)	
Prospective cohort		• Characteristics: Free from cancer at baseline		• G2 = 0.84 (0.60-1.19)	

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

D & B score = 14	• Nord-Trondelag Health Study	• G3 = 0.82 (0.58-1.17) • G4 = 0.81 (0.57-1.15) • G5 = 0.77 (0.54-1.09)
	Frequency of Recreational PA (times per week)	
	G1 = none	$p = 0.18$
	G2 = <1	HR (95% CI) by frequency of
	G3 = 1	recreational PA, women
	G4 = 2-3	• G1 = 1.00 (referent)
	G5 = ≥ 4	• G2 = 0.91 (0.66-1.25) • G3 = 0.79 (0.57-1.09) • G4 = 0.66 (0.47-0.92)
	Duration of recreational PA (min per exercise)	• G5 = 0.99 (0.72-1.36)
	G1 = none	$p = 0.35$
	G2 = <15	
	G3 = 15-30	HR (95% CI) by duration of
	G4 = 31-60	recreational
	G5 = >60	PA, men
		• G1 = 1.00 (referent) • G2 = 1.07 (0.71-1.60) • G3 = 0.80 (0.57-1.12)
	Intensity of recreational PA	• G4 = 0.68 (0.48-0.97) • G5 = 0.74 (0.50-1.08)
	G1 = none	$p = 0.02$
	G2 = Low	HR (95% CI) by duration of
	G3 = Moderate/High	recreational PA, women
	Summary score for recreational PA	• G1 = 1.00 (referent)
	G1 = None	• G2 = 0.85 (0.59-1.23)
	G2 = Low	• G3 = 0.81 (0.60-1.09)
	G3 = High	• G4 = 0.73 (0.53-1.01)
	By subsite-specific (transverse colon, descending colon, sigmoid colon) CC	• G5 = 0.84 (0.53-1.34)
		$p = 0.10$
		HR (95% CI) by intensity of
		recreational PA, men
	Levels of REC PA:	

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

G1 = None	• G1 = 1.00 (referent)
G2 = < 1 x/wk	• G2 = 0.83 (0.62-1.12)
G3 = low score	• G3 = 0.74 (0.52-1.06)
G4 = high score	<i>p</i> = 0.11
Outcome Measure: incidence of fatal and non fatal CC	HR (95% CI) by intensity of recreational PA, women
Cox proportional HR	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.77 (0.59-1.01)</li> <li>• G3 = 0.89 (0.60-1.32)</li> </ul>
	<i>p</i> = 0.33
	HR (95% CI) by summary score for recreational PA, men
	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.85 (0.62-1.16)</li> <li>• G3 = 0.69 (0.48-0.98)</li> </ul>
	<i>p</i> = 0.06
	HR (95% CI) by summary score for recreational PA, women
	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.86 (0.64-1.01)</li> <li>• G3 = 0.72 (0.53-0.98)</li> </ul>
	<i>p</i> = 0.03
	HR (95% CI) by total CC and recreational PA, incidence
	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.88 (0.70-1.12)</li> <li>• G3 = 0.87 (0.70-1.08)</li> <li>• G4 = 0.73 (0.58-0.92)</li> </ul>
	<i>p</i> = 0.009
	HR (95% CI) by subsite specific CC and recreational PA, death
	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.87 (0.64-1.18)</li> <li>• G3 = 0.79 (0.59-1.04)</li> <li>• G4 = 0.56 (0.41-0.78)</li> </ul>
	<i>p</i> <0.001
	HR (95% CI) for transverse CC incidence and recreational PA
	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

				<ul style="list-style-type: none"> <li>• G2 = 0.75 (0.44-1.28)</li> <li>• G3 = 0.66 (0.41-1.08)</li> <li>• G4 = 0.44 (0.25-0.78)</li> </ul> <p><math>p = 0.004</math></p> <p>HR (95% CI) for transverse CC death and recreational PA</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.73 (0.36-1.49)</li> <li>• G3 = 0.40 (0.19-0.82)</li> <li>• G4 = 0.33 (0.14-0.76)</li> </ul> <p><math>p = 0.002</math></p> <p>HR (95% CI) for sigmoid CC incidence and recreational PA</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.88 (0.59-1.32)</li> <li>• G3 = 0.68 (0.46-1.01)</li> <li>• G4 = 0.48 (0.31-0.75)</li> </ul> <p><math>p &lt; 0.001</math></p> <p>HR (95% CI) for sigmoid CC death and recreational PA</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.78 (0.45-1.35)</li> <li>• G3 = 0.51 (0.30-0.87)</li> <li>• G4 = 0.29 (0.15-0.56)</li> </ul> <p><math>p &lt; 0.001</math></p>	<p>For the most active men, VPA was associated with a non-significant lower risk of CC.</p>
Schnohr et al 2005 [292]	To assess the association between LTPA and incidence of cancer in the general population.	<ul style="list-style-type: none"> <li>• n = 28,259 (15,043 men, 13,216 women)</li> </ul>	14 year follow-up	<ul style="list-style-type: none"> <li>• Number of cases: 215 men, 108 women</li> </ul>	
Denmark			PA assessment: Questionnaire for LTPA	Multivariate RR (95% CI), men	
Prospective cohort		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>	G1 = Low G2 = Moderate	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.08 (0.74-1.57)</li> </ul>	
D & B score = 13		<ul style="list-style-type: none"> <li>• Age: 20-93 yr</li> <li>• Characteristics: Free from cancer at baseline</li> </ul>	G3 = Vigorous	<ul style="list-style-type: none"> <li>• G3 = 0.72 (0.47-1.11)</li> </ul>	
		<ul style="list-style-type: none"> <li>• Copenhagen Heart Study, The Copenhagen County Centre of Preventive Medicine and the Copenhagen Male Study</li> </ul>	Outcome Measure: Incidence of CC	<p><math>p = 0.06</math></p> <p>Multivariate RR (95% CI), women</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

			Cox proportional HR	<ul style="list-style-type: none"> <li>• G2 = 1.02 (0.70-1.50)</li> <li>• G3 = 0.90 (0.56-1.46)</li> </ul>	
				$p = 0.68$	
Slattery et al 1988 [293]	To assess the relationship of PA and diet with the development of CC in Utah.	<ul style="list-style-type: none"> <li>• n = 229 cases, 384 controls</li> </ul>	PA assessment: Interview for the following variables	<ul style="list-style-type: none"> <li>• Number of cases: 229</li> </ul>	PA shows an inverse relationship with incidence of CC.
USA		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>		Multivariate OR (95% CI) by TPA, men	
Case control		<ul style="list-style-type: none"> <li>• Age: 40-79 yr</li> <li>• Characteristics: Case: Diagnosed with CC Controls: no history of cancer</li> </ul>	TPA	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 1.19 (0.67-2.13)</li> </ul>	
			Q1 = Low	<ul style="list-style-type: none"> <li>• Q3 = 0.88 (0.48-1.69)</li> </ul>	
			Q2	<ul style="list-style-type: none"> <li>• Q4 = .70 (0.38-1.29)</li> </ul>	
			Q3		
			Q4 = high	Multivariate OR (95% CI) by TPA, women	
			Intense PA	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> </ul>	
			G1 = none	<ul style="list-style-type: none"> <li>• Q2 = 0.97 (0.56-1.69)</li> </ul>	
			G2 = low	<ul style="list-style-type: none"> <li>• Q3 = 0.91 (0.52-1.60)</li> </ul>	
			G3 = high	<ul style="list-style-type: none"> <li>• Q4 = 0.48 (0.27-0.87)</li> </ul>	
			Non-intense PA	Multivariate OR (95% CI) by intense PA, men	
			Q1 = Low		
			Q2	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
			Q3	<ul style="list-style-type: none"> <li>• G2 = 0.83 (0.40-1.75)</li> </ul>	
			Q4 = high	<ul style="list-style-type: none"> <li>• G3 = 0.27 (0.11-0.65)</li> </ul>	
			Outcome Measure: Diagnosed with CC	Multivariate OR (95% CI) by intense PA, women	
			Multiple logistic regression analysis	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.55 (0.23-1.34)</li> </ul>	
				Multivariate OR (95% CI) by non-intense PA, men	
				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 1.40 (0.76-2.57)</li> <li>• Q3 = 0.93 (0.51-1.72)</li> <li>• Q4 = 1.25 (0.68-2.29)</li> </ul>	
				Multivariate OR (95% CI) by non-intense PA, women	
				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 1.09 (0.62-1.90)</li> </ul>	
D & B score = 13					

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

				<ul style="list-style-type: none"> <li>• Q3 = 0.94 (0.53-1.66)</li> <li>• Q4 = 0.53 (0.29-0.95)</li> </ul>	
Slattery et al 1997 [294]	To examine the relationship between weekly PA patterns (source, duration and frequency) and CC.	• n = 1,993 cases, 2,410 controls	PA Assessment: Interview, adapted CARDIA PA history	Number of cases: 1,993	High level of leisure time VPA during the past 20 yrs was associated with a reduced risk of CC in both men and women. The same associations were not observed with leisure time MPA.
USA		• Sex: Men and women		Multivariate OR (95% CI) by recent leisure time VPA, men	
Case control		• Age: 30-79 yr	Recent leisure time VPA	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.80 (0.64-1.01)</li> </ul>	
		• Characteristics: Cases: diagnosed with first primary CC. Controls: no history of CC			
D & B score = 14			Q1 = None	• Q3 = 0.84 (0.66-1.05)	
			Q2	• Q4 = 0.69 (0.55-0.87)	
			Q3		
			Q4 = High	Multivariate OR (95% CI) by recent leisure time VPA, women	The greatest inverse association was observed when activities were performed for longer periods of time per session.
		The Three Centered Diet, Activity and Lifestyle Colon Cancer Study	Leisure time VPA	• Q1 = 1.00 (referent)	
			Q1 = Low	• Q2 = 0.79 (0.61-1.02)	
			Q2	• Q3 = 0.83 (0.64-1.07)	
			Q3	• Q4 = 0.86 (0.67-1.10)	
			Q4 = High		
			Current PA (min)	Multivariate OR (95% CI) by leisure time VPA, men	
			G1 = <30		
			G2 = 30-60	• Q1 = 1.00 (referent)	
			G3 = ≥ 60	• Q2 = 0.97 (0.76-1.25)	
				• Q3 = 0.86 (0.67-1.09)	
			LTPA (ranked by time per session)	• Q4 = 0.61 (0.47-0.79)	
			Q1 = None	Multivariate OR (95% CI) by leisure time VPA, women	
			Q2 = Low - <30 min		
			Q3 = moderate - 30-60 min	• Q1 = 1.00 (referent)	
			Q4 = high ->60 min	• Q2 = 0.75 (0.59-0.95)	
				• Q3 = 0.68 (0.53-0.87)	
			Number of activity session per week	• Q4 = 0.63 (0.48-0.83)	



**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

	G1 = None	Multivariate OR (95% CI) by current MPA time per week
	G2 = 1	
	G3 = 2-4	• Q1 = 1.00 (referent)
	G4 = 5-7	• Q2 = 1.00 (0.83-1.21)
	G5 = >7	• Q3 = 0.90 (0.76-1.07)
		• Q4 = 0.92 (0.77-1.10)
Outcome Measure: Diagnosed with CC		Multivariate OR (95% CI) by current VPA time per week
Unconditional regression models		• Q1 = 1.00 (referent)
		• Q2 = 0.90 (0.73-1.12)
		• Q3 = 0.89 (0.71-1.10)
		• Q4 = 0.83 (0.69-0.98)
		Multivariate OR (95% CI) by leisure time MPA time per session
		• Q1 = 1.00 (referent)
		• Q2 = 1.20 (0.91-1.59)
		• Q3 = 1.09 (0.83-1.42)
		• Q4 = 1.08 (0.82-1.42)
		Multivariate OR (95% CI) by leisure time VPA time per session
		• Q1 = 1.00 (referent)
		• Q2 = 0.86 (0.74-0.99)
		• Q3 = 0.76 (0.64-0.90)
		• Q4 = 0.68 (0.52-0.87)
		Multivariate OR (95% CI) by number of MPA sessions/wk
		• G1 = 1.00 (referent)
		• G2 = 1.02 (0.79-1.30)
		• G3 = 0.86 (0.72-1.02)
		• G4 = 0.91 (0.81-1.14)
		• G5 = 1.02 (0.82-1.27)
		Multivariate OR (95% CI) by number of VPA sessions/wk
		• G1 = 1.00 (referent)
		• G2 = 0.72 (0.56-0.92)
		• G3 = 0.87 (0.73-1.03)
		• G4 = 1.00 (0.81-1.25)
		• G5 = 0.84 (0.61-1.15)

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

Slattery et al 1997 [295]	To determine how physical inactivity interacts with other components of energy balance in determining risk of CC.	• n = 1,993 cases, 2,410 controls	PA Assessment: Interview for lifetime VPA (PA index)	Number of cases: 1,993	These results support previous findings that physical inactivity is associated with an increased risk of developing CC.
USA		• Sex: Men and women • Age: 30-79 yr • Characteristics: Cases: diagnosed with first primary CC. Controls: no history of CC	Q1 = 10-12 Q2 = 7-9 Q3 = 4-6	Multivariate OR (95% CI), men • Q1 = 1.00 (referent) • Q2 = 1.60 (1.11-1.75)	
Case control		• The Three Centered Diet, Activity and Lifestyle Colon Cancer Study	Q4 = <4	• Q3 = 1.59 (1.26-2.01)	
D & B score = 14			Outcome Measure: Diagnosed with CC	Multivariate OR (95% CI), women • Q4 = 1.63 (1.26-2.12) • Q1 = 1.00 • Q2 = 1.14 (0.86-1.52) • Q3 = 1.13 (0.85-1.49)	
			Unconditional regression models	• Q4 = 1.59 (1.21-2.10)	
Takahashi et al 2007 [296]	To investigate the association between time spent walking each day and the risk of CRC.	• n = 20,519 men, 21,469 women	7 year follow-up	• Number of cases: 101	Time spent walking per day was associated with a lower risk of colon cancer in men but not in women.
		• Sex: Men and women	PA assessment: Questionnaire for time spent walking (h/day)	• Number of dropouts: 3.5%	
Japan		• Age: 40-64 yr • Characteristics: Free from cancer at baseline		Multivariate RR (95% CI), men • G1 = 1.00 (referent)	
Prospective cohort			G1 = <0.5 G2 = 0.5-1 G3 = >1	• G2 = 0.72 (0.43-1.21) • G3 = 0.38 (0.22-0.64) p < 0.001	
D & B score = 12			Outcome Measure: Incidence of CC	Time spent walking and incidence of CC	
			Cox proportional HR	Multivariate RR (95% CI), women • G1 = 1.00 • G2 = 2.68 (0.94-7.68) • G3 = 1.79 (0.64-4.96) p = 0.42	

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

Tang et al 1999 [297]	To investigate the association between PA, water intake and risk of CRC in a hospital based case controlled study.	<ul style="list-style-type: none"> <li>• n = 163 cases, 163 controls</li> </ul>	PA assessment: Interview	<ul style="list-style-type: none"> <li>• Number of cases: 163</li> </ul>	Found a negative association between LTPA and the risk of CC among men.
Taiwan		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>	LTPA METs	Multivariate RR (95% CI), men	
Case control		<ul style="list-style-type: none"> <li>• Age: 33-80 yr</li> <li>• Characteristics: Cases: Hospital patients diagnosed with colorectal cancer Controls: Hospital patients in hospital for other reasons, free of CRC.</li> </ul>	G1 = Sedentary	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 2.22 (0.68-7.21)</li> <li>• G3 = 0.19 (0.05-0.77)</li> </ul>	
D & B score = 14			G2 = Moderate (< 20 MET)		
			G3 = Active (≥20 MET)	p = 0.03	
			Outcome Measure: Diagnosis of CC	Multivariate RR (95% CI), women	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.52 (0.13-2.03)</li> <li>• G3 = 0.63 (0.18-2.18)</li> </ul>	
			Conditional logistic regression analysis	p = 0.48	
Tavani et al 1999 [298]	To investigate the relationship between PA and risk of CC in both sexes at different ages.	<ul style="list-style-type: none"> <li>• n = 5,379 (1,225 cases and 4,154 controls)</li> </ul>	PA assessment: Questionnaire on activity at work and during leisure time	<ul style="list-style-type: none"> <li>• Number of cases: 537 women, 688 men</li> </ul>	The study confirms that OPA is protective against CC.
Italy		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>		Multivariate OR (95% CI) for CC by OPA at age 15-19 yr, men	
Case control		<ul style="list-style-type: none"> <li>• Age: 19-74 yr</li> </ul>	G1 = Highest	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.89 (0.64-1.23)</li> <li>• G3 = 0.72 (0.54-0.97)</li> <li>• G4 = 0.54 (0.40-0.74)</li> <li>• G5 = 0.47 (0.31-0.71)</li> </ul>	
D & B score = 13			G2		
			G3		
			G4		
			G5 = Lowest		
				p < 0.01	
			OPA at 30-39 yrs old		
			Q1 = Highest	Multivariate OR (95% CI) for CC by OPA at age 15-19 yr, women	
			Q2	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.73 (0.55-0.96)</li> <li>• G3 = 0.91 (0.69-1.21)</li> <li>• G4 = 0.62 (0.44-0.89)</li> </ul>	
			Q3		
			Q4 = Lowest		
			Outcome Measure: Diagnosis of CC		

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

Unconditional multiple Logistic Regression	$p < 0.05$
	Multivariate OR (95% CI) for CC by OPA at age 30-39 yr, men
	• G1 = 1.00 (referent)
	• G2 = 1.01 (0.75-1.37)
	• G3 = 0.79 (0.59-1.06)
	• G4 = 0.71 (0.52-0.97)
	• G5 = 0.64 (0.44-0.93)
	$p < 0.01$
	Multivariate OR (95% CI) for CC by OPA at age 30-39 yr, women
	• G1 = 1.00 (referent)
	• G2 = 0.65 (0.46-0.93)
	• G3 = 0.57 (0.41-0.79)
	• G4 = 0.49 (0.33-0.72)
	$p < 0.01$
	Multivariate OR (95% CI) for CC by OPA at age 50-59 yr, men
	• G1 = 1.00 (referent)
	• G2 = 1.06 (0.78-1.43)
	• G3 = 0.85 (0.63-1.14)
	• G4 = 0.68 (0.49-0.95)
	• G5 = 0.69 (0.45-1.05)
	$p < 0.01$
	Multivariate OR (95% CI) for CC by OPA at age 50-59 yr, women
	• G1 = 1.00 (referent)
	• G2 = 0.69 (0.47-1.00)
	• G3 = 0.68 (0.46-1.00)
	• G4 = 0.75 (0.47-1.20)
	$p = > 0.05$
	Multivariate OR (95% CI) for ascending CC by OPA at age 30-39 yr No significant associations for men or women
	Multivariate OR (95% CI) for transverse and descending CC by OPA at age 30-39 yr, men
	• Q1 = 1.00 (referent)
	• Q2 = 0.92 (0.51-1.67)

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

				<ul style="list-style-type: none"><li>• Q3 = 0.76 (0.43-1.37)</li><li>• Q4 = 0.46 (0.24-0.87)</li></ul> <p><i>p</i> &lt; 0.05</p> <p>Multivariate OR (95% CI) for transverse and descending CC by OPA at age 30-39 yr, women</p> <ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li><li>• Q2 = 0.51 (0.23-1.10)</li><li>• Q3 = 0.39 (0.19-0.80)</li><li>• Q4 = 0.29 (0.12-0.71)</li></ul> <p><i>p</i> &lt; 0.01</p> <p>Multivariate OR (95% CI) for sigmoid CC by OPA at age 30-39 yr, men</p> <ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li><li>• Q2 = 1.02 (0.65-1.57)</li><li>• Q3 = 0.78 (0.51-1.20)</li><li>• Q4 = 0.54 (0.34-0.85)</li></ul> <p><i>p</i> &lt; 0.01</p> <p>Multivariate OR (95% CI) for sigmoid CC by OPA at age 30-39 yr, women</p> <ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li><li>• Q2 = 0.62 (0.36-1.05)</li><li>• Q3 = 0.71 (0.44-1.15)</li><li>• Q4 = 0.58 (0.32-1.03)</li></ul> <p><i>p</i> &gt; 0.05</p>	
Thune et al 1996 [299]	To examine the association between self-reported OPA and LTPA and the subsequent risk of CC.	<ul style="list-style-type: none"><li>• n = 81,516 (53,242 men, 28,274 women)</li></ul>	16.3 year follow up	Number of cases: 236 men, 99 women	An inverse dose-response relationship between TPA and risk of CC was observed in women. In men this inverse dose-response was found only for those 45 yrs or older at study entry.
			PA assessment: Questionnaire for TPA (OPA plus recreational PA (combined))	Multivariate RR (95% CI) for total CC, men	
Norway Prospective cohort		<ul style="list-style-type: none"><li>• Sex: Men and women</li><li>• Age: 20-49 yr</li></ul>	G1 = Sedentary G2 = Moderate	<ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 1.18 (0.76-1.82)</li></ul>	
		<ul style="list-style-type: none"><li>• Characteristics: Free from cancer at baseline</li></ul>	G3 = Active	<ul style="list-style-type: none"><li>• G3 = 0.97 (0.63-1.50)</li></ul>	
D & B score = 14				<i>p</i> = 0.49	

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

	Multivariate RR (95% CI) for total CC, women
Outcome Measure: Incidence of CC	• G1 = 1.00 (referent)
Cox proportional HR	• G2 = 0.97 (0.33-2.77)
	• G3 = 0.63 (0.39-1.04)
	$p = 0.04$
	Multivariate RR (95% CI) for proximal CC, men
	• G1 = 1.00 (referent)
	• G2 = 1.16 (0.57-2.34)
	• G3 = 0.96 (0.47-1.93)
	$p = 0.64$
	Multivariate RR (95% CI) for proximal CC, women
	• G1 = 1.00 (referent)
	• G2 = 1.22 (0.51-2.94)
	• G3 = 0.62 (0.30-1.28)
	$p = 0.10$
	Multivariate RR (95% CI) for distal CC, men
	• G1 = 1.00 (referent)
	• G2 = 1.29 (0.72-2.33)
	• G3 = 0.99 (0.55-1.80)
	$p = 0.53$
	Multivariate RR (95% CI) for distal CC, women
	• G1 = 1.00 (referent)
	• G2 = 0.84 (0.32-2.17)
	• G3 = 0.61 (0.30-1.23)
	$p = 0.15$
	Multivariate RR (95% CI) for total CC, men < 45 yrs at entry
	• G1 = 1.00 (referent)
	• G2 = 2.02 (0.78-5.21)
	• G3 = 2.23 (0.88-5.66)
	$p = 0.13$
	Multivariate RR (95% CI) for total CC, women < 45 yrs at entry
	• G1 = 1.00 (referent)
	• G2 = 0.96 (0.39-2.40)

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

				<ul style="list-style-type: none"> <li>• G3 = 0.62 (0.31-1.23)</li> </ul> <p><math>p = 0.13</math></p> <p>Multivariate RR (95% CI) for total CC, men <math>\geq 45</math> yrs at entry</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.96 (0.59-1.58)</li> <li>• G3 = 0.66 (0.40-1.10)</li> </ul> <p><math>p = 0.04</math></p> <p>Multivariate RR (95% CI) for total CC, women <math>\geq 45</math> yrs at entry</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.99 (0.41-2.39)</li> <li>• G3 = 0.66 (0.33-1.33)</li> </ul> <p><math>p = 0.19</math></p>	
Vena et al 1985 [300]	To assess the relationship between lifetime OPA and the risk of CC.	<ul style="list-style-type: none"> <li>• n = 1,641 (210 cases, 1,431 control)</li> </ul>	PA assessment: Questionnaire	<ul style="list-style-type: none"> <li>• Number of cases: 210</li> </ul>	CC risk increased with increasing amount and proportion of time in jobs involving only sedentary or light work.
USA		<ul style="list-style-type: none"> <li>• Sex: Men</li> </ul>	Number of work years in jobs with sedentary or light work (yr)	OR (95% CI) by number of work years in jobs with sedentary or light work	
Case control		<ul style="list-style-type: none"> <li>• Age: 30-79 yr</li> <li>• Characteristics: Cases: admitted to hospital. Diagnosis of CC Controls: Admitted to hospital. Diagnosed with non-neoplastic non-digestive diseases</li> </ul>	G1 = None G2 = 1-20	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.49</li> </ul>	
D & B score = 15			G3 = >20	<ul style="list-style-type: none"> <li>• G3 = 1.97</li> </ul> <p>OR (95% CI) by proportion of years in jobs with sedentary or light work</p>	
			Proportion of years in jobs with sedentary or light work	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.53</li> <li>• G3 = 1.58</li> <li>• G4 = 2.10</li> </ul>	
			G1 = None G2 = 0.01-0.50 G3 = 0.41-0.99 G4 = 1.00 (referent)	OR (95% CI) by proportion of life in jobs with sedentary or light work	

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

			Proportion of life in jobs with sedentary or light work	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.66</li> <li>• G3 = 1.83</li> </ul>	
			G1 = None G2 = 0.01-0.40 G3 = 0.41-1.00 Outcome Measure: diagnosed with CC Multiple logistic regression		
Vetter et al 1992 [301]	To investigate the influence of OPA on the risk of CC in a developing country.	• n = 87 men cases, 13 women cases, 371 controls	PA assessment: Questionnaire Job title and industry names	Number of cases: 87 men, 13 women	This study presents a weak inverse association between CC and PA.
USA		• Sex: Men and women		OR (95% CI) Sitting time and CC	Only 2 of the 4 measures of activity showed evidence of an increased CC risk for sedentary jobs (time spent sitting and occupational EE) though neither was statistically significant.
Case control		• Age: 14-97 yr	Levels (Sitting time, energy expenditure	• G1 = 1.00 (referent)	
D & B score = 11		• Characteristics: Cases: Diagnosed with CC Controls: cancer cases other than colon, rectum and lung cancer.	G1 = High	• G2 = 1.0 (0.5-2.0)	
			G2 = Moderate G3 = Sedentary	• G3 = 1.5 (0.7-2.9) $p = 0.145$	
			Outcome Measure: Diagnosed with CC	OR (95% CI) Energy Expenditure and CC	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.5 (0.7-3.3)</li> <li>• G3 = 1.6 (0.8-3.6)</li> </ul>	
				$p = 0.143$	
White et al 1996 [302]	To assess the relationship between PA and CC among men and women.	• n = 871 (251 men, 193 women cases, 233 men & 194 women controls)	PA assessment: Phone interview	• Number of cases: 251 men & 193 women	The results of this study show modest support that recreational PA is associated with a reduced risk of CC.
USA			Total PA (episodes/wk)	RR (95% CI) by total PA, men	
Case control		• Sex: Men and women	G1 = 0 G2 = <1 G3 = 1-<2 G4 = 2-< 4	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.81 (0.45-1.44)</li> <li>• G3 = 0.53 (0.30-0.94)</li> <li>• G4 = 0.57 (0.33-1.00)</li> </ul>	
D & B score = 14		• Age: 30-62 yr			



**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

<p>• Characteristics: Cases: Diagnosed with CC, no previous history or CC or inflammatory bowel</p> <p>Controls: No history of CC or inflammatory bowel</p>	G5 = $\geq 4$	• G5 = 0.57 (0.40-1.11)
	Moderate-high intensity PA (episodes/wk)	$p = 0.03$
		RR (95% CI) by total PA, women
	G1 = 0	• G1 = 1.00 (referent)
	G2 = <1	• G2 = 1.17 (0.57-2.40)
	G3 = 1-<2	• G3 = 1.27 (0.65-2.45)
	G4 = $\geq 2$	• G4 = 0.59 (0.34-1.04)
		• G5 = 1.09 (0.61-1.97)
	High intensity PA (episodes/wk)	$p = 0.52$
	G1 = 0	RR (95% CI) by total PA, men and women
	G2 = <1	• G1 = 1.00 (referent)
	G3 = $\geq 1$	• G2 = 0.94 (0.60-1.47)
	METS/wk	• G3 = 0.77 (0.50-1.19)
	Q1 = 0	• G4 = 0.57 (0.39-0.85)
	Q2 = <7.30	
	Q3 = 7.30-17.88	• G5 = 0.83 (0.57-1.22)
	Q4 = $\geq 17.88$	$p = 0.04$
	Outcome Measure: Diagnosed with CC	RR (95% CI) by moderate-high intensity PA, men
	Unconditional logistic regression	• G1 = 1.00 (referent)
		• G2 = 0.84 (0.49-1.43)
		• Q3 = 0.75 (0.42-1.36)
		• Q4 = 0.66 (0.41-1.05)
		$p = 0.07$
		RR (95% CI) by moderate-high intensity PA, women
		• G1 = 1.00 (referent)
		• G2 = 1.07 (0.58-1.97)
		• G3 = 1.00 (0.51-1.98)
		• G4 = 0.74 (0.42-1.29)
		$p = 0.37$
		RR (95% CI) by moderate-high intensity PA, men and women

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.93 (0.62-1.39)</li> <li>• Q3 = 0.86 (0.55-1.34)</li> <li>• Q4 = 0.70 (0.49-1.00)</li> </ul>
	$p = 0.05$
	RR (95% CI) by high intensity PA, men
	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.85 (0.48-1.52)</li> <li>• G3 = 0.57 (0.35-0.92)</li> </ul>
	$p = 0.02$
	RR (95% CI) by high intensity PA, Women
	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.02 (0.51-2.04)</li> <li>• G3 = 0.74 (0.43-1.28)</li> </ul>
	$p = 0.31$
	RR (95% CI) by high intensity PA, men and women
	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.93 (0.59-1.44)</li> <li>• G3 = 0.64 (0.45-0.92)</li> </ul>
	$p = 0.02$
	RR (95% CI) by METs/wk, men
	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.64 (0.38-1.07)</li> <li>• Q3 = 0.59 (0.37-0.96)</li> <li>• Q4 = 0.69 (0.42-1.13)</li> </ul>
	$p = 0.05$
	RR (95% CI) by METs/wk, women
	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.87 (0.51-1.49)</li> <li>• Q3 = 1.20 (0.69-2.08)</li> <li>• Q4 = 0.74 (0.41-1.34)</li> </ul>
	$p = 0.62$
	RR (95% CI) by METs/wk, women
	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.73 (0.50-1.06)</li> <li>• Q3 = 0.80 (0.56-1.16)</li> <li>• Q4 = 0.73 (0.50-1.06)</li> </ul>
	$p = 0.08$

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

Wolin et al 2007 [303]	To assess the relationship between PA and risk of CC in women.	<ul style="list-style-type: none"> <li>• n = 79,295</li> </ul>	16 year follow-up	Number of cases: 547 (245 distal, 302 proximal) Number of dropouts: 10%	A significant inverse association exists between PA, including moderate intensity, such as walking, and risk of CC in women that is more pronounced for distal tumours.
USA		<ul style="list-style-type: none"> <li>• Sex: Women</li> <li>• Age: 40-65 yr</li> <li>• Characteristics: Nurses, no history of CC, ulcerative colitis and Crohn's disease</li> </ul>	PA assessment: Questionnaire		
Prospective cohort			Level of PA	Multivariate RR (95% CI) for distal CC by level of PA	
			G1 = <2	• G1 = 1.00 (referent)	
			G2 = 2.1-4.5	• G2 = 0.93 (0.64-1.36)	
			G3 = 4.6-10.3	• G3 = 0.99 (0.68-1.44)	
			G4 = 10.4 - 21.4	• G4 = 0.87 (0.59-1.29)	
			G5 = ≥ 21.5	• G5 = 0.54 (0.34-0.84)	
D & B score = 14		Nurses' Health Study	MPA or VPA (hr/wk)	p = 0.004	
			G1 = 0	Multivariate RR (95% CI) for proximal CC by level of PA not significant p = 0.77	
			G2 = <1		
			G3 = 1-1.9		
			G4 = 2-3.9		
			G5 = ≥ 4	Multivariate RR (95% CI) for all CC by MPA or VPA	
			Outcome Measure: Fatal and non fatal CC	• G1 = 1.00 (referent)	
				• G2 = 0.85, (0.64-1.14)	
				• G3 = 0.74 (0.53-1.04)	
				• G4 = 0.56 (0.33-0.94)	
			Cox proportional HR	p = 0.01	
				Multivariate RR (95% CI) for distal CC by MPA or VPA	
				• G1 = 1.00 (referent)	
				• G2 = 1.10 (0.73-1.66)	
				• G3 = 0.63 (0.36-1.10)	
				• G4 = 0.51 (0.22-1.17)	
				p = 0.04	
				Multivariate RR (95% CI) for proximal CC by MPA or VPA not significant p = 0.12	

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

Zhang et al 2006 [304]	To examine the relationship between LTPA and OPA and the risk of CC by anatomic site and to evaluate their joint effect on the risk of CC.	<ul style="list-style-type: none"> <li>• n = 585 cases 2,172 controls</li> </ul>	PA assessment: Questionnaire for the following variables	Number of cases: 585	Found a significant inverse association between reported LTPA and risk of CC with a slightly stronger association for the right colon than the left in both men and women.
USA		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>		Multivariate OR (95% CI) by moderate- strenuous LTPA, men and women	
		<ul style="list-style-type: none"> <li>• Age: 40-85 yr</li> </ul>	Moderate-Strenuous LTPA	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
Case control		<ul style="list-style-type: none"> <li>• Characteristics: Case: diagnosed with CC Control: no history of CC.</li> </ul>	G1 = <1 month	<ul style="list-style-type: none"> <li>• G2 = 0.7 (0.5-1.1)</li> </ul>	
			G2 = 1-4 months	<ul style="list-style-type: none"> <li>• G3 = 0.6 (0.4-0.8)</li> </ul>	
D & B score = 15			G3 = ≥ 2 weeks	<p>p = 0.003</p> <p>Multivariate OR (95% CI) by moderate- strenuous LTPA, men</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.9 (0.5-1.7)</li> <li>• G3 = 0.5 (0.3-0.9)</li> </ul> <p>p = 0.02</p> <p>Unconditional logistic regression models</p> <p>Multivariate OR (95% CI) by moderate-strenuous LTPA, women</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G3 = 0.5 (0.3-1.0)</li> <li>• G3 = 0.6 (0.4-0.9)</li> </ul> <p>p = 0.02</p> <p>Multivariate OR (95% CI) by moderate-strenuous LTPA, men and women</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.7 (0.5-1.1)</li> <li>• G3 = 0.8 (0.6-1.1)</li> </ul> <p>p = 0.53</p> <p>Multivariate OR (95% CI) by moderate-strenuous LTPA, men</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.9 (0.5-1.5)</li> </ul>	The joint effect of OPA and LTPA suggested that the risk was lowest for those with high OPA and non-OPA.

**Table 15: Studies examining the relationship between physical activity and colon cancer. (Continued)**

<p>• G3 = 0.8 (0.6-1.2)</p> <p><math>p = 0.55</math></p> <p>Multivariate OR (95% CI) by moderate- strenuous LTPA, women</p> <p>• G1 = 1.00 (referent)</p> <p>• G2 = 0.6 (0.3-1.1)</p> <p>• G3 = 0.8 (0.5-1.2)</p> <p><math>p = 0.62</math></p> <p>Multivariate OR (95% CI) by moderate- strenuous LTPA and OPA, OPA-Low</p> <p>• G1 = 1.00 (referent)</p> <p>• G2 = 0.5 (0.3-0.9)</p> <p>• G3 = 0.8 (0.5-1.2)</p> <p><math>p = 0.41</math></p> <p>Multivariate OR (95% CI) by moderate-strenuous LTPA and OPA, OPA-Medium</p> <p>• G1 = 0.7 (0.5-1.1)</p> <p>• G2 = 0.7 (0.4-1.3)</p> <p>• G3 = 0.5 (0.3-0.8)</p> <p><math>p = 0.04</math></p> <p>Multivariate OR (95% CI) by moderate-strenuous LTPA and OPA, OPA-High</p> <p>• G1 = 0.9 (0.5-1.6)</p> <p>• G2 = 0.6 (0.3-1.3)</p> <p>• G3 = 0.5 (0.3-0.8)</p>
---

D & B score, Downs and Black quality score; YR, years; PA, physical activity; OPA, occupational physical activity; kJ/min, kilojoules per minute; G, groups; MET, metabolic equivalent; HR, hazard ratio; RR, risk ratio; OR, odds ratio; 95% CI, confidence interval; LTPA, leisure-time physical activity; CC, colon cancer; TPA, total physical activity; MDA, moderate physical activity; h/d, hours per day; VPA, vigorous physical activity; h/wk, hours per week.

**Table 16 Studies examining the relationship between physical activity and breast cancer.**

Publication Country Study Design Quality Score	Objective	Population	Methods	Outcome	Comments and Conclusions
Rockhill et al 1999 [106]	To examine the effect of PA on the risk for BC.	• n = 121,701	PA assessment: Self-reported LTPA, grouped into hr/wk	3,137 cases of BC	Women who engaged in 7 or more hours per week of MVPA had a 20% lower risk of BC. An inverse dose-response relationship existed between PA and BC incidence.
USA		• Sex: Women • Age: 30-55 yr • 16-yr follow-up • Characteristics: Free of BC	G1 = <1 G2 = 1.0-1.9 G3 = 2.0-3.9	RR (95% CI) for BC and LTPA • G1 = 1.00 (referent) • G2 = 0.88 (0.79-0.98)	
Prospective cohort		• The nurses Health Study	G4 = 4.0-6.9 G5 = ≥7 Multivariate pooled logistic regression	• G3 = 0.89 (0.81-0.99) • G4 = 0.85 (0.77-0.94) • G5 = 0.82 (0.70-0.97)	
D & B score = 13				Trend $p = 0.004$	
Sesso et al 1998 [107]	To examine the association between PA and BC among postmenopausal women.	• n = 1,566	31-yr follow-up	109 cases of BC	There is an inverse relationship between PA and BC in postmenopausal women.
USA		• Sex: Women • Age: 45.5	PA assessment: Questionnaire at baseline, divided into tertiles (kcal/wk)	RR (95% CI) for BC and PA	
Prospective cohort		• Characteristics: Free of BC	T1 = <500 T2 = 500-999 T3 = ≥ 1,000	• T1 = 1.00 (referent) • T2 = 0.92 (0.58-1.45) • T3 = 0.73 (0.46-1.14)	
D & B score = 14				RR (95% CI), postmenopausal women only • T1 = 1.00 (referent) • T2 = 0.95 (0.58-1.57) • T3 = 0.49 (0.28-0.86)	
Dosemeci et al 1993 [278]	To conduct a multiple-site case-control study of 15 cancers to examine associations between PA, SES, and these cancer sites among workers.	• n = 2,643 control group	Cases: obtained from an oncological treatment center from 1979-1984	31 men had BC and 241 women had BC	This study shows the sitting-time index showed an elevated risk of female BC for sedentary jobs without SES adjustment.

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

Turkey		<ul style="list-style-type: none"> <li>• n = 2,127 men and n = 244 women)</li> <li>• Sex: Men and women</li> <li>• Characteristics: Cases - diagnosed with one of the 15 cancers being examined. Control Group - subjects diagnosed with non-cancers, cancers of the buccal cavity, esophagus, liver, bone, soft tissue, brain, lymphoma and other cancer sites for which there is no suggestion on an association with PA</li> </ul>	Adjusted SES OR (95%CI), men		
Case control		Controls: pulled from the same hospital as the cases			
D & B score = 12		PA assessment: OPA (kJ/min)	• G1 = 1.40 (0.60-3.90)		
		G1 = <8	• G2 = 1.10 (0.40-3.10)		
			• G3 = 1.00 (referent)		The slightly elevated risk of male BC was based on a small number and disappeared when the risk was adjusted for SES.
		G2 = 8-12		Trend $p = 0.34$	
		G3 = >12			
		Gart's method and Mantel's chi-square test	Adjusted SES OR (95%CI), women		
			• G1 = 1.10 (0.60-2.10)		
			• G2 = 0.90 (0.50-1.80)		
			• G3 = 1.00 (referent)		
				Trend $p = 0.23$	
Bernstein et al 1994 [305]	To determine whether young women who regularly participate in PA during their reproductive years had a reduced risk of BC.	<ul style="list-style-type: none"> <li>• n = 1,090 (545 cases; 545 controls)</li> </ul>	PA assessment: Questionnaire for overall participation in PA after menarche (h/wk), PA within 10 years after menarche (h/wk), each divided into 5 groups:	Adjusted OR (95% CI) by PA after menarche	PA may substantially reduce a women's lifetime risk of BC.
USA		<ul style="list-style-type: none"> <li>• Sex: Women</li> <li>• Age: <math>\leq 40</math> yr</li> <li>• Characteristics: White women matched for age and parity</li> </ul>		<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.95 (0.64-1.41)</li> </ul>	
Case control				• G3 = 0.65 (0.45-0.96)	
D & B score = 15			G1 = none	• G4 = 0.80 (0.54-1.17)	
			G2 = 0.1-0.7		
			G3 = 0.8-1.6	• G5 = 0.42 (0.27-0.64)	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

				<p>G4 = 1.7-3.7 G5 = <math>\geq 3.8</math> Logistic regression</p> <p>Trend <math>p = 0.0001</math> Adjusted OR (95% CI) by PA within 10 years after menarche</p> <ul style="list-style-type: none"> <li>• None = 1.00 (referent)</li> <li>• 0.1-1.2 = 0.93 (0.63-1.38)</li> <li>• 1.3-2.9 = 0.78 (0.52-1.19)</li> <li>• 3.0-5.5 = 0.69 (0.45-1.05)</li> <li>• <math>\geq 5.6 = 0.70</math> (0.47-1.06)</li> </ul> <p>Trend <math>p = 0.027</math> Adjusted OR (95% CI) by PA after menarche, nulliparous women</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.81 (0.42-1.57)</li> <li>• G3 = 0.65 (0.35-1.21)</li> <li>• G4 = 0.94 (0.53-1.67)</li> <li>• G5 = 0.73 (0.38-1.41)</li> </ul> <p>Trend <math>p = 0.43</math> Adjusted OR (95% CI) by PA after menarche, parous women</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.06 (0.65-1.74)</li> <li>• G3 = 0.65 (0.40-1.06)</li> <li>• G4 = 0.70 (0.42-1.18)</li> <li>• G5 = 0.38 (0.16-0.50)</li> </ul> <p>Trend <math>p &lt; 0.0001</math></p>	
Bernstein et al 2005 [306]	To examine the relationship between BC risk and lifetime and time- or age-specific measures of LTPA among white and black women.	<ul style="list-style-type: none"> <li>• n = 9,187 (4,538 cases; 4,649 control)</li> <li>• Sex: Women</li> <li>• Age: 35-64</li> </ul>	Cases: histologically confirmed cases of BC	4,538 cases of BC	This study supports an inverse association between PA and BC among black women and among white women.
USA		<ul style="list-style-type: none"> <li>• Ethnicity: White (including Hispanics) or Black</li> </ul>	Controls: random-digit dialing methods	Multivariate adjusted OR (95%CI) annual MET h/wk, White participants	



**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

Case control	D & B score = 13	<ul style="list-style-type: none"> <li>• Characteristics: Case Group: histologically confirmed cases of invasive BC</li> <li>• Control Group: healthy</li> </ul>	PA assessment:	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.84 (0.71-0.99)</li> </ul>	The relationship appears to be similar between black and white women.
			Questionnaire for lifetime PA (MET h/wk), divided into 5 groups	<ul style="list-style-type: none"> <li>• G3 = 0.89 (0.75-1.04)</li> <li>• G4 = 0.82 (0.69-0.97)</li> </ul>	
			G1 = Inactive	<ul style="list-style-type: none"> <li>• G5 = 0.81 (0.69-0.96)</li> </ul>	
			G2 = ≤ 2.2	Trend $p = 0.09$	
			G3 = 2.3-6.6		
			G4 = 6.7-15.1		
			G5 = ≥ 15.2		
			Unconditional logistic regression modeling	Multivariate adjusted OR (95%CI) annual MET h/wk, Black participants	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.11 (0.91-1.35)</li> <li>• G3 = 0.83 (0.67-1.03)</li> <li>• G4 = 0.79 (0.63-0.99)</li> <li>• G5 = 0.77 (0.62-0.95)</li> </ul>	
				Trend $p = 0.003$	
Carpenter et al. 1999 [307]	To examine whether lifetime exercise activity is related to BC risk in post-menopausal women.	<ul style="list-style-type: none"> <li>• n = 2,027 (1,123 case; 904 control)</li> </ul>	Cases: diagnosed with primary invasive or in situ BC	1,123 cases of BC	Strenuous exercise appears to reduce BC risk among post-menopausal women who do not gain sizable amounts of weight during adulthood.
		<ul style="list-style-type: none"> <li>• Sex: Women</li> </ul>		Multivariate adjusted OR (95%CI)	
USA		<ul style="list-style-type: none"> <li>• Age: 55-64 yr</li> <li>• Ethnicity: White (including Hispanic)</li> </ul>	Controls: individually matched to each case patient based on birth date and race	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
Case control		<ul style="list-style-type: none"> <li>• Characteristics: post-menopausal, English-speaking, born in USA, Canada or Western Europe</li> </ul>		<ul style="list-style-type: none"> <li>• G2 = 0.88 (0.72-1.07)</li> </ul>	
				<ul style="list-style-type: none"> <li>• G3 = 0.55 (0.37-0.83)</li> </ul>	
D & B score = 15			PA assessment: Questionnaire for lifetime PA (MET hr/wk), divided into 3 groups	Trend $p = 0.01$	
			G1 = no activity		
			G2 = 0.1-17.59		

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

			G3 = $\geq 17.6$		
			Conditional logistic regression		
Carpenter et al 2003 [308]	To examine the effects of obesity and lifetime exercise patterns on post-menopausal BC risk according to family history.	<ul style="list-style-type: none"> <li>• n = 3,511 (cases n = 1,883, controls) n = 1,628</li> <li>• Sex: Women</li> </ul>	PA assessment: Interview for the following PA variables	1,883 cases of BC	Exercise independent of body size seemed to exert a protective effect primarily among women with a negative family history.
USA		<ul style="list-style-type: none"> <li>• Age: 55-72</li> <li>• Characteristics: Postmenopausal Women</li> </ul>	Lifetime exercise between menarche and reference date (MET hr/wk)	Adjusted OR (95% CI) by lifetime exercise between menarche and reference date (MET hr/wk)	
Case control			G1 = 0	• G1 = 1.00 (referent)	
D & B score = 15			G2 = 0.1-3.74	• G2 = 0.85 (0.71-1.03)	
			G3 = 3.75-8.74		
			G4 = 8.75-17.59	• G3 = 0.87 (0.69-1.10)	
			G5 = $\geq 17.60$	• G4 = 1.02 (0.79-1.30)	
			Average exercise activity in 10 years prior to reference date (MET hr/wk)		
			G1 = 0	• G5 = 0.66 (0.48-0.90)	
			G2 = 0.1-6.9	Trend $p = 0.07$	
			G3 = 7.0-13.9		
				Adjusted OR (95% CI) by average exercise activity in 10 years prior to reference date (MET hr/wk)	
			G4 = 14.0-24.4		
			G5 = $\geq 24.5$		
				• G1 = 1.00 (referent)	
				• G2 = 0.93 (0.71-1.22)	
				• G3 = 0.92 (0.70-1.19)	
				• G4 = 0.86 (0.65-1.11)	
				• G5 = 0.75 (0.55-1.02)	
				Trend $p = 0.05$	
Chang et al 2006 [309]	To address the independent and combined effects of energy intake, BMI, and PA on BC incidence in women.	• n = 27,541	9.3 year follow-up (median 4.9 yr)	764 women developed BC	The study suggests that energy intake, BMI and physical inactivity are each independently and positively associated with BC risk.

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

USA		<ul style="list-style-type: none"> <li>• Sex: Women</li> <li>• Age: 55-74</li> <li>• Characteristics: no history of any cancer (nonmelanoma skin cancer patients were included in the trial)</li> </ul>	PA assessment: Questionnaire for vigorous PA (h/wk), divided into 6 groups	Multivariate adjusted RR (95% CI)	
Prospective cohort		<ul style="list-style-type: none"> <li>• Prostate, Lung, colorectal, and Ovarian Cancer Screening Trial</li> </ul>	G1 = 0		
D & B score = 13			G2 = <1	• G1 = 1.00 (referent)	
			G3 = 1	• G2 = 0.89 (0.69-1.15)	
			G4 = 2	• G3 = 0.96 (0.73-1.26)	
			G5 = 3	• G4 = 0.90 (0.70-1.16)	
			G6 = ≥4	• G5 = 1.02 (0.79-1.30)	
				• G6 = 0.78 (0.61-0.99)	
			Cox proportional HR	Trend $p = 0.153$	
Colditz et al 2003 [310]	To evaluate the relationship between PA and risk of pre-menopausal BC by type of activity and within subgroups of adiposity and oral contraceptive use.	• n = 110,468	PA assessment: Self report on 8 activities (walking or hiking, jogging (>10 min mile), running, Biking, racquet sports, lap swimming, calisthenics/aerobics other aerobic activities) to calculate MET scores (MET hrwk), divided into 5 groups:	Total cases diagnosed n = 849	These data among pre-menopausal women suggest that there is no overall association between PA and risk of BC. The findings also suggest that the effect of PA could be substantially modified by the underlying degree of adiposity.
USA		<ul style="list-style-type: none"> <li>• Sex: Women</li> <li>• Age: 25-42</li> <li>• Characteristics: pre-menopausal, no history of cancer other than nonmelanoma skin cancer</li> </ul>		Multivariate adjusted RR (95% CI)	
Prospective cohort				• G1 = 1.00 (referent)	
D & B score = 12				• G2 = 1.05 (0.82-1.34)	
				• G3 = 0.96 (0.75-1.23)	
			G1 = <3	• G4 = 1.05 (0.80-1.37)	
			G2 = 3-8.9		
			G3 = 9-17.9	• G5 = 1.07 (0.84-1.36)	
			G4 = 18-26.9		
			G5 = ≥27		
			Cox proportional HR	Trend $p = 0.69$	
Coogan et al 1997 [311]	To evaluate the effect of OPA on BC risk.	• n = 11,646 (4,863 cases and 6,783 controls).	PA assessment: Telephone interview to estimate OPA, divided into tertiles:	4,863 cases of BC	There was evidence of a graded inverse relationship between the intensity of work related activity and the incidence of BC.
USA		• Sex: Women	T1 = Sedentary	OR (95% CI)	
				• T1 = 1.00 (referent)	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

Case control		• Age: <74 yr	T2 = Medium activity jobs T3 = Heavy jobs	• T2 = 0.86 (0.77-0.97) • T3 = 0.82 (0.63-1.08)	
D & B score = 14			Logistic regression models		
Coogan and Aschengrau 1999 [312]	To evaluate the effect of OPA on BC risk.	• n = 903 (233 case; 670 control)	PA assessment: Telephone interview to estimate OPA, divided into tertiles:	233 cases of BC	There was no evidence that holding a job of medium/heavy activity reduced BC.
USA		• Sex: Women		OR (95%CI)	The study was limited by OPA misclassification and by the lack of information on LTPA.
		• Age: <50 - 80+	T1 = Exclusively sedentary	• T1 = 1.00 (referent)	
		• Ethnicity: White, Black or Other	T2 = Exclusively light	• T2 = 1.20 (0.70-1.90)	
		• Characteristics: must have worked outside the home. Cases: All incident cases of BC reported to the Massachusetts Cancer Registry from 1983 to 1986 were eligible	T3 = Exclusively medium or heavy	• T3 = 0.90 (0.40-1.90)	
Case control				Trend $p = 0.63$	
D & B score = 14			Miettinen's test-based method and Fisher's exact method		
Dallal et al 2007 [313]	To examine the relationship between LTPA and invasive and in situ BC among women.	• n = 110,599	6.6 yr follow-up	2,649 cases of invasive	The results support a protective role of strenuous long-term exercise activity against invasive and in situ BC and suggest differing effects by hormone receptor status.
USA		• Sex: Women		BC	593 cases of in situ BC
		• Age: 20-79	PA assessment: Self-reported participation in moderate and strenuous activities to estimate annual strenuous physical activity (hr/wk), divided into quintiles		
Prospective cohort	D & B score = 12	• Ethnicity: White, Black, Hispanic, Asian, American Indian or other		Multivariate adjusted RR (95% CI) for invasive BC	
		• Characteristics: California resident at baseline and no history of BC		• Q1 = 1.00 (referent)	
		• California Teachers Study cohort		• Q2 = 0.93 (0.85-1.02)	
			Q1 = 0.00-0.50		
			Q2 = 0.51-2.00	• Q3 = 0.88 (0.78-0.99)	
			Q3 = 2.01-3.50		
			Q4 = 3.51-5.00	• Q4 = 1.02 (0.88-1.18)	
			Q5 = >5	• Q5 = 0.80 (0.69-0.94)	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

			Cox proportional HR		
				Trend $p = 0.02$ Multivariate adjusted RR (95% CI) for in situ BC	
				• Q1 = 1.00 (referent) • Q2 = 0.96 (0.79-1.17) • Q3 = 0.86 (0.67-1.11) • Q4 = 0.95 (0.70-1.30) • Q5 = 0.69 (0.48-0.98)	
				Trend $p = 0.40$	
Dirx et al 2001 [314]	To evaluate the relationship between PA and BC risk with specific emphasis on interaction with other aspects of energy balance.	• n = 62,537	7.3 yr follow-up	1,208 cases of incident BC	The current study supports the hypotheses that PA is related inversely to BC risk in postmenopausal women.
Netherlands		• Sex: Women • Age: 55-69	PA assessment: Questionnaire for total recreational PA (min/day), divided into quartiles		
Case study		• Characteristics: healthy, postmenopausal		Multivariate adjusted RR (95% CI)	
D & B score = 11			Q1 = <30 Q2 = 30-60 Q3 = 61-90 Q4 = >90	• Q1 = 1.00 (referent) • Q2 = 0.84 (0.67-1.07) • Q3 = 0.78 (0.60-1.00) • Q4 = 0.76 (0.58-0.99)	
			Exponentially distributed failure time regression models		
				Trend $p = 0.003$	
Dorn et al 2003 [315]	To examine the associations between LTPA and OPA across the lifespan and pre- and post-menopausal BC.	• n = 1,550 (740 case; 810 control)	Cases: women diagnosed and histologically confirmed with BC	740 cases of BC	The study supports the hypothesis that strenuous LTPA is associated with a reduced risk of BC risk in both pre- and post menopausal women
USA		• Sex: Women		Multivariate adjusted OR (95%CI), pre- menopausal	
Case control		• Age: 40-85 • Characteristics: Case Group – histologically confirmed incidence of BC. Control Group – healthy	Controls: randomly selected and frequency matched on age and county with the cases.		
				• G1 = 1.00 (referent)	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

D & B score = 13			<ul style="list-style-type: none"> <li>• G2 = 0.94 (0.64-1.38)</li> <li>• G3 = 0.73 (0.44-1.22)</li> <li>• G4 = 1.07 (0.57-2.02)</li> </ul>	
		PA assessment: Questionnaire for lifetime strenuous PA (hr/yr)		
		G1 = 0		Trend $p = 0.82$
		G2 = 1-273		
		G3 = 274-545	Multivariate adjusted OR (95%CI), post-menopausal	
		G4 = >546		
		Logistic regression	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.85 (0.61-1.19)</li> <li>• G3 = 0.73 (0.45-1.17)</li> <li>• G4 = 0.78 (0.47-1.29)</li> </ul>	
				Trend $p = 0.19$
Drake 2001 [316]	To evaluate PA as a predictor of BC and describe BC risk factors in this sample.	<ul style="list-style-type: none"> <li>• n = 4,520</li> </ul>	PA assessment: Self-report of type, intensity, duration and frequency of walking, jogging, biking, stationary biking, swimming, dancing, racket sports, stretching, participating in other exercise, calisthenics, weight-lifting and treadmill exercises, divided into groups	150 incident cases of breast cancer
		<ul style="list-style-type: none"> <li>• Sex: Women</li> <li>• Age: 21-86</li> <li>• Characteristics: no diagnosis of BC at entry</li> </ul>		Increased frequency of a specific PA (jogging) was found to have an important protective role in BC incidence.
USA				OR (95% CI) for BC and PA
Prospective cohort				
D & B score = 11		<ul style="list-style-type: none"> <li>• Aerobic Center Longitudinal Study</li> </ul>		Activity type
				<ul style="list-style-type: none"> <li>• G1 = 1.32</li> <li>• G2 = 1.08</li> <li>• G3 = 1.35</li> </ul>
				Trend $p = 0.05$
			G1 = Aerobic (job, bike, aerobic dance)	
			G2 = Moderate (golf, walk)	
			G3 = Weight training	
			Chi-square	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

Friedenreich et al 2001 [317]	To examine the type and dose of PA and the time periods in life when PA may be specifically associated with BC risk.	• n = 2,470 (1,233 case; 1,237 control)	Cases: in situ and invasive cases of BC from 1995-1997	1,233 cases of BC	This study provides evidence that lifetime PA reduces risk of post-menopausal BC.
Canada		• Sex: Women		OR (95%CI), pre-menopausal	
Case Control		• Age: ≤ 80	Controls: matched to cases on age and place of residence	• Q1 = 1.00 (referent)	
		• Characteristics: Case Group - Alberta residents, English speaking, capable of completing an in-person interview. Control Group - no history of cancer diagnoses excluding nonmelanoma skin cancer		• Q2 = 1.15 (0.78- 1.70)	
D & B score = 13			PA assessment: Questionnaire for lifetime PA (MET hr/wk/yr), divided into quartiles by menopausal status	• Q3 = 1.15 (0.78- 1.69)	
			Pre-menopausal	• Q4 = 1.07 (0.72- 1.61)	
			Q1 = <86.6	Trend $p = 0.50$	
			Q2 = 86.6-108.3	OR (95%CI), post-menopausal	
			Q3 = 108.3-134.9	• Q1 = 1.00 (referent)	
			Q4 = ≥ 134.9	• Q2 = 0.73 (0.55- 0.98)	
			Post-menopausal	• Q3 = 0.75 (0.56- 1.00)	
			Q1 = <104.8	• Q4 = 0.70 (0.52- 0.94)	
			Q2 = 104.8-128.1	Trend $p = 0.003$	
			Q3 = 128.1-160.9		
			Q4 = ≥ 160.9		
			Logistic regression		
Friedenreich et al 2001 [318]	To examine the influence of frequency, duration, and intensity of PA on risk of BC and to compare BC risks associated with self-reported versus assigned intensity of PA.	• n = 2,470 (1,233 case; 1,237 control)	Cases: in situ and invasive cases of BC	1,233 cases of BC	This study found that moderate-intensity activities were the major contributors to the decrease in BC risk found in this study.
Canada		• Sex: Women		Multivariate adjusted OR (95% CI), pre- menopausal	
Case control		• Age: ≤ 80	Controls: matched to cases on age and place of residence	• Q1 = 1.00 (referent)	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

D & B score = 13		<ul style="list-style-type: none"><li>• Characteristics: Case Group – resident of Alberta, English speaking and able to complete an in-person interview. Control Group – free of any cancer diagnosis excluding nonmelanoma skin cancer</li></ul>	PA assessment: Questionnaire for lifetime PA questionnaire (MET hr/wk/yr), divided into quartiles	<ul style="list-style-type: none"><li>• Q2 = 1.19 (0.80- 1.76)</li></ul>	
			Q1 = <28.8	<ul style="list-style-type: none"><li>• Q3 = 1.33 (0.90- 1.96)</li></ul>	
			Q2 = 28.8-35.4	<ul style="list-style-type: none"><li>• Q4 = 1.07 (0.71- 1.62)</li></ul>	
			Q3 = 35.4-42.7	Trend $p = 0.52$	
			Q4 = $\geq 42.7$	Multivariate adjusted OR (95% CI), post- Menopausal	
			Logistic regression	<ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li><li>• Q2 = 0.85 (0.64- 1.14)</li><li>• Q3 = 0.83 (0.62- 1.10)</li><li>• Q4 = 0.69 (0.51- 0.93)</li></ul>	
				Trend $p = 0.006$	
Friedenreich and Rohan 1995 [319]	To describe the association between LTPA and BC.	<ul style="list-style-type: none"><li>• n = 902 (451 case; 451 control)</li></ul>	Cases: first diagnosis of BC in 1982 and 1984	Adjusted OR (95%CI), pre-menopausal	This study found some evidence (of borderline statistical significance) that recreational PA is associated with decreased risk of BC.
Australia		<ul style="list-style-type: none"><li>• Sex: Women</li></ul>		<ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li></ul>	
Case control		<ul style="list-style-type: none"><li>• Age: 20-74 yr</li></ul>	Controls: Randomly selected from the electoral roll, matched on date of birth to each case	<ul style="list-style-type: none"><li>• Q2 = 0.77 (0.36- 1.65)</li></ul>	
D & B score = 13		<ul style="list-style-type: none"><li>• Characteristics: Australian women</li></ul>	PA assessment: Self reported PA (kcal/wk), divided into quartiles	<ul style="list-style-type: none"><li>• Q3 = 0.48 (0.22- 1.03)</li></ul>	
			Q1 = 0	<ul style="list-style-type: none"><li>• Q4 = 0.60 (0.30- 1.17)</li></ul>	
			Q2 = 1-2,000	Trend $p = 0.09$	
			Q3 = 2000-4000	Adjusted OR (95%CI), post-menopausal	
			Q4 = >4000	<ul style="list-style-type: none"><li>• Q1 = 1.00 (referent)</li></ul>	
			Logistic regression models	<ul style="list-style-type: none"><li>• Q2 = 0.74 (0.46- 1.18)</li></ul>	
				<ul style="list-style-type: none"><li>• Q3 = 0.88 (0.53- 1.48)</li></ul>	
				<ul style="list-style-type: none"><li>• Q4 = 0.73 (0.44- 1.20)</li></ul>	
				Trend $p = 0.32$	



**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

Gammon et al 1998 [320]	To examine the association between LTPA and BC among young women.	• n = 3,173 (1,668 case; 1,505 control)	Cases: women diagnosed with BC between 1990-1992	1,668 cases of BC	The study's data does not support the hypothesis of a reduced risk of BC among young women with increased recreational PA in adolescence, young adulthood or during the year prior to the interview, or with the average PA over the three time periods
USA		• Sex: Women		Multivariate adjusted OR (95%CI)	
Case control		• Age: <45	Controls: were matched to cases by age group and geographic center	• Q1 = 1.00 (referent)	
D & B score = 13		• Characteristics: Case Group – diagnosed with invasive or in situ BC. Control Group - healthy	PA assessment: Questionnaire for recreational PA, for ages 12- 13 yr, age 20 yr and 1 year prior to the interview. Divided into quartiles MET score Q1 = 1.62-18.07 Q2 = 18.08-30.00 Q3 = 30.01-42.95 Q4 = 42.96-98.00 Logistic regression	• Q2 = 0.79 (0.63- 0.98)  • Q3 = 0.98 (0.79- 1.22) • Q4 = 1.01 (0.81- 1.25) Trend $p = 0.42$	
Gilliland et al 2001 [321]	To investigate the relationship of PA with BC risk in Hispanic and non- Hispanic White women	• n = 1,556 (712 case; 844 control)	Cases: diagnosed with BC between 1992-1994	712 cases of BC	Hispanic and non- Hispanic women with high PA during non-OPA were at substantially reduced risk of BC.
USA		• Sex: Women		Adjusted OR (95%CI), pre-menopausal Hispanic	
Case control		• Age: between 35-74 at diagnosis	Controls: matched on ethnicity, age and seven health planning districts	• G1 = 1.00 (referent)	
D & B score = 13		• Ethnicity: Hispanic and non-Hispanic White • Characteristics: Case Group – diagnosed with in situ or invasive BC and residents of New Mexico at time of diagnosis. Control Group – healthy	PA assessment: Self-reported non-OPA (MET hr/wk score) G1 = <25  G2 = 25-50 G3 = 50-80 G4 = ≥ 80 Logistic regression	• G2 = 1.17 (0.53- 2.55)  • G3 = 0.49 (0.22- 1.07) • G4 = 0.29 (0.12- 0.72) Trend $p < 0.001$ Adjusted OR (95%CI), pre-menopausal non- Hispanic • G1 = 1.00 (referent) • G2 = 1.35 (0.64- 2.85)	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

				<ul style="list-style-type: none"> <li>• G3 = 1.44 (0.67- 3.10)</li> <li>• G4 = 1.13 (0.49- 2.61)</li> </ul> <p>Trend <math>p = 0.741</math> Adjusted OR (95%CI), post-menopausal Hispanic</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.74 (0.40- 1.36)</li> <li>• G3 = 0.37 (0.18- 0.75)</li> <li>• G4 = 0.38 (0.18- 0.77)</li> </ul> <p>Trend <math>p = 0.002</math> Adjusted OR (95%CI), post-menopausal non- Hispanic</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.45 (0.26- 0.78)</li> <li>• G3 = 0.49 (0.28- 0.86)</li> <li>• G4 = 0.45 (0.24- 0.85)</li> </ul> <p>Trend <math>p = 0.019</math></p>	
Hsing et al 1998 [322]	To evaluate the role of selected demographic, lifestyle, and anthropometric factors in the risk for male BC.	<ul style="list-style-type: none"> <li>• n = 690 (178 case; 512 control)</li> </ul>	Cases: selected from 18,733 decedents included in the 1986 NMFS conducted by the US	178 cases of BC	This study suggests that obesity increases the risk of male BC, possibly through hormonal mechanisms, while dietary factors, PA and SES indicators also deserve further investigation.
USA		<ul style="list-style-type: none"> <li>• Sex: Men</li> </ul>	National Center for Health Statistics (NCHS)	Adjusted OR (95%CI)	
Case control		<ul style="list-style-type: none"> <li>• Age: 25-74</li> <li>• Ethnicity: Black and White</li> <li>• Characteristics: Case Group – deceased. Control Group – dying (or deceased) of causes other than BC</li> </ul>	Controls: selected from male decedents dying of causes other than BC	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.60 (0.30- 1.10)</li> <li>• G3 = 1.30 (0.80- 2.00)</li> </ul>	
D & B score = 12			PA assessment: Questionnaire (frequency and intensity), divided into groups G1 = Regular G2 = Irregular G3 = Hardly any Logistic regression analysis		
Hu et al 1997 [323]	To study breast cancer focusing on breast-feeding, body weight, and PA as well as reproductive histories on pre- and post-menopausal Japanese women.	<ul style="list-style-type: none"> <li>• n = 526 (157 case; 369 control)</li> </ul>	Cases: Histologically confirmed cases of BC from 1989-1993.	157 cases of BC	Reduced risk of pre- menopausal BC was associated with high EE in PA during teenage years, although the trend was not statistically significant.

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

Japan		• Sex: Women		Unadjusted RR (95%CI), pre-menopausal	
Case control		• Age: 26-75		• G1 = 1.00 (referent)	
D & B score = 13		• Characteristics: Case Group – histologically confirmed cases of BC and resident of Gifu prefecture at time of diagnosis. Control Group – no breast disease or hormone-related (ovarian, endometrial and thyroid) cancers	Controls: individuals who had the screening test for BC during the same period	• G2 = 0.74 (0.38- 1.44)	
			PA assessment: Questionnaire for TPA (kcal/wk), divided into groups	• G3 = 1.01 (0.54- 1.87)	
			G1 = 0	Trend $p = 0.876$	
			G2 = 1-649	Unadjusted RR (95%CI), post-menopausal:	
			G3 = $\geq 650$	• G1 = 1.00 (referent)	
				• G2 = 1.53 (0.69- 3.54)	
			Logistic regression models	• G3 = 1.39 (0.61- 3.13)	
John et al 2003 [324]	To examine BC risk in relation to lifetime histories of MPA and VPA including LTPA, transportation household and outdoor chores, and OPA in a multiethnic population.	• n = 2,870 (1,277 case; 1,593 control)	Cases: diagnosed between 1995-1998	1,277 cases of BC	This study supports previous reports of a reduced risk of BC in physically active women.
USA		• Sex: Women		Multivariate adjusted OR (95%CI), pre- menopausal Latinas	
Case control		• Age: 35-79		• G1 = 1.00 (referent)	
D & B score = 12		• Ethnicity: Latina, African-American and White	Controls: randomly selected according race/ethnicity and age distribution of cases	• G2 = 0.84 (0.49- 1.45)	
			PA assessment: In-person interview for lifetime PA (hr/wk), divided into groups	• G3 = 0.73 (0.42- 1.28)	
			Pre-menopausal	Multivariate adjusted OR (95%CI), pre- menopausal African Americans	
			G1 = <9.1	• G1 = 1.00 (referent)	
			G2 = 9.1-20.7	• G2 = 1.00 (0.55- 1.84)	
			G3 = $\geq 20.7$	• G3 = 0.68 (0.35- 1.34)	
			Post-menopausal		
			G1 = <9.6		
			G2 = 9.6-21.6		
			G3 = $\geq 21.7$		

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

			Logistic regression modeling	<p>Multivariate adjusted OR (95%CI), pre- menopausal Whites</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.82 (0.42- 1.58)</li> <li>• G3 = 0.76 (0.36- 1.61)</li> </ul> <p>Multivariate adjusted OR (95%CI), post- menopausal Latinas</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.82 (0.55- 1.24)</li> <li>• G3 = 0.81 (0.54- 1.22)</li> </ul> <p>Multivariate adjusted OR (95%CI), post- menopausal African Americans</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.78 (0.52- 1.17)</li> <li>• G3 = 0.71 (0.47- 1.07)</li> </ul> <p>Multivariate adjusted OR (95%CI), post- menopausal Whites</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.94 (0.64- 1.37)</li> <li>• G3 = 0.91 (0.60- 1.41)</li> </ul>	
Kruk 2007 [325]	To examine the association between all types of PA and BC risk among Polish women.	<ul style="list-style-type: none"> <li>• n = 590 (268 case; 322 control)</li> </ul>	PA assessment: Questionnaire for lifetime PA (MET hr/wk/yr), divided into groups	268 cases of BC	The results of this study provide evidence of an inverse association between PA and the risk of BC.
Poland		<ul style="list-style-type: none"> <li>• Sex: Women</li> </ul>	G1 = <110	Multivariate adjusted OR (95%CI), pre- menopausal	
Case control		<ul style="list-style-type: none"> <li>• Age: 35-75 yr</li> </ul>	G2 = 110-150	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
D & B score = 13		<ul style="list-style-type: none"> <li>• Characteristics: Polish women. Cases: identified from the Szczecin Regional Cancer Registry. Controls: matched on age and place of residence</li> </ul>	G3 = >150	<ul style="list-style-type: none"> <li>• G2 = 0.45 (0.14- 1.44)</li> </ul>	
			Logistic regression analysis	<ul style="list-style-type: none"> <li>• G3 = 0.44 (0.14- 1.37)</li> </ul> <p>Trend <math>p = 0.42</math></p> <p>Multivariate adjusted OR (95%CI), post- menopausal</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.60 (0.33- 1.09)</li> <li>• G3 = 0.31 (0.21- 0.70)</li> </ul> <p>Trend <math>p = 0.002</math></p>	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

Kruk 2007 [326]	To examine the relationship between LTPA and BC risk.	• n = 822 (cases n = 257, control n = 565)	PA assessment: Questionnaire for LTPA (METs), divided into groups	Adjusted OR (95% CI)	The findings provide further support to the hypothesis that increased LTPA throughout life is associated with a decreased risk of BC.
Poland		• Sex: Women	G1 = Low	• G1 = 1.00 (referent)	
Case control		• Age: 35-93 yr	G2 = Medium	• G2 = 0.57 (0.36- 0.89)	
D & B score = 13			G3 = High	• G3 = 0.22 (0.14- 0.35)	
				Trend $p < 0.0001$	
Lahmann et al 2007 [327]	To examine the association of PA with pre- and post-menopausal BC risk.	• n = 218,169	Baseline and 6.4 year follow-up	3,423 cases of BC	Increasing PA reduces BC risk.
Europe (9 countries)		• Sex: Women		Multivariate adjusted HR (95% CI) by TPA, pre-menopausal	
Prospective cohort		• Age: 20-80		• Q1 = 1.00 (referent)	
D & B score = 12		• The European Prospective Investigation into Cancer and nutrition study	PA assessment: Interviews and questionnaire for TPA and recreational PA, each divided into quartiles	• Q2 = 1.02 (0.84- 1.24)	
			TPA Index	• Q3 = 0.84 (0.68- 1.04)	
			Q1 = Inactive	• Q4 = 1.02 (0.77- 1.36)	
			Q2 = Moderately inactive	Trend $p = 0.267$	
			Q3 = Moderately active	Multivariate adjusted HR (95% CI) by TPA, Post-menopausal	
			Q4 = Active	• Q1 = 1.00 (referent)	
			Recreational PA (MET hr/wk)	• Q2 = 0.89 (0.79- 1.00)	
			Q1 = <14	• Q3 = 0.84 (0.74- 0.96)	
			Q2 = 14-24	• Q4 = 0.92 (0.76- 1.12)	
			Q3 = 25-42	Trend $p = 0.06$	
			Q4 = >42	Multivariate adjusted HR (95% CI) by recreational PA, pre-menopausal	
			Cox proportional index	• Q1 = 1.00 (referent)	
				• Q2 = 0.91 (0.75- 1.10)	
				• Q3 = 0.95 (0.78- 1.14)	
				• Q4 = 0.94 (0.76- 1.15)	
				Trend $p = 0.580$	
				Multivariate adjusted HR (95% CI) by recreational PA, post-menopausal	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 1.05 (0.94- 1.17)</li> <li>• Q3 = 0.92 (0.83- 1.03)</li> <li>• Q4 = 0.96 (0.85- 1.08)</li> </ul>	
				Trend $p = 0.176$	
Lee et al 2001 [328]	To examine the association between PA and BC risk.	• n = 39,322	Baseline and 4 year follow- up	411 cases of BC	The data suggest that PA during middle age and older is not uniformly associated with decreased BC risk. Among post-menopausal women only, higher levels of PA may decrease the risk of BC.
USA		<ul style="list-style-type: none"> <li>• Sex: Women</li> <li>• Age: <math>\geq 45</math> yr</li> </ul>		Multivariate adjusted RR (95% CI) by PA, all women	
Prospective cohort		• Characteristics: Healthy women	PA assessment: Questionnaire	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> </ul>	
D & B score = 12		Women's Health Study	PA (kJ/wk), divided into quartiles	<ul style="list-style-type: none"> <li>• Q2 = 1.04 (0.77- 1.40)</li> <li>• Q3 = 0.86 (0.64- 1.17)</li> <li>• Q4 = 0.80 (0.58- 1.12)</li> </ul>	
			Q1 = <840		
			Q2 = 840-2519		
			Q3 = 2520-6299	Trend $p = 0.11$	
			Q4 = $\geq 6300$	Multivariate adjusted RR (95% CI) by PA, post- menopausal only	
			VPA (kJ/wk), divided into quintiles	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.97 (0.68- 1.39)</li> <li>• Q3 = 0.78 (0.54- 1.12)</li> <li>• Q4 = 0.67 (0.44- 1.02)</li> </ul>	
			Q1 = none		
			Q2 = 1-839		
			Q3 = 840-2099		
			Q4 = 2100-4199	Trend $p = 0.03$	
			Q5 = $\geq 4200$	Multivariate adjusted RR (95% CI) by VPA, all women	
			Proportional hazard regression	<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 1.02 (0.70- 1.48)</li> <li>• Q3 = 1.11 (0.78- 1.58)</li> <li>• Q4 = 0.97 (0.66- 1.44)</li> <li>• Q5 = 0.98 (0.69- 1.40)</li> </ul>	
				Trend $p = 0.98$	
				Multivariate adjusted RR (95% CI) by VPA, post- menopausal only	
				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.93 (0.57- 1.50)</li> </ul>	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

				<ul style="list-style-type: none"> <li>• Q3 = 0.91 (0.57- 1.47)</li> <li>• Q4 = 0.93 (0.57- 1.50)</li> <li>• Q5 = 0.76 (0.47- 1.24)</li> </ul>	
				Trend $p = 0.29$	
Magnusson et al 2005 [329]	To report the relationship between pre-menopausal BC, body fatness at age 10 years and in adulthood, and sports participation during puberty, late adolescence and early adulthood from three related case-control studies.	<ul style="list-style-type: none"> <li>• n = 3,108 (1,560 cases; 1,548 controls)</li> </ul>	PA assessment: Interview for sports participation (h/wk in the following age categories (12-14 yr, 16-18 yr, 20-30 yr, 12-30 yr, around age of diagnosis)	Adjusted RR (95% CI), 12-14 yr	An inverse association between body fatness but not PA at a young age and the risk of BC in pre-menopausal women.
UK		<ul style="list-style-type: none"> <li>• Sex: Women</li> </ul>		<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
Case control		<ul style="list-style-type: none"> <li>• Age: Study 1 = 36 yr, study 2 = 36-45 yr, study 3 = 46-54 yr</li> </ul>		<ul style="list-style-type: none"> <li>• G2 = 1.04 (0.93- 1.17)</li> </ul>	
D & B score = 13		<ul style="list-style-type: none"> <li>• Characteristics: White women with no previous malignancy, mental handicap or illness</li> </ul>	Sports participation (h/wk)	<ul style="list-style-type: none"> <li>• G3 = 1.03 (0.93- 1.14)</li> </ul>	
			G1 = 0-1	Trend $p = 0.95$	
			G2 = 2-3	Adjusted RR (95% CI), 16-18 yr	
			G3 = $\geq 4$	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.95 (0.83- 1.09)</li> <li>• G3 = 0.89 (0.79- 1.02)</li> </ul>	
				Trend $p = 0.20$	
				Adjusted RR (95% CI), 20-30 yr	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.90 (0.76- 1.08)</li> <li>• G3 = 1.01 (0.81- 1.26)</li> </ul>	
				Trend $p = 0.73$	
				Adjusted RR (95% CI), 12-30 yr	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.99 (0.89- 1.11)</li> <li>• G3 = 1.01 (0.88- 1.16)</li> </ul>	
				Trend $p = 0.94$	
				Adjusted RR (95% CI), around age of diagnosis	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.84 (0.71- 1.00)</li> <li>• G3 = 1.06 (0.86- 1.32)</li> </ul>	
				Trend $p = 0.82$	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

Malin et al 2005 [330]	To evaluate a pattern of behavioral exposures indicating positive energy balance would be associated with increased BC risk.	• n = 3,015 (1,459 cases; 1,556 control)	PA assessment: Questionnaire for PA (MET hr/d/yr), divided into groups	OR (95% CI)	The study suggests that promotion of behavioral patterns that optimize energy balance maybe a viable option for BC prevention.
China		• Sex: Women	G1 = 0	• G1 = 1.86 (1.44- 2.41)	
Case control		• Age: Mean ~47 yr	G2 = 0.1-2.92	• G2 = 1.33 (0.96- 1.83)	
D & B score = 12		• Characteristics: Residents of urban Shanghai Shanghai Breast Cancer Study	G3 = >2.92	• G3 = 1.00 (referent)	
Margolis et al 2005 [331]	To study the association between PA and incident invasive BC.	• n = 99,504	Baseline and 9.1 year follow-up	1,166 cases of BC	No evidence of a protective effect of PA on BC risk was found.
Norway/Sweden		• Sex: Women			
Prospective cohort		• Age: 30-49 (mean 41 yr)	PA assessment: Questionnaire for PA using a 5 point scale and for competitive PA (years of participation), each divided into groups	Multivariate adjusted RR (95% CI) by PA level, at enrollment	
D & B score = 13		• The Norwegian- Swedish Women's Lifestyle and Health Study	PA level (5 point scale)	• G1 = 1.00 (referent)	
			G1 = None	• G2 = 1.35 (0.96- 1.90)	
			G2 = Low	• G3 = 1.26 (0.91- 1.74)	
			G3 = Moderate	• G4 = 1.19 (0.85- 1.67)	
			G4 = High	• G5 = 1.24 (0.85- 1.82)	
			G5 = Vigorous	Trend $p = 0.85$	
			Competitive PA (years)	Multivariate adjusted RR (95% CI) by PA level, at age 30	
			G1 = None	• G1 = 1.00 (referent)	
			G2 = 1-4	• G2 = 1.03 (0.64- 1.66)	
			G3 ≥ 5	• G3 = 1.16 (0.74- 1.81)	
				• G4 = 1.06 (0.67- 1.68)	
				• G5 = 1.20 (0.77- 1.95)	
				Trend $p = 0.60$	
				Multivariate adjusted RR (95% CI) by PA level, at age 14	
				• G1 = 1.00 (referent)	
				• G2 = 0.93 (0.62- 1.39)	
				• G3 = 0.94 (0.65- 1.35)	
				• G4 = 1.07 (0.73- 1.55)	
				• G5 = 1.05 (0.72- 1.54)	
				Trend $p = 0.14$	



**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

				<p>Multivariate adjusted RR (95% CI) by years of competitive PA</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.21 (0.95- 1.54)</li> <li>• G3 = 0.95 (0.75- 1.19)</li> </ul> <p>Trend <math>p = 0.96</math></p>	
McTiernan et al 1996 [332]	To investigate the relationship between LTPA and BC.	<ul style="list-style-type: none"> <li>• n = 1,029 (cases n = 537, controls n = 492)</li> </ul>	PA assessment: Questionnaire (Minnesota LTPA Questionnaire) for LTPA (hr/wk), divided into groups	Adjusted OR (95% CI) by LTPA during adulthood, all ages and menopausal status	The results indicate a weak negative association between PA and risk of BC in middle-aged women.
USA		<ul style="list-style-type: none"> <li>• Sex: Women</li> </ul>	G1 = None	<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> </ul>	
Case control		<ul style="list-style-type: none"> <li>• Age: 50-64</li> </ul>	G2 = 0.1-1.5	<ul style="list-style-type: none"> <li>• G2 = 1.1 (0.7-1.6)</li> </ul>	
D & B score = 13			G3 = 1.6-2.5	<ul style="list-style-type: none"> <li>• G3 = 0.7 (0.4-1.1)</li> </ul>	
			G4 = 2.6-3.5	<ul style="list-style-type: none"> <li>• G4 = 0.7 (0.4-1.1)</li> </ul>	
			G5 = 3.6-5.0	<ul style="list-style-type: none"> <li>• G5 = 0.6 (0.4-0.9)</li> </ul>	
			G6 = >5	<ul style="list-style-type: none"> <li>• G6 = 1.1 (0.7-1.6)</li> </ul>	
			Calculated categories of EE (total time x intensity code)	Trend $p = 0.29$	
			G1 = Lowest	Adjusted OR (95% CI) by LTPA during adulthood, aged $\geq 55$ yr, post-menopausal only	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.8 (0.5-1.3)</li> <li>• G3 = 0.5 (0.3-0.9)</li> <li>• G4 = 0.6 (0.4-1.1)</li> <li>• G5 = 0.4 (0.2-0.8)</li> <li>• G6 = 0.8 (0.5-1.3)</li> </ul>	
			G6 = Highest	Trend $p = 0.03$	
				Adjusted OR (95% CI) by category of total EE in adulthood, all ages and menopausal status	
				<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 1.2 (0.8-2.0)</li> <li>• G3 = 0.9 (0.6-1.3)</li> <li>• G4 = 0.6 (0.4-0.9)</li> <li>• G5 = 0.9 (0.6-1.5)</li> <li>• G6 = 0.9 (0.6-1.4)</li> </ul>	
				Trend $p = 0.25$	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

			Adjusted OR (95% CI) by category of total EE in adulthood, aged ≥ 55 yr, post-menopausal only		
			<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.8 (0.4-1.4)</li> <li>• G3 = 0.7 (0.4-1.2)</li> <li>• G4 = 0.5 (0.3-0.8)</li> <li>• G5 = 0.8 (0.5-1.3)</li> <li>• G6 = 0.6 (0.4-1.0)</li> </ul>		
			Trend $p = 0.009$		
McTiernan et al 2003 [333]	To examine the association between current and past LTPA and incidence of BC in post-menopausal women.	• n = 74,171	Baseline and mean follow-up of 4.7 years	1,780 cases of BC	Increased PA is associated with reduced risk for BC in post-menopausal women.
USA		• Sex: Women	PA assessment: Questionnaire for TPA (MET hr/wk), moderate or strenuous PA (hr/wk) and strenuous PA (hr/wk), each divided into groups	Adjusted RR (95% CI) by TPA	Longer duration provides the most benefit however need not be strenuous.
Prospective cohort		• Age: 50-79	TPA (MET hr/wk)	• G1 = 1.00 (referent)	
D & B score = 13		• Characteristics: Women from the Women's Health Initiative Observational Study	G1 = none	• G2 = 0.90 (0.77-1.07)	
			G2 = 0-5.0	• G3 = 0.82 (0.68-0.97)	
			G3 = 5.1-10.0	• G4 = 0.89 (0.76-1.00)	
			G4 = 10.1-20.0	• G5 = 0.83 (0.70-0.98)	
			G5 = 20.1-40	• G6 = 0.78 (0.62-1.00)	
			G6 = ≥ 40.0	Trend $p = 0.03$	
			Moderate or strenuous PA (hr/wk)	Adjusted RR (95% CI) by TPA, BMI ≤ 24.13	
			G1 = none	• G1 = 1.00 (referent)	
			G2 = ≤ 1	• G2 = 0.78 (0.57-1.10)	
			G3 = 1.1-2.0	• G3 = 0.70 (0.51-0.97)	
			G4 = 2.1-3.0	• G4 = 0.80 (0.60-1.10)	
			G5 = 3.1-4.0	• G5 = 0.68 (0.51-0.92)	
			G6 = 4.1-7.0	• G6 = 0.63 (0.43-0.93)	
			G7 = >7.0	Trend $p = 0.03$	
			Strenuous PA (hr/wk)	Adjusted RR (95% CI) by TPA, BMI 24.14-28.44	
			G1 = none	• G1 = referent	
			G2 = ≤ 1.0		

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

				<p>G3 = 1.1-2.0</p> <p>G4 = 2.1-4.0</p> <p>G5 = &gt;4.0</p> <p>Cox proportional hazard ratio</p> <p>G2 = 0.72 (0.53-0.98)</p> <p>G3 = 0.78 (0.57-1.10)</p> <p>G4 = 0.77 (0.58-1.00)</p> <p>G5 = 0.85 (0.64-1.10)</p> <p>G6 = 0.78 (0.52-1.20)</p> <p>Trend <math>p = 0.74</math></p> <p>Adjusted RR (95% CI) by TPA, BMI &gt;28.44</p> <p>G1 = 1.00 (referent)</p> <p>G2 = 1.10 (0.88-1.50)</p> <p>G3 = 0.90 (0.67-1.20)</p> <p>G4 = 1.00 (0.79-1.30)</p> <p>G5 = 0.89 (0.65-1.20)</p> <p>G6 = 0.94 (0.57-1.60)</p> <p>Trend <math>p = 0.30</math></p> <p>Adjusted RR (95% CI) by current moderate or strenuous PA</p> <p>G1 = 1.00 (referent)</p> <p>G2 = 0.92 (0.78-1.10)</p> <p>G3 = 0.91 (0.79-1.10)</p> <p>G4 = 0.94 (0.81-1.10)</p> <p>G5 = 0.99 (0.83-1.20)</p> <p>G6 = 0.91 (0.78-1.10)</p> <p>G7 = 0.79 (0.63-0.99)</p> <p>Trend <math>p = 0.12</math></p> <p>Adjusted RR (95% CI) by current strenuous PA</p> <p>G1 = 1.00 (referent)</p> <p>G2 = 0.94 (0.80-1.10)</p> <p>G3 = 0.95 (0.80-1.10)</p> <p>G4 = 0.93 (0.78-1.10)</p> <p>G5 = 0.91 (0.67-1.20)</p> <p>Trend <math>p = 0.25</math></p>	
Navarro Silvera et al 2006 [334]	To study the independent and combined associations of VPA, energy consumption and BMI with risk of subsequent BC.	<ul style="list-style-type: none"> <li>n = 40,318 in analysis (49,613 prior to exclusion)</li> </ul>	Baseline and 16.4 year follow-up	1,673 cases of BC from the 40,318 included in the analysis (2,545 cases from total prior to exclusion)	The results of the study suggest that BC risk may vary according to various combinations of the components of energy balance.
Canada		<ul style="list-style-type: none"> <li>Sex: Women</li> </ul>	PA assessment: Questionnaire by VPA (min/d), divided into groups	Adjusted HR (95% CI) by VPA	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

Prospective cohort D & B score = 13		• Age: 40-59	G1 = none	• G1 = 1.00 (referent)	
		• Characteristics: Canadian women with no history of BC	G2 = Any	• G2 = 0.98 (0.85-1.13)	
		• National Breast Screening Study (NBSS)	G3 = 0-30	• G3 = 1.06 (0.88-1.27)	
			G4 = 30-60	• G4 = 0.98 (0.83-1.16)	
			G5 > 60	• G5 = 0.93 (0.78-1.10)	
			Cox proportional hazard ratio	Trend $p = 0.38$	
				Adjusted HR (95% CI) by VPA, pre-menopausal	
				• G1 = 1.00 (referent)	
				• G2 = 0.91 (0.75-1.10)	
				• G3 = 1.02 (0.80-1.31)	
		• G4 = 0.88 (0.70-1.11)			
		• G5 = 0.87 (0.68-1.09)			
		Trend $p = 0.23$			
		Adjusted HR (95% CI) by VPA, post- menopausal			
		• G1 = 1.00 (referent)			
		• G2 = 1.06 (0.87-1.30)			
		• G3 = 1.08 (0.81-1.42)			
		• G4 = 1.11 (0.87-1.41)			
		• G5 = 1.00 (0.78-1.29)			
		Trend $p = 0.96$			
Patel et al 2003 [335]	To examine the association between various measures of PA and post-menopausal BC risk.	• n = 72,608	Baseline and 5 year follow-up	1,520 cases of breast cancer	The study shows a lower risk of post- menopausal BC is associated with regular PA.
USA		• Sex: Women	PA assessment: Questionnaire for LTPA (METs hr/wk) at various times during life, divided into groups	Adjusted RR (95% CI), LTPA at study entry	
Prospective cohort D & B score = 14		• Age: 50-74	G1 = none	• G1 = 0.86 (0.70- 1.04)	
		• Characteristics: Postmenopausal women	G2 = 0.1-6.9	• G2 = 1.00 (referent)	
		• The American Cancer Society Cancer Prevention Study II (CPS-II) Nutritional Cohort	G3 = 7.0-17.5	• G3 = 0.92 (0.81- 1.04)	
			G4 = 17.6-31.5	• G4 = 0.94 (0.81- 1.09)	
			G5 = 31.6-42.0	• G5 = 0.77 (0.56- 1.06)	
			G6 = >42.0	• G6 = 0.71 (0.49- 1.02)	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

			LTPA at 10 years prior to study, calculated MET score and categorized into groups: None Slight Moderate Heavy Cox proportional hazard ratio	Trend $p = 0.08$ (among active women $p = 0.03$ )  Adjusted RR (95% CI), LTPA at age 40 yr • G1 = 1.03 (0.88- 1.21) • G2 = 1.00 (referent) • G3 = 1.05 (0.92- 1.20) • G4 = 1.01 (0.87-1.18) • G5 = 1.16 (0.92- 1.46) • G6 = 0.79 (0.61- 1.03)  Trend $p = 0.31$ (among active women $p = 0.36$ ) Adjusted RR (95% CI), LTPA at 10 years prior to study entry • None = 0.80 (0.51- 1.25) • Slight = 1.00 (referent) • Moderate = 0.93 (0.83- 1.04) • Heavy = 0.87 (0.68- 1.13)  Trend $p = 0.32$ (among active women, trend $p = 0.16$ )	
Patel et al 2003 [336]	To evaluate the association between lifetime LTPA and BC risk.	• n = 1,183 (cases n = 616) n = 567, controls n = 616)	PA assessment: Calendar reporting for lifetime exercise activity (MET h/wk), divided into groups	Adjusted OR (95% CI)	The findings suggest that PA may modify the risk of in situ BC particularly in women without a family history of BC.
USA Case control D & B score = 14		• Sex: Women • Age: 35-64 • Characteristics: White and Black women	G1 = None G2 = 0.0-3.0 G3 = 3.0-8.0  G4 = 8.0-16.0 G5 = 16.0-32.0 G6 = >32.0 Unconditional logistical regression	• G1 = 1.00 (referent) • G2 = 0.70 (0.48- 1.03) • G3 = 0.65 (0.44- 0.96)  • G4 = 0.61 (0.41- 0.92) • G5 = 0.63 (0.40- 0.98) • G6 = 0.65 (0.39- 1.08) Trend $p = 0.27$ (among exercisers only $p = 0.81$ )	
Rintala et al 2002 [337]	To obtain an estimate of BC incidence in association with self-rated OPA.	• n = 680,000	PA assessment: Self-reported OPA in 5 classes (1=low, 5=high)	17,986 cases of BC	The results support the hypothesis that OPA, if high enough, markedly reduced BC risk.

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

Finland		• Sex: Women	Class 1 = Jobs sitting and light hand tasks	Adjusted RR (95% CI), age 25-39 years	
Prospective cohort		• Age: Women born in 1930-1969	Class 2 = Handling of heavier items (conveyor belt)	• C1+2 = 1.00 (referent)	
D & B score = 11		• Characteristics: Finish women	Class 3 = Jobs involving body motion	• C3 = 0.99 (0.85- 1.17)	
			Class 4 = Jobs involving walking up stairs or long distances, bending and carrying	• C4 = 0.90 (0.76- 1.07)	
			Class 5 = Same as class 4 except heavy tasks were performed for most of the day	• C5 = 0.68 (0.51- 1.93)	
			Poisson regression models	Trend	
				Adjusted RR (95% CI), age 40-54 years	
				• C1+2 = 1.00 (referent)	
				• C3 = 1.02 (0.94- 1.11)	
				• C4 = 0.99 (0.91- 1.09)	
				• C5 = 0.84 (0.70- 1.00)	
				Trend	
				Adjusted RR (95% CI), age ≥ 55 years	
				• C1+2 = 1.00 (referent)	
				• C3 = 1.01 (0.96- 1.07)	
				• C4 = 1.04 (0.98- 1.11)	
				• C5 = 0.82 (0.71- 0.94)	
				Trend	
Rockhill et al 1998 [338]	To examine the association between PA at two different times in life and BC risk.	• n = 372	Baseline and 6 year follow-up	372 cases of BC	The findings do not support a link between PA in late adolescence or in the recent past and BC risk among young adult women.
USA		• Sex: Women	PA assessment: Questionnaire for MVPA (h/wk)	Multivariate adjusted RR (95% CI)	
Prospective cohort		• Age: 25-42	G1 = <1	• G1 = 1.00 (referent)	
D & B score = 12		• Characteristics: Nurses	G2 = 1.0-1.9	• G2 = 1.1 (0.8-1.4)	
		• The Nurses Health Study	G3 = 2.0-3.9	• G3 = 1.1 (0.8-1.4)	
			G4 = 4.0-6.9	• G4 = 1.0 (0.7-1.4)	
			G5 = ≥ 7	• G5 = 1.1 (0.8-1.5)	
			Logistic regression		

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

Slattery et al 2007 [339]	To evaluate the BC risk associated with TPA and VPA at ages 15, 30 and 50 years and the referent year prior to diagnosis/selection.	<ul style="list-style-type: none"> <li>• n = 4,850 Non-Hispanic white: n = 3,128 (cases n = 1,527 controls n = 1601); Hispanic American Indian: n = 1,722 (cases n = 798, controls n = 924)</li> <li>• Sex: Women</li> <li>• Age: &lt;50 yr</li> <li>• Characteristics: Non- Hispanic white and Hispanic American Indian</li> </ul>	PA assessment: Questionnaire for TPA (activity score) and lifetime VPA (h/wk)	1527 cases of BC (non-Hispanic white; 798 cases of BC (Hispanic American Indian)	The data suggest that PA is important in reducing risk of BC in non-Hispanic white and Hispanic American Indian women.
USA			TPA score		
Case control			G1 = 0-3		
D & B score = 12			G2 = 4-6	OR (95% CI) by TPA score, non-Hispanic white	
			G3 = 7-9	• G1 = 1.00 (referent)	
			G4 = 10-12	• G2 = 0.78 (0.52- 1.17)	
			Lifetime VPA	• G3 = 0.84 (0.57- 1.22)	
			G1 = None	• G4 = 0.70 (0.44- 1.12)	
			G2 = <1.0	Trend $p = 0.26$	
			G3 = 1.0-2.9	OR (95% CI) by TPA score, Hispanic American Indian	
			G4 = $\geq 3.0$	• G1 = 1.00 (referent)	
			Multivariable logistic regression	• G2 = 1.49 (0.98- 2.26)	
				• G3 = 1.21 (0.80- 1.84)	
				• G4 = 0.97 (0.53- 1.76)	
				Trend $p = 0.90$	
				OR (95% CI) by lifetime VPA, non-Hispanic white	
				• G1 = 1.00 (referent)	
				• G2 = 0.66 (0.36- 1.23)	
				• G3 = 0.73 (0.40- 1.34)	
				• G4 = 0.69 (0.37- 1.27)	
				Trend $p = 0.68$	
				OR (95% CI) by lifetime VPA, Hispanic American Indian	
				• G1 = 1.00 (referent)	
				• G2 = 1.15 (0.67- 1.96)	
				• G3 = 1.19 (0.70- 2.03)	
				• G4 = 1.09 (0.62- 1.90)	
				Trend $p = 0.84$	
Sprague et al 2007 [340]	To investigate the relationship between LTPA and strenuous OPA and BC risk.	<ul style="list-style-type: none"> <li>• n = 15,710 (1,689 cases in situ; 6,391 invasive and 7,630 controls)</li> <li>• Sex: Women</li> <li>• Age: 20-69</li> </ul>	PA assessment: Questionnaire for lifetime TPA (hr/wk and MET hr/wk), divided into groups	Adjusted OR (95% CI) for in situ BC by lifetime TPA (hr/wk)	The results provide further evidence that for most women, PA may reduce the risk of invasive BC.
USA			Lifetime total PA (hr/wk)	• G1 = 1.00 (referent)	
Case control			G1 = 0	• G2 = 0.92 (0.72-1.19)	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

D & B score = 13		• The Collaborative Breast Cancer Study	G2 = 0.1-15.0	• G3 = 0.83 (0.62-1.13)	
			G3 = 15.1-30.0	• G4 = 0.86 (0.59- 1.24)	
			G4 = > 30.0	Trend $p = 0.22$	
			MET hr/wk	Adjusted OR (95% CI) for in situ BC by lifetime TPA (MET hr/wk)	
			G1 = 0.0	• G1 = 1.00 (referent)	
			G2 = 0.1-62.5	• G2 = 0.93 (0.72- 1.20)	
			G3 = 62.6-125.0	• G3 = 0.82 (0.61- 1.10)	
			G4 = >125.0	• G4 = 0.82 (0.57- 1.17)	
				Trend $p = 0.10$	
				Adjusted OR (95% CI) for invasive BC by lifetime TPA (hr/wk)	
				• G1 = 1.00 (referent)	
				• G2 = 0.88 (0.76- 1.03)	
				• G3 = 0.87 (0.73- 1.05)	
				• G4 = 0.85 (0.67- 1.07)	
				Trend $p = 0.22$	
				Adjusted OR (95% CI) for invasive BC by lifetime TPA (MET hr/wk)	
				• G1 = 1.00 (referent)	
				• G2 = 0.89 (0.76- 1.04)	
				• G3 = 0.82 (0.68- 0.99)	
				• G4 = 0.88 (0.71- 1.09)	
	Trend $p = 0.12$				
Steindorf et al 2003 [341]	To clarify the relationship between PA and BC risk.	• n = 1,246 (360 cases; 886 controls)	PA assessment: Computer assisted telephone interview for TPA (MET hr/wk) at various ages	360 cases of BC	The data do not suggest an inverse association between PA and BC risk in pre-menopausal women.
Germany		• Sex: Women	TPA at age 12-19 yr	Multivariate adjusted OR (95% CI) by TPA at age 12-19 yr	
Case control		• Age: Mean. cases 41.9 yr; controls 42.5 yr	G1 = 13.0-55.7	• G1 = 1.00 (referent)	
D & B score = 13			G2 = 55.8-88.7	• G2 = 1.07 (0.75- 1.52)	
			G3 = 88.8-134.0	• G3 = 1.00 (0.70- 1.42)	
			G4 = 134.1-695.9	• G4 = 0.73 (0.50- 1.07)	
			TPA at age 20-30 yr	Trend $p = 0.44$	



**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

			G1 = 6.4-69.0	Multivariate adjusted OR (95% CI) by TPA at age 20-30 yr	
			G2 = 69.1-109.0	• G1 = 1.00 (referent)	
			G3 = 109.1-160.4	• G2 = 0.95 (0.67- 1.37)	
			G4 = 160.5-728.8	• G3 = 0.85 (0.59- 1.23)	
			TPA at age 12-30 yr (both)	• G4 = 0.96 (0.67- 1.39)	
			G1 = 17.2-70.4	Trend $p = 0.32$	
			G2 = 70.5-104.0	Multivariate adjusted OR (95% CI) by TPA at age 12-30 yr	
			G3 = 104.1-145.5	• G1 = 1.00 (referent)	
			G4 = 145.6-564.4	• G2 = 0.97 (0.68- 1.38)	
			Logistic regression	• G3 = 0.68 (0.46- 0.99)	
				G4 = 0.94 (0.65- 1.35)	
				Trend $p = 0.29$	
Tehard et al 2006 [342]	To investigate the type, duration, frequency and intensity of PA required to reduce the risk of BC.	• n = 90,509	Baseline and follow-up every 2 years for 12 years	3,424 cases of BC	BC risk was reduced, especially with VPA.
France		• Sex: Women	PA assessment: Questionnaire for various PA variables, all divided into groups	Multivariate adjusted RR (95% CI) by TPA	
Prospective cohort		• Age: 40-65	TPA (MET hr/wk)	• G1 = 1.00 (referent)	
D & B score = 13		• Characteristics: French women insured with Mutuelle Generale de l'Education Nationale	G1 = <28.3	• G2 = 1.05 (0.93- 1.17)	
		• E3N Cohort Study	G2 = 28.3-41.8	• G3 = 0.94 (0.83- 1.05)	
			G3 = 41.8-57.8	• G4 = 0.90 (0.80- 1.02)	
			G4 = ≥ 57.8	Trend $p < 0.05$	
			Total recreational PA (MET hr/wk)	Multivariate adjustedRR (95% CI) by total recreational PA	
			G1 = Inactive	• G1 = 1.00 (referent)	
			G2 = <16.0	• G2 = 0.82 (0.71- 0.93)	
			G3 = 16.0-22.3	• G3 = 0.94 (0.84- 1.06)	
			G4 = 22.3-33.8	• G4 = 0.88 (0.79- 0.98)	
			G5 = ≥33.8 = 0.81	• G5 = 0.81 (0.72- 0.92)	
			Walking (min/d)	Trend $p < 0.01$	
			G1 = <500	Multivariate adjusted RR (95% CI) by walking duration	
			G2 = 500-2000	• G1 = 1.00 (referent)	
				• G2 = 1.03 (0.95- 1.11)	


**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

			G3 = >2000	• G3 = 0.91 (0.81- 1.02)	
			MPA (hr/wk)	Trend $p = 0.45$	
			G1 = Inactive	Multivariate adjusted RR	
			G2 = 0	(95% CI) by MPA	
			G3 = 1-4	• G1 = 1.00 (referent)	
			G4 = 5-13	• G2 = 0.80 (0.60- 1.05)	
			G5 = 14	• G3 = 0.87 (0.79- 0.94)	
			VPA (hr/wk)	• G4 = 0.86 (0.74- 0.99)	
			G1 = Inactive	• G5 = 0.89 (0.65- 1.24)	
			G2 = 0	Trend $p < 0.01$	
			G3 = 1-2	Multivariate adjusted RR (95% CI) by VPA	
			G4 = 3-4	• G1 = 1.00 (referent)	
			G5 = 5	• G2 = 0.90 (0.81- 0.99)	
			Cox proportional hazard ratio	• G3 = 0.88 (0.79- 0.97)	
				• G4 = 0.82 (0.71- 0.95)	
				• G5 = 0.62 (0.49- 0.78)	
				Trend $p < 0.0001$	
Thune et al 1997 [343]	To investigate whether everyday exercise is related to the risk of BC.	• n = 25,624	Baseline and mean follow- up of 14 years	351 cases of BC (110 pre-menopausal and 251 post-menopausal women)	LTPA and OPA are associated with a reduced risk of BC.
Norway		• Sex: Women	PA assessment: Self-reported LTPA and OPA, divided into groups	• G1 = 1.00 (referent)	Adjusted RR (95% CI) by LTPA
Prospective cohort		• Age: 20-54	LTPA	• G2 = 0.93 (0.71- 1.22)	
D & B score = 14			G1 = Sedentary	• G3 = 0.63 (0.42- 0.95)	
			G2 = Moderate	Trend $p = 0.04$	
			G3 = Regular exercise OPA	Adjusted RR (95% CI) by LTPA, pre- menopausal	
			G1 = Sedentary	• G1 = 1.00 (referent)	
			G2 = Walking	• G2 = 0.77 (0.46- 1.27)	
			G3 = Lifting	• G3 = 0.53 (0.25- 1.14)	
			G4 = Heavy manual labor	Trend $p = 0.10$	
			During work	Adjusted RR (95% CI) by LTPA, post- menopausal	
			Pre-menopausal	• G1 = 1.00 (referent)	
			G1 = Sedentary	• G2 = 1.00 (0.72- 1.39)	
			G2 = Walking	• G3 = 0.67 (0.41- 1.10)	
			G3 = Lifting or heavy manual labor	Trend $p = 0.15$	
				Adjusted RR (95% CI) by OPA	

**Table 16: Studies examining the relationship between physical activity and breast cancer. (Continued)**

				<ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.84 (0.63- 1.12)</li><li>• G3 = 0.74 (0.52- 1.06)</li><li>• G4 = 0.48 (0.25- 0.92)</li></ul> <p>Trend <i>p</i> = 0.02</p> <p>Adjusted RR (95% CI) by OPA, pre- menopausal</p> <ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.82 (0.50- 1.34)</li><li>• G3 = 0.48 (0.24- 0.95)</li></ul> <p>Trend <i>p</i> = 0.03</p> <p>Adjusted RR (95% CI) by OPA, post- menopausal</p> <ul style="list-style-type: none"><li>• G1 = 1.00 (referent)</li><li>• G2 = 0.87 (0.61- 1.24)</li><li>• G3 = 0.78 (0.52- 1.18)</li></ul> <p>Trend <i>p</i> = 0.24</p>	
Zheng et al 1993 [344]	To assess the role of OPA in the risk of BC.	<ul style="list-style-type: none"><li>• n= 3,783 (BC = 2,736)</li></ul>	PA assessment: Interview for OPA, divided into groups	2,736 cases of BC	Women with low OPA had an increased risk of BC; the incidence of BC was reduced in women with high-activity jobs.
China		<ul style="list-style-type: none"><li>• Sex: Women</li></ul>	G1 = Low	Standardized incidence ratios	
D & B score = 9		<ul style="list-style-type: none"><li>• Age: 30</li></ul>	G2 = Moderate		
			G3 = High		
				<ul style="list-style-type: none"><li>• G1= 131</li><li>• G2 = 95</li><li>• G3 = 79</li></ul>	

D & B score, Downs and Black quality score; YR, years; PA, physical activity; BC, breast cancer; LTPA, leisure-time physical activity; g, group; HR, hazard ratio; RR, risk ratio; OR, odds ratio; 95% CI, confidence interval; T, tertile; MET, metabolic equivalent; MET/wk, metabolic equivalent per week; OPA, occupational physical activity; MET h/wk/yr, metabolic equivalent per hour per week per year; kcal/wk, kilocalories per week; TPA, total physical activity; VPA, vigorous physical activity.



relationship in one or more measures of occupational and/or leisure-time physical activity and the risk for breast cancer. Moreover, the majority of studies demonstrated the greatest risk reduction at the highest activity level. With respect to the minimal and optimal volume of exercise required, Lee [105] stated that 30-60 min/day of moderate-to-vigorous physical activity is required to decrease the risk for breast cancer. This belief is strongly supported by the literature. However, others have shown significant risk reductions at lower exercise volumes. For instance, Rockhill et al. [106] showed significant reductions (12% or greater) in the risk for breast cancer in women who accumulated at least 1 hr of moderate or vigorous physical activity per week.

Similarly, Sesso et al. (1998) revealed that there was an 8% reduction in the risk for breast cancer with a relatively small energy expenditure of 500-999 kcal/wk. Further risk reductions were observed with higher energy expenditures (= 1000 kcal/wk = 51% reduction in the risk). As discussed above, Monnikhof et al. revealed a 6% decrease in breast cancer risk for each additional hour of physical activity per week [104]. Taken as a whole, it would therefore appear that Canada's guidelines for physical activity are more than appropriate for reducing the risk for breast cancer. Further research however is required to determine the minimal volume of exercise that is effective in the primary prevention of breast cancer.

#### Implications

There is a preponderance of data linking physical inactivity to site-specific cancers, particularly of the breast and colon [31,104-109]. The protective effects of physical activity also appear with other forms of cancer (such as endometrial cancer) [110]. In an important review of the literature Lee revealed that physically active women have a 20-30% lower risk of breast cancer, and physically active men and women have a 30-40% lower risk of colon cancer [105]. A more recent systematic review of the literature revealed a 20-80% lower risk of breast cancer in post-menopausal women [104], with a weaker association in pre-menopausal women. Considering data from both pre- and post-menopausal women the authors demonstrated that physically active individuals had a 15-20% lower risk of breast cancer. Monnikhof et al. also reported a 6% lower risk of breast cancer for each additional hour of physical activity per week [104]. This level of risk reduction was also supported by the U.S. Department of Health and Human Services during its recent evaluation of the literature [31].

Our current reviews of the literature support previous work in the field including the finding of a dose-response relationship between physical activity and cancers of the breast and colon [104,105,109]. It would appear that 30-60 min/day of moderate-to-vigorous physical activity is associated with a lower risk of breast and colon cancer.


#### Recommendation #5

*For a reduced risk for site specific cancers (such as colon cancer and breast cancer), it is recommended that individuals should participate in 30 min or more of moderate to vigorous exercise on most days of the week. [Level 2, Grade A]*

#### Primary Prevention of Type 2 Diabetes

In comparison to other chronic conditions, there is relatively limited literature examining the relationship between multiple levels of physical activity/fitness and the incidence of type 2 diabetes. All of the literature

examining the dose-response (for at least three levels of physical activity/fitness) involved prospective cohort analyses. A total of 3655 citations were identified during the electronic database search (Figure 9). Of these citations, 2038 were identified in MEDLINE, 1116 in EMBASE, 118 in Cochrane, and 372 in the CINAHL/SportDiscus/PsychInfo search. A total of 614 duplicates were found, leaving a total of 3041 unique citations. A total of 2865 articles were excluded after scanning, leaving a total of 176 articles for full review. From these articles 156 were excluded after full review leaving 20



**Figure 9** Results of the Literature Search for Diabetes.

articles for inclusion in the systematic review of the literature regarding the relationship between physical activity and type 2 diabetes. The reasons for exclusion included non-experimental/weak design (N = 18), three levels of physical activity not reported (N = 16), reviews, summaries, or meta-analyses (N = 41), not related to type 2 diabetes (N = 71), and other (N = 10).

As shown in Table 17, 20 investigations examined the dose-response (i.e., three or more levels) relationship between physical activity and the incidence of type 2 diabetes. This involved a total of 624,952 subjects, averaging 32,892 subjects per study (range 1,543-87,907). There were a total of 19,325 cases of type 2 diabetes (ranging per study from 78-4,030). The total length of follow-up averaged 9.3 yr (ranging from 3 -16.8 yr). The articles were published over a 16 yr period ranging from 1991 to 2007.

Of these studies 100% revealed an inverse relationship between type 2 diabetes and levels of physical activity or fitness. When comparing the most active/fit group versus the least active/fit group we found an average risk reduction of 42% (median = 44%). Therefore in our analyses the most physically active/fit had a 42% lower risk of developing type 2 diabetes. The majority (84%) of these studies revealed incremental reductions in the risk for type 2 diabetes with increasing activity/fitness levels. Therefore, the health benefits with respect to type 2 diabetes prevention appear to continue across the physical activity/fitness continuum. Similar to other clinical conditions, the dose-response relationship is such that small changes in activity levels yield marked reductions in the risk for type 2 diabetes. The health benefits of exercise appear to be particularly prevalent in individuals at high risk for developing type 2 diabetes (e.g., those with a high body mass index, the metabolic syndrome, a history of hypertension and/or a family history of type 2 diabetes). *The level of evidence relating physical activity to the primary prevention of type 2 diabetes would be considered to be Level 2A.* The quality of the investigations was generally high with a mean (and median) Downs and Black score of 13 (range 11-14).

As with other conditions is it difficult to separate the effects of volume and intensity of exercise. However, small changes in activity levels clearly can have a large effect on the risk for and incidence of type 2 diabetes. For instance, Hu and coworkers [111] revealed that nurses (n = 68,497) who engaged in 1 hr/day of brisk walking had 24% less obesity and 34% less type 2 diabetes (over a 6-year follow-up). These authors estimated that approximately 30% of new cases of obesity and 43% of new cases of type 2 diabetes could be prevented by adopting an active lifestyle including less than 10 hr/wk of television watching and  $\geq 30$  min/d of brisk walking. Similarly, over a 5-year period, male physicians who

**Table 17 The relationship between physical activity and the development of type 2 diabetes.**

Publication Country Study Design Quality Score	Objective	Population	Methods	Outcome	Comments and Conclusions
Haapanen et al 1997 [77]	To examine the association of PA and the risk of CHD, hypertension and T2D.	• n = 1,340 men, 1,500 women	10 yr follow-up	Number of cases: 118	LTPA has a preventive effect on T2D.
Finland		• Age: 35-63 yr	PA assessment: Self-reported LTPA (kcal/wk), divided into groups	Age-adjusted RR (95% CI), men • G1 = 1.54 (0.83-2.84) • G2 = 1.21 (0.63-2.31) • G3 = 1.00 (referent)	
Prospective cohort				$p = 0.374$	
D & B score = 14			Men G1 = 0-1100 G2 = 1101-1900 G3 = >1900  Women (kcal/wk) G1 = 0-900 G2 = 901-1500 G3 = >1500 Cox proportional HR	Age-adjusted RR (95% CI), women • G1 = 2.64 (1.28-5.44) • G2 = 1.17 (0.50-2.70) • G3 = 1.00 (referent) $p < 0.006$	
Hu et al 2003 [111]	To examine the relationship between sedentary behaviours (particularly prolonged television watching) and risk of obesity and T2D in women.	• n = 68,497 (diabetes specific analyses)	6 yr follow-up	Number of cases: 1515	Sedentary behaviours (especially television watching) are associated with an increased risk for obesity and T2D.
USA		• n = 50,277 (obesity specific analyses)	PA assessment: Self-reported PA and sedentary behaviour	Each 2-h/d increment in TV watching was associated with a 23% (95% CI, 17%-30%) increase in obesity and a 14% (95% CI, 5%- 23%) increase in risk of T2D	
Prospective cohort		• Age: 30-55 yr	Outcome measure: onset of obesity and T2D	Each 2-h/d increment in sitting at work was associated with a 5% (95% CI, 0%-10%) increase in obesity and a 7% (95% CI, 0%- 16%) increase in T2D	Light to moderate PA was associated with a significantly lower risk for obesity and T2D.
D & B score = 13		• Sex: Women	Multivariate analyses adjusting for age, smoking, dietary factors, and other covariates	Standing or walking around at home (2 h/d) was associated with a 9% (95% CI, 6%-12%) reduction in obesity and a 12% (95% CI, 7%- 16%) reduction in T2D	

**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

		<ul style="list-style-type: none"> <li>• Characteristics:</li> </ul>		Each 1 hour per day of brisk walking was associated with a 24% (95% CI, 19%-29%) reduction in obesity and a 34% (95% CI, 27%- 41%) reduction in T2D	
		Free of T2D, CVD, or cancer at baseline			
		• Nurses' Health Study			
Manson et al 1992 [112]	To examine the association between regular exercise and the subsequent development of T2D.	• n = 21,271	5 yr follow-up	Number of cases: 285	Exercise appears to reduce the development of T2D even after adjusting for BMI.
		• Sex: Men	PA assessment: Questionnaire Fpr VPA (enough to develop sweat)		
		• Age: 40-84 yr		The age-adjusted incidence of T2D:	
USA		• Characteristics:		<ul style="list-style-type: none"> <li>• 369 cases per 100,000 person- years in men who engaged in VPA less than once weekly</li> <li>• 214 cases per 100,000 person- years in those exercising at least five times per week (p trend &lt; 0.001)</li> </ul>	
		Free of diagnosed diabetes, CVD and cancer at baseline			
Prospective cohort					
D & B score = 14			Exercise frequency (times/wk)		
			G1 = < Weekly		
			G2 = At least weekly		
			Times per week		
			G1 = 0	• G1 = 1.00 (referent)	
			G2 = 1	• G2 = 0.64 (0.51- 0.82)	
			G3 = 2-4		
			G4 = >5		
			Outcome measure: Incidence T2D	Age-adjusted RR (95% CI) by exercise frequency	
				• G1 = 1.00 (referent)	
				• G2 = 0.77 (0.55-1.07)	
				• G3 = 0.62 (0.46-0.82)	
				• G4 = 0.58 (0.40-0.84)	
				Age- and BMI-adjusted RR (95% CI) by exercise frequency	
				• G1 = 1.00 (referent)	

**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

				<ul style="list-style-type: none"> <li>• G2 = 0.71 (0.56- 0.91)</li> </ul> <p>Age- and BMI-adjusted RR (95% CI) by exercise frequency</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.78 (0.56-1.09)</li> <li>• G3 = 0.68 (0.51-0.90)</li> <li>• G4 = 0.71 (0.49-1.03)</li> </ul>	
Hu et al. 2001 [114]	To examine the relationship between dietary and lifestyle factors in relation to the risk for T2D.	<ul style="list-style-type: none"> <li>• n = 84,941</li> </ul>	16 yr follow-up	Number of cases: 3300	The majority of T2D could be prevented through healthy living.
USA		<ul style="list-style-type: none"> <li>• Sex: Women</li> <li>• Age: 40-75 yr</li> <li>• Characteristics: participants had no history of diabetes, CVD, or cancer.</li> </ul>	PA assessment: Questionnaire For PA (h/wk), divided into groups	Multivariate-adjusted RR (95%)	
Retrospective cohort				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.89 (0.77-1.02)</li> <li>• Q3 = 0.87 (0.75-1.00)</li> <li>• Q4 = 0.83 (0.71-0.96)</li> <li>• Q5 = 0.71 (0.56-0.90)</li> </ul>	
D & B score = 13		Nurses' Health Study	<p>Q1 = &lt;0.5</p> <p>Q2 = 0.5-1.9</p> <p>Q3 = 2.0-3.9</p> <p>Q4 = 4.0-6.9</p> <p>Q5 = ≥7.0</p> <p>Outcome measure: Incidence of T2D</p> <p>Cox regression</p>		
Sato et al 2007 [116]	To examine the relationship between walking to work and the development of T2D.	<ul style="list-style-type: none"> <li>• n = 8,576</li> </ul>	4 yr follow-up	Number of cases: 878	The duration of a walk to work is an independent predictor of the risk for T2D.
Japan		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: 40-55 yr</li> <li>• Kansai Healthcare Study</li> </ul>	PA assessment: For time spent walking to work, divided into tertiles	OR (95% CI)	
Prospective cohort			<p>T1 = 0-10 min</p> <p>T2 = 11-20 min</p> <p>T3 = ≥20 min</p>	<ul style="list-style-type: none"> <li>• T1 = 1.00 (referent)</li> <li>• T2 = 0.86 (0.70-1.06)</li> <li>• T3 = 0.73 (0.58-0.92)</li> </ul> <p>Significant difference was seen between ≤ 10 min and ≤ 20 min only (p = 0.007)</p>	





**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

				<ul style="list-style-type: none"> <li>• G3 = 0.57 (0.34-0.96)</li> </ul> <p>Multivariate adjusted HR (95% CI) for commuting PA, men and women</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.96 (0.74-1.25)</li> <li>• G3 = 0.64 (0.45-0.92)</li> </ul> <p>Multivariate adjusted HR (95% CI) for LTPA, men</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.78 (0.57-1.06)</li> <li>• G3 = 0.84 (0.52-1.37)</li> </ul> <p>Multivariate adjusted HR (95% CI) for LTPA, women</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.81 (0.58-1.15)</li> <li>• G3 = 0.85 (0.43 -1.66)</li> </ul> <p>Multivariate adjusted HR (95% CI) for LTPA, men and women</p> <ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.81 (0.64-1.20)</li> <li>• G3 = 0.84 (0.57-1.25)</li> </ul>	
Hsia et al 2005 [118] USA	To evaluate the relationship between PA and the incidence of T2D in a large, diverse group of older women.	<ul style="list-style-type: none"> <li>• n = 87,907</li> </ul>	PA assessment: Questionnaire for frequency and duration of 4 walking speeds and 3 other activities classified by intensity (light, moderate, strenuous)	Number of cases: 2,271	There is a strong inverse relationship between PA and T2D. There is a stronger relationship between PA and T2D in Caucasian women than in minority women. This may be explained by less precise risk estimates in minority women.
		<ul style="list-style-type: none"> <li>• Sex: Women</li> <li>• Age: White 63.8 ± 7.3, African American 61.9 ± 7.3, Hispanic 60.5 ± 7.1, Asian/Pacific Islander 63.7 ± 7.6, American Indian 61.5 ± 8.0</li> </ul>		Multivariate adjusted HR (95% CI) by walking, Caucasian	
Prospective cohort				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.85 (0.74-0.87)</li> <li>• Q3 = 0.87 (0.75-1.01)</li> <li>• Q4 = 0.75 (0.64-0.89)</li> <li>• Q5 = 0.74, (0.62-0.89)</li> </ul> <p>Trend <math>p &lt; 0.001</math></p>	
D & B score = 11			<p>Q1 = Low</p> <p>Q2 =</p> <p>Q3 =</p>		

**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

			Q4 =	Multivariate adjusted HR (95% CI) by TPA, Caucasian	
			Q5 = High		
		<ul style="list-style-type: none"> <li>Ethnicity: White n = 74,240; African American n = 6,465; Hispanic n = 3,231; Asian/Pacific Islander 2,445; American Indian n = 327</li> </ul>		<ul style="list-style-type: none"> <li>Q1 = 1.00 (referent)</li> </ul>	
			Cox proportional HR		
		<ul style="list-style-type: none"> <li>Characteristics: participants had no history of diabetes, were not on any antidiabetic medications</li> <li>Women's Health Initiative</li> </ul>		<ul style="list-style-type: none"> <li>Q2 = 0.88 (0.76- 1.01)</li> <li>Q3 = 0.74 (0.64- 0.87)</li> <li>Q4 = 0.80 (0.68- 0.94)</li> <li>Q5 = 0.67 (0.56- 0.81) Trend <math>p = 0.002</math></li> </ul>	
Wannamethee et al 2000 [120]	To examine the role of components of the insulin resistance syndrome in the relationship between PA and the incidence of T2D and CHD.	<ul style="list-style-type: none"> <li>n = 5,159</li> </ul>	16.8 yr follow-up	Number of cases: 196	The relationship between PA and T2D appears to be mediated by serum insulin and components of the insulin resistance syndrome. However, these factors do not appear to explain the inverse relationship between PA and T2D.
		<ul style="list-style-type: none"> <li>Sex: Men</li> <li>Age: 40-59 yr</li> </ul>	PA assessment: Questionnaire for TPA Physical activity groups were identified and scored:	Multivariate adjusted RR (95% CI)	
England, Wales and Scotland		<ul style="list-style-type: none"> <li>Characteristics: No history of heart disease, diabetes or stroke</li> </ul>		Q1 = 1.00 (referent)	
				Q2 = 0.66 (0.42-1.02)	
				Q3 = 0.65 (0.41-1.03)	
				Q4 = 0.48 (0.28-0.83)	
Prospective cohort			Q1 = None	Q5 = 0.46 (0.27-0.79)	
			Q2 = Occasional	$p < 0.005$	
			Q3 = Light		
			Q4 = Moderate		
D & B score = 14					

**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

		Q5 = Moderately vigorous/vigorous		MPA (sporting activity once a week or frequent lighter- intensity activities such as walking, gardening, do-it yourself projects) are sufficient to produce a significant reduction in risk of both CHD and T2D.	
		The men were classified according to current smoking status, alcohol consumption, and social class			
		Cox proportional HR			
Manson et al 1991 [121]	To examine the association between regular VPA and the incidence of T2D.	• n = 87,253	8 yr follow-up	Number of cases: 1303 Women who engage in VPA at least once per week had reduced adjusted RR of T2D RR = 0.66 (0.6- 0.75)	PA is promising in the primary prevention of T2D.
USA		• Sex: Women • Age: 34-59 yr • Characteristics: Free of diagnosed diabetes, cardiovascular disease and cancer	PA assessment: Questionnaire		
Prospective cohort			Frequency of weekly exercise (0-+4)		
D & B score = 13				The reduction in risk remained significant after adjustment for BMI RR = 0.84 (0.75-0.95)	
			Analysis also restricted to the first 2 yr after the assessment of PA level and to symptomatic diabetes	When analysis was restricted to the first 2 years after ascertainment of PA level and to symptomatic disease as the outcome, the age- adjusted RR of those who exercised was 0.50, and age and body-mass index adjusted RR was 0.69 (0.48-1.0)	
			Multivariate adjustments for age, body-mass index, family history of diabetes, and other variables did not alter the reduced risk found with exercise		
			Multivariate analysis	Family history of diabetes did not modify the effect of exercise, and risk reduction with exercise was evident among both obese and non-obese women	

**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

Helmrich et al 1994 [122]	To examine the relationship between PA and the development of T2D.	<ul style="list-style-type: none"> <li>• n = 5,990</li> </ul>	98,524 man-years of follow-up (1962- 1976)	Number of cases: 202	Increased PA is effective in preventing T2D.
USA		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: 39-68 yr</li> <li>• Characteristics: healthy, asymptomatic</li> </ul>	PA assessment: Questionnaire for LTPA (walking, stair climbing, sports etc; kcal/wk) Blocks walked/day	RR (95% CI) by blocks walked per day	The protective benefit is especially pronounced in those individuals who have the highest risk of disease.
Further review of the data reported by Helmich et al. 1991				<ul style="list-style-type: none"> <li>• T1 = 1.00 (referent)</li> <li>• T2 = 1.30</li> </ul>	
		University of Pennsylvania Alumni Health Study		<ul style="list-style-type: none"> <li>• T3 = 0.92</li> </ul>	
			LTPA (kcal/wk) kcal were assigned to each activity and added together	$p = 0.80$ LTPA was inversely related to the development of T2D	
Prospective cohort				Same findings to that reported in 1991	
D & B score = 14			Lowest < 500 Highest ≥ 3500 Blocks walked/day T1 = <5 T2 = 5-14 T3 = ≥15 Cox proportional HR		
Helmrich et al 1991 [123]	To examine the Relationship between PA and the Subsequent development of T2D.	<ul style="list-style-type: none"> <li>• n = 5,990</li> </ul>	98,524 man-years of follow-up (1962- 1976)	Number of cases: 202	Increased PA is effective in preventing T2D.
USA		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: 39-68 yr</li> <li>• Characteristics: healthy, asymptomatic</li> </ul>	PA assessment: Questionnaire for LTPA kcal/wk: stairs climbed/day and blocks walked/day, divided into groups	LTPA was inversely related to the development of type 2 diabetes	The protective benefit is especially pronounced in those individuals who have the highest risk of disease.

**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

Prospective cohort		RR (95% CI) by sports played
	• University of Pennsylvania Alumni Health Study	• G1 = 1.00 (referent)
		• G2 = 0.90
		• G3 = 0.69
		• G4 = 0.65
D & B score = 13		Trend $p = 0.02$
	All activities LTPA	
	Q1 = <500	
	Q2 = 500-999	RR (95% CI) by Flights of stairs climbed/day
	Q3 = 1000-1499	
	Q4 = 1500-1999	• T1 = <5 = 1.00 (referent)
	Q5 = 2000-2499	• T2 = 0.78
	Q6 = 2500-2999	
		• T3 = 0.75
	Q7 = 3000-3499	
		Trend $p = 0.07$
	Q8 = $\geq 3500$	
		RR (95% CI) by Blocks walked/day
	Sports played	
	G1 = None	• T1 = 1.00 (referent0)
		• T2 = 1.31
	G2 = Moderate	
	G3 = Vigorous	• T3 = 0.93
		Trend $p = 0.80$
	G4 = Moderate and Vigorous	
		Age adjusted RR (95% CI) by all activities
	Stairs climbed per day	
	T1 = <5	• Q1 = 1.00 (referent)
	T2 = 5-14	• Q2 = 0.94
	T3 = $\geq 15$	• Q3 = 0.79
		• Q4 = 0.78
	Blocks walked per day	• Q5 = 0.68
	T1 = <5	• Q6 = 0.90

**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

			T2 = 5-14 T3 = ≥ 15  Cox proportional HR	<ul style="list-style-type: none"> <li>• Q7 = 0.86</li> <li>• Q8 = 0.52</li> </ul> <p><math>p = 0.01</math> for trend</p> <p>Age adjusted RR (95% CI) by all activities except vigorous sports</p> <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.97</li> <li>• Q3 = 0.87</li> <li>• Q4 = 0.92</li> <li>• Q5 = 0.75</li> <li>• Q6 = 1.29</li> <li>• Q7 = 1.03</li> <li>• Q8 = 0.48</li> </ul> <p>Trend <math>p = 0.07</math></p> <p>Age adjusted RR (95% CI) by vigorous sports only</p> <ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.69</li> <li>• Q3 = N/A</li> <li>• Q4 = 0.53</li> <li>• Q5 = 0.86</li> <li>• Q6 = 0.56</li> <li>• Q7 = 0.40</li> <li>• Q8 = 0.46</li> </ul> <p>Trend <math>p = 0.05</math></p>	
Wei et al 1999 [124]	To determine whether PF is associated with risk for impaired fasting glucose and T2D.	<ul style="list-style-type: none"> <li>• n = 8,633</li> </ul>	6 yr follow-up	Number of cases: 149	High PF is associated with a reduced risk for impaired fasting glucose and T2D.
USA		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: 43.5 yr</li> </ul>	PF assessment: Maximal treadmill exercise test (METs), divided into 3 groups	593 patients developed impaired fasting glucose	
Prospective cohort		<ul style="list-style-type: none"> <li>• Characteristics: Non-diabetic men</li> </ul>	T1 = Low  T2 = Moderate	<p>OR (95% CI) for developing glucose intolerance</p> <ul style="list-style-type: none"> <li>• T1 = 1.9 (1.5–2.4)</li> <li>• T2 = 1.5 (1.2–1.8)</li> </ul>	

**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

D & B score = 12			T3 = High Outcome measure: Incidence of impaired fasting glucose and T2D	<ul style="list-style-type: none"><li>• T3 = 1.00 (referent)</li><li>OR (95% CI) for developing T2D</li><li>• T1 = 3.7 (2.4 –5.8)</li><li>• T2 = 1.7 (1.1–2.7)</li><li>• T3 = 1.00 (referent)</li></ul>	
Statistics: GLM					
Katzmarzyk et al 2007 [126]	To examine the relationships among adiposity, PA, PF and the development of T2D in a diverse sample of Canadians.	<ul style="list-style-type: none"><li>• n = 1,543 (709 men and 834 women)</li></ul>	6 yr follow-up	Number of cases: 78 (37 in men, 41 in women)	Adiposity and PF are important predictors of the development of T2D.
Canada		<ul style="list-style-type: none"><li>• Sex: Men and women</li></ul>	PF assessment: Questionnaire	PA was associated with 23% lower odds of developing diabetes and maximal METs was also associated with significantly lower odds of developing diabetes (OR = 0.28)	
Prospective cohort		<ul style="list-style-type: none"><li>• Age: 36.8 - 37.5</li></ul>	PA assessment: LTPA Questionnaire		
D & B score = 13		<ul style="list-style-type: none"><li>• Characteristics: Free of diabetes at baseline</li><li>• Canadian Physical Activity Longitudinal Study</li></ul>			
Burchfiel et al 1995 [345]	To examine the relationship between PA and T2D.	<ul style="list-style-type: none"><li>• n = 6,815</li></ul>	6 yr follow-up	Number of cases: 391	PA is associated inversely and independently with incident T2D.
USA		<ul style="list-style-type: none"><li>• Sex: Men (Japanese-American)</li></ul>	PA assessment: Questionnaire PA index (based on intensity and duration of activity)	The age-adjusted 6-year cumulative incidence of diabetes decreased progressively with increasing quintile of physical activity from 73.8 to 34.3 per 1,000 (p < 0.0001, trend)	
Prospective cohort		<ul style="list-style-type: none"><li>• Age: 45-68 yr</li><li>• Characteristics: Free of diabetes at entry</li><li>• The Honolulu Heart Program</li></ul>	Levels of activity: Q1 = Basal - Sleeping reclining Q2 = Sedentary Q3 = Slight - Casual walking Q4 = Moderate – Gardening Q5 = Heavy - Lifting, shoveling Outcome measure: Self-reported T2D (clinically recognized)		
D & B score = 13					



**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

Dziura et al 2004 [346]	To determine the prospective relation between reports of habitual PA, 3-year change in body weight, and the subsequent risk of T2D in an older cohort.	<ul style="list-style-type: none"> <li>• n = 2,135</li> </ul>	PA assessment: Questionnaire for 4 types of activities (walking, gardening/housework, physical exercises, active sports or swimming) and frequency of participation measured with a PA score:	118 cases of T2D	Observation of an inverse relationship between reported PA and rate of T2DM.
USA		<ul style="list-style-type: none"> <li>• Sex: Men and women</li> </ul>		Incident density of T2D = 6.6/1000 person years	
Prospective cohort		<ul style="list-style-type: none"> <li>• Age: <math>\geq 65</math> yr</li> <li>• Ethnicity: 83% White, 15% African American, 2% Non-white</li> </ul>		Diabetes (n = 118) PA score: $2.17 \pm 1.7$ 'Some' PA: 78%	Subjects reporting some PA at baseline experienced a rate of T2D over 50% lower relative to those reporting no PA.
D & B score = 12		<ul style="list-style-type: none"> <li>• Characteristics: Healthy asymptomatic</li> </ul>	Never (score 0) Sometimes (score 1) Often (score 2) Pearson product moment correlation coefficient and Cox proportional HR	Non-Diabetes (n = 2017) PA score: $2.34 \pm 1.7$ 'Some' PA: 84%	
Hu et al. 1999 [347]	To quantify the dose-response relationship between total PA and incidence of T2D in women.	<ul style="list-style-type: none"> <li>• n = 70,102</li> </ul>	8 yr of follow-up	Number of cases: 1419	Increased PA is associated with substantial reduction in risk of T2D including PA of moderate intensity and duration.
USA		<ul style="list-style-type: none"> <li>• Sex: Women</li> </ul>	PA assessment: Questionnaire for TPA (MET hr/wk) and VPA (6 METs)	Multivariate-adjusted RR (95% CI) of by TPA	
Prospective cohort	To examine the health benefits of walking in comparison to more vigorous activity.	<ul style="list-style-type: none"> <li>• Age: 40-65 yr</li> <li>• Characteristics: participants had no history of diabetes, CVD, or cancer</li> </ul>	TPA (MET hr/wk)	<ul style="list-style-type: none"> <li>• Q1 = 1.0 (referent)</li> <li>• Q2 = 0.77 (0.66-0.90)</li> </ul>	
D & B score = 12		Nurses' Health Study	Q1 = 0-2.0 Q2 = 2.1-4.6 Q3 = 4.7-10.4 Q4 = 10.5-21.7 <ul style="list-style-type: none"> <li>• Q5 = <math>\geq 21.8</math></li> </ul> MET score	<ul style="list-style-type: none"> <li>• Q3 = 0.75 (0.65-0.88)</li> <li>• Q4 = 0.62 (0.52-0.73)</li> <li>• Q5 = 0.54 (0.45-0.64)</li> </ul> Trend $p < 0.001$	
			Q1 = $\leq 0.5$ Q2 = 0.6-2.0 Q3 = 2.1-3.8 Q4 = 3.9-9.9 Q5 = $\geq 10.0$	Multivariate-adjusted RR (95% CI) among women who did not perform vigorous exercise (MET's): <ul style="list-style-type: none"> <li>• Q1 = 1.0 (referent)</li> <li>• Q2 = 0.91 (0.75-1.09)</li> <li>• Q3 = 0.73 (0.59-0.90)</li> <li>• Q4 = 0.69 (0.56-0.86)</li> <li>• Q5 = 0.58 (0.46-0.73)</li> </ul>	

**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

			Outcome measures: Incidence of T2D	Trend p < 0.001	
Hu et al 2001 [348]	To examine the role of prolonged television watching on the risk for T2D.	• n = 37,918	10 year follow-up	Number of cases: 1058	Increasing PA is associated with a significant reduction in risk for T2D, whereas a sedentary lifestyle indicated by prolonged TV watching is related directly to increased risk.
USA		• Sex: Men • Age: 40-75 yr	PA assessment: Questionnaire for PA (MET hr/wk) and TV watching (h/wk), each divided into quintiles	Multivariate-adjusted RR (95% CI) by PA	
Prospective cohort		• Characteristics: participants had no history of diabetes, CVD, or cancer	Q1 = 0-5.9	• Q1 = 1.00 (referent)	
D & B score = 11		• Health Professionals' Follow-up Study	Q2 = 6.0-13.7	• Q2 = 0.78 (0.66 - 0.93)	
			Q3 = 13.8-24.2	• Q3 = 0.65 (0.54 - 0.78)	
			Q4 = 24.3-40.8	• Q4 = 0.58 (0.48 - 0.70)	
			Q5 = ≥ 40.9	• Q5 = 0.51 (0.41 - 0.63)	
			Time spent watching television per week (h/wk)	Trend p < 0.001	
			Q1 = 0-1	Multivariate-adjusted RR (95% CI) by TV time	
			Q2 = 2-10	• Q1 = 1.00 (referent)	
			Q3 = 11-20	• Q2 = 1.66 (1.15 - 2.39)	
			Q4 = 21-40	• Q3 = 1.64 (1.12 - 2.41)	
			Q5 = >40	• Q4 = 2.16 (1.45 - 3.22)	
				• Q5 = 2.87 (1.46 - 5.65)	
				Trend p < 0.001	
Rana et al 2007 [349]	To examine the individual and combined association of obesity and physical inactivity with the incidence of T2D.	• n = 68,907	16 yr follow-up	Number of cases: 4,030	This study found that obesity and physical inactivity independently contributed to the development of T2D.
USA		• Sex: Women			
Prospective cohort		• Age: 30-55 years age range in 1976 (note: 1986 was the baseline year for the study)	PA assessment: Questionnaire for average amount of time/week MET hours per week spent in MVPA (≥ 3 METs), divided into quintiles	Multivariate-adjusted RR (95% CI) by MVPA:	The benefits of PA were not limited to lean women; among those who were overweight and obese, physically active women tended to be at lower risk for T2D than sedentary women.

**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

D & B score = 12		<ul style="list-style-type: none"> <li>• Characteristics: No history of diabetes, CVD or cancer</li> <li>• Nurses' Health Study</li> </ul>	Q1 = <2.1  Q2 = 2.1-4.6 Q3 = 4.7-10.4 Q4 = 10.5-21.7 Q5 = ≥ 21.8	<ul style="list-style-type: none"> <li>• Q1 = 2.37 (2.15-2.16)</li> <li>• Q2 = 1.92 (1.73-2.13)</li> <li>• Q3 = 1.48 (1.34-1.64)</li> <li>• Q4 = 1.40 (1.26-1.55)</li> <li>• Q5 = 1.00 (referent)</li> </ul> Trend $p < 0.001$	
			Cox proportional HR		
Sawada et al 2003 [350]	To examine the association between PF and the incidence of T2D.	• n = 4,747	14 yr follow-up	Number of cases: 280	Low PF is associated with a higher risk for the development of T2D.
Japan		<ul style="list-style-type: none"> <li>• Sex: Men</li> <li>• Age: 20-40 yr</li> </ul>	PF assessment: Maximal aerobic power estimate ml/kg/min using a submaximal cycle ergometer test, divided into quartiles	Age-adjusted RR (95% CI)	
Prospective cohort		<ul style="list-style-type: none"> <li>• Characteristics: Free of diabetes, CVD, hypertensin, tuberculosis, and gastrointestinal disease at baseline</li> </ul>	Q1 = 32.4 ± 3.1	• Q1 = 1.00 (referent)	
D & B score = 13			Q2 = 38.0 ± 2.5 Q3 = 42.4 ± 3.0 Q4 = 51.1 ± 6.2	<ul style="list-style-type: none"> <li>• Q2 = 0.56 (0.42- 0.75)</li> <li>• Q3 = 0.35 (0.25- 0.50)</li> <li>• Q4 = 0.25 (0.17- 0.37)</li> </ul> Trend $p < 0.001$	
			Outcome measure: Incidence of T2D	Multivariate adjusted RR (95% CI)	
				<ul style="list-style-type: none"> <li>• Q1 = 1.00 (referent)</li> <li>• Q2 = 0.78 (0.58-1.05)</li> <li>• Q3 = 0.63 (0.45-0.89)</li> <li>• Q4 = 0.56 (0.37-0.84)</li> </ul> Trend $p = 0.001$	
			Cox proportional HR		
Weinstein et al 2004 [351]	To examine the relative contributions and joint association of PA and BMI with T2D.	• n = 37,878	6.9 year follow up	Number of cases: 1,361	Although BMI and physical inactivity are independent predictors of incident diabetes, the magnitude of the association with BMI was greater than with PA in combined analyses. These findings underscore the critical importance of adiposity as a determinant of T2D.

**Table 17: The relationship between physical activity and the development of type 2 diabetes. (Continued)**

USA	<ul style="list-style-type: none"> <li>• Sex: Women</li> </ul>	PA assessment: Questionnaire for walking per week (h/wk) and TPA (kcal/wk), divided into groups and quartiles respectively	Multivariate-adjusted HR (95% CI) by time spent walking
Prospective cohort	<ul style="list-style-type: none"> <li>• Age: 45+ years</li> <li>• Health care professionals</li> <li>• Characteristics: No history of CVD, cancer or diabetes</li> </ul>		<ul style="list-style-type: none"> <li>• G1 = 1.00 (referent)</li> <li>• G2 = 0.95 (0.82-1.10)</li> <li>• G3 = 0.87 (0.73 -1.02)</li> </ul>
D & B score = 12		Walking per week (h/wk)	• G4 = 0.66 (0.54-0.81)
		G1 = no walking	• G5 = 0.89 (0.73-1.09)
		G2 = <1	Trend $p = 0.004$
		G3 = 1-1.5	Multivariate-adjusted HR (95% CI) by TPA
		G4 = 2-3	• Q1 = 1.00 (referent)
		G5 = $\geq 4$	• Q2 = 0.91 (0.79-1.06)
		TPA (kcal/wk)	• Q3 = 0.86 (0.74-1.01)
		Q1 < 200	• Q4 = 0.82 (0.70-0.97)
		Q2 = 200-599	Trend $p = 0.01$
		Q3 = 600-1,499	
		Q4 $\geq 1500$	
		Cox proportional HR	

D & B score, Downs and Black quality score; YR, years; PA, physical activity; CHD, coronary heart disease; T2D, type 2 diabetes; LTPA, leisure-time physical activity; g, group; kcal/wk, kilocalories per week; HR, hazard ratio; RR, risk ratio; OR, odds ratio; 95% CI, confidence interval; CVD, cardiovascular disease; OPA, occupational physical activity; PF, physical fitness; MET, metabolic equivalent; MET/wk, metabolic equivalent per week.

exercised vigorously at least once weekly had a 29% lower incidence of type 2 diabetes than individuals who did not exercise regularly [112]. These authors also revealed that physical activity that was sufficient to cause sweating was associated with a lower incidence of type 2 diabetes. Other research has also demonstrated that moderate-to-vigorous physical activity ( $\geq 5.5$  METs for at least 40 min per week) and/or aerobic fitness levels above  $31 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  are associated with a lower risk of type 2 diabetes in middle-aged men [113] with the effect being the greatest in high-risk individuals. Therefore, it would appear that Canada's recommendations for physical activity are sufficient to reduce the risk for type 2 diabetes.

In 2001, Hu et al. [114] reported very interesting and compelling research regarding the role of lifestyle factors in the development of type 2 diabetes. Using data from the Nurses' Health Study, they defined a low-risk group according to five lifestyle factors including BMI, a healthy diet, the participation in moderate-to-vigorous physical activity for at least 30 min per day, no current smoking, and the consumption of an average of at least one-half serving of an alcoholic beverage per day. They revealed that the women in the low risk group had a RR for type 2 diabetes of only 0.09 (CI 0.05-0.17) in comparison to the rest of the cohort. They also found that 91% of the cases of type 2 diabetes in this cohort (CI 83-95%) could be attributed to the five lifestyle factors. This research provided compelling evidence that the majority of type 2 diabetes could be prevented through healthy living [115].

As reviewed in Table 17 there is evidence that leisure-time, occupational and commuting-related leisure time activities significantly reduce the risk for the development of type 2 diabetes. For instance, a recent study by Sato and colleagues [116] revealed that the walking distance to work was directly related to the incidence of type 2 diabetes in 8,576 Japanese men followed for 4 years. The risk reduction was approximately 27% in the participants who walked to work for  $\geq 21$  min compared to those who did so for  $\geq 10$  min. These findings are similar to that found by Hu et al. who reported that moderate occupational, commuting and leisure-time physical activities all had a significant inverse relationship to risk in middle-aged men and women [117].

Although ethnicity is often not reported in the current research, the studies examined in our systematic review came from a variety of countries and regions. Data was obtained from studies from the USA, Canada, UK, Japan, and Finland. For instance, Hsia et al. (2005) conducted a prospective 5-year study of 87,907 post-menopausal women, finding a strong graded inverse relationship between physical activity and type 2 diabetes. The relationship was stronger in "Caucasian" than

in minority (African-American, Hispanic or Asian) women. The authors postulated this finding might reflect less precise risk assessments in minority women [118]. As we have outlined previously, further research is clearly warranted that examines the relationship between physical activity and type 2 diabetes in persons of different ethnicities. Moreover, further research is needed to determine the effects of socio-economic status on the observed relationships. Nonetheless, the research is compelling, habitual physical activity appears to be highly effective in the primary prevention of type 2 diabetes.

### Implications

In 1992, the consensus panel from the International Conference on Physical Activity, Fitness and Health (held in Toronto, Canada) [17] indicated that physical activity can effectively reduce the risk for, and incidence of, type 2 diabetes. Over 15 years later, the research is compelling; habitual physical activity is an effective primary preventative strategy against the development of type 2 diabetes [111-113,118-123]. As shown in our analyses, numerous observational studies have revealed that regular physical activity is associated with a lower risk of developing type 2 diabetes [111-113,118-123]. Moreover, increased aerobic fitness is inversely associated with the risk of type 2 diabetes [113,124]. It is also apparent that both aerobic and resistance type activities reduce the risk for type 2 diabetes [125,126].

Although it is difficult to determine the dose-response between physical activity and type 2 diabetes in the majority of the current randomized controlled trials, these trials have revealed important findings. Influential exercise and/or lifestyle intervention trials have demonstrated clearly the health benefits of physical activity/exercise in the prevention of type 2 diabetes. For instance, in the Diabetes Prevention Program (US), 3,234 high-risk participants were randomly assigned to one of three groups: a) a placebo control, b) metformin drug therapy (850 mg twice daily), and c) a lifestyle intervention. The authors revealed that the lifestyle intervention (including physical activity for at least 150 minutes per week) was more effective than metformin (alone) (respective reductions in incidence 58% and 31%) [127]. Similarly, Tuomilehto et al. (2001) conducted a randomized controlled trial with middle-aged, overweight subjects with impaired glucose tolerance (172 males and 350 females). The authors reported a significant reduction in the incidence of type 2 diabetes in the intervention group (which received advice regarding moderate intensity exercise (30 min/day) and dietary control). The lifestyle intervention reduced the risk of type 2 diabetes by approximately 54% in women and 63% in men [128]. In a review of the literature, Williams et al. revealed modest weight loss via diet and

physical activity reduced the incidence of type 2 diabetes in high risk individuals by 40-60% over a 3-4 year period [129]. Collectively, the epidemiological and randomized controlled trials provide compelling evidence supporting the role of habitual physical activity in the primary prevention of type 2 diabetes.

#### **Recommendation #6**

***For a reduced risk for type 2 diabetes, it is recommended that individuals should participate in 30 min or more of moderate to vigorous exercise on most days of the week. [Level 2, Grade A]***

#### **Primary Prevention of Osteoporosis**

The protective effects of physical activity and exercise training on bone health are well documented. In fact, the relationship between indicators of bone health (such as bone mineral density or bone mineral content) and physical activity have been evaluated extensively (see Table 18). Numerous exercise intervention trials have revealed that aerobic and resistance activities have a beneficial effect on bone mineral density across the lifespan [16]. In fact, several systematic reviews of the literature [130-135] and major consensus statements [136] have shown clearly the potential benefits of both aerobic and resistance training on bone health (particularly in post-menopausal women). It has been estimated that exercise interventions prevent or reverse at least 1% of bone loss per year in the lumbar spine and the femoral neck of pre- and post-menopausal women [130,137].

Exercise has also been shown to significantly reduce the risk and/or number of falls in comparison to inactive controls [138-142]. Moreover, fracture risk and/or incidence has been shown to be reduced in active individuals [143-145]. Case-control studies of older persons who suffered a hip fracture have revealed that these individuals had significantly lower physical activity levels throughout adulthood [136,146]. Observational studies have also revealed an inverse relationship between the incidence of fractures and physical activity [147-149]. For instance, Joakimsen et al. revealed lower fracture rates in individuals who performed more weight-bearing activities [148]. Similarly, Kujala et al. [147] in a 21-year prospective study revealed that intense activity was associated with a lower incidence of hip fracture (Hazard Ratio = 0.38, 95% CI = 0.16-0.91). Feskanich et al. (2002) also revealed that moderate physical activity was inversely related to the risk of hip fracture in postmenopausal women [149]. In a review of observational trials, Katzmarzyk and Janssen [20] revealed that the fracture risk was markedly higher in habitually inactive individuals (RR = 1.59 (95% CI = 1.40-1.80)) with a population attributable risk of 24% in Canada.

There is clear evidence that exercise training is of benefit for bone health and accordingly reduces the risk for

osteoporosis. However, remarkably limited research has actually examined the relationship between routine physical activity and the prevalence and/or incidence of osteoporosis (Figure 10). In our systematic search of the osteoporosis literature, a total of 3655 citations were identified during the electronic database search (Figure 7). Of these citations, 1888 were identified in MEDLINE, 236 in EMBASE, 82 in Cochrane, and 481 in the CINAHL/SportDiscus/PsychInfo search. A total of 276 duplicates were found, leaving a total of 2411 unique citations. A total of 2059 articles were excluded after screening, leaving a total of 352 articles for full review. From these articles all 352 were excluded after full-text review. The reasons for exclusion included non-experimental/weak design (n = 87), did not contain three levels of physical activity or not possible to determine dose-response relationship (n = 38), reviews, summaries, meta-analyses (n = 39), not dealing specifically with osteoporosis (n = 21), only on change in bone mineral density (N = 123), clinical population (N = 10), bone metabolism (N = 13), fractures (N = 3), population < 18 yrs (N = 11), and other (N = 7). An additional 2 articles were found through the authors' knowledge of the field.

As identified in our systematic search, the majority of the literature has dealt with the relationship between physical activity and indicators of bone health and/or the incidence of fractures. However, a recent observational trial [150] has provided evidence supporting the ability of physical activity to reduce the incidence of osteoporosis. For instance, Robitaille et al. revealed a dose-response relationship between physical activity level and the prevalence of reported osteoporosis in 8073 women aged  $\geq 20$  yr in the National Health and Nutrition Examination Survey, 1999-2004 [150]. Those performing no physical activity were at a higher risk than those who engaged in moderate (<30 MET hr/wk) and high (>30 MET hr/wk) levels of physical activity. There was a dose-response relationship with the highest physical activity group having the lowest prevalence of osteoporosis. Similarly, Keramat et al. [151] in a case-control investigation revealed that physical activity was protective against the development of osteoporosis.

At this time it is difficult to define clearly the actual dose-response required to cause a reduction in the incidence of osteoporosis. It is clear that bone adaptations to exercise are load dependent and site specific [9,10,16,152]. As such, physical activities that involve significantly loading/impact are often advocated for the prevention of osteoporosis. It has been shown that running 15-20 miles per week is associated with bone mineral accrual or maintenance. Longer distances however may be associated with reduced bone mineral density [136].

Feskanich et al. reported that the risk of hip fracture was lowered by 6% for each increase of 3 MET-hours

**Table 18 Studies examining the relationship between physical activity and osteoporosis.**


Publication Country Study Design Quality Score	Objective	Population	Methods	Outcome	Comments and Conclusions
Robitaille et al 2008 [150]	To assess the relationship between the physician- diagnosed osteoporosis and family history and examine whether osteoporosis risk factors account for this relationship.	• n = 8,073  • Sex: Women • Age: ≥ 20 yrs • Characteristics: American women • Study: NHANES (1999-2004)	PA assessment: Questionnaire. Level of PA was expressed in MET (hr/wk)	Prevalence of reported osteoporosis in US women by PA level	Prevalence of osteoporosis declines with increasing PA in a dose-response manner.
USA			G1 = 0 G2 = <30 G3 = ≥ 30	PA level (% prevalence) • G1 = 11.0 (9.8-12.4) • G2 = 7.1 (6.0-8.4) • G3 = 3.9 (2.8-5.4)  Muscle strengthening activities were expressed in frequency/wk Times/week  p < 0.001	
Cross-sectional			G1 = 0 G2 = <2 G3 = ≥ 2	PA level (age adjusted) • G1 = 8.9 (7.7-10.1) • G2 = 8.4 (7.3-9.7) • G3 = 6.2 (4.4-8.5)  p < 0.01	
D & B score = 10			Criteria for diagnosis of osteoporosis: Self-reported physician diagnosed  Chi-square	Muscle strengthening (% prevalence) • G1 = 8.1 (7.2-9.1) • G2 = 3.1 (1.7-5.5) • G3 = 7.4 (5.8-9.4)  p < 0.001	
				Muscle strengthening (age adjusted) • G1 = 7.8 (6.9-8.7) • G2 = 6.7 (3.8-11.8) • G3 = 9.5 (7.6-11.9)  p < 0.05	
Keramat et al 2008 [151]	To assess risk factors for osteoporosis in postmenopausal women from selected BMD centers in Iran and India.	• Iran n = 363; 178 case, 185 control	Study period 2002 – 2005	OR (95% CI) of osteoporosis in exercisers vs. non-exercisers. Iran (age adjusted)	Exercise was shown as protective factor in both countries and it remained significant after adjustment for age weight and height in Iran.

**Table 18: Studies examining the relationship between physical activity and osteoporosis. (Continued)**

Iran and India	• India n = 354; 203 case, 151 control	PA assessment: Questionnaire. PA was categorized as exercises, other exercises (e.g., swimming, aerobics, weight training) and walking	
	• Sex: Women		• Exercises = 0.4 (0.2-0.7)
Case control	• Age: Iran Case = 58.2 (7.1) yr; Iran Control = 55.7 (6.0) yr; India Case = 58.9 (8.1) yr; India Control = 56.4 (7.5) yr	BMD assessment: DEXA	• Other exercises = 0.4 (0.2-0.6)
	• Characteristics: Cases had BMD > 2.5 SD below average of young normal bone density in L1-L4 spine region and/ or total femoral region. Controls had BMD < 1 SD below normal	Multinomial logistic regression	• Regular Walking = 0.5 (0.3- 0.8)
D & B score = 11			Walking and other exercises were shown as protective factors in Iranian subjects.
			Iran (age, weight, height adjusted)
			• Exercises = 0.4 (0.2-0.7)
			• Other exercises = 0.3 (0.2-0.6)
			• Regular Walking = 0.4 (0.2- 0.8)
			India (age adjusted)
			• Exercises = 0.4 (0.3-0.9)
			• Other exercises = NS
			• Regular Walking = NS
			India (age, weight, height adjusted)
			• Exercises = NS
			• Other exercises = NS
			• Regular Walking = 0.4 (0.2- 0.8)
	• Ethnicity: Indian and Iranian		

D & B score, Downs and Black quality score; YR, years; MET/wk, metabolic equivalent per week; G, groups; PA, physical activity; BMD, bone mineral density; SD, standard deviation; DEXA, dual energy x-ray absorptiometry; NS, not significant.





**Figure 10 Results of the Literature Search for Osteoporosis.**

per week of activity (or 1 hr/wk of walking at an average pace) [149]. There was a linear reduction with increasing physical activity level. Walking for at least 4 hr/wk was also associated with a 41% lower risk of hip fracture compared to less than 1 hr/wk [149]. The work of Robitaille et al. also indicated that moderate levels of physical activity are sufficient to reduce the prevalence of osteoporosis [150].

In summary, there is preliminary evidence to indicate that the current Canadian physical activity guidelines are sufficient to maintain and/improve bone health. However,

further research is clearly required, in particular research that examines the relationship between physical activity and the incidence of osteoporosis in both men and women from varied ethnic backgrounds. Currently, the level of evidence would be considered to be at a Level 3A. The quality of the investigations was generally low with a mean (and median) Downs and Black score of 11.

#### *Recommendation #7*

*For a reduced risk for osteoporosis, it is recommended that individuals should participate in load bearing activities for 30 min or more on most days of the week. [Level 3, Grade A]*

#### **Other Considerations**

##### *Musculoskeletal Fitness and Health*

In the present analyses, all indices of physical activity/fitness were incorporated into our systematic reviews. Although the majority of the data is related to aerobic activities, it should be noted that many of these activities also had a significant musculoskeletal component. Moreover, direct measurements of musculoskeletal fitness were included in various studies included in our review. Although there is limited information available (in comparison to aerobic activities) there is compelling evidence that musculoskeletal fitness is also positively associated with health status [9,10,16].

Warburton and colleagues [9,10] in two reviews of the literature reported that enhanced musculoskeletal fitness is associated positively with glucose homeostasis, bone health, functional independence, mobility, psychological well-being, and overall quality of life and negatively associated with fall risk, morbidity and premature mortality. They also reported that interventions that increase musculoskeletal fitness also have a significant positive effect on the health status of the individuals with a low musculoskeletal reserve (e.g., the frail elderly).

In an evaluation of the current literature some key findings emerge. Grip strength has particularly been shown to be inversely related to premature mortality and/or morbidity (e.g., functional limitations) [153-156]. Rantanen et al. (1998) reported that those individuals with the lowest grip strength had a higher rate of mortality at a younger age (over a 27- year period) than their counterparts with higher muscular strength. Furthermore, they revealed that those with a faster rate of decline in muscular strength (>1.5% per year) or a very low grip strength (<21 kg) had a greater incidence of chronic diseases, such as type 2 diabetes, stroke, arthritis, coronary heart disease, and pulmonary disorders. It was shown that those in the lowest grip strength tertile had an 8-fold increased risk for disability. Individuals with high muscular strength have also been shown to develop less functional limitations in comparison to their counterparts with lower strength over a 5-year period [157].

Katzmarzyk and colleagues [126,154,158] in Canada have also demonstrated a positive relationship between musculoskeletal fitness and health status. For instance, Katzmarzyk and Craig (2002) revealed that there was a significantly higher risk of premature mortality in the lower quartile of sit-ups in both men ( $RR = 2.72$ ) and women ( $RR = 2.26$ ). Grip strength was also predictive of mortality in men ( $RR = 1.49$ ), but not women. In a recent study, Mason et al. revealed that musculoskeletal fitness was a significant predictor of weight gain over a 20-year period [158]. Importantly, they also reported that individuals with low musculoskeletal fitness had 78% greater odds of significant weight gain ( $\geq 10$  kg) compared to those with high musculoskeletal fitness. These studies provide direct support for the inclusion of resistance and flexibility training in Canada's physical activity guidelines for adults [3,159].

#### **Recommendation #8**

***For improved health status and reduced risk for chronic disease and disability, it is recommended that individuals should include daily activities that tax the musculoskeletal system [Level 2, Grade A]***

#### **Limitations**

It is important to note that for each chronic condition, the methods used to determine the relationship between physical activity and the specific clinical outcome were often quite varied. For instance, early work in the field generally controlled for few confounding variables (such as age). In comparison, current literature often controls for a myriad of potential confounding variables. These discrepancies make the comparison of the relative risk reductions between studies and across clinical conditions more difficult. Moreover, the multivariate analyses (controlling for various potential confounding factors) may inappropriately decrease the level of risk reduction associated with physical activity and the clinical endpoint [31]. This is owing to the fact that some of the health benefits associated with physical activity may be mediated through these variables [31].

There was often considerable variability in the findings and major conclusions of the studies examined. Often the available literature was limited by the lack of a clear standard for assessing physical activity. In many instances, it was extremely difficult to determine the actual dosage of physical activity used to group the participants. This lack of clarity makes it very difficult to clearly define the dose-response relationship between physical activity and various chronic conditions.

#### **Conclusions**

There is incontrovertible evidence that regular exercise is an effective preventative strategy against premature mortality, cardiovascular disease, stroke, hypertension,

colon cancer, breast cancer, and type 2 diabetes. There is also compelling indirect evidence to support the protective effects of physical activity with respect to osteoporosis. In many instances the dose-response relationship is linear with further health benefits with increasing levels of activity. The current Canadian physical activity guidelines for adults are sufficient to reduce the risk for multiple chronic diseases simultaneously. The acknowledgement that every bit of exercise counts towards health benefits (with greater benefits at higher energy expenditures) is consistent with the literature and a reasonable message to promote to the general population. However, further investigation is likely required to evaluate the relationship between physical activity and health status in non-Caucasian populations.

#### **Acknowledgements**

Production of this paper has been made possible through a financial contribution from the Public Health Agency of Canada. The views expressed herein do not necessarily represent the views of the Public Health Agency of Canada. The leadership and administrative assistance was provided by the Canadian Society for Exercise Physiology (CSEP). Dr. Warburton is supported by a Canadian Institutes of Health Research New Investigator award and a Michael Smith Foundation for Health Research Clinical Scholar award. We are indebted to the work conducted by the staff from the CSEP Health & Fitness Program of BC and Physical Activity Support Line (PAL; <http://www.physicalactivityline.com>) in the systematic review of the literature and the development of tables for this manuscript and the companion paper by Paterson and Warburton [160].

#### **Author details**

<sup>1</sup>Cardiovascular Physiology and Rehabilitation Laboratory, University of British Columbia, Vancouver, Canada. <sup>2</sup>Experimental Medicine Program, Faculty of Medicine, University of British Columbia, Vancouver, Canada. <sup>3</sup>Cognitive and Functional Learning Laboratory, University of British Columbia, Vancouver, Canada.

#### **Authors' contributions**

DW was responsible for the conceptualization and design of the systematic review, the generation of the systematic review terms, oversaw the data collection, evaluated each article included in the review, and was responsible for creating and revising the manuscript. SC was involved in the data collection, the critical review of the articles, the creation of the tables contained in the article and the revision of the manuscript. AI assisted with the data collection, the critical review of the articles, and the creation and the revision of tables in the manuscript. IN assisted with the generation of the systematic review terms, the retrieval of articles, and the generation and revision of the tables. SB was involved in the conceptualization and design of the systematic review, the generation of the systematic review terms, oversaw the data collection, the review of the articles, and was responsible for creating and revising the manuscript. All authors have read and approved the final manuscript.

#### **Competing interests**

The authors declare that they have no competing interests.

Received: 24 July 2009 Accepted: 11 May 2010 Published: 11 May 2010

#### **References**

1. Bouchard C, Shephard RJ: **Physical activity fitness and health: the model and key concepts.** *Physical activity fitness and health: International proceedings and consensus statement* Champaign, IL: Human Kinetics; Bouchard C, Shephard RJ, Stephens T 1994, 77-88.

2. Blair SN, Brodney S: **Effects of physical inactivity and obesity on morbidity and mortality: current evidence and research issues.** *Med Sci Sports Exerc* 1999, **31**:5646-5652.
3. American College of Sports Medicine: **Position stand: Exercise and physical activity for older adults.** *Med Sci Sports Exerc* 1998, **30**:992-1008.
4. McAuley E: **Physical activity and psychosocial outcomes.** *Physical activity, fitness and health: the consensus knowledge* Champaign, IL: Human Kinetics; Bouchard C, Shephard RJ, Stephens T 1994, 551-568.
5. Taylor RS, Brown A, Ebrahim S, Jolliffe J, Noorani H, Rees K, Skidmore B, Stone JA, Thompson DR, Oldridge N: **Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials.** *Am J Med* 2004, **116**:682-692.
6. Blair SN, Cheng Y, Holder JS: **Is physical activity or physical fitness more important in defining health benefits?** *Med Sci Sports Exerc* 2001, **33**: 5379-399, discussion 5419-320.
7. Blair SN, Kohl HW, Paffenbarger RS Jr, Clark DG, Cooper KH, Gibbons LW: **Physical fitness and all-cause mortality. A prospective study of healthy men and women.** *JAMA* 1989, **262**:2395-2401.
8. Paffenbarger RS Jr, Hyde RT, Hsieh CC, Wing AL: **Physical activity, other life-style patterns, cardiovascular disease and longevity.** *Acta Med Scand Suppl* 1986, **771**:85-91.
9. Warburton DE, Gledhill N, Quinney A: **Musculoskeletal fitness and health.** *Can J Appl Physiol* 2001, **26**:217-237.
10. Warburton DE, Gledhill N, Quinney A: **The effects of changes in musculoskeletal fitness on health.** *Can J Appl Physiol* 2001, **26**:161-216.
11. U.S. Department of Health and Human Services: **Healthy People 2000: National Health Promotion and Disease Prevention Objectives.** Washington, D.C.: U.S. Department of Health and Human Services 1991.
12. Puett DW, Griffin MR: **Published trials of nonmedicinal and noninvasive therapies for hip and knee osteoarthritis.** *Ann Intern Med* 1994, **121**:133-140.
13. Shephard RJ: **Absolute versus relative intensity of physical activity in a dose-response context.** *Med Sci Sports Exerc* 2001, **33**:S400-418, discussion S419-420.
14. Lee IM, Skerrett PJ: **Physical activity and all-cause mortality: what is the dose-response relation?** *Med Sci Sports Exerc* 2001, **33**:S459-471, discussion S493-454.
15. Warburton DE, Nicol C, Bredin SS: **Prescribing exercise as preventive therapy.** *CMAJ* 2006, **174**:961-974.
16. Warburton DE, Nicol C, Bredin SS: **Health benefits of physical activity: the evidence.** *CMAJ* 2006, **174**:801-809.
17. Bouchard C, Shephard RJ, Stephens T: **The consensus statement. Physical activity fitness and health: international proceedings and consensus statement** Champaign, IL: Human Kinetics; Bouchard C, Shephard RJ, Stephens T 1994, 9-76.
18. Warburton DER, Katzmarzyk PT, Rhodes RE, Shephard RJ: **Evidence-informed physical activity guidelines for Canadian adults.** *Appl Physiol Nutr Metab* 2007, **32**:S17-74.
19. Warburton DER, Katzmarzyk PT, Rhodes RE, Shephard RJ: **Evidence-informed physical activity guidelines for Canadian adults.** *Can J Pub Health* 2007, **98**:S16-568.
20. Katzmarzyk PT, Janssen I: **The economic costs associated with physical inactivity and obesity in Canada: an update.** *Can J Appl Physiol* 2004, **29**:90-115.
21. Health Canada and Canadian Society for Exercise Physiology: **Canada's Physical Activity Guide to Healthy Active Living.** Ottawa, ON: Health Canada (Cat. No. H39-429/1998-1E) 1998 [http://www.paguide.com].
22. Health Canada and Canadian Society for Exercise Physiology: **Canada's Physical Activity Guide to Healthy Active Living for Older Adults.** Ottawa, ON: Health Canada 1999 [http://www.paguide.com].
23. Health Canada and Canadian Society for Exercise Physiology: **Canada's Physical Activity Guide to Healthy Active Living for Children and Youths.** Ottawa, ON: Health Canada 2002 [http://www.paguide.com].
24. Canadian Society for Exercise Physiology: **Canadian Physical Activity, Fitness and Lifestyle Approach** Ottawa: Canadian Society for Exercise Physiology, 3 2003.
25. Lau DC, Douketis JD, Morrison KM, Hamik IM, Sharma AM, Ur E: **2006 Canadian clinical practice guidelines on the management and prevention of obesity in adults and children (summary).** *CMAJ* 2007, **176**: S1-13.
26. Gotber SC, Tremblay M, Moher D, Gotber B: **A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review.** *Obes Rev* 2007, **8**:307-326.
27. Downs SH, Black N: **The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions.** *J Epidemiol Community Health* 1998, **52**:377-384.
28. Prince SA, Adamo KB, Hamel ME, Hader J, Gotber SC, Tremblay M: **A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review.** *Int J Behav Nutr Phys Act* 2008, **5**:56.
29. Macera CA, Hootman JM, Sniezek JE: **Major public health benefits of physical activity.** *Arthritis Rheum* 2003, **49**:122-128.
30. Macera CA, Powell KE: **Population attributable risk: implications of physical activity dose.** *Med Sci Sports Exerc* 2001, **33**:S635-639, discussion 640-631.
31. Physical Activity Guidelines Advisory Committee: **Physical Activity Guidelines Advisory Committee Report.** Washington, DC: U.S. Department of Health and Human Services 2008, 683.
32. Myers J, Kaykha A, George S, Abella J, Zahner N, Lear S, Yamazaki T, Froelicher V: **Fitness versus physical activity patterns in predicting mortality in men.** *Am J Med* 2004, **117**:912-918.
33. Williams PT: **Physical fitness and activity as separate heart disease risk factors: a meta-analysis.** *Med Sci Sports Exerc* 2001, **33**:754-761.
34. Eriksson G: **Physical fitness and changes in mortality: the survival of the fittest.** *Sports Med* 2001, **31**:571-576.
35. Eriksson G, Liestol K, Bjornholt J, Thaulow E, Sandvik L, Eriksson J: **Changes in physical fitness and changes in mortality.** *Lancet* 1998, **352**:759-762.
36. Blair SN, Kohl HW, Barlow CE, Paffenbarger RS Jr, Gibbons LW, Macera CA: **Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men.** *JAMA* 1995, **273**:1093-1098.
37. Bijnen EC, Feskens EJ, Caspersen CJ, Nagelkerke N, Mosterd WL, Kromhout D: **Baseline and previous physical activity in relation to mortality in elderly men: the Zutphen Elderly Study.** *Am J Epidemiol* 1999, **150**:1289-1296.
38. Johansson SE, Sundquist J: **Change in lifestyle factors and their influence on health status and all-cause mortality.** *Int J Epidemiol* 1999, **28**:1073-1080.
39. Gregg EW, Cauley JA, Stone K, Thompson TJ, Bauer DC, Cummings SR, Ensrud KE: **Relationship of changes in physical activity and mortality among older women.** *JAMA* 2003, **289**:2379-2386.
40. Wannamethee SG, Shaper AG, Walker M: **Changes in physical activity, mortality, and incidence of coronary heart disease in older men.** *Lancet* 1998, **351**:1603-1608.
41. Kujala UM, Kaprio J, Koskenvuo M: **Modifiable risk factors as predictors of all-cause mortality: the roles of genetics and childhood environment.** *Am J Epidemiol* 2002, **156**:985-993.
42. Wessel TR, Arant CB, Olson MB, Johnson BD, Reis SE, Sharaf BL, Shaw LJ, Hanberg E, Sopko G, Kelsey SE, Pepine CJ, Merz NB: **Relationship of physical fitness vs body mass index with coronary artery disease and cardiovascular events in women.** *JAMA* 2004, **292**:1179-1187.
43. Katzmarzyk PT, Church TS, Blair SN: **Cardiorespiratory fitness attenuates the effects of the metabolic syndrome on all-cause and cardiovascular disease mortality in men.** *Arch Intern Med* 2004, **164**:1092-1097.
44. Morris JN, Heady JA: **Mortality in relation to the physical activity of work: a preliminary note on experience in middle age.** *Br J Ind Med* 1953, **10**:245-254.
45. Morris JN, Heady JA, Raffle PA, Roberts CG, Parks JW: **Coronary heart-disease and physical activity of work.** *Lancet* 1953, **265**:1111-1120.
46. Paffenbarger RS Jr, Brand RJ, Sholtz RJ, Jung DL: **Energy expenditure, cigarette smoking, and blood pressure level as related to death from specific diseases.** *Am J Epidemiol* 1978, **108**:12-18.
47. Paffenbarger RS, Hale WE: **Work, activity and coronary heart mortality.** *N Engl J Med* 1975, **292**:545-550.
48. United States Department of Health and Human Services: **Physical activity and health: a report of the Surgeon General** Atlanta, GA: Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion 1996.
49. Booth FW, Gordon SE, Carlson CJ, Hamilton MT: **Waging war on modern chronic diseases: primary prevention through exercise biology.** *J Appl Physiol* 2000, **88**:774-787.

50. Katzmarzyk PT, Gledhill N, Shephard RJ: **The economic burden of physical inactivity in Canada.** *CMAJ* 2000, **163**:1435-1440.
51. Oguma Y, Sesso HD, Paffenbarger RS Jr, Lee IM: **Physical activity and all cause mortality in women: a review of the evidence.** *Br J Sports Med* 2002, **36**:162-172.
52. Katzmarzyk P: **Physical Activity Status and Chronic Diseases.** *ACSM's Resource Manual for Guidelines for Exercise Testing and Prescription* Philadelphia: Lippincott, Williams and Wilkins; 2005, 122-135.
53. Statistics Canada: **Deaths, 2002.** Ottawa: Statistics Canada 2004 [http://www.statcan.ca/Daily/English/040927/d040927a.htm].
54. World Health Organization: **The World Health Report: Reducing risks, promoting healthy life.** Geneva: World Health Organization 2002.
55. Emberson JR, Wincup PH, Morris RW, Wannamethee SG, Shaper AG: **Lifestyle and cardiovascular disease in middle-aged British men: the effect of adjusting for within-person variation.** *Eur Heart J* 2005, **26**:1774-1782.
56. Manson JE, Greenland P, LaCroix AZ, Stefanek ML, Mouton CP, Oberman A, Petri MG, Sheps DS, Pettinger MB, Siscovick DS: **Walking compared with vigorous exercise for the prevention of cardiovascular events in women.** *N Engl J Med* 2002, **347**:716-725.
57. Oguma Y, Shinoda-Tagawa T: **Physical activity decreases cardiovascular disease risk in women: review and meta-analysis.** *Am J Prev Med* 2004, **26**:407-418.
58. Wisloff U, Nilsen TI, Droyvold WB, Morkved S, Stordahl SA, Vatten LJ: **A single weekly bout of exercise may reduce cardiovascular mortality: how little pain for cardiac gain? The HUNT study, Norway.** *Eur J Cardiovasc Prev Rehabil* 2006, **13**:798-804.
59. Lee IM, Rexrode KM, Cook NR, Manson JE, Buring JE: **Physical activity and coronary heart disease in women: is "no pain, no gain" passe?** *JAMA* 2001, **285**:1447-1454.
60. Beilin JA, Colditz GA: **A meta-analysis of physical activity in the prevention of coronary heart disease.** *Am J Epidemiol* 1990, **132**:612-628.
61. Bauman AE: **Updating the evidence that physical activity is good for health: an epidemiological review 2000-2003.** *J Sci Med Sport* 2004, **7**:6-19.
62. Kohl HW: **Physical activity and cardiovascular disease: evidence for a dose response.** *Med Sci Sports Exerc* 2001, **33**:S472-483, discussion S493-474.
63. Paffenbarger RS Jr, Hyde RT, Wing AL, Hsieh CC: **Physical activity, all-cause mortality, and longevity of college alumni.** *N Engl J Med* 1986, **314**:605-613.
64. Schnohr P, Scharfing H, Jensen JS: **Intensity versus duration of walking, impact on mortality: the Copenhagen City Heart Study.** *Eur J Cardiovasc Prev Rehabil* 2007, **14**:72-78.
65. Kushi LH, Flee RM, Folsom AR, Mink PJ, Anderson KE, Sellers TA: **Physical activity and mortality in postmenopausal women.** *JAMA* 1997, **277**:1287-1292.
66. Leon AS, Connett J, Jacobs DR Jr, Rautamaa R: **Leisure-time physical activity levels and risk of coronary heart disease and death. The Multiple Risk Factor Intervention Trial.** *JAMA* 1987, **258**:2388-2395.
67. Paffenbarger RS Jr, Hyde RT, Wing AL, Lee IM, Jung DL, Kampert JB: **The association of changes in physical-activity level and other lifestyle characteristics with mortality among men.** *N Engl J Med* 1993, **328**:538-545.
68. Heart and Stroke Foundation of Canada: **Press Releases - Canadian leads effort to raise awareness of stroke as a global health epidemic.** Ottawa: Heart and Stroke Foundation of Canada 2007.
69. Abbott RD, Curb JD, Rodriguez BL, Masaki KH, Popper JS, Ross GW, Petrovitch H: **Age-related changes in risk factor effects on the incidence of thromboembolic and hemorrhagic stroke.** *J Clin Epidemiol* 2003, **56**:479-486.
70. Gillum RF, Mussolino ME, Ingram DD: **Physical activity and stroke incidence in women and men. The NHANES I Epidemiologic Follow-up Study.** *Am J Epidemiol* 1996, **143**:860-869.
71. Lee CD, Blair SN: **Cardiorespiratory fitness and stroke mortality in men.** *Med Sci Sports Exerc* 2002, **34**:592-593.
72. Hu FB, Stamper MJ, Colditz GA, Ascherio A, Rexrode KM, Willett WC, Manson JE: **Physical activity and risk of stroke in women.** *JAMA* 2000, **283**:2961-2967.
73. Lee IM, Paffenbarger RS Jr: **Physical activity and stroke incidence: the Harvard Alumni Health Study.** *Stroke* 1998, **29**:2049-2054.
74. Lee IM, Hennekens CH, Berger K, Buring JE, Manson JE: **Exercise and risk of stroke in male physicians.** *Stroke* 1999, **30**:1-6.
75. Rankinen T, Church TS, Rice T, Boucard C, Blair SN: **Cardiorespiratory fitness, BMI, and risk of hypertension: the HYPGENE study.** *Med Sci Sports Exerc* 2007, **39**:1687-1692.
76. Pereira MA, Folsom AR, McGovern PG, Carpenter M, Arnett DK, Liao D, Szklo M, Hutchinson RG: **Physical activity and incident hypertension in black and white adults: the Atherosclerosis Risk in Communities Study.** *Prev Med* 1999, **28**:304-312.
77. Haapaneen N, Mänttälä S, Vuori I, Oja P, Pasanen M: **Association of leisure time physical activity with the risk of coronary heart disease, hypertension and diabetes in middle-aged men and women.** *Int J Epidemiol* 1997, **26**:739-747.
78. Paffenbarger RS Jr, Wing AL, Hyde RT, Jung DL: **Physical activity and incidence of hypertension in college alumni.** *Am J Epidemiol* 1983, **117**:245-257.
79. Paffenbarger RS Jr, Lee IM: **Intensity of physical activity related to incidence of hypertension and all-cause mortality: an epidemiological view.** *Blood Press Monit* 1997, **2**:115-123.
80. Hemeleht M, Kujala UM, Kaprio J, Sarna S: **Long-term vigorous training in young adulthood and later physical activity as predictors of hypertension in middle-aged and older men.** *Int J Sports Med* 2002, **23**:178-182.
81. Hu G, Barengo NC, Tuomilehto J, Lakka TA, Nissinen A, Jousilahti P: **Relationship of physical activity and body mass index to the risk of hypertension: a prospective study in Finland.** *Hypertension* 2004, **43**:25-30.
82. Gu D, Wildman RP, Wu X, Reynolds K, Huang J, Chen CS, He J: **Incidence and predictors of hypertension over 8 years among Chinese men and women.** *J Hypertens* 2007, **25**:517-523.
83. Hayashi T, Tsumura K, Suematsu C, Okada K, Fujii S, Endo G: **Walking to work and the risk for hypertension in men: the Osaka Health Survey.** *Ann Intern Med* 1999, **131**:21-26.
84. Nakanishi N, Suzuki K: **Daily life activity and the risk of developing hypertension in middle-aged Japanese men.** *Arch Intern Med* 2005, **165**:214-220.
85. Foy CG, Foley KL, D'Agostino RB Jr, Goff DC Jr, Mayer-Davis E, Wagenknecht LE: **Physical activity, insulin sensitivity, and hypertension among US adults: findings from the Insulin Resistance Atherosclerosis Study.** *Am J Epidemiol* 2006, **163**:921-928.
86. Haennel RG, Lemire F: **Physical activity to prevent cardiovascular disease. How much is enough?** *Can Fam Physician* 2002, **48**:65-71.
87. Fagard RH: **Exercise characteristics and the blood pressure response to dynamic physical training.** *Med Sci Sports Exerc* 2001, **33**:S484-492, discussion S493-484.
88. Pescatello LS, Franklin BA, Fagard R, Farquhar WB, Kelley GA, Ray CA: **American College of Sports Medicine position stand. Exercise and hypertension.** *Med Sci Sports Exerc* 2004, **36**:533-553.
89. Cornelissen VA, Fagard RH: **Effect of resistance training on resting blood pressure: a meta-analysis of randomized controlled trials.** *J Hypertens* 2005, **23**:251-259.
90. National Center for Health Statistics: **Fast Stats: Hypertension.** Hyattsville, MD: Centers for Disease Control and Prevention 2008.
91. Statistics Canada: **CAHSIM.** Ottawa: Statistics Canada 2007.
92. McAlister FA, Woolferton E, Campbell NR: **The Canadian Hypertension Education Program (CHEP) recommendations: launching a new series.** *CMAJ* 2005, **173**:508-509.
93. Joffres MR, Gadhian P, Fodor JG, Petrasovits A, Chockalingam A, Hamet P: **Awareness, treatment, and control of hypertension in Canada.** *Am J Hypertens* 1997, **10**:1097-1102.
94. Fagard RH: **Physical activity in the prevention and treatment of hypertension in the obese.** *Med Sci Sports Exerc* 1999, **31**:562-630.
95. Kelley GA: **Aerobic exercise and resting blood pressure among women: a metaanalysis.** *Prev Med* 1999, **28**:264-275.
96. Whelton SP, Chin A, Xin X, He J: **Effect of aerobic exercise on blood pressure: a meta-analysis of randomized, controlled trials.** *Ann Intern Med* 2002, **136**:493-503.
97. Dickinson HO, Mason JM, Nicolson DJ, Campbell F, Beyer FR, Cook JV, Williams B, Ford GA: **Lifestyle interventions to reduce raised blood pressure: a systematic review of randomized controlled trials.** *J Hypertens* 2006, **24**:215-233.
98. Hamer M, Taylor A, Steptoe A: **The effect of acute aerobic exercise on stress related blood pressure response: a systematic review and meta-analysis.** *Biol Psychol* 2006, **71**:183-190.

99. Fagard RH, Cornelissen VA: **Effect of exercise on blood pressure control in hypertensive patients.** *Eur J Cardiovasc Prev Rehabil* 2007, **14**:12-17.
100. Fagard RH: **Exercise is good for your blood pressure: effects of endurance training and resistance training.** *Clin Exp Pharmacol Physiol* 2006, **33**:853-856.
101. Fagard RH: **Effects of exercise, diet and their combination on blood pressure.** *J Hum Hypertens* 2005, **19**(Suppl 3):S20-24.
102. Cornelissen VA, Fagard RH: **Effects of endurance training on blood pressure, blood pressure-regulating mechanisms, and cardiovascular risk factors.** *Hypertension* 2005, **46**:667-675.
103. Thompson PD, Crouse SF, Goodpaster B, Kelley D, Moyna N, Pescatello L: **The acute versus the chronic response to exercise.** *Med Sci Sports Exerc* 2001, **33**:S438-445, discussion S452-453.
104. Mommhof EM, Elias SG, Vlems FA, Tweel van der I, Schuit AJ, Voskuil DW, van Leeuwen EE: **Physical activity and breast cancer: a systematic review.** *Epidemiology* 2007, **18**:137-157.
105. Lee IM: **Physical activity and cancer prevention—data from epidemiologic studies.** *Med Sci Sports Exerc* 2003, **35**:1823-1827.
106. Rockhill B, Willett WC, Hunter DJ, Manson JE, Hankinson SE, Colditz GA: **A prospective study of recreational physical activity and breast cancer risk.** *Arch Intern Med* 1999, **159**:2290-2296.
107. Sesso HD, Paffenbarger RS Jr, Lee IM: **Physical activity and breast cancer risk in the College Alumni Health Study (United States).** *Cancer Causes Control* 1998, **9**:433-439.
108. Shephard RJ, Fuchser R: **Physical activity and cancer: how may protection be maximized?** *Crit Rev Oncog* 1997, **8**:219-272.
109. Thunne I, Furberg AS: **Physical activity and cancer risk: dose-response and cancer, all sites and site-specific.** *Med Sci Sports Exerc* 2001, **33**:5530-5550, discussion S609-510.
110. Cust AE, Armstrong BK, Friedreich CM, Sliaman N, Bauman A: **Physical activity and endometrial cancer risk: a review of the current evidence, biologic mechanisms and the quality of physical activity assessment methods.** *Cancer Causes Control* 2007, **18**:243-58.
111. Hu FB, Li TY, Colditz GA, Willett WC, Manson JE: **Television watching and other sedentary behaviors in relation to risk of obesity and type 2 diabetes mellitus in women.** *JAMA* 2003, **289**:1785-1791.
112. Manson JE, Nathan DM, Krolewski AS, Stampfer MJ, Willett WC, Hennekens CH: **A prospective study of exercise and incidence of diabetes among US male physicians.** *JAMA* 1992, **268**:63-67.
113. Lynch J, Helmerich SP, Lakka TA, Kaplan JA, Cohen RD, Salonen R, Salonen JT: **Moderately intense physical activities and high levels of cardiorespiratory fitness reduce the risk of non-insulin-dependent diabetes mellitus in middle-aged men.** *Arch Intern Med* 1996, **156**:1307-1314.
114. Hu FB, Manson JE, Stampfer MJ, Colditz G, Liu S, Solomon CG, Willett WC: **Diet, lifestyle, and the risk of type 2 diabetes mellitus in women.** *N Engl J Med* 2001, **345**:790-797.
115. Schulze MB, Hu FB: **Primary prevention of diabetes: what can be done and how much can be prevented?** *Annu Rev Public Health* 2005, **26**:445-467.
116. Saito KK, Hayashi T, Kambe H, Nakamura Y, Harita N, Endo G, Yoneeda T: **Walking to work is an independent predictor of incidence of type 2 diabetes in Japanese men: the Kansai Healthcare Study.** *Diabetes Care* 2007, **30**:2296-2298.
117. Hu G, Qiao Q, Silventoinen K, Eriksson JG, Jousilahti P, Lindstrom J, Valle TT, Nissinen A, Tuomilehto J: **Occupational, commuting, and leisure-time physical activity in relation to risk for Type 2 diabetes in middle-aged Finnish men and women.** *Diabetologia* 2003, **46**:322-329.
118. Hsia J, Wu L, Allen C, Oberman A, Lawson WE, Torrens J, Saford M, Limacher MC, Howard BV: **Physical activity and diabetes risk in postmenopausal women.** *Am J Prev Med* 2005, **28**:19-25.
119. Fosom AR, Kushi LH, Hong CP: **Physical activity and incident diabetes mellitus in postmenopausal women.** *Am J Public Health* 2000, **90**:134-138.
120. Wannamethee SG, Shaper AG, Alberti KG: **Physical activity, metabolic factors, and the incidence of coronary heart disease and type 2 diabetes.** *Arch Intern Med* 2000, **160**:2108-2116.
121. Manson JE, Rimm EB, Stampfer MJ, Colditz GA, Willett WC, Krolewski AS, Rosner B, Hennekens CH, Speizer FE: **Physical activity and incidence of non-insulin-dependent diabetes mellitus in women.** *Lancet* 1991, **338**:774-778.
122. Helmerich SP, Ragland DR, Paffenbarger RS Jr: **Prevention of non-insulin-dependent diabetes mellitus with physical activity.** *Med Sci Sports Exerc* 1994, **26**:824-830.
123. Helmerich SP, Ragland DR, Leung RW, Paffenbarger RS Jr: **Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus.** *N Engl J Med* 1991, **325**:147-152.
124. Wei M, Gibbons LW, Mitchell JB, Kampert JB, Lee CD, Blair SN: **The association between cardiorespiratory fitness and impaired fasting glucose and type 2 diabetes mellitus in men.** *Ann Intern Med* 1999, **130**:89-96.
125. Eriksson J, Tuominen J, Valle T, Sundberg S, Sovijarvi A, Lindholm H, Tuomilehto J, Koivisto V: **Aerobic endurance exercise or circuit-type resistance training for individuals with impaired glucose tolerance?** *Horm Metab Res* 1998, **30**:37-41.
126. Katzmarzyk PJ, Craig CL, Gauvin L, Adiposity, physical fitness and incident diabetes: the physical activity longitudinal study. *Diabetologia* 2007, **50**:538-544.
127. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM: **Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin.** *N Engl J Med* 2002, **346**:393-403.
128. Tuomilehto J, Lindstrom J, Eriksson JS, Valle TT, Hämäläinen H, Ilanne-Parikka P, Keinänen-Kiukkaanniemi S, Laakso M, Louheranta A, Rastas M, Salminen V, Uusitupa M: **Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance.** *N Engl J Med* 2001, **344**:1343-1350.
129. Williamson DF, Vinitor F, Bowman BA: **Primary prevention of type 2 diabetes mellitus by lifestyle intervention: implications for health policy.** *Ann Intern Med* 2004, **140**:951-957.
130. Wolff I, van Croonenborg JJ, Kemper HC, Kostense PJ, Twisk JW: **The effect of exercise training programs on bone mass: a meta-analysis of published controlled trials in pre- and postmenopausal women.** *Osteoporos Int* 1999, **9**:1-12.
131. Berard A, Bravo G, Gauthier P: **Meta-analysis of the effectiveness of physical activity for the prevention of bone loss in postmenopausal women.** *Osteoporos Int* 1997, **7**:331-337.
132. Kelley GA: **Exercise and regional bone mineral density in postmenopausal women: a meta-analytic review of randomized trials.** *Am J Phys Med Rehabil* 1998, **27**:76-87.
133. Kelley GA: **Aerobic exercise and bone density at the hip in postmenopausal women: a meta-analysis.** *Prev Med* 1998, **27**:798-807.
134. Kelley GA, Kelley KS: **Efficacy of resistance exercise on lumbar spine and femoral neck bone mineral density in premenopausal women: a meta-analysis of individual patient data.** *J Womens Health (Larchmt)* 2004, **13**:293-300.
135. Bonaïuti D, Shea B, Iovine R, Negrini S, Robinson V, Kemper HC, Wells G, Tugwell P, Cranney A: **Exercise for preventing and treating osteoporosis in postmenopausal women.** *Cochrane Database Syst Rev* 2002, CD000333.
136. Brown JP, Josse RG: **2002 clinical practice guidelines for the diagnosis and management of osteoporosis in Canada.** *CMAJ* 2002, **167**:S1-34.
137. Wallace BA, Cumming RG: **Systematic review of randomized trials of the effect of exercise on bone mass in pre- and postmenopausal women.** *Calcif Tissue Int* 2000, **67**:10-18.
138. Tinetti ME, Baker DI, McAvay G, Claus EB, Garrett P, Gottschalk M, Koch ML, Trainor K, Horwitz RJ: **A multifactorial intervention to reduce the risk of falling among elderly people living in the community.** *N Engl J Med* 1994, **331**:821-827.
139. Wolf SL, Barnhart HX, Kutner NG, McNeely E, Coogler C, Xu T: **Reducing frailty and falls in older persons: an investigation of Tai Chi and computerized balance training.** *Atlanta FICSIT Group. Frailty and Injuries: Cooperative Studies of Intervention Techniques.* *J Am Geriatr Soc* 1996, **44**:489-497.
140. Carter ND, Khan KM, Pett MA, Heinonen A, Waterman C, Donaldson MG, Janssen PA, Mallinson A, Riddell L, Kruse K, Prior JC, Flicker L, McKay HA: **Results of a 10 week community based strength and balance training programme to reduce fall risk factors: a randomised controlled trial in 65-75 year old women with osteoporosis.** *Br J Sports Med* 2001, **35**:348-351.
141. Liu-Ambrose T, Khan KM, Eng JJ, Janssen PA, Lord SR, McKay HA: **Resistance and agility training reduce fall risk in women aged 75 to 85 with low bone mass: a 6-month randomized, controlled trial.** *J Am Geriatr Soc* 2004, **52**:657-665.

142. Shaw JM, Snow CM: **Weighted vest exercise improves indices of fall risk in older women.** *J Gerontol A Biol Sci Med Sci* 1998, **53**:M53-58.
143. Gregg EW, Pereira MA, Caspersen CJ: **Physical activity, falls, and fractures among older adults: a review of the epidemiologic evidence.** *J Am Geriatr Soc* 2000, **48**:883-893.
144. Stevens JA, Powell KE, Smith SM, Wingo PA, Sartir RW: **Physical activity, functional limitations, and the risk of fall-related fractures in community-dwelling elderly.** *Ann Epidemiol* 1997, **7**:54-61.
145. Garter ND, Kannus P, Khan KM: **Exercise in the prevention of falls in older people: a systematic literature review examining the rationale and the evidence.** *Sports Med* 2001, **31**:427-438.
146. Boyce WJ, Vessey MP: **Habitual physical inertia and other factors in relation to risk of fracture of the proximal femur.** *Age Ageing* 1988, **17**:319-327.
147. Kujala UM, Kaprio J, Kannus P, Sarna S, Koskenvuo M: **Physical activity and osteoporotic hip fracture risk in men.** *Arch Intern Med* 2000, **160**:705-708.
148. Jokimsen RM, Fonnebo V, Magnus JH, Stormer J, Tøllan A, Sogaard AJ: **The Tromsø Study: physical activity and the incidence of fractures in a middle-aged population.** *J Bone Miner Res* 1998, **13**:1149-1157.
149. Feskanich D, Willett W, Colditz G: **Walking and leisure-time activity and risk of hip fracture in postmenopausal women.** *JAMA* 2002, **288**:2300-2306.
150. Roubaille J, Yoon PW, Moore CA, Liu T, Irazary-DeLacruz M, Looker AC, Khoury MJ: **Prevalence, family history, and prevention of reported osteoporosis in U.S. women.** *Am J Prev Med* 2008, **35**:47-54.
151. Karamat A, Panwardhan B, Larjani B, Chopra A, Mithal A, Chakravarty D, Adibi H, Khosravi A: **The assessment of osteoporosis risk factors in Iranian women compared with Indian women.** *BMC Musculoskelet Disord* 2008, **9**:28.
152. Kerr D, Morton A, Dick I, Prince R: **Exercise effects on bone mass in postmenopausal women are site-specific and load-dependent.** *J Bone Miner Res* 1996, **11**:218-225.
153. Rantanen T, Masaki K, Foley D, Izmitlian G, White L, Gurakik JN: **Grip strength changes over 27 yr in Japanese-American men.** *J Appl Physiol* 1998, **85**:2047-2053.
154. Kuzmarzyk PT, Craig CL: **Musculoskeletal fitness and risk of mortality.** *Med Sci Sports Exerc* 2002, **34**:740-744.
155. Metter EJ, Talbot LA, Schrager M, Conwit R: **Skeletal muscle strength as a predictor of all-cause mortality in healthy men.** *J Gerontol A Biol Sci Med Sci* 2002, **57**:359-365.
156. Fujita Y, Nakamura Y, Hiraoaka J, Kobayashi K, Sakata K, Nagai M, Yangawa H: **Physical-strength tests and mortality among visitors to health-promotion centers in Japan.** *J Clin Epidemiol* 1995, **48**:1349-1359.
157. Ball PA, Macera CA, Davis DR, Blair SN, Gordon N: **Muscular strength and physical function.** *Med Sci Sports Exerc* 2000, **32**:412-416.
158. Mason C, Breen SE, Craig CL, Gavin L, Kuzmarzyk PT: **Musculoskeletal fitness and weight gain in Canada.** *Med Sci Sports Exerc* 2007, **39**:38-43.
159. Blair SN, LaMonte MJ, Nichaman MZ: **The evolution of physical activity recommendations: how much is enough?** *Am J Clin Nutr* 2004, **79**:913S-920S.
160. Paterson DH, Warburton DER: **Physical activity and functional limitations in older adults: a systematic review related to Canada's Physical Activity Guidelines.** *Int J Behav Nutr Phys Act* 2010, **7**:38.
161. Stephenson JB, Armstrong A, Smith T, Bellaw B: **The costs of illness attributable to physical inactivity in Australia: A preliminary study.** *Commonwealth Department of Health and Aged Care and the Australian Sports Commission* 2000 [http://www.health.gov.au/internet/main/Publishing.nsf/Content/health/pubhlt-publicat-document-phys\_costofillness-ent.htm/\$FILE/phys\_costofillness.pdf].
162. Colditz GA: **Economic costs of obesity and inactivity.** *Med Sci Sports Exerc* 1999, **31**:5663-667.
163. Andersen LB, Schnohr P, Schroll M, Hein HO: **All-cause mortality associated with physical activity during leisure time, work, sports, and cycling to work.** *Arch Intern Med* 2000, **160**:1621-1628.
164. Barengo NC, Hu G, Lakka TA, Pekkarinen H, Nissinen A, Tuomilehto J: **Low physical activity as a predictor for total and cardiovascular disease mortality in middle-aged men and women in Finland.** *Eur Heart J* 2004, **25**:2204-2211.
165. Bath PA: **Differences between older men and women in the self-rated healthmortality relationship.** *Gerontologist* 2003, **43**:387-395. discussion 372-385.
166. Bijnen FC, Caspersen CJ, Feskens EJ, Smit WH, Mosterd WL, Kromhout D: **Physical activity and 10-year mortality from cardiovascular diseases and all causes: The Zutphen Elderly Study.** *Arch Intern Med* 1998, **158**:1499-1505.
167. Blair SN, Kohl HW, Barlow CE: **Physical activity, physical fitness, and all-cause mortality in women: do women need to be active?** *J Am Coll Nutr* 1993, **12**:368-371.
168. Blair SN, Kampert JB, Kohl HW, Barlow CE, Macera CA, Paffenbarger RS Jr, Gibbons LW: **Influences of cardiorespiratory fitness and other precursors on cardiovascular disease and all-cause mortality in men and women.** *JAMA* 1996, **276**:205-210.
169. Boyle PA, Buchman AS, Wilson RS, Bienias JL, Bennett DA: **Physical activity is associated with incident disability in community-based older persons.** *J Am Geriatr Soc* 2007, **55**:195-201.
170. Bucksch J: **Physical activity of moderate intensity in leisure time and the risk of all cause mortality.** *Br J Sports Med* 2005, **39**:632-638.
171. Bucksch J, Helmert U: **Leisure time sports activity and all-cause-mortality in West-Germany (1984-1998).** *Z Gesundheitswiss* 2004, **12**:351-358.
172. Carlsson S, Andersson T, Wolk A, Ahlbom A: **Low physical activity and mortality in women: baseline lifestyle and health as alternative explanations.** *Scand J Public Health* 2006, **34**:480-487.
173. Crespo CJ, Palmieri MR, Perdomo RP, McGee DL, Smit E, Semplos CT, Lee IM, Sorlie PD: **The relationship of physical activity and body weight with all-cause mortality: results from the Puerto Rico Heart Health Program.** *Ann Epidemiol* 2002, **12**:543-552.
174. Davey Smith G, Shipley MJ, Batty GD, Morris JN, Marmot M: **Physical activity and cause-specific mortality in the Whitehall study.** *Public Health* 2000, **114**:308-315.
175. Eaton CB, Medaile JH, Flocke SA, Zyzanski SJ, Yaari S, Goldbourt U: **Self-reported physical activity predicts long-term coronary heart disease and all-cause mortalities: Twenty-one-year follow-up of the Israeli Ischemic Heart Disease Study.** *Arch Fam Med* 1995, **4**:323-329.
176. Fang J, Wylie-Rosett J, Alderman MH: **Exercise and cardiovascular outcomes by hypertensive status: NHANES I epidemiological follow-up study, 1971-1992.** *Am J Hypertens* 2005, **18**:751-758.
177. Fried LP, Kronmal RA, Newman AB, Bild DE, Mittelman MB, Polak JF, Robbins JA, Gardin JM: **Risk factors for 5-year mortality in older adults: the Cardiovascular Health Study.** *JAMA* 1998, **279**:585-592.
178. Fujita K, Takahashi H, Miura G, Ohkubo T, Sato Y, Ugajin T, Kurashina K, Tsubono Y, Tsuji I, Fukao A, Hisamichi S: **Walking and mortality in Japan: the Miyagi Cohort Study.** *J Epidemiol* 2004, **14**(Suppl 1):S26-32.
179. Glass TA, de Leon CM, Martotoli RA, Berkman LF: **Population based study of social and productive activities as predictors of survival among elderly Americans.** *BMJ* 1999, **319**:478-483.
180. Gulati M, Pandey DK, Amsdorf MF, Lauderdale DS, Thisted RA, Wicklund RH, Al-Hanti AJ, Black HR: **Exercise capacity and the risk of death in women: the St James Women Take Heart Project.** *Circulation* 2003, **108**:1554-1559.
181. Haapainen N, Mänttinen S, Vuori I, Oja P, Pasanen M: **Characteristics of leisure time physical activity associated with decreased risk of premature all-cause and cardiovascular disease mortality in middle-aged men.** *Am J Epidemiol* 1996, **143**:870-880.
182. Hakim AA, Petrovitch H, Burchfiel CM, Ross GW, Rodriguez BL, White LR, Yano K, Curb JD, Abbott RD: **Effects of walking on mortality among nonsmoking retired men.** *N Engl J Med* 1998, **338**:94-99.
183. Hillsdon M, Thorogood M, Murphy M, Jones L: **Can a simple measure of vigorous physical activity predict future mortality? Results from the OXCHECK study.** *Public Health Nutr* 2004, **7**:557-562.
184. Hu G, Tuomilehto J, Silventoinen K, Barengo NC, Pekkonen M, Jousilahti P: **The effects of physical activity and body mass index on cardiovascular, cancer and all-cause mortality among 47 212 middle-aged Finnish men and women.** *Int J Obes (Lond)* 2005, **29**:894-902.
185. Hu FB, Willett WC, Li T, Stampfer MJ, Colditz GA, Manson JE: **Adiposity as compared with physical activity in predicting mortality among women.** *N Engl J Med* 2004, **351**:2694-2703.
186. Kampert JB, Blair SN, Barlow CE, Kohl HW: **Physical activity, physical fitness, and all-cause and cancer mortality: a prospective study of men and women.** *Ann Epidemiol* 1996, **6**:452-457.
187. Kaplan GA, Strawbridge WJ, Cohen RD, Hungerford LR: **Natural history of leisuretime physical activity and its correlates: associations with mortality from all causes and cardiovascular disease over 28 years.** *Am J Epidemiol* 1996, **144**:793-797.

188. Klaw KT, Jakes R, Bingham S, Welch A, Luben R, Day N, Wareham N: **Work and leisure time physical activity assessed using a simple, pragmatic, validated questionnaire and incident cardiovascular disease and all-cause mortality in men and women: The European Prospective Investigation into Cancer in Norfolk prospective population study.** *Int J Epidemiol* 2006, **35**:1034-1043.
189. Kohn HW, Nickman MZ, Frankowski RF, Blair SN: **Maximal exercise hemodynamics and risk of mortality in apparently healthy men and women.** *Med Sci Sports Exerc* 1996, **28**:601-609.
190. Kujala UM, Kaprio J, Sarna S, Koskenvuo M: **Relationship of leisure-time physical activity and mortality: the Finnish twin cohort.** *JAMA* 1998, **279**:440-444.
191. LaCroix AZ, Leveille SG, Hecht JA, Grothaus LC, Wagner EH: **Does walking older adults? J Am Geriatr Soc** 1996, **44**:13-120.
192. Lam TH, Ho SY, Hedley AJ, Mak KH, Leung GM: **Leisure time physical activity and mortality in Hong Kong: case-control study of all adult deaths in 1998.** *Ann Epidemiol* 2004, **14**:391-398.
193. Lan TY, Chang HY, Tai TY: **Relationship between components of leisure physical activity and mortality in Taiwanese older adults.** *Prev Med* 2006, **43**:36-41.
194. Lahtanen JA, Laikka TA, Rauramaa R, Kuhanen R, Vandenlaenen JM, Salonen R, Salonen JT: **Cardiovascular fitness as a predictor of mortality in men.** *Arch Intern Med* 2001, **161**:825-831.
195. Lee IM, Paffenbarger RS Jr: **Associations of light, moderate, and vigorous intensity physical activity with longevity. The Harvard Alumni Health Study.** *Am J Epidemiol* 2000, **151**:293-299.
196. Lee IM, Hsieh CC, Paffenbarger RS Jr: **Exercise intensity and longevity in men. The Harvard Alumni Health Study.** *JAMA* 1995, **273**:1179-1184.
197. Lee IM, Sesso HD, Oguma Y, Paffenbarger RS Jr: **The "weekend warrior" and risk of mortality.** *Am J Epidemiol* 2004, **160**:636-641.
198. Leitmann MF, Park Y, Blair A, Ballard-Barbash R, Mowu T, Hollenbeck AR, Schatzkin A: **Physical activity recommendations and decreased risk of mortality.** *Arch Intern Med* 2007, **167**:2453-2460.
199. Leon AS, Myers MJ, Connett J: **Leisure time physical activity and the 16-year risks of mortality from coronary heart disease and all-causes in the Multiple Risk Factor Intervention Trial (MRFIT).** *Int J Sports Med* 1997, **18**(Suppl 3):S208-215.
200. Lissner L, Bengtsson C, Björkelund C, Wedel H: **Physical activity levels and changes in relation to longevity. A prospective study of Swedish women.** *Am J Epidemiol* 1996, **143**:54-62.
201. Manni TM, Everhart JE, Patel KV, Schoeller DA, Colbert LH, Visser M, Tykarsky F, Bauer DC, Goodpaster BH, Harris TB: **Daily activity energy expenditure and mortality among older adults.** *JAMA* 2006, **296**:171-179.
202. Matthews CE, Jui AL, Shu XO, Li HL, Yang G, Li Q, Gao YT, Zheng W: **Influence of exercise, walking, cycling, and overall nonexercise physical activity on mortality in Chinese women.** *Am J Epidemiol* 2007, **165**:1343-1350.
203. Menotti A, Secareccia F: **Physical activity at work and job responsibility as risk factors for fatal coronary heart disease and other causes of death.** *J Epidemiol Community Health* 1985, **39**:325-329.
204. Mensink GB, Dekeeth M, Mul MD, Schuit AJ, Hofmeister H: **Physical activity and its association with cardiovascular risk factors and mortality.** *Epidemiology* 1996, **7**:391-397.
205. Morgan K, Clarke D: **Customary physical activity and survival in later life: a study in Nottingham, UK.** *J Epidemiol Community Health* 1997, **51**:490-493.
206. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE: **Exercise capacity and mortality among men referred for exercise testing.** *N Engl J Med* 2002, **346**:793-801.
207. Ostbye T, Taylor DH, Jung SH: **A longitudinal study of the effects of tobacco smoking and other modifiable risk factors on ill health in middle-aged and old Americans: results from the Health and Retirement Study and Asset and Health Dynamics among the Oldest Old survey.** *Prev Med* 2002, **34**:334-345.
208. Paffenbarger RS Jr, Kampert JB, Lee IM, Hyde RT, Leung RW, Wing AL: **Changes in physical activity and other lifestyle patterns influencing longevity.** *Med Sci Sports Exerc* 1994, **26**:857-865.
209. Richardson CR, Krista AM, Lantz PM, Hayward RA: **Physical activity and mortality across cardiovascular disease risk groups.** *Med Sci Sports Exerc* 2004, **36**:1923-1929.
210. Rockhill B, Willett WC, Manson JE, Leitzmann MF, Stampfer MJ, Hunter DJ, Colditz GA: **Physical activity and mortality: a prospective study among women.** *Am J Public Health* 2001, **91**:578-583.
211. Rosengren A, Wilhelmsen L: **Physical activity protects against coronary death and deaths from all causes in middle-aged men. Evidence from a 20-year follow-up of the primary prevention study in Goteborg.** *Ann Epidemiol* 1997, **7**:69-75.
212. Schnohr P, Schalling H, Jensen JS: **Changes in leisure-time physical activity and risk of death: an observational study of 7,000 men and women.** *Am J Epidemiol* 2003, **158**:639-644.
213. Schnohr C, Holbjer L, Riegeis M, Leder L, Larsen T, Schultz-Larsen K, Petersen L, Prescott E, Gronbaek M: **Does educational level influence the effects of smoking, alcohol, physical activity, and obesity on mortality? A prospective population study.** *Scand J Public Health* 2004, **32**:250-256.
214. Schnohr P, Lange P, Schalling H, Jensen JS: **Long-term physical activity in leisure time and mortality from coronary heart disease, stroke, respiratory diseases, and cancer: The Copenhagen City Heart Study.** *Eur J Cardiovasc Prev Rehabil* 2006, **13**:173-179.
215. Schooling CM, Lam TH, Li ZB, Ho SY, Chan WM, Ho KS, Tham MK, Cowling BJ, Leung GM: **Obesity, physical activity, and mortality in a prospective Chinese elderly cohort.** *Arch Intern Med* 2006, **166**:1498-1504.
216. Sundquist K, Qvist J, Sundquist J, Johansson SE: **Frequent and occasional physical activity in the elderly: a 12-year follow-up study of mortality.** *Am J Prev Med* 2004, **27**:22-27.
217. Talbot LA, Morrell CH, Fleg JL, Metter EJ: **Changes in leisure time physical activity and risk of all-cause mortality in men and women: the Baltimore Longitudinal Study of Aging.** *Prev Med* 2007, **45**:169-176.
218. Trolle-Lageros Y, Mucci LA, Kunle M, Braaten T, Weiderpass E, Hsieh CC, Sandin S, Lagiou P, Trichopoulos D, Lund E, Adami HO: **Physical activity as a determinant of mortality in women.** *Epidemiology* 2005, **16**:780-785.
219. Villeneuve PJ, Morrison H, Craig CL, Schabuel DE: **Physical activity, physical fitness, and risk of dying.** *Epidemiology* 1998, **9**:626-631.
220. Weller I, Corey P: **The impact of excluding non-leisure energy expenditure on the relation between physical activity and mortality in women.** *Epidemiology* 1998, **9**:632-635.
221. Yu S, Yarnell JW, Sweetnam PM, Murray L: **What level of physical activity protects against premature cardiovascular death? The Caerphilly study.** *Heart* 2003, **89**:502-506.
222. Athien A, Tavani A, Gallus S, La Vecchia C: **Occupational and leisure time physical activity and the risk of nonfatal acute myocardial infarction in Italy.** *Ann Epidemiol* 2004, **14**:461-466.
223. Barry GD, Shipley MJ, Marmot MG, Smith GD: **Leisure time physical activity and disease-specific mortality among men with chronic bronchitis: evidence from the Whitehall study.** *Am J Public Health* 2003, **93**:817-821.
224. Chen J, Millar WJ: **Health effects of physical activity.** *Health Rep* 1999, **11**, **21**:30(Engl); **21**:31(Fre).
225. Conroy MB, Cook NR, Manson JE, Buring JE, Lee IM: **Past physical activity, current physical activity, and risk of coronary heart disease.** *Med Sci Sports Exerc* 2005, **37**:1251-1256.
226. Dorn JP, Cerny FJ, Epstein LH, Naughton J, Vena JE, Winkelstein W Jr, Schisterman E, Trevisan M: **Work and leisure time physical activity and mortality in men and women from a general population sample.** *Ann Epidemiol* 1999, **9**:366-373.
227. Folsom AR, Arnett DK, Hutchinson RG, Liao F, Clegg LX, Cooper LS: **Physical activity and incidence of coronary heart disease in middle-aged women and men.** *Med Sci Sports Exerc* 1997, **29**:901-909.
228. Fransson E, de Faire U, Ahlbom A, Reuterwall C, Hallqvist J, Alfredsson L: **The risk of acute myocardial infarction: interactions of types of physical activity.** *Epidemiology* 2004, **15**:573-582.
229. Fransson E, de Faire U, Ahlbom A, Reuterwall C, Hallqvist J, Alfredsson L: **The effect of leisure-time physical activity on the risk of acute myocardial infarction depending on body mass index: a population-based case-control study.** *BMC Public Health* 2006, **6**:296.
230. Haapainen-Nieminen N, Mäntyselkä S, Räsänen M, Vuori I, Oja P, Malmberg J: **Body mass index, physical inactivity and low level of physical fitness as determinants of all-cause and cardiovascular disease mortality-16 y follow-up of middle-aged and elderly men and women.** *Int J Obes Relat Metab Disord* 2000, **24**:1465-1474.
231. Kannel WB, Belanger A, D'Agostino R, Israel I: **Physical activity and physical demand on the job and risk of cardiovascular disease and death: the Framingham Study.** *Am Heart J* 1986, **112**:820-825.



232. Kaprio J, Kujala UM, Koskenvuo M, Sarna S: **Physical activity and other risk factors in male twin-pairs discordant for coronary heart disease.** *Atherosclerosis* 2000, **150**:193-200.
233. Lakka TA, Venalainen JM, Rauramaa R, Salonen R, Tuomilehto J, Salonen JT: **Relation of leisure-time physical activity and cardiorespiratory fitness to the risk of acute myocardial infarction.** *N Engl J Med* 1994, **330**:1549-1554.
234. Laakkanen JA, Kurl S, Salonen R, Rauramaa R, Salonen JT: **The predictive value of cardiorespiratory fitness for cardiovascular events in men with various risk profiles: a prospective population-based cohort study.** *Eur Heart J* 2004, **25**:1428-1437.
235. Lee IM, Sesso HD, Paffenbarger RS Jr: **Physical activity and coronary heart disease risk in men: does the duration of exercise episodes predict risk?** *Circulation* 2000, **102**:981-986.
236. Lee IM, Sesso HD, Oguma Y, Paffenbarger RS Jr: **Relative intensity of physical activity and risk of coronary heart disease.** *Circulation* 2003, **107**:1110-1116.
237. Lemaire RN, Siscovick DS, Raghunathan TE, Weinmann S, Arbogast P, Lin DY: **Leisure-time physical activity and the risk of primary cardiac arrest.** *Arch Intern Med* 1999, **159**:686-690.
238. Lemaire RN, Heckbert SR, Psaty BM, Siscovick DS: **Leisure-time physical activity and the risk of nonfatal myocardial infarction in postmenopausal women.** *Arch Intern Med* 1995, **155**:2302-2308.
239. Li TY, Pava JS, Manson JE, Willett WC, Stampfer MJ, Colditz GA, Rexrode KM, Hu FB: **Obesity as compared with physical activity in predicting risk of coronary heart disease in women.** *Circulation* 2006, **113**:499-506.
240. Lopes C, Santos AC, Azevedo A, Maciel MJ, Barros H: **Physical activity and risk of myocardial infarction after the fourth decade of life.** *Rev Port Cardiol* 2005, **24**:1191-1207.
241. Lovasi GS, Lemaire RN, Siscovick DS, Dublin S, Bis JC, Lumley T, Heckbert SR, Smith NL, Psaty BM: **Amount of leisure-time physical activity and risk of nonfatal myocardial infarction.** *Ann Epidemiol* 2007, **17**:410-416.
242. Manson JE, Hu FB, Rich-Edwards JW, Colditz GA, Stampfer MJ, Willett WC, Speizer FE, Hennekens CH: **A prospective study of walking as compared with vigorous exercise in the prevention of coronary heart disease in women.** *N Engl J Med* 1999, **341**:650-658.
243. Mora S, Cook N, Buring JE, Ricker PM, Lee IM: **Physical activity and reduced risk of cardiovascular events: potential mediating mechanisms.** *Circulation* 2007, **116**:2110-2118.
244. O'Connor GT, Hennekens CH, Willett WC, Goldhaber SZ, Paffenbarger RS Jr, Breslow JL, Lee IM, Buring JE: **Physical exercise and reduced risk of nonfatal myocardial infarction.** *Am J Epidemiol* 1995, **142**:1147-1156.
245. Rastogi T, Vaz M, Spiegelman D, Reddy KS, Bhargava AV, Stampfer MJ, Willett WC, Ascherio A: **Physical activity and risk of coronary heart disease in India.** *Int J Epidemiol* 2004, **33**:759-767.
246. Rodriguez BL, Gub JD, Burchfiel CM, Abbott RD, Petrovitch H, Masaki K, Chiu D: **Physical activity and 23-year incidence of coronary heart disease morbidity and mortality among middle-aged men.** *The Honolulu Heart Program.* *Circulation* 1994, **89**:2540-2544.
247. Rothenbacher D, Hoffmeister A, Brenner H, Koenig W: **Physical activity, coronary heart disease, and inflammatory response.** *Arch Intern Med* 2003, **163**:1200-1205.
248. Secaneccia F, Menotti A: **Physical activity, physical fitness and mortality in a sample of middle aged men followed-up 25 years.** *J Sports Med Phys Fitness* 1992, **32**:206-213.
249. Sesso HD, Paffenbarger RS Jr, Lee IM: **Physical activity and coronary heart disease in men: The Harvard Alumni Health Study.** *Circulation* 2000, **102**:975-980.
250. Sundquist K, Qvist J, Johansson SE, Sundquist J: **The long-term effect of physical activity on incidence of coronary heart disease: a 12-year follow-up study.** *Prev Med* 2005, **41**:219-225.
251. Talbot LA, Morrell CH, Metter EJ, Flieg JL: **Comparison of cardiorespiratory fitness versus leisure time physical activity as predictors of coronary events in men aged < or = 65 years and > 65 years.** *Am J Cardiol* 2002, **89**:1187-1192.
252. Tanasescu M, Letzmann MF, Rimm EB, Willett WC, Stampfer MJ, Hu FB: **Exercise type and intensity in relation to coronary heart disease in men.** *JAMA* 2002, **288**:1994-2000.
253. Vatten LJ, Nilsen TI, Romundstad PR, Droyvold WB, Holmen J: **Adiposity and physical activity as predictors of cardiovascular mortality.** *Eur J Cardiovasc Prev Rehabil* 2006, **13**:909-915.
254. Wagner A, Simon C, Evans A, Ferneres J, Montaye M, Ducimetiere P, Anveiler D: **Physical activity and coronary event incidence in Northern Ireland and France: the Prospective Epidemiological Study of Myocardial Infarction (PRIME).** *Circulation* 2002, **105**:2247-2252.
255. Agnarsson U, Thorgerisson G, Sigvaldsson H, Sigfusson N: **Effects of leisure-time physical activity and ventilatory function on risk for stroke in men: the Reykjavik Study.** *Ann Intern Med* 1999, **130**:987-990.
256. Ellekjaer H, Holmen J, Ellekjaer E, Vatten L: **Physical activity and stroke mortality in women. Ten-year follow-up of the Nord-Trøndelag health survey, 1984-1986.** *Stroke* 2000, **31**:14-18.
257. Evenson KR, Rosamond WD, Cai J, Toole JF, Hutchinson RG, Shahar E, Folsom AR: **Physical activity and ischemic stroke risk. The atherosclerosis risk in communities study.** *Stroke* 1999, **30**:1333-1339.
258. Haheim LL, Holme I, Hjermann I, Leren P: **Risk factors of stroke incidence and mortality. A 12-year follow-up of the Oslo Study.** *Stroke* 1993, **24**:1484-1489.
259. Hu G, Sarti C, Jousilahti P, Silventoinen K, Barengo NC, Tuomilehto J: **Leisure time, occupational, and commuting physical activity and the risk of stroke.** *Stroke* 2005, **36**:1994-1999.
260. Kely DK, Wolf PA, Cupples LA, Beiser AS, Kannel WB: **Physical activity and stroke risk: the Framingham Study.** *Am J Epidemiol* 1994, **140**:608-620.
261. Karup LH, Truelsen T, Pedersen A, Løkke H, Lindahl M, Hansen L, Schnohr P, Boysen G: **Level of physical activity in the week preceding an ischemic stroke.** *Cerebrovasc Dis* 2007, **24**:296-300.
262. Kurl S, Laakkanen JA, Rauramaa R, Lakka TA, Sivenius J, Salonen JT: **Cardiorespiratory fitness and the risk for stroke in men.** *Arch Intern Med* 2003, **163**:1682-1688.
263. Wynt PK, Luben RN, Wareham NJ, Welch AA, Bingham SA, Day NE, Khaw KT: **Combined work and leisure physical activity and risk of stroke in men and women in the European prospective investigation into Cancer-Norfolk Prospective Population Study.** *Neuroepidemiology* 2006, **27**:122-129.
264. Noda H, Iso H, Toyoshima H, Date C, Yamamoto A, Kikuchi S, Koizumi A, Kondo T, Watanabe Y, Wada Y, Inaba Y, Tanakoshi A: **Walking and sports participation and mortality from coronary heart disease and stroke.** *J Am Coll Cardiol* 2005, **46**:1761-1767.
265. Paganini-Hill A, Perez Barreto M: **Stroke risk in older men and women: aspirin, estrogen, exercise, vitamins, and other factors.** *J Gen Specif Med* 2001, **4**:18-28.
266. Pitsavos C, Panagiotakos DB, Chrysohou C, Kokkinos P, Menotti A, Singh S, Dontas A: **Physical activity decreases the risk of stroke in middle-age men with left ventricular hypertrophy: 40-year follow-up (1961-2001) of the Seven Countries Study (the Corfu cohort).** *J Hum Hypertens* 2004, **18**:495-501.
267. Sacco RL, Gan R, Boden-Albala B, Lin IF, Kaigman DE, Hauser WA, Shea S, Paik MC: **Leisure-time physical activity and ischemic stroke risk: the Northern Manhattan Stroke Study.** *Stroke* 1998, **29**:380-387.
268. Simonsick EM, Lafferty ME, Phillips CL, Mendes de Leon CF, Kasl SV, Seeman TE, Fillenbaum G, Hebert P, Lenke JH: **Risk due to inactivity in physically capable older adults.** *Am J Public Health* 1993, **83**:1443-1450.
269. Thrift AG, Donnan GA, McNeil JJ: **Reduced risk of intracerebral hemorrhage with dynamic recreational exercise but not with heavy work activity.** *Stroke* 2002, **33**:559-564.
270. Folsom AR, Prineas RJ, Kaye SA, Munger RG: **Incidence of hypertension and stroke in relation to body fat distribution and other risk factors in older women.** *Stroke* 1990, **21**:701-706.
271. Levenstein S, Smith MW, Kaplan GA: **Psychosocial predictors of hypertension in men and women.** *Arch Intern Med* 2001, **161**:1341-1346.
272. Hou L, Ji BT, Blair A, Dai Q, Gao YT, Chow WH: **Commuting physical activity and risk of colon cancer in Shanghai, China.** *Am J Epidemiol* 2004, **160**:860-867.
273. Bouton-Ruault MC, Senesse P, Meance S, Belgitit C, Faivre J: **Energy intake, body mass index, physical activity, and the colorectal adenoma-carcinoma sequence.** *Nutr Cancer* 2001, **39**:50-57.
274. Brownson RC, Chang JC, Davis JR, Smith CA: **Physical activity on the job and cancer in Missouri.** *Am J Public Health* 1991, **81**:639-642.
275. Calton BA, Lacey JV Jr, Schatzkin A, Schaffer C, Colbert LH, Albanes D, Letzmann MF: **Physical activity and the risk of colon cancer among women: a prospective cohort study (United States).** *Int J Cancer* 2006, **119**:385-391.



276. Chao A, Connell CJ, Jacobs EJ, McCullough ML, Patel AV, Calle EE, Cokkinides VE, Thun MJ: Amount, type, and timing of recreational physical activity in relation to colon and rectal cancer in older adults: the Cancer Prevention Study II Nutrition Cohort. *Cancer Epidemiol Biomarkers Prev* 2004, **13**:2187-2195.
277. Colbert LH, Hartman TJ, Mallia N, Limburg PJ, Pietinen P, Virtamo J, Taylor PR, Albanes D: Physical activity in relation to cancer of the colon and rectum in a cohort of male smokers. *Cancer Epidemiol Biomarkers Prev* 2001, **10**:265-268.
278. Dosemeci M, Hayes RB, Vetter R, Hoover RN, Tucker M, Engin K, Unsal M, Blair A: Occupational physical activity, socioeconomic status, and risks of 15 cancer sites in Turkey. *Cancer Causes Control* 1993, **4**:313-321.
279. Friedenreich C, Norat T, Steindorf K, Boutton-Ruault MC, Pischon T, Mazur M, Clavel-Chapelon F, Linseisen J, Boeing H, Bergman M, Johansen NF, Tjønneland A, Overvad K, Mendez M, Quirós JR, Martinez C, Dorosio M, Navarro C, Guirra AB, Bingham S, Kraw KT, Allen N, Key T, Tichopoulos A, Tichopoulos D, Orfanou N, Krogh V, Palli D, Tumino R, Panico S, Vineis P, Bueno-de-Mesquita HB, Peeters PH, Monninkhof E, Beigund G, Manjer J, Ferrari P, Slimani N, Kaaks R, Riboli E: Physical activity and risk of colon and rectal cancers: the European prospective investigation into cancer and nutrition. *Cancer Epidemiol Biomarkers Prev* 2006, **15**:2398-2407.
280. Giovannucci E, Ascherio A, Rimm EB, Colditz GA, Stampfer MJ, Willett WC: Physical activity, obesity, and risk for colon cancer and adenoma in men. *Ann Intern Med* 1995, **122**:327-334.
281. Isomura K, Kono S, Moore MA, Toyomura K, Nagano J, Mizoue T, Milbu R, Tanaka M, Kakeji Y, Maehara Y, Okamura T, Ikejiri K, Futami K, Yasunari Y, Maekawa T, Takenaka K, Ichimya H, Imazumi NI: Physical activity and colorectal cancer: the Fukuoka Colorectal Cancer Study. *Cancer Sci* 2006, **97**:1099-1104.
282. Johnsen NF, Christensen J, Thomsen BL, Olsen A, Loft S, Overvad K, Tjønneland A: Physical activity and risk of colon cancer in a cohort of Danish middle-aged men and women. *Eur J Epidemiol* 2006, **21**:877-884.
283. Larsen IK, Goetzel T, Almendingen K, Hoff G: Lifestyle as a predictor for colonic neoplasia in asymptomatic individuals. *BMC Gastroenterol* 2006, **6**:5.
284. Larsson SC, Ruteigard J, Bergkvist L, Wolk A: Physical activity, obesity, and risk of colon and rectal cancer in a cohort of Swedish men. *Eur J Cancer* 2006, **42**:2590-2597.
285. Lee IM, Paffenbarger RS Jr: Physical activity and its relation to cancer risk: a prospective study of college alumni. *Med Sci Sports Exerc* 1994, **26**:831-837.
286. Lee IM, Manson JE, Ajani U, Paffenbarger RS Jr, Hennekens CH, Buring JE: Physical activity and risk of colon cancer: the Physicians' Health Study (United States). *Cancer Causes Control* 1997, **8**:565-574.
287. Lee KJ, Inoue M, Otani T, Iwasaki M, Sasaki S, Tsugane S: Physical activity and risk of colorectal cancer in Japanese men and women: the Japan Public Health Center-based prospective study. *Cancer Causes Control* 2007, **18**:199-209.
288. Longnecker MP, Gerhardsson le Verdier M, Funkin H, Carpenter C: A case-control study of physical activity in relation to risk of cancer of the right colon and rectum in men. *Int J Epidemiol* 1995, **24**:42-50.
289. Mai PL, Sullivan-Halley J, Ursin G, Stram DO, Deapen D, Villaluna D, Horn-Ross PL, Clarke CA, Reynolds P, Ross RK, West DW, Anton-Culver H, Zogas A, Bernstein L: Physical activity and colon cancer risk among women in the California Teachers Study. *Cancer Epidemiol Biomarkers Prev* 2007, **16**:517-525.
290. Martínez ME, Giovannucci E, Spiegelman D, Hunter DJ, Willett WC, Colditz GA: Leisure-time physical activity, body size, and colon cancer in women. Nurses' Health Study Research Group. *J Natl Cancer Inst* 1997, **89**:948-955.
291. Nilsen TI, Romundstad PR, Petersen H, Gunnell D, Vatten LJ: Recreational physical activity and cancer risk in subites of the colon (the Nord-Trøndelag Health Study). *Cancer Epidemiol Biomarkers Prev* 2008, **17**:183-188.
292. Schnohr P, Gronbaek M, Petersen L, Hein HO, Sorensen TI: Physical activity in leisure-time and risk of cancer: 14-year follow-up of 28,000 Danish men and women. *Scand J Public Health* 2005, **33**:244-249.
293. Slatery ML, Schumacher MC, Smith KR, West DW, Abd-Eghany N: Physical activity, diet, and risk of colon cancer in Utah. *Am J Epidemiol* 1988, **128**:989-999.
294. Slatery ML, Edwards SL, Ma KN, Friedman GD, Potter JD: Physical activity and colon cancer: a public health perspective. *Ann Epidemiol* 1997, **7**:137-145.
295. Slatery ML, Potter J, Cagan B, Edwards S, Coates A, Ma KN, Berry TD: Energy balance and colon cancer-beyond physical activity. *Cancer Res* 1997, **57**:75-80.
296. Takahashi H, Kuriyama S, Tsubono Y, Nakaya N, Fujita K, Nishino Y, Shibuya D, Tsuji I: Time spent walking and risk of colorectal cancer in Japan: the Miyagi Cohort study. *Eur J Cancer Prev* 2007, **16**:403-408.
297. Tang R, Wang JY, Lo SK, Hsieh LL: Physical activity, water intake and risk of colorectal cancer in Taiwan: a hospital-based case-control study. *Int J Cancer* 1999, **82**:484-489.
298. Tavani A, Braga C, La Vecchia C, Conti E, Filiberti R, Montella M, Amadori D, Russo A, Franceschi S: Physical activity and risk of cancers of the colon and rectum: an Italian case-control study. *Br J Cancer* 1999, **79**:1912-1916.
299. Thune I, Lund E: Physical activity and risk of colorectal cancer in men and women. *Br J Cancer* 1996, **73**:1134-1140.
300. Vena JE, Graham S, Zilezny M, Swanson MK, Barnes RE, Nolan J: Lifetime occupational exercise and colon cancer. *Am J Epidemiol* 1985, **122**:357-365.
301. Vetter R, Dosemeci M, Blair A, Wacholder S, Unsal M, Engin K, Fraumeni JF Jr: Occupational physical activity and colon cancer risk in Turkey. *Eur J Epidemiol* 1992, **8**:845-850.
302. White E, Jacobs EJ, Daling JR: Physical activity in relation to colon cancer in middle-aged men and women. *Am J Epidemiol* 1996, **144**:42-50.
303. Wolin KY, Lee IM, Colditz GA, Glynn RJ, Fuchs G, Giovannucci E: Leisure-time physical activity patterns and risk of colon cancer in women. *Int J Cancer* 2007, **121**:2776-2781.
304. Zhang Y, Cantor KP, Dosemeci M, Lynch CF, Zhu Y, Zheng T: Occupational and leisure-time physical activity and risk of colon cancer by subtype. *J Occup Environ Med* 2006, **48**:236-243.
305. Bernstein L, Henderson BE, Hanisch R, Sullivan-Halley J, Ross RK: Physical exercise and reduced risk of breast cancer in young women. *J Natl Cancer Inst* 1994, **86**:1403-1408.
306. Bernstein L, Patel AV, Ursin G, Sullivan-Halley J, Press MF, Deapen D, Berlin JA, Daling JR, McDonald JA, Norman SA, Malone KE, Strom BL, Liff J, Folger SG, Simon MS, Burkman RT, Marchbanks PA, Weiss LK, Spirtas R: Lifetime recreational exercise activity and breast cancer risk among black women and white women. *J Natl Cancer Inst* 2005, **97**:1671-1679.
307. Carpenter CL, Ross RK, Paganini-Hill A, Bernstein L: Lifetime exercise activity and breast cancer risk among post-menopausal women. *Br J Cancer* 1999, **80**:1852-1858.
308. Carpenter CL, Ross RK, Paganini-Hill A, Bernstein L: Effect of family history, obesity and exercise on breast cancer risk among postmenopausal women. *Int J Cancer* 2003, **106**:96-102.
309. Chang SC, Ziegler RG, Dunn B, Stolzenberg-Solomon R, Lacey JV Jr, Huang WY, Schatzkin A, Reding D, Hoover RN, Harige P, Leitzmann MF: Association of energy intake and energy balance with postmenopausal breast cancer in the prostate, lung, colorectal, and ovarian cancer screening trial. *Cancer Epidemiol Biomarkers Prev* 2006, **15**:334-341.
310. Colditz GA, Feskanich D, Chen WY, Hunter DJ, Willett WC: Physical activity and risk of breast cancer in premenopausal women. *Br J Cancer* 2003, **89**:847-851.
311. Coogan PF, Newcomb PA, Clapp RW, Trentham-Dietz A, Baron JA, Longnecker MP: Physical activity in usual occupation and risk of breast cancer (United States). *Cancer Causes Control* 1997, **8**:626-631.
312. Coogan PF, Aschengrau A: Occupational physical activity and breast cancer risk in the upper Cape Cod cancer incidence study. *Am J Ind Med* 1999, **36**:279-285.
313. Dallal CM, Sullivan-Halley J, Ross RK, Wang Y, Deapen D, Horn-Ross PL, Reynolds P, Stram DO, Clarke CA, Anton-Culver H, Zogas A, Peeli D, West DW, Wright W, Bernstein L: Long-term recreational physical activity and risk of invasive and in situ breast cancer: the California teachers study. *Arch Intern Med* 2007, **167**:408-415.
314. Dix MJ, Voorrips LE, Goldboom RA, Brandt van den PA: Baseline recreational physical activity, history of sports participation, and postmenopausal breast carcinoma risk in the Netherlands Cohort Study. *Cancer* 2001, **92**:1638-1649.
315. Dorn J, Vena J, Brasure J, Freudenheim J, Graham S: Lifetime physical activity and breast cancer risk in pre- and postmenopausal women. *Med Sci Sports Exerc* 2003, **35**:278-285.

316. Drake DA: A longitudinal study of physical activity and breast cancer prediction. *Cancer Nurs* 2001, **24**:371-377.
317. Friedenreich CM, Bryant HE, Courneya KS: Case-control study of lifetime physical activity and breast cancer risk. *Am J Epidemiol* 2001, **154**:336-347.
318. Friedenreich CM, Courneya KS, Bryant HE: Relation between intensity of physical activity and breast cancer risk reduction. *Med Sci Sports Exerc* 2001, **33**:1538-1545.
319. Friedenreich CM, Rohan TE: Physical activity and risk of breast cancer. *Eur J Cancer Prev* 1995, **4**:145-151.
320. Gammon MD, Schoenberg JB, Britton JA, Kelsey JL, Coates RJ, Brogan D, Pothsman N, Swanson CA, Daling JR, Stanford JL, Brinton JA: Recreational physical activity and breast cancer risk among women under age 45 years. *Am J Epidemiol* 1998, **147**:273-280.
321. Gilliland FD, Li YF, Baumgartner K, Cumley D, Samet JM: Physical activity and breast cancer risk in hispanic and non-hispanic white women. *Am J Epidemiol* 2001, **154**:442-450.
322. Hsing AW, McLaughlin JK, Cocco P, Co Chien HT, Fraumeni JF Jr: Risk factors for male breast cancer (United States). *Cancer Causes Control* 1998, **9**:269-275.
323. Hu YH, Nagata C, Shimizu H, Kaneda N, Kashiki Y: Association of body mass index, physical activity, and reproductive histories with breast cancer: a case-control study in Gifu, Japan. *Breast Cancer Res Treat* 1997, **43**:65-72.
324. John EM, Horn-Ross PL, Koo J: Lifetime physical activity and breast cancer risk in a multiethnic population: the San Francisco Bay area breast cancer study. *Cancer Epidemiol Biomarkers Prev* 2003, **12**:1143-1152.
325. Kruk J: Lifetime physical activity and the risk of breast cancer: a case-control study. *Cancer Detect Prev* 2007, **31**:18-28.
326. Kruk J: Leisure-time physical activity in relation to the risk of breast cancer. *European Journal of Sports Science* 2007, **7**:81-91.
327. Lahtman PH, Friedenreich C, Schuit AJ, Sakini S, Allen NE, Key TJ, Khaw KT, Bingham S, Peeters PH, Momminkhof E, Bueno-de-Mesquita HB, Wilfrat E, Manjer J, Gonzales CA, Ardanaz E, Amiano P, Quirós JR, Navarro C, Martinez C, Berrio F, Palli D, Turmino R, Panico S, Vineis P, Trichopoulos A, Bamia C, Trichopoulos D, Boeing H, Schulz M, Unseisen J, Chang-Claude J, Chapeau FC, Fournier A, Bourtou-Ruault MC, Tjønneland A, Fors Johnson N, Overvad K, Kaaks R, Riboli E: Physical activity and breast cancer risk: the European Prospective Investigation into Cancer and Nutrition. *Cancer Epidemiol Biomarkers Prev* 2007, **16**:36-42.
328. Lee JM, Rexrode KM, Cook NR, Henkens CH, Bunn JE: Physical activity and breast cancer risk: the Women's Health Study (United States). *Cancer Causes Control* 2001, **12**:137-145.
329. Magnusson CM, Roddam AW, Pike MC, Gyllvers C, Crossley B, Hermon C, McPherson K, Peto J, Vessey M, Beral V: Body fitness and physical activity at young ages and the risk of breast cancer in premenopausal women. *Br J Cancer* 2005, **93**:817-824.
330. Malin A, Matthews CE, Shu XO, Cai H, Dai Q, Jin F, Gao YT, Zheng W: Energy balance and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 2005, **14**:1496-1501.
331. Margolis KL, Mucci L, Braaten T, Kumle M, Trolle Lagerros Y, Adami HO, Lund E, Weiderpass E: Physical activity in different periods of life and the risk of breast cancer: the Norwegian-Swedish Women's Lifestyle and Health cohort study. *Cancer Epidemiol Biomarkers Prev* 2005, **14**:27-32.
332. McTiernan A, Stanford JL, Weiss NS, Daling JR, Voigt LF: Occurrence of breast cancer in relation to recreational exercise in women age 50-64 years. *Epidemiology* 1996, **7**:598-604.
333. McTiernan A, Kooperberg C, White E, Wilcox S, Coates R, Adams-Campbell LL, Woods N, Ockene J: Recreational physical activity and the risk of breast cancer in postmenopausal women: the Women's Health Initiative Cohort Study. *JAMA* 2003, **290**:1331-1336.
334. Silvera SA, Jain M, Howe GR, Miller AB, Rohan TE: Energy balance and breast cancer risk: a prospective cohort study. *Breast Cancer Res Treat* 2006, **97**:97-106.
335. Patel AV, Calle EE, Bernstein L, Wu AH, Thun MJ: Recreational physical activity and risk of postmenopausal breast cancer in a large cohort of US women. *Cancer Causes Control* 2003, **14**:519-529.
336. Patel AV, Press MF, Meskke K, Calle EE, Bernstein L: Lifetime recreational exercise activity and risk of breast carcinoma in situ. *Cancer* 2003, **98**:2161-2169.
337. Rintala PE, Pukkala E, Paakkulainen HT, Vihko V: Self-experienced physical workload and risk of breast cancer. *Scand J Work Environ Health* 2002, **28**:158-162.
338. Rockhill B, Willett WC, Hunter DJ, Manson JE, Hankinson SE, Spiegelman D, Colditz GA: Physical activity and breast cancer risk in a cohort of young women. *J Natl Cancer Inst* 1998, **90**:1155-1160.
339. Slattery ML, Edwards S, Murttaugh MA, Sweeney C, Herrick J, Byers T, Giuliano AR, Baumgartner KB: Physical activity and breast cancer risk among women in the southwestern United States. *Ann Epidemiol* 2007, **17**:342-353.
340. Sprague BL, Trentham-Dietz A, Newcomb PA, Titus-Ernstoff L, Hampton JM, Egan KM: Lifetime recreational and occupational physical activity and risk of in situ and invasive breast cancer. *Cancer Epidemiol Biomarkers Prev* 2007, **16**:236-243.
341. Steindorf K, Schmidt M, Kropp S, Chang-Claude J: Case-control study of physical activity and breast cancer risk among premenopausal women in Germany. *Am J Epidemiol* 2003, **157**:121-130.
342. Tehard B, Friedenreich CM, Oppert JM, Clavel-Chapelon F: Effect of physical activity on women at increased risk of breast cancer: results from the E3N cohort study. *Cancer Epidemiol Biomarkers Prev* 2006, **15**:57-64.
343. Thune I, Brenn T, Lund E, Gaard M: Physical activity and the risk of breast cancer. *N Engl J Med* 1997, **336**:1269-1275.
344. Zheng W, Shu XO, McLaughlin JK, Chow WH, Gao YT, Biot W: Occupational physical activity and the incidence of cancer of the breast, corpus uteri, and ovary in Shanghai. *Cancer* 1993, **71**:3620-3624.
345. Burchfiel CM, Sharp DS, Curb JD, Rodriguez BL, Huang LJ, Marcus EB, Yano K: Physical activity and incidence of diabetes: the Honolulu Heart Program. *Am J Epidemiol* 1995, **141**:360-368.
346. Dzura J, Kasi SV, DiPietro L: Physical activity reduces type 2 diabetes risk in aging independent of body weight change. *J Phys Activity Health* 2004, **1**:19-28.
347. Hu FB, Sigal RJ, Rich-Edwards JW, Colditz GA, Solomon CG, Willett WC, Speizer FE, Manson JE: Walking compared with vigorous physical activity and risk of type 2 diabetes in women: a prospective study. *JAMA* 1999, **282**:1433-1439.
348. Hu FB, Leitzmann MF, Stampfer MJ, Colditz GA, Willett WC, Rimm EB: Physical activity and television watching in relation to risk for type 2 diabetes mellitus in men. *Arch Intern Med* 2001, **161**:1542-1548.
349. Rana JS, Li TY, Manson JE, Hu FB: Adiposity compared with physical inactivity and risk of type 2 diabetes in women. *Diabetes Care* 2007, **30**:53-58.
350. Sawada SS, Lee JM, Muto T, Matsuzaki K, Blair SN: Cardiorespiratory fitness and the incidence of type 2 diabetes: prospective study of Japanese men. *Diabetes Care* 2003, **26**:2918-2922.
351. Weinstein AR, Sesso HD, Lee JM, Cook NR, Manson JE, Buring JE, Gaziano JT: Relationship of physical activity vs body mass index with type 2 diabetes in women. *JAMA* 2004, **292**:1188-1194.

doi:10.1186/1479-5868-7-39

**Cite this article as:** Warburton et al.: A systematic review of the evidence for Canada's Physical Activity Guidelines for Adults. *International Journal of Behavioral Nutrition and Physical Activity* 2010 **7**:39.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

