

Cardiovascular, Pulmonary and Renal Pathology

$\alpha 7$ Nicotinic Acetylcholine Receptor Regulates Airway Epithelium Differentiation by Controlling Basal Cell Proliferation

Kamel Maouche,^{*†‡} Myriam Polette,^{*†§}
Thomas Jolly,^{*†} Kahina Medjber,^{*†}
Isabelle Cloëz-Tayarani,[‡] Jean-Pierre Changeux,[‡]
Henriette Burlet,^{*†} Christine Terryn,[†]
Christelle Coraux,^{*†} Jean-Marie Zahm,^{*†}
Philippe Birembaut,^{*†§} and Jean-Marie Tournier^{*†}

From the Institut National de la Santé et de la Recherche Médicale (INSERM) U903,* Reims; IFR53, UMR-S903,† Université de Reims Champagne Ardenne, Reims; the Département de Neurosciences,‡ Institut Pasteur, Paris; and the Centre Hospitalier Universitaire Reims,§ Hôpital Maison Blanche, Laboratoire Pol Bouin, Reims, France

Airway epithelial basal cells are known to be critical for regenerating injured epithelium and maintaining tissue homeostasis. Recent evidence suggests that the $\alpha 7$ nicotinic acetylcholine receptor (nAChR), which is highly permeable to Ca^{2+} , is involved in lung morphogenesis. Here, we have investigated the potential role of the $\alpha 7$ nAChR in the regulation of airway epithelial basal cell proliferation and the differentiation of the human airway epithelium. *In vivo* during fetal development and *in vitro* during the regeneration of the human airway epithelium, $\alpha 7$ nAChR expression coincides with epithelium differentiation. Inactivating $\alpha 7$ nAChR function *in vitro* increases cell proliferation during the initial steps of the epithelium regeneration, leading to epithelial alterations such as basal cell hyperplasia and squamous metaplasia, remodeling observed in many bronchopulmonary diseases. The regeneration of the airway epithelium after injury in $\alpha 7^{-/-}$ mice is delayed and characterized by a transient hyperplasia of basal cells. Moreover, 1-year-old $\alpha 7^{-/-}$ mice more frequently present basal cells hyperplasia. Modulating nAChR function or expression shows that only $\alpha 7$ nAChR, as opposed to heteropentameric $\alpha_x\beta_y$ nAChRs, controls the proliferation of human airway epithelial basal cells. These findings suggest that $\alpha 7$ nAChR is a key regulator of the plasticity of the human airway epithelium by controlling basal cell proliferation and differentiation pathway

and is involved in airway remodeling during bronchopulmonary diseases. (Am J Pathol 2009, 175:1868–1882; DOI: 10.2353/ajpath.2009.090212)

The respiratory epithelium, which is constantly exposed to airborne pollutants, is frequently injured, which results in altered epithelial functions. To restore these functions, the respiratory epithelium must undergo rapid repair via epithelial cell spreading and migration and regenerate its structure via basal cell proliferation and differentiation.¹ These processes are tightly controlled to restore the pseudostratified architecture of the normal mucociliary epithelium. However, in most respiratory diseases, alterations of the regeneration processes induce epithelial remodeling such as hyperplasia, metaplasia, and fibrosis. Understanding the sequence of processes involved in cell proliferation and differentiation is therefore of crucial importance. Both *in vivo* and *in vitro*, human airway basal cells are able to proliferate and reconstitute a fully differentiated and functional epithelium.² These cells are such considered as progenitors of the human airway epithelium and important actors of the airway epithelium regeneration.

The nonneuronal cholinergic system is thought to be involved in the regulation of cell functions such as cell-cell interaction, apoptosis, and proliferation.³ It is now established that human bronchial epithelial cells contain all of the machinery for the production, storage, secretion, and degradation of acetylcholine, which acts as an autocrine or paracrine hormone.^{4,5} Acetylcholine exerts its effects through muscarinic and nicotinic acetylcholine receptors. Nicotinic acetylcholine receptors (nAChRs)

Supported by grants from the French Associations: Association pour la Recherche sur les Nicotianées and Ligue contre le Cancer (Comité de la Marne) (to J.M.T.), Vaincre la Mucoviscidose (to C.C.), the Fond National pour la Santé ACI-2004-2010 INCa (Cancéropôle Grand-Est project), the Lions Club of Soissons and 1 Euro Contre le Cancer (to P.B.), and by the Région Champagne-Ardenne (to T.J., K.M., and K.M.).

Accepted for publication July 22, 2009.

Address reprint requests to Jean-Marie Tournier, Ph.D., INSERM UMRS 903, 45 rue Cognacq-Jay, F-51092 Reims, France. E-mail: jm.tournier@univ-reims.fr.

are composed of five subunits, arranged as α/β heteromeric or α homomeric nAChRs, and assembled around a central ion channel, mediating the influx of Ca^{2+} .⁶ The airway epithelium expresses $\alpha 3$, $\alpha 4$, $\alpha 5$, $\alpha 7$, $\alpha 9$, $\beta 2$, and $\beta 4$ subunits for nAChRs.⁷⁻⁹

$\alpha 7$ nAChR is characterized by an elevated Ca^{2+} permeability¹⁰ and has been involved in several important biological processes such as cell proliferation, apoptosis, and angiogenesis in cancer.^{11,12} Prenatal nicotine exposure significantly increases pulmonary $\alpha 7$ nAChR expression and alters fetal lung development¹³ and subsequently pulmonary function in newborn.¹⁴ In particular, alteration of lung branching morphogenesis induced by nicotine is mediated by $\alpha 7$ nAChR.¹⁵ Altogether, these observations led us to investigate whether the $\alpha 7$ nAChR could be involved in the differentiation of the respiratory epithelium. In the human airway epithelium, we observed $\alpha 7$ nAChR expression in basal cells, which play a critical role in the epithelial regeneration. Both *in vivo* and *in vitro*, the $\alpha 7$ nAChR expression is associated with the airway epithelium differentiation. Moreover, *in vitro* inactivating $\alpha 7$ nAChR or *in vivo* disrupting genetic $\alpha 7$ nAChR expression induces airway epithelium remodeling by modulating basal cell proliferation. This study thus provides several lines of evidence that $\alpha 7$ nAChR is significant for airway epithelial differentiation and suggests that $\alpha 7$ nAChR is a key regulator of the plasticity of the airway epithelium.

Materials and Methods

Reagents

Unless otherwise stated, all reagents were from Sigma-Aldrich Chimie (L'Isle d'Abeau Chesnes, France): mouse anti-cytokeratin (CK)-13 (clone KS-1A3; 1/1000), mouse anti-CK14 (clone LL02; 0.7 $\mu\text{g}/\text{ml}$), mouse anti-Ki-67 (5 $\mu\text{g}/\text{ml}$) (Dako, Trappes, France), mouse anti- β -tubulin (clone KMX-1; 0.1 $\mu\text{g}/\text{ml}$) and mouse anti-glyceraldehyde-3-phosphate dehydrogenase (MAB374; 1 $\mu\text{g}/\text{ml}$) (Millipore, St. Quentin en Yvelines, France), mouse anti-human cystic fibrosis transmembrane conductance regulator (clone 24-1; 2 $\mu\text{g}/\text{ml}$) (R&D Systems, Minneapolis, MN), mouse anti-zonula occludens (ZO)-1 (clone ZO1-1A12; 1/20) (Zymed Laboratories, San Francisco, CA), mouse anti-desmoplakin 1-2 (clone DP1 and 2-2.15; 20 $\mu\text{g}/\text{ml}$) (Harlan Sera-Lab, Loughborough, UK), rabbit polyclonal antibodies anti-nAChR $\alpha 7$ (H-302, lot A2203; 1 $\mu\text{g}/\text{ml}$), anti-caveolin-1 (N-20; 2 $\mu\text{g}/\text{ml}$) and anti- α -ENaC (H-95; 1 $\mu\text{g}/\text{ml}$) and mouse anti-Foxj1 (clone 3-19; 2 $\mu\text{g}/\text{ml}$) (Santa Cruz Biotechnology, Santa Cruz, CA), rabbit anti-transglutaminase-1 (0.2 $\mu\text{g}/\text{ml}$) (Covalab, Cambridge, UK), 4',6'-diamino-2-phenylindole (200 ng/ml), *Griffonia simplicifolia* isolectin B₄-FITC (GSI-B4; 4 $\mu\text{g}/\text{ml}$), Alexa Fluor 488 and Alexa Fluor 594 goat anti-mouse or anti-rabbit (IgG H+L; 1/200) secondary antibodies, α BTX-Alexa Fluor 488 conjugate, rabbit anti-Alexa Fluor 488 (5 $\mu\text{g}/\text{ml}$) and donkey anti-rabbit IgG-Alexa Fluor conjugate (10 $\mu\text{g}/\text{ml}$) (Molecular Probes, Eugene, OR), and α -conotoxin MII was synthesized by Geneprep (Montpellier, France) according to Ref.¹⁶

Human Airway Tissues and Ex Vivo Wound Repair Model

The use of human tissues was authorized by the bioethical law 94-654 of the Public Health Code of France, with a written consent from the patients or their families. Tracheas from normal human fetuses ranging from 10 to 25 weeks of development were obtained after medically induced abortions. Human airway tissues were collected after nasal polypectomy of patients who did not suffer from any other disease. Human bronchial tissues from patients undergoing surgery for bronchial carcinoma were obtained from microscopically normal areas distant from the tumor.

Human bronchial tissue samples were locally injured and the repair process was followed for 4 days as described previously.^{17,18}

Epithelial Cell Isolation and Culture

Human airway epithelial cells (HAECs) were isolated from polyps and bronchial tissues and cultured in an air-liquid interface culture condition as described previously.^{2,19,20} After 5, 7, 9, 11, 16, 23, 30, and 33 days, cells were detached with trypsin and counted with the ADAM-MC apparatus (Labtech, France).

Human airway basal epithelial cells were sorted and separated from secretory and ciliated cells by FACS as described previously.² The sorted cells were composed of 98.4% basal cells as demonstrated by the reanalysis of the sorted cells.

Immunocyto/histochemistry

An indirect immunofluorescence labeling technique was performed on frozen sections of bronchial tissues or cell cultures as described previously.¹⁹ All fluorescence-labeled preparations were examined with an Axiomager microscope (Zeiss, Oberkochen, Germany) equipped with a Coolsnap FX camera (Roper Scientific, Lisses, France). Images were processed with the AxioVision (Zeiss) and Photoshop (Adobe Systems, San Jose, CA) software.

Localization of Binding Sites for α -Bungarotoxin

Binding sites for α -bungarotoxin (α BTX) were studied in bronchial tissue samples according to Ref.²¹ with some modifications, including the following steps: 1) overnight incubation at 4°C with 1 $\mu\text{mol}/\text{L}$ α BTX-Alexa Fluor 488 conjugate, 2) fixation with 4% paraformaldehyde for 20 minutes at room temperature, 3) overnight incubation at 4°C with 5 $\mu\text{g}/\text{ml}$ rabbit anti-Alexa Fluor 488, 4) incubation for 1 hour at room temperature with 10 $\mu\text{g}/\text{ml}$ donkey anti-rabbit IgG-Alexa Fluor 488 conjugate, and 5) counterstaining with hematoxylin. Controls were performed by omitting α BTX-Alexa Fluor 488 conjugate in the first incubation.

Cell Proliferation Assay

Freshly isolated HAECs or FACS sorted basal epithelial cells were seeded on 96-well microplates in a proliferation medium, adapted from Refs.¹⁹ and ²² at a density of 20×10^3 cells/cm², and cultured in the presence of 0–10 μ mol/L α BTX, methyllycaconitine, α -conotoxin MII, or mecamylamine. Cells were then incubated for 1 hour in 1 mg/ml 3-(4,5-dimethylthiazol-2-yl)-2,5-dimethyltetrazolium bromide. The dye was extracted with propanol-2 and the OD at 560 nm was read in a Xenius spectrophotometer (Safas, Monaco).

Transfection of Small Interfering RNA and nAChR α 7 Expression Vector and RT-PCR Analyses

Two α 7 subunit-specific 19-nt sequences were selected in the coding sequence of nAChR α 7 subunit (GenBank accession no. NM000746) to generate 21-nt sense and 21-nt antisense strands of the type (19N) TT (N, any nucleotide). The selected 19-nt sequences were as follows: α 7si1, 5'-CAUCCGAUUCGGUACCAUU-3'; and α 7si2, 5'-GGACAGAUACAUUUACA-3' (Eurogentec, Seraing, Belgium). Two duplexes, which do not recognize any sequence of human genome, were used as control. A total of 20,000 16HBE14o-cells were transiently transfected with 20 nmol/L siRNA duplexes by using the calcium phosphate-precipitation method. Twenty-four hours after transfection, cells were counted and collected for RT-PCR analysis.

Using FuGENE 6 reagent (Roche Diagnostics, Meylan, France), 20,000 16HBE14o-cells were transiently transfected with 1 μ g/ml pCMV6 empty vector or 1 μ g/ml pCMV6-cDNA α 7 (OriGene, Rockville, MD), according to the manufacturer's procedure. The cells were counted and collected for RT-PCR analysis at 1, 3, and 6 days after transfection.

RT-PCR analyses were performed as previously described,²³ using primers for human α 7 nAChR (5'-CAGTCTTACTCTCTTACCGTCT-3' and 5'-GCAC-CAGTTCAGAAGGATGACTC-3'; Eurogentec).

Western Blotting

Quantification of α 7 nAChR by Western blotting was performed as previously described,²³ using the H-302 anti- α 7 nAChR antibody. Subsequent detection of glyceraldehyde-3-phosphate dehydrogenase was performed on the same filters as a control.

Transgenic Mice

All experiments and procedures were performed in compliance with the French Ministry of Agriculture regulations for animal experimentation. Mice lacking the α 7 subunit of the nicotinic receptor (C57BL/6 background) and wild-type littermates were generated as previously reported,²⁴ shipped from Institut Pasteur (Paris, France) and housed

in a sterile animal care facility. Mice, used for the study of the tracheal epithelium regeneration process, were ~8 weeks of age. The phenotype of tracheal epithelium was also investigated in 1-year-old mice.

Induction of Tracheal Epithelial Damage in Mice

Epithelial desquamation was induced in mouse trachea after instillation of povidocanol.²⁵ Thirty-five microliters of 2% povidocanol in PBS was slowly instilled into the nostrils. Animals were sacrificed on days 0 (before polydocanol treatment), 2, 4, 6, 7, and 8, and trachea was removed. The morphology of the tracheal epithelium was studied in 10 hematoxylin-stained sections covering the entire proximal region of the trachea extending 1–3 mm beneath the larynx. The epithelial height was measured in three different areas of the tracheal epithelium.

Characterization of the Tracheal Epithelium Histology in α 7^{+/+} and α 7^{-/-} 1-Year-Old Mice

The morphology of the tracheal epithelium was studied in α 7^{+/+} and α 7^{-/-} 1-year-old mice, in ~10–15 hematoxylin-stained sections covering the entire proximal region of the trachea extending 1–3 mm beneath the larynx. In all sections derived from α 7^{+/+} and α 7^{-/-} mice, three different phenotypes of the surface epithelium were observed. In each section, we assessed the percentage of each phenotype along the total length of the tracheal epithelium covering cartilage.

Results

α 7 nAChR Expression Is Associated with the Differentiation of the Airway Epithelium

When using the H-302 polyclonal antibody or α BTX, a competitive antagonist with a high affinity for the α 7 nAChR,^{10,26,27} we localized the α 7 nAChR in the normal human airway epithelium, both at the apical plasma membrane of columnar ciliated cells and in the pericellular plasma membrane of basal cells (Figure 1, A and B).

To study the relationship between α 7 nAChR expression and the differentiation of the airway epithelium, we studied the expression of α 7 nAChR both *in vivo* in human tracheal tissue samples during fetal development and *in vitro* during the regeneration of human airway epithelium. The human fetal airway epithelium acquires its morphological differentiation at ~12 to 15 weeks of development.²⁸ We observed that, during the fetal development of human airways, α 7 nAChR expression starts at 13 weeks of development, simultaneously with the expression of ZO-1, a marker of epithelial cohesion, of β -tubulin, a marker of the ciliary differentiation and of cystic fibrosis transmembrane conductance regulator, a marker of epithelium functionality for Cl⁻ ion secretion at the apical membrane (Figure 2). The human airway epithelium can be fully regenerated *in vitro* from isolated HAECs maintained in culture in an air-liquid interface condition.²⁰ In




Figure 1. Localization of nAChR $\alpha 7$ in the airway epithelium. nAChR $\alpha 7$ was localized with the H-302 antibody at 1 $\mu\text{g/ml}$ (A) or with AlexaFluor488- αBTX at 1 $\mu\text{mol/L}$ (B) in human bronchial tissue samples, both at the apical side of the epithelium (arrows) and in basal epithelial cells (arrowheads).

this model, we observed a progressive regeneration of pseudostratified epithelium after 4 to 6 weeks (Figure 3A). By using immunofluorescence and RT-PCR techniques, we observed that the expression of $\alpha 7$ nAChR in the basal layers of the epithelium was maximal in steps II and III (appearance of the first ciliated cells) and paralleled the expression of CK14, a marker of airway basal cells,²⁹ suggesting that $\alpha 7$ nAChR expression in basal cells is essential during the initial step of epithelial differentiation and the establishment of the basal cell layer. Thereafter, the expression became progressively restricted to one single layer of basal cells for $\alpha 7$ nAChR or isolated basal cells for CK14 at a time where the epithelium was pseudostratified and continuously expressed β -tubulin at the apical membrane (steps IV and V). Quantification of RT-PCR showed that the amount of $\alpha 7$ nAChR mRNA progressively increased during steps I to III ($P = 0.0183$) and then decreased during steps III to V ($P = 0.0388$; Friedman test) (Figure 3B).

Altogether, these observations indicate that the $\alpha 7$ nAChR expression is related to the differentiation of the airway epithelium and provide a rationale for investigating the implication of $\alpha 7$ nAChR in the differentiation of the airway epithelium.

$\alpha 7$ nAChR Expression/Activity Is Associated in Vivo with the Remodeling of the Airway Epithelium

The knockout mice model gives a good opportunity to investigate the effect of the $\alpha 7$ nAChR absence on the

airway epithelium morphology and differentiation. In all tracheal sections from 1-year-old control and $\alpha 7^{-/-}$ mice, three different phenotypes of the surface epithelium were observed. Phenotype 1 corresponds to a normal epithelium with two layers of nuclei and an epithelium height in the range of 25 to 35 μm . Phenotype 2 corresponds to a normal epithelium with three to four layers of nuclei and an epithelium height in the range of 40 to 55 μm . Phenotype 3 corresponds to a disorganized epithelium, with four to six layers of nuclei, an absence of the superficial columnar cell layer and an epithelium height in the range of 30 to 55 μm (Figure 4A). By using the *Griffonia simplicifolia* isolectin B4 (GSI-B4), which binds specifically to basal cells in the mouse tracheal epithelium,³⁰ we confirmed that phenotype 1 is characterized by one monolayer of basal cells, phenotype 2 contains one/two layers of basal cells, whereas GSI-B4 recognizes three/four layers of basal cells in phenotype 3 (Figure 4B). The percentage of phenotype 2 in the tracheal epithelium was decreased in $\alpha 7^{-/-}$ mice as compared with control mice ($P = 0.0373$), whereas phenotype 3 was 3.8 times more expressed in $\alpha 7^{-/-}$ mice ($P = 0.0174$) (Figure 4C), suggesting that the absence of the $\alpha 7$ nAChR in mice is associated with a more frequently observed hyperplasia of basal cells in the tracheal epithelium.

The allosteric model for nAChRs implies the equilibrium between channel resting, opening and desensitization.³¹ Chronic exposure to nicotine results in both peaks of activation and long-term desensitization of nAChRs, but it is now believed that the overall effect of sustained smoking may result in nAChR desensitization and that chronic smoking maintains this desensitization, even if chronic nicotine exposure is also known to up-regulate nAChRs.³² The $\alpha 7$ nAChR has been shown to be particularly susceptible to desensitization³³⁻³⁵ and to up-regulation on nicotine exposure.^{7,13,36} We thus postulated that in heavy smokers the potential nicotine-induced inactivation of the $\alpha 7$ nAChR may result in alterations in the airway epithelium differentiation. Indeed, in patients with chronic obstructive pulmonary disease (COPD) and with a significant smoking history, the airway epithelium is characterized by secretory cell or basal cell hyperplasia and squamous metaplasia (Figure 5). Hyperplasia of basal cells (area B and D) and squamous metaplasia (area C) are characterized by an overexpression of CK13 and CK14, desmoplakins 1 and 2, and by the absence of β -tubulin at the apex of the epithelium, confirming the absence of ciliated and polarized cells at the luminal surface. Whereas $\alpha 7$ nAChR expression is restricted to one basal cell layer at the margin of the pseudostratified epithelium (Figure 5 D, arrow), $\alpha 7$ nAChR is present up through most basal layers in areas of basal cell hyperplasia or squamous metaplasia (Figure 5, D and C, arrowheads). Similarly to what we observed during the *in vitro* epithelium differentiation (Figure 3 A), $\alpha 7$ nAChR expression in remodeled areas of the airway epithelium is mainly restricted to CK14-expressing basal cells.

These results suggest that the airway epithelium differentiation-associated $\alpha 7$ nAChR expression implicates the differentiation pathway of airway basal cells.




Figure 2. Expression of nAChR $\alpha 7$ during the fetal development of the human airway epithelium. nAChR $\alpha 7$, the protein ZO-1, β -tubulin, and CFTR were localized with Alexa 488 conjugates in human tracheal tissue samples at 11, 13, and 24 weeks of development and in samples from adult patients.

The $\alpha 7$ nAChR Is Involved in the Differentiation Pathway of the Airway Epithelium

To investigate the function of the $\alpha 7$ nAChR in the differentiation of the airway epithelium, we first studied the effect of α BTX, a toxin with a high affinity for the $\alpha 7$ nAChR, on human bronchial tissues undergoing repair. After inducing a local wound in fresh human bronchial tissues (*ex vivo* wound repair model), we observed that basal cells, characterized by the expression of CK13, are able to spread and migrate to progressively cover the wounded area. Similarly, we observed that the $\alpha 7$ nAChR is continuously expressed in basal cells in the unwounded pseudostratified epithelium at the lesion edge and in the repairing area (Figure 6A). Exposure of cultures to α BTX induced a progressive thickening of the epithelium in the repairing areas, with two to three and then four to six cell layers after 3 and 4 days, respectively, as compared with one to three cell layers observed in the control cultures. In both control- and α BTX-exposed cultures, all repairing cells contained CK13, but more cells expressed Ki-67, a proliferation marker, in their nuclei when exposed to α BTX, especially after 4 days in culture (Figure 6B). These observations suggest that the pharmacological inactivation of $\alpha 7$ nAChR up-regulates basal cell proliferation and modulates the early events of epithelium regeneration. To analyze in more details and to quantify these effects, we

used an *in vitro* air-liquid interface culture model of airway epithelium regeneration.

When analyzing the effect of the chronic exposure to α BTX on the regeneration of the human airway epithelium (*in vitro* air-liquid interface culture model), we first observed that α BTX induced a dramatic increase in cell proliferation in the early phases of epithelial regeneration. Indeed, in the presence of 5 μ mol/L α BTX, we observed a significant increase of the cell number ($P < 0.05$) at day 5 (+626%), day 7 (+206%), day 9 (+123%), day 11 (+23%), day 16 (+11%), and day 23 (+10%) (Figure 7A). After 2 days in culture, we identified more Ki-67-positive nuclei in α BTX-exposed cells than in control cells (Figure 7B), suggesting that α BTX was likely to favor cell cycle progression rather than inhibiting apoptosis. Moreover, we observed that all cells expressing Ki-67 in the nucleus, were those expressing CK13, a marker of airway basal cells (Figure 7B), suggesting that α BTX induced basal cell proliferation during the initial step of epithelial regeneration. Figure 7C (control) depicts the evolution of the epithelium morphology during the regeneration process in the absence of α BTX. On day 16, the control cultures exhibit three to four layers of cuboidal undifferentiated cells. On day 23, columnar cells and some ciliated cells (arrowheads) are observed in the upper part of the epithelium. Finally on day 33, the epithelium is pseudostratified, ciliated cells are present all along the





Figure 3. Expression of nAChR $\alpha 7$ during the *in vitro* regeneration of the human airway epithelium. **A:** nAChR $\alpha 7$, CK14 and β -tubulin were localized during the different steps of the *in vitro* differentiation of HAEC cultures in air-liquid interface condition: step I corresponds to 4 to 7 days in culture, step II corresponds to the appearance of the first ciliated cells (day 15–20), step III corresponds to the presence of 50% ciliated cells (day 22–27) and steps IV and V correspond to 1 and 2 weeks, respectively, after step III. **B:** RT-PCR gel (left) and quantification of mRNA band for nAChR $\alpha 7$ subunit, normalized for 28S rRNA values (right) during steps I to V in air-liquid interface cultures. The results are expressed as mean band intensity \pm SD for four different experiments conducted on airway epithelial cell cultures derived from four different patients.

epithelium, CK13 and CK14 and desmoplakins 1 and 2 are essentially expressed in the basal layer of basal cells while β -tubulin and α subunit of the epithelial sodium channel (α -ENaC) are expressed all along the apical membrane of columnar cells, suggesting that the regenerated epithelium is now fully differentiated (Figure 7D, control). When epithelial regeneration was performed in the presence of α BTX, added from day 1, the epithelium was pluristratified and composed of five to six and seven to eight layers of cuboidal cells at days 16 and 23, respectively, with some very flat cells observed at the top of the epithelium on day 23. After 33 days of α BTX exposure, the epithelium still consisted in 8 to 10 layers of cuboidal and polyhedral cells, most of them overexpressing CK13 and CK14 and desmoplakins 1 and 2 in the basal layers whereas the apical expression of β -tubulin and α -ENaC was greatly reduced (Figure 7, C and D, α BTX d1), a situation similar to that observed *in vivo* in

squamous metaplasia of the airway epithelium from smoking patients with COPD (Figure 5C). Moreover, p63, present in airway basal cells and whose expression is decreased during squamous cell differentiation,³⁷ was expressed in the control basal cells but absent in α BTX-exposed cultures. Transglutaminase-1, a well-known marker of squamous differentiation of the respiratory tract,³⁸ is expressed at low level in the control cultures, but overexpressed in the presence of α BTX (Figure 7D). The forkhead transcription factor Foxj1 is required for ciliogenesis and also involved in the organization of the apical membrane.³⁹ In the differentiated pseudostratified epithelium, Foxj1 is present and expressed at the same level in most cell nuclei of the epithelium. In α BTX-exposed cultures, Foxj1 is still present in the basal progenitor cells of the epithelium, but its expression is much lower in the intermediate cells and finally absent in the upper part of the epithelium (Figure 7D). In summary,




Figure 4. Morphological characterization of the tracheal epithelium in $\alpha 7^{+/+}$ and $\alpha 7^{-/-}$ mice. **A:** Histological hematoxylin-stained sections of the upper trachea illustrating the three different epithelial phenotypes observed in 1-year-old mice. **B:** Binding of the FITC-conjugated GSI-B4 lectin to cryosections of mouse trachea. Each panel is matched with the corresponding hematoxylin-stained sections presented at the same level in **A**. **C:** Quantification (mean \pm SD) of each phenotype 1, 2, and 3, expressed as percentage of the total length of the tracheal epithelium in cross-sections of trachea from control $\alpha 7^{+/+}$ mice (white bars, $n = 9$) and from $\alpha 7^{-/-}$ mice (black bars, $n = 6$). The significance of the differences between control $\alpha 7^{+/+}$ mice and $\alpha 7^{-/-}$ mice was determined using the Mann-Whitney test.

adding α BTX from day 1-stimulated cell proliferation during the initial steps of the regeneration and prevented the differentiation of the normal pseudostratified airway epithelium. To investigate in more details the relationship between the initial stimulation of cell proliferation and the subsequent alteration of epithelium differentiation, we tested the effect of adding chronically α BTX from day 11, ie, after $\sim 50\%$ of cell proliferation has already occurred in the control culture (Figure 7A). When adding α BTX from day 11, we did not observe any subsequent modification of cell proliferation and the evolution of epithelium morphology and the distribution of differentiation markers were similar to those observed in the control culture, even after a 22-day exposure to α BTX (Figure 7 A, C, and D, α BTX d11), suggesting that cell proliferation needs to be altered by α BTX, during the first steps of the epithelium regeneration, to induce an alteration in epithelium differentiation.

Finally, we investigated the effect of $\alpha 7$ nAChR absence on the regeneration of the tracheal epithelium in $\alpha 7^{+/+}$ and $\alpha 7^{-/-}$ mice. Desquamation of the mouse tracheal epithelium was induced after instillation of a polydocanol solution.²⁵ Figure 8A depicts the morphology of regenerating tracheal epithelium in $\alpha 7^{+/+}$ and $\alpha 7^{-/-}$ mice from days 0 to 8 after polydocanol instillation. Before injury, both $\alpha 7^{+/+}$ and $\alpha 7^{-/-}$ 8-week-old mice presented the same pseudostratified morphology of the tracheal epithelium with one to two layers of nuclei. On day 2, only flat cells spreading over the denuded basement membrane are observed. On day 4, 3 to 4 and 8 to 10 cell layers are present in the control $\alpha 7^{+/+}$ mice and in $\alpha 7^{-/-}$ mice, respectively, suggesting that an active cell proliferation has occurred with a more significant effect observed in $\alpha 7^{-/-}$ mice. The first columnar cells are observed in $\alpha 7^{+/+}$ mice, whereas the epithelium in $\alpha 7^{-/-}$ mice is still undifferentiated. On day 6, the first basal cells are observed only in $\alpha 7^{+/+}$ mice. On day 7, the epithelium in $\alpha 7^{+/+}$ mice is almost fully differentiated whereas that of $\alpha 7^{-/-}$ mice begins to pseudostratify with the appearance of the first basal cells. On day 8, the epithelium in $\alpha 7^{+/+}$ mice and in $\alpha 7^{-/-}$ mice has a similar pseudostratified structure. We measured the epithelium height during the whole process of regeneration (Figure 8B). On day 4, the epithelium is thicker both in $\alpha 7^{+/+}$ and $\alpha 7^{-/-}$ mice ($P = 0.0339$) as compared with day 0. Then, we observed a progressive and significant decrease of the epithelium height between day 4 and day 8 both in $\alpha 7^{+/+}$ mice ($P = 0.0315$) and in $\alpha 7^{-/-}$ mice ($P = 0.0067$), with a thicker epithelium in $\alpha 7^{-/-}$ mice on day 4 and day 6 ($P = 0.0209$). Finally on day 8, the epithelium height in $\alpha 7^{-/-}$ mice ($33.9 \pm 3.4 \mu\text{m}$) is still higher as compared with day 0 (27.5 ± 1.3 ; $P = 0.0339$), whereas the epithelium height in $\alpha 7^{+/+}$ mice ($30.0 \pm 4 \mu\text{m}$) is similar to that observed on day 0 (28.8 ± 0.8 ; $P > 0.99$). Immunostaining for caveolin-1, a marker of basal cells in the mouse tracheal epithelium,⁴⁰ confirmed a basal cell hyperplasia more pronounced in $\alpha 7^{-/-}$ mice than in $\alpha 7^{+/+}$ mice both on days 4 and 6, whereas a single basal cell layer is observed on day 8 (Figure 8C). These results suggest that the regeneration of the airway epithelium in $\alpha 7^{-/-}$ mice is altered and characterized by a transient hyperplasia of basal cells.

$\alpha 7$ nAChR Regulates the Proliferation of Airway Epithelial Basal Cells

We have observed that $\alpha 7$ nAChR regulates the proliferation of HAECs in air-liquid interface culture and that the proliferating cells are mainly composed of basal epithelial cells (Figure 7, A and B). Because $\alpha 3\alpha 5\beta 2$ nAChR is also expressed in basal epithelial cells and involved in the wound repair process of the injured airway epithelium,¹⁹ we investigated the respective and potential role of $\alpha 7$ nAChR and $\alpha 3\alpha 5\beta 2$ nAChR in basal cell proliferation by studying the effect of different nAChR antagonists, α BTX^{10,26,27} and methyllycaconitine⁴¹ for the $\alpha 7$ nAChR, α -conotoxin MII for the $\alpha 3\beta 2$ nAChR,¹⁶ and mecamylamine for all $\alpha_x\beta_y$ heteropentameric nAChRs,⁴² on the *in vitro*




Figure 5. nAChR $\alpha 7$ in a smoker-derived remodeled human airway epithelium. The localization of nAChR $\alpha 7$, CK13 and CK14, desmoplakins 1 and 2 (Desmo), and β -tubulin was studied by immunofluorescence with Alexa 488 conjugates (green color in dark panels) in a human bronchial tissue sample derived from a patient with a significant smoking history and presenting different types of remodeling: normal pseudostratified epithelium with hyperplasia of secretory cells (A), hyperplasia of basal cells and squamous metaplasia (B), and squamous metaplasia (C). D reports $\alpha 7$ nAChR distribution in the area of basal cell hyperplasia. $\alpha 7$ nAChR is expressed in the basal cell layer in the pseudostratified epithelium (D, arrow) and present in most basal layers in areas of basal cell hyperplasia or squamous metaplasia (D and C, arrowheads).

proliferation of isolated HAECs (Figure 9A) and of sorted human airway basal epithelial cells (Figure 9B). We did not observe any significant effect of both α -conotoxin MII and mecamylamine on cell proliferation when tested from 0.1 to 10 $\mu\text{mol/L}$. On the contrary, αBTX exposure dose-dependently increased cell proliferation both in isolated HAECs (Figure 9A) and in sorted human airway basal epithelial cells (Figure 9B), with a maximal effect observed for a 10 $\mu\text{mol/L}$ concentration. Both αBTX and

methyllycaconitine, two competitive antagonists of $\alpha 7$ nAChRs, stimulated basal cell proliferation (Figure 9B). Similarly, exposure of 16HBE14o-cells, a normal human airway cell line⁴³ expressing the basal airway cell markers CK13 and CK14 (Figure 10 A), to 0.1–10 $\mu\text{mol/L}$ αBTX significantly increased ($P = 0.0074$, Friedman test) cell proliferation in a dose-dependent way (Figure 10B). Transfection of the 16HBE14o-cell line with two different small interfering RNAs (siRNAs), Si1 and Si2, targeted to




Figure 6. Effect of α BTX on the *ex vivo* wound repair of the human respiratory epithelium. Normal human bronchial tissue samples were locally wounded (*ex vivo* wound repair model) and maintained in culture. **A:** After 2 days in culture, the localization of CK13 and nAChR $\alpha 7$ was studied by immunofluorescence with Alexa 488 conjugates (green color in dark panels) in stationary cells in the unwounded pseudostratified normal epithelium (left column), in the transition area between the normal unwounded epithelium and the migrating cells (middle column) and in migrating cells in the wounded area (right column). Arrowheads denote expression of nAChR $\alpha 7$ in basal cells present in the unwounded pseudostratified epithelium and at the lesion edge. **B:** After 3 and 4 days in culture in the presence (right panels) or absence (control, left panels) of 5 $\mu\text{mol/L}$ α BTX, the localization of CK13 (Alexa 594, red color) and of the Ki-67 protein (Alexa 488, green color) was studied by immunofluorescence in the regenerating epithelium, ie, in cells located in the initially wounded area of the injured epithelium.

mRNA encoding the $\alpha 7$ nAChR subunit, significantly inhibited protein level for $\alpha 7$ nAChR by 56 and 41%, respectively (Figure 10C) and stimulated cell proliferation over a 1- to 6-day period (Figure 10D), as compared with scramble siRNA controls. On the contrary, transfection of the 16HBE14o-cell line with the $\alpha 7$ nAChR cDNA significantly increased by 157% $\alpha 7$ nAChR protein level (Figure 10E) and decreased cell proliferation by 49 and 25%, 3 and 6 days after transfection, respectively, as compared with transfection with the pCMV6 empty vector (Figure 10F).

Discussion

The remodeling of the airway epithelium, described as abnormal plasticity, includes basal cell and goblet cell

hyperplasia, and squamous cell metaplasia.⁴⁴ These changes involve that any delay in the repair of the injured epithelium, or disruption in proliferation and differentiation, may affect the dynamic process of regeneration. In this study, we demonstrate that the $\alpha 7$ nAChR, expressed in human airway basal cells, is essential for the regeneration of a normal pseudostratified epithelium by regulating basal cells proliferation *in vivo* and *in vitro*. Our findings also suggest that alteration of $\alpha 7$ nAChR expression or activity in airway basal cells is involved in the airway epithelium remodeling, namely basal cell hyperplasia and squamous metaplasia.

In animal models, it has been shown that airway basal cells have the potential to proliferate and to differentiate into most cell types of the mucociliary epithelium to re-




Figure 7. Effect of $\alpha 7$ nAChR on the *in vitro* regeneration of the human respiratory epithelium. HARECs were isolated and cultured in air-liquid interface in the absence (control) or in the presence of 5 μ mol/L α BTX added from day 1 (α BTX d1) or from day 11 (α BTX d11). **A:** Cells in each condition were counted over a 30-day period in culture. The results are expressed as mean cell number \pm SD for five different experiments conducted on primary cultures of HARECs derived from five different patients. The significance of the differences between α BTX-exposed cultures and control cultures was determined using Wilcoxon rank test ($^*P < 0.05$). **B:** CK13 (red color: Alexa 594 conjugate), Ki-67 (green color: Alexa 488 conjugate), and cell nuclei (blue color: 4',6'-diamino-2-phenylindole) immunostaining of HARECs after 2 days in control and α BTX-exposed cultures; arrowheads point to Ki-67 immunoreactivity in CK13-positive cells. **C:** Morphology of the regenerated epithelium after 16, 23, and 33 days in culture in the different conditions (control, α BTX d1 and α BTX d11). Arrowheads point to ciliated cells. **D:** CK13, CK14, desmoplakins 1 and 2 (Desm), P63, β -tubulin (β -Tub), α -ENaC, Foxj1, and transglutaminase-1 (TG-1) immunostaining with Alexa 488 conjugates of HARECs after 33 days in culture in air-liquid interface condition.

generate its structure.⁴⁵⁻⁴⁸ More recently, *in vivo* and *in vitro* studies demonstrated that human adult airway basal cells, as opposed to ciliated and secretory cells, are able to proliferate and to reconstitute a fully differentiated and functional mucociliary airway epithelium and may be considered as progenitors of the human airway epithelium.² We have observed that airway epithelial cells, migrating to repair a wound (*ex vivo* wound repair model) or adhering and proliferating to further regenerate the epithelium (air-liquid interface culture model), present basal cell characteristics and continuously express $\alpha 7$ nAChR. Al-

though $\alpha 7$ nAChR has been shown to be expressed at the apex of the rat tracheal epithelium⁴⁹ or of nicotine-treated monkey airway epithelium,¹³ the expression of $\alpha 7$ nAChR in airway basal cells has never been reported before. Although $\alpha 7$ nAChR is expressed in migrating basal cells repairing a wound, we previously observed that $\alpha 7$ nAChR is not involved in the first step of epithelium regeneration, ie, cell spreading and migration, which implies the $\alpha 3\alpha 5\beta 2$ nAChR.¹⁹ Here, we show that $\alpha 7$ nAChR expressed in airway basal cells is involved in the epithelium differentiation.




Figure 8. Regeneration of the tracheal epithelium after exposure to polydocanol in $\alpha 7^{+/+}$ and $\alpha 7^{-/-}$ mice. **A:** Histological hematoxylin-stained sections of upper trachea from day 0 (just before polydocanol treatment) to day 8 in 8-week-old mice. **Arrows** in day 2 denote only flat cells spreading over the denuded basement membrane are observed both in $\alpha 7^{+/+}$ mice and in $\alpha 7^{-/-}$ mice. **Arrows** in day 6 denote the first basal cells observed in $\alpha 7^{+/+}$ mice. **Arrows** in day 7 denote the appearance of the first basal cells in $\alpha 7^{-/-}$ mice. **B:** Variation of epithelium height during the epithelium regeneration. Results are expressed as mean epithelium height \pm SD for four to five mice for each time. The significance of the differences between control $\alpha 7^{+/+}$ mice and $\alpha 7^{-/-}$ mice was determined at each time using the Mann-Whitney test. **C:** Identification of caveolin-1 in cryosections by immunofluorescence with an Alexa 594 conjugate.

The expression of $\alpha 7$ nAChR is definitely associated with the differentiation state of the airway epithelium. The highest detectable $\alpha 7$ nAChR expression, both during the human airway fetal development and during the *in vitro* regeneration of the human pseudostratified epithelium,

is observed when airway epithelial differentiation takes place. Otherwise, $\alpha 7^{-/-}$ mice are viable and the airway epithelium morphology is similar in $\alpha 7^{-/-}$ mice and in control mice, at least during the first weeks of life, suggesting that $\alpha 7$ nAChR expression is not essential for




Figure 9. Effect of nicotinic antagonists on the *in vitro* proliferation of HAECs. The proliferation of HAECs, isolated from normal human bronchial tissues (A) or of basal cells, sorted by FACS (tetraspanin CD151⁺ and tissue factor⁺) from a normal human bronchial tissue sample (B) was studied with the 3-(4,5-dimethylthiazol-2-yl)-2,5-dimethyltetrazolium bromide (MTT) technique on multiwell plates in the presence of increasing concentrations of α BTX, methyllycaconitine, α -connotoxin MII, and mecamylamine (A and B). The results are expressed as mean OD at 560 nm \pm SD for 10 different experiments conducted on primary cultures of HRECs derived from 10 different patients (A) or for one experiment with sorted basal epithelial cells, run in sextuplicates (B). The significance of the differences between control and antagonist exposed-cells was determined using the Wilcoxon rank test (A) and the Mann-Whitney test (B).

normal airway development or that the missing function of $\alpha 7$ nAChR in $\alpha 7^{-/-}$ mice is compensated by up-regulated expression of other nAChR subunits as described in keratinocytes.⁵⁰ Nevertheless, chronic prenatal nicotine exposure increases pulmonary $\alpha 7$ nAChR expression and alters fetal lung development in monkeys.¹³ Chronic exposure to submicromolar concentration of nicotine is also known to irreversibly inactivate many nAChRs, including $\alpha 7$ nAChR,³⁵ and inactivation of $\alpha 7$ nAChR alters pulmonary morphogenesis.¹⁵ Similarly, we have observed that the chronic inactivation of $\alpha 7$ nAChR by α BTX prevents the *in vitro* differentiation of the normal human pseudostratified epithelium and that the absence of $\alpha 7$ nAChR in mice induces modifications of the epithelium phenotype and an altered epithelial regeneration after wounding. We can therefore infer that $\alpha 7$ nAChR plays a potential role in the airway epithelium differentiation. Surprisingly, in all these *in vivo* and *in vitro* approaches, we observed that the absence or the inactivation of $\alpha 7$ nAChR is characterized by either a transient or an established hyperplasia of basal epithelial cells, which led us to investigate the potential role of the $\alpha 7$ nAChR in the regulation of airway epithelial basal cell proliferation.




Figure 10. Effect of α BTX, of siRNA, and of cDNA specific for nAChR $\alpha 7$ on the proliferation of the normal airway cell line 16HBE140-. **A:** CK13 and CK14 immunostaining of 16HBE140-cells in culture. **B:** 16HBE140-cells were cultured for 6 days in multiwell plates in the presence of increasing concentrations of α BTX, and the cell number was determined using the 3-(4,5-dimethylthiazol-2-yl)-2,5-dimethyltetrazolium bromide (MTT) method. Results are expressed as mean OD at 560 nm \pm SD for four different α BTX-treated cultures. **C–F:** 16HBE140-cells were transiently transfected either with two different siRNA sequences against nAChR $\alpha 7$ subunit (Si 1 and Si 2), or with two scrambled control siRNAs (Ctrl 1 and Ctrl 2) (C and D), or with the $\alpha 7$ cDNA expression vector, or the empty control vector (pCMV6) (E and F). **C and E:** Immunoblots (top panels) and quantification of nAChR $\alpha 7$ subunit, normalized for glyceraldehyde-3-phosphate dehydrogenase (GAPDH) values (bottom panels) 6 days after transfection. Data are expressed as fold induction for $\alpha 7$ (Si1, Si2, or cDNA) transfectants relative to the control (Ctrl1, Ctrl2, or pCMV6) transfectants. The significance of the differences between transfected cells and the corresponding controls was determined using the one simple *t*-test (**P* < 0.05; ***P* < 0.01). **D and F:** Transfected cells were counted on days 1, 3, and 6 after transfection. The results are expressed as mean cell number \pm SD for four different transfected cultures. The significance of the differences between $\alpha 7$ siRNA or $\alpha 7$ cDNA-transfected cells and their corresponding controls was determined using the Mann-Whitney test (**P* = 0.0339; ***P* = 0.0209).

$\alpha 7$ nAChR appears to be a key negative regulator of airway epithelial basal cell proliferation: 1) inactivating $\alpha 7$ nAChR *in vitro* with α BTX stimulates cell proliferation in the early phases of the epithelial regeneration and the corresponding proliferating cells have phenotypic characteristics of basal epithelial cells, 2) regeneration of the airway epithelium in mice lacking $\alpha 7$ nAChR is characterized by a transient hyperplasia of basal cells, 3) airway epithelium in aged $\alpha 7^{-/-}$ mice more frequently presents areas of basal cell hyperplasia, and 4) *in vitro*, the proliferating cells have phenotypic characteristics of basal epithelial cells.

eration of sorted human airway basal cells is stimulated by α BTX and methyllycaconitine but not by α -Conotoxin MII nor mecamylamine, suggesting that only $\alpha 7$ nAChR regulates the proliferation of basal cells in the normal human airway epithelium. Similarly, inhibition of the $\alpha 7$ nAChR pathway favored cell cycle progression in epidermal keratinocytes.⁵⁰ On the contrary, many authors have noticed that $\alpha 7$ is the main nAChR subunit that mediates the proliferative effects of nicotine in cancer cell lines.¹² This $\alpha 7$ nAChR-mediated stimulatory effect of nicotine was also observed in commercialized normal bronchial epithelial cells, after these cells have been amplified after isolation.⁵¹ At the opposite, we always observed a stimulatory proliferative action of α BTX on primary cultures of freshly isolated HAECs and on purified human airway basal cells. We also observed the same proliferation stimulatory effect of both α BTX and $\alpha 7$ nAChR-specific siRNAs on the 16HBE14o-cell line, a normal human airway cell line expressing the basal airway cell markers CK13 and CK14, whereas transfection with the $\alpha 7$ nAChR cDNA inhibited cell proliferation. We observed a stimulatory proliferative effect of nicotine both on HAECs and isolated human airway basal cells only when cells were incubated for 7 to 15 days with high nicotine concentrations (10 to 100 μ mol/L), conditions which may have favored $\alpha 7$ nAChR inactivation (data not shown). This antagonist-like effect of nicotine, on over exposure of HAECs to nicotine, has been already described by Zia et al.⁷ These results suggest that the regulation of cell proliferation by $\alpha 7$ nAChR may be quite different in the normal airway epithelium and in primary cultures of normal HAECs, on the one hand, and in lung cancer and cell lines derived from lung cancers, on the other hand.

Our results raise the concept of coupling between cell growth and differentiation. In the normal pseudostratified tracheobronchial epithelium, a strict balance exists between cellular proliferation and differentiation.^{44,52} This balance is likely maintained via equilibrium between positive and negative growth regulatory factors and factors controlling differentiation.⁵³ It is generally accepted that an inverse relationship exists between growth and differentiation in the airway epithelium.^{52,54} For example, after mechanical or toxic injury, the balance between proliferation and differentiation is disturbed, and the epithelium undergoes hyperplasia followed by transient squamous metaplasia,⁵⁵⁻⁶⁰ a situation similar to our observation after α BTX exposure: that is, after an increased cell proliferation, the regenerated airway epithelium ends up with squamous metaplasia. Interestingly, we have observed that, although the airway epithelium undergoes squamous metaplasia *in vitro*, Foxj1, a transcription factor involved in ciliated cells differentiation,^{61,62} is down-regulated. Regulation of Foxj1 expression may thus be a key element in the α BTX-induced epithelium squamous metaplasia. We have observed that α BTX-induced squamous metaplasia takes place only when adding the $\alpha 7$ nAChR antagonist during the proliferation step of the airway epithelium regeneration. It has been proposed that the induction of squamous metaplasia after injury of the tracheobronchial epithelium involves enhanced proliferation of a p63⁺/CK14⁺/CK5⁺/BS-I-B4⁺ subpopulation of basal cells.³⁷

Indeed, we have observed that $\alpha 7$ nAChR blockade induces expression of squamous metaplasia phenotype in cultured HAECs, which was preceded by CK14⁺ basal cell hyperplasia. Interestingly, constitutive expression of the human CK14 gene in the mouse airway epithelium induced tracheal basal cell hyperplasia and squamous metaplasia.⁶³ Of note, the lack of p63 expression in α BTX-induced squamous metaplasia is in agreement with the hypothesis that down-regulation of p63 might be necessary for basal cells to differentiate into squamous cells.³⁷ Among the many factors that have been identified and regulate positively or negatively the growth and differentiation of airway epithelial cells by autocrine or paracrine mechanisms, transforming growth factor β (TGF- β), a major cytokine involved in driving squamous metaplasia, is of particular interest. Jetten et al.³⁸ and Masui et al.⁶⁴ have indeed shown that TGF- β exposure or overexpression in airway epithelial cells induces terminal cell division and favors the expression of a squamous phenotype. Similarly in the present study, α BTX-induced hyperproliferation is followed by a period in which cells stop proliferating and undergo squamous differentiation, suggesting that these changes in the differentiation pathway, when $\alpha 7$ nAChR is absent or inactivated, may be related to TGF- β overexpression. In addition, we have observed that the $\alpha 7$ nAChR blockade enhances the expression of transglutaminase, a marker of squamous differentiation known to be up-regulated by TGF- β .³⁸ Moreover, up-regulation of TGF- β by nicotine is mediated by $\alpha 7$ nAChR during atrial remodeling,⁶⁵ TGF- β up-regulates CK14 expression and promotes the expression of basal cell phenotype⁶⁶ and inhibits p63 function.⁶⁷ Taken together, our results suggest that $\alpha 7$ nAChRs inactivation can initiate an alternative program of the basal cell differentiation process leading to squamous differentiation and that this process may be associated to down-regulation of Foxj1 and/or up-regulation of TGF- β .

Our data suggest that $\alpha 7$ nAChR is critical for the renewal of the normal respiratory epithelium by repressing basal cell proliferation and stimulating their differentiation. After an acute injury, the regeneration of the airway epithelium in $\alpha 7^{-/-}$ mice is delayed, transiently abnormal, and characterized by a basal cell hyperplasia. On the long term and after many epithelium renewals, absence or inactivation of the $\alpha 7$ nAChR may progressively induce the frequent remodeling of the airway epithelium observed in old $\alpha 7^{-/-}$ mice: hyperplasia of basal cells and squamous metaplasia. $\alpha 7$ nAChR inactivation, during the *in vitro* regeneration of the human airway epithelium, also induces dramatic morphological changes leading to squamous metaplasia. Many inflammatory bronchopulmonary diseases are characterized by a bronchial epithelium remodeling⁶⁸ and by a progressive and altered ability of the airway epithelium to regenerate a normal and functional epithelium.⁶⁹ Chronic obstructive pulmonary diseases are also characterized by a permanent inflammatory context. The $\alpha 7$ nAChR is also now known as an essential regulator of inflammation.^{21,70} Its inactivation or down-expression may contribute to the excessive proinflammatory pathway, the increased fragility of the airway epithelium and the altered epithelium

regeneration observed in COPD. Most patients with COPD have a history of chronic smoking. In these patients, $\alpha 7$ nAChR inactivation may result from chronic exposure to nicotine. Indeed, the $\alpha 7$ nAChR has been shown to be particularly susceptible to inactivation^{33–35} and over exposure of bronchial epithelium cells to nicotine *in vitro* produced an antagonist-like effect (7). Interestingly, $\alpha 7^{-/-}$ mice and α BTX-exposed HAEC cultures present the same airway epithelium remodelings, basal cell hyperplasia, and squamous metaplasia, as those observed in these smoking COPD patients.

In summary, $\alpha 7$ nAChR appears to be a key regulator of the plasticity of the human airway epithelium in controlling basal cell proliferation. Our results also suggest that $\alpha 7$ nAChR may be involved in the basal cell differentiation process. These effects may be determinant in the airway epithelium remodeling associated to cigarette smoke as well as in the development of COPD and lung cancers.

Acknowledgments

We thank Edith Puchelle and Marina Pretolani (INSERM U700, Paris, France) for helpful discussions and Anne L. Leblond (INSERM U533, Nantes, France) for her help in developing the mouse model of tracheal desquamation. We thank all of the surgeons and doctors who provided us with human airway tissues: Dr. Pascal Corlieu (Hôpital Tenon, Paris, France), Prof. Jean-Michel Klossek (Hôpital Jean Bernard, Poitiers, France), Dr. Christophe Ruau (Clinique Mutualiste de la Sagesse, Rennes, France), Drs. Michel Monteau and Talal Nasser (Clinique Courlancy, Reims, France), and Drs. Elisabeth Alanio, Matthieu Capovilla, Anne Durlach, Dominique Gaillard, Karine Joseph, and Dominique Zachar (Hôpital Maison Blanche, Reims, France).

References

1. Coraux C, Hajj R, Lesimple P, Puchelle E: [Repair and regeneration of the airway epithelium]. *French. Med Sci* 2005, 21:1063–1069
2. Hajj R, Baranek T, Le NR, Lesimple P, Puchelle E, Coraux C: Basal cells of the human adult airway surface epithelium retain transit-amplifying cell properties. *Stem Cells* 2007, 25:139–148
3. Grando SA, Kawashima K, Kirkpatrick CJ, Wessler I: Recent progress in understanding the non-neuronal cholinergic system in humans. *Life Sci* 2007, 80:2181–2185
4. Wessler I, Kirkpatrick CJ, Racke K: Non-neuronal acetylcholine, a locally acting molecule, widely distributed in biological systems: expression and function in humans. *Pharmacol Ther* 1998, 77:59–79
5. Kummer W, Lips KS, Pfeil U: The epithelial cholinergic system of the airways. *Histochem Cell Biol* 2008, 130:219–234
6. Steinbach JH: Mechanism of action of the nicotinic acetylcholine receptor. *Ciba Found Symp* 1990, 152:53–61
7. Zia S, Ndoye A, Nguyen VT, and Grando SA: Nicotine enhances expression of the $\alpha 3$, $\alpha 4$, $\alpha 5$, and $\alpha 7$ nicotinic receptors modulating calcium metabolism and regulating adhesion and motility of respiratory epithelial cells. *Res Commun Mol Pathol Pharmacol* 1997, 97:243–262
8. Maus AD, Pereira EF, Karachunski PI, Horton RM, Navaneetham D, Macklin K, Cortes WS, Albuquerque EX, Conti-Fine BM: Human and rodent bronchial epithelial cells express functional nicotinic acetylcholine receptors. *Mol Pharmacol* 1998, 54:779–788
9. Carlisle DL, Hopkins TM, Gaitner-Davis A, Silhanek MJ, Luketich JD, Christie NA, Siegfried JM: Nicotine signals through muscle-type and

neuronal nicotinic acetylcholine receptors in both human bronchial epithelial cells and airway fibroblasts. *Respir Res* 2004, 5:27

10. Seguela P, Wadiche J, Ney Miller K, Dani JA, Patrick JW: Molecular cloning, functional properties, and distribution of rat brain $\alpha 7$: a nicotinic cation channel highly permeable to calcium. *J Neurosci* 1993, 13:596–604
11. Catassi A, Servent D, Paleari L, Cesario A, Russo P: Multiple roles of nicotine on cell proliferation and inhibition of apoptosis: implications on lung carcinogenesis. *Mutat Res* 2008, 659:221–231
12. Egleton RD, Brown KC, Dasgupta P: Nicotinic acetylcholine receptors in cancer: multiple roles in proliferation and inhibition of apoptosis. *Trends Pharmacol Sci* 2008, 29:151–158
13. Sekhon HS, Jia Y, Raab R, Kuryatov A, Pankow JF, Whitsett JA, Lindstrom J, Spindel ER: Prenatal nicotine increases pulmonary $\alpha 7$ nicotinic receptor expression and alters fetal lung development in monkeys. *J Clin Invest* 1999, 103:637–647
14. Sekhon HS, Keller JA, Benowitz NL, Spindel ER: Prenatal nicotine exposure alters pulmonary function in newborn rhesus monkeys. *Am J Respir Crit Care Med* 2001, 164:989–994
15. Wongtrakool C, Roser-Page S, Rivera HN, Roman J: Nicotine alters lung branching morphogenesis through the $\alpha 7$ nicotinic acetylcholine receptor. *Am J Physiol* 2007, 293:L611–L618
16. Cartier GE, Yoshikami D, Gray WR, Luo S, Olivera BM, McIntosh JM: A new α -conotoxin which targets $\alpha 3\beta 2$ nicotinic acetylcholine receptors. *J Biol Chem* 1996, 271:7522–7528
17. Buisson AC, Gilles C, Polette M, Zahm JM, Birembaut P, Tournier JM: Wound repair-induced expression of a stromelysins is associated with the acquisition of a mesenchymal phenotype in human respiratory epithelial cells. *Lab Invest* 1996, 74:658–669
18. Legrand C, Gilles C, Zahm JM, Polette M, Buisson AC, Kaplan H, Birembaut P, Tournier JM: Airway epithelial cell migration dynamics: MMP-9 role in cell-extracellular matrix remodeling. *J Cell Biol* 1999, 146:517–529
19. Tournier JM, Maouche K, Coraux C, Zahm JM, Cloez-Tayarani I, Nawrocki-Raby B, Bonnomet A, Bulet H, Lebargy F, Polette M, Birembaut P: $\alpha 3\alpha 5\beta 2$ -Nicotinic acetylcholine receptor contributes to the wound repair of the respiratory epithelium by modulating intracellular calcium in migrating cells. *Am J Pathol* 2006, 168:55–68
20. Hajj R, Lesimple P, Nawrocki-Raby B, Birembaut P, Puchelle E, Coraux C: Human airway surface epithelial regeneration is delayed and abnormal in cystic fibrosis. *J Pathol* 2007, 211:340–350
21. Wang H, Yu M, Ochani M, Amella CA, Tanovic M, Susarla S, Li JH, Wang H, Yang H, Ulloa L, Al-Abed Y, Czura CJ, Tracey KJ: Nicotinic acetylcholine receptor $\alpha 7$ subunit is an essential regulator of inflammation. *Nature* 2003, 421:384–388
22. Lechner JF, LaVeck MA: A serum-free method for culturing normal human bronchial epithelial cells at clonal density. *J Tissue Cult Meth* 1985, 9:43–48
23. Bonnomet A, Polette M, Strumane K, Gilles C, Dalstein V, Kileztky C, Berx G, van RF, Birembaut P, Nawrocki-Raby B: The E-cadherin-repressed hNanos1 gene induces tumor cell invasion by up-regulating MT1-MMP expression. *Oncogene* 2008, 27:3692–3699
24. Orr-Urtreger A, Goldner FM, Saeki M, Lorenzo I, Goldberg L, De BM, Dani JA, Patrick JW, Beaudet AL: Mice deficient in the $\alpha 7$ neuronal nicotinic acetylcholine receptor lack α -bungarotoxin binding sites and hippocampal fast nicotinic currents. *J Neurosci* 1997, 17:9165–9171
25. Borthwick DW, Shahbazian M, Krantz QT, Dorin JR, Randell SH: Evidence for stem-cell niches in the tracheal epithelium. *Am J Respir Cell Mol Biol* 2001, 24:662–670
26. Rangwala F, Drisdel RC, Rakhilin S, Ko E, Atluri P, Harkins AB, Fox AP, Salman SS, Green WN: Neuronal α -bungarotoxin receptors differ structurally from other nicotinic acetylcholine receptors. *J Neurosci* 1997, 17:8201–8212
27. Drisdel RC, Green WN: Neuronal α -bungarotoxin receptors are $\alpha 7$ subunit homomers. *J Neurosci* 2000, 20:133–139
28. Jeffery PK, Gaillard D, Moret S: Human airway secretory cells during development and in mature airway epithelium. *Eur Respir J* 1992, 5:93–104
29. Dupuit F, Kalin N, Brezillon S, Hinnrasky J, Tummler B, Puchelle E: CFTR and differentiation markers expression in non-CF and $\delta F 508$ homozygous CF nasal epithelium. *J Clin Invest* 1995, 96:1601–1611
30. Flint FF, Schulte BA, Spicer SS: Glycoconjugate with terminal α -galactose: a property common to basal cells and a subpopulation

- of columnar cells of numerous epithelia in mouse and rat. *Histochemistry* 1986, 84:387–395
31. Changeux JP, villiers-Thierry A, Chemouilli P: Acetylcholine receptor: an allosteric protein. *Science* 1984, 225:1335–1345
 32. Picciotto MR, Addy NA, Mineur YS, Brunzell DH: It is not "either/or": activation and desensitization of nicotinic acetylcholine receptors both contribute to behaviors related to nicotine addiction and mood. *Prog Neurobiol* 2008, 84:329–342
 33. Couturier S, Bertrand D, Matter JM, Hernandez MC, Bertrand S, Millar N, Valera S, Barkas T, Ballivet M: A neuronal nicotinic acetylcholine receptor subunit ($\alpha 7$) is developmentally regulated and forms a homo-oligomeric channel blocked by α -BTX. *Neuron* 1990, 5:847–856
 34. Giniatullin R, Nistri A, Yakel JL: Desensitization of nicotinic ACh receptors: shaping cholinergic signaling. *Trends Neurosci* 2005, 28:371–378
 35. Olale F, Gerzanich V, Kuryatov A, Wang F, Lindstrom J: Chronic nicotine exposure differentially affects the function of human $\alpha 3$, $\alpha 4$, and $\alpha 7$ neuronal nicotinic receptor subtypes. *J Pharmacol Exp Ther* 1997, 283:675–683
 36. Arredondo J, Chernyavsky AI, Jolkovsky DL, Pinkerton KE, Grando SA: Receptor-mediated tobacco toxicity: acceleration of sequential expression of $\alpha 5$ and $\alpha 7$ nicotinic receptor subunits in oral keratinocytes exposed to cigarette smoke. *FASEB J* 2008, 22:1356–1368
 37. Daniely Y, Liao G, Dixon D, Linnoila RI, Lori A, Randell SH, Oren M, Jetten AM: Critical role of p63 in the development of a normal esophageal and tracheobronchial epithelium. *Am J Physiol* 2004, 287:C171–C181
 38. Jetten AM, Shirley JE, Stoner G: Regulation of proliferation and differentiation of respiratory tract epithelial cells by TGF- β . *Exp Cell Res* 1986, 167:539–549
 39. Pan J, You Y, Huang T, Brody SL: RhoA-mediated apical actin enrichment is required for ciliogenesis and promoted by Foxj1. *J Cell Sci* 2007, 120:1868–1876
 40. Krasteva G, Pfeil U, Drab M, Kummer W, Konig P: Caveolin-1 and -2 in airway epithelium: expression and in situ association as detected by FRET-CLSM. *Respir Res* 2006, 7:108
 41. Davies AR, Hardick DJ, Blagbrough IS, Potter BV, Wolstenholme AJ, Wonnacott S: Characterisation of the binding of [3 H]methyllycaconitine: a new radioligand for labelling $\alpha 7$ -type neuronal nicotinic acetylcholine receptors. *Neuropharmacology* 1999, 38:679–690
 42. Chavez-Noriega LE, Crona JH, Washburn MS, Urrutia A, Elliott KJ, Johnson EC: Pharmacological characterization of recombinant human neuronal nicotinic acetylcholine receptors h $\alpha 2\beta 2$, h $\alpha 2\beta 4$, h $\alpha 3\beta 2$, h $\alpha 3\beta 4$, h $\alpha 4\beta 2$, h $\alpha 4\beta 4$ and h $\alpha 7$ expressed in *Xenopus oocytes*. *J Pharmacol Exp Ther* 1997, 280:346–356
 43. Cozens AL, Yezzi MJ, Kunzelmann K, Ohri T, Chin L, Eng K, Finkbeiner WE, Widdicombe JH, Gruenert DC: CFTR expression and chloride secretion in polarized immortal human bronchial epithelial cells. *Am J Respir Cell Mol Biol* 1994, 10:38–47
 44. Basbaum C, Jany B: Plasticity in the airway epithelium. *Am J Physiol* 1990, 259:L38–L46
 45. Inayama Y, Hook GE, Brody AR, Cameron GS, Jetten AM, Gilmore LB, Gray T, Nettesheim P: The differentiation potential of tracheal basal cells. *Lab Invest* 1988, 58:706–717
 46. Liu JY, Nettesheim P, Randell SH: Growth and differentiation of tracheal epithelial progenitor cells. *Am J Physiol* 1994, 266:L296–L307
 47. Hong KU, Reynolds SD, Watkins S, Fuchs E, Stripp BR: Basal cells are a multipotent progenitor capable of renewing the bronchial epithelium. *Am J Pathol* 2004, 164:577–588
 48. Ford JR, Terzaghi-Howe M: Basal cells are the progenitors of primary tracheal epithelial cell cultures. *Exp Cell Res* 1992, 198:69–77
 49. Wang Y, Pereira EF, Maus AD, Ostlie NS, Navaneetham D, Lei S, Albuquerque EX, Conti-Fine BM: Human bronchial epithelial and endothelial cells express $\alpha 7$ nicotinic acetylcholine receptors. *Mol Pharmacol* 2001, 60:1201–1209
 50. Arredondo J, Nguyen VT, Chernyavsky AI, Bercovich D, Orr-Urtreger A, Kummer W, Lips K, Vetter DE, Grando SA: Central role of $\alpha 7$ nicotinic receptor in differentiation of the stratified squamous epithelium. *J Cell Biol* 2002, 159:325–336
 51. Dasgupta P, Chellappan SP: Nicotine-mediated cell proliferation and angiogenesis: new twists to an old story. *Cell Cycle* 2006, 5:2324–2328
 52. Ayers MM, Jeffery PK: Proliferation and differentiation in mammalian airway epithelium. *Eur Respir J* 1988, 1:58–80
 53. Jetten AM: Growth and differentiation factors in tracheobronchial epithelium. *Am J Physiol* 1991, 260:L361–L373
 54. Jetten AM, Rearick JI, Smits HL: Regulation of differentiation of airway epithelial cells by retinoids. *Biochem Soc Trans* 1986, 14:930–933
 55. Harris CC, Silverman T, Smith JM, Jackson F, Boren HG: Proliferation of tracheal epithelial cells in normal and vitamin A-deficient Syrian golden hamsters. *J Natl Cancer Inst* 1973, 51:1059–1062
 56. Mossman BT, Ley BW, Craighead JE: Squamous metaplasia of the tracheal epithelium in organ culture. I. Effects of hydrocortisone and β -retinyl acetate. *Exp Mol Pathol* 1976, 24:405–414
 57. Lane BP, Gordon RE: Regeneration of vitamin A deficient rat tracheal epithelium after mild mechanical injury: growth kinetics and cellular differentiation. *Differentiation* 1979, 14:87–93
 58. Keenan KP, Combs JW, McDowell EM: Regeneration of hamster tracheal epithelium after mechanical injury. III. Large and small lesions: comparative stathmokinetic and single pulse and continuous thymidine labeling autoradiographic studies. *Virchows Arch B Cell Pathol Incl Mol Pathol* 1982, 41:231–252
 59. Keenan KP, Wilson TS, McDowell EM: Regeneration of hamster tracheal epithelium after mechanical injury. IV. Histochemical, immunocytochemical and ultrastructural studies. *Virchows Arch B Cell Pathol Incl Mol Pathol* 1983, 43:213–240
 60. Woodworth CD, Mossman BT, Craighead JE: Squamous metaplasia of the respiratory tract: possible pathogenic role in asbestos-associated bronchogenic carcinoma. *Lab Invest* 1983, 48:578–584
 61. Chen J, Knowles HJ, Hebert JL, Hackett BP: Mutation of the mouse hepatocyte nuclear factor/forkhead homologue 4 gene results in an absence of cilia and random left-right asymmetry. *J Clin Invest* 1998, 102:1077–1082
 62. Brody SL, Yan XH, Wuerffel MK, Song SK, Shapiro SD: Ciliogenesis and left-right axis defects in forkhead factor HFH-4-null mice. *Am J Respir Cell Mol Biol* 2000, 23:45–51
 63. Dakir EL, Feigenbaum L, Linnoila RI: Constitutive expression of human keratin 14 gene in mouse lung induces premalignant lesions and squamous differentiation. *Carcinogenesis* 2008, 29:2377–2384
 64. Masui T, Wakefield LM, Lechner JF, LaVeck MA, Sporn MB, Harris CC: Type β transforming growth factor is the primary differentiation-inducing serum factor for normal human bronchial epithelial cells. *Proc Natl Acad Sci USA* 1986, 83:2438–2442
 65. Shan H, Zhang Y, Lu Y, Zhang Y, Pan Z, Cai B, Wang N, Li X, Feng T, Hong Y, Yang B: Down-regulation of miR-133 and miR-590 contributes to nicotine-induced atrial remodelling in canines. *Cardiovasc Res* 2009, 83:465–472
 66. Maki M, Saitoh K, Kaneko Y, Fukayama M, Morohoshi T: Expression of cytokeratin 1, 5, 14, 19 and transforming growth factors $\beta 1$, $\beta 2$, [bet]a3 in osteofibrous dysplasia and adamantinoma: a possible association of transforming growth factor-beta with basal cell phenotype promotion. *Pathol Int* 2000, 50:801–807
 67. Adorno M, Cordenonsi M, Montagner M, Dupont S, Wong C, Hann B, Solari A, Bobisse S, Rondina MB, Guzzardo V, Parenti AR, Rosato A, Biciato S, Balmain A, Piccolo S: A Mutant-p53/Smad complex opposes p63 to empower TGF β -induced metastasis. *Cell* 2009, 137:87–98
 68. Jeffery PK: Remodeling in asthma and chronic obstructive lung disease. *Am J Respir Crit Care Med* 2001, 164:S28–S38
 69. Puchelle E, Zahn JM, Tournier JM, Coraux C: Airway epithelial repair, regeneration, and remodeling after injury in chronic obstructive pulmonary disease. *Proc Am Thorac Soc* 2006, 3:726–733
 70. Fujii YX, Fujigaya H, Moriwaki Y, Misawa H, Kasahara T, Grando SA, Kawashima K: Enhanced serum antigen-specific IgG1 and proinflammatory cytokine production in nicotinic acetylcholine receptor $\alpha 7$ subunit gene knockout mice. *J Neuroimmunol* 2007, 189:69–74