

# ***Ab initio* fragment molecular orbital studies of influenza virus hemagglutinin–sialosaccharide complexes toward chemical clarification about the virus host range determination**

Toshihiko Sawada · Tomohiro Hashimoto ·  
Hiroaki Tokiwa · Tohru Suzuki · Hirofumi Nakano ·  
Hideharu Ishida · Makoto Kiso · Yasuo Suzuki

Received: 14 March 2008 / Revised: 28 April 2008 / Accepted: 29 April 2008 / Published online: 24 June 2008  
© The Author(s) 2008

**Abstract** If we predict the host range of new or mutant influenza virus in advance, we are able to measure against pandemic human influenza immediately after the new virus emerges somewhere. Influenza viral hemagglutinin(HA)–sialoside receptor interaction is a target event for *in silico* chemical prediction studies about the virus host range determination. We theoretically studied avian and human influenza A virus HA H3 subtype complexed with avian or human type receptor Neu5Ac $\alpha$ (2-3 or 2-6)Gal analogues

by *ab initio* fragment molecular orbital (FMO) method at the second order Møller–Plesset (MP2)/6–31G level, which can evaluate correctly not only electrostatic interactions but also lipophilic interactions based on van der Waals dispersion force. Avian H3 bound to avian  $\alpha$ 2-3 11.4 kcal/mol stronger than to human  $\alpha$ 2-6 in the model complexes with taking account of intermolecular lipophilic interaction. A substitution at the position 226 between Gln (avian) and Leu(human) on influenza H3 HA1 has altered

---

T. Sawada · H. Tokiwa · M. Kiso · Y. Suzuki  
Core Research for Evolutional Science and Technology (CREST),  
Japan Science and Technology Agency (JST),  
Kawaguchi, Saitama, 332-0012, Japan

T. Sawada  
e-mail: sawada-t@aist.go.jp

T. Sawada · Y. Suzuki (✉)  
College of Life and Health Sciences, Chubu University,  
1200 Matsumoto-cho,  
Kasugai, Aichi, 487-8501, Japan  
e-mail: suzukiy@isc.chubu.ac.jp

T. Hashimoto (✉)  
Faculty of Regional Studies, Gifu University,  
1-1 Yanagido,  
Gifu, 501-1193, Japan  
e-mail: thashi@gifu-u.ac.jp

H. Tokiwa  
Department of Chemistry, Faculty of Science, Rikkyo University,  
3-34-1 Nishi-Ikebukuro, Tokyo, 171-8501, Japan

T. Suzuki  
Life Science Research Center,  
Gifu University,  
Gifu, Japan

H. Nakano  
Department of Chemistry,  
Aichi University of Education,  
Igaya, Kariya, Aichi, 448–8542, Japan

H. Ishida · M. Kiso  
Department of Applied Bioorganic Chemistry,  
Gifu University,  
Gifu, Japan

Y. Suzuki  
Japan and Global COE Program for Innovation  
in Human Health Sciences,  
University of Shizuoka School of Pharmaceutical Sciences,  
52-1 Yada,  
Shizuoka 422-8526, Japan

*Present Address:*

T. Sawada  
Research Institute for Computational Science (RICS),  
National Institute of Advanced Industrial Science  
and Technology (AIST),  
AIST Tsukuba Central 2, 1-1-1 Umezono,  
Tsukuba, Ibaraki, 305–8568, Japan

its virus host range between avian and human. In the *ab initio* FMO studies, binding energy of avian Gln226Leu H3–human  $\alpha$ 2-6 was quite similar to that in the human H3–human  $\alpha$ 2-6 complex with amino acid sequence differences at nine positions in the models. This similarity indicates that avian Gln226Leu H3 virus can infect human with the same level as human H3 virus. Opposite mutation Leu226Gln in the human H3 gave the moderate binding energies to avian  $\alpha$ 2-3 with similarity to avian H3– $\alpha$ 2-3 complex that supported our previous virus-sialoside binding assay. *Ab initio* FMO studies have revealed the relationship between influenza H3 virus host range and H3– $\alpha$ (2-3 or 2-6) receptors binding. Our theoretical approach may predict the infectious level of new viruses and point out some unknown dangerous mutation positions on HA in advance.

**Keywords** Influenza virus · Hemagglutinin · Sialosaccharide · Virus host range · *Ab initio* FMO

## Introduction

Avian H5N1 virus has a high potential for pandemic influenza. The original H5N1 avian viruses can bind to avian type receptors on human lower respiratory tract [1, 2], however, this infection mechanism does not cause pandemic human influenza. We should always take precaution against when single or double point mutations occur in H5N1 viral hemagglutinin and its higher binding affinity to human type receptor [3–6].

If we predict the host range of new or mutant influenza virus in advance, we are able to measure against pandemic human influenza immediately after the new virus emerges somewhere. Influenza virus host range is mainly determined by HA binding affinity to host cell surface receptors sialo-glycolipids and -glycoproteins terminating in sialic acid  $\alpha$ 2-3 or  $\alpha$ 2-6 galactose [7–10], therefore the HA–sialoside interaction is a target event for *in silico* chemical prediction studies about the virus host range determination. Quantum chemical calculations will provide various chemical foresights about HA–sialosaccharide interaction without treatments of unknown dangerous influenza virus mutants. We are attempting to construct HA–sialoside complex models in order to explain the HA binding specificity using H3 subtype system [11–12]. Our current interest is focused on avian virus HA–Neu5Ac $\alpha$ (2-3/6)Gal bindings and comparison between avian HA mutant–human  $\alpha$ 2-6 interaction and the original human HA–human  $\alpha$ 2-6 interaction.

Avian influenza A virus H3 subtype has Gln226 as one of the amino acids on sialoside binding site HA1 that binds to avian type receptor *N*-acetylneuraminic acid (Neu5Ac)  $\alpha$ 2-3 galactose (Gal) oligosaccharide stronger than to human type receptor Neu5Ac $\alpha$ (2-6)Gal [13]. We remark

that avian H3 weakly and certainly recognize human receptor  $\alpha$ 2-6 [14] on TLC-virus binding assay using synthetic sialylparaglobosides or corresponding B30 derivatives [14–18] that is supported by X-ray crystallographic structure of avian H3–human  $\alpha$ 2-6 complex [19]. Thus, it is significant to analyze the relationship between avian H3 binding affinity and the corresponding avian H3–Neu5Ac $\alpha$ (2-3 or 2-6)Gal complexes. Human H3 virus has Leu226 instead of Gln that strongly recognizes human receptor  $\alpha$ 2-6 sialosaccharide [13, 20–24]. In the human H3 subtype, point mutations on the sialoside binding site at Tyr98Phe, His183Phe, and Leu194Ala decrease their hemagglutination to human erythrocytes [25]. Human Ser193Ile H3 agglutinates  $\alpha$ 2-6 sialoside expressed erythrocytes stronger than the original human H3 [26]. In particular, a substitution from Leu226 to Gln at the position 226 in human H3 HA1 changes its binding specificity from human  $\alpha$ 2-6 to avian  $\alpha$ 2-3 [16, 27–29]. Weis *et al.* have reported X-ray crystal structure of human Leu226Gln H3 complexed with avian type receptor  $\alpha$ 2-3 sialyllactose [20].

For the purpose of theoretical clarification about the H3–sialoside binding properties, we plausibly investigate the binding affinities of H3–Neu5Ac $\alpha$ (2-3 or 2-6)Gal complexes by *ab initio* molecular orbital studies. It is worth while researching the energy profile on all steps in the HA–sialoside interaction composed of several events: encounter the sialoside binding site with sialoside receptor, leaving needless water solvent in the interaction process, thermodynamical relaxation on HA–sialoside complex, and multivalent effect on HA–sialoside interaction. However, we just urgently request a simple and essential approach with a reasonable cost. *Ab initio* fragment molecular orbital (FMO) calculations [30–39] to evaluate the binding energies between HA and Neu5Ac $\alpha$ (2-3/6)Gal receptors are one of the promising strategies.

*Ab initio* FMO method has been applied in large biochemical systems for quantum chemical analysis to molecular interaction [40–45]. In this method, HA–sialoside complex is divided into fragments, and MO calculations are carried out on each fragment and fragment pairs. The total energy on the HA–sialoside complex is obtained as a summation of the fragment energies and interfragment interaction energies (IFIEs) [30–32, 34, 46, 47].

Our previous *ab initio* FMO studies estimated the binding energies between influenza H3 and Neu5Ac $\alpha$ (2-3/6)Gal receptors at the FMO-Hartree-Fock(HF)/STO–3G level, and analyzed IFIEs between Neu5Ac-Gal receptor and amino acid residues on the sialoside binding site to confirm qualitatively avian H3–avian  $\alpha$ 2-3 binding affinity based on electrostatic interactions [11]. This approach would be valid qualitatively, but the FMO-HF/STO–3G method cannot evaluate significantly weak interactions such as intermolecular lipophilic stabilization which play

important roles in biochemical systems. Taking account of electron correlation effects is necessary for more quantitative treatment of these stabilizations based on van der Waals dispersion force. Recently, the FMO method has been extended to correlated calculations, which can be performed at the MP2 method [36, 48]. Including electron correlation effect, we applied the FMO-MP2 method to HA–sialoside model complexes and analyzed the hydrophilic and lipophilic interactions between Neu5Ac–Gal and amino acid residues on the sialoside binding site HA1 using IFIEs.

## Methods


Energy minimized structures of avian H3 complexed with Neu5Ac $\alpha$ (2-3 or 2-6)Gal disaccharide analogues were prepared from the corresponding X-ray crystallographic structures [19] by molecular mechanics (MM) energy calculation with the CFF force field [49] using Discovery Studio 1.5.1 program as shown in our previous works [11, 12]. We mutated *in silico* avian H3 Gln226 to Leu in the avian H3–human  $\alpha$ 2-6 disaccharide complex, changed  $\alpha$ 2-6 bond dihedral angle to human Leu226 H3 type orientation referring the crystal structure of Neu5Ac $\alpha$ (2-6)Gal $\beta$ (1-4)GlcNAc $\beta$ (1-3)Gal $\beta$ (1-4)Glc: LSTc complexed with human H3 [50], and optimized its geometry by MM calculation to make the energy minimum avian Gln226Leu H3–human  $\alpha$ 2-6 complex. Optimum human H3–human  $\alpha$ 2-6 complex was given *in silico* by modification of crystallographic structure of human H3–Neu5Ac $\alpha$ (2-3)Gal $\beta$ (1-4)Glc complex [51]. We replaced the  $\alpha$ 2-3 trisaccharide receptor with Neu5Ac $\alpha$ (2-6)Gal disaccharide based on the common Neu5Ac residue and changed its  $\alpha$ 2-6 bond dihedral angle referring LSTc conformation complexed with human H3 [50] that was similar manner to the previous study [12]. Human Leu226Gln H3–avian  $\alpha$ 2-3 complex was prepared by *in silico* point mutation from Leu226 to Gln in the human H3–avian  $\alpha$ 2-3 complex [51], changed  $\alpha$ 2-3 bond dihedral angle to avian Gln226 H3 type orientation [19] referring the crystal structure of human Leu226Gln–avian  $\alpha$ 2-3 complex [20] followed by geometry optimization. MM calculations were carried out *in vacuo* with structural determined water molecules in the crystal structures.

We cut out the H3–sialoside models (HA: N96–P99, G129–Y161, G181–V196, and N216–I232, 70 amino acid residues) from the optimum structures for *ab initio* FMO calculations (Fig. 1A). This approach covers the amino acid residues on the sialoside binding site especially position 226, besides whose range is outside of substitutions far away from the binding site. Peptide terminals in the models were treated as NH<sub>3</sub> and COO similar manner to the previous study [11]. Amino acid sequences of avian and human H3 differ at ten positions with same secondary

structure as shown in Fig. 1B. The differences at positions 137, 145, 226, and 228 are located on the direct interaction site to Neu5Ac–Gal disaccharides. Since there are few electron density of water molecules around the sialoside binding site–Neu5Ac $\alpha$ (2-3/6)Gal analogue complex in the H3 crystal structures, we computed the model complexes by the FMO method without water molecules.


Single point energies of the model complexes were computed at the FMO-HF level with STO–3G [11], 6–31G, and 6–31G(d) basis sets, and the correlated FMO-MP2/6–31G level of theory. The later method evaluates interfragment van der Waals type stabilization in the protein–ligand complex. Stabilizations of CH– $\pi$  interaction between tryptophan and carbohydrate in the  $\beta$ -galactosidase-substrates or products complex were calculated at the MP2/6–31+G(d) level as 2.4–5.2 kcal/mol [52]. Stabilizing interaction energy of the fucose-benzene complex was estimated 3.0 kcal/mol at the MP2/6–31G(d,p) level of theory [53].

The 70 amino acids on the H3–sialoside complexes were divided into one amino acid residue as a single fragment using automatic fragmentation program in the ABINIT-MP package, and Neu5Ac–Gal receptors were also treated as a single fragment. The receptors and sialoside binding sites were charged to –1 and +1. We calculated single point energies of the complexes ( $E_{\text{complex}}$ ), Neu5Ac–Gal ( $E_{\text{receptor}}$ ),



**Fig. 1** Interaction site of avian influenza A virus HA H3 subtype complexed with avian type receptor Neu5Ac $\alpha$ (2-3)Gal analogue for *ab initio* FMO studies. **A** Ribbon model; avian H3 sialoside binding site; N96–P99, G129–Y161, G181–V196, and N216–I232. CPK model; Neu5Ac $\alpha$ (2-3)Gal disaccharide, **B** sequence alignments of avian/human H3 in the model complexes. yellow; non-matching residues, green; weak matching residues, light blue; strong matching residues, red; helix, blue; sheet

**Fig. 2** Stereo depiction of the avian H3- $\alpha(2-3/6)$  complexes. *left*: avian H3-avian  $\alpha 2-3$  complex. *right*: avian H3-human  $\alpha 2-6$  complex. CPK: Neu5Ac $\alpha(2-3/6)$ Gal disaccharides




and the binding sites ( $E_{H3}$ ) to estimate binding energies ( $\Delta E$ ) between the receptor and H3 by the following expression;  $\Delta E = (E_{\text{receptor}} + E_{H3}) - E_{\text{complex}}$ . *Ab initio* FMO calculations were performed by using ABINIT-MP program [30–39].

## Results and discussion

Figures 2, 3, 4, and 5 show the direct interactions between Neu5Ac-Gal and sialoside binding site on H3 HA1.


Binding energies  $\Delta E$  and the selected IFIEs of Neu5Ac $\alpha(2-3$  or  $2-6)$ Gal with amino acid residues at the FMO-HF/STO-3G and FMO-MP2/6-31G levels are summarized in Tables 1, 2, 3, 4, and 5.

$\Delta E$ s are estimated with electron correlation at the FMO-MP2/6-31G level (Table 1). In the avian H3 complexes, MP2/6-31G  $\Delta E$ s are 43.5 and 40.3 kcal/mol larger than the corresponding HF/STO-3G energies to give  $\Delta E_{\alpha 2-3} - \Delta E_{\alpha 2-6}$  11.4 kcal/mol (entries 1, 3). Extension from minimal basis sets to valence double zeta basis sets at the HF level affords



**Fig. 3** Intermolecular interactions of Neu5Ac $\alpha(2-3/6)$ Gal with amino acid residues on the sialoside binding site in avian H3. **A–C**; avian H3-avian  $\alpha 2-3$  complex. **D–F**; avian H3-human  $\alpha 2-6$  complex. The

*red* and *black dotted lines* represent hydrogen bonds and long range interactions whose distances are given in angstrom



**Fig. 4** Stereo depiction of the human H3–human  $\alpha$ 2-6 complex

the differences  $\Delta E_{\alpha 2-3} - \Delta E_{\alpha 2-6}$  13.1 kcal/mol in the avian H3– $\alpha$ (2-3/6) complexes.


As shown in Tables 2 and 3, various intermolecular lipophilic interactions benefit forming H3–Neu5Ac $\alpha$ (2-3/6) Gal complexes at the FMO-MP2/6–31G level. Neu5-NHCOCH<sub>3</sub> makes CH– $\pi$  interaction network with Trp153 and Ile230 to afford MP2/6–31G IFIEs 8.5 and 8.6 kcal/mol in avian H3– $\alpha$ (2-3/6) complexes, 4.0 kcal/mol in human H3–human  $\alpha$ 2-6 complex (Figs. 3B, E, and 5B, Tables 2 and 3; entry 5), while the corresponding HF/STO–3G energies do not evaluate the CH– $\pi$  interaction stabilization. Gly134 and Thr155 directly interact with Neu 5-NHCOCH<sub>3</sub>, but do not contribute to the HA– $\alpha$ (2-3/6) binding (Tables 2 and 3; entries 6, 12). Hydrophobic association between Leu194 and Neu7-CH, 9-CH<sub>2</sub> has bonding IFIEs 2.4–3.7 kcal/mol at the MP2/6–31G level (Figs. 3C, F, and 5C, Tables 2 and 3; entry 13). Intermolecular hydrophilic interactions also advantage the forming HA–sialoside complex. In the avian H3 com-

plexes, MP2/6–31G IFIEs are 5.4–9.1 kcal/mol larger than the corresponding HF/STO–3G energies (Table 2; entries 1, 2, 4, 8, 10). Especially, Gln226 on avian H3 interacts with avian  $\alpha$ 2-3 whose MP2/6–31G IFIE is estimated 15.1 kcal/mol larger than its HF/STO–3G energy (Fig. 3A, Table 2; entry 3). In contrast, IFIE between Ser137 and Neu 1-CO1O1' has 33.9 kcal/mol at the MP2/6–31G which is 2.6 kcal/mol smaller than the HF/STO–3G energy (Fig. 3B, Table 2; entry 9). Gly135 clearly forms an intermolecular hydrogen bond with Neu 5-NHCOCH<sub>3</sub>, but whose IFIE are quite smaller than the others (Tables 2 and 3; entry 7).

In the following section, we discuss molecular interaction between Neu5Ac $\alpha$ (2-3/6)Gal and amino acid residues on the sialoside binding site in each complex with  $\Delta E$ s and IFIEs at the FMO-MP2/6–31G level.

#### Avian H3–Neu5Ac $\alpha$ (2-3 or 2-6)Gal complexes

Avian H3 binds to avian receptor Neu5Ac $\alpha$ (2-3)Gal 11.4 kcal/mol stronger than to human receptor Neu5Ac $\alpha$ (2-6)Gal with taking account of intermolecular lipophilic stabilization at the FMO-MP2/6–31G level (Table 1; entries 1, 3). In the avian H3–avian  $\alpha$ 2-3 complex, side chain NH<sub>2</sub>CO on Gln226 forms intermolecular hydrogen bond network with 8-OH, 1-COO on Neu5Ac, Gal 4-OH, and weakly with glycoside oxygen O3 to give IFIE 23.9 kcal/mol, which is 6.5 kcal/mol larger than the corresponding IFIE in avian H3–human  $\alpha$ 2-6 complex (Fig. 3A, D, Table 2; entry 3). Besides, hydrogen bond formations of Neu 1-COO with Ser136, Ser137 and Asn145 strongly stabilize the avian H3–avian  $\alpha$ 2-3 complex by IFIEs 27.1, 33.9, and 13.2 kcal/mol (Fig. 3B, E, Table 2; entries 8–10). The later interaction consists of intermolecular hydrogen bonds between Asn145 side chain CONH<sub>2</sub> and O1'O1C1 on Neu5Ac $\alpha$ (2-3)Gal with 2.78 Å. Gly228 on influenza A virus H3 subtype correlates with the shift of virus host range between avian and human. In our study, Gly228 interacts with Neu5Ac $\alpha$ (2-3/6)Gal disaccharides by IFIEs 4.7 and 4.3 kcal/mol (Fig. 3C, F, Table 2; entry 11).



**Fig. 5** Intermolecular interactions of human Neu5Ac $\alpha$ (2-6)Gal with amino acid residues on the sialoside binding site in human H3

**Table 1** Binding energies  $\Delta E$  between H3s and Neu5Ac $\alpha$ (2-3/6)Gal estimated by *ab initio* FMO calculations

Entry	$\Delta E^a$	H3	HF/STO-3G	MP2/6-31G
1	$\Delta E_{\alpha 2-3}$	Avian	136.9 <sup>b</sup>	180.4
2		Human Leu226Gln	137.6	179.7
3	$\Delta E_{\alpha 2-6}$	Avian	128.7 <sup>b</sup>	169.0
4		Avian Gln226Leu	118.2 <sup>b</sup>	157.6
5		Human	117.9	154.3

<sup>a</sup>  $\Delta E$ s were given in kilocalories per mole

<sup>b</sup> These data were previously reported in [11]

**Table 2** Interfragment interaction energies of Neu5Ac $\alpha$ (2-3/6)Gal with amino acid residues on the sialoside binding site in avian H3

Entry	Avian H3 amino acid	Interaction sites on Neu5Ac $\alpha$ (2-3/6)Gal		$\alpha 2-3$		$\alpha 2-6$	
		$\alpha 2-3$	$\alpha 2-6$	HF/STO-3G <sup>a</sup>	MP2/6-31G	HF/STO-3G <sup>a</sup>	MP2/6-31G
1	His183	Neu 8,9-OH		5.2	12.4	7.5	14.5
2	Tyr98	Neu 8-OH		6.7	14.6	5.1	14.2
3	Gln226	Neu 8-OH, 1-COIO1', Gal 4-OH	Neu 8-OH, 1-COIO1'	8.8	23.9	10.5	17.4
4	Ala138	Neu 1-COIO1'		10.1	15.8	11.1	16.5
5	Trp153	Neu 5-NHCOCH <sub>3</sub>		0.1	8.5	-0.4	8.6
6	Gly134	Neu 5-NHCOCH <sub>3</sub>		-1.4	-1.3	-1.9	-1.8
7	Gly135	Neu 5-NHCOCH <sub>3</sub>		1.9	2.6	2.7	3.9
8	Ser136	Neu 1-COIO1'		20.5	27.1	16.2	23.2
9	Ser137	Neu 1-COIO1'		36.5	33.9	27.7	28.7
10	Asn145	Neu 1-COIO1'		6.6	13.2	10.7	18.2
11	Gly228	–		1.7	4.7	1.8	4.3
12	Thr155	Neu 5-NHCOCH <sub>3</sub>		-1.8	0.1	-2.2	-0.3
13	Leu194	Neu 7-CH, 9-CH <sub>2</sub>		-1.2	3.7	-1.8	2.5
14	Sum				159.2		149.9

IFIEs were given in kilocalories per mole. IFIEs will be estimated lower by FMO-MP2/6-31G(d) energy calculations of the corresponding QM/MM optimum geometry

<sup>a</sup> IFIEs at the FMO-HF/STO-3G level were reported in [11]

**Table 3** Interfragment interaction energies of human Neu5Ac $\alpha$ (2-6)Gal with amino acid residues on the sialoside binding site in human H3

Entry	Human H3 amino acid	Interaction sites on Neu5Ac $\alpha$ (2-6)Gal	HF/STO-3G	MP2/6-31G
1	His183	Neu 8,9-OH	3.5	8.9
2	Tyr98	Neu 8-OH	6.4	16.2
3	Leu226	Gal 6-CH <sub>2</sub>	-1.3	6.1
4	Ala138	Neu 1-COIO1'	8.6	15.7
5	Trp153	Neu 5-NHCOCH <sub>3</sub>	-0.2	4.0
6	Gly134	Neu 5-NHCOCH <sub>3</sub>	-1.2	-0.9
7	Gly135	Neu 5-NHCOCH <sub>3</sub>	1.0	1.1
8	Ser136	Neu 1-COIO1'	24.2	29.3
9	Asn137	Neu 1-COIO1'	38.5	45.5
10	Ser145	Neu 1-COIO1'	0.0	2.6
11	Ser228	Neu 9-OH	6.8	14.8
12	Thr155	Neu 5-NHCOCH <sub>3</sub>	-1.6	-0.3
13	Leu194	Neu 7-CH, 9-CH <sub>2</sub>	-2.0	2.4
14	Sum			145.3

**Table 4** Interfragment interaction energies of human Neu5Ac $\alpha$ (2-6)Gal with amino acid residues on the sialoside binding site in avian Gln226Leu H3

Entry	Avian Q226L H3 amino acid	Interaction sites on Neu5Ac $\alpha$ (2-6)Gal	MP2/6–31G
1	His183	Neu 8,9-OH	14.3
2	Tyr98	Neu 8-OH	14.5
3	Leu226	Gal 6-CH <sub>2</sub>	4.8
4	Ala138	Neu 1-CO1O1'	16.5
5	Trp153	Neu 5-NHCOCH <sub>3</sub>	8.6
6	Gly134	Neu 5-NHCOCH <sub>3</sub>	-1.6
7	Gly135	Neu 5-NHCOCH <sub>3</sub>	3.6
8	Ser136	Neu 1-CO1O1'	25.0
9	Ser137	Neu 1-CO1O1'	26.2
10	Asn145	Neu 1-CO1O1'	18.1
11	Gly228	–	4.4
12	Thr155	Neu 5-NHCOCH <sub>3</sub>	-0.2
13	Leu194	Neu 7-CH, 9-CH <sub>2</sub>	2.7
14	Sum		136.9

Comparison between  $\Delta E$ s and sums of 13 IFIEs, the formers are about 20 kcal/mol larger than the 13 IFIEs sums (Table 1; entries 1, 3, Table 2; entry 14). IFIEs sum difference of 9.3 kcal/mol mainly causes the binding energy difference ( $\Delta E_{\alpha 2-3} - \Delta E_{\alpha 2-6}$ ) of 11.4 kcal/mol. In the X-ray crystal structure of avian H3 complexed with  $\alpha$ (2-3 or 2-6)-pentasaccharide receptors, the sialoside binding site reliably recognizes the non-reducing terminal Neu5Ac-Gal [19] that approximately allows our theoretical approach.

#### Human H3–Human Neu5Ac $\alpha$ (2-6)Gal complex


Binding energy  $\Delta E$  in the human H3–human Neu5Ac $\alpha$ (2-6)Gal disaccharide was estimated 154.3 kcal/mol at the FMO-MP2/6–31G level (Table 1; entry 5). Human H3 complex has an intramolecular hydrogen bond at Neu 8-OH $\cdots$ O1C1O1' instead of intermolecular Gln226 $\cdots$ Neu5Ac interaction observed in the avian H3 complex (Fig. 5A), thus  $\Delta E$  in the human H3 complex is smaller by 15 kcal/mol

than the corresponding  $\Delta E$  in the avian Gln226 H3 complex (Table 1; entries 3, 5). Neu 1-COO interacts with Ser136 by 29.3 kcal/mol, Asn137 by 45.5 kcal/mol, and Ser145 by 2.6 kcal/mol (Table 3; entries 8–10) whose summation is larger by 7.3 kcal/mol than that in the avian H3–human  $\alpha$ 2-6 complex (Table 2; entries 8–10). Intermolecular hydrogen bond network between His183, Tyr98 and Neu 9,8-OH stabilizes the human H3–human  $\alpha$ 2-6 complex with IFIEs of 8.9 and 16.2 kcal/mol (Fig. 5C, Table 3; entries 1, 2), therefore human Tyr98Phe or His183Phe H3 viruses lose this stabilization by hydrogen bonds decomposition to decrease severely their binding affinities [25]. Tyr98 forms a hydrogen bond with His183 by IFIE of 10.5 kcal/mol, and CH– $\pi$  interaction with Leu226 by 1.2 kcal/mol.

Inter- and intra-molecular lipophilic association is significant for the human H3–human  $\alpha$ 2-6 binding. Leu226 interacts with Gal 6-CH<sub>2</sub> on human  $\alpha$ 2-6 by lipophilic IFIE of 6.1 kcal/mol (Fig. 5A, Table 3; entry 3)


**Table 5** Interfragment interaction energies of avian Neu5Ac $\alpha$ (2-3)Gal with amino acid residues on the sialoside binding site in human Leu226Gln H3

Entry	Human L226Q H3 amino acid	Interaction sites on Neu 5Ac $\alpha$ (2-3)Gal	MP2/6–31G
1	His183	Neu 8,9-OH	8.6
2	Tyr98	Neu 8-OH	15.6
3	Gln226	Neu 8-OH, 1-CO1O1', Gal 4-OH	31.3
4	Ala138	Neu 1-CO1O1'	15.2
5	Trp153	Neu 5-NHCOCH <sub>3</sub>	7.9
6	Gly134	Neu 5-NHCOCH <sub>3</sub>	-1.3
7	Gly135	Neu 5-NHCOCH <sub>3</sub>	2.6
8	Ser136	Neu 1-CO1O1'	27.4
9	Asn137	Neu 1-CO1O1'	49.1
10	Ser145	Neu 1-CO1O1'	3.1
11	Ser228	Neu 9-OH	14.3
12	Thr155	Neu 5-NHCOCH <sub>3</sub>	-0.4
13	Leu194	Neu 7-CH, 9-CH <sub>2</sub>	1.6
14	Sum		175.0




**Fig. 6** Intermolecular interactions of human Neu5Ac $\alpha$ (2-6)Gal with amino acid residues on the sialoside binding site in avian Gln226Leu H3

that is supported by a lipophilic network beyond Leu226 consisted Tyr98, Pro99, Ala138, (CH<sub>2</sub>)<sub>3</sub> on Arg220 and Arg229, Ile230, and Trp153. Hydrophobic site on Neu 7-CH 9-CH<sub>2</sub> associates with lipophilic Leu194 by IFIE of 2.4 kcal/mol (Fig. 5C, Table 3; entry 13) supported by hydrophobic groups around Leu194 such as Thr155, Tyr195, (CH<sub>2</sub>)<sub>2</sub> on Glu190, and CH<sub>2</sub> on Ser193. Since human Leu194Ala H3 virus does not agglutinate human



**Fig. 7** Intermolecular interactions of avian Neu5Ac $\alpha$ (2-3)Gal with amino acid residues on the sialoside binding site in human Leu226Gln H3



**Fig. 8** Binding energy profile at the FMO-MP2/6–31G level on the relationship between influenza H3 virus host range and H3– $\alpha$ (2-3 or 2-6) receptors binding.  $\Delta E$ s are shown in Table 1

erythrocyte [25], the lipophilic association around Leu194 is significant for the human H3–human  $\alpha$ 2-6 binding affinity. A substitution at Ser193 to Ile in human H3 HA1 increases the hydrophobic capacity around Leu194 to give higher binding affinity to human  $\alpha$ 2-6 [26]. The two lipophilic networks around Leu226 $\cdots$ Gal 6-CH<sub>2</sub> and Leu194 $\cdots$ Neu 7-CH 9-CH<sub>2</sub> are connected by a hydrogen bond formation of indole NH Trp153 with OH Tyr195 (Fig. 5C). However, both substitutions Trp153Phe or Tyr195Phe partially decrease its binding affinity [8, 25].

Avian Gln226Leu H3–Human Neu5Ac $\alpha$ (2-6)Gal and human Leu226Gln H3–Avian Neu5Ac $\alpha$ (2-3)Gal complexes

Figures 6 and 7 show the interaction of Neu5Ac-Gal with avian Gln226Leu/human Leu226Gln H3s. Avian Gln226Leu H3 moderately binds to human  $\alpha$ 2-6 with  $\Delta E$  157.6 kcal/mol at the FMO-MP2/6–31G level (Table 1; entry 4). This binding energy is quite similar to  $\Delta E_{\alpha 2-6}$  154.3 kcal/mol in the human H3 complex with amino acid difference at nine positions (Fig. 1B), therefore avian Gln226Leu H3 virus can infect human with the same level as human H3 virus. In the avian Gln226Leu H3–human  $\alpha$ 2-6 complex, Leu226 associates with lipophilic Gal 6-CH<sub>2</sub> whose IFIE is smaller by 1.3 kcal/mol than that of Leu226 $\cdots$ Gal 6-CH<sub>2</sub> interaction in the human H3 complex (Tables 3 and 4; entry 3). Besides Neu 8-OH makes an intramolecular hydrogen bond with O1C1O1'-Neu similar manner to that in the human Leu226 H3 complex (Figs. 5A and 6).



Human Leu226Gln H3 interacts with avian  $\alpha$ 2-3 by  $\Delta E_{\alpha 2-3}$  of 179.7 kcal/mol that is almost the same value as  $\Delta E_{\alpha 2-3}$  of 180.4 kcal/mol in the avian H3 complex at the FMO-MP2/6–31G level (Table 1; entries 1, 2). In the human Leu226Gln H3–avian  $\alpha$ 2-3 complex, Gln226 makes intermolecular hydrogen bond network with 8-OH, 1-CO1O1' on Neu5Ac and Gal 4-OH to give IFIE 31.1 kcal/mol (Fig. 7, Table 5; entry 3). We previously reported that A/Udorn Leu226Gln (human Leu226Gln H3N2) bound to avian  $\alpha$ 2-3 receptor [14]. Our *ab initio* FMO studies have confirmed that human Leu226Gln H3 moderately interacts with avian  $\alpha$ 2-3 analogue at the correlated FMO-MP2/6–31G level.

## Conclusion

We theoretically studied influenza A virus hemagglutinin H3 subtype complexed with avian/human type receptor Neu5Ac $\alpha$ (2-3 or 2-6)Gal analogues by *ab initio* FMO method at the correlated MP2/6–31G level. Avian H3 bound to avian  $\alpha$ 2-3 11.4 kcal/mol stronger than to human  $\alpha$ 2-6 in the model complexes with taking account of intermolecular lipophilic interaction. Single point substitution at the position 226 on H3 subtype sialoside binding site HA1 changes its binding affinity between avian  $\alpha$ 2-3 and human  $\alpha$ 2-6. Our *ab initio* FMO studies showed that the binding energy of avian Gln226Leu H3 with human  $\alpha$ 2-6 was similar value to that in the human H3–human  $\alpha$ 2-6 complex at the FMO-MP2/6–31G level with amino acid differences at nine positions in our models. Thus avian Gln226Leu H3 virus can infect human with the same level as human H3 virus. Opposite mutation Leu226Gln in the human H3 gave the moderate binding energy to avian  $\alpha$ 2-3 that supported our previous virus-sialoside binding assay.

As shown in a schematic summary (Fig. 8), *ab initio* FMO studies revealed the relationship between influenza H3 virus host range and H3– $\alpha$ (2-3 or 2-6) receptors binding. Our theoretical approach will predict the infectious level of new viruses and point out some unknown dangerous mutation positions on HA in advance against human pandemic influenza.

**Acknowledgment** This work was supported in part by CREST, Japan Science and Technology Agency (JST), and a Grant-in-Aid (No. 17101007 to M. Kiso) for Scientific Research from Japan Society for the Promotion of Science.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

- Shinya, K., Ebina, M., Yamada, S., Ono, M., Kasai, N., Kawaoka, Y.: Avian flu: influenza virus receptors in the human airway. *Nature* **440**, 435–436 (2006)
- van Riel, D., Munster, V.J., de Wit, E., Rimmelzwaan, G.F., Fouchier, R.A.M., Osterhaus, A.D.M.E., Kuiken, T.: H5N1 virus attachment to lower respiratory tract. *Science* **312**, 399 (2006)
- Yamada, S., Suzuki, Y., Suzuki, T., Le, M.Q., Nidom, C.A., Sakai-Tagawa, Y., Muramoto, Y., Ito, M., Kiso, M., Horimoto, T., Shinya, K., Sawada, T., Kiso, M., Usui, T., Murata, T., Lin, Y., Hay, A., Haire, L.F., Stevens, D.J., Russell, R.J., Gamblin, S.J., Skehel, J.J., Kawaoka, Y.: Haemagglutinin mutations responsible for the binding of H5N1 influenza A viruses to human-type receptors. *Nature* **444**, 378–382 (2006)
- Stevens, J., Blixt, O., Tumpey, T.M., Taubenberger, J.K., Paulson, J.C., Wilson, I.A.: Structure and receptor specificity of the hemagglutinin from an H5N1 influenza virus. *Science* **312**, 404–410 (2006)
- Yang, Z.-Y., Wei, C.-J., Kong, W.-P., Wu, L., Xu, L., Smith, D.F., Nabel, G.J.: Mutants with altered receptor binding specificity immunization by avian H5 influenza hemagglutinin. *Science* **317**, 825–828 (2007)
- Auewarakul, P., Suptawiwat, O., Kongchanagul, A., Sangma, C., Suzuki, Y., Ungchusak, K., Louisirirotchanakul, S., Lerdsamran, H., Pooruk, P., Thitithanyanont, A., Pittayawonganon, C., Guo, C.-T., Hiramatsu, H., Jampangern, W., Chunsutthiwat, S., Puthavathana, P.: An avian influenza H5N1 virus that binds to a human-type receptor. *J. Virol.* **81**, 9950–9955 (2007)
- Suzuki, Y.: Sialobiology of influenza: molecular mechanism of host range variation of influenza viruses. *Biol. Pharm. Bull.* **28**, 399–408 (2005)
- Skehel, J.J., Wiley, D.C.: Receptor binding and membrane fusion in virus entry: the influenza hemagglutinin. *Ann. Rev. Biochem.* **69**, 531–569 (2000)
- Horimoto, T., Kawaoka, Y.: Influenza: lessons from past pandemics, warnings from current incidents. *Nat. Rev. Microbiol.* **3**, 591–600 (2005)
- Matrosovich, M.N., Klenk, H.D., Kawaoka, Y.: Receptor specificity, host-range, and pathogenicity of influenza viruses. In: Kawaoka, Y. (ed.) *Influenza Virology Current Topics*, pp. 95–137. Caister Academic Press, Norfolk (2006)
- Sawada, T., Hashimoto, T., Nakano, H., Suzuki, T., Ishida, H., Kiso, M.: Why does avian influenza A virus hemagglutinin bind to avian receptor stronger than to human receptor? *Ab initio* fragment molecular orbital studies. *Biochem. Biophys. Res. Commun.* **351**, 40–43 (2006)
- Sawada, T., Hashimoto, T., Nakano, H., Suzuki, T., Suzuki, Y., Kawaoka, Y., Ishida, H., Kiso, M.: Influenza viral hemagglutinin complicated shape is advantageous to its binding affinity for sialosaccharide receptor. *Biochem. Biophys. Res. Commun.* **355**, 6–9 (2007)
- Ito, T., Suzuki, Y., Mitnaul, L., Vines, A., Kida, H., Kawaoka, Y.: Receptor specificity of influenza A viruses correlates with the agglutination of erythrocytes from different animal species. *Virology* **227**, 493–499 (1997)
- Ito, T., Suzuki, Y., Suzuki, T., Takada, A., Horimoto, T., Wells, K., Kida, H., Otsuki, K., Kiso, M., Ishida, H., Kawaoka, Y.: Recognition of N-glycolylneuraminic acid linked to galactose by the  $\alpha$ 2,3 linkage is associated with intestinal replication of influenza A virus in ducks. *J. Virol.* **74**, 9300–9305 (2000)
- Suzuki, Y., Nakano, T., Ito, T., Watanabe, N., Toda, Y., Guiyun, X., Suzuki, T., Kobayashi, T., Kimura, Y., Yamada, A., Sugawara, K., Nishimura, H., Kitame, F., Nakamura, K., Deya, E., Kiso, M., Hasegawa, A.: Structural determination of gangliosides that bind to influenza A, B, and C viruses by an improved binding assay:

- strain-specific receptor epitopes in sialo-sugar chains. *Virology* **189**, 121–131 (1992)
16. Ito, T., Suzuki, Y., Takada, A., Kawamoto, A., Otsuki, K., Masuda, H., Yamada, M., Suzuki, T., Kida, H., Kawaoka, Y.: Differences in sialic acid–galactose linkages in the chicken egg amnion and allantois influence human influenza virus receptor specificity and variant selection. *J. Virol.* **71**, 3357–3362 (1997)
  17. Masuda, H., Suzuki, T., Sugiyama, Y., Horiike, G., Murakami, K., Miyamoto, D., Hidari, K.I.-P.J., Ito, T., Kida, H., Kiso, M., Fukunaga, K., Ohuchi, M., Toyoda, T., Ishihama, A., Kawaoka, Y., Suzuki, Y.: Substitution of amino acid residue in influenza A virus hemagglutinin affects recognition of sialyl-oligosaccharides containing N-glycolylneuraminic acid. *FEBS Lett.* **464**, 71–74 (1999)
  18. Shinya, K., Hatta, M., Yamada, S., Takada, A., Watanabe, S., Halfmann, P., Horimoto, T., Neumann, G., Hyun Kim, J., Lim, W., Guan, Y., Peiris, M., Kiso, M., Suzuki, T., Suzuki, Y., Kawaoka, Y.: Characterization of a human H5N1 influenza A virus isolated in 2003. *J. Virol.* **79**, 9926–9932 (2005)
  19. Ha, Y., Stevens, D.J., Skehel, J.J., Wiley, D.C.: X-ray structure of the hemagglutinin of a potential H3 avian progenitor of the 1968 Hong Kong pandemic influenza virus. *Virology* **309**, 209–218 (2003)
  20. Weis, W., Brown, J.H., Cusack, S., Paulson, J.C., Skehel, J.J., Wiley, D.C.: Structure of the influenza virus haemagglutinin complexed with its receptor, sialic acid. *Nature* **333**, 426–431 (1988)
  21. Connor, R.J., Kawaoka, Y., Webster, R.G., Paulson, J.C.: Receptor specificity in human, avian, and equine H2 and H3 influenza virus isolates. *Virology* **205**, 17–23 (1994)
  22. Gambaryan, A.S., Piskarev, V.E., Yamskov, I.A., Sakharov, A.M., Tuzikov, A.B., Bovin, N.V., Nifant'ev, N.E., Matrosovich, M.N.: Human influenza virus recognition of sialyloligosaccharides. *FEBS Lett.* **366**, 57–60 (1995)
  23. Gambaryan, A.S., Tuzikov, A.B., Piskarev, V.E., Yamnikova, S.S., Lvov, D.K., Robertson, J.S., Bovin, N.V., Matrosovich, M.N.: Specification of receptor-binding phenotypes of influenza virus isolates from different hosts using synthetic sialylglycopolymers: nonegg-adapted human H1 and H3 influenza A and influenza B viruses share a common high binding affinity for 6'-sialyl (N-acetyl)lactosamine. *Virology* **232**, 345–350 (1997)
  24. Nobusawa, E., Ishihara, H., Morishita, T., Sato, K., Nakajima, K.: Change in receptor-binding specificity of recent human influenza A viruses (H3N2): a single amino acid change in hemagglutinin altered its recognition of sialyloligosaccharides. *Virology* **278**, 587–596 (2000)
  25. Martin, J., Wharton, S.A., Lin, Y.P., Takemoto, D.K., Skehel, J.J., Wiley, D.C., Steinhauer, D.A.: Studies of the binding properties of influenza hemagglutinin receptor-site mutants. *Virology* **241**, 101–111 (1998)
  26. Daniels, R.S., Jeffries, S., Yates, P., Schild, G.C., Rogers, G.N., Paulson, J.C., Wharton, S.A., Douglas, A.R., Skehel, J.J., Wiley, D.C.: The receptor-binding and membrane-fusion properties of influenza virus variants selected using anti-hemagglutinin monoclonal antibodies. *EMBO J.* **6**, 1459–1465 (1987)
  27. Rogers, G.N., Paulson, J.C., Daniels, R.S., Skehel, J.J., Wilson, I.A., Wiley, D.C.: Single amino acid substitutions in influenza haemagglutinin change receptor binding specificity. *Nature* **304**, 76–78 (1983)
  28. Naeve, C.W., Hinshaw, V.S., Webster, R.G.: Mutations in the hemagglutinin receptor-binding site can change the biological properties of an influenza virus. *J. Virol.* **51**, 567–569 (1984)
  29. Rogers, G.N., Daniels, R.S., Skehel, J.J., Wiley, D.C., Wang, X.F., Higs, H.H., Paulson, J.C.: Host-mediated selection of influenza virus receptor variants. Sialic acid- $\alpha$ 2, 6 gal-specific clones of A/duck/Ukraine/1/63 revert to sialic acid- $\alpha$ 2,3Gal-specific wild type in ovo. *J. Biol. Chem.* **260**, 7362–7367 (1985)
  30. Kitaura, K., Sawai, T., Asada, T., Nakano, T., Uebayasi, M.: Pair interaction molecular orbital method: an approximate computational method for molecular interactions. *Chem. Phys. Lett.* **312**, 319–324 (1999)
  31. Kitaura, K., Ikeo, E., Asada, T., Nakano, T., Uebayasi, M.: Fragment molecular orbital method: an approximate computational method for large molecules. *Chem. Phys. Lett.* **313**, 701–706 (1999)
  32. Nakano, T., Kaminuma, T., Sato, T., Akiyama, Y., Uebayasi, M., Kitaura, K.: Fragment molecular orbital method: application to polypeptides. *Chem. Phys. Lett.* **318**, 614–618 (2000)
  33. Kitaura, K., Sugiki, S., Nakano, T., Komeiji, Y., Uebayasi, M.: Fragment molecular orbital method: analytical energy gradients. *Chem. Phys. Lett.* **336**, 163–170 (2001)
  34. Nakano, T., Kaminuma, T., Sato, T., Fukuzawa, K., Akiyama, Y., Uebayasi, M., Kitaura, K.: Fragment molecular orbital method: use of approximate electrostatic potential. *Chem. Phys. Lett.* **351**, 475–480 (2002)
  35. Mochizuki, Y., Nakano, T., Koikegami, S., Tanimori, S., Abe, Y., Nagashima, U., Kitaura, K.: A parallelized integral-direct second-order Møller–Plesset perturbation theory method with a fragment molecular orbital scheme. *Theor. Chem. Acc.* **112**, 442–452 (2004)
  36. Mochizuki, Y., Koikegami, S., Nakano, T., Amari, S., Kitaura, K.: Large scale MP2 calculations with fragment molecular orbital scheme. *Chem. Phys. Lett.* **396**, 473–479 (2004)
  37. Mochizuki, Y., Koikegami, S., Amari, S., Segawa, K., Kitaura, K., Nakano, T.: Configuration interaction singles method with multilayer fragment molecular orbital scheme. *Chem. Phys. Lett.* **406**, 283–288 (2005)
  38. Mochizuki, Y.: A size-extensive modification of super-CI for orbital relaxation. *Chem. Phys. Lett.* **410**, 165–171 (2005)
  39. Mochizuki, Y., Fukuzawa, K., Kato, A., Tanaka, S., Kitaura, K., Nakano, T.: A configuration analysis for fragment interaction. *Chem. Phys. Lett.* **410**, 247–253 (2005)
  40. Fedorov, D.G., Kitaura, K.: Extending the power of quantum chemistry to large systems with the fragment molecular orbital method. *J. Phys. Chem. A* **111**, 6904–6914 (2007)
  41. Fukuzawa, K., Kitaura, K., Uebayasi, M., Nakano, K., Kaminuma, T., Nakano, T.: *Ab initio* quantum mechanical study of the binding energies of human estrogen receptor with its ligands: an application of fragment molecular orbital method. *J. Comput. Chem.* **26**, 1–10 (2005)
  42. Fukuzawa, K., Komeiji, Y., Mochizuki, Y., Kato, A., Nakano, T., Tanaka, S.: Intra- and intermolecular interactions between cyclic-AMP receptor protein and DNA: *ab initio* fragment molecular orbital study. *J. Comput. Chem.* **27**, 948–960 (2006)
  43. Fukuzawa, K., Mochizuki, Y., Shigenori, T., Kitaura, K., Nakano, T.: Molecular interactions between estrogen receptor and its ligand studied by the *ab initio* fragment molecular orbital method. *J. Phys. Chem. B* **110**, 16102–16110 (2006)
  44. Yamamoto, K., Abe, D., Yoshimoto, N., Choi, M., Yamagishi, K., Tokiwa, H., Shimizu, M., Makishima, M., Yamada, S.: Vitamin D receptor: ligand recognition and allosteric network. *J. Med. Chem.* **49**, 1313–1324 (2006)
  45. Yamagishi, K., Yamamoto, K., Yamada, S., Tokiwa, H.: Functions of key residues in the ligand-binding pocket of vitamin D receptor: fragment molecular orbital-interfragment interaction energy analysis. *Chem. Phys. Lett.* **420**, 465–468 (2006)
  46. Nakano, T., Mochizuki, Y., Amari, S., Kobayashi, M., Fukuzawa, K., Tanaka, S.: Applications of the fragment molecular orbital (FMO) method to nano-bio field. *J. Comput. Chem. Jpn.* **6**, 173–184 (2007)
  47. Fukuzawa, K., Nakano, T., Kato, A., Mochizuki, Y., Tanaka, S.: Applications of the fragment molecular orbital method for biomacromolecules. *J. Comput. Chem. Jpn.* **6**, 185–197 (2007)
  48. Fedorov, D.G., Kitaura, K.: Second order Møller–Plesset perturbation theory based upon the fragment molecular orbital method (FMO). *J. Chem. Phys.*, **121**, 2483–2490 (2004)

49. Maple, J.R., Hwang, M.-J., Jalkanen, K.J., Stockfish, T.P., Hagler, A.T.: Derivation of class II force fields: quantum force field for amides, peptides, and related compounds. *J. Comput. Chem.* **19**, 430–458 (1998)
50. Eisen, M.B., Sabesan, S., Skehel, J.J., Wiley, D.C.: Binding of the influenza A virus to cell-surface receptors: structures of five hemagglutinin-sialyloligosaccharide complexes determined by x-ray crystallography. *Virology* **232**, 19–31 (1997)
51. Sauter, N.K., Hanson, J.E., Glick, G.D., Brown, J.H., Crowther, R.L., Park, S.J., Skehel, J.J., Wiley, D.C.: Binding of influenza virus hemagglutinin to analogs of its cell-surface receptor, sialic acid: analysis by proton nuclear magnetic resonance spectroscopy and x-ray crystallography. *Biochemistry* **31**, 9609–9621 (1992)
52. Spiwok, V., Lipovová, P., Skálová, T., Buchtelová, E., Hašek, J., Králova, B.: Role of CH/ $\pi$  interactions in substrate binding by *Escherichia coli*  $\beta$ -galactosidase. *Carbohydr. Res.* **339**, 2275–2280 (2004)
53. Fernández-Alonso, M.D.C., Cañada, F.J., Jiménez-Barbero, J., Gabriel Cuevas, G.: Molecular recognition of saccharides by proteins. insights on the origin of the carbohydrate–aromatic interactions. *J. Am. Chem. Soc.* **127**, 7379–7386 (2005)