

Title	ABELIAN SUBGROUPS OF THE MAPPING CLASS GROUPS FOR NON-ORIENTABLE SURFACES
Author(s)	Kuno, Erika
Citation	Osaka Journal of Mathematics. 56(1) P.91-P.100
Issue Date	2019-01
Text Version	publisher
URL	https://doi.org/10.18910/71137
DOI	10.18910/71137
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

ABELIAN SUBGROUPS OF THE MAPPING CLASS GROUPS FOR NON-ORIENTABLE SURFACES

ERIKA KUNO

(Received March 23, 2017, revised August 23, 2017)

Abstract

One of the basic and important problems to study algebraic structures of the mapping class groups is finding abelian subgroups included in the mapping class groups. Birman-Lubotzky-McCarthy gave the answer of this question for the orientable surfaces, namely, they proved that any abelian subgroup of the mapping class groups for orientable surfaces of genus g with b boundary components and c connected components is finitely generated and the maximal torsion-free rank of it is $3g + b - 3c$. In the present paper, we prove that any abelian subgroup of the mapping class group of a compact connected non-orientable surface N of genus $g \geq 1$ with $n \geq 0$ boundary components whose Euler characteristic is negative is finitely generated and the maximal torsion-free rank of it is $\frac{3}{2}(g - 1) + n - 2$ if g is odd and $\frac{3}{2}g + n - 3$ if g is even.

1. Introduction

Let S be a compact orientable surface of genus g with b boundary components and c connected components. Assume each connected component of S has a negative Euler characteristic. Let $S_{g,n}$ be a compact connected orientable surface of genus g with n boundary components. We also write S as $S_{g,n}$. We denote by $\mathcal{M}(S)$ the mapping class group of S , that is, the group of isotopy classes of orientation preserving self-homeomorphisms of S with isotopies fixing each boundary component of S setwise. Birman-Lubotzky-McCarthy [3] proved that any abelian subgroup of $\mathcal{M}(S)$ is finitely generated with torsion-free rank bounded by $3g + b - 3c$. Kim-Koberda [4] quoted it in their paper and renew it as follows: if S is a compact connected orientable surface and G is a torsion-free abelian subgroup of $\mathcal{M}(S)$, then G is isomorphic to a finitely generated subgroup of $\mathcal{M}(S)$ which consists of Dehn twists and pseudo-Anosov elements on some connected subsurface on S whose supports are pairwise disjoint. Let $N = N_{g,n}$ be a compact connected non-orientable surface of genus $g \geq 1$ with $n \geq 0$ boundary components whose Euler characteristic is negative, that is, $g + n \geq 3$. The mapping class group $\mathcal{M}(N)$ of N is the group consisting of the isotopy classes of self-homeomorphisms on N . Similarly, the isotopies fix each boundary component of N setwise. Let $p: S_{g-1,2n} \rightarrow N_{g,n}$ be the double covering map of $N_{g,n}$. Because $\mathcal{M}(N_{g,n})$ is a subgroup of $\mathcal{M}(S_{g-1,2n})$, it follows that any abelian subgroup of $\mathcal{M}(N_{g,n})$ is finitely generated with torsion-free rank bounded by $3(g - 1) + 2b$ by the result of Birman-Lubotzky-McCarthy ([3, Theorem A]). In this paper, we write $\iota: \mathcal{M}(N_{g,n}) \rightarrow \mathcal{M}(S_{g-1,2n})$ for the injective homomorphism. By Szepietowski [5] the image $\iota(\mathcal{M}(N_{g,n}))$ includes no Dehn twists in $\mathcal{M}(S_{g-1,2n})$. Therefore, our motivation in this paper is to detect finitely generated groups isomorphic

to torsion-free abelian subgroups of $\mathcal{M}(N_{g,n})$. Applying Birman-Lubotzky-McCarthy’s arguments to the mapping class groups of non-orientable surfaces, we obtain the following theorem:

Theorem 1.1. *Let N be a non-orientable surface whose Euler characteristic is negative and G a torsion-free abelian subgroup of $\mathcal{M}(N)$. Then, the torsion-free rank of G is $k \leq \frac{3}{2}(g - 1) + n - 2$ if g is odd and $k \leq \frac{3}{2}g + n - 3$ if g is even.*

The idea to prove Theorem 1.1 is to show G is isomorphic to a subgroup $\langle \tau_1, \dots, \tau_k \rangle < \mathcal{M}(N)$, where each τ_i is an isotopy class of a Dehn twist and the supports of τ_i and τ_j are disjoint for $i \neq j$ (see Figure 1).




Fig. 1. The maximal number of pairwise disjoint two-sided non-separating curves on odd genus surfaces (left-hand side) and even genus surfaces (right-hand side).

We remark that for even genus non-orientable surfaces Atalan [1, Proposition 3.1] found the maximal rank of the abelian subgroups which contain groups generated by Dehn twists about pairwise non-isotopic non-separating curves.

Thurston [7] proved that every mapping class $\tau \in \mathcal{M}(N)$ is either reducible or of finite order or pseudo-Anosov, and if τ is reducible, then it has a family \mathcal{A} of isotopy classes of essential simple closed curves such that $\tau(\mathcal{A}) = \mathcal{A}$ and each of the restrictions of τ is of finite order or pseudo-Anosov on each connected component of $N - A$, where A is a set of representatives of \mathcal{A} . We call this theorem Thurston’s theorem. In the theorem, the system A is not unique in general. In Section 2, we will introduce an “essential reduction system” on N in a similar way to Birman-Lubotzky-McCarthy [3]. Birman-Lubotzky-McCarthy proved that the essential reduction system satisfies the condition in Thurston’s theorem and it is a minimal reduction system among such systems and unique up to isotopy for only orientable surfaces. We show the same result for non-orientable surfaces:

Theorem 1.2. *A system A satisfying the conditions of Thurston’s theorem that is minimal among such systems is unique up to isotopy.*

Theorem 1.2 was first proven by Wu [8]. We give another proof by applying the arguments of Birman-Lubotzky-McCarthy.

Because every abelian subgroup G of $\mathcal{M}(N_{g,n})$ is a subgroup of $\mathcal{M}(S_{g-1,2n})$, we see G is finitely generated by [3, Theorem A] immediately. Hence, we obtain the following.

Corollary 1.3. *Let G be an abelian subgroup of $\mathcal{M}(N)$. Then G is finitely generated with torsion-free rank bounded by $\frac{3}{2}(g - 1) + n - 2$ if g is odd and $\frac{3}{2}g + n - 3$ if g is even.*

There are several differences from the case of orientable surfaces in the proofs of the lemmas to prove Theorems 1.1 and 1.2. First difference appears in the proof of (1) in Lemma 2.1. We use the result of Stukow [6] about the kernel of the inclusion homomorphism from the mapping class group of a subsurface to that of the ambient surface for the proof. Secondly, the statement of Lemma 2.6 is different from the orientable surface case ([3, Lemma 2.4]). Thirdly, because we use Lemma 2.6 to prove Lemma 2.7, we have some differences from the orientable surface case in the proof of Lemma 2.7. However, we obtain a similar result to the corresponding lemma ([3, Lemma 2.5]).

The paper is organized as follows. In Section 2 we will introduce essential reduction systems in accordance with orientable surface case by Birman-Lubotzky-McCarthy, and prove Theorem 1.2. In Section 3 we will accomplish the proof of Theorem 1.1.

2. Essential reduction classes

In this section, following Birman-Lubotzky-McCarthy we introduce essential reduction systems for non-orientable surfaces. A compact connected *non-orientable surface* of genus $g \geq 1$ with $n \geq 0$ boundary components is the connected sum of g projective planes with n open disks removed. We denote it by $N = N_{g,n}$. Note that N is homeomorphic to the surface obtained from a sphere by removing $g + n$ open disks and attaching g Möbius bands along their boundaries, and we call each of the Möbius bands the crosscap. A simple closed curve on N is essential if it does not bound a disk or a Möbius band, and is not parallel to a boundary component of N . We often refer to essential simple closed curves as curves. The collection of non-oriented isotopy classes of curves in N is denoted by the symbol $\mathcal{S}(N)$. We remark that admissible isotopies fix each boundary component setwise. If $\tau \in \mathcal{M}(N)$ and $\alpha \in \mathcal{S}(N)$, then $\tau(\alpha)$ denotes the class of $t(a)$, where $t \in \tau$ and $a \in \alpha$. A subset $\mathcal{A} \subset \mathcal{S}(N)$ is *admissible* if a set A of the representatives of \mathcal{A} can be chosen so that it consists of pairwise disjoint curves. Similarly we say that A is an *admissible set of representatives*. Let \mathcal{A} be an admissible subset of $\mathcal{S}(N)$. From now we use some notations which are the same as Birman-Lubotzky-McCarthy [3]. The symbol $\mathcal{M}_{\mathcal{A}}(N)$ denotes the stabilizer of \mathcal{A} in $\mathcal{M}(N)$ which preserves the set \mathcal{A} . We denote by $N_{\mathcal{A}}$ the natural compactification of $N - A$, where A is any admissible set of representatives of \mathcal{A} . If $\tau \in \mathcal{M}_{\mathcal{A}}(N)$, then we can choose an admissible set A of representatives of \mathcal{A} and a representative t of τ such that $t(A) = A$. Furthermore, $t|_{N-A}$ extends uniquely to $N_{\mathcal{A}}$. Note that this process determines a well defined class $\widehat{\tau} \in \mathcal{M}(N_{\mathcal{A}})$. We shall refer to this class $\widehat{\tau}$ as the *reduction* of τ along \mathcal{A} . The assignment $\tau \mapsto \widehat{\tau}$ yields a homomorphism $\Lambda: \mathcal{M}_{\mathcal{A}}(N) \rightarrow \mathcal{M}(N_{\mathcal{A}})$, which we shall refer to as the *reduction homomorphism*. Let a be a two-sided simple closed curve on N . We denote by t_a the *Dehn twist* along a , which is a homeomorphism on N defined by cutting N along a , twisting one side by 2π , and reglueing. Let τ_a be the isotopy class of t_a , where α is an isotopy class of a . Abusing the notation we often call τ_a the Dehn twist along a .

We remark that Λ is not an isomorphism in general, because according to our definition of the mapping class group each Dehn twist τ_a can be in the kernel $\text{Ker}(\Lambda)$ of Λ . A natural representation $\partial: \mathcal{M}(N) \rightarrow \text{Aut}(\partial N)$ arises from the permutation of boundary components. Let \mathcal{A}^{two} and \mathcal{A}^{one} be the subsets of \mathcal{A} which consist of all isotopy classes of two-sided curves and one-sided curves respectively. If we write $\text{card}(\mathcal{A})$, then it means the cardinality of \mathcal{A} .

Lemma 2.1. *Let \mathcal{A} be an admissible subset of $\mathcal{S}(N)$. Then the following occur.*

- (1) $\text{Ker}(\Lambda) = \langle \tau_\alpha \mid \tau_\alpha \in \mathcal{A}^{\text{two}} \rangle$.
- (2) $\text{Ker}(\Lambda) \subset \text{center}(\text{Ker}(\partial \circ \Lambda))$.

Proof of Lemma 2.1. We will prove only (1), since the proof of (2) is the same as that of [3, Lemma 2.1 (2)]. Let A be a set of representatives of \mathcal{A} which are pairwise disjoint, and $\widehat{N-A}$ the surface which we cap the components of ∂A in $N-A$ by once punctured disks. We put orientations in the new boundary components of $N_{\mathcal{A}}$. We denote by $\{p_i^+, p_i^-\}$ the pair of points on two new boundary components obtained by cutting N along a two-sided curve and compactifying it naturally, and by q_i a point on a new boundary component obtained by cutting N along a one-sided curve and compactifying it naturally. We define $\text{Homeo}'(N_{\mathcal{A}})$ as the set of homeomorphisms t which satisfy $t(\{p_i^+, p_i^-\}) = \{p_j^+, p_j^-\}$ and $t(q_i) = q_j$, and if the orientation of the boundary component which has p_i^+ is the same as (resp. opposite to) the boundary component which has $t(p_i^+)$, then the orientation of the boundary component which has p_i^- is the same as (resp. opposite to) the boundary component which has $t(p_i^-)$. Set $\mathcal{M}'(N_{\mathcal{A}}) = \pi_0(\text{Homeo}'(N_{\mathcal{A}}))$. We consider the capping homomorphism $\eta_1: \mathcal{M}'(N_{\mathcal{A}}) \rightarrow \mathcal{M}(\widehat{N-A})$ induced by the inclusion $N_{\mathcal{A}} \hookrightarrow \widehat{N-A}$. Note that the Dehn twists along the new boundary components are not isotopic to the identity in $\mathcal{M}'(N_{\mathcal{A}})$. We define a homomorphism $\nu_1: \mathcal{M}(\widehat{N-A}) \rightarrow \mathcal{M}(N_{\mathcal{A}})$ by $\nu_1(\tau) = \tau|_{N_{\mathcal{A}}}$ for any $\tau \in \mathcal{M}(\widehat{N-A})$. Let g be a regluing map from $N_{\mathcal{A}}$ to N . For any $\tau \in \mathcal{M}'(N_{\mathcal{A}})$, there exists $\sigma \in \mathcal{M}_{\mathcal{A}}(N)$ such that $\sigma \circ g = g \circ \tau$. This process determines the homomorphism ν_2 from $\mathcal{M}'(N_{\mathcal{A}})$ to $\mathcal{M}_{\mathcal{A}}(N)$ by $\nu_2(\tau) = \sigma$. Let ν_3 be a restriction of ν_2 to $\text{Ker}(\eta_1)$. Then, we obtain the following commutative diagram.

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \text{Ker}(\eta_1) & \longrightarrow & \mathcal{M}'(N_{\mathcal{A}}) & \xrightarrow{\eta_1} & \mathcal{M}(\widehat{N-A}) \\
 & & \downarrow \nu_3 & \cup & \downarrow \nu_2 & \cup & \cong \downarrow \nu_1 \\
 1 & \longrightarrow & \text{Ker}(\Lambda) & \longrightarrow & \mathcal{M}_{\mathcal{A}}(N) & \xrightarrow{\Lambda} & \mathcal{M}(N_{\mathcal{A}})
 \end{array}$$

The homomorphism $\nu_1: \mathcal{M}(\widehat{N-A}) \rightarrow \mathcal{M}(N_{\mathcal{A}})$ is isomorphism by our definition of the mapping class groups. The homomorphism $\nu_2: \mathcal{M}'(N_{\mathcal{A}}) \rightarrow \mathcal{M}_{\mathcal{A}}(N)$ is surjective. Actually, for any $\tau \in \mathcal{M}_{\mathcal{A}}(N)$ there exists a representative t of τ and an admissible subset A of representatives of \mathcal{A} such that $t(A) = A$ and t maps each two-sided (resp. one-sided) curve to a two-sided (resp. one-sided) curve. Hence $\tau|_{N_{\mathcal{A}}}$ is an element of $\mathcal{M}'(N_{\mathcal{A}})$, and so it is a lift of τ to $\mathcal{M}'(N_{\mathcal{A}})$. The homomorphism $\nu_3: \text{Ker}(\eta_1) \rightarrow \text{Ker}(\Lambda)$ is surjective by the five-lemma. Thus, $\text{Ker}(\Lambda)$ is generated by at most the projections of the generating set of $\text{Ker}(\eta_1)$. By [6, Theorem 3.6], we have $\text{Ker}(\eta_1) = \langle \tau_\alpha \mid \alpha \in \mathcal{A} \rangle \cong \mathbb{Z}^N$, where $N = 2\text{card}(\mathcal{A}^{\text{two}}) + \text{card}(\mathcal{A}^{\text{one}})$. We know that for each regular neighborhood α of a one-sided curve on N , the Dehn twist τ_α along α is not contained in $\text{Ker}(\Lambda)$ since it is a trivial element in $\mathcal{M}(N_{\mathcal{A}})$. Moreover, both two Dehn twists along the boundary components in $N_{\mathcal{A}}$ which come from the same $\alpha \in \mathcal{A}$ as the boundary components of the regular neighborhood of α in N project τ_α in $\text{Ker}(\Lambda)$. Therefore $\text{Ker}(\Lambda) = \langle \tau_\alpha \mid \alpha \in \mathcal{A}^{\text{two}} \rangle$. \square

Let $\Gamma(N_{\mathcal{A}}) = \{N_i \mid i \in I\}$ be the set of the connected components of $N_{\mathcal{A}}$, and so $N_{\mathcal{A}} = \coprod_{i \in I} N_i$. There is a natural representation $\varphi: \mathcal{M}(N_{\mathcal{A}}) \rightarrow \text{Aut}(\partial N_{\mathcal{A}})$ which arises from the permutation of components. The kernel $\text{Ker}(\varphi)$ of φ is isomorphic to $\bigoplus_{i \in I} \mathcal{M}(N_i)$. If

$\tau \in \mathcal{M}(N_{\mathcal{A}})$, then for some exponent n , τ^n is contained in $\text{Ker}(\varphi)$. For any such an exponent, we refer to the element of $\mathcal{M}(N_i)$ obtained by restricting τ^n as *restrictions* of τ .

Let S be the double covering orientable surface of N . Wu [8] proved that a mapping class $\tau \in \mathcal{M}(N)$ is of *finite order* (resp. *reducible*, *pseudo-Anosov*) if $\iota(\tau) \in \mathcal{M}(S)$ is of *finite order* (resp. *reducible*, *pseudo-Anosov*). Moreover according to Thurston [7], there is Nielsen-Thurston classification for $\mathcal{M}(N)$ (the proof is found in the paper of Wu [8]):

Theorem 2.2 ([7], [8, Theorem 2]). *A homeomorphism t on N is reduced and is not of finite order if and only if t is isotopic to a pseudo-Anosov homeomorphism.*

Then we can rephrase the definition of pseudo-Anosov and reducible mapping classes on non-orientable surfaces according to Birman-Lubotzky-McCarthy [3] as follows:

DEFINITION 2.3. A mapping class $\tau \in \mathcal{M}(N_{\mathcal{A}})$ is *pseudo-Anosov* if $\mathcal{S}(N_i) \neq \emptyset$ for every $i \in I$ and $\tau^n(\alpha) \neq \alpha$ for any $\alpha \in \mathcal{S}(N_{\mathcal{A}})$ and any $n \neq 0$. A mapping class $\tau \in \mathcal{M}(N_{\mathcal{A}})$ is *reducible* if there exists an admissible subset \mathcal{A} such that $\tau(\mathcal{A}) = \mathcal{A}$.

We call the admissible set \mathcal{A} as in Definition 2.3 a *reduction system* for τ , and we say each $\alpha \in \mathcal{A}$ as a *reduction class* for τ . A reduction system \mathcal{A} for τ is an *adequate reduction system* if the restrictions of τ to each component of $N_{\mathcal{A}}$ are either of finite order or pseudo-Anosov. If $\tau \in \mathcal{M}(N)$ is either of finite order or pseudo-Anosov mapping class on N , we call τ *adequately reduced*.

We restate Thurston's theorem by using adequate reduction system as follows:

Theorem 2.4 ([7]). *Every mapping class $\tau \in \mathcal{M}(N)$ is either reducible or adequately reduced. If τ is reducible, then there exists an adequate reduction system \mathcal{A} for τ .*

Let $i: \mathcal{S}(N) \times \mathcal{S}(N) \rightarrow \mathbb{N} \cup \{0\}$ be the geometric intersection number. A reduction class α for τ is *essential* if for each $\beta \in \mathcal{S}(N)$ such that $i(\alpha, \beta) \neq 0$ and each integer $m \neq 0$, the class $\tau^m(\beta)$ is distinct from β . We often say α is essential if α is an essential reduction system for some τ .

Proposition 2.5 ([3, Proposition 2.3]). *Let $\alpha \in \mathcal{A}$ and $\alpha' \in \mathcal{A}'$ be reduction classes for $\tau \in \mathcal{M}(N)$. Suppose that α is essential. Then $i(\alpha, \alpha') = 0$.*

The following lemma is different from the corresponding lemma ([3, Lemma 2.4]) by Birman-Lubotzky-McCarthy.

Lemma 2.6. *Let F be a compact connected orientable or non-orientable surface with $\chi(F) < 0$. Fix any isotopy class δ of properly embedded arc d on F , namely, ∂d is embedded in ∂F and the interior of d is embedded in the interior of F . We choose any $\tau \in \mathcal{M}(F)$ with $\tau(\delta) = \delta$. Then one of the following occurs.*

- (1) F is either $S_{0,3}$ or $N_{1,2}$ or $N_{2,1}$.
- (2) If δ is an isotopy class of an arc which connects distinct two boundary components, then there exists $\gamma \in \mathcal{S}(F)$ such that $\tau(\gamma) = \gamma$ and $i(\alpha, \gamma) \neq 0$ for any $\alpha \in \mathcal{S}(F)$ with $i(\alpha, \delta) \neq 0$.
- (3) If δ is an isotopy class of an arc which connects one boundary component, goes through crosscaps even number of times, and surrounds one crosscap, then for any $\alpha \in \mathcal{S}(F)$ excepting β_0 which is shown in Figure 2 with $i(\alpha, \delta) \neq 0$ there exists $\gamma \in \mathcal{S}(F)$ such that $\tau(\gamma) = \gamma$ and $i(\alpha, \gamma) \neq 0$.

- (4) If δ is an isotopy class of an arc which connects one boundary component, goes through crosscaps even number of times, and does not surround one crosscap, then for any $\alpha \in \mathcal{S}(F)$ with $i(\alpha, \delta) \neq 0$ there exists $\gamma \in \mathcal{S}(F)$ such that $\tau(\gamma) = \gamma$ and $i(\alpha, \gamma) \neq 0$.
- (5) If δ is an isotopy class of an arc which connects one boundary component and goes through crosscaps odd number of times, then for any $\alpha \in \mathcal{S}(F)$ excepting β_1 and β_2 which are shown in Figure 2 with $i(\alpha, \delta) \neq 0$ there exists $\gamma \in \mathcal{S}(F)$ such that $\tau(\gamma) = \gamma$ and $i(\alpha, \gamma) \neq 0$.

Proof. Let d be a properly embedded representative of δ , and $\eta(d)$ a regular neighborhood of d with the boundary components of F which have the end points of d . If d connects distinct two boundary components (Case (a)) or connects one boundary component and goes through crosscaps even number of times (Case (b)), then $\eta(d)$ is homeomorphic to $S_{0,3}$. If d connects one boundary component and goes through crosscaps odd number of times (Case (c)), then $\eta(d)$ is homeomorphic to $N_{1,2}$. In Case (a), two of the boundary components of $\eta(d)$ are those of F . We denote by γ the isotopy class of the other boundary component of $\eta(d)$. Then γ is not isotopic to a point since $\chi(F) < 0$. If γ is isotopic to a component of ∂F (resp. a crosscap), then F is homeomorphic to $S_{0,3}$ (resp. $N_{1,2}$). We suppose that γ is an essential curve. Since $\tau(\delta) = \delta$, it follows that $\tau(\gamma) = \gamma$. If $\alpha \in \mathcal{S}(F)$ intersects δ nontrivially, then $i(\alpha, \gamma) \neq 0$. In Case (b), one of the boundary component of $\eta(d)$ is that of F . We put γ_2 and γ_3 as the isotopy classes of the other two components. Neither γ_2 nor γ_3 is isotopic to a point. Firstly we suppose that γ_2 is parallel to ∂F . If γ_3 is isotopic to a component of ∂F (resp. a crosscap), then F is homeomorphic to $S_{0,3}$ (resp. $N_{1,2}$). We suppose that γ_3 is an essential curve. Then it follows that $\tau(\gamma_3) = \gamma_3$ and $i(\alpha, \gamma_3) \neq 0$ for any $\alpha \in \mathcal{S}(F)$ with $i(\alpha, \delta) \neq 0$. Next we suppose that γ_2 is isotopic to a crosscap. If γ_3 is isotopic to a component of ∂F (resp. a crosscap), then F is homeomorphic to $N_{1,2}$ (resp. $N_{2,1}$). We suppose that γ_3 is an essential curve. Unless α is isotopic β_0 in Figure 2, we see $i(\alpha, \gamma_3) \neq 0$ for any $\alpha \in \mathcal{S}(F)$ with $i(\alpha, \delta) \neq 0$. We can show similar results if γ_2 is essential. In Case (c), one of the boundary component of $\eta(d)$ is that of F . We denote by γ the isotopy class of the other boundary component of $\eta(d)$. We see γ is not isotopic to a point. If γ is isotopic to a component of ∂F (resp. a crosscap), then F is homeomorphic to $N_{1,2}$ (resp. $N_{2,1}$). We suppose that γ is an essential curve. Since $\tau(\delta) = \delta$, it follows that $\tau(\gamma) = \gamma$. Note that γ bounds $N_{1,1}$. We can take only β_1 and β_2 which intersect δ and do not intersect γ as shown in Figure 2. Unless α is isotopic to β_1 or β_2 in Figure 2, we see $i(\alpha, \gamma) \neq 0$ for any $\alpha \in \mathcal{S}(F)$ with $i(\alpha, \delta) \neq 0$. \square

The result of the following lemma is the same as the orientable surface case by Birman-Lubotzky-McCarthy ([3, Lemma 2.5]), while the statement of Lemma 2.6 is different from the orientable surface case by them ([3, Lemma 2.4]).

Lemma 2.7. *Let \mathcal{A} be an adequate reduction system for τ and let $\alpha \in \mathcal{A}$. Set $\mathcal{A}' = \mathcal{A} - \{\alpha\}$. Then the following are equivalent:*

- (1) α is essential.
- (2) \mathcal{A}' is not an adequate reduction system for τ^m for any $m \neq 0$.




Fig. 2. Curves $\beta_0, \beta_1,$ and β_2 .

Proof. We omit the proof that (1) implies (2) because it is the same as that of Birman-Lubotzky-McCarthy. We assume that $\alpha \in \mathcal{A}$ is not an essential reduction class for τ . Then there exists a class $\gamma \in \mathcal{S}(N)$ with $i(\alpha, \gamma) \neq 0$ and $n \neq 0$ such that $\tau^n(\gamma) = \gamma$. Let A be an admissible set of the representatives of \mathcal{A} . We cut N along A , then γ determines a finite set of pairwise disjoint isotopy classes of properly embedded arcs in $N_{\mathcal{A}}$, which we denote by $\widehat{\gamma}$. If a component of $N_{\mathcal{A}}$ has boundary components arising from α , then we shall say the component is bounded by α . We note that at least one element of $\widehat{\gamma}$ occurs in a component of $N_{\mathcal{A}}$ bounded by α . We put $\Lambda: \mathcal{M}_{\mathcal{A}}(N) \rightarrow \mathcal{M}(N_{\mathcal{A}})$, and set $\widehat{\tau} = \Lambda(\tau)$. Since $\tau^n(\gamma) = \gamma$, therefore $\widehat{\tau}^n(\widehat{\gamma}) = \widehat{\gamma}$. By choosing a larger exponent n if necessary, we may assume that $\widehat{\tau}^n$ preserves each component of $N_{\mathcal{A}}$, $\partial N_{\mathcal{A}}$, and $\widehat{\gamma}$. In particular, the restrictions of $\widehat{\tau}^n$ to the components of $N_{\mathcal{A}}$ bounded by α preserve a nontrivial isotopy class of a properly embedded arc. By Lemma 2.6, for each such component, either the corresponding restriction of $\widehat{\tau}^n$ is reducible or the component is $S_{0,3}$, $N_{1,2}$, or $N_{2,1}$. Now we suppose that $\widehat{\tau}^n$ is adequately reduced, and therefore all the restrictions of $\widehat{\tau}^n$ do not preserve any curves. Then it follows that each component bounded by α is either $S_{0,3}$ or $N_{1,2}$ or $N_{2,1}$. A pair of pants will not support a pseudo-Anosov mapping class. We can show that the mapping classes on $N_{1,2}$ (resp. $N_{2,1}$) which preserve a properly embedded essential arc on it and each component of $\partial N_{1,2}$ (resp. $\partial N_{2,1}$) are of finite order. Thus by choosing a larger exponent n if necessary, we may assume that the restrictions of $\widehat{\tau}^n$ to the component bounded α are trivial.

From now we consider the corresponding situation when we reduce along \mathcal{A}' . We define the reduction homomorphism $\Lambda': \mathcal{M}_{\mathcal{A}'}(N) \rightarrow \mathcal{M}(N_{\mathcal{A}'})$. Let $\widehat{\gamma}'$ and $\widehat{\alpha}'$ be the lift of γ and α on $N_{\mathcal{A}'}$, respectively. We see $\widehat{\alpha}'$ is an adequate reduction class for $\Lambda'(\tau^n)$. We denote by L the component of $N_{\mathcal{A}'}$ which includes $\widehat{\alpha}'$. We set $\nu = \Lambda'(\tau^n)|_L$. We also define the reduction homomorphism $\Lambda'': \mathcal{M}_{\widehat{\alpha}'}(L) \rightarrow \mathcal{M}(L_{\widehat{\alpha}'})$, and set $\nu'' = \Lambda''(\nu)$. Because the restriction of $\widehat{\tau}^n$ to the component of $N_{\mathcal{A}}$ bounded by α is the identity, ν'' is the identity. If $\widehat{\alpha}'$ is an isotopy class of one-sided curve, then ν is also identity by Lemma 2.1. If $\widehat{\alpha}'$ is an isotopy class of two-sided curve, then $\text{Ker}(\Lambda'') = \langle \tau_{\widehat{\alpha}'} \rangle$ by Lemma 2.1, and so there exists $k \neq 0$ such that $\nu = \tau_{\widehat{\alpha}'}^k$. We will prove that $k = 0$ from now. First we assume that γ intersects only α , that is, $\gamma \subset N - A'$ (A' is an admissible set of the representatives of \mathcal{A}'). We have $i(\widehat{\alpha}', \widehat{\gamma}') \neq 0$, so $\widehat{\gamma}' \in \mathcal{S}(L)$. Since $\tau^n(\gamma) = \gamma$, it follows that $\nu(\widehat{\gamma}') = \widehat{\gamma}'$. Then we see k should be 0. Next we assume that γ intersects other curves of \mathcal{A}' , that is, $i(\gamma, \mathcal{A}') \neq 0$. In this case $\widehat{\gamma}'$ is the family consisting of isotopy classes of arcs which go through L at least once. If $L = S_{0,3}$, then ν is the identity by our assumption. If $L = N_{1,2}$, then k has to be 0 because $\mathcal{M}(N_{1,2})$ is finite. If $L = N_{2,1}$, then there is only one kind of two-sided curve on L shown in Figure 3. By Lemma 2.6, $\widehat{\alpha}'$ can not be $\beta_0, \beta_1,$ or β_2 because $\widehat{\alpha}'$ is an isotopy class of two-sided curve. Then there exists $\delta \in \mathcal{S}(L)$ such that $i(\widehat{\alpha}', \delta) \neq 0$ and $\nu(\delta) = \delta$. Hence $k = 0$. Otherwise, similarly to the previous case $\widehat{\alpha}'$ can not be $\beta_0, \beta_1,$ or β_2 , and so there exists $\delta \in \mathcal{S}(L)$ such

that $i(\widehat{\alpha}', \delta) \neq 0$ and $v(\delta) = \delta$. Hence we obtain $k = 0$. From the above arguments, we prove \mathcal{A}' is an adequate reduction system for τ . \square

We set $\mathcal{A}_\tau = \{\alpha \in \mathcal{S}(N) \mid \alpha \text{ is an essential reduction class for } \tau\}$. Since we obtain the same result in Lemma 2.7 as [3, Lemma 2.5], we can prove the following lemma by similar arguments to that of Birman-Lubotzky-McCarthy. Hence, we omit the proof of Lemma 2.8 (we use Lemma 2.7 in the proofs of (3) and (4) in Lemma 2.8).

Lemma 2.8 ([3, Lemma 2.6]).

- (1) $\sigma(\mathcal{A}_\tau) = \mathcal{A}_{\sigma\tau\sigma^{-1}}$.
- (2) $\mathcal{A}_{\tau^m} = \mathcal{A}_\tau$ for all $m \neq 0$.
- (3) \mathcal{A}_τ is an adequate reduction system for τ .
- (4) $\mathcal{A}_\tau \subset \mathcal{A}$ for each adequate reduction system \mathcal{A} for τ .

At the end of this section, we prove Theorem 1.2. The proof is the same as the proof of [3, Theorem C].

Proof of Theorem 1.2. Let $\tau \in \mathcal{M}(N)$. Then, by Theorem 2.4 either τ is adequately reduced (that is the case $\mathcal{A} = \emptyset$) or τ is reducible, and if τ is reducible, then there exists an adequate reduction system. By Lemma 2.8, \mathcal{A}_τ is the intersection of all adequate reduction systems for τ . Hence \mathcal{A}_τ is canonical and unique. The desired curve system A is any representative of \mathcal{A}_τ . \square




Fig.3. The only two-sided curve $\widehat{\alpha}'$ on $N_{2,1}$.

3. Abelian subgroups of mapping class groups

In this section we prove Theorem 1.1. Let N be a compact connected non-orientable surface, and $\mathcal{A} \in \mathcal{S}(N)$ an admissible subset. Then $G < \mathcal{M}(N_{\mathcal{A}})$ is *adequately reduced* if each $\tau \in G$ is adequately reduced. Let G be an abelian subgroup of $\mathcal{M}(N_{\mathcal{A}})$. We denote by $\text{rank}(G)$ the torsion-free rank of G .

Lemma 3.1 ([3, Lemma 3.1]).

- (1) Let \mathcal{A}_G be the union of the essential reduction systems \mathcal{A}_τ for any $\tau \in G$. Then \mathcal{A}_G is an adequate reduction system for each $\tau \in G$.
- (2) If G is adequately reduced, then $\text{rank}(G) \leq C_0(N_{\mathcal{A}})$, where $C_0(N_{\mathcal{A}})$ is the number of components of $N_{\mathcal{A}}$ not homeomorphic to a pair of pants.

Proof. We can show (1) by a similar argument to Birman-Lubotzky-McCarthy. Therefore we only give the proof of (2). We set $N_{\mathcal{A}} = \coprod_{i=1}^k N_i$, where each N_i is a connected component of $N_{\mathcal{A}}$. If N_i is a non-orientable surface, then let $p_i: S_i \rightarrow N_i$ be the double covering map of

N_i , where S_i is the double covering orientable surface of N_i . If N_i is an orientable surface, then let $p_i: S_i \rightarrow N_i$ be the identity map on N_i . We define $p = \prod_{i=1}^k p_i: \prod_{i=1}^k S_i \rightarrow \prod_{i=1}^k N_i$ by $p(x) = p_i(x)$ for $x \in S_i$, then p is a covering map of $N_{\mathcal{A}}$. Let $\iota: \mathcal{M}(N_{\mathcal{A}}) \rightarrow \mathcal{M}(\prod_{i=1}^k S_i)$ be an injective homomorphism induced by p . We choose an adequately reduced abelian subgroup $G < \mathcal{M}(N_{\mathcal{A}}) = \mathcal{M}(\prod_{i=1}^k N_i)$. We set $\tilde{G} = \iota(G) < \mathcal{M}(\prod_{i=1}^k S_i)$. Then, \tilde{G} is an adequately reduced abelian subgroup, since $\tau \in \mathcal{M}(N_i)$ is finite order (resp. pseudo-Anosov) mapping class if and only if $\tilde{\tau} \in \mathcal{M}(S_i)$ is finite order (resp. pseudo-Anosov) mapping class. By [3, Lemma 3.1], we know $\text{rank}(\tilde{G}) \leq C_0(\prod_{i=1}^k S_i)$, where $C_0(\prod_{i=1}^k S_i)$ is the number of components of $\prod_{i=1}^k S_i$ not homeomorphic to a pair of pants. If N_i is a non-orientable surface, then S_i can not be a pair of pants, because the Euler characteristic of S_i should be even but that of a pair of pants is -1 . If N_i is an orientable surface, then N_i is a pair of pants if and only if S_i is a pair of pants. Hence $C_0(\prod_{i=1}^k S_i) = C_0(N_{\mathcal{A}})$. By $\tilde{G} \cong G$, it follows that $\text{rank}(G) \leq C_0(N_{\mathcal{A}})$. \square

Finally, we give the proof of Theorem 1.1.

Proof of Theorem 1.1. Let G be a torsion-free abelian subgroup of $\mathcal{M}(N)$. By (1) of Lemma 3.1, we see $G \in \mathcal{M}_{\mathcal{A}_G}(N)$. We denote by $\Lambda: \mathcal{M}_{\mathcal{A}_G}(N) \rightarrow \mathcal{M}(N_{\mathcal{A}_G})$ the reduction homomorphism along \mathcal{A}_G . We set $H = \Lambda(G)$. The sequence

$$1 \longrightarrow G \cap \text{Ker}(\Lambda) \longrightarrow G \longrightarrow H \longrightarrow 1$$

is exact. From this exact sequence, we have $\text{rank}(G) = \text{rank}(G \cap \text{Ker}(\Lambda)) + \text{rank}(H)$. By Lemma 2.1, $\text{rank}(G \cap \text{Ker}(\Lambda)) \leq \text{rank}(\text{Ker}(\Lambda)) \leq \text{card}(\mathcal{A}^{\text{two}})$. Therefore, we see $G \cap \text{Ker}(\Lambda)$ is isomorphic to the group generated by Dehn twists along pairwise disjoint curves whose cardinality is at most $\text{card}(\mathcal{A}^{\text{two}})$.

From (2) of Lemma 3.1, we see $\text{rank}(H) \leq C_0(N_{\mathcal{A}_G})$. By the arguments in the proof of [3, Lemma 3.1 (2)], we see H is isomorphic to the group generated by pseudo-Anosov mapping classes on connected subsurfaces, and the number is bounded by $C_0(N_{\mathcal{A}_G})$. However, we remark that any subsurface which supports a pseudo-Anosov mapping class must also support a non-trivial Dehn twist. Hence we can replace the pseudo-Anosov mapping class generators by Dehn twists. As shown in Figure 1 (see also [2, Proposition 2.3]), $\text{card}(\mathcal{A}^{\text{two}})$ is bounded by $\frac{3}{2}(g-1) + n - 2$ if g is odd and $\frac{3}{2}g + n - 3$ if g is even, so we are done. \square

ACKNOWLEDGEMENTS. The author is extremely grateful to Hisaaki Endo for his warm encouragement and helpful advice. She also wishes to thank Błażej Szepietowski. He told her some properties of the mapping class group of non-orientable surface, gave her a lot of valuable comments, and she can improve Theorem 1.1. She also thanks Genki Omori for the useful discussion with her particularly about Lemma 2.1. This work was supported by JSPS KAKENHI, the grant number 16J00397 of Research Fellowship for Young Scientists.

References

- [1] F. Atalan: *An algebraic characterization of a Dehn twist for nonorientable surfaces*, arXiv:1501.07183v2 [math.GT].
- [2] F. Atalan and M. Korkmaz: *Automorphisms of curve complexes on nonorientable surfaces*, Groups Geom. Dyn. **8** (2014), 39–68.
- [3] J.S. Birman, A. Lubotzky, and J. McCarthy: *Abelian and solvable subgroups of the mapping class groups*, Duke Math. J. **50** (1983), 1107–1120.
- [4] S. Kim and T. Koberda: *Right-angled Artin groups and finite subgraphs of curve graphs*, Osaka J. Math. **53**, 705–716.
- [5] B. Szepietowski: *Embedding the braid group in mapping class groups*, Publ. Mat. **54** (2010), 359–368.
- [6] M. Stukow: *Commensurability of geometric subgroups of mapping class groups*, Geom. Dedicata **143** (2009), 117–142.
- [7] W. Thurston: *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Amer. Math. Soc. (N.S.) **19** (1988), 417–431.
- [8] Y. Wu: *Canonical reducing curves of surface homeomorphism*, Acta Math. Sinica (N.S.) **3** (1987), 305–313.

Department of Mathematics
Graduate School of Science, Osaka University
1–1 Machikaneyama-cho, Toyonaka, Osaka 560–0043
Japan
e-mail: e-kuno@math.sci.osaka-u.ac.jp