

# ABI3 and PIL5 Collaboratively Activate the Expression of *SOMNUS* by Directly Binding to Its Promoter in Imbibed *Arabidopsis* Seeds <sup>W</sup>

Jeongmo Park,<sup>1</sup> Nayoung Lee,<sup>1</sup> Woohyun Kim, Soohwan Lim, and Giltso Choi<sup>2</sup>

Department of Biological Sciences, KAIST, Daejeon 305-701, Korea

A previous study showed that *SOMNUS* (*SOM*), which encodes a C3H-type zinc finger protein, is a key negative regulator of seed germination that acts downstream of PHYTOCHROME INTERACTING FACTOR3-LIKE5 (*PIL5*). However, it was not determined if *PIL5* is the sole regulator of *SOM* expression. Public microarray data suggest that the expression of *SOM* mRNA is regulated also by ABSCISIC ACID INSENSITIVE3 (*ABI3*), another key regulator of seed germination. By analyzing *abi3* mutants and *ABI3* overexpression lines, we show here that *ABI3* activates the expression of *SOM* mRNA collaboratively with *PIL5* in imbibed seeds. Chromatin immunoprecipitation analysis coupled with electrophoretic mobility shift assay indicate that *ABI3* activates the expression of *SOM* mRNA by directly binding to two RY motifs present in the *SOM* promoter *in vivo*, which is further supported by the greatly decreased expression of a reporter gene driven by a *SOM* promoter bearing mutated RY motifs. At the protein level, the *ABI3* protein interacts with the *PIL5* protein. The *ABI3*-*PIL5* interaction, however, does not affect targeting of *ABI3* and *PIL5* to *SOM* promoters. Taken together, our results indicate that *ABI3* and *PIL5* collaboratively activate the expression of *SOM* mRNA by directly binding to and interacting with each other at the *SOM* promoter.

## INTRODUCTION

The decision for a seed to germinate at a given time and space is determined by seed developmental status and environmental conditions. In *Arabidopsis thaliana*, freshly harvested seeds display seed dormancy, a property that inhibits germination even in favorable environmental conditions (Finch-Savage and Leubner-Metzger, 2006). Abscisic acid (ABA) plays important roles in both initiating and maintaining seed dormancy, as shown by the shallow seed dormancy of ABA synthetic mutants, such as *aba2*, and the deep seed dormancy of ABA catabolic mutants, such as a *cyp707a* triple mutant (Léon-Kloosterziel et al., 1996; Okamoto et al., 2010). Mutations in some of the ABA signaling genes, such as *ABSCISIC ACID INSENSITIVE3* (*ABI3*), which encodes a DNA binding protein that acts as a positive component of ABA signaling, also disrupts seed dormancy (Lopez-Molina et al., 2002). Other components, such as *DELAY OF GERMINATION1* and *HISTONE MONOUBIQUITINATION1*, have been identified as being important regulators of seed dormancy, but their relationship with ABA signaling has not been determined (Bentsink et al., 2006; Liu et al., 2007).

Freshly harvested seeds eventually lose dormancy when they are sufficiently dried or have undergone stratification. It is not

clear how seed dormancy breaks, but, once broken, nondormant seeds germinate if environmental conditions are favorable. Among various environmental factors, light and temperature play important roles in the decision to germinate. A seed must monitor and integrate various environmental conditions into cellular processes in order for germination to occur. Ultimately, favorable conditions activate germination-promoting hormone signaling, such as gibberellic acid (GA) signaling, and repress germination-inhibiting hormone signaling, such as ABA signaling. The molecular pathways involved in monitoring environmental conditions and integrating them with hormonal signaling are currently being actively investigated (Finkelstein et al., 2008; Holdsworth et al., 2008; Seo et al., 2009).

Phytochrome-mediated light signaling provides a good model system of how environmental conditions are perceived and integrated into internal cellular processes. Inside seed cells, phytochromes perceive red and far-red spectra of light. In *Arabidopsis*, phytochrome B (phyB), which is the major phytochrome present in dry seeds, perceives red light and promotes seed germination. Phytochrome A (phyA), which accumulates during seed imbibition, also perceives light and promotes seed germination. Unlike phyB, however, phyA perceives both very low fluences of all spectra of visible light and prolonged far-red light (Shinomura et al., 1996). Other minor phytochromes, such as phyE, also play a role in promoting seed germination in imbibed seeds (Hennig et al., 2002). When imbibed seeds perceive light, activated phytochromes enter the nucleus and transmit the light signal, partly by destabilizing a germination-inhibiting, phytochrome-interacting bHLH transcription factor, called *PIL5* (also known as *PIF1*) (Oh et al., 2004, 2006). The central role of *PIL5* in mediating phytochrome signaling can be

<sup>1</sup> These authors contributed equally to this work.

<sup>2</sup> Address correspondence to gchoi@kaist.edu.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Giltso Choi (gchoi@kaist.edu).

<sup>W</sup>Online version contains Web-only data.

www.plantcell.org/cgi/doi/10.1105/tpc.110.080721

inferred from the 100% germination frequency of the *pil5* mutant, even if phytochrome B is inactivated by far-red light. In addition, microarray analysis indicates that all genes in imbibed seeds that are differentially expressed in response to red light are either directly or indirectly regulated by PIL5 (Oh et al., 2009). Thus, the destabilization of PIL5 by activated phytochromes serves to release the repression of seed germination imposed by PIL5 and allows seeds to germinate.

PIL5 inhibits seed germination both by coordinating various hormone signaling cascades and by inhibiting cell wall loosening in imbibed seeds (Oh et al., 2009). Chromatin immunoprecipitation analysis coupled with microarray analysis shows that PIL5 regulates the expression of 166 genes by directly binding to their promoters. These direct target genes include cell wall loosening genes, such as expansin and xyloglucan endotransglycosylase genes, and various hormone signaling genes, such as GA signaling genes (*GA INSENSITIVE [GAI]* and *REPRESSOR OF GA [RGA]*), ABA signaling genes (*ABI5* and *ABI3*), auxin signaling genes (*AUXIN RESPONSE FACTOR18 [ARF18]* and *INDOLE ACETIC ACID-INDUCED PROTEIN16 [IAA16]*), cytokinin signaling genes (*CYTOKININ RESPONSE FACTOR1 [CRF1]*, *CRF2*, and *CRF3*), a JA signaling gene (*JAZ1*), and a BR signaling gene (*BES1-INTERACTING MYC-LIKE PROTEIN2*). The direction of expression indicates that PIL5 inhibits seed germination partly by activating the expression of *GAI* and *RGA*, which inhibit seed germination as negative GA signaling components, and of *ABI3* and *ABI5*, which also inhibit seed germination as positive ABA signaling components. The role of other signaling components in seed germination is not clearly defined. However, the repression of *JAZ1* and *IAA16*, negative signaling components of JA and auxin signaling, respectively, and the activation of *ARF8*, a positive auxin signaling component, are also consistent with the inhibitory role of PIL5 in seed germination. PIL5 also modulates hormone signaling by regulating the metabolism of various hormones. For example, PIL5 decreases the level of GA by repressing GA synthetic genes (*GA3ox1* and *GA3ox2*) and activating a GA catabolic gene (*GA2ox1*), while it increases the ABA level by activating ABA synthetic genes (*ABA DEFICIENT1 [ABA1]*, *NINE-CIS-EPOXYCAROTENOID DIOXYGENASE6 [NCED6]*, and *NCED9*) and repressing an ABA catabolic gene (*CYP707A2*). The reciprocal regulation of hormone synthetic and catabolic genes can also be seen in auxin metabolism, as PIL5 activates the expression of genes involved in auxin synthesis (*NITRILASE1 [NIT1]*, *NIT3*, *ALDEHYDE OXIDASE1*, and *AMIDASE1*) but represses the expression of a gene involved in auxin catabolism (*DWARF IN LIGHT1*). Curiously, PIL5 does not bind to the promoters of most genes involved in hormone metabolism, suggesting that PIL5 indirectly regulates these metabolic genes through other direct target genes. However, the degree of separation between PIL5 and genes involved in hormone metabolism is not known. Furthermore, genetic networks operating under PIL5 to regulate seed germination have not yet been identified. An analysis of the genes directly targeted by PIL5 might be useful to identify these genetic networks.

As a positive ABA signaling component, *ABI3* also regulates various aspects of seed development (Giraudat et al., 1992). It belongs to a plant-specific B3 domain protein family that includes *ABI3*, *FUSCA3 (FUS3)*, and *LEAFY COTYLEDON2 (LEC2)*

in *Arabidopsis* and *VIVIPAROUS1 (VP1)* in maize (*Zea mays*). *ABI3* contains four domains, dubbed A, B1, B2, and B3, which are named after the activation domain and the three basic amino acid clusters contained in it (Bies-Etheve et al., 1999). Among them, the B3 domain is known to bind to the RY motif (CATGCA) in vitro (Suzuki et al., 1997; Mönke et al., 2004; Braybrook et al., 2006), while the B1 and B2 domains interact with a set of bZIP transcription factors, including *ABI5*, *bZIP10*, *bZIP25*, and *TRAB1* (Hobo et al., 1999; Nakamura et al., 2001; Lara et al., 2003), that bind to either the G-box element (CACGTG) or ABA response elements (ABREs) (Choi et al., 2000). Consistent with its binding to the RY motif and its interaction with bZIP factors, promoters of many *ABI3*-regulated genes contain RY motifs coupled with a G-box or ABREs. Upon binding to these promoters either directly or indirectly, *ABI3* regulates the expression of genes involved in seed dormancy, seed germination, desiccation tolerance, the accumulation of seed storage proteins, and the breakdown of chlorophyll molecules in developing seeds. The genetic networks operating under *ABI3* to regulate these processes are not fully understood.


We previously showed that *SOMNUS (SOM)* is a key direct target of PIL5 that inhibits seed germination downstream of PIL5 (Kim et al., 2008). *SOM* encodes a C3H-type zinc finger protein of unknown function. The expression analysis suggests that *SOM* inhibits seed germination partly by activating the expression of ABA synthetic genes and by inhibiting the expression of GA synthetic genes downstream of PIL5 but not by activating the expression of *GAI* and *RGA* in imbibed seeds. In this study, we further investigated how the expression of *SOM* is regulated by *ABI3* and PIL5. We found that *ABI3* directly binds to RY motifs present in the *SOM* promoter and activates the expression of *SOM* independently of PIL5 in maturing seeds but collaboratively with PIL5 in imbibed seeds. The interaction between *ABI3* and PIL5 at the protein level further indicates that these two key regulators form a complex that regulates the expression of *SOM* in imbibed seeds.

## RESULTS

### *ABI3* Regulates the Expression of *SOM*

*SOM* inhibits seed germination by regulating genes downstream of PIL5 that are involved in both GA and ABA metabolism in imbibed seeds (Oh et al., 2007). We further investigated how the expression of *SOM* is regulated. The public microarray data compiled using the BAR HeatMapper tool suggested that *SOM* is a seed-specific gene whose expression increases during seed maturation (Figure 1A). The timing of *SOM* expression is slightly later than that of *ABI3* or *FUS3* but similar to that of *ABI5* or *Em1* during seed maturation. Unlike *ABI3* and *ABI5*, *PIL5* is expressed at a low level throughout seed maturation (Figure 1A).

Experimental analysis further supports that *SOM* is a seed-specific gene. We determined the expression levels of *SOM* in various tissues. *SOM* was expressed at very low levels in leaf, root, stem, and flower tissues but at high levels in seeds. During seed maturation, *SOM* began to be detected in siliques at 9 d postanthesis (DPA), and its expression increased beyond this



**Figure 1.** *SOM* Is Highly Expressed during Seed Maturation.

**(A)** Public microarray data showing the seed-specific expression of *SOM*. The expression levels are visualized by BAR HeatMapper. Numbers beneath the heat map indicate the relative expression levels, and the higher expression levels are indicated by more reddish colors. Ros., rosette; Caul., cauline; Aging, aging rosette. Hours/Wc, imbibition hours in continuous white light.

**(B)** Seed-specific expression of *SOM* during seed maturation. The *SOM* mRNA expression levels were quantified by quantitative RT-PCR and are indicated as relative expression levels compared with *PP2A* mRNA ( $SOM/PP2A$ ) ( $sd, n = 3$ ). An inset indicates the relative expression levels of *SOM* mRNA in seeds and other tissues (valve and replum) that are separated from the 12-DPA silique.

**(C)** Relative expression of *Em1* and *ABI5* mRNAs during seed maturation ( $sd, n = 3$ ).


**(D)** A diagram of the *SOM* promoter showing the RY motifs. G-box (CACGTG), PIL5-associated E-box (CANNTG), and ABRE (CACGTA) are also indicated.

time point (Figure 1B). The expression pattern of *SOM* during seed maturation was similar to that of *ABI5* and *Em1*, two seed-specific genes that are known to be regulated by *ABI3* (Figure 1C). When siliques were separated into developing seeds and other parts (valve and replum), *SOM* was detected only in the developing seeds but not in the valve and replum (Figure 1B, inset), further supporting the seed-specific expression of *SOM*.

We analyzed the upstream region of *SOM* and noticed that there were three RY motifs, in addition to other motifs, such as an ABRE and E-boxes, in the 2.7-kb intergenic region ranging from the transcription start site of *SOM* to its upstream neighboring gene (*At1g03780*) (Figure 1D). The three RY motifs are 0.3, 1.4, and 2.4 kb away from the transcription start site of *SOM*. Since the RY motif is bound by *ABI3* or its homologs, the presence of RY motifs in its promoter suggests that *SOM* is regulated by *ABI3* or related B3-domain proteins during seed maturation.

To experimentally test whether *ABI3* regulates the expression of *SOM* during seed maturation, we determined the *SOM* transcript levels in an *abi3* mutant. Since *PIL5* regulates the expression of *SOM* in imbibed seeds, we first tested if *PIL5* also regulates the expression of *SOM* during seed maturation. As in the wild type, however, the expression level of *SOM* increased during seed maturation and reached a similar level in the *pil5* mutant (Figure 2A), indicating that *PIL5* does not play a significant role in regulating the expression of *SOM* during seed maturation. In contrast with the *pil5* mutant, the expression level of *SOM* remained low in the *abi3* mutants during seed maturation and also in dry seeds (Figure 2A). These results indicate that *ABI3* regulates the expression of *SOM* during seed maturation.

Seed maturation is known to be regulated by three B3-domain proteins (i.e., *ABI3*, *FUS3*, and *LEC2*) and a HAP3 subunit of CCAAT binding protein (*LEC1*) (Giraudat et al., 1992; Luerssen



**Figure 2.** ABI3 Regulates the Expression of SOM.

- (A) Decreased expression of SOM mRNA (*SOM/PP2A*) in the *abi3-6* mutant but not in the *pil5* mutant seeds.  
 (B) Relative expression of SOM mRNA (*SOM/PP2A*) in freshly harvested *abi3-6* and other related mutant seeds. An inset indicates the relative expression levels of ABI3 mRNA (*ABI3/PP2A*) in *abi3-6* and its related mutant seeds.  
 (C) Increased expression of SOM mRNA (*SOM/PP2A*) in leaves of *ABI3-OX*. An inset indicates a magnified view of the relative expression levels of SOM mRNA in the absence of ABA treatment.  
 (D) Transient expression assay showing the activation of the firefly luciferase reporter gene linked to the *SOM* promoter by ABI3 but not by FUS3 or GFP in *Arabidopsis* protoplasts. Error bars are SD ( $n = 3$ ).

et al., 1998; Raz et al., 2001; Stone et al., 2001). To investigate if these other proteins also regulate the expression of SOM, we determined the expression levels of SOM transcript in *abi3*, *fus3*, *lec1*, and *lec2* mutant seeds. The *abi3* mutant seeds expressed the lowest level of SOM mRNA. The *fus3* and *lec1* mutant seeds also expressed lower levels of SOM than the wild type, while the *lec2* mutant expressed a similar level of SOM as the wild type (Figure 2B). Since ABI3 levels were lower in the *fus3* and *lec1* mutants than in the wild type (Figure 2B, inset), the reduced expression of SOM in the *fus3* and *lec1* mutants could be due to the lower levels of ABI3 in these mutants.

To further show that ABI3 is the major regulator of SOM, we generated transgenic lines expressing either ABI3 or FUS3 under the cauliflower mosaic virus (CaMV) 35S promoter. Transgenic lines expressing FUS3 exhibited inflorescences with flowers that opened prematurely and short and thick siliques. For the analysis, we selected homozygous transgenic lines that expressed similar levels of ABI3 and FUS3 (*ABI3-OX* and *FUS3-OX*) (see Supplemental Figure 1 online). We determined the expression levels of the SOM transcript in transgenic leaves to avoid the complex interlocked transcriptional regulation among ABI3 and other seed maturation genes in the seeds (To et al., 2006). SOM

mRNA was expressed at higher levels in *ABI3-OX* but only slightly higher levels in *FUS3-OX* (Figure 2C, inset). When the plants were subjected to ABA treatment, the SOM expression level increased dramatically only in the *ABI3-OX* line (Figure 2C), supporting the notion that ABI3 activates the expression of SOM. ABI3 also activates the expression of luciferase in a transient expression assay using *Arabidopsis* protoplasts and a 2-kb SOM promoter linked to the luciferase gene (Figure 2D). Taken together, these results suggest that the seed-specific expression of SOM is caused by the seed-specific expression of ABI3.

ABI3 is also required for the expression of SOM mRNA in imbibed seeds. When imbibed, the level of SOM transcript decreased drastically in the beginning of seed imbibition (after 3 h of seed imbibition) regardless of the light conditions. The initial decrease in SOM transcript is in agreement with the initial decrease in the majority of seed-specific mRNAs during seed imbibition. After imbibition, however, the expression of SOM depended on the light conditions. If a red light pulse was administered at the beginning of seed imbibition ( $\text{phyB}_{\text{on}}$ ), the SOM transcript level remained low, whereas it increased if a far-red pulse was given ( $\text{phyB}_{\text{off}}$ ) (Figures 3A and 3B). This light-dependent expression of SOM in imbibed seeds, as reported

before (Kim et al., 2008), was regulated by PIL5, as the *SOM* transcript levels remained low in the *pil5* mutant, regardless of the light conditions (Figure 3C). The *SOM* transcript levels remained even lower in the *abi3-sk11* mutant, regardless of the light conditions (Figure 3D). These results indicate that ABI3 is necessary for the high expression of *SOM* mRNA also in imbibed seeds.


### ABI3 Regulates the Expression of *SOM* by Directly Binding to RY Motifs in the *SOM* Promoter in Vivo

The presence of RY motifs in the *SOM* promoter suggested that ABI3 regulates the expression of *SOM* by directly binding to its promoter. To determine if ABI3 directly binds to the *SOM* promoter in vivo, we performed chromatin immunoprecipitation (ChIP) analysis using imbibed seeds of a transgenic line expressing flag-tagged ABI3 under the control of the CaMV 35S promoter. The flag-tagged ABI3 was functional, as it rescued the green desiccation-intolerant seed phenotype of the *abi3-6* mutant. After the ChIP with anti-flag antibody, the enrichments of specific *SOM* promoter fragments in the immunoprecipitates were determined by six primer pairs (F1 to F6; Figure 4A). A primer pair that amplifies the genomic region around the *5S rRNA* gene was included as a negative control, and two primer pairs that amplify two seed-specific gene promoters containing RY

motifs (*HsfA9* and *2S3*) were also included as potential ABI3 target sites (Kagaya et al., 2005; Kotak et al., 2007).

The ChIP-enriched *SOM* promoter fragments ranging from F3 to F5 fragments, with the strongest enrichment being at the F5 fragment (Figure 4A). The *HsfA9* and *2S3* promoters were also enriched to a greater extent than the negative control, indicating that ABI3 binds directly to the promoters of *SOM*, *HsfA*, and *2S3* in vivo. We noticed that the region of the *SOM* promoter that was enriched by ABI3 is relatively wide (>1 kb), as if ABI3 binds to a RY motif (pRY) at 0.3 kb upstream of the transcription start site in addition to a RY motif (dRY) at 1.4 kb upstream. However, not all fragments containing an RY motif were enriched. A fragment (F1) near an RY motif at 2.4 kb upstream of the transcription start site was not enriched much by ABI3, indicating that ABI3 binds to only some RY motifs.

To determine if ABI3 can directly bind to an RY motif present in the *SOM* promoter, we performed electrophoretic mobility shift assay (EMSA) analysis using a proximal RY motif-containing double-stranded oligomer from the F5 fragment region. Recombinant ABI3 fused to maltose binding protein (MBP-ABI3) bound to the oligomer, and its binding was effectively out-competed by nonlabeled oligomer (Figure 4B). Unlike unlabeled oligomer, however, a double-stranded oligomer containing a mutated RY motif (mpRY) did not compete for the binding to MBP-ABI3. Taken together, these results indicate that ABI3 binds to the




**Figure 3.** ABI3 Regulates the Light-Dependent Expression Levels of *SOM* mRNA in Imbibed Seeds.

**(A)** A diagram showing the light treatment schemes. Seeds were treated with either a far-red pulse (FR<sub>p</sub>, 2.8 μmol·m<sup>-2</sup>·s<sup>-1</sup> for 5 min) or a far-red pulse followed by a red pulse (R<sub>p</sub>, 18.7 μmol·m<sup>-2</sup>·s<sup>-1</sup> for 5 min) 1 h after the start of seed sterilization, before transferring the seeds to the dark.

**(B)** Light-dependent expression of *SOM* mRNA (*SOM/PP2A*) during imbibition of wild-type seeds (SD, *n* = 3).

**(C)** Abolition of light-dependent expression of *SOM* mRNA (*SOM/PP2A*) during imbibition of *pil5* mutant seeds (SD, *n* = 3).

**(D)** Decreased expression of *SOM* mRNA (*SOM/PP2A*) during imbibition of *abi3-sk11* mutant seeds (SD, *n* = 3).



**Figure 4.** ABI3 Directly Binds to the *SOM* Promoter through RY Motifs Both in Vivo and in Vitro.

**(A)** ChIP analysis showing the direct binding of ABI3 to the *SOM* promoter in vivo. Top diagram, the *SOM* promoter and three RY motifs (a proximal RY motif [pRY], a distal RY motif [dRY], and an RY motif near At1g03780 [RY]). Seeds imbibed for 12 h under phyB<sub>off</sub> conditions were used for ChIP analysis. F1 to F6 indicate genomic DNA fragments around the *SOM* promoter tested for enrichment by quantitative PCR (SD,  $n = 3$ ). IP indicates immunoprecipitated DNA, and IN indicates input DNA.

**(B)** EMSA analysis showing the direct binding of ABI3 to pRY in the *SOM* promoter in vitro. MBP-ABI3, ABI3 protein fused to MBP; cold probe, nonlabeled wild-type pRY (pRY) or mutated pRY (mpRY). The increased concentration of competitors (5 $\times$ , 10 $\times$ , and 20 $\times$ ) is indicated by triangles.

*SOM* promoter through RY motifs and activates the expression of *SOM*.


To investigate the role of these RY motifs further in vivo, we cloned a 2-kb region of the *SOM* promoter, fused it to a reporter gene ( $\beta$ -glucuronidase [*GUS*]), and generated transgenic lines (*SOM*<sub>pro</sub>:*GUS*). The 2-kb *SOM* promoter expressed the *GUS* reporter gene in seeds and mildly in hypocotyls, vegetative leaves, cauline leaves, and flowers (Figure 5A). When developing embryos were dissected out, we observed that *GUS* expression began to be detected in embryos 9 DPA and was strong in dry

seeds, consistent with the expression pattern of endogenous *SOM*. Unlike the *SOM* promoter, the CaMV 35S promoter drove expression of the *GUS* reporter gene throughout seedlings (Figure 5A). The expression of *SOM*-green fluorescent protein (GFP) under the same 2-kb *SOM* promoter also increased dramatically in developing seeds. In imbibed seeds, the *SOM*-GFP protein level decreased initially but increased again when phyB was in an inactive state, consistent with the expression pattern of endogenous *SOM* transcript in this condition (Figure 5B). These results indicate that the 2-kb region of the *SOM* promoter linked to a reporter gene functionally substitutes the endogenous *SOM* promoter.

To determine the role of RY motifs in regulating the expression of *SOM*, the 2-kb *SOM* promoter bearing mutations in two RY motifs, either individually or together, was fused to a *GUS* reporter gene and the respective transgenic lines were generated (*mpRY*:*GUS*, *mdRY*:*GUS*, and *mpdRY*:*GUS*). For each promoter construct, 10 randomly chosen independent transgenic lines were established, and the activities of promoters were determined by measuring the *GUS* transcript levels in the imbibed seeds. As expected, different transgenic lines expressed different levels of *GUS* transcript; thus, we determined the average expression levels of 10 transgenic lines. The average expression level of *GUS* mRNA driven by wild-type 2-kb *SOM* promoter was 38 relative to that of *PP2A* (Figure 5C). When a distal or a proximal RY motif was mutated, the average relative expression levels decreased to 10, whereas when both RY motifs were mutated, the average relative expression level decreased further to 5, indicating that two RY motifs contribute similarly to the high expression of *SOM*. Consistent with the view that both RY motifs contribute to *SOM* expression, the relative expression level of *SOM* was reduced when a T-DNA fragment was inserted between the proximal and distal RY motifs (*som-4* allele), whereas the level was further reduced when a T-DNA fragment was inserted between the proximal RY motif and the transcription start site (*som-3* allele) (Figure 5D). Taken together, these results indicate that both RY motifs are required for the high expression of *SOM*.

#### ABI3 Interacts with PIL5 to Potentiate the Expression of *SOM* in Imbibed Seeds

Our analysis indicates that the expression of *SOM* mRNA is regulated by both ABI3 and PIL5 in imbibed seeds. In imbibed seeds, public microarray data show that *PIL5* is expressed comparably in both tissues, while *ABI3* is expressed higher in endosperm. Consistent with the requirement of both ABI3 and PIL5 for the expression of *SOM*, *SOM* is also expressed higher in endosperms than in embryos of imbibed seeds (see Supplemental Figure 2 online). We further investigated the role of ABI3 and PIL5 in regulating the expression of *SOM* by comparing the expression levels of *SOM* in various mutants in imbibed seeds. In the wild type, the expression levels of *SOM* transcript were  $\sim$ 5-fold higher under the phyB<sub>off</sub> condition than the phyB<sub>on</sub> condition (Figure 6A). In the *pil5* mutant, the expression levels of *SOM* transcript were similarly low, regardless of the light conditions, consistent with a previous report that showed that PIL5 mediates light signaling to regulate the expression of *SOM* mRNA (Kim



**Figure 5.** RY Motifs Are Necessary for the High Expression of *SOM*.

**(A)** Seed-specific expression driven by a 2-kb region of the *SOM* promoter.  $SOM_{pro}:GUS$  indicates various tissues and organs of transgenic plants harboring  $SOM_{pro}:GUS$ , while  $35S_{pro}:GUS$  indicates transgenic plants harboring  $35S_{pro}:GUS$ . Numbers and DS on figures indicate DPA and dry seeds, respectively. Bar = 5 mm.

**(B)** Immunoblot analysis showing the increased expression of *SOM*-GFP protein by the *SOM* promoter during seed maturation and imbibition. hours, hours after imbibition.

**(C)** Decreased expression of *GUS* reporter transcripts (*GUS*/*PP2A*) by the *SOM* promoter with mutations in their RY motifs. Numbers indicate average  $\pm$  SD ( $n = 10$ ). Seeds imbibed for 24 h under phyB<sub>off</sub> condition were used for analysis. The inset shows the light-dependent expression of the *GUS* transcript in two transgenic lines. Error bars are SD ( $n = 3$ ).


**(D)** Decreased expression of *SOM* mRNA (*SOM*/*PP2A*) in the *som-4* allele. Seeds imbibed for 12 h were used for analysis. Triangles indicate RY motifs in the *SOM* promoter (SD,  $n = 3$ ).

et al., 2008). In the *abi3* mutants, the expression levels of *SOM* mRNA were greatly reduced relative to the wild type, indicating that *ABI3* is also necessary for the high expression of *SOM* (Figure 6A). However, the *SOM* mRNA was still expressed 6-fold higher in FR-treated seeds than in R-treated imbibed seeds of the *abi3* mutant, indicating that *ABI3*, although necessary for the high expression of *SOM*, is dispensable for the light-dependent expression of *SOM* mRNA. This reduced expression of *SOM* in the *abi3* mutant was not due to the lower levels of *PIL5* transcript (Figure 6B). Taken together, these results suggest that *PIL5* regulates the light-dependent expression of *SOM* mRNA even in the absence of *ABI3*, while *ABI3* collaborates with *PIL5* to increase the expression of *SOM* mRNA in imbibed seeds.

Numerous transcription factors have been shown to regulate the expression of target genes collaboratively by forming a complex on the promoters of the target genes. For example, *ABI3* has been shown to interact with a subset of bZIP transcription factors, including *ABI5*, bZIP10, and bZIP25, in *Arabi-*

*dopsis* (Nakamura et al., 2001; Lara et al., 2003). In maize, VP1, an *ABI3* homolog, also interacts with *TRAB1*, a bZIP transcription factor (Hobo et al., 1999). We investigated if *ABI3* and *PIL5* interact with each other at the protein level. We detected the interaction by an in vitro binding assay using MBP-tagged *ABI3* and His-tagged *PIL5*. As a positive control, we included His-tagged *ABI5*, a known *ABI3*-interacting transcription factor. When MBP-*ABI3* was precipitated, it coprecipitated *PIL5* (Figure 7A), whereas MBP alone did not coprecipitate *PIL5*, indicating that *ABI3* interacts with *PIL5*. Consistent with a previous report, MBP-*ABI3* but not MBP alone coprecipitated *ABI5*. The results indicate that *ABI3* interacts with various transcription factors, including *PIL5* and *ABI5*, at the protein level.

The physical interaction between *ABI3* and *PIL5* may suggest that they help each other to bind to the *SOM* promoter. Alternatively, they may bind to the *SOM* promoter independently but interact with each other to activate the expression of *SOM* mRNA. The light-dependent expression of *SOM* mRNA in the



**Figure 6.** PIL5 Requires ABI3 to Activate the Expression of SOM Efficiently.


(A) The expression levels of SOM mRNA (*SOM/PP2A*) in the wild type, the *pil5* and *abi3-sk11* mutants, and the *abi3-sk11 pil5* double mutant. Seeds were imbibed for 12 h. The inset indicates the fold difference in SOM mRNA expression between the phyB<sub>off</sub> and phyB<sub>on</sub> conditions (SD, *n* = 3). Student's *t* test: \*\**P* < 0.01.

(B) Expression levels of *PIL5* mRNA (*PIL5/PP2A*) in the wild type, the *pil5* and *abi3-sk11* mutants, and the *pil5 abi3-sk11* double mutant (SD, *n* = 3).

*abi3* mutant suggested that PIL5 binds to the *SOM* promoter even in the absence of ABI3 (Figure 6A). We further determined if ABI3 and PIL5 affect the binding of each other to the *SOM* promoter in vivo. ChIP analysis was used to determine the binding of ABI3 and PIL5 to the *SOM* promoter in vivo in the presence or absence of PIL5 or ABI3. The ChIP analysis showed that ABI3 binds to the *SOM* promoter similarly both in *ABI3-OX* and *ABI3-OX/pil5* (Figure 7B), while PIL5 also binds to the *SOM* promoter similarly both in *PIL5-OX* and *PIL5-OX/abi3* (Figure 7C), indicating that ABI3 and PIL5 bind to the *SOM* promoter even in the absence of each other. Taken together, these results indicate that PIL5 and ABI3 do not drastically affect each other's ability to bind to the *SOM* promoter in vivo. Rather, they suggest that they interact with each other to activate the expression of *SOM*.

**DISCUSSION**

We show that ABI3 binds to the *SOM* promoter through two RY motifs using ChIP analysis and EMSA. Binding of ABI3 to both RY motifs was necessary to activate the expression of *SOM* mRNA fully, as mutations in either of two RY motifs greatly reduced the expression of a reporter gene. We analyzed the relationship between ABI3 and PIL5, which also regulates the expression of *SOM* by directly binding to its promoter (Kim et al., 2008). We found that ABI3 and PIL5 regulate the expression of *SOM* either independently or collaboratively. In maturing seeds, ABI3, but not PIL5, was necessary for the high expression of *SOM* mRNA, whereas, in imbibed seeds, PIL5 mediated light-dependent expression of *SOM* mRNA and ABI3 potentiated the ability of PIL5 to increase the expression level of *SOM* mRNA. At the protein level, ABI3 interacts with PIL5. This protein-protein interaction, however, did not drastically affect the targeting of ABI3 to the *SOM* promoter, indicating that ABI3 and PIL5 interact with each other to regulate *SOM* transcription collaboratively in imbibed seeds. Since ABI3 and PIL5 are key signaling components of ABA signaling and light signaling (Lopez-Molina et al., 2002; Oh et al., 2007), respectively, our results suggest that the *SOM* promoter integrates ABA and light signaling to regulate seed germination.



**Figure 7.** ABI3 and PIL5 Interact at the Protein Level.

(A) In vitro binding assay showing the protein-protein interaction between ABI3 and PIL5. MBP-ABI3, MBP-fused ABI3 protein; PIL5H, His-tagged PIL5 protein; ABI5H, His-tagged ABI5 protein; IN, 5% of input proteins.

(B) ChIP assay showing the similar enrichments of *SOM* promoter fragments by ABI3-flag both in the *ABI3-OX* and *ABI3-OX pil5* lines. F1 and F5 are as indicated in Figure 4A. The values shown are immunoprecipitated DNA/input DNA relative to F1. Error bars are SD (*n* = 3).

(C) ChIP assay showing that PIL5-myc enriches *SOM* promoter fragments to a similar extent both in the *PIL5-OX3* and *PIL5-OX3 abi3-6* lines. F1 and F5 are as indicated in Figure 4A. The values shown are immunoprecipitated DNA/input DNA relative to F1 (SD, *n* = 3).



### **ABI3 Is Necessary for the High Expression of *SOM* Both in Maturing and Imbibed Seeds, but *PIL5* Is Necessary Only in Imbibed Seeds**

Our expression analysis shows that *ABI3* is necessary for the activation of *SOM* mRNA both in maturing seeds and in imbibed seeds. However, *PIL5* was necessary only in the imbibed seeds, but not in the maturing seeds. What determines this stage-specific role of *PIL5* is not clear. A few different possible explanations may account for the stage specificity. First, maturing seeds may express other *PIF* family members that activate the expression of *SOM* mRNA redundantly with *PIL5*. Public microarray data indicate that *PIL5* is expressed at lower levels than other *PIFs* (*PIF5* and *PIF6*) in maturing seeds, while *PIL5* is a dominant *PIF* in imbibed seeds (see Supplemental Figure 3 online). If other *PIFs* could activate the expression of *SOM* redundantly with *PIL5*, the mutation in *PIL5* would be phenotypically manifested more severely in imbibed seeds than in maturing seeds. Second, other transcription factors that are known to interact with *ABI3* may functionally substitute for *PIL5* in maturing seeds. The same public microarray data indicate that *ABI5*, *EEL*, *DPBF2*, and *AREB3* are expressed at high levels in maturing seeds (see Supplemental Figure 3 online) (Bensmihen et al., 2005). Since these bZIP transcription factors are known to activate the expression of seed storage-specific protein genes in maturing seeds, it is possible that these bZIP factors bind to the *SOM* promoter and activate the expression of *SOM* in maturing seeds. Alternatively, stage-specific modification or cofactors that enable *ABI3* to activate the expression of *SOM* mRNA alone may account for the dispensability of *PIL5* in maturing seeds. Further analysis is needed to clarify why *PIL5* is dispensable for the activation of *SOM* mRNA in maturing seeds.

### ***ABI3* Binds to the *SOM* Promoter through RY Motifs to Activate the Expression of *SOM* mRNA**

A previous study showed that *SOM* is a C3H-type zinc finger protein that inhibits seed germination downstream of *PIL5* partly by activating ABA biosynthesis and inhibiting GA biosynthesis in imbibed seeds (Oh et al., 2007; Kim et al., 2008). We further investigated how the expression of *SOM* is regulated in seeds. Public microarray data indicated that *SOM* is highly expressed in maturing seeds, which is reminiscent of the expression pattern of *ABI3*-regulated genes, such as *ABI5* (Figure 1A). The presence of RY motifs, which serve as a binding site of B3-domain proteins, such as *ABI3*, *FUS3*, and *LEC2*, further suggested that *SOM* is regulated by *ABI3* or its homologs. Consistent with this notion, the expression level of *SOM* was drastically reduced in *abi3* and weakly reduced in *fus3*. Transgenic lines overexpressing *ABI3* and *FUS3* showed that the expression of *SOM* mRNA is activated mainly by *ABI3*. Apparently, ABA strongly enhanced the expression of *SOM* by *ABI3*. Taken together, our results suggest that the high expression of *SOM* in maturing seeds is caused by the seed-specific expression of *ABI3* and the concomitant increase in ABA level during seed maturation.

Our ChIP analysis coupled with EMSA indicated that *ABI3* directly binds to two regions of the *SOM* promoter through RY

motifs and activates the expression of *SOM* mRNA. The binding of *ABI3* to both RY motifs was necessary for the high expression of *SOM* mRNA, as mutations in either one of the RY motifs greatly decreased the expression levels of a reporter gene. As in the *SOM* promoters, multiple RY motifs are also found in the promoters of many seed storage protein genes, including *OLEOSIN*, *CRUCIFERIN*, and *2S*, suggesting that promoters of *ABI3*-regulated genes recruit multiple copies of *ABI3*. Interestingly, an insertion of a 4.5-kb T-DNA fragment from *pBIN-pROK2* between two of the RY motifs also reduced the expression of *SOM* mRNA. Since the insertion of the T-DNA fragment increases the distance between the distal RY motif and the transcription start site, the reduced expression may indicate that *ABI3* must be in relatively close proximity to the transcription start site to activate the transcription. Alternatively, the insertion of T-DNA may disrupt the molecular interaction among proteins that bind to the proximal region and proteins that bind to distal regions of the *SOM* promoter.

The ChIP analysis further indicates that *ABI3* binds to only a subset of RY motifs in vivo. A genomic region we scanned by ChIP analysis included three RY motifs. Among these RY motifs, only two were highly enriched, and one was not enriched by *ABI3*. The enrichment was not correlated with the distance from the transcription start site, as two enriched RY motifs were 0.3 and 1.4 kb upstream of the *SOM* transcription start site, respectively, while one nonenriched RY motif was 0.3 kb upstream of the transcription start site of At1g03780 (Figure 1D). These results indicate that *ABI3* binds to only a subset of RY motifs in vivo and imply that in vivo *ABI3* binding sites will have to be determined experimentally, such as by ChIP analysis. Binding of the transcription factor to its target appeared to be more selective under in vivo conditions than in vitro, and this seems to be a rather general phenomenon, instead of an exception. A recent ChIP-Chip analysis showed that *PIL5* binds to only a subset of G-box elements in vivo (Oh et al., 2009). Similarly, a bZIP transcription factor, *HY5*, binds to only a subset of G-box elements (Lee et al., 2007). The molecular mechanisms that determine how the in vivo binding sites are selected for each transcription factor are not known.

### ***ABI3* and *PIL5* Interact and Activate the Expression of *SOM* mRNA Collaboratively in Imbibed Seeds**

*ABI3* and *PIL5* collaboratively activated the expression of *SOM* mRNA. The expression of *SOM* mRNA was greatly reduced in the *abi3* mutant. Mutations in RY motifs greatly decreased the expression of reporter genes, supporting the hypothesis that *ABI3* activates the expression of *SOM*. Nevertheless, the expression of *SOM* mRNA was still repressible by red light, as long as *PIL5* was present, indicating that *PIL5* mediates light signaling to activate the expression of *SOM* mRNA, while *ABI3* enhances the expression level.

Although the molecular mechanisms underlying the ability of *ABI3* to enhance the activity of *PIL5* are not known, the interaction between *ABI3* and *PIL5* at the protein level may contribute to this collaboration. *ABI3* and *PIL5* may form a protein complex that activates the transcription more efficiently. Alternatively, *ABI3* and *PIL5* may help each other bind to the *SOM* promoter.

Our data indicate that ABI3 and PIL5 do not drastically affect each other's targeting to the *SOM* promoter, implying that the synergism comes after their targeting to the promoter, rather than during the targeting itself (Figures 6 and 7). Our ChIP analysis indicates that ABI3 enriches the *SOM* promoter and that this enrichment is of a similar magnitude in the wild type and the *pil5* mutant. PIL5 also enriched the *SOM* promoter, and this enrichment was similar in the wild type and the *abi3* mutants. These results indicate that ABI3 and PIL5 can bind to their target sites in the absence of each other. Our results are similar to a previous report showing that ABRE binding factors bind to the maize *rab28* promoter even in the absence of VP1 (Busk and Pagès, 1997), while they are different to another report showing that ABI3 slightly enhances the binding between bZIP10/OPA-QUE2 and the G-box element in vitro (Lara et al., 2003). Since a slight increase or decrease in binding affinity is difficult to resolve in our ChIP assay, the results do not exclude the possibility that ABI3 mildly affects the ability of PIL5 or other interacting bZIP factors to bind to DNA. Therefore, even though ABI3 and PIL5 do not drastically affect each other's binding to *SOM* promoter, further analysis is needed to determine if they affect DNA binding activity more subtly.

The direct interaction between ABI3 and PIL5 indicates that three germination-inhibiting transcription factors (PIL5, ABI3, and ABI5) form a functional module, both through transcriptional regulation and protein-protein interaction, to inhibit seed germination. At the transcriptional level, PIL5 directly binds to promoters of *ABI3* and *ABI5* and activates the expression of their mRNAs, while ABI3 activates the expression of *ABI5* mRNA. At the protein level, ABI3 interacts with both its regulator (PIL5) and its regulatee (ABI5), which leads to the synergistic activation of their target gene expression. This type of functional module that is composed of interconnected transcription factors is likely to be useful to regulate the expression of wide sets of genes either independently, in combination with each other or with other proteins as well, or interdependently. Further studies are needed to decipher which target genes are regulated by these three factors independently, in combination, or interdependently.

## METHODS

### Plant Materials and Growth Conditions

*Arabidopsis thaliana* plants were grown in a growth room with a 16-h light/8-h-dark cycle at 22 to 24°C. T-DNA insertion lines (*abi3-sk11*, *som-4*, *lec1*, and *lec2*) were obtained from the Salk Institute (Salk\_023411, CS806302, Salk\_131219, and CS873714, respectively), and insertion sites of *abi3-sk11* and *som-4* were determined by amplifying and sequencing the flanking regions. The *lec1* and *lec2* mutants were confirmed by monitoring anthocyanin accumulation and desiccation intolerance in seeds. To generate *ABI3-OX* and *FUS3-OX* plants, full-length *ABI3* and *FUS3* cDNAs were amplified, cloned into binary vectors, and transformed into *Arabidopsis*, and then homozygous lines were selected (primers are presented in Supplemental Table 1 online). For transgenic lines harboring the GFP-tagged mini-gene construct of *SOM* (*SOM<sub>pro</sub>-SOM-GFP*) in the *som-2* background, *SOM* cDNA and its promoter region were sequentially cloned into the pbGFP1 binary vector and transformed into the *som-2* mutant. Homozygous lines were selected. For mutagenesis of the RY motifs in the *SOM* promoter, inverse PCR was performed with mutated

primer sets (see Supplemental Table 1 online) using a *SOM* promoter-containing plasmid as a template. Mutated *SOM* promoters were confirmed by sequencing, cloned into a pbGG4 binary vector for promoter-reporter construction, and transformed into Columbia-0 (Col-0). Ten independent transgenic lines showing the 3:1 segregation ratio were used to monitor the expression of *GUS* mRNA during imbibition. The *abi3-6* and *fus3-3* mutant seeds were kind gifts from F. Parcy at Centre National de la Recherche Scientifique, France. All the mutants used in this study (*som-2*, *som-3*, *som-4*, *pil5-1*, *PIL5-OX3*, *lec1*, *lec2*, *fus3-3*, *abi3-sk11*, and *abi3-6*) are in the Col-0 background. Desiccation-intolerant mutants were maintained by sowing seeds harvested from green siliques.

### Quantitative and Qualitative Gene Expression Analysis

For mRNA expression analysis, total RNAs were extracted either from dried seeds, imbibed seeds, or transgenic leaves using a Spectrum plant total RNA kit (Sigma-Aldrich) and converted to cDNAs using MMLV-RTase (Promega) according to the manufacturers' protocols. Transcript levels of each mRNA were determined by real-time PCR and normalized with the level of *PP2A* mRNA using the delta Ct method (Czechowski et al., 2005; Schmittgen and Livak, 2008). Amplification log curves of each primer set were confirmed to be parallel to that of *PP2A* amplification to validate the comparisons among samples and primer sets before use. Real-time PCR was performed with a protocol (45 cycles, each cycle consisting of 95°C/58°C/72°C for 20 s each) in an iCycler iQ5 cyclor (Bio-Rad). We tested more than three different seed batches to confirm the gene expression patterns and presented representative results in each figure.

Expression browser and Heat-Mapper tools provided by BAR (The Bio-Array Resource for *Arabidopsis* Functional Genomics; <http://bbc.botany.utoronto.ca/>) were employed to show the heat map of gene expression patterns.

To determine the tissue-specific expression of *SOM* by histochemical *GUS* reporter assay, seed coats, which inhibit penetration of X-Gluc into seeds, were carefully pinched off the mature or imbibed seeds using forceps. Decoated seed embryos were stained for *GUS* activity after having undergone fixing in 90% acetone for 30 min, as described previously (Jefferson et al., 1987).

For the effector activity assay of ABI3 and FUS3 using *Arabidopsis* protoplasts, mesophyll protoplasts were prepared from fully mature leaves as described previously (Yoo et al., 2007). Briefly, leaf slices were dipped in cell wall digestion solution (10 mM MES, pH 5.7, 1.5% cellulase R10, 0.4% macerozyme R10, 0.4 M mannitol, 20 mM KCl, 10 mM CaCl<sub>2</sub>, and 0.1% [w/v] BSA) and incubated for 3 h with mild shaking. One hundred microliters of protoplasts ( $2 \times 10^4$ ) was used for each transfection. After incubation for 1 d, luciferase activity was measured using a Dual Luciferase assay kit (Promega).

For the analysis of protein levels, samples were ground in protein extraction buffer (100 mM NaH<sub>2</sub>PO<sub>4</sub>, 8 M urea, and 10 mM TrisCl, pH 8.0) using Tissue-lyzer (Qiagen). Tissue lysates were cleared by centrifugation at 20,000g for 10 min and subjected to immunoblotting as described previously (Park et al., 2004).

### In Vitro Binding Assay

For the in vitro binding assay between PIL5 and ABI3, recombinant MBP-ABI3 protein was produced in *Escherichia coli* at 25°C using the pMAL-c2X-ABI3 expression vector and purified using amylose-resin following the manufacturer's protocol (NEB) (see Supplemental Table 1 online for primer sets). His-tagged PIL5 and ABI5 (PIL5H and ABI5H) were produced as described previously (Oh et al., 2007). MBP-resin and MBP-ABI3-resin were incubated with purified PIL5H and ABI5H at 4°C for 2 h, sedimented by spin down, and washed five times. MBP resin was used as

an MBP control as well as a nonspecific binding control. Then, 1× PBS with 0.1% Nonidet P-40, 0.5 mM DTT, 10% glycerol, 1 mM PMSF, and Protease inhibitor cocktail (Roche) was used for binding and washing. PIL5H and ABI5H bound to MBP-ABI3 were visualized by protein blotting using anti-His antibody.

#### ChIP Analysis and EMSA

ChIP was performed essentially as described previously, except that three separate steps (the elution of the DNA–protein complex from resin, the reversal of cross-linking, and the digestion with proteinase K) were combined into one step in this study (Kim et al., 2008). Immunoprecipitated DNA was eluted from resin in 200  $\mu$ L of elution buffer containing 20 mM Tris-HCl, pH 7.5, 5 mM EDTA, 50 mM NaCl, 1% SDS, and 50  $\mu$ g/mL proteinase K at 68°C for 2 h with rotary mixing. Residual DNA was further eluted by subsequent incubation in 100  $\mu$ L of elution buffer at 68°C for 5 min. Eluted DNA was purified with a PCR purification kit (Solgent) and subjected to real-time PCR. For ChIP of flag-tagged ABI3, anti-flag antibody-conjugated resin (Sigma-Aldrich) was used, while anti-myc Ab (Cell Signaling) and protein A-conjugated resin (Upstate) was used for ChIP of myc-tagged PIL5. For EMSA, previously described procedures were followed, except that 5' biotin-labeled probes were synthesized by chemical modification (Bioneer).

#### Accession Numbers

Sequence data from this article can be found in the Arabidopsis Genome Initiative or GenBank/EMBL databases under the following accession numbers: PIL5 (At2g20180), ABI3 (At3g24650), SOM (At1g03790), LEC1 (At1g21970), LEC2 (At1g28300), FUS3 (At3g26790), ABI5 (At2g36270), Em1 (At3g51810), Em6 (At2g40170), 2S3 (At4g27160), RGA (At2g01570), HsfA9 (At5g54070), 5S rRNA (At3g41979), and PP2A (At1g13320).

#### Supplemental Data

The following materials are available in the online version of this article.

**Supplemental Figure 1.** Expression Levels of *ABI3* and *FUS3* Transgenes.

**Supplemental Figure 2.** Tissue-Specific Expression of *ABI3*, *PIL5*, and *SOM*.

**Supplemental Figure 3.** Expression of *PIFs* and *bZIPs* during Seed Maturation and Imbibition.

**Supplemental Table 1.** Primers Used in This Study.

#### ACKNOWLEDGMENTS

This work was supported in part by grants from the National Research Foundation of Korea (2007-0056949 and 2009-0077873) to G.C.

Received October 25, 2010; revised February 12, 2011; accepted March 25, 2011; published April 5, 2011.

#### REFERENCES

- Bensmihen, S., Giraudat, J., and Parcy, F. (2005). Characterization of three homologous basic leucine zipper transcription factors (bZIP) of the ABI5 family during *Arabidopsis thaliana* embryo maturation. *J. Exp. Bot.* **56**: 597–603.
- Bentsink, L., Jowett, J., Hanhart, C.J., and Koornneef, M. (2006). Cloning of DOG1, a quantitative trait locus controlling seed dormancy in Arabidopsis. *Proc. Natl. Acad. Sci. USA* **103**: 17042–17047.
- Bies-Etheve, N., da Silva Conceicao, A., Giraudat, J., Koornneef, M., Léon-Kloosterziel, K., Valon, C., and Delseny, M. (1999). Importance of the B2 domain of the Arabidopsis ABI3 protein for Em and 2S albumin gene regulation. *Plant Mol. Biol.* **40**: 1045–1054.
- Braybrook, S.A., Stone, S.L., Park, S., Bui, A.Q., Le, B.H., Fischer, R.L., Goldberg, R.B., and Harada, J.J. (2006). Genes directly regulated by LEAFY COTYLEDON2 provide insight into the control of embryo maturation and somatic embryogenesis. *Proc. Natl. Acad. Sci. USA* **103**: 3468–3473.
- Busk, P.K., and Pagès, M. (1997). Protein binding to the abscisic acid-responsive element is independent of VIVIPAROUS1 in vivo. *Plant Cell* **9**: 2261–2270.
- Choi, H., Hong, J., Ha, J., Kang, J., and Kim, S.Y. (2000). ABFs, a family of ABA-responsive element binding factors. *J. Biol. Chem.* **275**: 1723–1730.
- Czechowski, T., Stitt, M., Altmann, T., Udvardi, M.K., and Scheible, W.R. (2005). Genome-wide identification and testing of superior reference genes for transcript normalization in Arabidopsis. *Plant Physiol.* **139**: 5–17.
- Finch-Savage, W.E., and Leubner-Metzger, G. (2006). Seed dormancy and the control of germination. *New Phytol.* **171**: 501–523.
- Finkelstein, R., Reeves, W., Ariizumi, T., and Steber, C. (2008). Molecular aspects of seed dormancy. *Annu. Rev. Plant Biol.* **59**: 387–415.
- Giraudat, J., Hauge, B.M., Valon, C., Smalle, J., Parcy, F., and Goodman, H.M. (1992). Isolation of the Arabidopsis ABI3 gene by positional cloning. *Plant Cell* **4**: 1251–1261.
- Hennig, L., Stoddart, W.M., Dieterle, M., Whitelam, G.C., and Schäfer, E. (2002). Phytochrome E controls light-induced germination of Arabidopsis. *Plant Physiol.* **128**: 194–200.
- Hobo, T., Kowayama, Y., and Hattori, T. (1999). A bZIP factor, TRAB1, interacts with VP1 and mediates abscisic acid-induced transcription. *Proc. Natl. Acad. Sci. USA* **96**: 15348–15353.
- Holdsworth, M.J., Bentsink, L., and Soppe, W.J. (2008). Molecular networks regulating Arabidopsis seed maturation, after-ripening, dormancy and germination. *New Phytol.* **179**: 33–54.
- Jefferson, R.A., Kavanagh, T.A., and Bevan, M.W. (1987). GUS fusions: Beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J.* **6**: 3901–3907.
- Kagaya, Y., Okuda, R., Ban, A., Toyoshima, R., Tsutsumida, K., Usui, H., Yamamoto, A., and Hattori, T. (2005). Indirect ABA-dependent regulation of seed storage protein genes by FUSCA3 transcription factor in Arabidopsis. *Plant Cell Physiol.* **46**: 300–311.
- Kim, D.H., Yamaguchi, S., Lim, S., Oh, E., Park, J., Hanada, A., Kamiya, Y., and Choi, G. (2008). SOMNUS, a CCH-type zinc finger protein in Arabidopsis, negatively regulates light-dependent seed germination downstream of PIL5. *Plant Cell* **20**: 1260–1277.
- Kotak, S., Vierling, E., Bäumllein, H., and von Koskull-Döring, P. (2007). A novel transcriptional cascade regulating expression of heat stress proteins during seed development of Arabidopsis. *Plant Cell* **19**: 182–195.
- Lara, P., Oñate-Sánchez, L., Abraham, Z., Ferrándiz, C., Díaz, I., Carbonero, P., and Vicente-Carbajosa, J. (2003). Synergistic activation of seed storage protein gene expression in Arabidopsis by ABI3 and two bZIPs related to OPAQUE2. *J. Biol. Chem.* **278**: 21003–21011.
- Lee, J., He, K., Stolc, V., Lee, H., Figueroa, P., Gao, Y., Tongprasit, W., Zhao, H., Lee, I., and Deng, X.W. (2007). Analysis of transcription factor HY5 genomic binding sites revealed its hierarchical role in light regulation of development. *Plant Cell* **19**: 731–749.

- Léon-Kloosterziel, K.M., Gil, M.A., Ruijs, G.J., Jacobsen, S.E., Olszewski, N.E., Schwartz, S.H., Zeevaart, J.A., and Koornneef, M. (1996). Isolation and characterization of abscisic acid-deficient *Arabidopsis* mutants at two new loci. *Plant J.* **10**: 655–661.
- Liu, Y., Koornneef, M., and Soppe, W.J. (2007). The absence of histone H2B monoubiquitination in the *Arabidopsis* hub1 (*rdo4*) mutant reveals a role for chromatin remodeling in seed dormancy. *Plant Cell* **19**: 433–444.
- Lopez-Molina, L., Mongrand, S., McLachlin, D.T., Chait, B.T., and Chua, N.H. (2002). ABI5 acts downstream of ABI3 to execute an ABA-dependent growth arrest during germination. *Plant J.* **32**: 317–328.
- Luerssen, H., Kirik, V., Herrmann, P., and Miséra, S. (1998). FUSCA3 encodes a protein with a conserved VP1/AB13-like B3 domain which is of functional importance for the regulation of seed maturation in *Arabidopsis thaliana*. *Plant J.* **15**: 755–764.
- Mönke, G., Altschmied, L., Tewes, A., Reidt, W., Mock, H.P., Bäumllein, H., and Conrad, U. (2004). Seed-specific transcription factors ABI3 and FUS3: Molecular interaction with DNA. *Planta* **219**: 158–166.
- Nakamura, S., Lynch, T.J., and Finkelstein, R.R. (2001). Physical interactions between ABA response loci of *Arabidopsis*. *Plant J.* **26**: 627–635.
- Oh, E., Kang, H., Yamaguchi, S., Park, J., Lee, D., Kamiya, Y., and Choi, G. (2009). Genome-wide analysis of genes targeted by PHYTOCHROME INTERACTING FACTOR 3-LIKE5 during seed germination in *Arabidopsis*. *Plant Cell* **21**: 403–419.
- Oh, E., Kim, J., Park, E., Kim, J.I., Kang, C., and Choi, G. (2004). PIL5, a phytochrome-interacting basic helix-loop-helix protein, is a key negative regulator of seed germination in *Arabidopsis thaliana*. *Plant Cell* **16**: 3045–3058.
- Oh, E., Yamaguchi, S., Hu, J., Yusuke, J., Jung, B., Paik, I., Lee, H.S., Sun, T.P., Kamiya, Y., and Choi, G. (2007). PIL5, a phytochrome-interacting bHLH protein, regulates gibberellin responsiveness by binding directly to the GAI and RGA promoters in *Arabidopsis* seeds. *Plant Cell* **19**: 1192–1208.
- Oh, E., Yamaguchi, S., Kamiya, Y., Bae, G., Chung, W.I., and Choi, G. (2006). Light activates the degradation of PIL5 protein to promote seed germination through gibberellin in *Arabidopsis*. *Plant J.* **47**: 124–139.
- Okamoto, M., et al. (2010). Genome-wide analysis of endogenous abscisic acid-mediated transcription in dry and imbibed seeds of *Arabidopsis* using tiling arrays. *Plant J.* **62**: 39–51.
- Park, E., Kim, J., Lee, Y., Shin, J., Oh, E., Chung, W.I., Liu, J.R., and Choi, G. (2004). Degradation of phytochrome interacting factor 3 in phytochrome-mediated light signaling. *Plant Cell Physiol.* **45**: 968–975.
- Raz, V., Bergervoet, J.H., and Koornneef, M. (2001). Sequential steps for developmental arrest in *Arabidopsis* seeds. *Development* **128**: 243–252.
- Schmittgen, T.D., and Livak, K.J. (2008). Analyzing real-time PCR data by the comparative C(T) method. *Nat. Protoc.* **3**: 1101–1108.
- Seo, M., Nambara, E., Choi, G., and Yamaguchi, S. (2009). Interaction of light and hormone signals in germinating seeds. *Plant Mol. Biol.* **69**: 463–472.
- Shinomura, T., Nagatani, A., Hanzawa, H., Kubota, M., Watanabe, M., and Furuya, M. (1996). Action spectra for phytochrome A- and B-specific photoinduction of seed germination in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **93**: 8129–8133.
- Stone, S.L., Kwong, L.W., Yee, K.M., Pelletier, J., Lepiniec, L., Fischer, R.L., Goldberg, R.B., and Harada, J.J. (2001). LEAFY COTYLEDON2 encodes a B3 domain transcription factor that induces embryo development. *Proc. Natl. Acad. Sci. USA* **98**: 11806–11811.
- Suzuki, M., Kao, C.Y., and McCarty, D.R. (1997). The conserved B3 domain of VIVIPAROUS1 has a cooperative DNA binding activity. *Plant Cell* **9**: 799–807.
- To, A., Valon, C., Savino, G., Guilleminot, J., Devic, M., Giraudat, J., and Parcy, F. (2006). A network of local and redundant gene regulation governs *Arabidopsis* seed maturation. *Plant Cell* **18**: 1642–1651.
- Yoo, S.D., Cho, Y.H., and Sheen, J. (2007). *Arabidopsis* mesophyll protoplasts: A versatile cell system for transient gene expression analysis. *Nat. Protoc.* **2**: 1565–1572.