

 Open access • Journal Article • DOI:10.1002/PATH.1661

Abnormal PcG protein expression in Hodgkin's lymphoma. Relation with E2F6 and NFκB transcription factors — [Source link](#)

Margarita Sánchez-Beato, Esther María Sánchez Sánchez, Juan F. García, Alberto Pérez-Rosado ...+10 more authors

Institutions: Museo Nacional Del Prado, Spanish National Research Council

Published on: 01 Dec 2004 - The Journal of Pathology (J Pathol)

Topics: Transcription factor, E2F Transcription Factors, Cell cycle, Polycomb-group proteins and BMI1

Related papers:

- [Unique polycomb gene expression pattern in Hodgkin's lymphoma and Hodgkin's lymphoma-derived cell lines.](#)
- [Coexpression of BMI-1 and EZH2 polycomb group genes in Reed-Sternberg cells of Hodgkin's disease.](#)
- [RYBP, a new repressor protein that interacts with components of the mammalian Polycomb complex, and with the transcription factor YY1.](#)
- [Variability in the expression of polycomb proteins in different normal and tumoral tissues. A pilot study using tissue microarrays.](#)
- [Loss of the B-lineage-specific gene expression program in Hodgkin and Reed-Sternberg cells of Hodgkin lymphoma.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/abnormal-pcg-protein-expression-in-hodgkin-s-lymphoma-3zajtj9c>

The requested paper is not presently available in its full-text version for it has not been supplied yet by the researcher in charge of the archiving.

*

*

*

*

El artículo seleccionado no se encuentra disponible por ahora a texto completo por no haber sido facilitado todavía por el investigador a cargo del archivo del mismo.