

## Accumulation of Amyloid $\beta$ and Tau and the Formation of Neurofilament Inclusions Following Diffuse Brain Injury in the Pig

D. H. SMITH, MD, X-H CHEN, MD, M. NONAKA, MD, J. Q. TROJANOWSKI, MD, PhD, VM-Y LEE, PHD, K. E. SAATMAN, PHD, M. J. LEONI, B-N XU, MD, J. A. WOLF, AND D. F. MEANEY, PHD

**Abstract.** Brain trauma in humans increases the risk for developing Alzheimer disease (AD) and may induce the acute formation of AD-like plaques containing amyloid  $\beta$  (A $\beta$ ). To further explore the potential link between brain trauma and neurodegeneration, we conducted neuropathological studies using a pig model of diffuse brain injury. Brain injury was induced in anesthetized animals via nonimpact head rotational acceleration of 110° over 20 ms in the coronal plane ( $n = 15$  injured,  $n = 3$  noninjured). At 1, 3, 7, and 10 days post-trauma, control and injured animals were euthanized and immunohistochemical analysis was performed on brain sections using antibodies specific for A $\beta$ ,  $\beta$ -amyloid precursor protein ( $\beta$ PP), tau, and neurofilament (NF) proteins. In addition to diffuse axonal pathology, we detected accumulation of A $\beta$  and tau that colocalized with immunoreactive  $\beta$ PP and NF in damaged axons throughout the white matter in all injured animals at 3–10 days post-trauma. In a subset of brain injured animals, diffuse A $\beta$ -containing plaque-like profiles were found in both the gray and white matter, and accumulations of tau and NF rich inclusions were observed in neuronal perikarya. These results show that this pig model of diffuse brain injury is characterized by accumulations of proteins that also form pathological aggregates in AD and related neurodegenerative diseases.

**Key Words:** Alzheimer disease; Amyloid precursor protein; Amyloid  $\beta$ ; Brain trauma; Neurodegeneration; Neurofilament; Tau.

### INTRODUCTION

While traumatic brain injury is one of the leading causes of death and disability (1, 2), mounting evidence also suggests that brain trauma may have prolonged effects and initiate insidiously progressive neurodegenerative processes. Previously, postmortem histopathologic analysis of brains from boxers with *dementia pugilistica* ("punch-drunk syndrome") revealed neurofibrillary tangles (NFTs) and diffuse plaques composed of amyloid  $\beta$  peptides (A $\beta$ s) similar to the hallmark lesions of Alzheimer disease (AD) (3, 4). Subsequently, a single incident of brain trauma was shown to induce the formation of A $\beta$  plaques within days following injury (5, 6). In addition, brain trauma patients have been shown to have accelerated cognitive decline during aging (7, 8) and an increased risk of developing AD, even if the injury occurred in the remote past (9–11). Moreover, we and others have recently observed that brain trauma in the rat induces substantially progressive neuron loss, axonal degeneration, and atrophy that proceeds unabated for at least one year following injury (12–13). Taken together, the results from these studies suggest that neurodegenerative changes triggered by brain trauma may follow a remarkably complex and prolonged temporal course. However, the mechanisms underlying the

---

From the Departments of Neurosurgery (DHS, X-HC, MN, KES, B-NX, ML, JAW), Pathology and Laboratory Medicine (JQT, VM-YL), and Biocengineering (DFM), University of Pennsylvania, Philadelphia, Pennsylvania.

Correspondence to: Douglas H. Smith, MD, Department of Neurosurgery, University of Pennsylvania, 3320 Smith Walk, 105 Hayden Hall, Philadelphia, PA 19104.

This study was funded, in part, by NIH grants NS08803, NS38104, AG 12527, and NS 35721, AG09215 and AG10124.

relationship between brain trauma and neurodegenerative processes remain unknown.

It has been observed that brain trauma in humans and experimental animals induces marked accumulations of  $\beta$ -amyloid precursor proteins ( $\beta$ PPs) (14–18), suggesting that ample substrate is available for A $\beta$  production. However, experimental investigations of this relationship have been hampered since accumulation of A $\beta$  has not been observed in standard rodent models of focal brain trauma (14–16, 18, 19). As previously suggested, the post-traumatic absence of A $\beta$  accumulation in rodents may reflect a difference in the processing of  $\beta$ PP compared with humans (15, 19).

To further explore the potential link between brain trauma and neurodegeneration, in the present study, we used a well-characterized and clinically relevant model of diffuse brain injury in the pig (20–22). The most salient feature of this pig model is the production of widespread axonal pathology in the white matter resulting from nonimpact rotational acceleration of the head. This head rotation induces inertial loading to the pig brain that replicates the forces experienced by the human brain during traumatic events such as automotive crashes. The rationale to use this model was based on the observation that many trauma patients who developed A $\beta$  plaques suffered from diffuse brain injury. In addition, this model affords an opportunity to evaluate neurodegenerative changes following diffuse brain trauma in an animal with a relatively high order, gyrencephalic brain.

### MATERIALS AND METHODS

In these studies, we carefully adhered to the animal welfare guidelines set forth in the *Guide for the Care and Use of Laboratory Animals*, US Department of Health and Human Services Publication, 85-23. All animal procedures were approved by

TABLE  
Summary of Antibodies Used for Immunohistochemistry

Antibody	Epitope protein/amino acids	Type	Dilution	Reference
369W	$\beta$ PP/645-694a	P	1:1000	(44, 45)
22C11	$\beta$ PP/60-100	M	1:5	(46-48)
13335	A $\beta$ /1-42	P	1:1000	(49)
Karen	$\beta$ PP/N-Terminal	P	1:800	(49)
2332	A $\beta$ /1-17	P	1:4000	(50-52)
BCO5	A $\beta$ /1-42(43)	M	1:200	(53, 54)
4G8	A $\beta$ /17-24	M	1:1000	(55, 56)
6E10	A $\beta$	M	1:200	(51, 56, 57)
10A5	A $\beta$ /1-28	M	1:50	(58, 59)
Tau-2	Tau/192-204	M	1:400	(60-62)
PHF-1	PHF-tau	M	1:200	(63-66)
PHF-6	PHF-tau	M	1:250	(67-69)
PHF-13.5	PHF-tau	M	1:1000	(67) (69)
NR4	NF-L	M	1:400	(70-72)
NS2	NF-H	M	1:400	(70, 73, 74)
Alz-50	A $\beta$ 8 protein	M	1:5	(75)

the University of Pennsylvania Institutional Animal Care and Use Committee.

#### Preinjury Preparation

Eighteen miniature young adult swine (4 months of age, Hanford and Hormel strains), both male and female, weighing 17-20 kg, were used for this study ( $n = 3$  noninjured,  $n = 15$  brain injured). The animals were fasted for 12 h, after which anesthesia was induced with an initial injection of midazolam (400-600 mg/kg, i.m.). Once sedated, animals received 2%-4% isoflurane via snout mask until they reached a plane of surgical anesthesia. A venous catheter was then inserted in the ear, and the animals were endotracheally intubated and maintained on 1.5%-2% isoflurane. Physiologic monitoring and apparatus included noninvasive ECG electrode leads affixed to the and chest and extremities, a pulse oximeter placed on the skin of the tail, a rectal thermometer, and sampling tubes for end tidal CO<sub>2</sub> measurement attached to the endotracheal tube. Arterial blood gases were also periodically evaluated pre- and postinjury. The pigs were continuously monitored and all data from physiologic monitoring were collected on a computer driven storage system. Intracranial pressure monitoring was not performed since previous studies demonstrated only small transient changes using the injury parameters applied in this study (20).

#### Brain Injury

Brain trauma was induced via head rotational acceleration as previously described in detail (20). Briefly, the head of each animal was secured to a padded snout clamp, which was mounted to the linkage assembly of pneumatic actuator device that converts the linear motion to an angular (rotational) motion. Rotation of the sidearm is triggered by the release of pressurized nitrogen into the actuator. For these experiments, the linkage was adjusted to produce a pure impulsive head rotation 110° in the coronal plane over a period of 20 ms, with the center of rotation close to the brain center of mass. Ten seconds prior to injury isoflurane anesthesia was withdrawn. The injury parameters were set to induce biphasic head rotational acceleration

with a predominant deceleration phase. Following injury the animals were released from the device. All animals received buprenorphine (0.1 mg/kg, i.m., q 12 h, p.r.n) for postoperative analgesia. It is important to note that previous studies with these techniques demonstrated that injured animals were awake and ambulatory within 8 hours of injury (20).

#### Histopathology

At 1-10 days after brain injury the animals received an overdose of pentobarbital (150 mg/kg, i.v.) and were transcardially perfused with saline following by 4% paraformaldehyde ( $n = 3$ , sham (no injury);  $n = 3$ , 1 day;  $n = 3$ , 3 days;  $n = 6$ , 7 days;  $n = 3$ , 10 days). The brains were removed, postfixated in 4% paraformaldehyde and stored in phosphate buffer saline and cryoprotected with sucrose. Subsequently, the brains were blocked into 0.5 cm coronal sections for gross examination and photography. A series of 40- $\mu$ m-frozen sections were cut from the front face of each block and mounted on microscope slides. Some blocks were cut into 3-5-mm-thick blocks and processed for paraffin embedding in an automated tissue processor (Shandon Hypercenter XP; Shandon Scientific Instruments, Cheshire, UK). Serial 6  $\mu$ m sections from these blocks were cut on a rotary microtome and mounted on poly-L-lysine-coated slides. Primary antibodies specific for NF proteins,  $\beta$ PP, A $\beta$ , and normal tau as well as paired helical filament (PHF) tau in AD NFTs used in these studies are outlined in the Table. Immunostaining was performed on free floating and paraffin-embedded sections using an avidin-biotin-immunoperoxidase complex method. The sections were incubated with primary antibody overnight at 4°C and then incubated at room temperature for 1 hour each with the appropriate secondary and tertiary antibodies, followed by enzymatic development with 3,3'-diaminobenzidine. Omission of the primary antibody or application of control serum on adjacent sections provided a negative control. Double labeling was performed with fluorescein isothiocyanate (FITC) and Texas red fluorescent secondary antibodies. Positive controls for NFTs and A $\beta$  plaques were performed on human AD tissue and run in parallel with the pig tissue.

## RESULTS

### Physiology and Behavior

Consistent with previous reports, immediately following trauma, no substantial changes in arterial blood gases, pulse oximetry, or end tidal CO<sub>2</sub> were observed following injury. All animals began to awaken within 15 min following injury. Although the animals were able to ambulate typically within 1 hour following injury, they appeared to have slightly sluggish responses to sensory stimuli (startle reflex, tactile response) for up to 8 hours post-trauma. However, by 24 hours postinjury, all of the animals appeared completely normal based on gross neurosensory examination (normal startle reflexes, gait, rooting behavior, eating, and drinking).

### Axonal and Neuronal Soma Pathology


Consistent with previous findings, axonal bulbs and varicose axonal swellings were observed following trauma at all timepoints evaluated (1 day–10 days) (Fig. 1). These axonal pathologies were identified by antibodies targeting  $\beta$ PP (Karen, 369W, 22C11) and NF proteins (NR4, N52). Axonal pathology was widespread throughout the brain and found in combination with gliosis most commonly in the root of gyri and at the interface of the gray and white matter (data not shown). No tissue tears and almost no vascular disruption were noted in regions of axonal injury. Modest neuronal damage was primarily found in the CA1 and CA3 subfields of the hippocampus as evidenced by pyknotic neurons and a general thinning of the pyramidal cell layers. No overt neuronal damage was observed in the cortex.

### Accumulation of A $\beta$ and Tau in Axonal Bulbs

At 3–10 days post-trauma we found that A $\beta$  and tau accumulated in most axonal bulbs found throughout the brains, demonstrated by all specific antibodies utilized (A $\beta$  immunostains: 2332, 13335, BCO5, 4G8, 6E10 and 10A5; Tau immunostains: Tau-2, PHF-1, PHF-6 and PHF-13.5) (Fig. 2). Positive staining with antibodies specific for PHF tau suggests that highly phosphorylated tau (like that in AD NFTs) accumulated in these brains. No A $\beta$  or tau accumulations in axons were detected at 1 day post-trauma. Colocalization of NF proteins,  $\beta$ PP, A $\beta$ , and tau was found in most, but not all, axonal bulbs demonstrated by multiple immunostains with fluorescence microscopy (Fig. 3A–H). However, no A $\beta$  or tau was found accumulating in axonal swellings (Fig. 3I, J). Therefore, A $\beta$  and tau accumulation was limited to the terminal ends of disconnected axons.


### A $\beta$ in Plaques

At 3–10 days postinjury, we found A $\beta$  containing plaque-like profiles in the pig brain tissue (stained with antibodies 2332, 13335 and BCO5) (Fig. 4). These



**Fig. 1.** Representative photomicrographs of pig brain sections demonstrating axonal pathologies following brain trauma. Numerous axonal bulbs were found in the subcortical white matter in sections stained with antibodies against  $\beta$ PP (A) and NF-H (B). NF-H immunostains also revealed varicose swellings of axons in the deep white matter (C). Scale bar = 50  $\mu$ m.

plaques were predominantly found in white matter with axonal pathology as well as in layer III of the cortex, and these findings were confirmed by positive staining in identical regions of adjacent thin (6  $\mu$ m) sections. However, the pig A $\beta$  plaques did not stain as robustly as plaques from positive control AD brains, and they were not very numerous (the most plaques found in any section was 10). In addition A $\beta$  containing plaques were only found in approximately one third of the injured animals, which also represented the group with the highest total amount of axonal pathology.




**Fig. 2.** Representative photomicrographs demonstrating A $\beta$  and tau accumulation in axonal bulbs of brain injured pigs. A $\beta$  was identified with several specific antibodies, including 10 $\Delta$ 5 (A), 13335 (B), 2332 (C), 4G8 (D), 6E10 (E), and BCO5 (F). Tau staining was also found with several specific antibodies including PHF-1 (G), PHF-13.5 (H), and PHF-6 (I). Scale bar = 50  $\mu$ m.

#### Tau Accumulation in Neurons

Accumulation of tau was found in the cytoplasm of neurons throughout the frontal, parietal, and temporal cortices at 3–10 days after brain injury (identified with antibodies; Tau-2, PHF-tau, PHF-1, PHF-6, and PHF-13.5) (Fig. 5C–E). The frequency and prevalence of this staining also appeared to correspond with the relative extent of axonal pathology. Our AD brain sections also demonstrated cytoplasmic staining with the same set of antibodies (Fig. 5A, B).

#### Neurofilament-rich Inclusions in Neurons

At 3–10 days post-trauma we found NF immunoreactive inclusions in the cytoplasm of neurons in the parietal and temporal cortices. These inclusions had a highly dense core surrounded by cytoplasm similar to the Lewy body (LB) inclusions found in human neurodegenerative diseases such as dementia with LBs (DLB) and Parkinson disease (PD) (Fig. 6A–G). Strong NF immunostaining of these inclusions was found in adjacent thin sections (6  $\mu$ m). However, we only found



**Fig. 3.** Representative photomicrographs demonstrating coaccumulation of A $\beta$ ,  $\beta$ PP, tau, and neurofilament in axonal bulbous following brain injury in the pig. Double immunolabeling with fluorescent staining demonstrated colocalization of A $\beta$  (A) with NF-H (B). Tau (C) with A $\beta$  (D), A $\beta$  (E) with  $\beta$ PP (F), and  $\beta$ PP (G) with tau (H). In another type of axonal pathology, varicose swellings, typical  $\beta$ PP accumulation was found, demonstrated by light microscopy (I). However, in the same section, A $\beta$  staining was not found in varicose axonal swellings, demonstrated by fluorescence microscopy (J). Scale bar = 50  $\mu$ m.

this pathology in the same subset of animals that demonstrated A $\beta$  plaque-like profiles (i.e. approximately one third of the injured animals).

#### DISCUSSION


In this study we found that inertial brain trauma in the pig produced diffuse axonal pathology in combination with several unique pathologic features that may be suggestive of neurodegenerative processes. The most remarkable and consistent finding was extensive A $\beta$  and tau accumulation in damaged axons following trauma. In addition, in a subset of brain injured animals, diffuse A $\beta$ -containing plaque-like profiles were found in both the gray and white matter, and accumulations of tau and NF-rich inclusions were observed in neuronal perikarya. To our knowledge, this is the first report of these collective findings in an animal model of brain trauma.

The observation of widespread A $\beta$  accumulation in damaged axons following inertial brain injury in the

pig may have important implications. It has previously been suggested that aberrant conversion of  $\beta$ PP to A $\beta$  at synapses may play a critical role in the evolution of AD. This same aberrant processing of  $\beta$ PP has been proposed to be initiated by brain trauma in humans, leading to the formation of diffuse A $\beta$  plaques in the gray matter within days following injury (5, 6). However, the most abundant accumulations of  $\beta$ PP resulting from brain trauma are found in damaged axons in the white matter (6, 17). Nonetheless, previous studies have not identified colocalization of A $\beta$  with the axonal pool of  $\beta$ PP in studies of brain injured humans or in rodent models of brain trauma (14–16, 18). The ability to identify A $\beta$  in damaged axons in the present study may reflect the use of highly specific A $\beta$  antibodies in conjunction with a gyrencephalic animal model of diffuse axonal pathology.

It is important to consider that this axonal pool of A $\beta$  may be released into the surrounding tissue from lysis or






**Fig. 4.** Representative photomicrographs of tissue sections immunostained for A $\beta$  demonstrating plaque formation. Typical A $\beta$  plaques in the gray matter of an Alzheimer disease brain (A and B). A $\beta$  plaques in pig brain following trauma in the gray matter (C and D), and white matter (E and F). Note the axonal bulb staining for A $\beta$  in proximity to a plaque (arrow, E). Scale bar = 50  $\mu$ m.

leakage of axonal bulbs. While A $\beta$  alone has not been shown to be substantially toxic in vivo (23, 24), we have previously proposed a "two hit" hypothesis whereby A $\beta$  may potentiate damage when combined with brain injury. Using transgenic mice that overexpress mutant human  $\beta$ PP and eventually develop A $\beta$  plaques (25), we found that brain trauma at an age prior to plaque formation induced massive neuron death accompanied by a marked increase in soluble A $\beta$  peptide levels (19). Moreover, a recent report has also described a large increase in A $\beta$

peptides in the cerebrospinal fluid of brain injured patients (26). Our present results suggest that damaged axons are one potential source for a massive increase in soluble A $\beta$  following brain trauma. Collectively, these data provide corroborative evidence that the production and release of A $\beta$  plays a role in the delayed pathogenesis of brain trauma.

The colocalization of  $\beta$ PP and A $\beta$  in damaged axons in the present study appeared to be limited to a specific morphologic subtype of axonal pathology. The major




**Fig. 5.** Representative photomicrographs of tau staining in neurons. Typical pathology of tau staining in neurons in Alzheimer disease brains (A and B). Neuronal staining with various specific antibodies targeting tau is also found in brain injured pigs (C–G). Scale bar = 50  $\mu$ m.

morphologic characteristics or “phenotypes” of post-traumatic axonal pathology include 1) varicose swellings encompassing long regions of injured axons, and 2) discrete axonal bulbs (also referred to as retraction balls and terminal clubs), characterized by individual rounded swellings at the terminal end of disconnected axons (20, 27). In the present study,  $\beta$ PP accumulated in both discrete axonal bulbs and in elongated varicose axonal swellings consistent with previous observations (15–17). However, A $\beta$  was only observed accumulating in axonal bulbs, i.e. only in axon regions proximal to clearly identified axotomy. These data suggest that axotomy induces a unique intra-axonal proteolytic milieu that favors the production and accumulation of A $\beta$  in axonal bulbs. Conversely, this process does not appear to occur in damaged

yet still connected axons, despite substantial swelling and  $\beta$ PP accumulation.

In addition to the widespread accumulation of A $\beta$  in axonal bulbs, we also found a limited number of A $\beta$  containing diffuse plaque-like profiles at 3–10 days following brain injury in the pig. These were found on adjacent sections and were identified with several highly specific anti-A $\beta$  antibodies. Although this finding may appear novel, it should be emphasized that the plaque-like profiles were relatively few in number and were primarily identified in the white matter, a location inconsistent with the distribution of A $\beta$  plaques described in recent studies of brain injured humans (5, 6). Nonetheless, the white matter location of the A $\beta$  plaques in brain injured pigs may reflect the release of A $\beta$  from damaged axons and



**Fig. 6.** Representative photomicrographs of neurofilament protein accumulation in neurons in brain injured pigs. Neurofilament immunoreactivity demonstrates neurofilament inclusions in neurons, seen as darkly stained profiles in the cytoplasm. Scale bar = 50  $\mu$ m.

subsequent extracellular aggregation in the same region. Furthermore, the formation of A $\beta$  plaques appeared to be related to the extent of axonal pathology.

Based on the identification of A $\beta$  plaques in brain injured humans, we and others have previously attempted to elucidate potential A $\beta$  plaque formation in rodent models of focal brain trauma without success (14–16, 18, 19). We did not even find acceleration or augmentation of A $\beta$  plaque formation following brain trauma in transgenic mice that otherwise go on to develop A $\beta$  plaques (19, 28). It is not presently clear whether the inability to replicate the human condition of A $\beta$  plaque formation in

rodent models of brain trauma is due to species effect or mechanisms of injury. While results from the present study may suggest that A $\beta$  plaques are produced following inertial diffuse brain injury in the pig, further studies are needed to confirm this potentially important finding.

Another potential link between neurodegenerative changes and diffuse brain injury in the pig is the accumulation of the microtubule-associated protein, tau, in damaged axons. Highly phosphorylated tau is the primary constituent of PHFs that form NFTs, one of the two major pathologic features of AD (29–31). Since tau is an integral structural protein in axons, it has been presumed



that it would accumulate in damaged axons with impaired transport. However, axonal accumulation of tau has not been previously observed following trauma despite several investigative efforts (32, 33). The colocalization of tau with A $\beta$  in damaged axons observed in the present study may have important implications since it is has been suggested that tau potentiates A $\beta$  toxicity in vitro and facilitates the polymerization of A $\beta$  peptides that may lead to A $\beta$  plaque formation (34, 35).

The accumulation of tau was also found in the cytoplasm of neurons of brain injured pigs in the present study. Since these profiles were identified with antibodies that recognize the highly phosphorylated forms of tau that form AD PHTs, these profiles most closely resemble the so-called "pre-tangle" somatodendritic tau lesions seen in AD brains. Indeed, it will be important to perform ultrastructural analyses of these lesions in future studies to determine if tau filaments are present in these perikaryal tau accumulations. Recently, neuronal staining for tau has also been found following brain injury in the rat (36). These findings may have important clinical implications since NFTs have been found in the brains of boxers, but no NFT-like lesions have been detected in the human brain following a single incident of brain trauma (3, 4).

Yet another unexpected finding in this study was that NF proteins, the building blocks of NFs, formed inclusion bodies in neurons following brain injury. While cytoplasmic NF rich inclusions, known as LBs, are signature lesions of DLB and PD, they have not previously been reported following trauma in humans. Nonetheless, accumulations of NFs are well documented in damaged axons following brain trauma in humans and experimental animals (20, 27, 37, 38). NF proteins are components of LBs, but alpha-synuclein may be the major building block of these lesions in PD and DLB (39). In addition, LBs are very common in the AD brain (40, 41). However, the mechanisms of development and the role of NF protein inclusions in neurodegenerative diseases have yet to be elucidated following brain trauma. Recent studies have also shown accumulation of NF protein in neuronal perikarya following trauma in a nontransgenic animal model of brain injury (42) and that LB-like inclusions may render neurons more vulnerable to degenerate following brain trauma in a transgenic mouse that overexpresses a NF hybrid protein (43). Since the predominant pathology in the pig diffuse brain injury model is damage to axons, not cortical neurons, our finding of cytoplasmic NF inclusions suggests that impaired axonal transport may play a role in the perikaryal accumulation of NF proteins.

Taken together, the results from the present study demonstrate that several pathologic characteristics of neurodegenerative diseases may also be found following diffuse brain injury in the pig. Accordingly, our results support the proposed link between brain trauma and the initiation of neurodegenerative processes. It is not clear

if the markers of neurodegenerative changes found accompanying diffuse axonal pathology in our pig model are consequences of injury specific mechanisms (i.e. inertial brain injury), or a general response of a gyrencephalic brain to trauma. Nonetheless, mechanisms of trauma-induced neurodegenerative processes may be further explored using this unique model.

## ACKNOWLEDGMENTS

We would like to thank Jeanne Marks for her excellent preparation of this manuscript.

## REFERENCES

1. Sosin DM, Sacks JJ, Smith SM. Head injury-associated deaths in the United States from 1979 to 1986. *JAMA* 1989;262:2251-55
2. Sosin DM, Sniezek JE, Waxweiler RJ. Trends in death associated with traumatic brain injury, 1979 through 1992. *JAMA* 1995;273:1778-80
3. Tokuda T, Ikeda S, Yanagisawa N, Ihara Y, Glenner GG. Re-amination of ex-boxers' brains using immunohistochemistry with antibodies to amyloid  $\beta$ -protein and tau protein. *Acta Neuropathol* 1991;82:281-85
4. Roberts GW, Allsop D, Bruton C. The occult aftermath of boxing. *J Neurol Neurosurg Psychiatry* 1990;53:373-78
5. Roberts GW, Gentleman SM, Lynch A, Graham DI.  $\beta$ A4 amyloid protein deposition in brain after head trauma. *Lancet* 1991;338:1422-23
6. Graham DI, Gentleman SM, Lynch A, Roberts GW. Distribution of  $\beta$ -amyloid protein in the brain following severe head injury. *Neuropath Appl Neurobiol* 1995;21:27-34
7. Stuss DT, Stehlem LL, Picton TW, Leech EE, Peichat G. Traumatic brain injury, aging and reaction time. *Can J Neurol Sci* 1989;16:161-67
8. Corkin S, Rosen JT, Sullivan EV, Clegg RA. Penetrating head injury in young adulthood exacerbates cognitive decline in later years. *J Neurosci* 1989;9:3876-83
9. Mortimer JA, Van Duijn CM, Chandra L, et al. Head trauma as a risk factor for Alzheimer's disease: A collaborative re-analysis of case-control studies. *Int J Epidemiol* 1991;20:S28-S35
10. Rasmussen DX, Brandt J, Martin DB, Folstein MF. Head injury as a risk factor in Alzheimer's disease. *Brain Inj* 1995;3:213-19
11. Mayeux R, Ottman R, Maestre G, et al. Synergistic effects of traumatic head injury and apolipoprotein- $\epsilon$ 4 in patients with Alzheimer's disease. *Neurology* 1995;45:555-57
12. Smith DH, Chen X-H, Pierce JES, et al. Progressive atrophy and neuronal death for one year following brain trauma in the rat. *J Neurotrauma* 1997;14(10):715-27
13. Bramlett HM, Dietrich WD, Green EJ, Busto R. Chronic histopathological consequences of fluid-percussion brain injury in rats: Effects of post-traumatic hypothermia. *Acta Neuropathol* 1997;93:190-99
14. Murali H, Pierce JES, Raghupathi R, et al. Two-fold overexpression of human  $\beta$ -amyloid precursor proteins in transgenic mice does not affect the neuromotor, cognitive, or neurodegenerative sequelae following experimental brain injury. *J Comp Neurol* 1998;392:428-38
15. Pierce JES, Trojanowski JQ, Graham DI, Smith DH, McIntosh TK. Immunohistochemical characterization of alterations in the distribution of amyloid precursor proteins and amyloid  $\beta$  peptide following experimental brain injury in the rat. *J Neurosci* 1996;16:1063-90
16. Lewen A, Li GL, Nilsson P, Olsson P, Hillered L. Traumatic brain injury in rat produces changes of  $\beta$ -amyloid precursor protein immunoreactivity. *NeuroReport* 1995;6:357-60

17. Sherriff FE, Bridges LR, Sivaloganathan S. Early detection of axonal injury after human head trauma using immunocytochemistry for  $\beta$ -amyloid precursor protein. *Acta Neuropathol* 1994;87:55–62
18. Bramlett HM, Kraydieh S, Green EJ, Dietrich WD. Temporal and regional patterns of axonal damage following traumatic brain injury: A beta-amyloid precursor protein immunocytochemical study in rats. *J Neuropathol Exp Neurol* 1997;56:1132–41
19. Smith DH, Nakamura M, McIntosh TK, et al. Brain trauma induces massive hippocampal neuron death linked to a surge in  $\beta$ -amyloid levels in mice overexpressing mutant amyloid precursor protein. *Am J Pathol* 1998;153:1005–110
20. Smith DH, Chen X-H, Xu B-N, McIntosh TK, Gennarelli TA, Meade DF. Characterization of diffuse axonal pathology and selective hippocampal damage following inertial brain trauma in the pig. *J Neuropathol Exp Neurol* 1997;56:822–34
21. Kimura H, Meaney DF, McGowan JC, et al. Magnetization transfer imaging of diffuse axonal injury following experimental brain injury in the pig: Characterization by magnetization transfer ratio with histopathologic correlation. *J Computer Assisted Tomography* 1996;20:540–46
22. Meaney DF, Thibault LE, Smith DH, Ross DT, Gennarelli TA. Diffuse axonal injury in the miniature pig: Biomechanical development and injury threshold. *ASME WAM* 1993;25:169–75
23. Stephenson DT, Clemens JA. In vivo effects of beta-amyloid implants in rodents: Lack of potentiation of damage associated with transient global forebrain ischemia. *Brain Res* 1992;586:235–46
24. Games D, Khan KM, Soriano FG, et al. Lack of Alzheimer pathology after  $\beta$ -amyloid protein injections in rat brain. *Neurobiol Aging* 1992;13:569–76
25. Games D, Adams D, Alessandrini R, et al. Alzheimer-type neuropathology in transgenic mice overexpressing V717F  $\beta$ -amyloid precursor protein. *Nature* 1995;373:523–27
26. Raby CA, Morganti-Kossmann MC, Kossmann T, et al. Traumatic brain injury increases beta-amyloid peptide 1–42 in cerebrospinal fluid. *J Neurochem* 1998;71:2505–9
27. Chen X-H, Xu B-N, Meaney DF, et al. Evolution of neurofilament subtype accumulation in axons following diffuse brain injury in the pig. *J Neuropathol Exp Neurol* 1999;58:588–96
28. Nakagawa Y, Nakamura M, McIntosh TK, et al. Traumatic brain injury in young PDAPP transgenic mice induces marked ipsilateral hippocampal atrophy and diminished A $\beta$  deposition during aging. *J Comp Neurol* 1999 (in press)
29. Wisniewski H, Terry RD, Hirano A. Neurofibrillary pathology. *J Neuropathol Exp Neurol* 1970;29:163–76
30. Kidd M. Paired helical filaments in electron microscopy of Alzheimer's disease. *Nature* 1963;197:192–93
31. Clark CM, Ewbank D, Lee VM-Y, Trojanowski JQ. Molecular pathology of Alzheimer's disease: Neuronal cytoskeletal abnormalities. In: Growdon J, Rossor M, eds. *Blue books of practical neurobiology: The dementias*. Boston: Butterworth Heinemann, 1998: 285–304
32. Sherriff FE, Bridges LR, Gentleman SM, Sivaloganathan S, Wilson S. Markers of axonal injury in post mortem human brain. *J Neuropath* 1994;88:433–39
33. Gultekin SH, Smith TW. Diffuse axonal injury in craniocerebral trauma. A comparative histologic and immunohistochemical study. *Arch Pathol Lab Med* 1994;118:168–71
34. Smith MA, Siedlak SL, Richey PL, et al. Tau protein directly interacts with the amyloid  $\beta$ -protein precursor: Implications for Alzheimer's disease. *Nature Med* 1995;1:365–69
35. Caputo CB, Sygowski LA, Scott CW, Sobel IRE. Role of tau in the polymerization of peptides from  $\beta$ -amyloid precursor protein. *Brain Res* 1992;597:227–32
36. Hoshino S, Tamaoka A, Takahashi M, et al. Emergence of immunoreactivities for phosphorylated tau and amyloid-beta protein in chronic stage of fluid percussion in rat brain. *NeuroReport* 1998;9: 1879–83
37. Grady MS, McLaughlin MR, Christman CW, Valadka AB, Fligner CL, Povlishock JT. The use of antibodies targeted against the neurofilament subunits for the detection of diffuse axonal injury in humans. *J Neuropathol Exp Neurol* 1993;52:143–52
38. Povlishock JT, Christman CW. Diffuse axonal injury. In: Waxman SC, Kocsis JD, Stys PK, eds. *The axon: Structure, function and pathophysiology*. New York: Oxford University Press, 1995:504–29
39. Trojanowski JQ, Goedert M, Iwatsubo T, Lee VM-Y. Fatal attractions: Abnormal protein aggregation and neuron death in Parkinson's disease and Lewy body dementia. *Cell Death Differentiation* 1998;5:832–37
40. Goldman JE, Yen SH. Cytoskeletal protein abnormalities in neurodegenerative diseases. *Ann Neurol* 1986;19:209–23
41. Trojanowski JQ, Lee VMY. Phosphorylation of neuronal cytoskeletal proteins in Alzheimer's disease and Lewy body dementia. *Ann NY Acad Sci* 1994;747:92–109
42. Kanayama G, Takeda M, Morihara T, et al. Temporal and regional profiles of cytoskeletal protein accumulation in the rat brain following traumatic brain injury. *Psychiatry Clin Neurosci* 1997;51:157–65
43. Nakamura M, Santman KE, Galvin JE, et al. Increased vulnerability of NFH-LacZ transgenic mouse to traumatic brain injury-induced behavioral deficits and cell loss. *J Cereb Blood Flow Metab* 1999;762–70
44. Buxbaum JD, Gandy SE, Cicchetti P, et al. Processing of Alzheimer  $\beta$ /A4 amyloid precursor protein: Modulation by agents that regulate protein phosphorylation. *Proc Natl Acad Sci* 1990;87:6003–6
45. Gandy SE, Bhusin R, Ramabhadran TV, et al. Alzheimer  $\beta$ /A4-amyloid precursor protein: Evidence for putative amyloidogenic fragment. *J Neurochem* 1992;58:383–86
46. Kang J, Lemaire H-G, Unterbeck A, et al. The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature* 1987;325:733–36
47. Hilbich C, Monning U, Grund C, Masters CL, Beyreuther K. Amyloid-like properties of flanking the epitope of amyloid precursor protein-specific monoclonal antibody 22c11. *J Biol Chem* 1993;268:26571–77
48. Weidemann A, König G, Bunke D, et al. Identification, biogenesis, and localization of precursors of Alzheimer's disease A4 amyloid protein. *Cell* 1989;57:115–26
49. Pierce JES, Smith DH, Trojanowski JQ, McIntosh TK. Enduring cognitive, neurobehavioral, and histopathological changes persist for up to one year following severe experimental brain injury in rats. *Neurosci* 1998;87:359–69
50. Cai X-D, Golde TE, Youngkin SG. Release of excess amyloid  $\beta$  protein from a mutant amyloid  $\beta$  protein precursor. *Science* 1993;259:514–16
51. Szendrei GI, Prammer KV, Vaski M, Lee VMY, Olivos L, Jr. The effects of aspartic acid-bond isomerization on in vitro properties of the amyloid beta peptide as modeled with N-terminal decapeptide fragments. *Int J Pept Protein Res* 1996;47:289–96
52. Schmidt ML, DiDario AG, Olivos L, Jr, et al. Plaque-associated neuronal proteins: A recurrent motif in neuritic amyloid deposits throughout diverse cortical areas of the Alzheimer's disease brain. *Exp Neurol* 1994;130:311–22
53. Iwatsubo T, Odaka A, Suzuki N, Mizusawa H, Nukina N, Ihara Y. Visualization of A $\beta$ 42(43) and A $\beta$ 40 in senile plaques with end-specific A $\beta$  monoclonals: Evidence that an initially deposited species is A $\beta$ 42(43). *Neuron* 1994;13:45–53
54. Suzuki N, Cheung TT, Cai X-D, et al. An increased percentage of long amyloid  $\beta$  protein secreted by familial amyloid  $\beta$  protein precursor (BAPP<sub>717</sub>) mutants. *Science* 1994;264:1336–40
55. Prittila T, Kim KS, Mehta PD, Frey H, Wisniewski HM. Soluble amyloid beta-protein in the cerebrospinal fluid from patients with

- Alzheimer's disease, vascular dementia and controls. *J Neurol Sci* 1994;127:90-95
56. Kim KS, Miller DL, Sapienza VJ, et al. Production and characterization of monoclonal antibodies reactive to synthetic cerebrovascular amyloid peptide. *Neurosci Res Comm* 1988;2:121-30
  57. Kim KS, Wen GY, Bancher C, et al. Detection and quantitation of amyloid  $\beta$ -peptide with 2 monoclonal antibodies. *Neurosci Res Comm* 1990;7:113-22
  58. Hyman BT, Tanzi RE, Marzloff K, Barbour R, Schenk D, Kunitz protease inhibitor-containing amyloid beta protein precursor immunoreactivity in Alzheimer's disease. *J Neuropathol Exp Neurol* 1992;51:76-83
  59. Frenkel D, Balass M, Solomon B. N-terminal EFRH sequences of Alzheimer's beta-amyloid peptide represents the epitope of its anti-aggregating antibodies. *J Neuroimmunol* 1998;88:85-90
  60. Papasozomenos SC, Binder LI. Phosphorylation determines two distinct species of tau in the central nervous system. *Cell Motil Cytoskeleton* 1987;8:210-26
  61. Papasozomenos SC. Tau protein immunoreactivity in dementia of the Alzheimer's type I. Morphology, evolution, distribution and pathogenic implication. *Lab Invest* 1989;60:123-37
  62. Papasozomenos SC. Tau protein immunoreactivity in dementia of the Alzheimer's type II. Electron microscopy and pathogenic implications. Effects of fixation on the morphology of the Alzheimer's abnormal filaments. *Lab Invest* 1989;60:375-89
  63. Greenberg SG, Davies P. A preparation of Alzheimer-paired helical filaments that displays distinct tau proteins by polyacrylamide gel electrophoresis. *Proc Natl Acad Sci USA* 1990;87:5827-31
  64. Bramblett GT, Goedert M, Jakes R, Merrick SE, Trojanowski JQ, Lee VMY. The abnormal phosphorylation of Tau at Ser<sup>36</sup> in Alzheimer's disease recapitulates phosphorylation during development and contributes to reduced microtubule binding. *Neuron* 1993;10:1089-99
  65. Greenberg SG, Davies P, Schein JD, Binder LI. Hydrofluoric acid-treated tau PHF proteins display the same biochemical properties as normal tau. *J Biol Chem* 1992;267:564-69
  66. Lang E, Szendrei GI, Lee VMY, Otvos L. Immunological and conformational characterization of a phosphorylated immunodominant epitope on the paired helical filaments found in Alzheimer's disease. *Biochem Biophys Res Commun* 1992;187:783-90
  67. Mawal-Dewan MY, Henley J, Van de Voorde A, Trojanowski JQ, Lee VM. The phosphorylation state of tau in the developing rat brain is regulated by phosphoprotein phosphatases. *J Biol Chem* 1994;269:30981-87
  68. Reed LA, Schmidt ML, Wszolek ZK, et al. The neuropathology of a chromosome 17-linked autosomal dominant Parkinsonism and dementia ("pallido-ponto-nigral degeneration"). *J Neuropathol Exp Neurol* 1998;57:588-601
  69. Hoffmann R, Lee VM, Leight S, Varga I, Otvos L, Jr. Unique Alzheimer's disease paired helical filament specific epitopes involve double phosphorylation at specific sites. *Biochem* 1997;36:8114-24
  70. Shaw G, Osborn M, Weber K. Reactivity of a panel of neurofilament antibodies on phosphorylated and dephosphorylated neurofilaments. *Eur J Cell Biol* 1986;42:1-9
  71. Debus E, Weber K, Osborn M. Monoclonal antibodies specific for glial fibrillary acidic (GFA) protein and for each of the neurofilament triplet polypeptides. Differentiation 1983;25:193-203
  72. Shaw G, Winiński L, Reier P. The effect of axotomy and deafferentation on phosphorylation-dependent antigenicity of neurofilaments in rat superior cervical ganglion neurons. *Brain Res* 1988;460:227-34
  73. Gotow T, Tanaka J. Phosphorylation of neurofilament H subunit as related to arrangement of neurofilaments. *J Neurosci Res* 1994;37:673
  74. Harris J, Ayyub C, Shaw G. A molecular dissection of the carboxy-terminal tails of the major neurofilament subunits NF-H. *J Neurosci Res* 1991;30:47-62
  75. Jicha GA, Bowser R, Kazam IG, Davies P. Alz-50 and MC-1, a new monoclonal antibody raised to paired helical filaments, recognize conformational epitopes on recombinant tau. *J Neurosci Res* 1997;48:128-32

Received March 10, 1999

Revision received April 30, 1999

Accepted May 4, 1999