

RESEARCH PAPER

Activation of endothelial and epithelial $K_{Ca}2.3$ calcium-activated potassium channels by NS309 relaxes human small pulmonary arteries and bronchioles

Christel Kroigaard¹, Thomas Dalsgaard¹, Gorm Nielsen²,
Britt E Laursen¹, Hans Pilegaard³, Ralf Köhler^{2,4} and Ulf Simonsen¹

¹Department of Biomedicine, Pulmonary and Cardiovascular Pharmacology, Faculty of Health Sciences, Aarhus University, Aarhus C, Denmark, ²Cardiovascular and Renal Research, Institute for Molecular Medicine, University of Southern Denmark, Odense, Denmark, ³Department of Thoracic Surgery, Aarhus University Hospital Skejby, Aarhus N, Denmark, and ⁴Aragon Institute of Health Sciences I+CS and ARAID, Unit for Translational Research, Hospital Miguel Servet, Zaragoza, Spain

Correspondence

Ulf Simonsen, Department of Biomedicine, Pulmonary and Cardiovascular Pharmacology, Faculty of Health Sciences, Aarhus University, Wilhelm Meyers Allé 4, 8000 Aarhus C, Denmark. E-mail: us@farm.au.dk

Keywords

calcium-activated potassium channels; pulmonary arteries; bronchioles; $K_{Ca}3.1$; $K_{Ca}2.3$; NS309

Received

20 January 2012

Revised

13 March 2012

Accepted

14 March 2012

BACKGROUND AND PURPOSE

Small ($K_{Ca}2$) and intermediate ($K_{Ca}3.1$) conductance calcium-activated potassium channels (K_{Ca}) may contribute to both epithelium- and endothelium-dependent relaxations, but this has not been established in human pulmonary arteries and bronchioles. Therefore, we investigated the expression of $K_{Ca}2.3$ and $K_{Ca}3.1$ channels, and hypothesized that activation of these channels would produce relaxation of human bronchioles and pulmonary arteries.

EXPERIMENTAL APPROACH

Channel expression and functional studies were conducted in human isolated small pulmonary arteries and bronchioles. $K_{Ca}2$ and $K_{Ca}3.1$ currents were examined in human small airways epithelial (HSAEpi) cells by whole-cell patch clamp techniques.

RESULTS

While $K_{Ca}2.3$ expression was similar, $K_{Ca}3.1$ protein was more highly expressed in pulmonary arteries than bronchioles. Immunoreactive $K_{Ca}2.3$ and $K_{Ca}3.1$ proteins were found in both endothelium and epithelium. K_{Ca} currents were present in HSAEpi cells and sensitive to the $K_{Ca}2.3$ blocker UCL1684 and the $K_{Ca}3.1$ blocker TRAM-34. In pulmonary arteries contracted by U46619 and in bronchioles contracted by histamine, the $K_{Ca}2.3$ / $K_{Ca}3.1$ activator, NS309, induced concentration-dependent relaxations. NS309 was equally potent in relaxing pulmonary arteries, but less potent in bronchioles, than salbutamol. NS309 relaxations were blocked by the $K_{Ca}2$ channel blocker apamin, while the $K_{Ca}3.1$ channel blocker, charybdotoxin failed to reduce relaxation to NS309 (0.01–1 μ M).

CONCLUSIONS AND IMPLICATIONS

$K_{Ca}2.3$ and $K_{Ca}3.1$ channels are expressed in the endothelium of human pulmonary arteries and epithelium of bronchioles. $K_{Ca}2.3$ channels contributed to endo- and epithelium-dependent relaxations suggesting that these channels are potential targets for treatment of pulmonary hypertension and chronic obstructive pulmonary disease.

Abbreviations

1-EBIO, 1-ethyl-2-benzimidazolinone; COPD, chronic obstructive pulmonary disease; CyPPA, cyclohexyl-[2-(3,5-dimethyl-pyrazol-1-yl)-6-methyl-pyrimidin-4-yl]-amine; EDHF, endothelium-derived hyperpolarizing factor; EET, epoxyeicosatrienoic acid; EpDHF, epithelium-derived hyperpolarizing factor; GEA 3175, 3-(2-chloro-3-methylphenyl)-5-[[4-methylphenyl]sulphonyl] amino]-hydroxide; K_{ATP} channel, ATP-sensitive potassium channel; $K_{Ca1.1}$ channel, large conductance calcium-activated potassium channel; K_{Ca2} channel, small conductance calcium-activated potassium channel; $K_{Ca3.1}$ channel, intermediate conductance calcium-activated potassium channel; NS309, 6,7-dichloro-1H-indole-2,3-dione 3-oxime; SKA-31, naphtho[1,2-d]thiazol-2-ylamine; TRAM-34, 1-[(2-chlorophenyl)diphenylmethyl]-1H-pyrazole; U46619, 9,11-dideoxy-9a,11a-epoxymethanoprostaglandin $F_{2\alpha}$; UCL1684, 6,12,19,20,25,26-hexahydro-5,27:13,18:21,24-trietheno-11,7-metheno-7H-dibenzo [*b,n*] [1,5,12,16]tetraazacyclotricosine-5,13-dium dibromide

Introduction

The respiratory epithelium and the pulmonary arterial endothelium release factors that are thought to be important in the regulation of pulmonary arterial pressure and bronchial tone (Flavahan *et al.*, 1985; Morrison *et al.*, 1990). In the pulmonary arterial system, endothelium-derived NO (Ignarro *et al.*, 1987; Feletou *et al.*, 1995), prostacyclin (Frantz *et al.*, 1989) and an endothelium-derived hyperpolarising factor (EDHF) (Feletou *et al.*, 1995; Zhang *et al.*, 2006) appear to be involved in regulating vascular tone. Activation of calcium-activated potassium channels of small conductance (K_{Ca2} , subtype $K_{Ca2.3}$) and intermediate conductance ($K_{Ca3.1}$) are required for initiation of EDHF-type relaxations in several vascular beds including human arteries (Edwards *et al.*, 1998; Buus *et al.*, 2000; Grgic *et al.*, 2009; Chadha *et al.*, 2011; channel nomenclature follows Alexander *et al.*, 2011). In the rat pulmonary circulation, we have recently found that the $K_{Ca3.1}$ channel and the Na^+/K^+ -ATPase are involved in this EDHF-type relaxation (Kroigaard *et al.*, 2010), but the contribution of $K_{Ca2.3}$ and $K_{Ca3.1}$ channels for EDHF-type relaxation in human pulmonary arteries is not known.

Even less is known about the contribution of $K_{Ca2.3}$ and $K_{Ca3.1}$ channels to epithelium-derived relaxation, and it is so far unknown whether an epithelium-derived hyperpolarizing factor (EpDHF) relaxation mechanism involving $K_{Ca2.3}$ and $K_{Ca3.1}$ channels exists. Since the pioneering work of Paul Vanhoutte and colleagues in the mid-eighties, it is well established that the epithelium can release factors producing bronchial dilation (Stuart-Smith and Vanhoutte, 1987; 1988; Vanhoutte, 1987; Morrison *et al.*, 1990). Indeed, removal of the epithelium in larger airways enhances the effect of contractile agonists and reduces the relaxing responses in smaller airways (Stuart-Smith and Vanhoutte, 1987; Kroigaard *et al.*, 2010). Both NO and PGE_2 serve as epithelium-derived relaxing factors in human airways (Folkerts and Nijkamp, 1998). In addition, epoxyeicosatrienoic acids were proposed to evoke non-NO, non-prostanoid EpDHF-type relaxation in guinea pig airways (Benoit *et al.*, 2001). Recently, we found that both $K_{Ca2.3}$ and $K_{Ca3.1}$ channels were involved in an EpDHF-type relaxation in rat bronchioles, also associated with activation of Na^+/K^+ -ATPase (Kroigaard *et al.*, 2010). However, it is unclear whether $K_{Ca2.3}$ and $K_{Ca3.1}$ channels are involved in EpDHF-type relaxation in human bronchioles.

The objective of this study was to investigate the expression, cellular localization and function of $K_{Ca2.3}$ and $K_{Ca3.1}$

channels in human pulmonary arteries and bronchioles. We hypothesized that activation of these calcium-activated potassium (K_{Ca}) channels by a $K_{Ca2.3}$ and $K_{Ca3.1}$ channel opener NS309 (6,7-dichloro-1H-indole-2,3-dione 3-oxime) (Strobaek *et al.*, 2004), leads to relaxation in human pulmonary arteries and bronchioles. To address the hypothesis, human small pulmonary arteries and bronchioles were obtained from patients undergoing lung surgery. Expression of $K_{Ca2.3}$ and $K_{Ca3.1}$ channels was examined by quantitative PCR and immunoblotting and cellular localization by immunohistochemistry. Functional expression of $K_{Ca2.3}$ and $K_{Ca3.1}$ channels was examined in human small airway epithelial (HSAEpi) cells by patch clamp techniques and use of a compound, naphtho[1,2-d]thiazol-2-ylamine (SKA-31), structurally related to NS309 (Sankaranarayanan *et al.*, 2009). Finally, the relaxant effects of NS309 were investigated in isolated human bronchioles and pulmonary arteries.

Methods

Human tissue

The research was carried out in accordance with the Declaration of Helsinki (2000) of the World Medical Association and the Local Ethics Committee approved the study (Permission number: 200440154). Each patient was informed about the purpose and nature of the project and gave informed consent. Experiments were performed with human pulmonary arteries and bronchiole branches from 30 patients, who had undergone surgery for lung carcinoma at the Department of Thoracic Surgery, Aarhus University Hospital, Skejby, Denmark. Mean age of the patients examined was 67 years (55–76 years), and all individuals were smokers or former smokers. Bronchioles and small pulmonary arteries were carefully dissected under a microscope by removing the surrounding tissue.

Cell culture

Human small airway epithelial cells (HSAEpi, Sciencell, third to sixth passage) were cultured in poly-L-lysine-coated flasks containing small airway epithelial cell medium (Sciencell), epithelial cell growth supplement (EpiCGS), penicillin and streptomycin. For patch-clamp experiments, cells were trypsinized and seeded on coverslips.

Quantitative PCR of *K_{Ca}2.3* and *K_{Ca}3.1* mRNA in human lung

Human intrapulmonary arteries and bronchioles were isolated in calcium free physiological saline solution (PSS) and transferred to RNA-later (Ambion, Foster City, CA). Purification of total RNA was achieved by using the RNeasy Mini Fibrous Tissue Kit from Qiagen (Hilden, Germany). Furthermore, removal of genomic DNA was achieved using a DNase I digestion step (Qiagen). The concentration of the RNA was estimated by OD measurements at 260 nm. Total RNA (100 µg·mL⁻¹) was reverse transcribed using Oligo-dT primer and Superscript™ SIII reverse transcriptase (Invitrogen, Hercules, CA, USA). Expression of *K_{Ca}* channels was assessed by TaqMan quantitative PCR (QPCR). 200 ng RNA was used in a 25 µL reaction and quantitative PCR conducted using Ex Taq™ (TaKaRa, Shiga, Japan). The following cycles were run: 1 cycle at 95°C for 2 min, 40 cycles at 95°C for 15 s, 55°C for 1 min and 70°C for 15 s on a Stratagene Mx3000P machine (La Jolla, CA). Primers were designed from the National Center for Biotechnical Information (NCBI) using PerlPrimer and ordered at MWG-Biotech AG (Ebersberg, Germany): *K_{Ca}2.3*, F: GATTGACCATGCCAAAGTGAG; *K_{Ca}2.3*, R: ACATGACATTCTGCATCTTGG; *K_{Ca}3.1*, F: GTTCTACAAACATAC TCGCAGGA; *K_{Ca}3.1*, R: GCGTGTCAATCTGTTTCTCAA. The fluorogenic probe contained a reporter dye, 6-carboxyfluorescein (FAM), at the 5'-end and a Blackhole Quencher 1 dye at the 3'-end: *K_{Ca}2.3*: TCCTCCAAGCTATCC ACCAGTTGAG; *K_{Ca}3.1*: TCAACGCGTCCGCCAGGTGCGG CTGAAA. The amount of cDNA was normalized to the reference gene by using the threshold Ct values: $\Delta Ct = Ct(\text{target}) - Ct(\text{GAPDH})$, and the results were analysed as a ratio to GAPDH expression: $\text{Ratio} = 2^{-\Delta Ct}$ and expressed as a percentage.

Reverse transcriptase-PCR of HSAEpi cells

Reverse transcriptase-PCR (RT-PCR) was done using RNeasy Mini Kit (Qiagen) for RNA purification, iScript cDNA Synthesis Kit (Bio-Rad, Hercules, CA, USA) for cDNA synthesis and Taq DNA polymerase (Invitrogen) for PCR. A standard PCR protocol was used with initial denaturing at 95°C for 3 min and subsequently 40 cycles of denaturing at 95°C for 25 s, annealing at 57°C for 20 s and extending at 72°C for 40 s. A final step of extension was done at 72°C for 3 min. PCR-products were analysed by gel electrophoresis using 1.5% agarose in TBE buffer and staining with GelRed™ (Biotium Hayward, CA, USA). Specific RT-PCR primers were designed to span intronic sequences. The expected product lengths were 159bp for *K_{Ca}3.1*, 333 bp for *K_{Ca}2.3*, 395 bp for *K_{Ca}2.2* and 391 bp for *K_{Ca}2.1*. Primer sequences: h*K_{Ca}3.1* F, 5'-CATCA CATTCTGACCATCG; h*K_{Ca}3.1* R, 5'-ACGTGCTTCTCTGCCT TGTT; h*K_{Ca}2.3* F, 5'-GCTCTCTTGGGGTTTGTACTCAA; h*K_{Ca}2.3* R, 5'-CGCAGGAACATGGGGATAGA; h*K_{Ca}2.2* F, 5'-TCTAAGC CCGAGCACAA; h*K_{Ca}2.2* R, 5'-TTGTCCACCATGAACA ACTGTA; h*K_{Ca}2.1* F, 5'-GGGCTCCAGGTGGTAGT; h*K_{Ca}2.1* R, 5'-CACCATGAACAGCTGGATCTC.

Immunohistochemistry of *K_{Ca}2.3* and *K_{Ca}3.1* protein in human lung

Evaluation of the distribution of *K_{Ca}3.1* and *K_{Ca}2.3* channels was achieved using immunohistochemistry with primary

antibodies against *K_{Ca}2.3* (1:200; Santa Cruz Biotechnology, Santa Cruz, CA) and *K_{Ca}3.1* (1:100 or 1:500; Cell Applications, San Diego, CA). Segments of isolated pulmonary arteries and bronchioles were fixed in 4% formalin. The tissue was processed through increasing alcohol percentages to paraffin wax blocks. Sections of 3 µm were cut with a microtome, collected on glass slides and heated in an oven (80°C). The sections were de-paraffinized, processed in decreasing alcohol percentages, transferred to distilled water, incubated with 0.3% hydrogen peroxide for 20 min and rinsed in PBS. Antigen retrieval was achieved by heating sections in a microwave in TEG-buffer for 2 × 5 min. They were then quenched in 0.2% Triton X in PBS for 15 min. Blocking of non-specific antibody binding was done with 10% fetal calf serum in 1% BSA in PBS for 20 min. Incubation with the primary antibody in PBS with 1% BSA was overnight at 4°C. To ensure that there was no non-specific staining of primary antibodies, sections were incubated with isotype-matched antibodies (rabbit IgG; Abcam, Cambridge, UK) instead of primary antibodies. The sections were rinsed in PBS and incubated for 1 h at room temperature with secondary antibody, goat anti-rabbit IgG coupled to HRP (1:2000; Invitrogen) in PBS containing 1% BSA followed by rinsing in PBS. Finally, DAB was applied for 5 min, the sections rinsed and for histology, and sections were stained with haematoxylin and 0.1% w/v eosin, and dehydrated in increasing concentrations of ethanol. Background staining was controlled by incubating sections only with secondary antibody. The sections were coverslipped and analysed by light microscopy.

Immunoblotting of *K_{Ca}2.3* and *K_{Ca}3.1* protein in human lung

Human pulmonary arteries and bronchioles were carefully dissected from the surrounding tissue, frozen in liquid nitrogen and transferred to -80°C. Protein was extracted in lysis buffer, and samples were placed in an ultrasonic bath for 45 s and centrifuged for 15 min at 13 250 g at 4°C. The supernatant was frozen at -80°C. Total protein was quantified using the Bio-Rad Protein Assay (Bio-Rad, Hercules, CA). Protein lysate was mixed with sample buffer and incubated at 99°C for 10 min. Samples and a pre-stain marker (Bio-Rad, Hercules, CA) were loaded onto a 4–12% Criterion XT Bis-Tris gel (Bio-Rad Hercules, CA) and separated by SDS-PAGE at 200 V. Proteins were transferred to membrane for 1 h at 100 V. The membrane was washed in TBS-T and blocked in 5% skimmed milk in TBS-T for 2 h. Incubation with primary antibody (β -actin (1:20 000), *K_{Ca}2.3* (1:200, Santa Cruz Biotechnology) and *K_{Ca}3.1* (1:200, Cell Applications, San Diego, CA) and in 5% skimmed milk in TBS-T was done overnight at 4°C. The membrane was washed in 5% skimmed milk in TBS-T before incubation with secondary antibody: goat anti-rabbit IgG conjugated to HRP (1:4000; Santa-Cruz Biotechnology), β -actin: goat anti-mouse IgG conjugated to HRP (1:4000; Sigma, St. Louis, MO). The membrane was developed by using the ECL-Plus kit (GE Healthcare, Copenhagen, Denmark). The blot was placed in an X-ray film cassette and developed in photographic developer. Amounts of *K_{Ca}2.3* and *K_{Ca}3.1* protein were normalized to β -actin, which was equally expressed in human pulmonary arteries and bronchioles.

Patch-clamp electrophysiology

Whole-cell membrane currents were recorded using an Axopatch patch-clamp amplifier (Axon Instruments, Foster City, CA) and the Clampex 9.2 data acquisition software and were analysed by the Clampfit 9.2 software. For activation of K_{Ca} currents, HSAEpi cells were dialysed with a KCl-pipette solution containing 3 μM $[\text{Ca}^{2+}]_{\text{free}}$ (in mM): 140 KCl, 1 Na_2ATP , 1 MgCl_2 , 2 EGTA, 1.91 CaCl_2 and 5 HEPES, pH 7.2. The NaCl bath solution contained (mM): 137 NaCl, 4.5 Na_2HPO_4 , 3 KCl, 1.5 KH_2PO_4 , 0.4 MgCl_2 , 0.7 CaCl_2 and 10 glucose (pH 7.4). For maximal and stable channel activation, the bath solution contained the $K_{Ca2}/K_{Ca3.1}$ channel opener SKA-31 (1 μM). For blocking experiments, the selective $K_{Ca2.3}$ channel blocker UCL1684 (1 μM) and the $K_{Ca3.1}$ channel blocker TRAM-34 (1 μM) were added to the bath. 1 mM stock solution of TRAM-34, UCL1684 and SKA-31 were prepared in DMSO. The final DMSO concentration did not exceed 0.3%.

Wire myography

For isometric tension recordings, small pulmonary arteries and bronchioles with a length of approximately 2 mm were mounted on two 40 μm steel wires in microvascular myographs (DMT, Aarhus, Denmark). The baths were heated to 37°C and equilibrated with 5% CO_2 to maintain the desired pH of 7.4. Segments were allowed to equilibrate for 10 min thereafter. For experiments on bronchioles, the organ bath contained calcium-free PSS during mounting and normalization to avoid development of spontaneous tension. For experiments on arteries, the organ bath contained normal PSS.

Segments were stretched to 2.4 kPa for optimal measurements, corresponding to a transmural pressure of 18 mmHg. The viability of the bronchial segments were examined by exposure to a potassium-rich PSS (60 mM KPSS). This was initially done twice and in the end of each experiment once. Viability of arterial segments was tested with a 124 mM KPSS. Preparations in which the final response to KPSS was not comparable to the two initial responses to KPSS were not further considered. After the initial test of viability, vessels were incubated for 30 min in PSS containing the COX inhibitor, indomethacin (3 μM), to block the synthesis of prostaglandins. Indomethacin remained present throughout the entire experiment. Arterial segments were pre-contracted with the thromboxane analogue, U46619 (9,11-dideoxy-9 α ,11 α -epoxymethanoprostaglandin $F_{2\alpha}$, 10 nM). After reaching a level of stable pre-contraction, we used acetylcholine (10 μM) to ensure preservation of endothelial function. Only segments exhibiting more than 50% relaxation to acetylcholine were included in the study. Bronchioles were pre-contracted with histamine (2 μM).

Concentration–response curves (0.001–10 μM) were constructed for the $K_{Ca2.3}$ and $K_{Ca3.1}$ channel opener NS309 (Neurosearch A/S, Ballerup, Denmark), the ‘classical’ $K_{Ca2.3}$ and $K_{Ca3.1}$ channel opener 1-ethyl-benzimidazolinone (1-EBIO; Sigma Aldrich, St. Louis, MO); a selective activator of K_{Ca2} channels, cyclohexyl-[2-(3,5-dimethyl-pyrazol-1-yl)-6-methyl-pyrimidin-4-yl]-amine (CyPPA); a NO donor, 3-(2-chloro-3-methylphenyl)-5-[[[4-methylphenyl]sulphonyl]amino]-hydroxide (GEA 3175; GEA A/S, Copenhagen, Denmark) and the β_2 -adrenoceptor agonist, salbutamol (Sigma Aldrich).

The involvement of $K_{Ca2.3}$ and $K_{Ca3.1}$ channels in NS309 relaxation was investigated by incubating the segments with a $K_{Ca2.3}$ channel blocker (apamin, 0.05 μM), a $K_{Ca3.1}$ channel blocker (charybdotoxin, 0.07 μM), charybdotoxin and apamin in combination and a large conductance calcium-activated potassium ($K_{Ca1.1}$) channel blocker (iberiotoxin 0.1 μM) 15 min before contraction and during construction of the concentration–response curves for NS309.

In preparations where the endothelium or epithelium was removed, this was achieved by rubbing the inner layer with a human hair. Endothelial function was evaluated by contracting the pulmonary arteries with U46619 (0.01 μM) followed by addition of ACh (10 μM). Endothelial removal was successful, when relaxation to ACh was below 10%. There is no functional test for evaluating the presence of bronchiolar epithelium; therefore, epithelial removal was evaluated by histology as previously described (Kroigaard *et al.*, 2010).

Data analysis

All pulmonary arteries or bronchioles originated from different patients, where n indicates the number of preparations (one preparation per patient) examined. The results are expressed as mean \pm SEM. The responses with or without blockers were analysed by two-way ANOVA. To compare differences in expression, one-way ANOVA was used. For cell culture studies, a paired Student's t -test was used. In all cases, a probability less than 5% ($P < 0.05$) was considered significant. For statistical analysis, we used the computer software Graph Pad Prism 5.02 (GraphPad Software, San Diego, CA).

Results

$K_{Ca3.1}$ and $K_{Ca2.3}$ channel expression in human pulmonary arteries and bronchioles

In pulmonary arteries and bronchioles, we detected considerable amounts of $K_{Ca3.1}$ mRNA by RT-PCR. Quantitative RT-PCR also revealed that mRNA expression levels were significantly lower in pulmonary arteries than in bronchioles (Figure 1A). Likewise, we detected $K_{Ca2.3}$ mRNA in pulmonary arteries and bronchioles, with expression levels being similar in both tissues (Figure 1B). If compared with $K_{Ca3.1}$ mRNA expression levels, $K_{Ca2.3}$ mRNA expression was found to be approximately 100 times higher in both tissues. Immunoblotting revealed bands for $K_{Ca2.3}$ protein at 70 kDa and $K_{Ca3.1}$ protein at 50 kDa (Figure S1). The $K_{Ca3.1}$ protein was more abundant in pulmonary arteries than in bronchioles (Figure 2A,C). This finding contrasted with the lower levels of mRNA for $K_{Ca3.1}$ protein found in pulmonary arteries and the higher mRNA expression in bronchioles. $K_{Ca2.3}$ protein levels were similar in pulmonary arteries and bronchioles (Figure 2B,D). Moreover, $K_{Ca2.3}$ and $K_{Ca3.1}$ protein levels were similar. These data showed $K_{Ca2.3}$ and $K_{Ca3.1}$ proteins were present in considerable amounts in both pulmonary arteries and bronchioles. In addition, the observed differences in mRNA expression levels, as determined by qRT-PCR, were not predictive of the actual protein levels.

Cellular localization of $K_{Ca}2.3$ and $K_{Ca}3.1$ proteins in human pulmonary arteries and bronchioles

Immunohistochemistry revealed immunoreactions for both $K_{Ca}2.3$ and $K_{Ca}3.1$ proteins in the epithelium of bronchioles (Figure 3D,E), and the signals obtained with immunoreactions were clearly different from background (Figure 3F). While immunoreactivities for $K_{Ca}3.1$ protein were similar




Figure 1

Quantitative RT-PCR analysis of $K_{Ca}3.1$ and $K_{Ca}2.3$ mRNA expression in human pulmonary arteries and bronchioles. Data (ratio of GAPDH) are means \pm SEM of $n = 8$ human pulmonary arteries or bronchioles originating from different individuals; * $P < 0.05$, one-way ANOVA.

in bronchioles and arteries (Figure 3A,D), immunoreactive $K_{Ca}2.3$ protein was more evident in the epithelium of bronchioles than in arteries (Figure 3B,E) and clearly different from background (Figure 3C). This immunohistochemistry thus showed that $K_{Ca}2.3$ and $K_{Ca}3.1$ channels were located in the human bronchial epithelium as well as in the endothelium of pulmonary arteries.

Patch-clamp electrophysiology

We investigated functional expression of $K_{Ca}2.3$ and $K_{Ca}3.1$ channels in HSAEpi cells by whole-cell patch-clamp analysis. Figure 4A shows representative recordings of composite and voltage-independent $K_{Ca}2$ and $K_{Ca}3.1$ currents elicited by Ca^{2+} dialysis in combination with the $K_{Ca}2 / K_{Ca}3.1$ opener SKA-31 for stable channel activity. Currents were blocked by half by the selective $K_{Ca}3.1$ channel blocker TRAM-34 and were further reduced by the $K_{Ca}2$ channel blocker UCL1684. These data indicated that both channels carry the voltage-independent K_{Ca} conductance in HSAEpi cells, with both channels contributing almost equally to the total current (Figure 4B). At the transcriptional level, expression of mRNA for $K_{Ca}3.1$ and $K_{Ca}2.3$ channels, but not for the other $K_{Ca}2.1$ and $K_{Ca}2.2$ channel subtypes was confirmed by RT-PCR as shown in Figure 4C.

Functional studies in human pulmonary arteries and bronchioles

The diameter of the pulmonary arteries averaged $649 \pm 44 \mu\text{m}$ ($n = 45$), whereas the diameters of accompanying stretched bronchioles were larger and averaged $1132 \pm 85 \mu\text{m}$




Figure 2

Immunoblotting of $K_{Ca}3.1$ and $K_{Ca}2.3$ protein. Representative immunoblots showing the amount of (A) $K_{Ca}3.1$ (50 kDa) (B) $K_{Ca}2.3$ (70 kDa) and β -actin (42 kDa) protein (lower panels) in human pulmonary arteries and bronchioles. (C and D) Graphs illustrating the protein level based on $n = 5$ human tissue samples originating from the same five patients. Data are given as means \pm SEM.




Figure 3

Representative immunostaining of human pulmonary arteries (a–c) and bronchioles (d–f). Immunoreaction for $K_{Ca}3.1$ (A, D), $K_{Ca}2.3$ (B, E) and control (C, F) without primary antibody. Arrows indicate immunoreactions and the scale bar 25 μm .

($n = 45$). In pulmonary arteries, contraction induced by U46619 (10 nM) was $5.3 \pm 0.5 \text{ N m}^{-1}$ and in bronchioles contraction to histamine (2 μM) was $1.4 \pm 0.2 \text{ N m}^{-1}$. We next tested the efficacy of either the $K_{Ca}2.3/K_{Ca}3.1$ channel openers, NS309 and 1-EBIO, the opener of ATP-sensitive potassium (K_{ATP}) channels, levcromakalim, the β_2 -adrenoceptor agonist, salbutamol or the NO donor, GEA 3175, to produce relaxation of pulmonary arteries and bronchioles. The results are shown in Figure 5A–D.

As expected the NO donor, GEA 3175, induced potent concentration-dependent relaxations in pulmonary arteries, starting at 10 nM (approx. 25%); and the greatest relaxation was observed at the highest concentration tested (approx. 70% at 10 μM ; Figure 5A). Salbutamol was less effective in pulmonary arteries (Figure 5A). In pulmonary arteries, levcromakalim induced concentration-dependent relaxations, and the maximum relaxation was observed at 10 μM (Figure 5B). In pulmonary arteries, NS309 induced concentration-dependent relaxations, reaching 70% at the highest concentration of NS309 (Figure 5B). 1-EBIO produced similar

relaxations (Figure 5B), with the exception that relaxations were smaller at 10 μM , than those obtained with 10 μM NS309. Thus, if compared with direct stimulation of smooth muscle by the endothelium-independent vasorelaxing agent, GEA 3175, NS309, 1-EBIO and salbutamol relaxed human pulmonary arteries with substantially comparable effects at $\leq 1 \mu\text{M}$.

In bronchioles, salbutamol was more efficient in producing relaxation, approximately 55% vs. 25% in pulmonary arteries at 1 μM (Figure 5C). As expected, GEA 3175 evoked strong relaxations (approx. 120% at 10 μM). Interestingly, these were apparently smaller at concentrations in the sub-micromolar range but larger at 10 μM when compared with pulmonary arteries. In bronchioles (Figure 5D), levcromakalim produced potent concentration-dependent relaxations, which reached 120% at 10 μM . NS309 also induced concentration-dependent relaxations in this tissue (approx. 65% at 10 μM), similar to that seen in the pulmonary arteries. The other dual channel opener, 1-EBIO, also relaxed bronchioles, although it was less potent at the highest dose tested




Figure 4

KCa3.1 and KCa2 currents in HSAEpi cells. (A) Upper panel on left illustrates base line currents (1), immediately after membrane rupture (to achieve electrical access) and current–voltage relationship of KCa currents (2; $n = 6$ experiments) activated during cell dialysis with a pipette solution containing $3 \mu\text{M}$ $[\text{Ca}^{2+}]_{\text{free}}$. To achieve maximal and stable channel activation, the KCa2/KCa3.1 channel opener SKA-31 ($1 \mu\text{M}$) was added to the bath solution. Upper panel on right: Currents were reduced by the KCa3.1 channel blocker TRAM-34 ($1 \mu\text{M}$, $n = 4$ experiments) and further reduced by additional application of the KCa2 channel blocker UCL1684 ($1 \mu\text{M}$, $n = 4$ experiments). (B) Summary data for K^+ -currents at 0 mV. Values are given as mean \pm SEM; * $P < 0.05$, paired Student's t -test. (C) RT-PCR of KCa2 channels on HSAEpi cells and negative controls (non-reverse transcribed mRNA, -RT).

(approx. 30% at $10 \mu\text{M}$; Figure 5D). Together, NS309 and 1-EBIO were found to be fairly potent bronchodilating agents in human bronchioles, relative to the established bronchodilating efficacy of salbutamol (approx. 100%) and of a NO donor, here GEA3175.

The KCa3.1 channel blocker charybdotoxin reduced the response to $10 \mu\text{M}$ NS309 in pulmonary arteries but failed to reduce the relaxations to 100 nM and $1 \mu\text{M}$ NS309 (Figure 6A,B). In contrast, the KCa2.3 channel blocker apamin clearly reduced relaxation to NS309 at 100 nM , $1 \mu\text{M}$ and $10 \mu\text{M}$ in pulmonary arteries and in bronchioles (Figure 6A,B); and a combination of charybdotoxin and apamin did not reduce this relaxation further. In human pulmonary arteries, the specific KCa2.2 and KCa2.3 channel activator, CyPPA, induced concentration-dependent relaxations, which were reduced by apamin (Figure S3A). In bronchioles, only $100 \mu\text{M}$ CyPPA induced relaxations which were reduced in the presence of apamin (Figure S3B). In both bron-

chioles and pulmonary arteries, blocking the KCa1.1 channels with iberiotoxin did not change NS309 relaxation ($n = 4$; data in Figure S2). This indicated a major contribution of KCa2.3 channels in the relaxation response. Notably, the marked relaxation caused by $10 \mu\text{M}$ NS309 was only reduced by the combination of apamin and charybdotoxin, indicating other effects of NS309 at this high concentration.

The relaxation induced by NS309 was reduced in pulmonary arteries without endothelium (Figure 7A). The relaxation to salbutamol was also blunted by removing the endothelium (Figure 7B), whereas relaxation to the NO-donor GEA3175 was unaffected by removing the endothelium (Figure 7C). In the bronchioles, removing the epithelium reduced the relaxation to NS309 to a similar degree as in the pulmonary arteries. Salbutamol-evoked relaxation was also blunted in bronchioles without epithelium (Figure 7E), whereas GEA3175-induced relaxations were unaltered in preparations without epithelium (Figure 7F).




Figure 5

The effect of vehicle, NS309, 1-EBIO, levromakalim, salbutamol and GEA 3175 in (A, B) human pulmonary arteries and (C, D) human bronchioles. Data points represent means \pm SEM, with $n = 4-12$ per data point.




Figure 6

The effect of NS309 in (A) human pulmonary arteries and (B) human bronchioles in the presence and absence of charybdotoxin (ChTX), apamin or a combination. Data points represent means \pm SEM, with $n = 4-12$ per data point. The concentration-response curves were compared by two-way ANOVA. $*P < 0.05$, versus control curve.

Figure 7

The effect of NS309, salbutamol and GEA3175 in (A-C) human pulmonary arteries with or without endothelium, (D-F) human bronchioles with or without epithelium. Data points represent means \pm SEM, with $n = 4-7$ per data point. The concentration-response curves were compared by two-way ANOVA. $*P < 0.05$, versus control curve.

Discussion

The main findings of the present study are that $K_{Ca}2.3$ and $K_{Ca}3.1$ channels are localized in the human pulmonary endothelium and epithelium and that the pharmacological activation of these channels by a selective activator, NS309, relaxed both human isolated pulmonary arteries and bronchioles.

Our data on tissue expression of the two channels support previous findings and provides additional insights regarding channel expression in human lung tissues. Thus, in trachea and lung, previous studies suggested a high expression of $K_{Ca}3.1$ channels using RT-PCR, Northern blotting and RNA array (Jensen *et al.*, 2001; MaoXiang *et al.*, 2004). $K_{Ca}3.1$ mRNA expression is well established in systemic arteries (Grgic *et al.*, 2009) and, in particular, human mesenteric endothelium (Kohler *et al.*, 2000; Chadha *et al.*, 2011). Here, we found that the mRNAs for $K_{Ca}3.1$ and $K_{Ca}2.3$ channels were expressed both in human pulmonary arteries and bronchioles. Notably, the expression level of $K_{Ca}3.1$ mRNA was higher in bronchioles than in pulmonary arteries. However, $K_{Ca}3.1$ protein expression was found to be higher in pulmonary arteries than in bronchioles, showing that mRNA expression level did not necessarily correlate with protein levels, as previously reported (Griffin *et al.*, 2002; Minagawa *et al.*, 2008). Nonetheless, the present findings showed that $K_{Ca}3.1$ channels were expressed in considerable amounts in both human pulmonary arteries and bronchioles, and that $K_{Ca}3.1$ channels were likely to be more prominent in human pulmonary arteries.

Previous studies reported that $K_{Ca}2.3$ channels were expressed not only in various tissues especially in brain but also in peripheral tissues, such as the lung (MaoXiang *et al.*, 2004) and in arteries (Grgic *et al.*, 2009). In agreement with these previous reports, our study found expression of mRNA for $K_{Ca}2.3$ protein in human pulmonary arteries and bronchioles. Moreover, protein levels of $K_{Ca}2.3$ in human pulmonary arteries and bronchioles were comparable.

Our immunohistological approaches revealed that $K_{Ca}3.1$ channels were localized to the endothelium and the epithelium of human pulmonary arteries and bronchioles. The $K_{Ca}2.3$ channels were found in human bronchiolar epithelium and arterial endothelium. Thus, these results, together with previous reports by our group (Kroigaard *et al.*, 2010), suggest that the $K_{Ca}2.3$ and $K_{Ca}3.1$ channels are preferentially localised in the epithelial cell layer of the bronchioles and the endothelial cell layer of human pulmonary arteries. That mRNA expression and proteins of both channels led to functional channels at the membrane level in human bronchial epithelium was supported by the detection of K_{Ca} currents with the electrophysiological and pharmacological fingerprints of $K_{Ca}2.3$ and $K_{Ca}3.1$ channels in HSAEpi cells.

Concerning endothelium-dependent vasodilatation, our present study demonstrated that NS309 and the classical but less potent channel opener 1-EBIO induced dose-dependent relaxations in human pulmonary arteries. Considering the higher potency of NS309 over 1-EBIO, NS309-induced relaxations were larger than those elicited by 1-EBIO at the highest dose tested. NS309 relaxation was reduced by endothelial cell removal, suggesting that it induced endothelium-dependent

relaxations. Relaxations induced by NS309 were indeed mediated by activation of $K_{Ca}2.3$ and $K_{Ca}3.1$ channels, as indicated by the blocking effects of apamin and of charybdotoxin at the highest NS309 concentration tested. However, at high concentrations (10 μ M) of NS309, this channel opener caused relaxations in preparations without endothelium and these relaxations were partly insensitive to charybdotoxin and apamin. These relaxations resistant to apamin and charybdotoxin were most likely caused by a blockade of L-type calcium channels (Morimura *et al.*, 2006; Dalsgaard *et al.*, 2009), thus indicating loss of specificity for $K_{Ca}2.3$ and $K_{Ca}3.1$ channels of NS309 at concentrations ≥ 10 μ M. Concerning the relaxations by lower NS309 concentrations, apamin was effective in virtually abolishing the response, while charybdotoxin was ineffective here.

In summary, these results suggest that NS309 mainly activated $K_{Ca}2.3$ channels and, to a lesser extent, $K_{Ca}3.1$ channels to elicit endothelium-dependent relaxations in human pulmonary arteries, irrespective of a similar expression of the two channels. This discrepancy may be explained by considering the mechanism by which NS309 stimulates $K_{Ca}2.3$ and $K_{Ca}3.1$ channels. We found that NS309 increased the open probability of the two channels at a given Ca^{2+} concentration but did not activate the closed channels. Thus, our finding of a predominant activation of $K_{Ca}2.3$ channels could indicate basal activity of $K_{Ca}2.3$, but not of $K_{Ca}3.1$, channels in this preparation, which was then potentiated by NS309 at concentrations in the submicromolar range. Concerning $K_{Ca}3.1$ channels, the lack of a contribution of these channels to relaxations induced by 0.01–1 μ M NS309 is in line with previous studies that showed that the related $K_{Ca}3.1/K_{Ca}2.3$ channel activator, SKA-31, *per se*, was also not able to produce $K_{Ca}3.1$ channel-mediated vasodilatation in murine carotid arteries; while SKA-31 potentiated $K_{Ca}3.1$ channel-mediated vasodilatation to acetylcholine (Sankaranarayanan *et al.*, 2009). Thus, activation of $K_{Ca}3.1$ channels by this class of activators appeared to require first pre-stimulation, for instance, by Ca^{2+} mobilization following receptor activation. Indeed, the contribution of $K_{Ca}3.1$ channels to vasodilatation was only seen during acetylcholine stimulation and genetic deficiency was shown to impair this vasodilatation, selectively (Brahler *et al.*, 2009). Despite these circumstances, our functional studies suggested that NS309 activation of $K_{Ca}2$ channels (and at high concentration also of $K_{Ca}3.1$ channels) produced relaxation in human pulmonary arteries, highlighting these channels as vasorelaxing effector proteins in the pulmonary circulation. This is furthermore supported by the observation that the selective opener of $K_{Ca}2$ channels, CyPPA, induced apamin-sensitive relaxations in human pulmonary arteries.

The epithelium generates more than one factor, inducing relaxation of the underlying bronchial smooth muscle layer (Stuart-Smith and Vanhoutte, 1987; Benoit *et al.*, 2001). Moreover, activation of the $K_{Ca}1.1$ channel led to relaxation of airway smooth muscle (Benoit *et al.*, 2001), suggesting a role of K_{Ca} channels in these responses. Our present study shows that the $K_{Ca}2.3$ and $K_{Ca}3.1$ channel activator NS309 evoked concentration-dependent relaxations in human bronchioles that were greater than those evoked by 1-EBIO, a finding that was explained by the higher potency of NS309, as discussed above. Moreover, NS309 relaxation was blunted

by epithelial removal, indicating epithelium-dependent relaxation.

As in pulmonary arteries without endothelium, NS309 at high concentrations also relaxed bronchioles without epithelium, suggesting other mechanisms may also contribute to NS309 relaxations. However, apamin blocked NS309 relaxation to a larger degree than epithelial cell removal (see Figure 6B vs. 7B). Based on histological examinations, it is unlikely that a remainder of the epithelium contributes to NS309 relaxations in bronchioles without epithelium, and these findings suggest that apamin-sensitive channels in other cells in the bronchioles contribute to the NS309 relaxations. $K_{Ca2.3}$ appeared to be the channel involved in the relaxations because apamin was more effective than charybdotoxin in blocking the response. Thus, similar to our findings in the pulmonary arteries, $K_{Ca2.3}$ channels (exhibiting some basal activity) appeared to be predominantly used by NS309 to produce relaxation in human bronchioles. There are relevant species differences, as in rats both apamin- and charybdotoxin-sensitive channels are involved in NS309 relaxation in bronchioles (Kroigaard *et al.*, 2010). $K_{Ca3.1}$ channels are clearly expressed in human bronchioles and rather than a role in tone regulation, these channels may control chloride secretion in bronchial cells to maintain liquid composition and volume (Bardou *et al.*, 2009). Also the choice of spasmogen/constrictor can influence the relaxation of bronchodilators (Hernandez *et al.*, 1998) and we cannot exclude the possibility that using another spasmogen would increase the relaxation to, for instance, NS309 and reveal contributions from $K_{Ca3.1}$ channels. Despite the differential contribution of the channels to epithelium-dependent relaxation, our study demonstrates that pharmacological activation of $K_{Ca2.3}$ channels was able to relax human bronchioles, most likely by mechanisms similar to those of EDHF-induced relaxations in arteries.

In terms of pulmonary disease, $K_{Ca3.1}$ channels were found to inhibit human airway smooth muscle proliferation (Shepherd *et al.*, 2007), and activation of these channels potentiated the degranulation of human lung mast cells (Duffy *et al.*, 2001; 2005). Hence blockade of $K_{Ca3.1}$ channels was proposed as a useful approach to the treatment of asthma (Bradding and Wulff, 2009). The present study on intact human bronchioles and pulmonary arteries suggested that blockade of $K_{Ca3.1}$ channels could reduce receptor-mediated relaxation responses of pulmonary arteries and could thereby add to dysregulation of arterial tone. While pharmacological modulators of $K_{Ca3.1}$ channels could have a Janus face, selective activators of $K_{Ca2.3}$ channels could serve as more potent and safe pharmacological targets to produce pulmonary vasodilatation and bronchodilatation.

In conclusion, pharmacological enhancement of the functions of epithelial and endothelial $K_{Ca2.3}$ channels may represent a novel approach to treat chronic obstructive pulmonary disease and associated pulmonary hypertension.

Acknowledgements

We thank Helle Zibrandtsen, Henriette Gram Johanson and Susie Mogensen for excellent technical assistance. Christel Kroigaard was supported by a grant from the Danish Heart

Foundation, while Ulf Simonsen was supported by a grant from the Novo Nordisk foundation.

Conflict of interest

The authors state no conflict of interest.

References

- Alexander SPH, Mathie A, Peters JA (2011). Guide to Receptors and Channels (GRAC), 5th Edition. Br J Pharmacol 164 (Suppl. 1): S1–S324.
- Bardou O, Trinh NT, Brochiero E (2009). Molecular diversity and function of K^+ channels in airway and alveolar epithelial cells. Am J Physiol Lung Cell Mol Physiol 296: L145–L155.
- Benoit C, Renaudon B, Salvail D, Rousseau E (2001). EETs relax airway smooth muscle via an EpDHF effect: BK(Ca) channel activation and hyperpolarization. Am J Physiol Lung Cell Mol Physiol 280: L965–L973.
- Bradding P, Wulff H (2009). The K^+ channels $K(Ca)3.1$ and $K(v)1.3$ as novel targets for asthma therapy. Br J Pharmacol 157: 1330–1339.
- Brahler S, Kaistha A, Schmidt VJ, Wolffe SE, Busch C, Kaistha BP *et al.* (2009). Genetic deficit of SK3 and IK1 channels disrupts the endothelium-derived hyperpolarizing factor vasodilator pathway and causes hypertension. Circulation 119: 2323–2332.
- Buus NH, Simonsen U, Pilegaard HK, Mulvany MJ (2000). Nitric oxide, prostanoid and non-NO, non-prostanoid involvement in acetylcholine relaxation of isolated human small arteries. Br J Pharmacol 129: 184–192.
- Chadha PS, Liu L, Rikard-Bell M, Senadheera S, Howitt L, Bertrand RL *et al.* (2011). Endothelium-dependent vasodilation in human mesenteric artery is primarily mediated by myoendothelial gap junctions intermediate conductance calcium-activated k^+ channel and nitric oxide. J Pharmacol Exp Ther 336: 701–708.
- Dalsgaard T, Kroigaard C, Bek T, Simonsen U (2009). Role of calcium-activated potassium channels with small conductance in bradykinin-induced vasodilation of porcine retinal arterioles. Invest Ophthalmol Vis Sci 50: 3819–3825.
- Duffy SM, Lawley WJ, Conley EC, Bradding P (2001). Resting and activation-dependent ion channels in human mast cells. J Immunol 167: 4261–4270.
- Duffy SM, Cruse G, Lawley WJ, Bradding P (2005). Beta2-Adrenoceptor regulation of the K^+ channel $iKCa1$ in human mast cells. FASEB J 19: 1006–1008.
- Edwards G, Dora KA, Gardener MJ, Garland CJ, Weston AH (1998). K^+ is an endothelium-derived hyperpolarizing factor in rat arteries. Nature 396: 269–272.
- Feletou M, Girard V, Canet E (1995). Different involvement of nitric oxide in endothelium-dependent relaxation of porcine pulmonary artery and vein: influence of hypoxia. J Cardiovasc Pharmacol 25: 665–673.
- Flavahan NA, Aarhus LL, Rimele TJ, Vanhoutte PM (1985). Respiratory epithelium inhibits bronchial smooth muscle tone. J Appl Physiol 58: 834–838.

- Folkerts G, Nijkamp FP (1998). Airway epithelium: more than just a barrier! *Trends Pharmacol Sci* 19: 334–341.
- Frantz E, Soifer SJ, Clyman RI, Heymann MA (1989). Bradykinin produces pulmonary vasodilation in fetal lambs: role of prostaglandin production. *J Appl Physiol* 67: 1512–1517.
- Grgic I, Kaistha BP, Hoyer J, Kohler R (2009). Endothelial Ca²⁺-activated K⁺ channels in normal and impaired EDHF-dilator responses – relevance to cardiovascular pathologies and drug discovery. *Br J Pharmacol* 157: 509–526.
- Griffin TJ, Gygi SP, Ideker T, Rist B, Eng J, Hood L *et al.* (2002). Complementary profiling of gene expression at the transcriptome and proteome levels in *Saccharomyces cerevisiae*. *Mol Cell Proteomics* 1: 323–333.
- Hernandez M, Elmedal B, Mulvany MJ, Simonsen U (1998). Mechanisms of relaxations of bovine isolated bronchioles by the nitric oxide donor, GEA 3175. *Br J Pharmacol* 123: 895–905.
- Ignarro LJ, Byrns RE, Buga GM, Wood KS (1987). Endothelium-derived relaxing factor from pulmonary artery and vein possesses pharmacologic and chemical properties identical to those of nitric oxide radical. *Circ Res* 61: 866–879.
- Jensen BS, Strobaek D, Olesen SP, Christophersen P (2001). The Ca²⁺-activated K⁺ channel of intermediate conductance: a molecular target for novel treatments? *Curr Drug Targets* 2: 401–422.
- Kohler R, Degenhardt C, Kuhn M, Runkel N, Paul M, Hoyer J (2000). Expression and function of endothelial Ca²⁺-activated K⁺ channels in human mesenteric artery: a single-cell reverse transcriptase-polymerase chain reaction and electrophysiological study in situ. *Circ Res* 87: 496–503.
- Kroigaard C, Dalsgaard T, Simonsen U (2010). Mechanisms underlying epithelium-dependent relaxation in rat bronchioles: analogy to EDHF-type relaxation in rat pulmonary arteries. *Am J Physiol Lung Cell Mol Physiol* 298: L531–L542.
- MaoXiang C, Gorman A, Bill B, Kuljit S, Paul H, Michel C *et al.* (2004). Small and intermediate conductance Ca²⁺-activated K⁺ channels confer distinctive patterns of distribution in human tissues and differential cellular localisation in the colon and corpus cavernosum. *Naunyn Schmiedebergs Arch Pharmacol* V369: 602–615.
- Minagawa H, Honda M, Miyazaki K, Tabuse Y, Teramoto R, Yamashita T *et al.* (2008). Comparative proteomic and transcriptomic profiling of the human hepatocellular carcinoma. *Biochem Biophys Res Commun* 366: 186–192.
- Morimura K, Yamamura H, Ohya S, Imaizumi Y (2006). Voltage-dependent Ca²⁺-channel block by openers of intermediate and small conductance Ca²⁺-activated K⁺ channels in urinary bladder smooth muscle cells. *J Pharmacol Sci* 100: 237–241.
- Morrison KJ, Gao Y, Vanhoutte PM (1990). Epithelial modulation of airway smooth muscle. *Am J Physiol* 258: L254–L262.
- Sankaranarayanan A, Raman G, Busch C, Schultz T, Zimin PI, Hoyer J *et al.* (2009). Naphtho[1,2-d]thiazol-2-ylamine (SKA-31), a new activator of KCa2 and KCa3.1 potassium channels, potentiates the endothelium-derived hyperpolarizing factor response and lowers blood pressure. *Mol Pharmacol* 75: 281–295.
- Shepherd MC, Duffy SM, Harris T, Cruse G, Schuliga M, Brightling CE *et al.* (2007). KCa3.1 Ca²⁺-activated K⁺ channels regulate human airway smooth muscle proliferation. *Am J Respir Cell Mol Biol* 37: 525–531.
- Strobaek D, Teuber L, Jorgensen TD, Ahring PK, Kjaer K, Hansen RS *et al.* (2004). Activation of human IK and SK Ca²⁺-activated K⁺ channels by NS309 (6,7-dichloro-1H-indole-2,3-dione 3-oxime). *Biochim Biophys Acta* 1665: 1–5.
- Stuart-Smith K, Vanhoutte PM (1987). Heterogeneity in the effects of epithelium removal in the canine bronchial tree. *J Appl Physiol* 63: 2510–2515.
- Stuart-Smith K, Vanhoutte PM (1988). Airway epithelium modulates the responsiveness of porcine bronchial smooth muscle. *J Appl Physiol* 65: 721–727.
- Vanhoutte PM (1987). Airway epithelium and bronchial reactivity. *Can J Physiol Pharmacol* 65: 448–450.
- Zhang RZ, Yang Q, Yim APC, Huang Y, He GW (2006). Role of NO and EDHF-mediated endothelial function in the porcine pulmonary circulation: comparison between pulmonary artery and vein. *Vascul Pharmacol* 44: 183–191.

Supporting information

Additional Supporting Information may be found in the online version of this article:

Figure S1 Immunoblotting of K_{Ca}2.3 and K_{Ca}3.1 protein. Representative immunoblots showing the amount of (A) K_{Ca}2.3 (70 kDa) and (B) K_{Ca}3.1 (50 kDa) protein in human pulmonary arteries (PA) and bronchioles (B).

Figure S2 The effect of NS309 in human pulmonary arteries (A) and human bronchioles (B) in the presence and absence of iberiotoxin (IbTX). Data points represent means ± SEM, with *n* = 4 per data point. The concentration–response curves were compared by two-way ANOVA, **P* < 0.05 versus control curve.

Figure S3 The effect of CyPPA in (A) human pulmonary arteries, in which apamin reduced the relaxation to CyPPA, and (B) human bronchioles, in which apamin reduced CyPPA relaxation at the highest concentration applied (100 μM, Bonferroni post test). Data points represent means ± SEM, with *n* = 3 per data point. The concentration–response curves were compared by two-way ANOVA, **P* < 0.05 versus control.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.