

## VU Research Portal

### Acupuncture for peripheral joint osteoarthritis

Manheimer, Eric; Cheng, K.; Linde, Klaus; Lao, Lixing; Yoo, Junghee; Wieland, Susan; van der Windt, Daniëlle Awm; Berman, Brian M.; Bouter, Lex M.

#### **published in**

Cochrane Database of Systematic Reviews  
2010

#### **DOI (link to publisher)**

[10.1002/14651858.CD001977.pub2](https://doi.org/10.1002/14651858.CD001977.pub2)

#### **document version**

Publisher's PDF, also known as Version of record

#### [Link to publication in VU Research Portal](#)

#### **citation for published version (APA)**

Manheimer, E., Cheng, K., Linde, K., Lao, L., Yoo, J., Wieland, S., van der Windt, D. A., Berman, B. M., & Bouter, L. M. (2010). Acupuncture for peripheral joint osteoarthritis. *Cochrane Database of Systematic Reviews*, 2010(1), [CD001977]. <https://doi.org/10.1002/14651858.CD001977.pub2>

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

#### **E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

# Acupuncture for peripheral joint osteoarthritis (Review)

Manheimer E, Cheng K, Linde K, Lao L, Yoo J, Wieland S, van der Windt DAWM, Berman BM, Bouter LM



**THE COCHRANE  
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2010, Issue 1

<http://www.thecochranelibrary.com>



[Intervention Review]

## Acupuncture for peripheral joint osteoarthritis

Eric Manheimer<sup>1</sup>, Ke Cheng<sup>2</sup>, Klaus Linde<sup>3</sup>, Lixing Lao<sup>4</sup>, Junghee Yoo<sup>5</sup>, Susan Wieland<sup>6</sup>, Daniëlle AWM van der Windt<sup>7</sup>, Brian M Berman<sup>1</sup>, Lex M Bouter<sup>8</sup>

<sup>1</sup>Center for Integrative Medicine, University of Maryland School of Medicine, Baltimore, Maryland, USA. <sup>2</sup>Acupuncture-Moxibustion and Tuina, Shanghai University of TCM, Shanghai Research Center of Acupuncture and Meridians, Shanghai, China. <sup>3</sup>Institut für Allgemeinmedizin / Institute of General Practice, Technische Universität München / Klinikum rechts der Isar, München, Germany. <sup>4</sup>Complementary Medicine Program, University of Maryland School of Medicine, Baltimore, Maryland, USA. <sup>5</sup>College of Oriental Medicine, Kyunghee University, Seoul, Korea, South. <sup>6</sup>Providence, Rhode Island, USA. <sup>7</sup>Department of Primary Care & Health Sciences, Keele University, Keele, UK. <sup>8</sup>Executive Board of VU University Amsterdam, Amsterdam, Netherlands

Contact address: Eric Manheimer, Center for Integrative Medicine, University of Maryland School of Medicine, Kernan Hospital Mansion, 2200 Kernan Drive, Baltimore, Maryland, 21207-6697, USA. [emanheimer@compmed.umm.edu](mailto:emanheimer@compmed.umm.edu).

**Editorial group:** Cochrane Musculoskeletal Group.

**Publication status and date:** New, published in Issue 1, 2010.

**Review content assessed as up-to-date:** 16 April 2008.

**Citation:** Manheimer E, Cheng K, Linde K, Lao L, Yoo J, Wieland S, van der Windt DAWM, Berman BM, Bouter LM. Acupuncture for peripheral joint osteoarthritis. *Cochrane Database of Systematic Reviews* 2010, Issue 1. Art. No.: CD001977. DOI: 10.1002/14651858.CD001977.pub2.

Copyright © 2010 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

### ABSTRACT

#### Background

Peripheral joint osteoarthritis is a major cause of pain and functional limitation. Few treatments are safe and effective.

#### Objectives

To assess the effects of acupuncture for treating peripheral joint osteoarthritis.

#### Search strategy

We searched the Cochrane Central Register of Controlled Trials (*The Cochrane Library* 2008, Issue 1), MEDLINE, and EMBASE (both through December 2007), and scanned reference lists of articles.

#### Selection criteria

Randomized controlled trials (RCTs) comparing needle acupuncture with a sham, another active treatment, or a waiting list control group in people with osteoarthritis of the knee, hip, or hand.

#### Data collection and analysis

Two authors independently assessed trial quality and extracted data. We contacted study authors for additional information. We calculated standardized mean differences using the differences in improvements between groups.

#### Main results

Sixteen trials involving 3498 people were included. Twelve of the RCTs included only people with OA of the knee, 3 only OA of the hip, and 1 a mix of people with OA of the hip and/or knee. In comparison with a sham control, acupuncture showed statistically significant, short-term improvements in osteoarthritis pain (standardized mean difference -0.28, 95% confidence interval -0.45 to -0.11; 0.9 point greater improvement than sham on 20 point scale; absolute percent change 4.59%; relative percent change 10.32%;

---

**Acupuncture for peripheral joint osteoarthritis (Review)**

Copyright © 2010 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

9 trials; 1835 participants) and function (-0.28, -0.46 to -0.09; 2.7 point greater improvement on 68 point scale; absolute percent change 3.97%; relative percent change 8.63%); however, these pooled short-term benefits did not meet our predefined thresholds for clinical relevance (i.e. 1.3 points for pain; 3.57 points for function) and there was substantial statistical heterogeneity. Additionally, restriction to sham-controlled trials using shams judged most likely to adequately blind participants to treatment assignment (which were also the same shams judged most likely to have physiological activity), reduced heterogeneity and resulted in pooled short-term benefits of acupuncture that were smaller and non-significant. In comparison with sham acupuncture at the six-month follow-up, acupuncture showed borderline statistically significant, clinically irrelevant improvements in osteoarthritis pain (-0.10, -0.21 to 0.01; 0.4 point greater improvement than sham on 20 point scale; absolute percent change 1.81%; relative percent change 4.06%; 4 trials; 1399 participants) and function (-0.11, -0.22 to 0.00; 1.2 point greater improvement than sham on 68 point scale; absolute percent change 1.79%; relative percent change 3.89%). In a secondary analysis versus a waiting list control, acupuncture was associated with statistically significant, clinically relevant short-term improvements in osteoarthritis pain (-0.96, -1.19 to -0.72; 14.5 point greater improvement than sham on 100 point scale; absolute percent change 14.5%; relative percent change 29.14%; 4 trials; 884 participants) and function (-0.89, -1.18 to -0.60; 13.0 point greater improvement than sham on 100 point scale; absolute percent change 13.0%; relative percent change 25.21%). In the head-on comparisons of acupuncture with the 'supervised osteoarthritis education' and the 'physician consultation' control groups, acupuncture was associated with clinically relevant short- and long-term improvements in pain and function. In the head-on comparisons of acupuncture with 'home exercises/advice leaflet' and 'supervised exercise', acupuncture was associated with similar treatment effects as the controls. Acupuncture as an adjuvant to an exercise based physiotherapy program did not result in any greater improvements than the exercise program alone. Information on safety was reported in only 8 trials and even in these trials there was limited reporting and heterogeneous methods.

### **Authors' conclusions**

Sham-controlled trials show statistically significant benefits; however, these benefits are small, do not meet our pre-defined thresholds for clinical relevance, and are probably due at least partially to placebo effects from incomplete blinding. Waiting list-controlled trials of acupuncture for peripheral joint osteoarthritis suggest statistically significant and clinically relevant benefits, much of which may be due to expectation or placebo effects.

## **PLAIN LANGUAGE SUMMARY**

### **Acupuncture for osteoarthritis**

This summary of a Cochrane review presents what we know from research about the effect of acupuncture on osteoarthritis.

#### **The review shows that in people with osteoarthritis,**

-Acupuncture may lead to small improvements in pain and physical function after 8 weeks.

-Acupuncture may lead to small improvements in pain and physical function after 26 weeks.

We often do not have precise information about side effects and complications. This is particularly true for rare but serious side effects. Possible side effects of acupuncture treatment include minor bruising and bleeding at the site of needle insertion.

#### **What is osteoarthritis and what is acupuncture?**

Osteoarthritis (OA) is a disease of the joints, such as your knee or hip. When the joint loses cartilage, the bone grows to try and repair the damage. Instead of making things better, however, the bone grows abnormally and makes things worse. For example, the bone can become misshapen and make the joint painful and unstable. This can affect your physical function or ability to use your knee.

According to the philosophy of traditional acupuncture, energy circulates in 'meridians' located throughout the body. Pain or ill health happens when something occurs to cause this meridian energy circulation to be blocked. The way to restore health is to stimulate the appropriate combination of acupuncture points in the body by inserting very thin needles. Sometimes in painful conditions, electrical stimulation along with the needles is also used. According to acupuncture theory, one way you can tell that acupuncture is relieving pain is that you may feel numbness or tingling, called de qi, where the needle is inserted.

#### **Best estimate of what happens to people with osteoarthritis who have acupuncture:**

---

**Acupuncture for peripheral joint osteoarthritis (Review)**

Copyright © 2010 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

**Pain after 8 weeks:**

- People who had acupuncture rated their pain to be improved by about 4 points on a scale of 0 to 20.
- People who received sham acupuncture rated their pain to be improved by about 3 points on a scale of 0 to 20.
- People who received acupuncture had a 1 point greater improvement on a scale of 0-20. (5% absolute improvement).

**Pain after 26 weeks:**

- People who had acupuncture rated their pain to be improved by slightly more than 3 points on a scale of 0 to 20.
- People who received sham acupuncture rated their pain to be improved by slightly less than 3 points on a scale of 0 to 20.
- People who received acupuncture had under a 1 point greater improvement on a scale of 0-20. (2% absolute improvement).

**Physical function after 8 weeks :**

- People who had acupuncture rated their function to be improved by about 11 points on a scale of 0 to 68.
- People who received sham acupuncture rated their function to be improved by about 8 points on a scale of 0 to 68.
- People who received acupuncture had about a 3 point greater improvement on a scale of 0-68. (4% absolute improvement)

**Physical function after 26 weeks :**

- People who had acupuncture rated their function to be improved by about 11 points on a scale of 0 to 68.
- People who received sham acupuncture rated their function to be improved by about 10 points on a scale of 0 to 68.
- People who received acupuncture had about a 1 point greater improvement on a scale of 0-68. (2% absolute improvement)