

Acute lower respiratory infections in childhood: opportunities for reducing the global burden through nutritional interventions

Daniel E Roth,^a Laura E Caulfield,^a Majid Ezzati^b & Robert E Black^a

Abstract Inadequate nutrition and acute lower respiratory infection (ALRI) are overlapping and interrelated health problems affecting children in developing countries. Based on a critical review of randomized trials of the effect of nutritional interventions on ALRI morbidity and mortality, we concluded that: (1) zinc supplementation in zinc-deficient populations prevents about one-quarter of episodes of ALRI, which may translate into a modest reduction in ALRI mortality; (2) breastfeeding promotion reduces ALRI morbidity; (3) iron supplementation alone does not reduce ALRI incidence; and (4) vitamin A supplementation beyond the neonatal period does not reduce ALRI incidence or mortality. There was insufficient evidence regarding other potentially beneficial nutritional interventions. For strategies with a strong theoretical rationale and probable operational feasibility, rigorous trials with active clinical case-finding and adequate sample sizes should be undertaken. At present, a reduction in the burden of ALRI can be expected from the continued promotion of breastfeeding and scale-up of zinc supplementation or fortification strategies in target populations.

Bulletin of the World Health Organization 2008;86:356–364.

Une traduction en français de ce résumé figure à la fin de l'article. Al final del artículo se facilita una traducción al español. الترجمة العربية لهذه الخلاصة في نهاية النص الكامل لهذه المقالة.

Introduction

Estimation of the global burden of child mortality attributable to undernutrition has played a crucial role in refocusing the attention of researchers and policy-makers on the importance of optimal maternal–child nutrition for promoting neonatal, infant and child survival,^{1–3} including the prevention of mortality due to severe acute lower respiratory infection (ALRI).^{2–5} To advance the public health application of knowledge about the interrelated burdens of childhood ALRI and poor nutrition in developing countries, we have critically reviewed available data regarding the efficacy and effectiveness of specific nutritional interventions for reducing global childhood ALRI incidence, morbidity and ALRI-specific mortality.

Methods

Search strategy

This review included meta-analyses and large-scale randomized controlled trials of micronutrient supplementation, breastfeeding promotion, complementary food provision or counselling, and antenatal nutritional interventions, in

which at least one childhood ALRI outcome (incidence, morbidity or mortality) was measured. Smaller studies or those with non-randomized designs were included where higher-quality data were unavailable. The literature review was based primarily on several systematic reviews that formed the evidence base for the *Lancet Undernutrition Series (LUS)* published earlier this year (available at: http://www.globalnutrition-series.org/web_appendices). To include articles published after completion of the systematic reviews, we searched PubMed (1990–January 2008) and reference lists of selected recent articles published on each topic. PubMed search terms included nutrient-specific keyword(s) and a string that broadly captured childhood ALRI-related articles (“ALRI” OR “ARI” OR “pneumonia” OR “lower respiratory tract infection” OR “lower respiratory infection” OR “bronchiolitis” OR “bronchopneumonia” OR “morbidity” OR “mortality”) without age or language restrictions. Titles/abstracts were scanned for relevant interventional studies or key supportive articles, for which full-text articles were retrieved.

ALRI outcome definition

As there is no standard definition of childhood ALRI,⁶ studies were included if they applied an outcome definition incorporating at least one specific lower respiratory tract sign reported by a caregiver or study personnel (fast or difficulty breathing, chest wall indrawing) and/or abnormal auscultatory findings (crackles/crepitations or bronchial breath sounds). Authors occasionally differentiated ALRI subtypes on the basis of wheeze versus crepitations/crackles (probable bronchiolitis versus pneumonia, respectively); however, viral diagnostics, isolation of pathogenic bacteria from a sterile fluid (i.e. blood culture, lung aspirate), or unequivocal radiographic findings (i.e. lobar consolidation or pleural effusion) were not documented in the reviewed trials.

Burden of disease

Estimates of the burden of ALRI attributable to selected nutritional factors were extracted from an analysis performed for LUS^{3,7} [where “burden” refers to ALRI-related deaths and disability-adjusted life years (DALYs) lost], and are discussed in the context of

^a Department of International Health, Johns Hopkins University, Bloomberg School of Public Health, 615 North Wolfe Street, Baltimore, MD, United States of America.

^b Department of Population and International Health and Department of Environmental Health, Harvard School of Public Health, Boston, MA, USA.

Correspondence to Laura E Caulfield (e-mail: lcaulfie@jhsph.edu).

doi:10.2471/BLT.07.049114

(Submitted: 31 October 2007 – Revised version received: 15 January 2008 – Accepted: 23 January 2008)

Table 1. Estimated ALRI deaths and disease burden in children under 5 years of age attributed to nutritional risk factors, in 2004

Nutritional risk factor	Estimates ^a	UN region			All developing countries
		Africa ^b	Asia ^b	Latin America ^b	
Underweight ^c	Deaths	238 234 (24.3)	203 104 (28.1)	1 542 (3.1)	442 879 (25.3)
	DALYs x 10 ³	8 335 (24.3)	7 262 (27.9)	65.69 (3.1)	15 662 (25.1)
Stunting ^c	Deaths	243 325 (24.9)	152 906 (21.1)	3 540 (7.0)	399 771 (22.8)
	DALYs x 10 ³	8 510 (24.9)	5 479 (21.1)	150.0 (7.1)	14 136 (22.7)
IUGR-LBW ^{d,e}	Deaths	31 866 (11.9)	97 297 (22.0)	1 550 (5.2)	130 713 (17.6)
	DALYs x 10 ³	1 092 (11.9)	3 330 (22.0)	53.97 (5.2)	4 475 (17.6)
Zinc deficiency ^f	Deaths	66 062 (6.8)	48 800 (6.7)	3 135 (6.2)	117 997 (6.7)
	DALYs x 10 ³	2 350 (6.9)	1 840 (7.1)	159.2 (7.5)	4 349 (7.0)
Suboptimal breastfeeding (0–28 days of age) ^{g,h}	Deaths	115 193 (42.9)	198 128 (44.8)	14 394 (48.2)	327 714 (44.2)
	DALYs x 10 ³	3 947 (42.9)	6 798 (44.8)	500.9 (48.2)	11 246 (44.3)
Suboptimal breastfeeding (> 1 month of age) ^g	Deaths	187 304 (19.1)	141 942 (19.6)	14 987 (29.8)	344 234 (19.6)
	DALYs x 10 ³	6 536 (19.1)	5 036 (19.4)	578.3 (27.3)	12 150 (19.5)

ALRI, acute lower respiratory infection; DALYs, disability-adjusted life years; IUGR, intrauterine growth restriction; LBW, low birth weight.

^a Values in parentheses are percentage of all under-five ALRI deaths or DALYs. Because nutritional risk factors are overlapping and often correlated, attributable fractions should not be directly summed across risk factors.

^b UN region composition available at: <http://unstats.un.org/unsd/methods/m49/m49regin.htm>

^c In these analyses, risks of children considered underweight [weight-for-age z score < -1 standard deviations (SD) on the WHO growth curve] or stunted (height-for-age z score < -1 SD) were compared with those classified as "normal" (> -1 SD on the respective growth curve).

^d Risks among term neonates weighing between 1500 g and 2499 g were compared with those of birth weight \geq 2500 g. Neonates weighing < 1500 g were excluded from the analysis because of the strong likelihood of preterm birth.

^e Estimates for IUGR-LBW and suboptimal breastfeeding in the newborn period only apply to neonates 0–28 days of age and were based on the relative risk of the aggregate outcome "perinatal infections". Absolute numbers of deaths and DALYs were calculated by assuming that 80% of perinatal infections are ALRIs.⁸¹

^f Estimates were applied to children aged 6–59 months of age, since the efficacy of zinc supplementation has not been demonstrated in infants < 6 months of age (see text).

^g For the 0–5 month age range, infants not exclusively breastfed were compared with those exclusively breastfed. For infants 6–23 months of age, those receiving any breast milk were compared with those receiving no breast milk.

related interventions. The ALRI-specific attributable fractions (Table 1) were not reported in *LUS*.

Results

Breastfeeding promotion

A lack of exclusive breastfeeding in the first half of infancy is a risk factor for ALRI incidence, morbidity and death.^{3,5} In the *LUS* analysis,³ approximately 44% of infection-related neonatal deaths/DALYs (including those due to ALRI) and 20% of postnatal ALRI deaths/DALYs lost were attributed to suboptimal breastfeeding (Table 1). A causal effect of breastfeeding is plausible given the maternal-infant transfer of innate immune effectors (e.g. lactoferrin, lysozyme, secretory IgA, leukocytes)⁸ and influences of breast milk on immune-system maturation.⁹ Breastfeeding may enhance the antibody response to important pneumonia-causing pathogens (e.g. pneumococci, *Haemophilus influenzae*),¹⁰ but the specific mechanisms by which breastfeed-

ing ameliorates ALRI resistance are less obvious than those that underlie diarrhoea risk reduction.

Eliminating the fraction of the ALRI burden due to suboptimal breastfeeding relies on effective breastfeeding promotion and education; however, few studies have quantified the effect of breastfeeding programmes on ALRI risk reduction.¹¹ Four studies that reported respiratory outcomes did not distinguish upper and lower respiratory tract infections.^{12–15} PROBIT, a large cluster-randomized trial of breastfeeding promotion (based on the WHO baby-friendly hospital initiative) in Belarus,¹⁶ was the only study from which an effect of breastfeeding promotion on ALRI outcomes could be inferred. Success of the promotion efforts was evidenced by increased breastfeeding continuation and a significant 40% decrease in diarrhoea incidence in the intervention group. The trial showed a 15% decrease in respiratory-disease-related hospitalizations (presumably due

to ALRI), but despite the large sample size ($n = 17\ 046$) and high event rate (20.5% of infants in control arm had ≥ 1 respiratory disease hospitalization), the reported confidence intervals crossed the null after adjustment for confounders and design effects (Table 2).

The overall benefits of breastfeeding promotion are now widely accepted, and it seems doubtful that future similar trials will be deemed ethical or be more rigorously conducted than PROBIT, even though the magnitude of the effect might differ across populations or alternative behaviour-change strategies. Therefore, we concluded that the benefit of breastfeeding promotion on ALRI morbidity in PROBIT was consistent with observational data, and alongside the other known health benefits of breastfeeding (e.g. diarrhoea prevention, lactational amenorrhoea), reinforces the need to continue to support breastfeeding promotion policies and programmes in resource-poor settings.

Table 2. Estimated effects of selected nutritional interventions on ALRI incidence, morbidity or mortality

Nutritional intervention	ALRI outcome	Effect size ^a	Source	Comments
Vitamin A supplementation	Incidence	0.95 (0.89–1.01)	Meta-analysis of five trials ²³	Recent systematic review confirmed the null effects on incidence and mortality ²⁴
	Mortality	0.98 (0.75–1.28)	Meta-analysis of five trials ²³	
Zinc supplementation	Incidence	0.80 (0.70–0.92)	Meta-analysis of four trials ⁴⁴	Included trials of supplementation for at least 3 months
	Mortality	0.85 (0.65–1.11)	Unpublished meta-analysis of three trials ³	Effect size (95% CI) in individual trials: Bangladesh – cannot calculate relative rate [10 deaths in placebo group ($n = 812$) and 0 in zinc arm ($n = 809$)]; Nepal 0.90 (0.64–1.26); United Republic of Tanzania 0.88 (0.5–1.5)
Breastfeeding promotion	Hospitalization	0.85 (0.57–1.27)	Single large trial in Belarus ¹⁶	Outcome ascertained by maternal recall
Iron supplementation	Incidence	0.97 (0.83–1.23)	Meta-analysis of eight trials ³¹	Trials published since the meta-analysis have confirmed the null effect ³²
Iron and folic acid supplementation	Incidence	0.92 (0.7–1.09)	Single large trial in Nepal ³⁴	In a companion trial conducted in a malaria-endemic region (Pemba), iron-folic acid (with or without zinc) significantly increased the risk of death due to non-malarial infections (HR 1.61, 95% CI: 1.03–2.52), although the specific effect on ALRI mortality was not estimated ⁵⁸
	Mortality	0.88 (0.50–1.46)	Single large trial in Nepal ³⁴	
Iron, folic acid and zinc supplementation	Incidence	0.91 (0.76–1.08)	Single large trial in Nepal ³⁴	Single large trial in Nepal ³⁴
	Mortality	0.80 (0.45–1.34)	Single large trial in Nepal ³⁴	
Multiple micronutrient-fortified milk	Incidence	0.74 (0.57–0.97)	Single trial in India ⁶⁰	Experimental milk contained higher concentrations of vitamins A, E, C, iron, zinc, selenium and copper than the control milk

ALRI, acute lower respiratory infection; CI, confidence interval; HR, hazard ratio.

^a Relative risk or relative rate, odds ratio or hazard ratio, compared with placebo. Values in parentheses are 95% CI.

Complementary food supplementation or counselling

A large body of literature has established strong links between malnutrition, immune dysfunction and infectious diseases.^{4,17,18} The *LUS* analysis estimated that one-quarter of under-five deaths and DALYs lost were attributable to undernutrition (represented by underweight or stunting) and could theoretically be prevented by dietary interventions that normalize anthropometric indices in early childhood (Table 1). These estimates suggest that in addition to an optimal prenatal environment and breastfeeding practices, improvements in the density, diversity and quality of complementary foods may reduce the risk of ALRI incidence or mortality. However, there is a surprising lack of intervention research addressing this question. We found only one study that measured the impact of complementary feeding education or food supplements on ALRI incidence,¹⁹ but the low statistical power limited interpretation of its null findings. There were also few published studies of the

effect of comprehensive community-based nutrition programmes on ALRI outcomes. Two studies from Viet Nam reported ALRI outcomes,^{20,21} but interpretation of their findings was limited by methodological weaknesses (i.e. small sample size, quasi- or non-randomized designs, non-specific outcome ascertainment and incomplete presentation of findings).

Single micronutrient interventions

Vitamin A

Vitamin A is involved in immune function and respiratory epithelial cell differentiation,²² yet the impact of high-dose supplementation on under-five mortality cannot be accounted for by its effects on ALRI outcomes. A rigorous meta-analysis of five large trials concluded that routine vitamin A supplementation did not affect ALRI incidence or mortality (Table 2),²³ inferences that have not changed over time.²⁴ Estimates of the effect of routine early neonatal vitamin A supplementation remain imprecise and inconsistent,^{25–27} and await clarification by as-yet unpub-

lished findings from a large study of early neonatal vitamin A supplementation in Bangladesh. In the treatment setting, infants and children with non-measles-related ALRI do not benefit from adjunctive high-dose vitamin A administration.^{24,28,29}

Iron and folic acid

The interplay between iron and infection has been the subject of enduring debate in nutritional immunology, primarily because iron deficiency impairs components of cell-mediated immunity but can also inhibit the growth of bacterial pathogens.³⁰ Nonetheless, the epidemiological data regarding the effect of iron supplementation on ALRI incidence quite convincingly reveal its effect to be null (Table 2).^{31,32}

A recent prospective study in India found serum folate status to be independently associated with the risk of ALRI,³³ but we did not identify any trials assessing the effect of folic acid supplementation alone on ALRI outcomes. However, in a large trial in Nepal, the combination of daily iron and folic

acid (IFA) supplementation modestly reduced ALRI incidence and mortality compared to placebo (Table 2).³⁴ Given the lack of effect of iron supplementation alone and biological mechanisms by which folates may be implicated in immune function,³⁵ the benefit may have been attributable to improvements in folate status; however, the wide confidence intervals that included the null indicate that the observation requires confirmation.

Vitamin D and calcium

Nutritional rickets, caused by dietary deficiencies of vitamin D and/or calcium, has been strongly associated with severe ALRI in hospitalized children in some settings.^{36,37} Case-control studies in developing countries have found low vitamin D status to be associated with the risk of ALRI in Indian children³⁸ and Turkish neonates,³⁹ but not with the risk of viral bronchiolitis in young Canadian children.⁴⁰ The active form of vitamin D affects a variety of immune functions,³⁵ including the synthesis of endogenous antimicrobial peptides.⁴¹ However, interventional trials of the effect of maternal or infant supplementation with vitamin D, with or without calcium, on the incidence of infant ALRI have not yet been reported.

Zinc

Infant and child zinc supplementation has been intensively studied because dietary zinc deficiency is highly prevalent in developing countries⁴² and zinc is involved in the maintenance of skin and mucous membranes, leukocyte function and cytokine expression.⁴³ A recent meta-analysis of randomized efficacy trials conducted in South Asia confirmed that routine oral daily or weekly zinc supplementation for at least 3 months significantly reduces the incidence of childhood ALRI (Table 2),⁴⁴ an effect that may be specific to non-wheezy ALRI (i.e. cases that are more likely to be pneumonia than viral bronchiolitis);⁴⁵ this effect appears to increase with greater severity of disease,⁴⁵ and was not evident in infants less than 6 months of age.⁴⁶

Several studies not included in the published meta-analysis have shown null or non-significant effects of routine zinc on ALRI incidence in Latin America,^{47,48} South Africa⁴⁹ and Nepal.⁵⁰ Although it is possible that the heterogeneity is due to characteristics of

the study populations, it is notable that ALRI was ascertained on the basis of either maternal report or WHO criteria alone (elevated respiratory rate) in these null studies. In an updated meta-analysis of routine zinc supplementation trials, we found that incidence of ALRI was substantially reduced by zinc supplementation when the diagnosis was established by active prospective case-finding and a clinical exam that included pulmonary auscultation (relative rate, RR 0.78; 95% confidence interval, CI: 0.68–0.91; based on three of the four trials included in the published meta-analysis), whereas the effect was absent in trials that ascertained ALRI on the basis of either elevated respiratory rate alone (RR 0.99; 95% CI: 0.89–1.09) or caregiver recall of ALRI signs or symptoms (RR 1.02; 95% CI: 0.97–1.09) (Roth & Black, unpublished findings). Null findings may thus be biased due to non-specific outcome definitions⁵¹ or “contamination” by episodes of upper respiratory tract infections.

If the magnitude of the benefit of zinc prophylaxis on ALRI incidence is limited to bacterial pneumonia or increases with greater disease severity,⁴⁵ a large relative decrease in ALRI-specific mortality would be expected; however, the estimated reduction upon pooling available data for children aged 1–59 months was only 15% (Table 2).³ The discrepancy could be purely methodological (i.e. misclassification of causes of death), or perhaps reflects the failure of zinc to prevent disease in those children with the highest baseline risk of ALRI fatality (i.e. young infants). An alternative explanation is that zinc supplements may increase the case fatality rate in a small subset of children who acquire ALRIs of certain bacterial etiologies, as suggested by Coles et al. who hypothesized that specific zinc–pathogen interactions may deleteriously alter the host response in severe pneumonia.⁵²

In summary, routine zinc supplementation to young children prevents about one-quarter of ALRI cases, which likely translates into a modest reduction in ALRI-related mortality. Scaling-up zinc supplementation to all regions with high prevalence of zinc deficiency could avert an estimated 7% of ALRI deaths and DALYs lost (Table 1).

In contrast to its prophylactic effect, there is no strong evidence of a benefit of zinc supplementation in the

treatment context. Of three published randomized controlled trials evaluating zinc supplementation as adjunctive treatment in children presenting to hospital with ALRIs, one found a relative reduction in the duration of severe disease equivalent to about one day in hospital,⁵³ one study showed no benefit,⁵⁴ and a third study found that zinc caused a slightly reduced duration of fever and severe illness in boys, but not in girls.⁵⁵

Multiple micronutrient interventions

Children in resource-poor settings often exhibit concurrent deficiencies of multiple micronutrients,⁵⁶ many of which function as cofactors or regulatory molecules in immune or inflammatory cascades (e.g. vitamins C, D, selenium).⁵⁵ Therefore, multiple micronutrient supplements may theoretically augment the effect of zinc on infectious disease resistance in resource-poor settings.

Addition of zinc to IFA supplementation

Routine infant/child IFA supplementation is currently recommended by WHO for regions with a high prevalence of iron-deficiency anaemia,⁵⁷ so it is of primary interest to know whether including zinc in a daily IFA regimen would reduce ALRI-associated morbidity or mortality. In a large factorial trial in Nepal, zinc minimally strengthened the effect of IFA on ALRI incidence and mortality relative to placebo (Table 2);³⁴ however, in the United Republic of Tanzania, the addition of zinc to IFA did not notably alter the risk of non-malarial infection-related adverse events (hospitalization or death) compared to placebo.⁵⁸ In a third study involving children aged 1–23 months in Delhi ($n = 94\ 359$), but in which a placebo arm was not included, adding zinc to routine daily IFA supplementation for up to 12 months did not significantly impact the risk of ALRI-related hospitalizations (RR 1.09; 95% CI: 0.94–1.25) compared to IFA alone.⁵⁹ In the latter trial, adding zinc was associated with reduced ALRI-related mortality (RR 0.73; 95% CI: 0.55–0.96; confidence limits unadjusted for the likely negligible effect of intrahousehold clustering), but the authors emphasized that

overall mortality and the distribution of causes of death did not significantly differ between the IFA + zinc and IFA-treatment arms, suggesting this estimate should be interpreted with caution.

Multiple micronutrient supplements or fortificants

In a recent trial involving children aged 1–3 years in Delhi, India, a commercial milk powder fortified with additional vitamins A, C, E, iron, zinc (7.8 mg more than control group), selenium and copper led to a significant reduction in the incidence of clinically confirmed ALRI compared to the standard milk powder (Table 2).⁶⁰ Three other studies of multiple micronutrient supplements or fortificants had null results but were inadequately powered to detect moderate differences in ALRI incidence between treatment and control arms.^{49,61,62}

In summary, the only multiple micronutrient intervention with a clear benefit on ALRI incidence or mortality was a milk powder fortified with zinc and six other micronutrients, but the effect size was comparable to that achieved by supplementation with zinc alone. Null findings in trials using non-food-based supplements that combined iron and zinc (i.e. attenuation of the expected zinc effect) may be partially attributable to the inhibitory effects of supplemental iron on absorption of zinc supplements.⁶³ Where food fortification is not feasible, a proposed approach to avoid intrainestinal mineral interactions is alternate-day zinc and iron supplementation.

Antenatal nutritional interventions

Low birth weight (LBW: birth weight < 2500 g) in term-infants is a surrogate marker of intrauterine growth restriction (IUGR) and an established epidemiological risk factor for ALRI incidence and hospitalization.⁵ In the *LUS* analysis,³ an estimated 17.6% of ALRI-related deaths and DALYs lost in term neonates (birth to 1 month of age) were attributable to LBW (Table 1).

The biological plausibility of the association between LBW and respiratory morbidity is supported by observations of impaired immunocompetence in infants born small-for-gestational^{64–66} and the association of fetal growth restriction with structural alterations that may affect lung anatomy and func-

tion.^{67–69} However, the causal nature of the association between LBW and ALRI is complex, particularly given that LBW represents the cumulative and aggregative effect of a range of nutritional and non-nutritional prenatal exposures.^{70,71} Wilcox⁷² and others⁷³ have persuasively questioned the use of LBW as a biomarker of the effect of antenatal exposures on neonatal morbidity and mortality, suggesting it is of more practical relevance to directly consider the modifiable maternal nutritional and dietary factors that are hypothesized to “programme” the fetal immune system.⁹ Unfortunately, at present there are insufficient data to recommend antenatal nutritional interventions on the basis of their effects on infant ALRI outcomes. Trials of antenatal zinc,⁷⁴ multiple micronutrients⁷⁵ and vitamin A⁷⁶ did not have statistically significant impacts, but may have lacked sufficient power to demonstrate subtle, but potentially meaningful, effects.

Discussion

Observational evidence of an association between inadequate growth and the risk of experiencing or dying from ALRI in early childhood is strong and consistent,^{2,3,5} suggesting there could be a potentially large impact of nutritional interventions on ALRI morbidity and mortality. However, evidence-based public health policy and practice ideally rely on efficacy and effectiveness data from well conducted trials.

Based on a critical review of the literature, we were able to reach conclusions about only a few of the potential nutritional interventions. First, oral zinc supplementation in children > 6 months of age was the only intervention for which high-quality trial evidence has demonstrated a significant reduction in childhood ALRI incidence and a trend towards a reduction in mortality. Important feasibility issues remain to be resolved, including attenuation of the effects by concurrent iron supplementation, potential effects on case-fatality due to ALRIs of specific etiologies, and the optimal mode of community-based delivery (i.e. fortification, supplementation). Large-scale cluster-randomized effectiveness trials in Bangladesh⁷⁷ and India⁷⁸ have demonstrated that programmatic implementation of the WHO recommendation to incorporate a two-week

course of zinc supplementation into case management algorithms for children with acute gastroenteritis has the added benefit of reducing the burden of severe ALRI in the community. Scale-up of this approach is therefore suggested as a feasible and effective strategy for improving zinc status in at-risk populations.

Second, despite the limited available trial data, there should be little hesitation in including ALRI risk reduction in the list of benefits of exclusive breastfeeding, given the strength and consistency of observational findings and the evidence of a modest reduction in ALRI-related hospitalizations in the only rigorous trial of breastfeeding promotion. Third, there was strong evidence that vitamin A supplementation (after the neonatal period), or iron supplementation alone, do not reduce the risk of ALRI incidence or mortality.

We found insufficient evidence of efficacy of other specific nutritional interventions for ALRI risk reduction. However, the strength of the association between anthropometric indices and ALRI mortality suggests that a range of interventions (not limited to dietary changes) that reduce the prevalence of LBW, underweight or stunting⁷⁹ would be expected to consequentially diminish the burden of ALRI. Rigorous assessments of the effect on ALRI outcomes of improved feeding practices or comprehensive nutrition education programmes should be considered, since multicomponent interventions are most likely to improve overall nutritional status. In future trials, two major methodological issues need to be addressed. First, non-specific outcome ascertainment based on maternal recall or WHO criteria alone may nullify true effects, strongly suggesting that lower respiratory tract morbidity be prospectively assessed by active case-finding and clinical criteria. Second, trials need to have adequate power to detect meaningful intergroup differences in ALRI incidence. Even if a relatively large effect size is expected (e.g. hazard ratio = 0.7), over 800 child-years of follow-up are required in each group (e.g. 800 children treated and monitored for one year), under optimal conditions of a high baseline incidence rate (i.e. 0.3 episodes per child-year), minimal loss to follow-up, and conventional statistical assumptions (i.e.

5% risk of type I error, 80% power). Sufficient sample sizes are of particular importance in trials of multiple micronutrient supplements, food fortification or integrated nutrition programmes, because these trials cannot readily be pooled post-hoc due to non-uniformity of the interventions. The logistical, ethical and financial challenges of conducting large trials with active surveillance emphasizes the importance of building a strong scientific rationale for testing a specific nutritional intervention,⁸⁰ based on translational research (e.g. effects on immune function biomarkers), preliminary dose-finding trials (e.g. efficacy in improving biomarkers of micronutrient status), and qualitative feasibility studies (i.e.

to establish that the intervention is acceptable to communities and could reasonably be scaled-up if shown to be effective).

Conclusion

As with other interventions to prevent ALRI discussed elsewhere in this issue of the *Bulletin*, nutritional interventions such as micronutrient supplementation or breastfeeding promotion can have a broad range of child health benefits.⁷⁹ Implementation and rigorous evaluation of multicomponent nutritional intervention programmes should be a high priority – to reduce the incidence and mortality associated with severe ALRI, as well as the overall disease burden that

persists among children in developing countries. ■

Acknowledgements

We thank the authors of the *Lancet Undernutrition Series* papers for their contribution to the burden of disease estimates, WHO and others for the development of prevalence estimates for the risk factors, the scientists who provided their data for use in pooled analyses, those who conducted the pooled analyses to develop risk estimates, and Rodrigo Dias at the Harvard Initiative for Global Health, who assisted with calculating the burden of nutrition risk factors.

Competing interests: None declared.

Résumé

Infections aiguës des voies respiratoires inférieures chez l'enfant : possibilités de réduire la charge mondiale d'ALRI par des interventions nutritionnelles

La malnutrition et les infections aiguës des voies respiratoires inférieures (ALRI) sont des problèmes sanitaires interdépendants et imbriqués, qui affectent les enfants des pays en développement. Sur la base d'une analyse critique d'essais randomisés étudiant les effets d'interventions nutritionnelles sur la morbidité et la mortalité par ALRI, nous avons conclu que (1) la supplémentation en zinc des populations déficitaires en ce métal prévient environ un quart des épisodes d'ALRI, d'où éventuellement une diminution modeste de la mortalité associée ; (2) la promotion de l'allaitement fait baisser la morbidité due aux ALRI ; (3) la supplémentation en fer ne suffit pas seule à réduire l'incidence des ALRI ; et (4) la supplémentation en vitamine A au-delà de la période néonatale

ne diminue ni l'incidence des ALRI, ni la mortalité qui leur est imputable. Les éléments disponibles à propos des autres interventions nutritionnelles potentiellement bénéfiques étaient insuffisants. Concernant les stratégies bénéficiant d'une justification théorique solide et probablement réalisables sur le plan opérationnel, il conviendrait d'entreprendre des essais rigoureux, avec une recherche des cas active et des échantillons de taille appropriée. On pourrait à présent s'attendre à une baisse de la charge d'ALRI comme conséquence de la promotion permanente de l'allaitement et du passage à l'échelle supérieure des stratégies de supplémentation ou d'enrichissement en zinc, menées parmi les populations cibles.

Resumen

Infecciones agudas de las vías respiratorias inferiores en la niñez: oportunidades para reducir la carga mundial de morbilidad mediante intervenciones nutricionales

Problemas interrelacionados que se solapan, la mala nutrición y las infecciones agudas de las vías respiratorias inferiores (IAVRI) afectan a los niños de los países en desarrollo. A partir de una reseña crítica de los ensayos aleatorizados sobre el efecto de las intervenciones nutricionales en la morbilidad y mortalidad por IAVRI, llegamos a la conclusión de que: (1) la administración de suplementos de zinc en las poblaciones con déficit de ese elemento previene aproximadamente una cuarta parte de los episodios de IAVRI, lo que puede traducirse en una reducción moderada de la mortalidad por IAVRI; (2) el fomento de la lactancia materna reduce la morbilidad por IAVRI; (3) la administración de suplementos de hierro por sí sola no reduce la incidencia de IAVRI; y (4) los suplementos de vitamina A

superado el periodo neonatal no reducen la incidencia de IAVRI ni la mortalidad por esa causa. La evidencia respecto a otras intervenciones nutricionales potencialmente beneficiosas era insuficiente. Cuando las estrategias tengan una justificación teórica sólida y probabilidades de ser operacionalmente viables, deberían emprenderse ensayos rigurosos con procedimientos de búsqueda activa de casos clínicos y tamaños de muestra adecuados. Por el momento, cabe prever una reducción de la carga de IAVRI como consecuencia del fomento continuo de la lactancia materna y de la extensión masiva de las estrategias de uso de zinc como suplemento o como enriquecimiento de los alimentos en las poblaciones objetivo.

ملخص

التهابات الجهاز التنفسي السفلي لدى الأطفال: فرص خفض العبء المرضي العالمي بالتدخلات التغذوية

التنفسي السفلي. (4) إضافة مكملات الفيتامين A بعد مضي الدورة الوليدية لا يقلص حدوث التهاب الجهاز التنفسي السفلي أو الوفيات الناجمة عنه. وهناك بيانات يعتد بها إحصائياً حول المنافع المحتملة للتدخلات التغذوية. وينبغي إجراء تجارب ذات مصداقية عالية وتجر فعال للحالات السريرية مع توفير عينات ذات أحجام مناسبة وذلك من أجل وضع استراتيجيات تركز على أساس نظري قوي وتكون ممكنة التطبيق من الناحية العملية. ويمكن في الوقت الحاضر توقع حدوث نقص في العبء المرضي الناجم عن التهاب الجهاز التنفسي السفلي بفعل التعزيز المتواصل للرضاعة من الثدي وبفعل النهوض باستراتيجية إضافة مكملات الزنك لدى السكان المستهدفين.

ثمة علاقة متبادلة ومتراكبة بين قصور التغذية والتهاب الجهاز التنفسي السفلي وبين المشكلات الصحية التي تصيب الأطفال في البلدان النامية. وبناءً على مراجعة دقيقة للتجارب المعشاة التي أجريت على تأثير التدخلات التغذوية على المراضة والوفيات الناجمة عن التهاب الجهاز التنفسي السفلي، وصلنا إلى الاستنتاجات التالية: (1) إن إضافة مكملات الزنك إلى غذاء السكان الذين يعانون من عوز الزنك يقي ما يقرب من ربع السكان من هجمات التهاب الجهاز التنفسي السفلي، والذي قد يظهر في شكل انخفاض طفيف في الوفيات الناجمة عن التهاب الجهاز التنفسي السفلي. (2) تشجيع وتعزيز الرضاعة من الثدي يقلص المراضة الناجمة عن التهاب الجهاز التنفسي السفلي. (3) إضافة مكملات الحديد لوحده لا تقلص حدوث التهاب الجهاز

References

1. Pelletier DL, Frongillo EA Jr, Habicht JP. Epidemiologic evidence for a potentiating effect of malnutrition on child mortality. *Am J Public Health* 1993;83:1130-3. PMID:8342721
2. Fishman SM, Caulfield LE, de Onis M, Blossner M, Hyder AA, Mullany L, et al. Childhood and maternal underweight. In: Ezzati M, Lopez AD, Rodgers A, Murray CJL, eds. *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*. Geneva: WHO; 2004. pp. 39-161.
3. Black RE, Allen LH, Bhutta ZA, Caulfield LE, de Onis M, Ezzati M, et al. Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet* 2008;371:243-60. PMID:18207566 doi:10.1016/S0140-6736(07)61690-0
4. Rice AL, Sacco L, Hyder A, Black RE. Malnutrition as an underlying cause of childhood deaths associated with infectious diseases in developing countries. *Bull World Health Organ* 2000;78:1207-21. PMID:11100616
5. Victora CG, Kirkwood BR, Ashworth A, Black RE, Rogers S, Sazawal S, et al. Potential interventions for the prevention of childhood pneumonia in developing countries: improving nutrition. *Am J Clin Nutr* 1999;70:309-20. PMID:10479192
6. Lanata CF, Rudan I, Boschi-Pinto C, Tomaskovic L, Cherian T, Weber M, et al. Methodological and quality issues in epidemiological studies of acute lower respiratory infections in children in developing countries. *Int J Epidemiol* 2004;33:1362-72. PMID:15166188 doi:10.1093/ije/dyh229
7. Lopez AD. *Disease control priorities project. Global burden of disease and risk factors*. New York, NY, and Washington, DC: Oxford University Press and World Bank; 2006.
8. Newburg DS, Walker WA. Protection of the neonate by the innate immune system of developing gut and of human milk. *Pediatr Res* 2007;61:2-8. PMID:17211132 doi:10.1203/01.pdr.0000250274.68571.18
9. Moore SE, Collinson AC, N'Gom PT, Prentice AM. Maternal malnutrition and the risk of infection in later life. The impact of maternal nutrition on the offspring. *Nestle Nutr Workshop Ser Pediatr Program* 2005;55:153-67. PMID:16632933
10. Silfverdal SA, Ekholm L, Bodin L. Breastfeeding enhances the antibody response to Hib and pneumococcal serotype 6B and 14 after vaccination with conjugate vaccines. *Vaccine* 2007;25:1497-502. PMID:17097198 doi:10.1016/j.vaccine.2006.10.025
11. Giugliani ERJ, Victora CG. Breastfeeding promotion and infant growth. *Lancet Nutrition Series* 2008. [web appendix #1] Available from: http://www.globalnutritionseries.org/web_appendices [accessed on 12 March 2008].
12. Alvarado R, Zepeda A, Rivero S, Rico N, Lopez S, Diaz S. Integrated maternal and infant health care in the postpartum period in a poor neighborhood in Santiago, Chile. *Stud Fam Plann* 1999;30:133-41. PMID:16617547 doi:10.1111/j.1728-4465.1999.00133.x
13. Burkhalter BR, Marin PS. A demonstration of increased exclusive breastfeeding in Chile. *Int J Gynaecol Obstet* 1991;34:353-9. PMID:1674483 doi:10.1016/0020-7292(91)90604-4
14. Froozani MD, Permezhadeh K, Motlagh AR, Golestan B. Effect of breastfeeding education on the feeding pattern and health of infants in their first 4 months in the Islamic Republic of Iran. *Bull World Health Organ* 1999;77:381-5. PMID:10361754
15. Vitolo MR, Bortolini GA, Feldens CA, Drachler Mde L. Impacts of the 10 steps to healthy feeding in infants: a randomized field trial. *Cad Saude Publica* 2005;21:1448-57. PMID:16158151
16. Kramer MS, Chalmers B, Hodnett ED, Sevkovskaya Z, Dzikovich I, Shapiro S, et al. Promotion of breastfeeding intervention trial (PROBIT): a randomized trial in the Republic of Belarus. *JAMA* 2001;285:413-20. PMID:11242425 doi:10.1001/jama.285.4.413
17. Chandra RK. 1990 McCollum award lecture. Nutrition and immunity: lessons from the past and new insights into the future. *Am J Clin Nutr* 1991;53:1087-101. PMID:1902345
18. Cunningham-Rundles S, McNeeley D, Moon A. Mechanisms of nutrient modulation of the immune response. *J Allergy Clin Immunol* 2005; 115:1119-28. PMID:15940121 doi:10.1016/j.jaci.2005.04.036
19. Bhandari N, Bahl R, Nayyar B, Khokhar P, Rohde J, Bhan MK. Food supplementation with encouragement to feed it to infants from 4 to 12 months of age has a small impact on weight gain. *J Nutr* 2001;131:1946-51. PMID:11435512
20. Sripaipan T, Schroeder DG, Marsh DR, Pachon H, Dearden KA, Ha TT, et al. Effect of an integrated nutrition program on child morbidity due to respiratory infection and diarrhea in northern Viet Nam. *Food Nutr Bull* 2002;23 Suppl;70-7. PMID:12503234
21. English RM, Badcock JC, Giay T, Ngu T, Waters AM, Bennett SA. Effect of nutrition improvement project on morbidity from infectious diseases in preschool children in Vietnam: comparison with control commune. *BMJ* 1997; 315:1122-5. PMID:9374884
22. Villamor E, Fawzi WW. Effects of vitamin A supplementation on immune responses and correlation with clinical outcomes. *Clin Microbiol Rev* 2005; 18:446-64. PMID:16020684 doi:10.1128/CMR.18.3.446-464.2005
23. Potential interventions for the prevention of childhood pneumonia in developing countries: a meta-analysis of data from field trials to assess the impact of vitamin A supplementation on pneumonia morbidity and mortality. The Vitamin A and Pneumonia Working Group. *Bull World Health Organ* 1995;73:609-19. PMID:8846487
24. Sachdev HPS, Gogia S. Vitamin A. *Lancet Nutrition Series* 2008. [web appendix #10] Available from: http://www.globalnutritionseries.org/web_appendices [accessed on 12 March 2008]
25. Humphrey JH, Agoestina T, Wu L, Usman A, Nurachim M, Subardja D, et al. Impact of neonatal vitamin A supplementation on infant morbidity and mortality. *J Pediatr* 1996;128:489-96. PMID:8618182 doi:10.1016/S0022-3476(96)70359-1
26. Coutsooudis A, Adhikari M, Pillay K, Kuhn L, Coovadia HM. Effect of vitamin A supplementation on morbidity of low-birth-weight neonates. *S Afr Med J* 2000;90:730-6. PMID:10985138

27. Tielsch JM, Rahmathullah L, Thulasiraj RD, Katz J, Coles C, Sheeladevi S, et al. Newborn vitamin A dosing reduces the case fatality but not incidence of common childhood morbidities in South India. *J Nutr* 2007;137:2470-4. PMID:17951487
28. Ni J, Wei J, Wu T. Vitamin A for non-measles pneumonia in children. *Cochrane Database Syst Rev* 2005; (3):CD003700. PMID:16034908
29. Brown N, Roberts C. Vitamin A for acute respiratory infection in developing countries: a meta-analysis. *Acta Paediatr* 2004;93:1437-42. PMID:15513568 doi:10.1080/08035250410022143
30. Oppenheimer SJ. Iron and its relation to immunity and infectious disease. *J Nutr* 2001;131:616S-35S. PMID:11160594
31. Gera T, Sachdev HP. Effect of iron supplementation on incidence of infectious illness in children: systematic review. *BMJ* 2002;325:1142. PMID:12433763 doi:10.1136/bmj.325.7373.1142
32. Sachdev HPS, Gogia S. Iron. *Lancet Nutrition Series* 2008. [web appendix #8] Available from: http://www.globalnutritionseries.org/web_appendices [accessed on 12 March 2008].
33. Strand TA, Taneja S, Bhandari N, Refsum H, Ueland PM, Gjessing HK, et al. Folate, but not vitamin B-12 status, predicts respiratory morbidity in north Indian children. *Am J Clin Nutr* 2007;86:139-44. PMID:17616773
34. Tielsch JM, Khatri SK, Stoltzfus RJ, Katz J, LeClerq SC, Adhikari R, et al. Effect of routine prophylactic supplementation with iron and folic acid on preschool child mortality in southern Nepal: community-based, cluster-randomised, placebo-controlled trial. *Lancet* 2006;367:144-52. PMID:16413878 doi:10.1016/S0140-6736(06)67963-4
35. Maggini S, Wintergerst ES, Beveridge S, Hornig DH. Selected vitamins and trace elements support immune function by strengthening epithelial barriers and cellular and humoral immune responses. *Br J Nutr* 2007;98 Suppl 1:S29-35. PMID:17922955 doi:10.1017/S0007114507832971
36. Muhe L, Lulseged S, Mason KE, Simoes EA. Case-control study of the role of nutritional rickets in the risk of developing pneumonia in Ethiopian children. *Lancet* 1997;349:1801-4. PMID:9269215 doi:10.1016/S0140-6736(96)12098-5
37. Najada AS, Habashneh MS, Khader M. The frequency of nutritional rickets among hospitalized infants and its relation to respiratory diseases. *J Trop Pediatr* 2004;50:364-8. PMID:15537725 doi:10.1093/tropej/50.6.364
38. Wayne V, Yousafzai A, Mogale K, Filteau S. Association of subclinical vitamin D deficiency with severe acute lower respiratory infection in Indian children under 5 years. *Eur J Clin Nutr* 2004;58:563-7. PMID:15042122 doi:10.1038/sj.ejcn.1601845
39. Karatekin G, Kaya A, Salihoğlu O, Balci H, Nuhoğlu A. Association of subclinical vitamin D deficiency in newborns with acute lower respiratory infection and their mothers. *European Journal of Clinical Nutrition* 2007. E-published.
40. Roth DE, Jones AB, Prosser C, Robinson JL, Vohra S. Vitamin D status is not associated with the risk of hospitalization for acute bronchiolitis in early childhood. *European Journal of Clinical Nutrition* 2007. E-published.
41. Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutik SR, et al. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* 2006;311:1770-3. PMID:16497887 doi:10.1126/science.1123933
42. Wuehler SE, Peerson JM, Brown KH. Use of national food balance data to estimate the adequacy of zinc in national food supplies: methodology and regional estimates. *Public Health Nutr* 2005;8:812-9. PMID:16277796 doi:10.1079/PHN2005724
43. Shankar AH, Prasad AS. Zinc and immune function: the biological basis of altered resistance to infection. *Am J Clin Nutr* 1998;68 Suppl:447S-63S. PMID:9701160
44. Aggarwal R, Sentz J, Miller MA. Role of zinc administration in prevention of childhood diarrhea and respiratory illnesses: a meta-analysis. *Pediatrics* 2007;119:1120-30. PMID:17545379 doi:10.1542/peds.2006-3481
45. Brooks WA, Santosham M, Naheed A, Goswami D, Wahed MA, Diener-West M, et al. Effect of weekly zinc supplements on incidence of pneumonia and diarrhoea in children younger than 2 years in an urban, low-income population in Bangladesh: randomised controlled trial. *Lancet* 2005;366:999-1004. PMID:16168782 doi:10.1016/S0140-6736(05)67109-7
46. Osendarp SJ, Santosham M, Black RE, Wahed MA, van Raaij JM, Fuchs GJ. Effect of zinc supplementation between 1 and 6 mo of life on growth and morbidity of Bangladeshi infants in urban slums. *Am J Clin Nutr* 2002; 76:1401-8. PMID:12450909
47. Long KZ, Montoya Y, Hertzmark E, Santos J, Rosado J. A double-blind, randomized, clinical trial of the effect of vitamin A and zinc supplementation on diarrheal disease and respiratory tract infections in children in Mexico City, Mexico. *Am J Clin Nutr* 2006;83:693-700. PMID:16522919
48. Richard SA, Zavaleta N, Caulfield LE, Black RE, Witzig RS, Shankar AH. Zinc and iron supplementation and malaria, diarrhea, and respiratory infections in children in the Peruvian Amazon. *Am J Trop Med Hyg* 2006;75:126-32. PMID:16837718
49. Luabeya KK, Mpontshane N, Mackay M, Ward H, Elson I, Chhagan M, et al. Zinc or multiple micronutrient supplementation to reduce diarrhea and respiratory disease in South African children: a randomized controlled trial. *PLoS ONE* 2007;2:e541. PMID:17593956 doi:10.1371/journal.pone.0000541
50. Tielsch JM, Khatri SK, Stoltzfus RJ, Katz J, LeClerq SC, Adhikari R, et al. Effect of daily zinc supplementation on child mortality in southern Nepal: a community-based, cluster randomised, placebo-controlled trial. *Lancet* 2007;370:1230-9. PMID:17920918 doi:10.1016/S0140-6736(07)61539-6
51. Smith PG, Morrow RH, eds. *Field trials of health interventions in developing countries: a toolbox*. London: Macmillan Education; 1996.
52. Coles CL, Bose A, Moses PD, Mathew L, Agarwal I, Mammen T, et al. Infectious etiology modifies the treatment effect of zinc in severe pneumonia. *Am J Clin Nutr* 2007;86:397-403. PMID:17684211
53. Brooks WA, Yunus M, Santosham M, Wahed M, Nahar K, Yeasmin S, et al. Zinc for severe pneumonia in very young children: double-blind placebo-controlled trial. *Lancet* 2004;363:1683-8. PMID:15158629 doi:10.1016/S0140-6736(04)16252-1
54. Bose A, Coles CL, Gunavathi, John H, Moses P, Raghupathy P, Kirubakaran C et al. Efficacy of zinc in the treatment of severe pneumonia in hospitalized children < 2 y old. *Am J Clin Nutr* 2006;83:1089-96. PMID:16685051
55. Mahalanabis D, Lahiri M, Paul D, Gupta S, Gupta A, Wahed MA, et al. Randomized, double-blind, placebo-controlled clinical trial of the efficacy of treatment with zinc or vitamin A in infants and young children with severe acute lower respiratory infection. *Am J Clin Nutr* 2004;79:430-6. PMID:14985218
56. Sivakumar B, Nair KM, Sreeramulu D, Suryanarayana P, Ravinder P, Shatrugna V, et al. Effect of micronutrient supplement on health and nutritional status of schoolchildren: biochemical status. *Nutrition* 2006;22 Suppl:S15-25. PMID:16426959 doi:10.1016/j.nut.2005.07.012
57. Stoltzfus R, Dreyfuss M. *Guidelines for the use of iron supplements to prevent and treat iron deficiency anemia*. International Nutritional Anemia Consultative Group, WHO and UNICEF. Washington, DC: International Life Sciences Institute; 1998.
58. Sazawal S, Black RE, Ramsan M, Chhaya HM, Stoltzfus RJ, Dutta A, et al. Effects of routine prophylactic supplementation with iron and folic acid on admission to hospital and mortality in preschool children in a high malaria transmission setting: community-based, randomised, placebo-controlled trial. *Lancet* 2006;367:133-43. PMID:16413877 doi:10.1016/S0140-6736(06)67962-2
59. Bhandari N, Taneja S, Mazumder S, Bahl R, Fontaine O, Bhan MK; Zinc Study Group. Adding zinc to supplemental iron and folic acid does not affect mortality and severe morbidity in young children. *J Nutr* 2007;137:112-7. PMID:17182810
60. Sazawal S, Dhingra U, Dhingra P, Hiremath G, Kumar J, Sarkar A, et al. Effects of fortified milk on morbidity in young children in north India: community based, randomised, double masked placebo controlled trial. *BMJ* 2007;334:140. PMID:17132678 doi:10.1136/bmj.39035.482396.55
61. Bégin F, Santizo MC, Peerson JM, Torún B, Brown KH. Effects of bovine serum concentrate, with or without supplemental micronutrients, on the growth, morbidity, and micronutrient status of young children in a low-income, peri-urban Guatemalan community. *Eur J Clin Nutr* 2008;62:39-50. PMID:17299460 doi:10.1038/sj.ejcn.1602682
62. Penny ME, Marin RM, Duran A, Peerson JM, Lanata CF, Lönnerdal B, et al. Randomized controlled trial of the effect of daily supplementation with zinc or multiple micronutrients on the morbidity, growth, and micronutrient status of young Peruvian children. *Am J Clin Nutr* 2004;79:457-65. PMID:14985222
63. Wieringa FT, Berger J, Dijkhuizen MA, Hidayat A, Ninh NX, Utomo B, et al.; for the SEAMTIZI. (South-East Asia Multi-country Trial on Iron and Zinc supplementation in Infants) Study Group. Combined iron and zinc supplementation in infants improved iron and zinc status, but interactions reduced efficacy in a multicountry trial in southeast Asia. *J Nutr* 2007; 137:466-71. PMID:17237328

64. Chandra RK. Serum thymic hormone activity and cell-mediated immunity in healthy neonates, preterm infants, and small-for-gestational age infants. *Pediatrics* 1981;67:407-11. PMID:6972515
65. Ferro-Luzzi A, Ashworth A, Martorell R, Scrimshaw N. Report of the IDECG working group on effects of IUGR on infants, children and adolescents: immunocompetence, mortality, morbidity, body size, body composition, and physical performance. *Eur J Clin Nutr* 1998;52 Suppl 1:S97-9. PMID:9511026
66. Saha K, Kaur P, Srivastav G, Chaudhury DS. A six-months' follow-up study of growth, morbidity and functional immunity in low birth weight neonates with special reference to intrauterine growth retardation in small-for-gestational-age infants. *J Trop Pediatr* 1983;29:278-82. PMID:6685774
67. Harding R, Tester ML, Moss TJ, Davey MG, Louey S, Joyce B, et al. Effects of intra-uterine growth restriction on the control of breathing and lung development after birth. *Clin Exp Pharmacol Physiol* 2000;27:114-9. PMID:10696539 doi:10.1046/j.1440-1681.2000.03191.x
68. Rees S, Ng J, Dickson K, Nicholas T, Harding R. Growth retardation and the development of the respiratory system in fetal sheep. *Early Hum Dev* 1991; 26:13-27. PMID:1914984 doi:10.1016/0378-3782(91)90039-6
69. Hoo AF, Stocks J, Lum S, Wade AM, Castle RA, Costeloe KL, et al. Development of lung function in early life: influence of birth weight in infants of nonsmokers. *Am J Respir Crit Care Med* 2004;170:527-33. PMID:15172896 doi:10.1164/rccm.200311-15520C
70. Kramer MS. Determinants of low birth weight: methodological assessment and meta-analysis. *Bull World Health Organ* 1987;65:663-737. PMID:3322602
71. Ramakrishnan U. Nutrition and low birth weight: from research to practice. *Am J Clin Nutr* 2004;79:17-21. PMID:14684392
72. Wilcox AJ. On the importance – and the unimportance – of birthweight. *Int J Epidemiol* 2001;30:1233-41. PMID:11821313 doi:10.1093/ije/30.6.1233
73. Christian P, West KP, Khatri SK, Leclercq SC, Pradhan EK, Katz J, et al. Effects of maternal micronutrient supplementation on fetal loss and infant mortality: a cluster-randomized trial in Nepal. *Am J Clin Nutr* 2003;78:1194-202. PMID:14668283
74. Osendarp SJ, van Raaij JM, Darmstadt GL, Baqui AH, Hautvast JG, Fuchs GJ. Zinc supplementation during pregnancy and effects on growth and morbidity in low birthweight infants: a randomised placebo controlled trial. *Lancet* 2001;357:1080-5. PMID:11297959 doi:10.1016/S0140-6736(00)04260-4
75. Christian PS, Darmstadt GL, Wu L, Khatri SK, LeClerq SC, Katz J, et al. The impact of maternal micronutrient supplementation on early neonatal morbidity in rural Nepal: a randomized, controlled, community trial. *Arch Dis Child Fetal Neonatal Ed* 2007; 3. E-published.
76. Schmidt MK, Muslimatun S, Schultink W, West CE, Hautvast JG. Randomised double-blind trial of the effect of vitamin A supplementation of Indonesian pregnant women on morbidity and growth of their infants during the first year of life. *Eur J Clin Nutr* 2002;56:338-46. PMID:11965510 doi:10.1038/sj.ejcn.1601318
77. Baqui AH, Black RE, El Arifeen S, Yunus M, Chakraborty J, Ahmed S, et al. Effect of zinc supplementation started during diarrhoea on morbidity and mortality in Bangladeshi children: community randomised trial. *BMJ* 2002; 325:1059. PMID:12424162 doi:10.1136/bmj.325.7372.1059
78. Bhandari N, Mazumder S, Taneja S, Dube B, Agarwal D, Fontaine O, et al. The effectiveness of zinc supplementation plus ORS compared to ORS alone as a treatment acute diarrhea in a primary health care setting: a cluster randomised trial. *Pediatrics* 2008. In press.
79. Bhutta ZA, Ahmed T, Black RE, Cousens S, Dewey K, Giugliani E, et al. What works? Interventions for maternal and child undernutrition and survival. *Lancet* 2008;371:417-40. PMID:18206226 doi:10.1016/S0140-6736(07)61693-6
80. Prentice AM, Ghattas H, Cox SE. Host–pathogen interactions: can micronutrients tip the balance? *J Nutr* 2007;137:1334-7. PMID:17449601
81. Baqui AH, Darmstadt GL, Williams EK, Kumar V, Kiran TU, Panwar D, et al. Rates, timing and causes of neonatal deaths in rural India: implications for neonatal health programmes. *Bull World Health Organ* 2006;84:706-13. PMID:17128340 doi:10.2471/BLT.05.026443