

Acute visceral pain relief mediated by A₃AR agonists in rats: involvement of N-type voltage-gated calcium channels

Elena Lucarini^a, Elisabetta Coppi^a, Laura Micheli^a, Carmen Parisio^a, Alessia Vona^a, Federica Cherchi^a, Anna M. Pugliese^a, Felicità Pedata^a, Paola Failli^a, Seph Palomino^b, Jared Wahl^b, Tally M. Largent-Milnes^b, Todd W. Vanderah^b, Dilip K. Tosh^c, Kenneth A. Jacobson^c, Daniela Salvemini^{d,e}, Carla Ghelardini^a, Lorenzo Di Cesare Mannelli^{a,*}

Abstract

Pharmacological tools for chronic visceral pain management are still limited and inadequate. A₃ adenosine receptor (A₃AR) agonists are effective in different models of persistent pain. Recently, their activity has been related to the block of N-type voltage-gated Ca²⁺ channels (Ca_v2.2) in dorsal root ganglia (DRG) neurons. The present work aimed to evaluate the efficacy of A₃AR agonists in reducing postinflammatory visceral hypersensitivity in both male and female rats. Colitis was induced by the intracolonic instillation of 2,4-dinitrobenzenesulfonic acid (DNBS; 30 mg in 0.25 mL 50% EtOH). Visceral hypersensitivity was assessed by measuring the visceromotor response and the abdominal withdrawal reflex to colorectal distension. The effects of A₃AR agonists (MRS5980 and CI-IB-MECA) were evaluated over time after DNBS injection and compared to that of the selective Ca_v2.2 blocker PD173212, and the clinically used drug linaclotide. A₃AR agonists significantly reduced DNBS-evoked visceral pain both in the postinflammatory (14 and 21 days after DNBS injection) and persistence (28 and 35 days after DNBS) phases. Efficacy was comparable to effects induced by linaclotide. PD173212 fully reduced abdominal hypersensitivity to control values, highlighting the role of Ca_v2.2. The effects of MRS5980 and CI-IB-MECA were completely abolished by the selective A₃AR antagonist MRS1523. Furthermore, patch-clamp recordings showed that A₃AR agonists inhibited Ca_v2.2 in dorsal root ganglia neurons isolated from either control or DNBS-treated rats. The effect on Ca²⁺ current was PD173212-sensitive and prevented by MRS1523. A₃AR agonists are effective in relieving visceral hypersensitivity induced by DNBS, suggesting a potential therapeutic role against abdominal pain.

Keywords: A₃ adenosine receptor, DRG, Abdominal pain, IBS, IBD

1. Introduction

Visceral pain management is a major clinical problem, because of the lack of effective and safe drugs.¹¹ Abdominal pain is the most common form of visceral hypersensitivity and is often the result of inflammatory bowel diseases (IBDs).^{14,51,84,85} The poor correlations between reported abdominal pain intensity and IBD activity

raise close similarities with the irritable bowel syndrome (IBS), a symptom-based clinical condition defined by the presence of persistent abdominal pain and altered bowel habits, in the absence of any other disease able to account for these symptoms.^{24,35,51,58,84,85} Frequently, IBS results from a previous intestinal damage caused by severe infections or prolonged inflammatory processes.^{24,83} In the inflamed gut, there is breakdown of intestinal barrier function, abnormal secretion, changes in the patterns of motility, and altered visceral sensation, which altogether contribute to generation of symptoms (diarrhoea, cramping, and pain). Pain, in particular, persists beyond the relief of inflammation, thus revealing a peculiar kind of chronic hypersensitivity thought to be due to inflammatory, immune, and neuropathic mechanisms.^{9,26,40,53,86} Currently, the most efficacious therapies against visceral hypersensitivity are mainly directed toward treating bowel dysfunction, whereas drugs able to directly target the associated pain are still unsatisfactory.¹¹

The neuromodulator adenosine exerts potent and long-lasting pain suppression in preclinical models as well as in human studies.¹⁰⁵ Moreover, adenosinergic signalling is known to modulate intestinal functionality.²⁻⁴ For decades, it was thought that the analgesic effects of adenosine were mediated by A₁ adenosine receptor (A₁AR) activation,^{8,54,65,77,105} but research efforts over the past several years have also implicated a key role for the A₃AR subtype.^{43,75} For example, A₃AR agonists are able to block the development of trauma- and chemotherapeutic-

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

^a Department of Neuroscience, Psychology, Drug Research and Child Health-Neurofarba-Division of Pharmacology and Toxicology, Florence, Italy, ^b Department of Pharmacology, University of Arizona, College of Medicine Tucson, Tucson, AZ, United States, ^c Molecular Recognition Section, Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD, United States, ^d Department of Pharmacology and Physiology, Saint Louis University School of Medicine, St. Louis, MO, United States, ^e Henry and Amelia Nasrallah Center for Neuroscience, Saint Louis University, St. Louis, MO, United States

*Corresponding author. Address: Department of Neuroscience, Psychology, Drug Research and Child Health-Neurofarba-Pharmacology and Toxicology Section, University of Florence, Viale Pieraccini 6, 50139 Florence, Italy. Tel.: 00390552758395. E-mail address: lorenzo.mannelli@unifi.it (L. Di Cesare Mannelli).

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.painjournalonline.com).

PAIN 161 (2020) 2179–2190

© 2020 International Association for the Study of Pain

<http://dx.doi.org/10.1097/j.pain.0000000000001905>

induced neuropathic pain^{17,42} and are also effective in reducing inflammatory and cancer-related pain.^{50,93,100,101} Recently, we found that selective A₃AR stimulation inhibits the opening of N-type voltage-gated Ca²⁺ channels (Ca_v2.2) and decreases the electrically evoked excitation of isolated rat dorsal root ganglia (DRG) neurons. This mechanism could explain the antihyperalgesic effect of A₃AR agonists across different models.¹⁹ Therapeutic targeting of A₁AR has been limited to severe cardiovascular side effects; by contrast, A₃AR agonists are already in advanced clinical trials for different indications with a good safety profile.^{29,80}

The aims of this study were to: (1) evaluate the efficacy of selective A₃AR agonists in postinflammatory visceral hypersensitivity induced in male and female rats by the intrarectal administration of 2,4-dinitrobenzenesulfonic acid (DNBS); (2) investigate the electrophysiological effects of selective A₃AR stimulation on DRG neurons isolated from control and DNBS-treated animals.

2. Materials and methods

An extensive description of materials and methods has been reported in the supplemental digital content (available at <http://links.lww.com/PAIN/B10>).

2.1. Animals

We used male and female Sprague-Dawley rats (Envigo, Varese, Italy) weighing approximately 220 to 250 g at the beginning of the experimental procedure. All animal manipulations were conducted according to the Directive 2010/63/EU of the European Parliament and of the European Union Council (September 22, 2010) on the protection of animals used for scientific purposes. The ethical policy of the University of Florence complies with the Guide for the Care and Use of Laboratory Animals of the US National Institutes of Health (NIH Publication No. 85-23, revised 1996; University of Florence assurance number: A5278-01). Formal approval to conduct the described experiments was obtained from the Animal Subjects Review Board of the University of Florence. Experiments involving animals have been reported according to ARRIVE guidelines.⁵⁷ Gastrointestinal motility assays (GI) were conducted using Sprague-Dawley male rats (Envigo, 250-350 g) in accordance with the University of Arizona Institutional Animal Care and Use Committee (approval 06-110). Experiments were performed on male rats when not otherwise stated.

2.2. Determination of female estrous cycle phases

Estrous cycle was assessed in rats by analysing rat vaginal smears as previously described.³⁹ This screening was performed before the behavioral test; tests were selectively performed in female rats in estrous/proestrous phase to avoid sensitive difference related to the estrous cycle. Moreover, in most studies, this phase is reported to be related to the highest pain sensitivity,^{45,46,59,60,74} therefore suitable for studying pain-relieving compounds.

2.3. Induction of colitis

Colitis was induced using the method described previously by Fornai et al.³⁰ with minor changes. In brief, during a short period of anaesthesia with isoflurane (2%), 30 mg of DNBS in 0.25 mL of 50% ethanol was administered intrarectally through a polyethylene PE-60

catheter inserted 8 cm proximal to the anus. Control rats received 0.25 mL of saline solution. Subsequent experimental procedures began 14 days after DNBS administration to allow for recovery from the initial colonic inflammation.

2.4. Assessment of visceral sensitivity by visceromotor response and abdominal withdrawal reflex

The visceromotor response (VMR) to colorectal distension (CRD) was used as an objective measure of visceral sensitivity. Visceromotor response assessment was conducted in animals under light anaesthesia (2% isoflurane) 14 and 21 days after DNBS administration. The amplitude of the abdominal contraction consequent to colorectal stimulation (a balloon inflated with 0.5, 1, 2, and 3 mL referred to as distension volume) was quantified by electromyography recordings as reported by Parisio et al.⁶⁴ Behavioural responses to CRD (0.5, 1, 2, and 3 mL referred to as distension volume) were assessed through abdominal withdrawal reflex (AWR) measurement in conscious animals by using a semiquantitative score (0-4) as described previously.¹⁶ The measurements were conducted 14, 21, 28, and 35 days after DNBS administration.

2.5. Upper gastrointestinal transit treatment and harvesting

All experiments were performed as previously described.^{66,76,95} Before experiment, rats were fasted for 12 hours, maintaining the access to water ad libitum. After 15 minutes had elapsed, rats were then given an oral gavage with fluorescein isothiocyanate dextran (300 µL of 5 mg/mL 70 kDa FITC-dextran; Sigma, Milan, Italy). Forty min after completing the gavage, rats were anaesthetized with isoflurane and whole blood harvested through cardiac puncture. After blood harvest, the upper portion of the gastrointestinal tract was carefully removed, being cut proximally at the pyloric sphincter and distally at the cecum. The intestine was sectioned into 10 equal lengths, which were homogenized in 1 mL 1X PBS and then centrifugated. Fluorescence was determined in the supernatants collected from intestinal lining (10 µL) and content diluted into 90 µL 1X PBS; serum fluorescence was determined after dilution of 10 µL serum into 40 µL 1X PBS using a ClarioStar (BMG Labtech, Milan, Italy) at 483 to 530 nm. Readings were then collected and prepared for statistical analyses.

2.6. Drug administration

The selective A₃AR agonist, 2-chloro-N⁶-(3-iodobenzyl)-adenosine-5'-N-methyluronamide (Cl-IB-MECA; Tocris Bioscience, Bristol, United Kingdom), and the selective A₃ receptor antagonist, 3-propyl-6-ethyl-5-[(ethylthio)carbonyl]-2-phenyl-4-propyl-3-pyridine carboxylate (MRS1523; Sigma-Aldrich, Milan, Italy), were dissolved in 5% DMSO and 5% TWEEN 20 saline solution for *in vivo* administration. The new, highly selective (10,000-fold vs each of the other 3 receptor subtypes) A₃AR agonist, MRS5980, (1S,2R,3S,4R,5S)-4-(2-((5-chlorothiophen-2-yl)ethynyl)-6-(methylamino)-9H-purin-9-yl)-2,3-dihydroxy-N-methylbicyclo[3.1.0]hexane-1-carboxamide⁸⁹ was dissolved in 5% DMSO saline solution for *in vivo* administration. As already reported,⁵⁰ the antiallodynic effects of the A₃AR agonists used in this work were lost in the A₃AR knockout mouse and in the presence of the A₃AR antagonist MRS1523.⁴⁹ N-[[4-(1,1-dimethylethyl)phenyl]methyl]-N-methyl-L-leucyl-N-(1,1-dimethylethyl)-O-(phenylmethyl)-L-tyrosinamide (PD173232) was purchased from Alomone Labs (Jerusalem, Israel) and dissolved in 5% DMSO and 5% TWEEN 20 saline solution for

in vivo administration. Both the vehicle and the DNBS control groups received an i.p. injection of the vehicle used for the administration of the respective molecules. The acute injection of the vehicles (5% DMSO or 5% DMSO + 5% TWEEN 20) did not alter visceral pain threshold and did not induce side effects in the animals. The guanylate cyclase-C (GC-C) agonist linaclotide (Allergan, Buckinghamshire, United Kingdom) was dissolved in water and orally administered 1 hour before starting the behavioural tests.²⁵ CI-IB-MECA, MRS5980, and PD173232 were administered i.p. 15 minutes before the test. MRS1523 was administered i.p. 15 minutes before CI-IB-MECA and MRS5980. Gastrointestinal transit study: the compound MRS5980 was dissolved in 5% DMSO, and then suspended in saline. MRS5980 and vehicle were administered through i.p. injection to all animals in study at 2.4 $\mu\text{mol}\cdot\text{kg}^{-1}$ body weight (1 mL/kg injection volume). Seventy kDa FITC-dextran (Sigma) was dosed by oral gavage at a concentration of 5 mg/mL, 300 μL per rat. Additional pharmacological characteristics of A_3 AR agonists are shown in the Supplemental material (available at <http://links.lww.com/PAIN/B10>).

2.7. Cell cultures

Primary DRG neurons (related to colon-sensitive innervations: T12, T13, L4, L5, L6, S1, S2) were isolated from vehicle- or DNBS-treated (at 14th day) rats and cultured for 1 to 2 days before being used for experiments as described.¹⁹

2.8. Electrophysiology

Whole-cell patch-clamp recordings were performed as previously described.¹⁹ Passive membrane properties of DRG neurons isolated from control or DNBS-treated rats were investigated under physiological-like conditions by using the following K-gluconate-based pipette solution (mM): K₂gluconate 130; NaCl 4.8; KCl 10; MgCl₂ 2; CaCl₂ 1; Na₂-ATP 2; Na₂-GTP 0.3; EGTA 3; HEPES 10 (pH 7.4 with KOH). Resting membrane potential (V_m) was measured immediately after seal breakthrough by switching the amplifier to the current-clamp mode. The calculated liquid junction potential for K-gluconate pipettes in our experimental conditions was 15.0 mV, and V_m values reported in the present research have been corrected accordingly.

Voltage-dependent Ca^{2+} currents (VDCCs) were recorded by using a Cs⁺-based pipette solution having the following composition (mM): CsCl (130); NaCl (4.8); KCl (10); MgCl₂ (2); CaCl₂ (1); Na₂-ATP (2); Na₂-GTP (0.3); EGTA (3); and HEPES (10—pH 7.4 with CsOH). The extracellular solution was (in mM): NaCl (147); CsCl (4); MgCl₂ (1); and CaCl₂ (5); HEPES (10); D-glucose (10); pH 7.4 with NaOH. Tetrodotoxin (TTX; 1 μM) and 5-(4-butoxy-3-chlorophenyl)-N-[[2-(4-morpholinyl)-3-pyridinyl]methyl]-3-pyridinecarboxamide (A887826; 200 nM) were added to the extracellular solution to block TTX-sensitive $\text{Na}_v1.1$, 1.2, 1.3, 1.4, 1.6, 1.7 channels and TTX-resistant $\text{Na}_v1.8$, respectively. VDCC currents were evoked by a 0 mV step depolarization (200 ms) once every 30 seconds to minimize Ca^{2+} current run down. Peak Ca^{2+} current (I_{Ca} peak) was measured as the peak current amplitude reached during the first 50 ms of voltage step. Steady-state Ca^{2+} current (I_{Ca} st-state) was measured as the averaged current amplitude measured between 160 and 190 ms of voltage step. The current-to-voltage relationship (I-V plot) of Ca^{2+} currents was obtained by eliciting 10 depolarizing voltage steps (200-ms duration, 10-mV increments, 5-second interval) from -50 to $+50$ mV starting from a holding potential (V_h) of -65 mV.

Data were acquired with an Axopatch 200B amplifier (Axon Instruments, San Jose, CA), low-pass filtered at 10 kHz, and stored and analysed with pClamp 9.2 software (Axon Instruments). Membrane resistance (R_m) and membrane capacitance (C_m) were routinely measured by fast hyperpolarizing voltage pulses (from -60 to -70 mV, 40-ms duration). Averaged currents were normalized to cell capacitance and expressed as pA/pF.

Cell capacitance was used to estimate neuronal diameter by assuming an approximated spherical cell shape according to the calculated C_m for all biological membranes of 1 $\mu\text{F}/\text{cm}^2$ and to the equation of the sphere surface: $A = 4\pi r^2$.

The in vitro concentrations were chosen on the base of our previous work: the A_3 AR agonists CI-IB-MECA and MRS5980 were applied at 30 nM, a concentration able to produce maximal inhibition of Ca^{2+} currents in rat DRG neurons.¹⁹

2.9. Statistical analysis

Behavioural measurements were performed on 6 animals for each treatment. All the experimental procedures were performed by a researcher blind to the treatment. Results were expressed as mean \pm SEM. The analysis of variance (ANOVA) of behavioural data was performed by one-way ANOVA with Bonferroni's significant difference procedure used for post hoc comparisons. Statistics of electrophysiological data was performed by Student paired or unpaired t test or by one-way ANOVA followed by Bonferroni's post hoc test, as appropriate. P values of less than 0.05 were considered significant. Data were analyzed using the "Origin 9" software (OriginLab, Northampton, MA). Data from gastrointestinal transit measures were analyzed in GraphPad Prism 7. Standard curves were fit using a nonlinear, hyperbolic equation for individual plates. A two-way ANOVA with Tukey post hoc test was performed using the variables of GI segment and treatment. Serum samples were compared using an unpaired t test. Data were considered statistically significant when $P < 0.05$ to detect 20% difference with 80% power (necessary n -values determined using G.Power3.1).

3. Results

3.1. A_3 AR agonists reduce colitis-induced visceral hypersensitivity in rats

The visceromotor response and the abdominal withdrawal reflex (VMR and AWR) to CRD were measured by using progressively increasing balloon volumes as pressuring stimuli on the colon (highest volume: 3 mL, to avoid tissue damage). Fourteen and 21 days after DNBS injection, both VMR and AWR were significantly higher in comparison to controls (vehicle + vehicle) as shown, respectively, in **Figures 1A–B** (day 14) and **1C–D** (day 21). On day 14, the acute administration of MRS5980 (0.3, 1.2, and 2.4 $\mu\text{mol}\cdot\text{kg}^{-1}$; i.p.) dose-dependently reduced the postinflammatory visceral hypersensitivity induced by DNBS; the magnitude of the reduction was similar for VMR and AWR. The highest dose (2.4 $\mu\text{mol}\cdot\text{kg}^{-1}$) completely reversed the sensitivity alteration back to the value of control rats. MRS5980 1.2 $\mu\text{mol}\cdot\text{kg}^{-1}$ partially but significantly reduced the response of the animals to CRD. The lowest dose of MRS5980 (0.3 $\mu\text{mol}\cdot\text{kg}^{-1}$) was ineffective in both tests (**Figs. 1A and B**). On day 21, the effect of MRS5980 (0.3, 1.2, and 2.4 $\mu\text{mol}\cdot\text{kg}^{-1}$) was confirmed (VMR and AWR to CRD, **Figs. 1C and D**, respectively).

Figure 2A shows the result of the pretreatment with the selective A_3 AR antagonist MRS1523⁴⁹ on the antihyperalgesic effect of MRS5980 on day 14. MRS1523 (5 $\mu\text{mol}\cdot\text{kg}^{-1}$) completely




Figure 1. Effect of MRS5980 on postinflammatory visceral pain induced by DNBS. Effect of MRS5980 (0.3, 1.2, and 2.4 $\mu\text{mol}\cdot\text{kg}^{-1}$; i.p.) on visceromotor response (VMR) to CRD (left column) and abdominal withdrawal reflex (right column) on day 14 (A and B) and day 21 (C and D) after DNBS-induced colonic inflammation. Each value is the mean \pm SEM of 6 rats per group. ^^ $P < 0.01$ vs vehicle-treated normal controls (blue). * $P < 0.05$ and ** $P < 0.01$ vs DNBS + vehicle-treated group (red). CRD, colorectal distension.

abolished the pain-relieving effect of MRS5980 (2.4 $\mu\text{mol}\cdot\text{kg}^{-1}$) confirming a A_3 AR mechanism of action. Furthermore, the pain-relieving effect evoked by A_3 AR stimulation was confirmed by using another selective A_3 AR agonist, CI-IB-MECA. On day 14, the acute administration of CI-IB-MECA dose-dependently relieved visceral pain in DNBS-treated animals, reducing their VMR to the value of controls (dosed at 1.2 $\mu\text{mol}\cdot\text{kg}^{-1}$, **Fig. 2B**). The lowest dose of 0.3 $\mu\text{mol}\cdot\text{kg}^{-1}$ evoked a weaker effect only with the 2 mL stimulus. The CI-IB-MECA (1.2 $\mu\text{mol}\cdot\text{kg}^{-1}$) effect was blocked by MRS1523 (5 $\mu\text{mol}\cdot\text{kg}^{-1}$, **Fig. 2B**).

3.2. N-type voltage-gated Ca^{2+} channel block relieves colitis-induced visceral pain in rats

A_3 AR agonists have been recently reported to inhibit $\text{Ca}_v2.2$ -mediated currents in DRG neurons, thus suggesting a possible mechanism for pain relief.¹⁹ To verify this hypothesis, we evaluated the effect of acute administration of the selective N-type $\text{Ca}_v2.2$ blocker PD173212³⁸ in our visceral pain model. The test (VMR assessment) was performed on day 14 after DNBS injection. As shown in **Figure 3C**, PD173212 (0.0017–1.7 $\mu\text{mol}\cdot\text{kg}^{-1}$, i.p.) dose-dependently reduced the visceral hypersensitivity induced by DNBS. The compound started to be effective at a dose of 0.017 $\mu\text{mol}\cdot\text{kg}^{-1}$ and completely relieved abdominal pain when administered at a ten-fold higher dose (**Fig. 2C**).

3.3. Comparison between the effects of MRS5980 and linaclotide in male and female rats

The effects of MRS5980 against the postinflammatory hypersensitivity induced by DNBS were evaluated in both male and female

rats and compared with the effects of linaclotide, a guanylate cyclase-C (GC-C) agonist that represents the reference drug in the management of pain in patients affected by IBS with predominant constipation (IBS-C).^{18,97} On day 14, 0.2 and 0.066 $\text{nmol}\cdot\text{kg}^{-1}$ linaclotide strongly reduced the VMR to CRD in male DNBS-treated animals, whereas the ten-fold lower dose was ineffective (**Fig. 2D**).

Similar to males, 14 days after DNBS injection in female rats, both VMR and AWR to colorectal distension were significantly higher in comparison to controls (**Figs. 3A and B**). The increase of visceral sensitivity was even more pronounced in females than in males as DNBS-treated females showed a significantly increased response also when stimulated with the 0.5 mL volume of distension in both the test performed (**Figs. 3A and B**). The acute i.p. administration of MRS5980 dose-dependently reduced visceral hyperalgesia induced by DNBS in female rats. In particular, the highest doses (1.2 and 2.4 $\mu\text{mol}\cdot\text{kg}^{-1}$) reversed visceral sensitivity alteration back to the value of controls in both VMR and AWR tests (**Figs. 3A and B**), showing similar efficacy to linaclotide (0.2 $\text{nmol}\cdot\text{kg}^{-1}$, **Fig. 3A**). The lowest dose of MRS5980 (0.3 $\mu\text{mol}\cdot\text{kg}^{-1}$) was ineffective in females like in males (**Figs. 3A and B**). To note, tests were performed in female animals synchronized for the estrous cycle choosing the proestrous/estrous phases to eliminate any possible confounding factor in the evaluation of A_3 receptor agonists' activity.

3.4. MRS5980 reduces persistent visceral pain

This type of pain has the peculiar tendency to become chronic in patients remitting from an intestinal damage.^{84,85} Unlike VMR, the




Figure 2. Evaluation of the role of adenosine A_3 ARs and N-type voltage-gated calcium channels in visceral pain relief and comparison with the effect of reference drugs. Tests were performed 14 days after DNBS treatment by measuring the visceromotor response (VMR) to colorectal distension (CRD) after compound administration. (A) Effect of pretreatment with the selective A_3 antagonist MRS1523 ($5 \mu\text{mol}\cdot\text{kg}^{-1}$ i.p.) on the visceral pain-relieving effect of the highly selective A_3 AR agonist, MRS5980 ($2.4 \mu\text{mol}\cdot\text{kg}^{-1}$ i.p.). (B) Effect of the selective A_3 agonist, CI-IB-MECA (0.03 , 0.3 , and $2.4 \mu\text{mol}\cdot\text{kg}^{-1}$; i.p.). (C) Effect of the selective N-type voltage-gated calcium channel ($\text{Ca}_v2.2$) blocker, PD173212 (0.0017 - $1.7 \mu\text{mol}\cdot\text{kg}^{-1}$ i.p.). (D) Effect of linaclotide (0.0066 - $0.2 \text{nmol}\cdot\text{kg}^{-1}$ p.o.). Each value is the mean \pm SEM of 6 rats per group. $\wedge\wedge P < 0.01$ vs vehicle-treated normal controls (blue). $*P < 0.05$ and $**P < 0.01$ vs DNBS + vehicle-treated group (red). $^{\circ}P < 0.01$ vs DNBS + MRS5980 ($2.4 \mu\text{mol}\cdot\text{kg}^{-1}$, black triangles in A) or CI-IB-MECA ($1.2 \mu\text{mol}\cdot\text{kg}^{-1}$, black triangles in B).

assessment of AWR allows the evaluation of DNBS-induced hypersensitivity for a long time permitting the measurement of persistent pain. The behavioural response was still altered 28 and 35 days after the colonic irritation (Figs. 4A and B, respectively). Hence, we used this test to evaluate the effect of MRS5980 on visceral pain in a more delayed phase after the initial insult. On Day 28 and 35, the pain-relieving effect of MRS5980 (0.3 , 1.2 , and $2.4 \mu\text{mol}\cdot\text{kg}^{-1}$) was as potent as that seen earlier. The effect on chronic hypersensitivity was dose-dependent, and the highest dose ($2.4 \mu\text{mol}\cdot\text{kg}^{-1}$) was again able to reverse the pain threshold of DNBS-treated animals to the value of controls (Figs. 4A and B).

3.5. MRS5980 does not impair gastrointestinal transit

To determine whether the A_3 AR agonist MRS5980 influenced gut motility, we assessed total gastrointestinal transit. Groups of 6 rats were treated with either vehicle (5% DMSO) or MRS5980 ($2.4 \mu\text{mol}\cdot\text{kg}^{-1}$). Fifteen min later, animals received an oral gavage of 70 kDa FITC dextran (5mg/mL , 300mL per rat). Samples were harvested 40 min after 70 kDa FITC-dextran administration. Total fluorescence of 70 kDa FITC-dextran was determined for each of the 10 segments (Fig. 5A). 70 kDa FITC was detected in vehicle-treated samples in segments 5 ($8.9 \pm 2.5 \mu\text{g/mL}$), 6 ($3.0 \pm 2.2 \mu\text{g/mL}$), and 8 ($6.3 \pm 3.2 \mu\text{g/mL}$). MRS5980 treatment showed a similar pattern of GI transit of 70 kDa FITC; segment 5: $11.17 \pm 5.0 \mu\text{g/mL}$,




Figure 3. Effect of MRS5980 and linaclotide on visceral pain induced by DNBS in female rats. Effect of MRS5980 (0.3 , 1.2 , and $2.4 \mu\text{mol}\cdot\text{kg}^{-1}$; i.p.) and linaclotide ($0.2 \text{nmol}\cdot\text{kg}^{-1}$ p.o.) on visceromotor response (VMR, A) to CRD on day 14 after DNBS injection. Effect of MRS5980 (0.3 , 1.2 , and $2.4 \mu\text{mol}\cdot\text{kg}^{-1}$; i.p.) on abdominal withdrawal reflex (AWR, B) to CRD on day 14 after DNBS injection. Each value is the mean \pm SEM of 6 rats per group. $\wedge\wedge P < 0.01$ vs vehicle-treated normal controls (violet). $*P < 0.05$ and $**P < 0.01$ vs DNBS + vehicle-treated group (red). CRD, colorectal distension.




Figure 4. Effect of MRS5980 on persistent visceral pain induced by DNBS injection. Effect of MRS5980 (0.3, 1.2, and 2.4 $\mu\text{mol kg}^{-1}$; i.p.) was observed 28 days (A) and 35 days (B) after DNBS treatment. The compound was administered 15 minutes before the first colorectal distension (CRD). Visceral pain was assessed by measuring the animal abdominal withdrawal reflex (AWR) to CRD. Each value is the mean \pm SEM of 6 rats per group. ^^ $P < 0.01$ vs vehicle-treated normal controls (blue). * $P < 0.05$ and ** $P < 0.01$ vs DNBS + vehicle-treated group (red).

segment 6: $11.2 \pm 6.3 \mu\text{g/mL}$, and segment 8: $15.1 \pm 14.3 \mu\text{g/mL}$. Detection of 70 kDa FITC-dextran in all other segments was $<4 \mu\text{g/mL}$ (2.1–3.7 $\mu\text{g/mL}$). No significant difference between treatments ($P = 0.453$) or the interaction of treatment and GI segment ($F(9,92) = P = 0.8676$) was found. FITC levels were significantly different between the GI segments ($P = 0.0010$) indicative of intestinal movement as anticipated. These data indicate that A₃AR agonism with MRS5980 does not statistically reduce GI transit as compared to vehicle.

To determine whether MRS5980 influenced paracellular uptake of 70 kDa from the GI tract, we measured fluorescence in serum collected from the same animals (Fig. 5B). Serum levels of 70 kDa FITC-dextran were minimal (vehicle: 0.82 ± 0.27 ; MRS5980: 1.38 ± 0.57) and not significantly different ($P = 0.3955$). These data suggest that MRS5980 does not influence the paracellular integrity of GI epithelium alone as compared to vehicle.

3.6. A₃R agonists inhibit N-type voltage-gated Ca²⁺ channels in dorsal root ganglia neurons isolated from control or DNBS-treated rats

To gain insight into mechanisms underlying the antihyperalgesic role of A₃ARs, we explored electrophysiological properties and CI-IB-MECA effects in DRG primary sensory neurons isolated from vehicle-treated (control group) or DNBS-treated rats (14

days after treatment). Concerning passive membrane properties (Fig. 6A), DNBS neurons presented a more depolarized V_m in comparison to those from the vehicle-treated animals (Fig. 6A, left panel: from -61.4 ± 2.2 mV in vehicle-treated to -52.0 ± 2.6 mV in DNBS group, $n = 23$ and $n = 33$, respectively, $P = 0.0143$, unpaired Student *t* test), whereas no obvious differences were found in R_m or C_m (Fig. 6A, central and right panels). C_m was, on average, 27.6 ± 4.1 pF in vehicle-treated animals and 28.3 ± 2.7 pF in DNBS-treated animals (Fig. 6A, right panel), corresponding to a cell diameter of 29.9 and 30.7 μm , respectively, thus confirming that present data were collected from small–medium-sized DRG neurons. Of note, VDCCs, activated by a depolarizing voltage step protocol (from -50 to $+50$ mV, 200 ms; see insert in Fig. 6B), showed some differences in DNBS-treated vs vehicle-treated rats. Although if total peak currents were unchanged, either in amplitude (Fig. 6C) or time to peak (Fig. 6D), Ca²⁺ currents measured at the steady state in DNBS neurons were significantly smaller in amplitude (Fig. 6E right panel: from -21.5 ± 8.1 pA/pF in vehicle-treated to -1.4 ± 4.0 pA/pF in DNBS group, $n = 15$ and $n = 14$, respectively, $P = 0.0393$, unpaired Student *t* test).

Consistent with our previous work¹⁹ and with the above in vivo results (Fig. 2), we confirmed that CI-IB-MECA (30 nM) inhibited Ca²⁺ currents in DRG neurons isolated either from vehicle- or DNBS-treated rats (Figs. 7A and B) and the effect was prevented




Figure 5. Gastrointestinal transit is not reduced by acute A₃AR agonism with MRS5980. (A) Amount of 70 kDa FITC-dextran in small intestine segments 15 minutes after MRS5980 or vehicle dosing and 40 minutes after oral gavage of the marker. Two-way ANOVA, interaction: $F(9,92) = 0.8676$; $P = 0.9553$; GI Section: $P = 0.0010$; Treatment: $P = 0.4584$ (B) Amount of 70,000 kDa FITC-dextran in serum samples of 15 minutes after MRS5980 or vehicle dosing and 40 minutes after oral gavage of the marker (unpaired *t* test, $P = 0.3955$ MRS5980 compared to vehicle.) Each value is the mean \pm S.E.M. of 6 rats. ANOVA, analysis of variance.




Figure 6. Electrophysiological properties of DRG neurons isolated from vehicle- or DNBS-treated rats. (A) Pooled data (mean \pm SEM) of passive membrane properties (resting membrane potential: V_m , left panel; membrane resistance: R_m , central panel, and cell capacitance: C_m , right panel) measured in DRG neurons isolated from vehicle- (veh) or DNBS-treated rats (DNBS). Unpaired Student t test, $n = 23$ and $n = 33$, respectively. (B) Original Ca^{2+} current traces elicited in representative DRG neurons isolated from a vehicle- (black traces) or DNBS-treated (red traces) rat by applying a depolarizing voltage step protocol (from -50 to $+50$ mV, 200 ms duration, $V_h = -65$ mV; see inset). Averaged I-V plots of peak Ca^{2+} currents (C), and pooled data of time to peak of Ca^{2+} currents (D), measured at different step potentials in 17 cells isolated from vehicle-treated rats and 15 cells isolated from DNBS-treated animals. No significant differences were found between groups at any step potential, unpaired Student t test. (E) Left panel: averaged I-V plot of steady-state Ca^{2+} currents measured in 17 cells isolated from vehicle-treated rats and 15 cells isolated from DNBS-treated animals. Right panel: pooled data of the same currents measured at the 0 mV step. (A) significant reduction of steady-state Ca^{2+} currents was found in the DNBS-treated group, $P = 0.0393$, unpaired Student t test. DRG, dorsal root ganglia.

by the A_3AR antagonist MRS1523 (100 nM; **Fig. 7B**) and by the N-type Ca^{2+} channel blocker PD173232 (1 μ M; **Figs. 7C–E**). No differences were found in the percentage of CI-IB-MECA-inhibited Ca^{2+} current in neurons isolated from vehicle- or DNBS-treated rats ($20.0 \pm 4.1\%$ and $22.5 \pm 6.1\%$ of I_{Ca} inhibition in CI-IB-MECA, respectively, $n = 7$ in both groups; $P = 0.7332$ unpaired Student t test; **Fig. 7C**) nor in the percentage of PD173232-blocked currents ($72.3 \pm 4.8\%$ and $77.7 \pm 8.2\%$, of I_{Ca} inhibition in PD173232, respectively; $n = 4$ in both groups; $P = 0.5867$ unpaired Student t test, data not shown). Similar results to those obtained with the prototypical A_3AR agonist CI-IB-MECA were recorded with MRS5980 (30 nM): the compound inhibited, to a similar extent, Ca^{2+} currents either in neurons isolated from vehicle- or DNBS-treated rats and the effect was prevented by MRS1523 and by the N-type channel blocker PD173232 (1 μ M; **Figs. 7C–E**).

4. Discussion

In the present work, we studied the role of A_3AR as a possible new pharmacological target for relieving visceral pain. The novel highly

selective second-generation A_3AR agonist, MRS5980, as well as the first-generation A_3AR agonist, CI-IB-MECA, were effective against postinflammatory visceral hypersensitivity in rats. The intracolonic injection of DNBS in rodents induces a long-lasting visceral hypersensitivity,^{1,32,33} which can be measured with high reproducibility as a lowered sensory threshold to CRD.^{52,61,82} DNBS-induced visceral hypersensitivity persists after the resolution of the acute inflammatory phase,^{1,40,53} making this model suitable to investigate visceral pain related not only related to IBD but also to IBS, a chronic disease characterized by a marked abdominal pain in the absence of histopathological explanations.^{84,90}

The administration of A_3AR agonists in the model of post-inflammatory visceral pain showed an efficacy equivalent to that of linaclotide, an approved treatment for reducing abdominal pain (and also abdominal bloating and bowel symptoms) in adult patients suffering from IBS with predominant constipation (IBS-C).^{13,18}

The efficacy of MRS5980 and linaclotide was confirmed in female rats. These data have a clinical significance because IBS affects more women than men.¹² Abundant evidence from




Figure 7. Effects of the adenosine A_3R agonists CI-IB-MECA and MRS5980 on voltage-gated Ca^{2+} currents in isolated DRG neurons. (A) Time course (left panel) and representative Ca^{2+} current traces (right panel) measured in a representative DRG neuron isolated from a DNBS-treated rat (DNBS) before and after the application of the A_3R agonist CI-IB-MECA (30 nM) or the Ca^{2+} channel blocker Cd^{2+} (100 μ M). (B) Averaged time courses of peak Ca^{2+} currents during CI-IB-MECA (30 nM) superfusion in vehicle-treated or DNBS-treated animals in the absence or presence of the A_3R antagonist, MRS1523 (1523; 100 nM). Pooled data (mean \pm SEM) of % Ca^{2+} current inhibition induced by CI-IB-MECA (C) or MRS5980 (D) applied alone or in combination with MRS1523 or PD173232 (PD; 1 μ M) in vehicle-treated (white columns) or DNBS-treated (red columns) animals. In all panels: * $P < 0.05$ vs respective control, paired Student t test. (E) Averaged time course of peak Ca^{2+} currents before or after MRS5980 (30 nM) superfusion DNBS rats in the presence of the A_3R antagonist MRS1523 or the Ca^{2+} channel blocker PD173232 (1 μ M). DRG, dorsal root ganglia.

epidemiologic studies clearly demonstrates that women are at substantially greater risk for many clinical pain conditions.⁵ In accordance, the present results showed a higher visceral hypersensitivity in female animals than in males. This phenomenon is not limited to abdominal/pelvic pain (such as IBS, painful bladder syndrome, and dyspareunia), but also includes conditions associated with neuropathic pain, musculoskeletal pain, orofacial pain, and headache/migraine.²⁷ Women are therefore more likely than men to pursue a variety of treatments for many painful conditions.

The trinitrobenzenesulfonic acid (TNBS)-induced model was used in preclinical studies to identify the potential clinical utility of linaclotide.²⁵ Our results with linaclotide in the DNBS-induced model support previous findings. The analgesic mechanism of linaclotide involves the activation of guanylate cyclase-C (GC-C) expressed on mucosal epithelial cells, resulting in the production and release of cyclic guanosine-3',5'-monophosphate (cGMP). The extracellular cGMP acts on and inhibits nociceptors, thereby reducing nociception.¹³ Although linaclotide is an efficacious, well-tolerated treatment option for improving both bowel symptoms and abdominal pain in IBS-C, it acts as a secretagogues and this implies that most patients reported episodes of diarrhea within the first 2 to 4 weeks of treatment.¹⁸ This adverse effect also makes linaclotide unsuitable for patients affected by diarrhoea-predominant IBS or IBS with alternating constipation and diarrhoea. In healthy condition, adenosine modulated colonic cholinergic motility through activation of A_3 receptors in the myenteric plexus. Antonioli et al. demonstrated that A_3 receptor-mediated tonic inhibitory control by adenosine was impaired in inflamed bowel, despite the increased density of functioning and pharmacologically recruitable A_3 receptors found in the gut.² To verify a possible interaction of MRS5980 with gut motility, we

performed focused experiments. Our findings using 70 kDa FITC-dextran revealed that MRS5980 does not impair gastrointestinal transit as compared to vehicle. Importantly, these results indicate that the antihyperalgesic efficacy of MRS5980 in visceral pain condition is not associated to motility alteration, suggesting a mechanism related to sensitivity modulation rather than spasmolytic activity. This profile is different, eg, from the opioid analgesic drugs, mainly used in patients affected by IBS-D, that in the same experiment reduce intestinal transit leading to constipation in humans.^{10,47,66,76} Anyway, the impact of A_3 receptor agonists' repeated administrations on intestinal motility and sensitivity is an aspect that needs to be explored further in the perspective of developing these molecules as drugs and of establishing a therapeutic regime in patients. The lack of significant long-term effects on intestinal motility could privilege A_3 adenosine agonists over all the other drugs currently in use. Anyway, in clinical trials investigating the efficacy of A_3 receptor agonist on rheumatoid arthritis and psoriasis, no relevant gastrointestinal side effects were reported.^{20,41,80} The efficacy of A_3R agonists against visceral pain is consistent with data showing that the A_3R is involved with multiple pain mechanisms at peripheral, spinal, and supraspinal levels.^{43,50} There is evidence that A_3R agonists are antihyperalgesic against neuropathic pain through inhibition of the astrocyte-associated neuroinflammatory response in the spinal cord,^{42,44,96} a phenomenon strongly involved in pain persistence.^{22,23} A_3R activation has been reported to enhance the formation of anti-inflammatory cytokines^{36,42,44,96} and the production of glial-derived neuroprotective substances.⁹⁹ Furthermore, in vitro and in vivo studies demonstrate that A_3R produces its effects by inhibiting the p38 MAPK and NF- κ B signaling pathways^{42,55,92,94} and inflammatory activity.⁹⁶ All these mechanisms may contribute to the relief

of colitis-evoked visceral pain consistently with the reports that A₃AR activation reduced colitis-induced tissue injury by modulating the NF- κ B signalling pathway and through inhibiting NLRP3 inflammasome activation and pyroptosis in human colonic epithelial cells.^{70,71}

A₃ARs may limit excitatory neurotransmission, which is altered in chronic³⁴ as well as in visceral pain.^{31,48,98} For example, the protective role of A₃AR in the first phase of ischemia seems to be at least partly related to a decrease in synaptic transmission.^{67,68,72} Moreover, A₃AR activation protects against the neurotoxic intracellular Ca²⁺ rise mediated by P2X7 or NMDA receptors.^{103,104} All the mechanisms described above could account for a therapeutic effect of repeated administrations of A₃ receptor agonists, which is another aspect interesting to study. Anyway, the acute efficacy showed by A₃ receptor agonists in reducing visceral hypersensitivity can be only partially explained by the mechanisms of action previously cited. This consideration has aroused our interest in elucidating further the pharmacodynamic mechanisms of A₃ receptor agonists on pain.

As recently demonstrated,¹⁹ selective A₃AR stimulation inhibits N-type Ca_v2.2 opening in isolated rat DRG neurons. Here, we found that both CI-IB-MECA and MRS5980 significantly inhibited Ca_v2.2 activation in DRG neurons of DNBS-treated rats. The effect was prevented by the selective A₃AR antagonist, MRS1523, and by the Ca_v2.2 blocker, PD173212, confirming the involvement of Ca_v2.2 in the A₃AR-mediated effect. We did not measure a significant difference between A₃AR-mediated inhibition in DRG neurons isolated from DNBS-treated vs vehicle-treated animals. However, alterations in passive membrane properties and steady-state Ca²⁺ currents were observed between the 2 groups. The resting membrane potential was significantly more depolarized in DRG neurons isolated from DNBS-treated animals, suggesting a hyperexcitable state. Interestingly, steady-state Ca²⁺ currents, but not peak Ca²⁺ currents, were markedly reduced in DRG neurons isolated from DNBS animals. In an attempt to explain the relationship between steady-state Ca²⁺ current reduction in DRG neurons and visceral hypersensitivity, we hypothesize that a decrease in sustained Ca²⁺ influx into the cell can lead to membrane potential instability and depolarization through a decrease of Ca²⁺-activated K⁺ channel (K_{Ca}) opening. Ca²⁺-activated K⁺ channel is known to stabilize the membrane potential and participate in repolarizing neurons after action potential firing, thus avoiding bursting activity.^{37,63,102}

Moreover, a recent publication demonstrates that Ca_v2.2 induces a voltage-dependent, but Ca²⁺-independent ATP secretion from the soma of DRG neurons.¹⁵ ATP is a powerful mediator of pain, for example, ATP-induced P2X3 receptor activation is involved in visceral pain, as shown by the ability of P2X3 antagonist A-317491 to potently reduce hypersensitivity.²¹

The hypothesis that Ca_v2.2-mediated modulation contributes to A₃AR-mediated antihyperalgesia is supported by other evidence. Modulation of Ca_v2.2 channel activity is implied in the pharmacodynamics of pain-relieving compounds, such as other Gi-coupled receptor agonists, eg, opioids, cannabinoids, neuropeptide Y, and substance P.^{7,78,81,87,91} Also, the neuropathic pain analgesics gabapentin and pregabalin²⁸ inhibit Ca_v2.2-mediated synaptic transmission.^{6,88} Gabapentinoids are effective in reducing visceral pain and preventing spinal neuronal activation associated with CRD in animals^{62,69,79} and have been suggested to be treatments for IBS.¹¹ A direct inhibitor of Ca_v2.2 channel activity, ziconotide, is currently used clinically for pain therapy, although it is limited to the intrathecal route.⁵⁶ In our hands, the i.p. administration of the specific Ca_v2.2 inhibitor

PD173212^{38,56,73} significantly decreased visceral hypersensitivity in DNBS-treated animals. In conclusion, A₃AR agonists seem to be a promising resource for visceral pain management, an unmet medical need that impairs quality of life.

Conflict of interest statement

K.A. Jacobson thanks the NIDDK Intramural Research Program (ZIA DK031117). D. Salvemini is a cofounder of BioIntervene, Inc, which licensed related intellectual property from Saint Louis University. The remaining authors have no conflicts of interest to declare.

Acknowledgments

This research was supported by the Italian Ministry of Instruction, University and Research and by the University of Florence. The authors are grateful to Dr G.J. Bennett (Department of Anesthesiology, University of California, San Diego, La Jolla, CA) for his input in our work and for editorial contribution to the manuscript.

Appendix A. Supplemental digital content

Supplemental digital content associated with this article can be found online at <http://links.lww.com/PAIN/B10>.

Article history:

Received 31 January 2020

Received in revised form 27 March 2020

Accepted 14 April 2020

Available online 4 May 2020

References

- [1] Adam B, Liebrechts T, Gschossmann JM, Krippner C, Scholl F, Ruwe M, Holtmann G. Severity of mucosal inflammation as a predictor for alterations of visceral sensory function in a rat model. *PAIN* 2006;123: 179–86.
- [2] Antonioli L, Fornai M, Colucci R, Ghisu N, Tuccori M, Awwad O, Bin A, Zoppellaro C, Castagliuolo I, Gaion RM, Giron MC, Blandizzi C. Control of enteric neuromuscular functions by purinergic A(3) receptors in normal rat distal colon and experimental bowel inflammation. *Br J Pharmacol* 2010;161:856–71.
- [3] Asano T, Takenaga M. Adenosine A2B receptors: an optional target for the management of irritable bowel syndrome with diarrhea? *J Clin Med* 2017;6:E104.
- [4] Asano T, Tanaka KI, Tada A, Shimamura H, Tanaka R, Maruoka H, Takenaga M, Mizushima T. Aminophylline suppresses stress-induced visceral hypersensitivity and defecation in irritable bowel syndrome. *Sci Rep* 2017;7:40214.
- [5] Bartley EJ, Fillingim RB. Sex differences in pain: a brief review of clinical and experimental findings. *Br J Anaesth* 2013;111:52–8.
- [6] Bauer CS, Nieto-Rostro M, Rahman W, Tran-Van-Minh A, Ferron L, Douglas L, Kadurin I, Sri Ranjan Y, Fernandez-Alacid L, Millar NS, Dickenson AH, Lujan R, Dolphin AC. The increased trafficking of the calcium channel subunit alpha2delta-1 to presynaptic terminals in neuropathic pain is inhibited by the alpha2delta ligand pregabalin. *J Neurosci* 2009;29:4076–88.
- [7] Beedle AM, Zamponi GW. Modulation of high voltage-activated calcium channels by G protein-coupled receptors. In: McDonough SI, editor. *Calcium channel pharmacology*. Boston: Springer US, 2004. pp. 331–67.
- [8] Borea PA, Gessi S, Merighi S, Vincenzi F, Varani K. Pharmacology of adenosine receptors: the state of the art. *Physiol Rev* 2018;98: 1591–625.
- [9] Brierley SM, Linden DR. Neuroplasticity and dysfunction after gastrointestinal inflammation. *Nat Rev Gastroenterol Hepatol* 2014;11: 611–27.

- [10] Burks TF, Fox DA, Hirning LD, Shook JE, Porreca F. Regulation of gastrointestinal function by multiple opioid receptors. *Life Sci* 1988;43:2177–81.
- [11] Camilleri M, Boeckstaens G. Dietary and pharmacological treatment of abdominal pain in IBS. *Gut* 2017;66:966–74.
- [12] Canavan C, West J, Card T. The epidemiology of irritable bowel syndrome. *Clin Epidemiol* 2014;6:71–80.
- [13] Castro J, Harrington AM, Hughes PA, Martin CM, Ge P, Shea CM, Jin H, Jacobson S, Hannig G, Mann E, Cohen MB, MacDougall JE, Lavins BJ, Kurtz CB, Silos-Santiago I, Johnston JM, Currie MG, Blackshaw LA, Brierley SM. Linaclotide inhibits colonic nociceptors and relieves abdominal pain via guanylate cyclase-C and extracellular cyclic guanosine 3',5'-monophosphate. *Gastroenterology* 2013;145:1334–46.e1331–1311.
- [14] Cervero F, Laird JM. Spinal mechanisms of visceral pain and hyperalgesia. In: *Synaptic plasticity in pain*. New York: Springer, 2009. pp. 289–306.
- [15] Chai Z, Wang C, Huang R, Wang Y, Zhang X, Wu Q, Wang Y, Wu X, Zheng L, Zhang C, Guo W, Xiong W, Ding J, Zhu F, Zhou Z. CaV2.2 gates calcium-independent but voltage-dependent secretion in mammalian sensory neurons. *Neuron* 2017;96:1317–26.e1314.
- [16] Chen Y, Lin C, Tang Y, Chen AQ, Liu CY, Lu DL. ZD 7288, an HCN channel blocker, attenuates chronic visceral pain in irritable bowel syndrome-like rats. *World J Gastroenterol* 2014;20:2091–7.
- [17] Chen Z, Janes K, Chen C, Doyle T, Bryant L, Tosh DK, Jacobson KA, Salvemini D. Controlling murine and rat chronic pain through A3 adenosine receptor activation. *FASEB J* 2012;26:1855–65.
- [18] Chey WD, Lembo AJ, Lavins BJ, Shiff SJ, Kurtz CB, Currie MG, MacDougall JE, Jia XD, Shao JZ, Fitch DA, Baird MJ, Schneier HA, Johnston JM. Linaclotide for irritable bowel syndrome with constipation: a 26-week, randomized, double-blind, placebo-controlled trial to evaluate efficacy and safety. *Am J Gastroenterol* 2012;107:1702–12.
- [19] Coppi E, Cherchi F, Fusco I, Failli P, Vona A, Dettori I, Gaviano L, Lucarini E, Jacobson KA, Tosh DK, Salvemini D, Ghelardini C, Pedata F, Di Cesare Mannelli L, Pugliese AM. Adenosine A3 receptor activation inhibits pronociceptive N-type Ca²⁺ currents and cell excitability in dorsal root ganglion neurons. *PAIN* 2019;160:1103–18.
- [20] David M, Gospodinov DK, Gheorghe N, Mateev GS, Rusinova MV, Hristakieva E, Solovastru LG, Patel RV, Giurcaneanu C, Hitova MC, Purcaru AI, Horia B, Tsingov II, Yankova RK, Kadurina MI, Ramon M, Rotaru M, Simionescu O, Benea V, Demerdjieva ZV, Cosgarea MR, Morariu HS, Michael Z, Cristodor P, Nica C, Silverman MH, Bristol DR, Harpaz Z, Farbstein M, Cohen S, Fishman P. Treatment of plaque-type psoriasis with oral CF101: data from a phase II/III multicenter, randomized, controlled trial. *J Drugs Dermatol* 2016;15:931–8.
- [21] Deiteren A, van der Linden L, de Wit A, Ceuleers H, Bückinx R, Timmermans JP, Moreels TG, Pelckmans PA, De Man JG, De Winter BY. P2X3 receptors mediate visceral hypersensitivity during acute chemically-induced colitis and in the post-inflammatory phase via different mechanisms of sensitization. *PLoS One* 2015;10:e0123810.
- [22] Di Cesare Mannelli L, Pacini A, Bonaccini L, Zanardelli M, Mello T, Ghelardini C. Morphologic features and glial activation in rat oxaliplatin-dependent neuropathic pain. *J Pain* 2013;14:1585–600.
- [23] Di Cesare Mannelli L, Pacini A, Micheli L, Tani A, Zanardelli M, Ghelardini C. Glial role in oxaliplatin-induced neuropathic pain. *Exp Neurol* 2014;261:22–33.
- [24] Enck P, Aziz Q, Barbara G, Farmer AD, Fukudo S, Mayer EA, Niesler B, Quigley EM, Rajilic-Stojanovic M, Schemann M, Schwille-Kiuntke J, Simren M, Zipfel S, Spiller RC. Irritable bowel syndrome. *Nat Rev Dis Primers* 2016;2:16014.
- [25] Eutamene H, Bradesi S, Larauche M, Theodorou V, Beaufrand C, Ohning G, Fioramonti J, Cohen M, Bryant AP, Kurtz C, Currie MG, Mayer EA, Bueno L. Guanylate cyclase C-mediated antinociceptive effects of linaclotide in rodent models of visceral pain. *Neurogastroenterol Motil* 2010;22:312–e84.
- [26] Farrokhyar F, Marshall JK, Easterbrook B, Irvine EJ. Functional gastrointestinal disorders and mood disorders in patients with inactive inflammatory bowel disease: prevalence and impact on health. *Inflamm Bowel Dis* 2006;12:38–46.
- [27] Fillingim RB, King CD, Ribeiro-Dasilva MC, Rahim-Williams B, Riley JL III. Sex, gender, and pain: a review of recent clinical and experimental findings. *J Pain* 2009;10:447–85.
- [28] Finnerup NB, Attal N, Haroutunian S, McNicol E, Baron R, Dworkin RH, Gilron I, Haanpaa M, Hansson P, Jensen TS, Kamerman PR, Lund K, Moore A, Raja SN, Rice AS, Rowbotham M, Sena E, Siddall P, Smith BH, Wallace M. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. *Lancet Neurol* 2015;14:162–73.
- [29] Fishman P, Bar-Yehuda S, Liang BT, Jacobson KA. Pharmacological and therapeutic effects of A3 adenosine receptor agonists. *Drug Discov Today* 2012;17:359–66.
- [30] Fornai M, Blandizzi C, Antonoli L, Colucci R, Bernardini N, Segnani C, De Ponti F, Del Tacca M. Differential role of cyclooxygenase 1 and 2 isoforms in the modulation of colonic neuromuscular function in experimental inflammation. *J Pharmacol Exp Ther* 2006;317:938–45.
- [31] Gebhart GF, Bielefeldt K. Physiology of visceral pain. *Compr Physiol* 2016;6:1609–33.
- [32] Greenwood-Van Meerveld B, Prusator DK, Johnson AC. Animal models of gastrointestinal and liver diseases. Animal models of visceral pain: pathophysiology, translational relevance, and challenges. *Am J Physiol Gastrointest Liver Physiol* 2015;308:G885–903.
- [33] Gschostmann JM, Liebrechts T, Adam B, Buenger L, Ruwe M, Gerken G, Holtmann G. Long-term effects of transient chemically induced colitis on the visceromotor response to mechanical colorectal distension. *Dig Dis Sci* 2004;49:96–101.
- [34] Gwak YS, Kang J, Unabia GC, Hulsebosch CE. Spatial and temporal activation of spinal glial cells: role of gliopathy in central neuropathic pain following spinal cord injury in rats. *Exp Neurol* 2012;234:362–72.
- [35] Halpin SJ, Ford AC. Prevalence of symptoms meeting criteria for irritable bowel syndrome in inflammatory bowel disease: systematic review and meta-analysis. *Am J Gastroenterol* 2012;107:1474–82.
- [36] Hasko G, Szabo C, Nemeth ZH, Kvetan V, Pastores SM, Vizi ES. Adenosine receptor agonists differentially regulate IL-10, TNF-alpha, and nitric oxide production in RAW 264.7 macrophages and in endotoxemic mice. *J Immunol* 1996;157:4634–40.
- [37] Hogan QH. Role of decreased sensory neuron membrane calcium currents in the genesis of neuropathic pain. *Croat Med J* 2007;48:9–21.
- [38] Hu LY, Ryder TR, Rafferty MF, Dooley DJ, Geer JJ, Lotarski SM, Miljanich GP, Millerman E, Rock DM, Stoehr SJ, Szoke BG, Taylor CP, Vartanian MG. Structure-activity relationship of N-methyl-N-aryl-peptidylamines as novel N-type calcium channel blockers. *Bioorg Med Chem Lett* 1999;9:2151–6.
- [39] Hubscher CH, Brooks DL, Johnson JR. A quantitative method for assessing stages of the rat estrous cycle. *Biotech Histochem* 2005;80:79–87.
- [40] Hughes PA, Brierley SM, Martin CM, Brookes SJ, Linden DR, Blackshaw LA. Post-inflammatory colonic afferent sensitization: different subtypes, different pathways and different time courses. *Gut* 2009;58:1333–41.
- [41] Jacobson KA, Merighi S, Varani K, Borea PA, Baraldi S, Aghazadeh Tabrizi M, Romagnoli R, Baraldi PG, Ciancetta A, Tosh DK, Gao ZG, Gessi S. A3 adenosine receptors as modulators of inflammation: from medicinal chemistry to therapy. *Med Res Rev* 2018;38:1031–72.
- [42] Janes K, Esposito E, Doyle T, Cuzzocrea S, Tosh DK, Jacobson KA, Salvemini D. A3 adenosine receptor agonist prevents the development of paclitaxel-induced neuropathic pain by modulating spinal glial-restricted redox-dependent signaling pathways. *PAIN* 2014;155:2560–7.
- [43] Janes K, Symons-Liguori AM, Jacobson KA, Salvemini D. Identification of A3 adenosine receptor agonists as novel non-narcotic analgesics. *Br J Pharmacol* 2016;173:1253–67.
- [44] Janes K, Wahlman C, Little JW, Doyle T, Tosh DK, Jacobson KA, Salvemini D. Spinal neuroimmune activation is independent of T-cell infiltration and attenuated by A3 adenosine receptor agonists in a model of oxaliplatin-induced peripheral neuropathy. *Brain Behav Immun* 2015;44:91–9.
- [45] Ji Y, Murphy AZ, Traub RJ. Estrogen modulates the visceromotor reflex and responses of spinal dorsal horn neurons to colorectal stimulation in the rat. *J Neurosci* 2003;23:3908.
- [46] Ji Y, Tang B, Traub RJ. The visceromotor response to colorectal distention fluctuates with the estrous cycle in rats. *Neuroscience* 2008;154:1562–7.
- [47] Jiang Q, Sheldon RJ, Porreca F. A comparison of the central gastrointestinal antitransit effects of morphine and bombesin in the mouse. *Life Sci* 1987;41:2455–61.
- [48] Lapointe TK, Basso L, Iftinca MC, Flynn R, Chapman K, Dietrich G, Vergnolle N, Altier C. TRPV1 sensitization mediates postinflammatory visceral pain following acute colitis. *Am J Physiol Gastrointest Liver Physiol* 2015;309:G87–99.
- [49] Li AH, Moro S, Melman N, Ji XD, Jacobson KA. Structure-activity relationships and molecular modeling of 3, 5-diacyl-2,4-dialkylpyridine derivatives as selective A3 adenosine receptor antagonists. *J Med Chem* 1998;41:3186–201.
- [50] Little JW, Ford A, Symons-Liguori AM, Chen Z, Janes K, Doyle T, Xie J, Luongo L, Tosh DK, Maione S, Bannister K, Dickenson AH, Vanderah TW, Porreca F, Jacobson KA, Salvemini D. Endogenous adenosine A3

- receptor activation selectively alleviates persistent pain states. *Brain* 2015;138:28–35.
- [51] Longstreth GF, Thompson WG, Chey WD, Houghton LA, Mearin F, Spiller RC. Functional bowel disorders. *Gastroenterology* 2006;130:1480–91.
- [52] Lowen MB, Mayer E, Tillisch K, Labus J, Naliboff B, Lundberg P, Thorell LH, Strom M, Engstrom M, Walter S. Deficient habituation to repeated rectal distensions in irritable bowel syndrome patients with visceral hypersensitivity. *Neurogastroenterol Motil* 2015;27:646–55.
- [53] Lucarini E, Di Cesare Mannelli L, Micheli L, Trallori E, Antonioli L, Fornai M, Blandizzi C, Ghelardini C. P060 Post-inflammatory visceral pain induced by DNBS: preclinical features for novel therapeutics. *J Crohn's Colitis* 2018;12(suppl_1):S123.
- [54] Luongo L, Petrelli R, Gatta L, Giordano C, Guida F, Vita P, Franchetti P, Grifantini M, de Novellis V, Cappellacci L, Maione S. 5'-Chloro-5'-deoxy-(+/-)-ENBA, a potent and selective adenosine A(1) receptor agonist, alleviates neuropathic pain in mice through functional glial and microglial changes without affecting motor or cardiovascular functions. *Molecules* 2012;17:13712–26.
- [55] Madi L, Cohen S, Ochayin A, Bar-Yehuda S, Barer F, Fishman P. Overexpression of A3 adenosine receptor in peripheral blood mononuclear cells in rheumatoid arthritis: involvement of nuclear factor-kappaB in mediating receptor level. *J Rheumatol* 2007;34:20–6.
- [56] McDowell GC II, Pope JE. Intrathecal ziconotide: dosing and administration strategies in patients with refractory chronic pain. *Neuromodulation* 2016;19:522–32.
- [57] McGrath JC, Lilley E. Implementing guidelines on reporting research using animals (ARRIVE etc.): new requirements for publication in *BJP*. *Br J Pharmacol* 2015;172:3189–93.
- [58] Minderhoud IM, Oldenburg B, Wismeijer JA, van Berge Henegouwen GP, Smout AJ. IBS-like symptoms in patients with inflammatory bowel disease in remission; relationships with quality of life and coping behavior. *Dig Dis Sci* 2004;49:469–74.
- [59] Moloney RD, Sajjad J, Foley T, Felice VD, Dinan TG, Cryan JF, O'Mahony SM. Estrous cycle influences excitatory amino acid transport and visceral pain sensitivity in the rat: effects of early-life stress. *Biol Sex Differ* 2016;7:33.
- [60] Ness TJ. Cardiovascular and visceromotor responses to noxious visceral stimuli vary with the estrous cycle in rats. *Reg Anesth Pain Med* 1999;24(suppl 1):26.
- [61] Ness TJ, Gebhart GF. Colorectal distension as a noxious visceral stimulus: physiologic and pharmacologic characterization of pseudodiffuse reflexes in the rat. *Brain Res* 1988;450:153–69.
- [62] Ohashi K, Kawai M, Ninomiya N, Taylor C, Kurebayashi Y. Effect of a new alpha 2 delta ligand PD-217014 on visceral hypersensitivity induced by 2,4,6-trinitrobenzene sulfonic acid in rats. *Pharmacology* 2008;81:144–50.
- [63] Pagadala P, Park CK, Bang S, Xu ZZ, Xie RG, Liu T, Han BX, Tracey WD, Wang F, Ji RR. Loss of NR1 subunit of NMDARs in primary sensory neurons leads to hyperexcitability and pain hypersensitivity: involvement of Ca2+-activated small conductance potassium channels. *J Neurosci* 2013;33:13425.
- [64] Parisio C, Lucarini E, Micheli L, Toti A, Di Cesare Mannelli L, Antonini G, Panizzi E, Maidecchi A, Giovagnoni E, Lucci J, Ghelardini C. Researching new therapeutic approaches for abdominal visceral pain treatment: preclinical effects of an assembled system of molecules of vegetal origin. *Nutrients* 2019;12:E22.
- [65] Petrelli R, Scortichini M, Belardo C, Boccella S, Luongo L, Capone F, Kachler S, Vita P, Del Bello F, Maione S, Lavecchia A, Klotz KN, Cappellacci L. Structure-based design, synthesis, and in vivo antinociceptive effects of selective A1 adenosine receptor agonists. *J Med Chem* 2018;61:305–18.
- [66] Porreca F, Burks TF. The spinal cord as a site of opioid effects on gastrointestinal transit in the mouse. *J Pharmacol Exp Ther* 1983;227:22–7.
- [67] Pugliese AM, Coppi E, Spalluto G, Corradetti R, Pedata F. A3 adenosine receptor antagonists delay irreversible synaptic failure caused by oxygen and glucose deprivation in the rat CA1 hippocampus in vitro. *Br J Pharmacol* 2006;147:524–32.
- [68] Pugliese AM, Latini S, Corradetti R, Pedata F. Brief, repeated, oxygen-glucose deprivation episodes protect neurotransmission from a longer ischemic episode in the in vitro hippocampus: role of adenosine receptors. *Br J Pharmacol* 2003;140:305–14.
- [69] Ravnefjord A, Brusberg M, Larsson H, Lindstrom E, Martinez V. Effects of pregabalin on visceral pain responses and colonic compliance in rats. *Br J Pharmacol* 2008;155:407–16.
- [70] Ren T, Tian T, Feng X, Ye S, Wang H, Wu W, Qiu Y, Yu C, He Y, Zeng J, Cen J, Zhou Y. An adenosine A3 receptor agonist inhibits DSS-induced colitis in mice through modulation of the NF-kappaB signaling pathway. *Sci Rep* 2015;5:9047.
- [71] Ren T, Zhou Y, Wu W. Su1831—activation of adenosine A3 receptor inhibits Nlrp3 inflammasome and pyroptosis of colonic epithelial cells of patients with ulcerative colitis. *Gastroenterology* 2019;156:S-627.
- [72] Rivera-Oliver M, Diaz-Rios M. Using caffeine and other adenosine receptor antagonists and agonists as therapeutic tools against neurodegenerative diseases: a review. *Life Sci* 2014;101:1–9.
- [73] Ryder TR, Hu LY, Rafferty MF, Millerman E, Szoke BG, Tarczy-Hornoch K. Multiple parallel synthesis of N,N-dialkylidipeptidylamines as N-type calcium channel blockers. *Bioorg Med Chem Lett* 1999;9:1813–18.
- [74] Sajjad J, Felice VD, Golubeva AV, Cryan JF, O'Mahony SM. Sex-dependent activity of the spinal excitatory amino acid transporter: role of estrous cycle. *Neuroscience* 2016;333:311–19.
- [75] Salvemini D, Jacobson KA. Highly selective A3 adenosine receptor agonists relieve chronic neuropathic pain. *Expert Opin Ther Pat* 2017;27:967.
- [76] Santos FA, Rao VS. Quinine-induced inhibition of gastrointestinal transit in mice: possible involvement of endogenous opioids. *Eur J Pharmacol* 1999;364:193–7.
- [77] Sawynok J. Adenosine receptor targets for pain. *Neuroscience* 2016;338:1–18.
- [78] Shapiro MS, Hille B. Substance P and somatostatin inhibit calcium channels in rat sympathetic neurons via different G protein pathways. *Neuron* 1993;10:11–20.
- [79] Sikandar S, Dickenson AH. Visceral pain: the ins and outs, the ups and downs. *Curr Opin Support Palliat Care* 2012;6:17–26.
- [80] Silverman MH, Strand V, Markovits D, Nahir M, Reitblat T, Molad Y, Rosner I, Rozenbaum M, Mader R, Adawi M, Caspi D, Tishler M, Langevit P, Rubinow A, Friedman J, Green L, Tanay A, Ochaion A, Cohen S, Kerns WD, Cohn I, Fishman-Furman S, Farbstein M, Yehuda SB, Fishman P. Clinical evidence for utilization of the A3 adenosine receptor as a target to treat rheumatoid arthritis: data from a phase II clinical trial. *J Rheumatol* 2008;35:41–8.
- [81] Soldo BL, Moises HC. mu-opioid receptor activation inhibits N- and P-type Ca2+ channel currents in magnocellular neurones of the rat supraoptic nucleus. *J Physiol* 1998;513:787–804.
- [82] Spaziani R, Bayati A, Redmond K, Bajaj H, Mazzadi S, Bienenstock J, Collins SM, Kamath MV. Vagal dysfunction in irritable bowel syndrome assessed by rectal distension and baroreceptor sensitivity. *Neurogastroenterol Motil* 2008;20:336–42.
- [83] Spiller R, Garsed K. Infection, inflammation, and the irritable bowel syndrome. *Dig Liver Dis* 2009;41:844–9.
- [84] Spiller R, Major G. IBS and IBD—separate entities or on a spectrum? *Nat Rev Gastroenterol Hepatol* 2016;13:613–21.
- [85] Srinath A, Young E, Szigethy E. Pain management in patients with inflammatory bowel disease: translational approaches from bench to bedside. *Inflamm Bowel Dis* 2014;20:2433–49.
- [86] Srinath AI, Walter C, Newara MC, Szigethy EM. Pain management in patients with inflammatory bowel disease: insights for the clinician. *Therap Adv Gastroenterol* 2012;5:339–57.
- [87] Sun L, Miller RJ. Multiple neuropeptide Y receptors regulate K+ and Ca2+ channels in acutely isolated neurons from the rat arcuate nucleus. *J Neurophysiol* 1999;81:1391–403.
- [88] Sutton KG, Martin DJ, Pinnock RD, Lee K, Scott RH. Gabapentin inhibits high-threshold calcium channel currents in cultured rat dorsal root ganglion neurones. *Br J Pharmacol* 2002;135:257–65.
- [89] Tosh DK, Finley A, Paoletta S, Moss SM, Gao ZG, Gizewski ET, Auchampach JA, Salvemini D, Jacobson KA. In vivo phenotypic screening for treating chronic neuropathic pain: modification of C2-arylethynyl group of conformationally constrained A3 adenosine receptor agonists. *J Med Chem* 2014;57:9901–14.
- [90] Tsang SW, Auyeung KKW, Bian ZX, Ko JKS. Pathogenesis, experimental models and contemporary pharmacotherapy of irritable bowel syndrome: story about the brain-gut Axis. *Curr Neuropharmacol* 2016;14:842–56.
- [91] Twitchell W, Brown S, Mackie K. Cannabinoids inhibit N- and P/Q-type calcium channels in cultured rat hippocampal neurons. *J Neurophysiol* 1997;78:43–50.
- [92] Varani K, Maniero S, Vincenzi F, Targa M, Stefanelli A, Maniscalco P, Martini F, Tognon M, Borea PA. A(3) receptors are overexpressed in pleura from patients with mesothelioma and reduce cell growth via Akt/nuclear factor-kappaB pathway. *Am J Respir Crit Care Med* 2011;183:522–30.
- [93] Varani K, Vincenzi F, Targa M, Paradiso B, Parrilli A, Fini M, Lanza G, Borea PA. The stimulation of A(3) adenosine receptors reduces bone-residing breast cancer in a rat preclinical model. *Eur J Cancer* 2013;49:482–91.

- [94] Varani K, Vincenzi F, Tosi A, Targa M, Masieri FF, Ongaro A, De Mattei M, Massari L, Borea PA. Expression and functional role of adenosine receptors in regulating inflammatory responses in human synoviocytes. *Br J Pharmacol* 2010;160:101–15.
- [95] Vilz TO, Overhaus M, Stoffels B, Websky M, Kalff JC, Wehner S. Functional assessment of intestinal motility and gut wall inflammation in rodents: analyses in a standardized model of intestinal manipulation. *J Vis Exp* 2012;67:4086.
- [96] Wahlman C, Doyle TM, Little JW, Luongo L, Janes K, Chen Z, Esposito E, Tosh DK, Cuzzocrea S, Jacobson KA, Salvemini D. Chemotherapy-induced pain is promoted by enhanced spinal adenosine kinase levels through astrocyte-dependent mechanisms. *PAIN* 2018;159:1025–34.
- [97] Waldman SA, Camilleri M. Guanylate cyclase-C as a therapeutic target in gastrointestinal disorders. *Gut* 2018;67:1543–52.
- [98] Willert RP, Woolf CJ, Hobson AR, Delaney C, Thompson DG, Aziz Q. The development and maintenance of human visceral pain hypersensitivity is dependent on the N-methyl-D-aspartate receptor. *Gastroenterology* 2004;126:683–92.
- [99] Wittendorp MC, Boddeke HW, Biber K. Adenosine A3 receptor-induced CCL2 synthesis in cultured mouse astrocytes. *Glia* 2004;46:410–18.
- [100] Wu WP, Hao JX, Halldner-Henriksson L, Xu XJ, Jacobson MA, Wiesenfeld-Hallin Z, Fredholm BB. Decreased inflammatory pain due to reduced carrageenan-induced inflammation in mice lacking adenosine A3 receptors. *Neuroscience* 2002;114:523–7.
- [101] Yoon MH, Bae HB, Choi JI. Antinociception of intrathecal adenosine receptor subtype agonists in rat formalin test. *Anesth Analg* 2005;101:1417–21.
- [102] Zahn PK, Straub H, Wenk M, Pogatzki-Zahn EM. Adenosine A1 but not A2a receptor agonist reduces hyperalgesia caused by a surgical incision in rats: a pertussis toxin-sensitive G protein-dependent process. *Anesthesiology* 2007;107:797–806.
- [103] Zhang M, Hu H, Zhang X, Lu W, Lim J, Eysteinnsson T, Jacobson KA, Laties AM, Mitchell CH. The A3 adenosine receptor attenuates the calcium rise triggered by NMDA receptors in retinal ganglion cells. *Neurochem Int* 2010;56:35–41.
- [104] Zhang X, Zhang M, Laties AM, Mitchell CH. Balance of purines may determine life or death of retinal ganglion cells as A3 adenosine receptors prevent loss following P2X7 receptor stimulation. *J Neurochem* 2006;98:566–75.
- [105] Zylka MJ. Pain-relieving prospects for adenosine receptors and ectonucleotidases. *Trends Mol Med* 2011;17:188–96.