



Additive Main Effect and Multiplicative Interactions (AMMI) and Regression Analysis in Sorghum [*Sorghum bicolor* (L.) Moench] Varieties

Gebeyehu Chala^{1*}, Bulti Tesso², Dagnachew Lule³ and Kebede Dessalegn⁴

¹Oromia Agricultural Research Institute, Mechara Agricultural Research Center

²Haromaya University, Alemaya, Ethiopia, ³Oromia Agricultural Research Institute, Addis Ababa, Ethiopia,

⁴Bako Agricultural Research Center, Bako, Ethiopia

*Corresponding author: gebeyehuchal@gmail.com

Original submitted in on 11th January 2019. Published online at www.m.elewa.org/journals/ on 30th April 2019
<https://dx.doi.org/10.4314/jab.v136i1.4>

ABSTRACT

Background and justification: Ethiopia is the third largest sorghum producer in Africa next to Nigeria and Sudan. Shortage of widely adapted and stable high yielding variety is one of the major bottlenecks for production and productivity of sorghum in the country. Grain yield performance is not the only parameter for selection as a genotype with the highest grain yield would not necessarily mean stable and adaptable across location and years. Eberhart and Russell (1996) and AMMI model could be the preferable tools to identify stable, high yielding and adaptable genotype (s) for wider or specific environments.

Objectives: To identify stable high yielding sorghum varieties that could be adapted for wider and/or specific environments and make recommendations for further demonstration and production in the test environments and similar agro ecologies.

Material and methods: A total of 21 released sorghum varieties and a local check were evaluated at three locations in western Ethiopia (Bako, Gute, Biloboshe) and two locations in eastern Ethiopia (Mechara and Mieso) in 2017 main cropping seasons. The trial was arranged in a randomized complete block design (RCBD) in three replications.

Summary of results and application of the study: The combined analysis of variance revealed highly significant effect of environment and genotype by environment interactions for grain yield. This indicated that the tested varieties showed inconsistent grain yield performance across locations. Birmash variety gave the highest grain yield with average yield of 3.5 ton ha⁻¹ but specifically adapted to Gute, Biloboshe and Mechara. Baji was the second high yielding variety with mean grain yield of 3.3 ton ha⁻¹ and relatively with wider adaptability. The first two IPCAs accounted for a total of 88.64% of the interaction sum square. In general, deviation from regression coefficient, AMMI stability value and genotype selection index revealed that Baji, Birmash, Emahoy, IS9302 and Gambella-1107 were relatively stable varieties with optimum grain yield and therefore recommended for further demonstration and popularization in the test locations and areas with similar agro-ecologies.

Keywords: AMMI, ASV, IPCAs, Sorghum

INTRODUCTION

Sorghum [*Sorghum bicolor* (L.) Moench] is a C4 cereal crop belonging to the family *Gramineae*. It has $2n = 20$ chromosomes and an estimated genome size of 750 Mb being twice the genome of rice and six times the genome of *Arabidopsis* (Passardi *et al.*, 2004). Sorghum is a dryland cereal crop grown on approximately 44 million hectares of land (Prakash *et al.*, 2010) in 99 countries (ICRISAT, 2009) with an annual production of 60 million tons (Iqbal *et al.*, 2010). It is widely cultivated in different parts of Ethiopia. Several authors reported that Ethiopia is the primary center of origin and hence, center of diversity for sorghum (De Wet and Huckabay, 1967; Doggett, 1988; Smith and Frederiksen, 2000). Sorghum is widely grown in the dry areas of Africa, Asia, America and Australia (Dickon *et al.*, 2006). Although sorghum is cultivated both in tropical and temperate climates, it is best known for its adaptation to the drought-prone semi-arid tropical (SAT) regions of the world (Baummhardt, 2000) and used for food for the poorest people who live in semiarid regions of the world (Jiang *et al.*, 2013). It can be adapted to environments with 400-600 mm annual rainfall that are too dry for other cereals to grow (Dickon *et al.*, 2006). Sorghum is one of the most important cereal crops planted as food insurance, especially in the moisture deficit lowlands of eastern, northern and north-eastern parts of Ethiopia where the climate is characterized by unpredictable drought and erratic rainfall (Degu *et al.*, 2009). It is also one of the most important cereal crops of the tropics grown extensively over wider areas with altitude ranging from 400 to 3000 meters above sea level (m.a.s.l) due to its ability to adapt the adverse environmental conditions. It is the major source of energy and protein for millions of people living in arid and semi-arid region of the world. It occupied third position in terms of production in Africa after wheat and maize and fifth in the world after wheat, maize, rice and barley (FAO, 2017). Moreover, it is widely used as a source of nutrition, fodder, biofuel, fiber and confection (Abubakar and Bubuche, 2013). It is able to grow under severe stress conditions. Sorghum can be cultivated successfully on almost all soils and in the temperature range of 16–40°C (Abubakar and Bubuche, 2013). Ethiopia is the third largest sorghum producer in Africa next to Nigeria and Sudan (FAOSTAT, 2012). In Ethiopia, a total of 4.34 million tons of sorghum is being produced per annum. The mean yield level in the country is estimated at 2.4 t ha⁻¹. The crop is the major food cereal after maize and tef in terms of number of growers, area coverage and grain production in the country (FAO, 2017). Oromia, Amhara and Tigray regions are the three major sorghum producers in the country (CSA, 2016). Out of the total sorghum area harvested in 2014 main cropping season, Oromia region accounts 39.92% (669,575.97 hectares), Amhara and Tigray regions contributed 33.31% (558,827.95 hectares) and 12.82% (215,111.82 hectares), respectively. In multi-environment trials, the phenotype of an individual is a measure of an environment main effect, a genotype main effect, and the genotype by environment interaction (GEI) (Yan and Tinker, 2005). More than forty sorghum varieties were released in the country from different regional and national research centers during the last 40 years (MoA, 2017). However, most of the varieties were not evaluated for their specific and wider adaptability and thus exhibit fluctuating yields when grown in different environments or agro-climatic zones. To this end, multi-environment adaptability and stability test is crucial to identify stable high yielding and adaptable varieties and discover sites that best represent the target environment (Yan *et al.*, 2000). Adaptability is the result of genotype, environment and genotype by environment interaction. A given variety could perform at an acceptable level in a range of environments, referred to as general adaptability, but others could perform well only in desirable environments, known as specific adaptability (Farshadfar and Sutka, 2008). Therefore, the present study aims to identify stable high yielding sorghum varieties that could be adapted for wider and/or specific environments and make recommendations for further demonstration and production in the test environments and similar agro ecologies.

MATERIALS AND METHODS

Description of the Study Area: Field experiment was conducted during 2017 main cropping season at five locations in Ethiopia where sorghum is widely grown. The locations were Bako, Gute, Biloboshe (Western

Oromia), Mechara, and Mieso (Eastern Oromia). The detailed agro-ecological conditions of the locations are presented in Table 1.

Table 1: Agro-ecological features of the experimental locations.

Locations	Altitude (m.a.s.l)	Ave. Rain fall (mm)	Soil Type	Geographic coordinates		Ave. Temp. (°C)	
				Latitude	Longitude	Max.	Min.
Gute	1906	1633.5	Alfisoils	9°00'N	36°38'E	21.6	14.3
Biloboshe	1758	1568.6	Sandy Loam	9°00'N	38°10'E	21.4	14.2
Bako	1650	1425.3	Alfisoils	9°6' N	37°09'E	20.4	13.5
Mechara	1760	871	Sandy loam	8°36'N	40°18 'E	23.4	8.9
Mieso	1470	856.8	Vertisol	16°06'N	37° 08'E	35.0	8.3

Source: Bako and Mechara Agricultural Research Centers

Plant Materials: A total of 21 sorghum varieties released from different research centers in Ethiopia were evaluated against one local check. The detailed

information about the experimental materials is presented in Table 2.

Table 2: Description of different sorghum varieties tested at five locations.

# No	Variety	Pedigree	Year of Release	Adaptation area (m.a.s.l.)	Breeder/Maintainer
1	Baji	85 MW 5334	1996	1600-1900	MARC/EIAR
2	Birmash	NA	1989	1600-1900	MARC/EIAR
3	Geremew	87 BK -4122	2007	1600-1900	MARC/EIAR
4	Lalo	BRC-245	2006	>1600	BARC/OARI
5	Teshale	3443-2-0P	2002	1450-1850	SRARC/ARARI and MARC/EIAR
6	Melkam	WSV 387	2009	<1600	MARC/EIAR
7	Gobiye	P-9401	1999	<1850	MARC/EIAR
8	Abshir	P-9403	2000	<1850	MARC/EIAR
9	Dagim	IS10892XRS/R-20-8614-2 x IS	2011	1600-1900	SRARC
10	IS9302	NA	1981	1600-1900	MARC/EIAR
11	ESH-1	P-9501 A x ICSR14	2009	<1600	MARC/EIAR
12	Birhan	Key#8566	2002	<1850	SRARC/ARARI
13	Gambella-1107	NA	1981	1450-1850	MARC/EIAR
14	Emahoy	Pw01-092	2007	1600-1900	PARC/EIAR
15	Dekeba	ICSR 24004	2012	<1600	MARC/EIAR
16	Chemeda	Acc-BCC-5	2013	>1600	BARC/OARI
17	Local Check	-	-	-	Farmers
18	07MW6035	(89MW4122*85MW5552)*85MW5340	2016	1600-1900	MARC/EIAR
19	07MW6002	(89MW4122*85MW5552)*85MW5340	2016	1600-1900	MARC/EIAR
20	Assosa_1	Bambasi # 9	2015	1500-1850	AARC
21	Adukara	NA	2015	1500-1850	AARC
22	07MW6052	(89MW4122*85MW5552)*85MW5340	2016	1600-1900	MARC/EIAR

Key: EIAR=Ethiopian Institute of Agricultural Research, MARC=Melkasa Agricultural Research Center, BARC= Bako Agricultural Research Center, SRARC= Sirinka Agricultural Research Center, ARARI=Amhara Regional Agricultural Research Institute, OARI= Oromia Agricultural Research Institute, PARC= Pawe Agricultural Research Center, AARC= Assosa Agricultural Research Center, NA= Not Available

Experimental procedures: The trial was laid out in Randomized Complete Block Design (RCBD) with three replications. The experimental plot consisted of two rows, each 5 m in length with 75 cm between row spacing and 15 cm spacing between plants. Seeds were sown by hand drilling at the rate of 12 kg ha⁻¹ as per the recommendation for row planting in sorghum. Thinning was done two weeks after emergence to adjust plant to plant spacing. 100 kg ha⁻¹ NPS fertilizer was applied at planting. Urea was applied as top

dressing at the rate of 50 kg ha⁻¹ at knee height stage. The field was kept free of weeds by hand weeding during the whole growing period.

Stability analysis

Eberhart and Russell's model: Yield stability was determined following the Eberhart and Russell (1966) model by regressing of the mean grain yield of individual genotypes on environmental index and calculating the deviation from the regression.

$$Y_{ij} = \mu_i + \beta_{ij}l_j + \delta_{ij} + \epsilon_{ij}$$

Where:

Y_{ij} = the mean of the ith genotype in the jth environment,
 μ_i = the grand mean,
 β_i = the regression coefficient of the ith genotype on environmental index,
 l_j = the environmental index obtained by the difference between the mean of each environment and the grand

mean, δ_{ij} = the regression deviation of the ith cultivar in the jth environment

Additive Main effect and Multiplicative Interaction (AMMI) model :

The AMMI model was calculated using the formula suggested by Zobel (*et al.*, 1988):

$$Y_{ij} = \mu + \alpha_i + \beta_j + \sum_{n=1}^N \lambda_n Y_{in} \delta_{jn} + \theta_{ij} + \epsilon_{ij}$$

Where: Y_{ij} = the mean yield of genotype i in environment j, μ = the grand mean, α_i = the deviation of the genotype mean from the grand mean, β_j = the deviation of the environment mean from the grand mean, λ_n = the singular value for the IPCA n, N = the number of PCA axis retained in the model, Y_{in} = the PCA score of a genotype for PCA axis n, δ_{jn} = the

environmental PCA score for PCA axis n, θ_{ij} = the AMMI residual and ε_{ij} = the residuals.

AMMI's stability value (ASV): The AMMI model is essential to quantify and rank genotypes according to their yield stability. This value was calculated according to Purchase (1997) as follow:

$$(ASV) = \sqrt{\left[\left(\frac{IPCA1SS}{IPCA2SS}\right)(IPCA1Score)\right]^2 + (IPCA2Score)^2}$$

In effect, the ASV is the distance from zero in a two dimensional scatter graph of IPCA1 (Interaction Principal Component Analysis axis 1) scores against IPCA 2 scores. Since the IPCA1 score contributes more to G x E sum of squares, it has to be weighted by the proportional difference between IPCA1 and IPCA2 scores to compensate for the relative contribution of IPCA1 and IPCA2 to the total G x E sum of squares.

Genotype Selection Index (GSI): Stability is not the only parameter for selection as the most stable genotypes would not necessarily give the best yield performance. Therefore, based on the rank of mean grain yield of genotypes (RY_i) across environments and rank of AMMI stability value (RASV_i), genotype selection index (GSI) was calculated for each genotype as: RY_i = RASV_i + GS_i. A genotype with the least GSI is considered as the most stable (Farshadfar, 2008).

RESULTS AND DISCUSSION

Analysis of variance The mean grain yield of varieties averaged over environments indicated that, Birmash, Baji and IS9302 followed by Emahoy gave higher grain yield (3.52, 3.34, 3.21 and 3.19 ton ha⁻¹), respectively and the lowest for Abshir (1.52 ton ha⁻¹) (Table 3). The result of the combined ANOVA showed that, the total variation in grain yield was attributed due to

environmental (19.34%), genotypic (19.78%) and GEI (47.85%) effects (Table 5). This implied that the largest proportion of the variation was due to the interaction effect than the main effect. Similar results were reported by Asfaw (2007); Hagos and Fetien (2011); Mahnaz *et al.* (2013) and Sewagegne *et al.* (2013).

Table 3. Mean grain yield (tons ha⁻¹) across different locations in 2017 cropping season

# No	Varieties	Grain yield (ton ha ⁻¹) of testing locations					Mean GY
		Bako	Biloboshe	Gute	Mechara	Miesso	
1	Baji	3.123 ^{ab}	4.136 ^{ab}	4.311 ^b	3.863 ^{a-d}	1.287 ^{cd}	3.34
2	Birmash	2.674 ^{b-e}	4.229 ^a	4.795 ^{ab}	4.159 ^{a-c}	1.721 ^{a-d}	3.52
3	Geremew	2.808 ^{a-c}	3.294 ^{a-d}	4.655 ^{ab}	2.271 ^{f-k}	1.037 ^d	2.81
4	Lalo	2.909 ^{a-c}	4.051 ^{ab}	4.778 ^{ab}	2.110 ^{g-l}	1.680 ^{b-d}	3.17
5	Teshale	2.598 ^{b-e}	2.378 ^{de}	1.295 ^{ij}	4.331 ^{a-c}	2.500 ^a	2.62
6	Melkam	2.720 ^{a-d}	1.512 ^e	1.487 ^{g-j}	3.717 ^{b-e}	1.693 ^{a-d}	2.23
7	Gobiye	2.096 ^{d-f}	2.339 ^{de}	0.951 ^{ij}	2.280 ^{f-j}	1.476 ^{b-d}	1.78
8	Abshir	1.609 ^f	1.503 ^e	0.597 ^j	2.325 ^{f-i}	1.571 ^{b-d}	1.52
9	Dagim	1.675 ^f	4.034 ^{a-c}	4.419 ^{ab}	2.615 ^{d-h}	1.232 ^{cd}	2.79
10	IS9302	2.987 ^{a-c}	3.731 ^{a-c}	4.461 ^{ab}	3.490 ^{b-g}	1.368 ^{b-d}	3.21
11	ESH-1	2.801 ^{a-c}	1.538 ^e	1.735 ^{f-i}	2.333 ^{f-i}	1.832 ^{a-d}	2.05
12	Birhan	2.056 ^{ef}	1.477 ^e	0.682 ^j	2.429 ^{e-h}	1.650 ^{b-d}	1.61
13	Gambella-1107	2.452 ^{c-e}	2.994 ^{b-d}	2.634 ^{d-f}	4.121 ^{a-c}	1.747 ^{a-d}	2.97
14	Emahoy	2.970 ^{a-c}	2.884 ^{cd}	3.251 ^{cd}	4.871 ^{ab}	1.975 ^{a-c}	3.19
15	Dekeba	1.597 ^f	2.344 ^{de}	1.895 ^{f-i}	4.090 ^{a-c}	1.442 ^{b-d}	2.27
16	Chemeda	2.366 ^{c-e}	1.589 ^e	2.177 ^{e-h}	3.593 ^{b-f}	2.200 ^{ab}	2.38
17	Local check	2.635 ^{b-e}	1.357 ^e	4.445 ^{ab}	0.721 ^m	1.293 ^{cd}	2.12
18	07MW6035	2.750 ^{a-c}	3.425 ^{a-d}	3.099 ^{c-e}	0.941 ^{j-m}	1.241 ^{cd}	2.35
19	07MW6002	2.988 ^{a-c}	3.247 ^{a-c}	5.444 ^a	0.99 ^{j-m}	1.212 ^{cd}	2.82
20	Assosa_1	2.988 ^{a-c}	1.440 ^e	1.232 ^{h-j}	3.433 ^{c-g}	1.559 ^{b-d}	2.01
21	Adukara	2.379 ^f	1.468 ^e	2.823 ^{e-g}	5.120 ^a	1.559 ^{b-d}	2.76
22	07MW6052	2.834 ^{a-c}	3.693 ^{a-c}	3.929 ^{bc}	1.952 ^{h-m}	1.025 ^d	2.6
Mean		2.525	2.68	2.921	2.98	1.559	2.553
CV%		13.2	31.5	18.9	24.4	27.5	
LSD (0.05)		0.551	1.382	0.91	1.199	0.708	

Stability Analysis using Eberhart and Russell Regression Model; According to Eberhart and Russell (1966), a stable genotype should have high yield, unit regression coefficient (b_i) and deviation from regression (Sd_i^2) close to zero. Accordingly, varieties such as Gambella-1107 and Emahoy had regression coefficient closer to unity, deviation from regression very close to zero with and mean grain yield greater than the average and hence could be considered as stable

varieties (Table 4). Varieties such as Dekeba and Lalo showed regression coefficient close to unity but Dekeba was specifically adapted to Mechara and Lalo to Gute & Billo Boshe site. Varieties such as Chemeda, ESH-1 and Gobiye had regression coefficients less than one, implying their specific adaptability to marginal environments. Miesso site was low yielding environment in 2017.

Table 4: Mean yield, regression coefficients and deviation from regression

Varieties	Yield (ton ha ⁻¹)	Rank	b _i	Ranks	S ² d _i	Ranks	ri ²	MS-TXL	MS-REG
07MW6002	2.777	10	1.338	5	3.50	22	0.01	2.67	0.15
07MW6035	2.291	14	0.637	6	1.49	17	0.04	1.16	0.18
07MW6052	2.785	9	1.407	7	1.15	14	0.06	0.92	0.22
Abshir	1.521	22	-0.060	18	0.50	7	0.50	0.75	1.49
Adukara	2.664	11	3.830	22	2.69	20	0.17	2.16	1.11
Assosa_1	1.651	21	2.513	21	1.16	15	0.12	0.88	0.32
Baji	3.344	2	2.029	20	0.16	1	0.74	0.47	1.41
Birhan	1.659	20	-0.064	17	0.58	9	0.46	0.81	1.50
Birmash	3.515	1	1.966	19	0.41	5	0.50	0.61	1.24
Chemeda	2.385	13	0.375	10	0.66	10	0.21	0.62	0.52
Dagim	2.795	7	1.675	11	1.35	16	0.13	1.17	0.61
Dekeba	2.273	15	1.107	1	0.97	11	0.01	0.73	0.02
Emahoy	3.190	4	1.425	8	0.55	8	0.13	0.47	0.24
ESH-1	2.048	18	0.077	16	0.34	3	0.52	0.54	1.13
Gambella-1107	2.789	8	1.173	2	0.38	4	0.03	0.29	0.04
Geremew	2.813	6	1.680	12	1.06	12	0.16	0.95	0.61
Gobiye	1.828	19	0.184	13	0.45	6	0.39	0.56	0.88
IS9302	3.208	3	1.843	15	0.24	2	0.57	0.41	0.95
Lalo	3.166	5	1.303	3	1.68	18	0.02	1.29	0.12
Local check	2.091	17	0.692	4	2.73	21	0.02	2.08	0.13
Melkam	2.226	16	0.577	9	1.11	13	0.07	0.89	0.24
Teshale	2.584	12	0.182	14	1.73	19	0.15	1.52	0.89

Key: MS-TXL = contribution of each variety to interaction MS, MS-REG = contribution of each variety to the regression component of the treatment by location interaction, MS-DEV (sd²) =deviations from regression component of interaction, ri² = squared correlation between residuals from the main effects model and the site index, b_i = regression coefficient

Additive main effect and multiplicative interactions (AMMI) model:

The combined AMMI ANOVA of the twenty two sorghum varieties over five locations for grain yield (ton ha⁻¹) is presented in Table 5. The ANOVA table indicated highly significant differences (p<0.01) for the environments, genotypes, GEI and the first two IPCA. The total variation explained was 86.45% for treatments (20.08% was due to the genotypes, 17.71% was due to the environment and 48.66% was due to their interaction) and the remaining 13.55% for error. The greater contribution of the treatment over the error indicated the reliability of this multi-location experiment. The lion share of variation was largely due to GEI. As discussed earlier, the high percentage of GEI is an indication for the highly variable sorghum growing environment. This also revealed that there was a differential yield performance among the released varieties across testing

environments. As G x E interaction was significant, further calculation of genotype stability was conducted. In the AMMI ANOVA, the GEI was further partitioned using PCA. The number of PCA axis to be retained is determined by testing the mean square of each axis with the estimate of residual using the F-statistics. The result of ANOVA showed that the first two IPCA were highly significant at (P<0.01) implying the inclusion of the first two interactions PCA axes in the model. Hence, the best fit AMMI model for this multi-location yield trial data was AMMI-2 (Table 5). Gauch and Zobel (1996) suggested that the most accurate model for AMMI can be predicted by using the first two IPCAs. Several authors took the first two IPCAs such as for bread wheat (Asnake *et al.*, 2013), common bean (Abeya *et al.*, 2008) and finger millet (Dagnachew *et al.*, 2014, Kebede *et al.*, 2018).

Table 5: AMMI analysis of variance for grain yield (ton/ha) of sorghum varieties tested at five locations during 2017 main cropping season.

Source	DF	Sum of squares explained				MS
		SS	%Total	%Contribution to the variation	% G x E	
Total	329	523				1.59
Treatments	109	452.1	86.44			4.148**
Genotypes	21	105	20.08			5.001**
Environments	4	92.6	17.71			23.14**
Block	10	6.61				0.661*
G x E	84	254.5	48.66			3.029**
IPCA 1	24	186.2			73.16	7.76**
IPCA 2	22	39.4			15.48	1.79**
Residuals	36	28.9			11.36	0.803
Error	206	64.3	12.29			0.312

Key: DF = degree of freedom, SS =sum of squares, MS = mean of squares and, GxE = Genotype by Environment, * significant (P<0.05), ** = highly significant (P<0.01).

AMMI Stability Values (ASV) and Genotype Selection Index : In AMMI stability value analysis (ASV), a genotype with least ASV score is the most stable across environments and the larger the ASV value, either negative or positive, the more specifically adapted a genotype is to certain environments (Purchase, 1997). Accordingly, Gambella-1107 and Bajji showed the least ASV than the others (Table 6).

Stability is not the only parameter for selection of high yielding genotypes as the most stable genotypes would not necessarily give the best yield performance. As such, the genotype selection index (GSI) revealed that Bajji, IS9302, Birmash, Gambella-1107 and Emahoy were among the top-ranking genotypes integrating both stability and grain yield performance parameters (Table 6).

Table 6. Mean grain yield, IPCA scores and ASV of 22 sorghum varieties evaluated at five locations during 2017 main cropping season.

Varieties	Yield (ton ha ⁻¹)	Rank	IPCA1	IPCA2	IPCA3	IPCA4	ASV	ASV Rank	GSI
07MW6002	2.777	10	-1.134	-0.269	0.309	-0.079	2.4791	22	32
07MW6035	2.291	14	-0.609	-0.491	-0.537	0.190	1.4122	15	29
07MW6052	2.785	9	-0.615	-0.142	-0.212	0.578	1.3440	13	22
Abshir	1.521	22	0.509	-0.401	-0.317	-0.270	1.1769	9	31
Adukara	2.664	11	0.781	0.297	0.680	0.266	1.7236	19	30
Assosa_1	1.651	21	0.543	0.275	0.066	0.154	1.2125	11	32
Baji	3.344	2	-0.287	-0.526	-0.108	0.378	0.8146	3	5
Birhan	1.659	20	0.526	-0.504	-0.224	-0.058	1.2483	12	32
Birmash	3.515	1	-0.304	0.681	-0.029	-0.186	0.9480	7	8
Chemeda	2.385	13	0.484	-0.194	0.339	-0.365	1.0690	8	21
Dagim	2.795	7	-0.604	0.544	-0.322	-0.545	1.4216	16	23
Dekeba	2.273	15	0.513	0.440	-0.102	-0.287	1.1983	10	25
Emahoy	3.190	4	0.380	0.393	0.264	0.079	0.9161	6	10
ESH-1	2.048	18	0.272	-0.648	0.126	0.097	0.8773	4	22
Gambella-1107	2.789	8	0.318	0.306	-0.118	-0.030	0.7552	1	9
Geremew	2.813	6	-0.678	0.130	0.192	0.121	1.4802	17	23
Gobiye	1.828	19	0.328	-0.370	-0.572	0.059	0.8046	2	21
IS9302	3.208	3	-0.362	0.392	0.074	0.179	0.8793	5	8
Lalo	3.166	5	-0.783	0.017	-0.328	-0.124	1.7020	18	23
Local check	2.091	17	-0.772	-0.639	0.858	-0.294	1.7960	20	37
Melkam	2.226	16	0.636	-0.197	0.234	0.241	1.3958	14	30
Teshale	2.584	12	0.858	-0.144	-0.275	-0.107	1.8694	21	33
Mean	2.553								

Key: IPCA=Interaction Principal Component Axis 1, 2, 3 and 4; ASV=AMMI Stability Value

SUMMARY AND CONCLUSIONS

Multi-location trials are very important for selecting the best genotype for wider or specific environments before advising or recommending crop varieties for commercial production. A total of 21 sorghum varieties were evaluated against the local check across five locations in 2017. The combined analysis of variance revealed highly significant effect of environment and genotype by environment interactions for grain yield. About 48.68% of the total variation was explained by G x E interaction effect. This implied that the environments were variable and varietal performance

across locations was inconsistent. Most of the tested varieties were adapted to specific location. For instance, 07MW6002, Adukara and Teshale variety showed better performance at Gute, Mechara and Mieso, respectively. Generally, the mean grain yield performance, ASV, deviation from regression coefficient and GSI revealed that Baji, Birmash, Emahoy, IS9302 and Gambella-1107 were relatively stable varieties with optimum grain yield and therefore recommended for further popularization.

REFERENCE

Abeya Temesgen, Chemedaba Daba, Girma Mengistu, Dagnachew Lule and Negash Geleta (2008). Regression and Additive mean effect and multiple interactions (AMMI) in common bean (*Phaseolus vulgaris* (L.) genotypes. *Ethiopian Journal of Biological Science*. 7(1):45-53

Abubakar, L. and Bubuche, T. S. 2013. Genotype x environment interaction on yield and its

component of sorghum (*Sorghum bicolor* (L.) Moench) genotypes of some selected states in North-Western Nigeria. *International Journal of Current Agricultural Research*, 1(6): 27-29. <http://www.wrpjournals.com/journals/V1/IJCAR> Accessed on June 1, 2014.

- Asfaw Adugna. 2007. Assessment of yield stability in sorghum. *African Crop Science Journal*, 15 (2): 83 – 92.
- Asnake W, Henry M, Temesgen Z, Girma T, 2013. Additive main effects and multiplicative interactions model (AMMI) and genotype main effect and genotype by environment interaction (GGE) biplot analysis of multienvironmental wheat variety trials. *African J. Agric. Res.* 8(12):1033-1040.
- CSA (Central Statistical Agency). 2016. Agricultural sample survey 2010/2011: report on area and production of crops (private peasant holdings, main season), vol. 1. Addis Ababa: Federal Democratic Republic of Ethiopia, Central Statistical Agency.
- Dagnachew Lule, Masresha Fetene, Santie de Villiers and Kassahun Tesfaye (2014). Additive Main Effects and Multiplicative Interactions (AMMI) and genotype by environment interaction (GGE) biplot analyses aid selection of high yielding and adapted finger millet varieties. *Journal of Applied Biosciences.* 76:6291–6304.
- De Wet JMJ, Huckabay JP (1967). The origin of *Sorghum bicolor*. II. Distribution and domestication. *Evolution* 21 (4):787-802.
- Doggett H (1988) *Sorghum*. Longman Scientific & Technical, London.
- Eberhart, S.A. and Russel, W.A. 1966. Stability parameters for comparing varieties. *Crop Science*, 6:36-40.
- FAOSTAT. 2012. Database of agricultural production. Rome: Food and Agriculture Organization of the United Nations. Available at <http://faostat.fao.org/default.aspx> accessed November 2015.
- FAOSTAT. 2017. Food and Agriculture Organization of the United Nations Data base of Agricultural Production. FAO Statistical Database s. Available at <http://faostat.fao.org/site/339/default.aspx>.
- Farshadfar, E. (2008). Incorporation of AMMI Stability Value and Grain Yield in a Single NonParametric Index (Genotype Selection Index) in Bread Wheat. *Pakistan Journal of Biological Sciences* 11:1791–1796.
- Gauch HG, Zobel RW, 1996. AMMI analysis of yield trials. In: Genotype by environment interaction. pp. 85-122 (Kang, M. and Gauch, H. eds.). Boca Raton. CRC press, New York.
- Hagos Tadesse and Fetien Abay. 2011. Additive Main Effects and Multiplicative Interactions analysis of yield performance of sesame genotypes across environments in Northern Ethiopia. *Journal of the dry lands*, 4(1): 259-266.
- ICRISAT (International Crops Research Institute for the Semi-Arid Tropics). 2009. online Available on the website <http://www.icrisat.org/> Assessed on July 5, 2011.
- Iqbal, A., Sadia, B., Khan, A.I., Awan, F.S., Kainthand, R.A. and Sadaqat, H.A. 2010. Biodiversity in the sorghum (*Sorghum bicolor* L. Moench) germplasm of Pakistan. *Genetics and Molecular Research*, 9 (2): 756-764.
- Kebede Dessalegn, Dagnachew Lule, Girma Mengistu, Kassa Mamo, Chemedda Birhanu, Girma Chemedda & Gudeta Bedada (2018). Genotype by Environment Interaction & Grain Yield Stability Analysis for Finger Millet Genotypes from Western Oromia. *Ethio. J Crop Sci.* 6 (1):119-134
- Mahnaz, R., Ezatollah, F. and Mohammad, M. J. 2013. Additive Main Effect and Multiplicative Interaction Analysis of phenotypic stability in chickpea genotypes over stress and non-stress environments. *International Journal of Agriculture and Crop Sciences*, 5(36):253-260.
- MOA (Ministry of Agriculture). 2017. Annual action plan report on cereals. Amhara National Regional State Bureau of Agriculture, Bahir Dar, Ethiopia.
- Prakash, R., Ganesamurthy, K., Nirmalakumari, A. and Nagarajan, P. 2010. Heterosis for fodder yield in sorghum (*Sorghum bicolor* L. Moench). *Electronic Journal of Plant Breeding*, 1(3): 319-327.
- Passardi, F., Longet, D., Penel, C. and Dunand, C. 2004. The class III peroxidase multigenic family in rice and its evolution in land plants. *Phytochemistry*, 65: 1879–1893.
- Purchase, J.L. 1997. Parametric analysis to describe genotype by environment interaction and stability in winter wheat. PhD.thesis. Department of Agronomy, Faculty of Agriculture, University of the Orange Free State, Bloemfonten, South Africa.
- Sewagegne Tariku, Tadesse Lakew, Mulugeta Bitew and Mitiku Asfaw, 2013. Genotype by environment interaction and grain yield stability analysis of rice (*Oryza sativa* L.) genotypes evaluated in north western

- Ethiopia. *Net Journal of Agricultural Science*, 1(1): 10-16.
- Smith CW, Frederiksen RA (2000) Sorghum: origin, history, technology, and production, vol 2. John Wiley & Sons, USA.
- Yan, W., Hunt, L.A., Sheng, Q. and Szlavnic, Z. 2000. Cultivar evaluation and mega environment investigation based on the GGE biplot. *Crop Science*, 40: 597–605.
- Yan, W. and Tinker, N.A. 2005. An integrated biplot analysis system for displaying, interpreting and exploring genotype by environment interactions. *Crop Science*, 45: 1004–16.
- Zobel, W.R., Wright, M.J. and Gauch, H.G. 1988. Statistical analysis of a yield trial. *Agronomy Journal*, 80:388-393