

Review

Adenosine signaling and function in glial cells

D Boison^{*1}, J-F Chen² and BB Fredholm³

Despite major advances in a variety of neuroscientific research fields, the majority of neurodegenerative and neurological diseases are poorly controlled by currently available drugs, which are largely based on a neurocentric drug design. Research from the past 5 years has established a central role of glia to determine how neurons function and, consequently, glial dysfunction is implicated in almost every neurodegenerative and neurological disease. Glial cells are key regulators of the brain's endogenous neuroprotectant and anticonvulsant adenosine. This review will summarize how glial cells contribute to adenosine homeostasis and how glial adenosine receptors affect glial function. We will then move on to discuss how glial cells interact with neurons and the vasculature, and outline new methods to study glial function. We will discuss how glial control of adenosine function affects neuronal cell death, and its implications for epilepsy, traumatic brain injury, ischemia, and Parkinson's disease. Eventually, glial adenosine-modulating drug targets might be an attractive alternative for the treatment of neurodegenerative diseases. There are, however, several major open questions that remain to be tackled.

Cell Death and Differentiation (2010) 17, 1071–1082; doi:10.1038/cdd.2009.131; published online 18 September 2009

It is becoming increasingly clear that inflammatory processes play an important role in neurodegenerative disease, just as inflammation is becoming increasingly implicated in various systemic diseases elsewhere in the body.¹ As the immune responses in brain show uncommon features, it was long considered to be 'immune privileged'. However, cells that are involved in adaptive immune reactions do enter the brain, and this can result in major CNS pathology. Multiple sclerosis is an important example. Furthermore, immune reactions, often attributable to the innate immune system, feature in many, if not all, neurodegenerative diseases. Microglial cells, astrocytes, endothelial cells, oligodendrocytes, and neurons all produce signals to orchestrate these reactions. Arachidonic acid metabolites, nitric oxide, cytokines, and chemokines all appear to play some role. Recently, the role of ATP and adenosine as critically important signaling molecules has become appreciated.^{2,3} This awareness that a multiplicity of signals other than those principally involved in nerve–nerve communication play a role in neurodegenerative disease has also meant that we have to consider other cells than neurons as critically important players. The present brief review will focus on adenosine (and to a lesser extent ATP) signaling, and the role of glial cells in neurodegenerative disease. The review will feature results from the authors' own work more prominently than would be motivated from an objective standpoint.

Glial Control of Adenosine Homeostasis

In mammals, purine *de novo* synthesis proceeds via formation of IMP, which is then converted into AMP; however, there is no *de novo* synthesis pathway for adenosine. Physiologically, intracellular adenosine can be formed by either dephosphorylation of AMP by 5'-nucleotidase, or, alternatively, by hydrolysis of *S*-adenosylhomocysteine, whereas extracellular adenosine can be formed from released adenine nucleotides by a cascade of ectonucleotidases.⁴ Two metabolic pathways are responsible for the removal of adenosine: deamination into inosine via adenosine deaminase (ADA; EC 3.5.4.4) and phosphorylation into AMP via adenosine kinase (ADK; EC 2.7.1.20). Based on its low K_M for adenosine, ADK is considered to be the primary route of adenosine metabolism.⁵ Recent findings indicate that extracellular levels of adenosine, and consequently the levels close to synapses, are largely regulated by astrocytes,^{6–10} and an astrocyte-based adenosine cycle has been proposed.^{11,12}

Glial release of ATP as source for extracellular adenosine. ATP can be released from neurons and astrocytes, is identified as a neurotransmitter in both CNS and PNS, and exerts a multitude of largely excitatory effects by activation of specific ATP receptors (P2X and P2Y receptors).¹³ Vesicular release has also been clearly

¹Robert Stone Dow Neurobiology Laboratories, Legacy Research, Portland, OR 97232, USA; ²Department of Neurology, Boston University School of Medicine, Boston, MA 02118, USA and ³Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, S-17177, Sweden

*Corresponding author: D Boison, Robert Stone Dow Neurobiology Laboratories, Legacy Research, 1225 NE 2nd Avenue, Portland, OR 97232, USA.

Tel.: +1 503 413 1754; Fax: +1 503 413 5465; E-mail: dboison@downeurobiology.org

Keywords: astrocyte; adenosine receptor; adenosine kinase; epilepsy; excitotoxicity; neurodegeneration

Abbreviations: AAT, adenosine augmentation therapy; ADA, adenosine deaminase; ADK, adenosine kinase; AMPA, α -amino-3-hydroxyl-5-methyl-4-isoxazole-propionate; AR, adenosine receptor; CBF, cerebral blood flow; CCI, controlled cortical impact; EGFP, enhanced green fluorescent protein; FACS, fluorescence activated cell sorting; MAP, microtubule associated protein; NMDA, *N*-methyl-D-aspartic acid; MPTP, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; NOS, nitric oxide synthase; NTPDase, nucleoside triphosphate diphosphohydrolase; PD, Parkinson's disease; SNARE, soluble NSF attachment protein; SOD, superoxide dismutase

Received 10.6.09; revised 21.7.09; accepted 27.7.09; Edited by RA Knight; published online 18.9.09

demonstrated from endocrine cells, and here release of ATP may differ in several ways from that of the hormone, because of so-called kiss-and-run release.¹⁴ Thus, ATP can be released even in situations when the vesicle fusion is too incomplete and transient to allow release of the stored hormone. Perhaps this can occur also in nerves. However, under physiological conditions the vesicular release of ATP, not from neurons, but from astrocytes, has been identified as a major source of synaptic adenosine.¹⁵ Transgenic mice that express a dominant-negative soluble NSF attachment protein (SNARE) domain selectively in astrocytes were characterized by the loss of the adenosine A₁ receptor-mediated tonic inhibition in synaptic slices, indicating that under physiological conditions astrocytic release of ATP (followed by degradation into adenosine via ectonucleotidases) is a major source of adenosine¹⁵ that affects synaptic transmission. A kiss-and-run-like exocytotic release of ATP has been detected in astrocytes.¹⁶ Another, possibly related, mechanism involves release of ATP from a lysosome pool in astrocytes.¹⁷ However, astrocytes appear to use also other mechanisms to release ATP as a signal. One of those involve connexin hemichannels,¹⁸ but recent work provides strong reason to assume that pannexin-1, rather than connexin-43, provides the most responsible channel.¹⁹ Another proposal is that ATP is released via maxi-anion channels, and this may be particularly important in astrocytic swelling.²⁰ The P2X₇ receptor may in some conformations allow ATP to be released. Finally, ATP may be released whenever there are increases in membrane volume via incorporation of intracellular vesicles into the membrane, or when there is a shedding of small vesicles.²¹ In addition to the mechanisms described above, astrocytes can directly release adenosine, especially in response to hypoxic stimulation,^{6,22} even though release of adenosine *per se* is more typical of neurons.²³ In that case the release depends on export of adenosine via the equilibrative nucleoside transporters.

Since astrocytes can contact thousands of synapses and coordinate synaptic networks,^{8,24} it is conceivable that astrocytic release of ATP and its subsequent degradation into adenosine has a major regulatory function in setting a global adenosine-mediated inhibitory tone within a neuronal network. In addition, other glial cells can also contribute. For example, in retinal tissue, stimulation of glial cells leads to an adenosine-mediated inhibition of neuronal activity, but in this case Muller cells rather than astrocytes appear to be most important.²⁵

The distribution of adenosine formed from breakdown of released ATP will obviously also depend on the distribution of the enzymes that degrade the nucleotide. It was shown long ago that 5'nucleotidase tended to accumulate in areas of a lesion,²⁶ and we now know that CD73, the ecto-5'nucleotidase, is highly expressed in microglial cells, as is the ecto-NTPDase (nucleoside triphosphate diphosphohydrolase), CD39.²⁷

Elimination of adenosine via astrocytic adenosine kinase. Several lines of evidence indicate that astrocytic ADK is the key regulator for ambient levels of adenosine: (i) In adult brain, ADK is predominantly expressed in astrocytes;¹⁰ (ii) pharmacological inhibition of ADK is

sufficient to prevent seizures in various models of epilepsy;²⁸ (iii) genetic knockout or knockdown of ADK in cultured cells induces secretion of adenosine into the medium;^{29–32} (iv) transgenic overexpression of ADK triggers seizures by reduction of ambient adenosine;³³ (v) inhibition of ADK in hippocampal slices increases endogenous adenosine and depresses neuronal firing, whereas inhibition of ADA had no effect;³⁴ (vi) a substrate cycle between AMP and adenosine, which involves ADK and 5'-nucleotidase, enables minor changes in ADK activity to rapidly translate into major changes in adenosine; and³⁵ (vii) ADK activity is regulated in response to brain injury and is subject to developmental regulation.^{10,36,37} Based on these considerations, and based on the lack of a classical transporter-regulated reuptake system for adenosine and the ubiquitous presence of bidirectional equilibrative nucleoside transporters,³⁸ ADK likely fulfills the role of a metabolic reuptake system for adenosine. Thus, tight regulation of ADK expression levels and of its specific activity becomes a necessity. Therefore, it is not surprising that ADK is highly conserved in evolution, that no naturally occurring mutations of the *Adk* gene are known, and that a genetic disruption of the *Adk* gene is lethal.^{5,39}

The fact that astrocytic ADK is of critical importance in regulating extracellular adenosine concentrations implies that the nucleoside transporters that facilitate adenosine uptake into astrocytes are important. Astrocytes express one concentrating (sodium dependent) and two equilibrative nucleoside transporters.⁴⁰ Inhibition of this transport could potentially be used to elevate brain adenosine levels under conditions where extracellular adenosine is derived from extracellular ATP, but they would be less useful under conditions when adenosine is derived from intracellular production.⁴¹ It is interesting to note that cannabinoids can block the equilibrative transporter and this effect can partly explain the immunosuppressive effects of these compounds.⁴² The central role of astrocytes in regulating extracellular levels of adenosine is demonstrated in Figure 1.



Figure 1 Extracellular adenosine levels are thought to be regulated by an astrocyte-based adenosine cycle. Astrocytes can release ATP via vesicular release and/or by direct release through hemichannels (h-ch). Extracellular ATP is rapidly degraded into adenosine (ADO) by a series of ectonucleotidases. Adenosine can also be released directly via equilibrative nucleoside transporters (nt). Intracellularly adenosine levels are largely controlled by adenosine kinase, which is part of a substrate cycle between adenosine and AMP. Small changes in adenosine kinase activity rapidly translate into major changes in adenosine. Intracellular adenosine kinase is considered to be a metabolic reuptake system for adenosine. Only selected mechanisms and pathways are shown; for details please refer to the main text

Glial Adenosine Receptors

There are four types of evolutionarily conserved and pharmacologically well-characterized adenosine receptors (ARs) called A_1 , A_{2A} , A_{2B} , and A_3 ⁴³ (Figure 2). Adenosine is the endogenous agonist at all these receptors, but at A_1 and A_3 receptors inosine can act as a partial agonist.^{44,45} The A_1 and A_3 receptors couple to the G_i family of G proteins and, thus, stimulate K^+ channels, reduce transient voltage-dependent Ca^{2+} channels, and inhibit cAMP formation; A_{2A} receptors couple to members of the G_s family (G_{olf} in striatal neurons), whereas A_{2B} receptors couple to many G proteins including G_s , G_q , and G_{12} . Adenosine is approximately equipotent on A_1 , A_{2A} , and A_3 receptors, whereas A_{2B} receptors require higher agonist concentrations, if cAMP changes is the readout,⁴⁵ but if microtubule associated protein (MAP) kinase activation is used to measure receptor activation adenosine is virtually equipotent on all four receptors.⁴⁶ All four ARs are detected in astrocytes,²² and all have been reported to be expressed in microglial cells or microglial cell lines.^{47–49}

A_1 receptors. A_1 receptors on astrocytes reduce their proliferation rate in culture.⁵⁰ As in many other types of cells, activation of A_1 receptors can not only decrease cAMP accumulation but also stimulate phospholipase C, especially if this pathway is simultaneously activated by other stimuli.^{51–53} A_1 receptors help protect astrocytes from damage and cell death,^{22,54,55} partly via activation of PI3K and Erk 1/2 phosphorylation.

Nerve activity promotes myelination and it has been shown that this response is dependent on ATP,⁵⁶ but ATP acts on oligodendrocytes indirectly, because ATP acts on astrocytes to release leukemia inhibitory factor. This response may be modified by adenosine, as A_1 receptors are present in oligodendroglia⁵⁷ and stimulate their migration. It has been shown that A_1 receptor activation leads to white matter loss,

and that A_1 receptors contribute to hypoxia-induced white matter loss.⁵⁸ A_1 receptors on microglial cells are reported to reduce excessive activation of microglial cells upon immune activation.⁵⁹ Activation of these microglial receptors may secondarily affect oligodendroglial cells⁵⁹ and also astrocyte proliferation,⁶⁰ which emphasizes the possibility of an extended glial network of signaling. A_1 receptors on neurons (especially at nerve terminals) are critically important in mediating the dampening effect on neuronal activity mediated by adenosine generated from ATP released from astrocytes.^{6,61} The highly abundant A_1 receptors at nerve endings may preferentially signal via G_o proteins to inhibit transient calcium channels, whereas the same receptors in nerve cell bodies and dendrites may preferentially regulate potassium channel conductance via G_i proteins.

A_{2A} receptors. In brain, A_{2A} Rs are expressed at high levels in striatal neurons and at low levels in neurons outside of the striatum and in glial cells.^{62,63} Many functional measurements (such as cAMP levels and cytokine release) coupled with pharmacological tools have clearly demonstrated the presence and function of A_{2A} Rs in glial cells.⁶⁴ A_{2A} R binding densities are at the range ~30–60 fmole/mg protein in primary cultured microglial cells or in sorted microglial cells derived from striatum, as estimated by ³H-ligand binding studies.^{65,66} Furthermore, the expression of the A_{2A} R in glial elements in both the striatum and the solitary tract is confirmed by electronmicroscopic studies.^{63,67} It should be noted that A_{2A} R expression in microglia and astrocytes is usually low under physiological conditions and frequently below the detection limit of histological methods (i.e., immunohistochemistry, autoradiography, or *in situ* hybridization).^{62,64,68}

Importantly, expression of A_{2A} Rs in glial cells is induced following brain insults. For example, LPS treatment induced A_{2A} R mRNA and protein in primary cultures of (mixed) glial cells (mainly in microglial cells) at 16 h and peaked at 48 h after the treatment.⁶⁵ Recently, using double immunohistochemistry analysis, we demonstrated that A_{2A} R expression is induced in microglial cells and astrocytes of mouse substantia nigra at 24 h after 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) intoxication. The induction of A_{2A} Rs in glial cells by brain insults and inflammatory signals, coupled with local increase in adenosine and proinflammatory cytokine levels (such as IL-1 β , which further induces A_{2A} R expression), may serve as part of an important feed-forward mechanism to locally control neuroinflammatory responses in the brain. It has been shown that adenosine, acting on A_{2A} receptors, can increase extracellular levels of glutamate, both by reducing glutamate uptake via GLT-1 and by direct release.^{69,70} Thus, some of the reported effects of A_{2A} receptors on glutamate release may be based on mechanisms mediated by astrocytes rather than by neurons.

A_{2A} Rs in glial cells may exert complex actions on neuronal cell death (both, potentially deleterious as well as neuroprotective) and possibly other functions such as modulation of synaptic transmission. In astrocytes, activation of A_{2A} Rs by extracellular adenosine increases astrocyte proliferation and activation,^{71,72} but inhibits the expression of iNOS and the production of NO,⁷³ and regulates glutamate efflux by




Figure 2 ARs, their coupling to G-proteins, and some of the downstream consequences of receptor activation

astrocytes.⁶⁹ Thus, modulation of astrogliosis by A_{2A} Rs is likely involved in brain repair processes, possibly via formation of tissue scar. In microglial cells, activation of A_{2A} Rs has mixed effects on microglial proliferation, but has clear facilitating effects on the release of cytokines, including upregulation of cyclooxygenase 2 and release of prostaglandin E2 (PGE2),⁷⁴ and on increases in nitric oxide synthase (NOS) activity and NO release⁶⁵ and nerve growth factor expression.⁷⁵

A_{2B} receptors. It was shown early on that cyclic AMP accumulation in brain slices was due to an AR different than that responsible for adenylate cyclase stimulation in striatum, and it is now clear that a major part of the brain slice cAMP response is due to A_{2B} receptor activation on astrocytes.⁷⁶ A_{2B} receptors can couple with two different classes of G proteins, G_q and G_s . G_q regulates intracellular calcium and vesicular release, whereas G_s affects a plethora of cAMP-dependent signaling pathways. The A_{2B} receptor may also activate phospholipase C⁷⁷ and appears to be responsible for the adenosine-induced stimulation of IL-6 from astrocytes.⁷⁸ In airways, A_{2B} receptors, via cAMP, regulate chloride channels.⁷⁹ Since this occurs also in the intestine, it is fair to predict that cells in the CNS likewise alter chloride flux via A_{2B} R signaling. In particular, the possibility exists that adenosine is an important regulator of astrocytic swelling via modulation of volume-regulated anion channels. A_{2B} receptors may also play an important role in the development of the nervous system; it was initially proposed that the important regulator, netrin, required signaling via A_{2B} receptors,⁸⁰ but later studies showed that netrin does not require simultaneous A_{2B} R signaling,⁸¹ but that netrin signaling requires prior A_{2B} activation because the receptor is thereby regulated.⁸²

A_3 receptors. Astrocytic A_3 receptors appear to regulate chemokine release.⁸³ Activation of A_3 receptors by endogenous adenosine protects astrocytes from cell death induced, for example, by hypoxia.²² Microglial cells have functional A_3 receptors coupled to MSAP kinase and p38 signaling.^{47,84} Although adenosine itself has little effect on microglial migration, it was shown that migration induced by ATP,⁸⁵ perhaps predominantly via P2Y12 receptors,⁸⁶ is lost when ATP hydrolysis via CD39 is eliminated, and can be restored by adenosine.²⁷ Although it is not absolutely certain that the relevant AR is the A_3 R, it is tempting to speculate that there is a similarity to the situation in neutrophil leucocytes where ATP acting on a P2Y receptor acts in concert with adenosine acting on A_3 receptors to stimulate migration.⁸⁷ Microglial cells (and A_3 receptors) may also be of particular importance in the regulation of chemokine release and chemokine actions.⁸⁸

Glial Cells and Neurovascular Coupling

Glial cells play important roles in coupling neuronal function to the cerebral microvasculature that controls cerebral blood flow (CBF),^{89,90} in the sense that increased neuronal activity requires corresponding increases in CBF. Apart from large processes that stain for intermediate filaments and give

astrocytes their stellar appearance, astrocytes have a multitude of fine processes that have little overlap with processes from other astrocytes and that define individual astrocytic domains, which each contain 300–600 neuronal dendrites and 10^5 synapses in rodent hippocampus.^{8,91–93} Thus, a single astrocyte can sense the activity, and integrate the function, of hundreds of neurons within its domain. In addition, each astrocyte extends at least one process with endfeet surrounding blood vessels of the microvasculature. Therefore, astrocytes are uniquely located to adjust regional CBF to regional energy metabolism.

The vasodilator adenosine has been identified as an important mediator that couples CBF to neuronal activation.⁹⁴ Thus, adenosine was demonstrated to mediate glutamate-induced vasodilation in the cerebral cortex.^{95,96} Topical application of glutamate dilated pial arterioles, an effect that could be reversed by an A_{2A} R antagonist, but not by an A_{2B} R antagonist.⁹⁵ Likewise topical superfusion AMPA on the cortical surface through a closed cranial window resulted in increases in pial arteriolar diameter, an effect that could be reversed by A_{2A} and A_{2B} R blockade, but not by inhibition of NO synthase, cyclooxygenase-2, or cytochrome P-450 epoxygenase.⁹⁶ Apart from the activation of vascular ARs, adenosine can exert important regulatory functions by activation of astrocytic ARs.⁹⁷ Thus, the adenosine-evoked calcium response in acutely isolated astrocytes was found to be coupled to the A_{2B} receptor;⁷⁷ based on these findings adenosine could be implicated in promoting the propagation of calcium increases throughout astrocytic processes. Increased Ca^{2+} in turn is also associated with the release of ATP through connexin hemichannels, a process that is potentiated by A_{2B} R activation.^{98,99} Through these mechanisms ATP release and degradation into adenosine via ectonucleotidases appears to mediate arteriolar dilation in response to neuronal activation.¹⁰⁰ This process was dependent on astrocytes, since application of the selective gliotoxin, L-AAA, led to complete loss of arteriolar dilation in response to neuronal activation.¹⁰¹

Glial Control of Glutamate and Excitotoxicity

Astrocytes play a fundamental role in the pathogenesis of ischemic neuronal death.¹⁰² A large body of evidence indicates that astrocytes are involved in the control of glutamate homeostasis and susceptibility of the brain to excitotoxic injury.¹⁰³ Glutamate transporters are expressed in many different types of brain cells, but astrocytes are primarily responsible for glutamate uptake. Studies using genetic deletion or antisense oligonucleotides-mediated knockdown of the astroglial glutamate transporter GLT-1 have demonstrated that this transporter is the predominant subtype responsible for the clearance of extracellular glutamate in the brain.^{104,105} Affected animals were highly susceptible to glutamate-dependent excitotoxicity and developed epileptic seizures.^{104,105} After uptake of glutamate into astrocytes, the enzyme glutamine synthetase converts glutamate into glutamine, which is then transported into neurons where it is converted back into glutamate. Interestingly, a loss of glutamine synthetase was found in the sclerotic hippocampus of human patients with temporal lobe epilepsy¹⁰⁶ and the

authors of that study concluded that reduced activity of the glutamate–glutamine cycle led to accumulation of extracellular glutamate.

Apart from the mechanisms described above, astrocytes themselves can be a significant source of extracellular glutamate, which can be released by a variety of mechanisms.¹⁰⁷ It has been demonstrated that Ca^{2+} elevations in astrocytes induce the excitotoxic release of glutamate from these cells.¹⁰⁸ Most importantly, the Ca^{2+} -dependent astrocytic release of glutamate was also dependent on the vesicular glutamate transporters (VGLUT1/2) and the vesicular SNARE protein, cellubrevin, and was consistent with a vesicular release mechanism of glutamate that was similar to synaptic release of glutamate.¹⁰⁸ Finally, it was shown that astrocyte-derived glutamate targets synaptic *N*-methyl-D-aspartic acid (NMDA) receptors,¹⁰⁹ providing a rational explanation for the astrocyte-based control of neurotoxicity. Given the emerging roles of astrocytes in the control of neuronal excitotoxicity, neuroprotective efforts targeting the functional integrity of astrocytes may constitute a superior strategy for future neuroprotection.

Novel Methods for Studying the Role of Glia

Co-culture of glia and neurons. Many original insights into gliotransmission were discovered using co-culture systems and then extended to brain slices or the intact brain. For example, the first evidence for gliotransmission came from studies of mixed cultures of astrocytes and neurons demonstrating that experimentally evoked Ca^{2+} elevation in astrocytes evoked elevation of Ca^{2+} in adjacent neurons.^{110,111} In this method, cultured cells from postnatal day 1–4 rodent brain are first enriched with one population of either glial or neuronal origin using specific culture conditions and then confirmed by immunohistochemistry. Glial or neuronal cells are then plated separately or together onto a coated substrate. The distinct morphology of glial cells and neurons allows identification of these distinct cell types and permits direct application of electric field potentials, micropipette tips, or neurochemical substrates to astrocytes in order to evoke a specific elevation of calcium in glial cells (e.g., in astrocytes).^{110,111} The spatiotemporal control over mechanical stimuli has permitted the selective stimulation of single astrocytes in mixed cultures of rat forebrain astrocytes and neurons. Using this strategy it was demonstrated that the elevation of calcium, triggered by focal electric field potentials in single astrocytes, induced a wave of calcium increase that was propagated from astrocyte to astrocyte, and importantly, this wave of calcium increase also evoked large increases in the concentration of cytosolic calcium in neurons depending on those astrocytes.^{110,111} More recently, mixed co-cultures in multi-compartment dishes were equipped with the capability to provide electronic stimuli selectively to neurons in mixed cultures of oligodendrocytes and dorsal root ganglion neurons. This approach led to the finding that ATP and adenosine, released from neurons, act as potent neuron–glial transmitters that inhibited oligodendrocyte progenitor cell proliferation, stimulated their differentiation, and promoted the formation of myelin.^{56,112}

Transgenic overexpression and targeted knockout of genes in defined glial populations. Glial cells have been shown to release gliotransmitters, including ATP,^{9,17} glutamate,^{111,113} and D-serine,¹¹⁴ to coordinate synaptic networks. However, neurons and glia share these same chemical signaling molecules, making it difficult to define the role of gliotransmitters. To molecularly dissect the role of glial signaling molecules, genetic approaches have been developed to selectively manipulate the SNARE-dependent release of gliotransmitters using a glia-specific promoter. Pascual and Haydon developed a transgenic mouse line, which uses the tet-off system to allow conditional expression of the cytosolic portion of the SNARE domain of synptobrevin-2 [dominant-negative SNARE (dn-SNARE)] selectively in astrocytes.¹⁵ The selective expression of dn-SNARE was achieved by the use of an astrocyte-specific glial fibrillary acid protein promoter. To confirm the cell type selectivity of astrocytic transgene expression, Pascual and co-workers used enhanced green fluorescent protein (EGFP) as a reporter system and showed that EGFP was visually detectable in 97% of the dn-SNARE-transgene-expressing cultured astrocytes, and that EGFP-positive cells colocalized specifically with the astrocytic marker, but not with neuronal, NG2–glial, or oligodendroglial markers. This transgenic line has been used successfully to demonstrate the functional significance of gliotransmission on synaptic plasticity in hippocampus,¹⁵ and more recently in the sleep–wake cycle.⁶¹ Similarly, transgenic overexpression of a mutant disease-causing gene in a defined glial cell population has been used to study the role of superoxide dismutase-1 (SOD1) in astrocytes in the development of motor neuron death. Nagai *et al.*¹¹⁵ demonstrated that the astrocyte-selective expression of mutant human SOD1 (but not in spinal motor neurons, microglia or fibroblasts) killed spinal primary and embryonic mouse stem cell-derived motor neurons.

As a complementary genetic approach, the selective deletion of signaling molecules in defined glial cell populations can be accomplished using the Cre-loxP strategy. For example, Boillee *et al.*¹¹⁶ generated a transgenic line (LoxSOD1^{G37R}) that carried a mutant human SOD1 gene flanked at both ends by a 34-bp LoxP sequence. These 'floxed' mice were then cross-bred to two transgenic lines with expression of the Cre protein under control of (i) the promoter from the *Islet-1* transcription factor, which directs the expression exclusively in progenitors of motor and dorsal root ganglion neurons, and (ii) the *CD11b* promoter, which directs the expression exclusively in the myeloid lineage (including macrophages and microglial cells). The establishment of these two novel transgenic lines allowed to demonstrate that the SOD1 mutation in motor neurons and microglial cells contributes distinctly to the onset and progression of amyotrophic lateral sclerosis: whereas expression of SOD1 in motor neurons is the primary signal for the initiation of motor neurodegeneration and an early sign of disease progression, the genetic inactivation of SOD in microglial cells had little effect on the early disease phase, but markedly attenuated disease progression.¹¹⁶ Thus, these genetic approaches to selectively manipulate signaling molecules in defined glial (or neuronal) populations provide

critical insights into the distinct role of glial cells in the development of neurodegeneration.

Flow cytometric analysis and FACS. The structural complexity of brain tissues hampers the dissection of unique roles of glial cells under various physiological and pathological conditions. Glial cells are characterized by a unique morphology, which permits distinction of neuronal versus glial cells in intact brain by immunohistochemistry. However, it is difficult to quantify immunohistochemical changes without performing labor-intensive stereological analysis. Furthermore, there is a critical need to isolate large numbers of pure glial cells from intact brain for detailed molecular analysis such as qPCR and microarray analysis. Flow cytometric and fluorescence-activated cell sorting (FACS) can be adapted to partially circumvent these limitations for the study of glial cell functions in brain.

In the first application, defined glial cell populations (such as astrocytes, microglia, or oligodendrocytes) are identified by labeling with a fluorescent antibody directed against specific cell-surface markers. Quantitative changes of glial populations in normal and injured brains are determined by flow cytometry. For example, we recently used this analysis to evaluate the change of CD11b+ (a cell-surface marker for microglial cells) and GFAP+ (a marker for astrocytes) cells in mouse striatum after treatment with MPTP and the A_{2A}R antagonist KW6002.¹¹⁷ This analysis not only provided an improved quantitative assessment of the effect of the A_{2A}R antagonist on microglial activation at the very early phase of MPTP intoxication, but also identified a specific microglial cell population (i.e., CD11b+ cells with a large cell size representing fully activated microglial cells), which are most sensitive to KW6002 treatment in the brain.¹¹⁷

In the second application, FACS permits isolation and purification of distinct populations of glial cells from neurons from brain tissues using fluorescent antibodies directed against cell-surface markers. The sorted glial cell populations can then be used for detailed molecular analyses such as quantitative PCR and microarray analysis. For example, Lovatt *et al.*¹¹⁸ successfully performed microarray profiling of sorted astrocytes from mouse cortex using FACS and (surprisingly) demonstrated that most enzymes in the tricarboxylic acid cycle are expressed at higher relative levels in astrocytes than in neurons.

Adenosine Signaling in Glial Cells, Excitotoxicity, and Cell Death

As outlined above, synaptic levels of adenosine are largely controlled by an astrocyte-based adenosine cycle and the activity of the astrocyte-based enzyme ADK. Consequently, adenosine signaling in glial cells effects excitotoxicity and cell death in a variety of experimental paradigms. In addition, several forms of brain insult activate microglial cells. Here, ATP release is critically important,⁸⁵ but the ATP response in microglial cells is markedly enhanced by adenosine generated from ATP.²⁷

Epilepsy. The adenosine kinase hypothesis of epileptogenesis implies that dysregulation of ADK is a major

contributing factor to the epileptogenic cascade¹² (Figure 3). Consequently, ADK expression levels (that determine levels of ambient adenosine) determine the brain's susceptibility to acute seizure-induced cell death. Mice with only moderate transgenic overexpression of ADK in the brain (141% of normal) were highly susceptible to acute seizure-induced cell death and did not survive beyond 3 days following status epilepticus,³⁷ whereas engineered mice with reduced levels of ADK in the forebrain (62% of normal) were completely resistant to seizure-induced cell death.³⁷ Resistance to seizure-induced excitotoxic cell death in ADK-deficient mice was dependent on adenosine and increased adenosine A₁R activation, since blockade of A₁Rs with its selective antagonist 8-cyclopentyl-1,3-dipropylxanthine restored wild-type-like seizure-induced excitotoxic cell death.³⁷

Astrogliosis is a pathological hallmark of the epileptic brain and contributes to seizure generation by a variety of mechanisms.^{12,119,120} A recent study in our laboratory has identified the enzyme ADK in astrocytes as a molecular link between astrogliosis and neuronal dysfunction in epilepsy.³⁷ In a mouse model of CA3-selective epileptogenesis, we found spatio-temporal colocalization of astrogliosis, upregulated ADK, and focal spontaneous electrographic seizures that were all restricted to the CA3 region, the site of the epileptogenesis precipitating acute injury; importantly, seizures could be suppressed pharmacologically by ADK inhibition.³⁷ In this model, the seizures remained highly localized and restricted to the astrogliotic scar, presumably due to normal adenosinergic control of the surrounding brain tissue. Transgenic overexpression of ADK, as well as genetic disruption of the A₁R, was sufficient to trigger spontaneous seizures, indicating that adenosine dysfunction rather than astrogliosis *per se* was responsible for seizure generation.¹²¹




Figure 3 Role of the adenosine (ADO)/ ADK system in regulating acute and chronic responses to injury. Left: Within hours after brain injury (e.g., stroke, trauma, prolonged seizures) a surge in micromolar levels of ADO results that protects the brain from further injury and from seizures. Hypoxia and trauma can directly lead to a rise in extracellular ATP that is rapidly degraded into adenosine. High levels of adenosine are known to inhibit ADK, further amplifying the adenosine surge. Right: The acute adenosine surge contributes to trigger astrogliosis via a variety of mechanisms that include modulation of astrocytic ARs, modulation of inflammatory processes, and release of cytokines. Astrogliosis leads to overexpression of ADK, resulting in adenosine deficiency, which contributes to seizure generation

Conversely, mice with genetically induced reduction of ADK in forebrain were completely resistant to the development of spontaneous seizures.³⁷ *In vitro* studies performed on hippocampal slices have subsequently demonstrated that reduction of the basal tone of adenosine by ADK is permissive to seizure generation, whereas ADK did not limit activity-dependant adenosine release.¹²² Together, these findings provide a neurochemical rationale for adenosine augmentation therapies (AATs). Consequently, several focal AAT-approaches, based on intracerebral adenosine-releasing implants, have demonstrated robust anticonvulsive and, possibly, antiepileptogenic efficacy in a variety of experimental paradigms that have been reviewed elsewhere.^{123,124}

Traumatic brain injury. Traumatic brain injury triggers an acute surge in adenosine, presumably as a consequence of ATP release, and this may represent an endogenous neuroprotective mechanism. In one study, adenosine levels increased 61-fold following controlled cortical impact (CCI) in rats and peaked at 20 min following the impact.¹²⁵ The existence of an endogenous protective action of adenosine at A_1 receptors early after experimental TBI was further corroborated by the finding of lethal status epilepticus in A_1 R-knockout mice subjected to either controlled cortical impact.¹²⁶ or to kainic acid-induced hippocampal injury.¹²⁷ In contrast to A_1 R-knockout mice, A_{2A} R-knockout mice were largely protected from the adverse effects of CCI.¹²⁸ In line with these findings, increases in cerebrospinal fluid caffeine concentration were associated with favorable outcome after severe traumatic brain injury in humans, likely due to caffeine-mediated inhibition of A_{2A} Rs.¹²⁹ Likewise, chronic, but not acute, caffeine attenuated the consequences of TBI in the mouse CCI model.¹³⁰ Although it probably does not contribute to the surge in adenosine following brain injury, acute downregulation of ADK in astrocytes has been described as a consequence of stroke³⁶ or acute seizures,¹³¹ and this may prolong the adenosine increase.

Parkinson's disease. The adenosine A_{2A} receptor is a leading non-dopaminergic therapeutic target in Parkinson's disease (PD) (Figure 4). Interest in this receptor within the context of PD derives primarily from two lines of experimental

and clinical investigations: first, decade-long preclinical studies demonstrated a unique colocalization of A_{2A} Rs and dopamine D_2 Rs in striatopallidal neurons. Antagonistic interactions between A_{2A} Rs and D_2 Rs at the molecular, neurochemical, and behavioral level explain the motor-stimulant effects of A_{2A} R blockade.^{132–134} Thus, A_{2A} R antagonists, such as KW-6002 (istradefylline) and SCH420814, have now completed clinical phase IIB–III trials. Despite some limitations of these clinical trials and admittedly modest effects ('OFF' time reduced by one hour), these studies confirm that selective A_{2A} R antagonists can stimulate motor activity by potentiating the L-dopa effect in advanced PD patients.^{135,136} Second, in addition to symptomatic relief, A_{2A} R antagonists appear to more directly attenuate dopaminergic neurodegeneration, as suggested by convergent epidemiological and experimental evidence. Following an initial report from the Honolulu Heart Program¹³⁷ by Ross and co-workers, several large-cohort, prospective studies have confirmed a similar inverse relationship between the consumption of caffeinated coffee and the risk of developing PD. Including the Health Professionals' Follow-Up Study and the Nurses' Health Study these studies involved a total of 47 351 men and 88 565 women,¹³⁸ whereas a more recent study conducted by the Finnish Mobile Clinic Health Examination Survey included 19 518 men and women.¹³⁹ These studies firmly established a relationship between increased caffeine consumption and decreased risk of developing PD in males. In addition, studies with animal models of PD provide a compelling clue about the potentially protective effects of caffeine, by demonstrating that pharmacological blockade (by caffeine or selective A_{2A} R antagonists) or genetic depletion of the A_{2A} R attenuates dopaminergic neurotoxicity and neurodegeneration.^{140–142} These studies provide a neurobiological basis for the inverse relationship between increased caffeine consumption and reduced risk of developing PD.

Despite consistent demonstration that A_{2A} R antagonists afford neuroprotection against MPTP or 6-hydroxydopamine-induced dopaminergic neurotoxicity, the mechanism by which A_{2A} R inactivation protects against the loss of dopaminergic neurons remains unknown. The particular challenge lies in




Figure 4 The dual functions of A_{2A} R antagonists in PD models: A_{2A} R antagonists act at the A_{2A} R in striatal neurons to stimulate motor activity. Furthermore, it is postulated that A_{2A} R antagonists may modulate microglial activation in substantia nigra to exert a possible neuroprotective effect in an animal model of PD

explaining the apparent dichotomy between restricted expression of the $A_{2A}R$ in striatopallidal neurons and neuroprotection against degeneration of dopaminergic neurons in the substantia nigra where only a scattered expression of $A_{2A}Rs$ is detected. An additional challenge is to identify the cellular mechanism, which allows $A_{2A}R$ inactivation to protect neurons against a broad spectrum of brain insults, from ischemia to excitotoxicity to mitochondrial toxicity.¹⁴³ In this context, the involvement of the glial $A_{2A}R$ becomes an attractive possibility, since glial function and neuroinflammation is commonly associated with diverse pathological conditions as mentioned above. Indeed, we recently demonstrated that MPTP treatment markedly upregulates $A_{2A}R$ expression in microglial cells at 24–48 h after treatment,^{65,66} which may result in further amplification of the $A_{2A}R$ -mediated modulation of neuroinflammation in PD models. Consistent with this notion, an immunohistochemical study showed that KW-6002 reduced the loss of striatal dopamine contents and nigral cell bodies, and this coincided with inhibition of microglial activation.¹⁴⁴ Furthermore, we demonstrated by flow cytometry that KW-6002 attenuated MPTP-induced microglial activation at 48 h after MPTP treatment.¹¹⁷ In conclusion, $A_{2A}R$ antagonists may confer neuroprotection by acting at $A_{2A}Rs$ in glial cells, at least in the MPTP model of PD.

Ischemia. During ischemia an imbalance between ATP degradation and resynthesis brings about a rapid and marked increase in extracellular level of adenosine in the brain during ischemia.^{145–154} In addition hypoxia will increase ATP release, resulting in further adenosine production. Although no clinical reports with purinergic compounds in human stroke exist, it is widely believed that adenosine and its receptors function as an endogenous neuroprotectant under these conditions.^{155–161} Indeed, adenosine¹⁶² or adenosine-potentiating agents (such as inhibitors of ADA or ADK,^{163–167} or of adenosine transport^{151,166,168–173}) offer protection against ischemic neuronal damage in different *in vivo* ischemia models. Furthermore, transgenic overexpression of ADK aggravates cell death, whereas reduction of ADK level in the hippocampus increases protection after transient focal ischemia.^{36,174}

Such a protective effect is attributed to stimulation of adenosine A_1 receptors that exert a protective role in ischemia by presynaptic reduction of Ca^{2+} influx, by inhibition of the release of excitatory neurotransmitters,^{175,176} and by postsynaptic hyper-polarization and reduction of neuronal activity through increases in K^+ and Cl^- ion conductances.¹⁷⁷ The efficacy of A_1 receptor stimulation on neuroprotection depends on the model used and no protective effect was observed in a global ischemia model.¹⁷⁸ Since adenosine does influence glutamate release, it is suggested that this is not critically important in some ischemic models. However, adenosine (probably acting at the A_{2A} receptor) may in fact contribute to neurotoxicity, neuronal damage, and cell death. The potential neuroprotection by $A_{2A}R$ antagonists was first reported in a global ischemia model with the less selective antagonist CGS 15943.^{179,180} Further studies substantiated this finding in different models of ischemia with the selective A_{2A} receptor antagonist 8-(3-chlorostyryl)caffeine (CSC)

and SCH 58261^{181,147,182} in various animal models of stroke.^{183,184} Studies of genetically manipulated mice confirmed the neuroprotective role of A_{2A} receptor antagonists on ischemic brain damage.¹⁸⁵ Major protective effects of A_{2A} receptor antagonists in stroke have been attributed to reduced glutamate outflow.^{183,186,187} It should be considered however, that in several studies, A_{2A} receptor agonists have been found protective in the global ischemia model in the gerbil¹⁸⁰ and that A_{2A} -knockout mice show aggravated hypoxic ischemic injury in neonatal mice.¹⁸⁸ Possible mechanisms are not clear yet, but include $A_{2A}R$ -mediated protection via inhibition of platelet aggregation, vasodilation,^{163,189} or anti-inflammatory actions. Lastly, activation of A_3Rs produced mixed results and the exact contribution of A_3Rs to ischemic brain injury is not clear.¹⁹⁰

So far, support for a role for glia in the neuroprotective effect of $A_{2A}R$ antagonism in ischemia comes from the observation that the A_{2A} receptor antagonist SCH 58261 reduces p38 MAPK activation in microglial cells¹⁸⁴ and phospho-JNK in neurons and oligodendrocytes¹⁹¹ in the ischemic hemisphere 24 h after permanent MCAO. Since p38 MAPK and JNK are activated up to 24 h after ischemia^{192,193} and are involved in neuronal death,^{194,195} this correlation indicates that $A_{2A}R$ antagonists may confer neuroprotection against ischemic brain injury through modulation of glial function and neuroinflammation. However, reduced MAPK activation might be secondary to a reduction in the excitotoxic cascade that primes p38 and JNK activation, since reduction glutamate outflow in the ischemic brain by $A_{2A}R$ blockade is believed to be one of the main underlying mechanisms.¹⁹⁶ Further studies with selective manipulation of glial ARs or glial function *in vivo* are critical to our understanding to what extent adenosine regulation of glial signaling and function is responsible for ischemic brain injury.

Conclusions and Major Open Questions

The discussion above has shown that much is now known about a role of glial cells in mediating the effects of adenosine (and other purines) in different neurodegenerative states. It is clear that many of the actions of endogenous or exogenous adenosine (and ATP) are in fact due to actions on glial cells. This makes it much more complicated to understand precisely how adenosine acts, and it has become apparent that the often diverse actions reported are due to the fact that several different receptors, located on many cell types, are involved. In order to get a better understanding, there are several major questions that require an answer. Among the questions that need to be addressed in future research we find:

1. Is the acute adenosine surge that follows brain injury a trigger for subsequent astrocyte activation?
2. Do all the proposed mechanisms of ATP release from glial cells in fact occur, and are the triggers for astrocytic ATP release via these mechanisms different?
3. To what extent does regional variation in the expression of ecto-nucleotidases control the distribution of adenosine in brain?
4. Can adenosine signals propagate within the brain via astrocyte–astrocyte communication?

5. Given the major increase in glial cells in humans, how far can we extrapolate from rodents to man?
6. When several ARs, with at least partly opposing signaling, appear to regulate a single biological response, are the receptors located on different cells or on very different parts of the same cell?
7. Given that two GPCR molecules can form dimers, but apparently only one of them can actually signal, is there any functional significance of heterodimers between two types of purine receptors?
8. Can activation of endothelial cells at the vascular interphase signal to synapses?
9. Are microglial cells directly involved in synaptic transmission or do they mainly help prune synaptic contacts?
10. Why are effects of A2ARs on inflammation different in CNS and peripheral organs?

Acknowledgements. This work was supported by NIH Grants NS058780, MH083973, NS061844, NS057538, and NS057475, and by the CURE Foundation in collaboration with the Department of Defense (to DB), by NIH Grants NS48995, NS41083, and DA19362, and a grant from the Department of Defense W81XWH-071-1-0012 (to JFC), and by grants from the Swedish Science Research Council and Hjärfonden (to BBF).

1. Zipp F, Aktas O. The brain as a target of inflammation: common pathways link inflammatory and neurodegenerative diseases. *Trends Neurosci* 2006; **29**: 518–527.
2. Geiger JD, Buscemi L, Fotheringham JA. Role of adenosine in the control of inflammatory events associated with acute and chronic neurodegenerative disorders In: Hasko G, Cronstein BN, Szabo C (eds). *Adenosine Receptors. Therapeutic Aspects for Inflammatory and Immune Diseases*. Taylor and Francis: CRC, 2007 pp 213–236.
3. Di Virgilio F, Ceruti S, Bramanti P, Abbracchio MP. Purinergic signalling in inflammation of the central nervous system. *Trends Neurosci* 2009; **32**: 79–87.
4. Zimmermann H. Extracellular metabolism of ATP and other nucleotides. *Naunyn-Schmiedeberg's Arch Pharmacol* 2000; **362**: 299–309.
5. Boison D. Adenosine kinase, epilepsy and stroke: mechanisms and therapies. *Trends Pharmacol Sci* 2006; **27**: 652–658.
6. Martin ED, Fernandez M, Perea G, Pascual O, Haydon PG, Araque A *et al*. Adenosine released by astrocytes contributes to hypoxia-induced modulation of synaptic transmission. *Glia* 2007; **55**: 36–45.
7. Halassa MM, Fellin T, Haydon PG. The tripartite synapse: roles for gliotransmission in health and disease. *Trends Mol Med* 2007; **13**: 54–63.
8. Halassa MM, Fellin T, Takano H, Dong JH, Haydon PG. Synaptic islands defined by the territory of a single astrocyte. *J Neurosci* 2007; **27**: 6473–6477.
9. Haydon PG, Carmignoto G. Astrocyte control of synaptic transmission and neurovascular coupling. *Physiol Rev* 2006; **86**: 1009–1031.
10. Studer FE, Fedele DE, Marowsky A, Schwerdel C, Wernli K, Vogt K *et al*. Shift of adenosine kinase expression from neurons to astrocytes during postnatal development suggests dual functionality of the enzyme. *Neuroscience* 2006; **142**: 125–137.
11. Boison D. Adenosine as a neuromodulator in neurological diseases. *Curr Opin Pharmacol* 2008; **8**: 2–7.
12. Boison D. The adenosine kinase hypothesis of epileptogenesis. *Prog Neurobiol* 2008; **84**: 249–262.
13. Abbracchio MP, Burnstock G, Verkhratsky A, Zimmermann H. Purinergic signalling in the nervous system: an overview. *Trends Neurosci* 2009; **32**: 19–29.
14. MacDonald PE, Braun M, Galvanovskis J, Rorsman P. Release of small transmitters through kiss-and-run fusion pores in rat pancreatic beta cells. *Cell Metab* 2006; **4**: 283–290.
15. Pascual O, Casper KB, Kubera C, Zhang J, Revilla-Sanchez R, Sul JY *et al*. Astrocytic purinergic signaling coordinates synaptic networks. *Science* 2005; **310**: 113–116.
16. Chen X, Wang L, Zhou Y, Zheng LH, Zhou Z. 'Kiss-and-run' glutamate secretion in cultured and freshly isolated rat hippocampal astrocytes. *J Neurosci* 2005; **25**: 9236–9243.
17. Zhang Z, Chen G, Zhou W, Song A, Xu T, Luo Q *et al*. Regulated ATP release from astrocytes through lysosome exocytosis. *Nat Cell Biol* 2007; **9**: 945–953.
18. Kang J, Kang N, Lovatt D, Torres A, Zhao Z, Lin J *et al*. Connexin 43 hemichannels are permeable to ATP. *J Neurosci* 2008; **28**: 4702–4711.
19. Iglesias R, Dahl G, Qiu F, Spray DC, Scemes E. Pannexin 1: the molecular substrate of astrocyte 'hemichannels'. *J Neurosci* 2009; **29**: 7092–7097.
20. Liu HT, Sabirov RZ, Okada Y. Oxygen–glucose deprivation induces ATP release via maxi-anion channels in astrocytes. *Purinergic Signal* 2008; **4**: 147–154.
21. Cocucci E, Racchetti G, Meldolesi J. Shedding microvesicles: artefacts no more. *Trends Cell Biol* 2009; **19**: 43–51.
22. Bjorklund O, Shang MM, Tonazzini I, Dare E, Fredholm BB. Adenosine A(1) and A(3) receptors protect astrocytes from hypoxic damage. *Eur J Pharmacol* 2008; **596**: 6–13.
23. Parkinson FE, Sinclair CJ, Othman T, Haughey NJ, Geiger JD. Differences between rat primary cortical neurons and astrocytes in purine release evoked by ischemic conditions. *Neuropharmacology* 2002; **43**: 836–846.
24. Ventura R, Harris KM. Three-dimensional relationships between hippocampal synapses and astrocytes. *J Neurosci* 1999; **19**: 6897–6906.
25. Newman EA. Glial cell inhibition of neurons by release of ATP. *J Neurosci* 2003; **23**: 1659–1666.
26. Kreutzberg GW, Barron KD. 5'-Nucleotidase of microglial cells in the facial nucleus during axonal reaction. *J Neurocytol* 1978; **7**: 601–610.
27. Farber K, Markworth S, Pannasch U, Nolte C, Prinz V, Kronenberg G *et al*. The ectonucleotidase cd39/ENTPDase1 modulates purinergic-mediated microglial migration. *Glia* 2008; **56**: 331–341.
28. Kowaluk EA, Jarvis MF. Therapeutic potential of adenosine kinase inhibitors. *Expert Opin Invest Drugs* 2000; **9**: 551–564.
29. Ren G, Li T, Lan JQ, Wilz A, Simon RP, Boison D. Lentiviral RNAi-induced downregulation of adenosine kinase in human mesenchymal stem cell grafts: a novel perspective for seizure control. *Exp Neurol* 2007; **208**: 26–37.
30. Güttinger M, Padrun V, Pralong W, Boison D. Seizure suppression and lack of adenosine A₁ receptor desensitization after focal long-term delivery of adenosine by encapsulated myoblasts. *Exp Neurol* 2005; **193**: 53–64.
31. Fedele DE, Koch P, Brüstle O, Scheurer L, Simpson EM, Mohler H *et al*. Engineering embryonic stem cell derived glia for adenosine delivery. *Neurosci Lett* 2004; **370**: 160–165.
32. Huber A, Padrun V, Deglon N, Aebischer P, Mohler H, Boison D. Grafts of adenosine-releasing cells suppress seizures in kindling epilepsy. *Proc Natl Acad Sci USA* 2001; **98**: 7611–7616.
33. Fedele DE, Gouder N, Güttinger M, Gabernet L, Scheurer L, Rulicke T *et al*. Astroglial gliosis in epilepsy leads to overexpression of adenosine kinase resulting in seizure aggravation. *Brain* 2005; **128**: 2383–2395.
34. Pak MA, Haas HL, Decking UKM, Schrader J. Inhibition of adenosine kinase increases endogenous adenosine and depresses neuronal activity in hippocampal slices. *Neuropharmacology* 1994; **33**: 1049–1053.
35. Arch JR, Newsholme EA. Activities and some properties of 5'-nucleotidase, adenosine kinase and adenosine deaminase in tissues from vertebrates and invertebrates in relation to the control of the concentration and the physiological role of adenosine. *Biochem J* 1978; **174**: 965–977.
36. Pignataro G, Maysami S, Studer FE, Wilz A, Simon RP, Boison D. Downregulation of hippocampal adenosine kinase after focal ischemia as potential endogenous neuroprotective mechanism. *J Cereb Blood Flow Metab* 2008; **28**: 17–23.
37. Li T, Ren G, Lusardi T, Wilz A, Lan JQ, Iwasato T *et al*. Adenosine kinase is a target for the prediction and prevention of epileptogenesis in mice. *J Clin Invest* 2008; **118**: 571–582.
38. Baldwin SA, Beal PR, Yao SY, King AE, Cass CE, Young JD. The equilibrative nucleoside transporter family, SLC29. *Pharmacol Ther* 2004; **447**: 735–743.
39. Boison D, Scheurer L, Zumsteg V, Rulicke T, Litynski P, Fowler B *et al*. Neonatal hepatic steatosis by disruption of the adenosine kinase gene. *Proc Natl Acad Sci USA* 2002; **99**: 6985–6990.
40. Peng L, Huang R, Yu AC, Fung KY, Rathbone MP, Hertz L. Nucleoside transporter expression and function in cultured mouse astrocytes. *Glia* 2005; **52**: 25–35.
41. King AE, Ackley MA, Cass CE, Young JD, Baldwin SA. Nucleoside transporters: from scavengers to novel therapeutic targets. *Trends Pharmacol Sci* 2006; **27**: 416–425.
42. Carrier EJ, Auchampach JA, Hillard CJ. Inhibition of an equilibrative nucleoside transporter by cannabidiol: A mechanism of cannabinoid immunosuppression. *Proc Natl Acad Sci USA* 2006; **103**: 7895–7900.
43. Fredholm BB, Ijzerman AP, Jacobson KA, Klotz KN, Linden J. International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol Rev* 2001; **53**: 527–552.
44. Jin X, Shepherd RK, Duling BR, Linden J. Inosine binds to A3 adenosine receptors and stimulates mast cell degranulation. *J Clin Invest* 1997; **100**: 2849–2857.
45. Fredholm BB, Irenius E, Kull B, Schulte G. Comparison of the potency of adenosine as an agonist at human adenosine receptors expressed in Chinese hamster ovary cells. *Biochem Pharmacol* 2001; **61**: 443–448.
46. Schulte G, Fredholm BB. Human adenosine A(1), A(2A), A(2B), and A(3) receptors expressed in Chinese hamster ovary cells all mediate the phosphorylation of extracellular-regulated kinase 1/2. *Mol Pharmacol* 2000; **58**: 477–482.
47. Hammarberg C, Schulte G, Fredholm BB. Evidence for functional adenosine A3 receptors in microglia cells. *J Neurochem* 2003; **86**: 1051–1054.
48. van Calker D, Biber K. The role of glial adenosine receptors in neural resilience and the neurobiology of mood disorders. *Neurochem Res* 2005; **30**: 1205–1217.
49. Dare E, Schulte G, Karovic O, Hammarberg C, Fredholm BB. Modulation of glial cell functions by adenosine receptors. *Physiol Behav* 2007; **92**: 15–20.

50. Ciccarelli R, Di Iorio P, Ballerini P, Ambrosini G, Giuliani P, Tiboni GM *et al*. Effects of exogenous ATP and related analogues on the proliferation rate of dissociated primary cultures of rat astrocytes. *J Neurosci Res* 1994; **39**: 556–566.
51. Peakman MC, Hill SJ. Adenosine A1 receptor-mediated changes in basal and histamine-stimulated levels of intracellular calcium in primary rat astrocytes. *Br J Pharmacol* 1995; **115**: 801–810.
52. Biber K, Klotz KN, Berger M, Gebicke-Harter PJ, van Calker D. Adenosine A1 receptor-mediated activation of phospholipase C in cultured astrocytes depends on the level of receptor expression. *J Neurosci* 1997; **17**: 4956–4964.
53. Cormier RJ, Mennerick S, Melbostad H, Zorumski CF. Basal levels of adenosine modulate mGluR5 on rat hippocampal astrocytes. *Glia* 2001; **33**: 24–35.
54. Ciccarelli R, D'Alimonte I, Ballerini P, D'Auro M, Nargi E, Buccella S *et al*. 2007 Molecular signalling mediating the protective effect of A1 adenosine and mGlu3 metabotropic glutamate receptor activation against apoptosis by oxygen/glucose deprivation in cultured astrocytes. *Mol Pharmacol* 2007; **71**: 1369–1380.
55. D'Alimonte I, Ballerini P, Nargi E, Buccella S, Giuliani P, Di Iorio P *et al*. Staurosporine-induced apoptosis in astrocytes is prevented by A1 adenosine receptor activation. *Neurosci Lett* 2007; **418**: 66–71.
56. Ishibashi T, Dakin KA, Stevens B, Lee PR, Kozlov SV, Stewart CL *et al*. Astrocytes promote myelination in response to electrical impulses. *Neuron* 2006; **49**: 823–832.
57. Othman T, Yan H, Rivkees SA. Oligodendrocytes express functional A1 adenosine receptors that stimulate cellular migration. *Glia* 2003; **44**: 166–172.
58. Kim M, Yu ZX, Fredholm BB, Rivkees SA. Susceptibility of the developing brain to acute hypoglycemia involving A1 adenosine receptor activation. *Am J Physiol Endocrinol Metab* 2005; **289**: E562–E569.
59. Tsutsui S, Schnermann J, Noorbakhs F, Henry S, Yong VW, Winston BW *et al*. A1 adenosine receptor upregulation and activation attenuates neuroinflammation and demyelination in a model of multiple sclerosis. *J Neurosci* 2004; **24**: 1521–1529.
60. Synowitz M, Glass R, Farber K, Markovic D, Kronenberg G, Herrmann K *et al*. A1 adenosine receptors in microglia control glioblastoma–host interaction. *Cancer Res* 2006; **66**: 8550–8557.
61. Halassa MM, Florian C, Fellin T, Munoz JR, Lee SY, Abel T *et al*. Astrocytic modulation of sleep homeostasis and cognitive consequences of sleep loss. *Neuron* 2009; **61**: 213–219.
62. Svenningsson P, Le Moine C, Fisone G, Fredholm BB. Distribution, biochemistry and function of striatal adenosine A2A receptors. *Prog Neurobiol* 1999; **59**: 355–396.
63. Hettlinger BD, Lee A, Linden J, Rosin DL. Ultrastructural localization of adenosine A2A receptors suggests multiple cellular sites for modulation of GABAergic neurons in rat striatum. *J Comp Neurol* 2001; **431**: 331–346.
64. Hasko G, Pacher P, Vizi ES, Illes P. Adenosine receptor signaling in the brain immune system. *Trends Pharmacol Sci* 2005; **26**: 511–516.
65. Saura J, Angulo E, Ejarque A, Casado V, Tusell JM, Moratalla R *et al*. Adenosine A2A receptor stimulation potentiates nitric oxide release by activated microglia. *J Neurochem* 2005; **95**: 919–929.
66. Yu L-Q, Shen H-Y, Coelho J, Araújo I, Huang Q-Y, Day Y-J *et al*. A2A receptor antagonists exert motor and protective effects by distinct cellular mechanism in MPTP model of PD. *Ann Neurol* 2008; **63**: 338–346.
67. Pickel VM, Chan J, Linden J, Rosin DL. Subcellular distributions of adenosine A1 and A2A receptors in the rat dorsomedial nucleus of the solitary tract at the level of the area postrema. *Synapse* 2006; **60**: 496–509.
68. Cunha RA. Neuroprotection by adenosine in the brain: From A1 receptor activation to A2A receptor blockade. *Purinergic Signal* 2005; **1**: 111–134.
69. Li XX, Nomura T, Aihara H, Nishizaki T. Adenosine enhances glial glutamate efflux via A2a adenosine receptors. *Life Sci* 2001; **68**: 1343–1350.
70. Nishizaki T, Nagai K, Nomura T, Tada H, Kanno T, Tozaki H *et al*. A new neuromodulatory pathway with a glial contribution mediated via A(2a) adenosine receptors. *Glia* 2002; **39**: 133–147.
71. Hindley S, Herman MA, Rathbone MP. Stimulation of reactive astrogliosis *in vivo* by extracellular adenosine diphosphate or an adenosine A2 receptor agonist. *J Neurosci Res* 1994; **38**: 399–406.
72. Brambilla R, Cottini L, Fumagalli M, Ceruti S, Abbracchio MP. Blockade of A2A adenosine receptors prevents basic fibroblast growth factor-induced reactive astrogliosis in rat striatal primary astrocytes. *Glia* 2003; **43**: 190–194.
73. Brodie C, Blumberg PM, Jacobson KA. Activation of the A2A adenosine receptor inhibits nitric oxide production in glial cells. *FEBS Lett* 1998; **429**: 139–142.
74. Fiebich BL, Biber K, Lieb K, van Calker D, Berger M, Bauer J *et al*. Cyclooxygenase-2 expression in rat microglia is induced by adenosine A2a-receptors. *Glia* 1996; **18**: 152–160.
75. Heese K, Fiebich BL, Bauer J, Otten U. Nerve growth factor (NGF) expression in rat microglia is induced by adenosine A2a-receptors. *Neurosci Lett* 1997; **231**: 83–86.
76. Peakman MC, Hill SJ. Adenosine A2B-receptor-mediated cyclic AMP accumulation in primary rat astrocytes. *Br J Pharmacol* 1994; **111**: 191–198.
77. Pilitsis JG, Kimelberg HK. Adenosine receptor mediated stimulation of intracellular calcium in acutely isolated astrocytes. *Brain Res* 1998; **798**: 294–303.
78. Schwaninger M, Neher M, Viegas E, Schneider A, Spranger M. Stimulation of interleukin-6 secretion and gene transcription in primary astrocytes by adenosine. *J Neurochem* 1997; **69**: 1145–1150.
79. Lazarowski ER, Boucher RC. Purinergic receptors in airway epithelia. *Curr Opin Pharmacol* 2009; **9**: 262–267.
80. Corset V, Nguyen-Ba-Charvet KT, Forcet C, Moysé E, Chedotal A, Mehlen P. Netrin-1-mediated axon outgrowth and cAMP production requires interaction with adenosine A2b receptor. *Nature* 2000; **407**: 747–750.
81. Stein E, Zou Y, Poo M, Tessier-Lavigne M. Binding of DCC by netrin-1 to mediate axon guidance independent of adenosine A2B receptor activation. *Science* 2001; **291**: 1976–1982.
82. McKenna WL, Wong-Staal C, Kim GC, Macias H, Hinck L, Bartoe JL. Netrin-1-independent adenosine A2b receptor activation regulates the response of axons to netrin-1 by controlling cell surface levels of UNC5A receptors. *J Neurochem* 2008; **104**: 1081–1090.
83. Wittendorp MC, Boddeke HW, Biber K. Adenosine A3 receptor-induced CCL2 synthesis in cultured mouse astrocytes. *Glia* 2004; **46**: 410–418.
84. Hammarberg C, Fredholm BB, Schulte G. Adenosine A3 receptor-mediated regulation of p38 and extracellular-regulated kinase ERK1/2 via phosphatidylinositol-3'-kinase. *Biochem Pharmacol* 2004; **67**: 129–134.
85. Davalos D, Grutzendler J, Yang G, Kim JJ, Zuo Y, Jung S *et al*. ATP mediates rapid microglial response to local brain injury *in vivo*. *Nat Neurosci* 2005; **8**: 752–758.
86. Haynes SE, Hoppeler G, Yang G, Kurpius D, Dailey ME, Gan WB *et al*. The P2Y12 receptor regulates microglial activation by extracellular nucleotides. *Nat Neurosci* 2006; **9**: 1512–1519.
87. Chen Y, Corriden R, Inoue Y, Yip L, Hashiguchi N, Zinkernagel A *et al*. ATP release guides neutrophil chemotaxis via P2Y2 and A3 receptors. *Science* 2006; **314**: 1792–1795.
88. Miller RJ, Rostene W, Apartis E, Banisadr G, Biber K, Milligan ED *et al*. Chemokine action in the nervous system. *J Neurosci* 2008; **28**: 11792–11795.
89. Iadecola C, Nedergaard M. Glial regulation of the cerebral microvasculature. *Nat Neurosci* 2007; **10**: 1369–1376.
90. Koehler RC, Roman RJ, Harder DR. Astrocytes and the regulation of cerebral blood flow. *Trends Neurosci* 2009; **32**: 160–169.
91. Wilhelmsson U, Bushong EA, Price DL, Smarr BL, Phung V, Terada M *et al*. Redefining the concept of reactive astrocytes as cells that remain within their unique domains upon reaction to injury. *Proc Natl Acad Sci USA* 2006; **103**: 17513–17518.
92. Oberheim NA, Tian GF, Han X, Peng W, Takano T, Ransom B *et al*. Loss of astrocytic domain organization in the epileptic brain. *J Neurosci* 2008; **28**: 3264–3276.
93. Oberheim NA, Wang X, Goldman S, Nedergaard M. Astrocytic complexity distinguishes the human brain. *Trends Neurosci* 2006; **29**: 547–553.
94. Dimagl U, Niwa K, Lindauer U, Villringer A. Coupling of cerebral blood flow to neuronal activation: role of adenosine and nitric oxide. *Am J Physiol* 1994; **267**: H296–H301.
95. Iliff JJ, D'Ambrosio R, Ngai AC, Winn HR. Adenosine receptors mediate glutamate-evoked arteriolar dilation in the rat cerebral cortex. *Am J Physiol Heart Circ Physiol* 2003; **284**: H1631–H1637.
96. Ohata H, Cao S, Koehler RC. Contribution of adenosine A2A and A2B receptors and heme oxygenase to AMPA-induced dilation of pial arterioles in rats. *Am J Physiol Regul Integr Comp Physiol* 2006; **291**: R728–R735.
97. Fields RD, Burnstock G. Purinergic signalling in neuron–glia interactions. *Nat Rev Neurosci* 2006; **7**: 423–436.
98. Jimenez AI, Castro E, Mirabet M, Franco R, Delicado EG, Miras-Portugal MT. Potentiation of ATP calcium responses by A2B receptor stimulation and other signals coupled to Gs proteins in type-1 cerebellar astrocytes. *Glia* 1999; **26**: 119–128.
99. Alloisio S, Cugnoli C, Ferroni S, Nobile M. Differential modulation of ATP-induced calcium signalling by A1 and A2 adenosine receptors in cultured cortical astrocytes. *Br J Pharmacol* 2004; **141**: 935–942.
100. Xu HL, Pelligrino DA. ATP release and hydrolysis contribute to rat pial arteriolar dilatation elicited by neuronal activation. *Exp Physiol* 2007; **92**: 647–651.
101. Xu HL, Mao L, Ye S, Paisansathan C, Vetri F, Pelligrino DA. Astrocytes are a key conduit for upstream signaling of vasodilation during cerebral cortical neuronal activation *in vivo*. *Am J Physiol Heart Circ Physiol* 2008; **294**: H622–H632.
102. Takano T, Oberheim N, Cotrina ML, Nedergaard M. Astrocytes and ischemic injury. *Stroke* 2009; **40**: S8–12.
103. Eid T, Williamson A, Lee TS, Petroff OA, de Lanerolle NC. Glutamate and astrocytes – key players in human mesial temporal lobe epilepsy? *Epilepsia* 2008; **49** (Suppl 2): 42–52.
104. Tanaka K, Watase K, Manabe T, Yamada K, Watanabe M, Takahashi K *et al*. Epilepsy and exacerbation of brain injury in mice lacking the glutamate transporter GLT-1. *Science* 1997; **276**: 1699–1702.
105. Rothstein JD, Dykes-Hoberg M, Pardo CA, Bristol LA, Jin L, Kuncl RW *et al*. Knockout of glutamate transporters reveals a major role for astroglial transport in excitotoxicity and clearance of glutamate. *Neuron* 1996; **16**: 675–686.
106. Eid T, Thomas MJ, Spencer DD, Runden-Pran E, Lai JC, Malthankar GV *et al*. Loss of glutamine synthetase in the human epileptogenic hippocampus: possible mechanism for raised extracellular glutamate in mesial temporal lobe epilepsy. *Lancet* 2004; **363**: 28–37.
107. Volterra A, Meldolesi J. Astrocytes, from brain glue to communication elements: the revolution continues. *Nat Rev Neurosci* 2005; **6**: 626–640.
108. Bezzi P, Gunderson V, Galbete JL, Seifert G, Steinhauser C, Pilati E *et al*. Astrocytes contain a vesicular compartment that is competent for regulated exocytosis of glutamate. *Nat Neurosci* 2004; **7**: 613–620.

109. Lee SY, Haydon PG. Astrocytic glutamate targets NMDA receptors. *J Physiol* 2007; **581**: 887–888.
110. Nedergaard M. Direct signaling from astrocytes to neurons in cultures of mammalian brain cells. *Science* 1994; **263**: 1768–1771.
111. Parpura V, Basarsky TA, Liu F, Jęftinija K, Jęftinija S, Haydon PG. Glutamate-mediated astrocyte-neuron signalling. *Nature* 1994; **369**: 744–747.
112. Stevens B, Porta S, Haak LL, Gallo V, Fields RD. Adenosine: a neuron–glial transmitter promoting myelination in the CNS in response to action potentials. *Neuron* 2002; **36**: 855–868.
113. Jourdain P, Bergersen LH, Bhaukaurally K, Bezzi P, Santello M, Domercq M *et al*. Glutamate exocytosis from astrocytes controls synaptic strength. *Nat Neurosci* 2007; **10**: 331–339.
114. Yang Y, Ge W, Chen Y, Zhang Z, Shen W, Wu C *et al*. Contribution of astrocytes to hippocampal long-term potentiation through release of D-serine. *Proc Natl Acad Sci USA* 2003; **100**: 15194–15199.
115. Nagai M, Re DB, Nagata T, Chalazonitis A, Jessell TM, Wichterle H *et al*. Astrocytes expressing ALS-linked mutated SOD1 release factors selectively toxic to motor neurons. *Nat Neurosci* 2007; **10**: 615–622.
116. Boillee S, Vande Velde C, Cleveland DW. ALS: a disease of motor neurons and their nonneuronal neighbors. *Neuron* 2006; **52**: 39–59.
117. Yu L, Shen HY, Coelho JE, Araujo IM, Huang QY, Day YJ *et al*. 2008 Adenosine A2A receptor antagonists exert motor and neuroprotective effects by distinct cellular mechanisms. *Ann Neurol* 63: 338–346.
118. Lovatt D, Sonnewald U, Waagepetersen HS, Schousboe A, He W, Lin JH *et al*. The transcriptome and metabolic gene signature of protoplasmic astrocytes in the adult murine cortex. *J Neurosci* 2007; **27**: 12255–12266.
119. Tian GF, Azmi H, Takano T, Xu QW, Peng WG, Lin J *et al*. An astrocytic basis of epilepsy. *Nat Med* 2005; **11**: 973–981.
120. Binder DK, Steinhilber C. Functional changes in astroglial cells in epilepsy. *Glia* 2006; **54**: 358–368.
121. Li T, Lan JQ, Fredholm BB, Simon RP, Boison D. Adenosine dysfunction in astrogliosis: cause for seizure generation? *Neuron Glia Biol* 2007; **3**: 353–366.
122. Etherington LA, Patterson GE, Meechan L, Boison D, Irving AJ, Dale N *et al*. Astrocytic adenosine kinase regulates basal synaptic adenosine levels and seizure activity but not activity-dependent adenosine release in the hippocampus. *Neuropharmacology* 2009; **56**: 429–437.
123. Boison D. Adenosine augmentation therapies (AATs) for epilepsy: prospect of cell and gene therapies. *Epilepsies Res* 2009; **85**: 131–141.
124. Boison D. Engineered adenosine-releasing cells for epilepsy therapy: human mesenchymal stem cells and human embryonic stem cells. *Neurotherapeutics* 2009; **6**: 278–283.
125. Bell MJ, Kochanek PM, Carcillo JA, Mi Z, Schiding JK, Wisniewski SR *et al*. Interstitial adenosine, inosine, and hypoxanthine are increased after experimental traumatic brain injury in the rat. *J Neurotrauma* 1998; **15**: 163–170.
126. Kochanek PM, Vagni VA, Janesko KL, Washington CB, Crumrine PK, Garman RH *et al*. Adenosine A1 receptor knockout mice develop lethal status epilepticus after experimental traumatic brain injury. *J Cereb Blood Flow Metab* 2006; **26**: 565–575.
127. Fedele DE, Li T, Lan JQ, Fredholm BB, Boison D. Adenosine A₁ receptors are crucial in keeping an epileptic focus localized. *Exp Neurol* 2006; **200**: 184–190.
128. Li W, Dai S, An J, Xiong R, Li P, Chen X *et al*. Genetic inactivation of adenosine A2A receptors attenuates acute traumatic brain injury in the mouse cortical impact model. *Exp Neurol* 2009; **215**: 69–76.
129. Sachse KT, Jackson EK, Wisniewski SR, Gillespie DG, Puccio AM, Clark RS *et al*. Increases in cerebrospinal fluid caffeine concentration are associated with favorable outcome after severe traumatic brain injury in humans. *J Cereb Blood Flow Metab* 2007; **28**: 395–401.
130. Li W, Dai S, An J, Li P, Chen X, Xiong R *et al*. Chronic but not acute treatment with caffeine attenuates traumatic brain injury in the mouse cortical impact model. *Neuroscience* 2008; **151**: 1198–1207.
131. Gouder N, Scheurer L, Fritschy J-M, Boison D. Overexpression of adenosine kinase in epileptic hippocampus contributes to epileptogenesis. *J Neurosci* 2004; **24**: 692–701.
132. Ferre S, von Euler G, Johansson B, Fredholm BB, Fuxe K. Stimulation of high-affinity adenosine A2 receptors decreases the affinity of dopamine D2 receptors in rat striatal membranes. *Proc Natl Acad Sci USA* 1991; **88**: 7238–7241.
133. Richardson PJ, Gubitz AK, Freeman TC, Dixon AK. Adenosine receptor antagonists and Parkinson's disease: actions of the A2A receptor in the striatum. *Adv Neurol* 1999; **80**: 111–119.
134. Schwarzschild MA, Agnati L, Fuxe K, Chen JF, Morelli M. Targeting adenosine A2A receptors in Parkinson's disease. *Trends Neurosci* 2006; **29**: 647–654.
135. Jenner P, Mori A, Hauser R, Morelli M, Fredholm BB, Chen JF. Adenosine, adenosine A(2A) antagonists, and Parkinson's disease. *Parkinsonism Relat Disord* 2009; **15**: 406–413.
136. Schwarzschild MA. Adenosine A2A antagonists as neurotherapeutics: crossing the bridge. *Prog Neurobiol* 2007; **83**: 261–262.
137. Ross GW, Abbott RD, Petrovitch H, Morens DM, Grandinetti A, Tung KH *et al*. Association of coffee and caffeine intake with the risk of Parkinson disease. *JAMA* 2000; **283**: 2674–2679.
138. Ascherio A, Zhang SM, Hernan MA, Kawachi I, Colditz GA, Speizer FE *et al*. Prospective study of caffeine consumption and risk of Parkinson's disease in men and women. *Ann Neurol* 2001; **50**: 56–63.
139. Saaksjarvi K, Knekt P, Rissanen H, Laaksonen MA, Reunanen A, Mannisto S. Prospective study of coffee consumption and risk of Parkinson's disease. *Eur J Clin Nutr* 2008; **62**: 908–915.
140. Chen JF, Xu K, Petzer JP, Staal R, Xu YH, Beilstein M *et al*. Neuroprotection by caffeine and A(2A) adenosine receptor inactivation in a model of Parkinson's disease. *J Neurosci* 2001; **21**: RC143.
141. Ikeda K, Kurokawa M, Aoyama S, Kuwana Y. Neuroprotection by adenosine A2A receptor blockade in experimental models of Parkinson's disease. *J Neurochem* 2002; **80**: 262–270.
142. Kelly S, Zhang ZJ, Zhao H, Xu L, Giffard RG, Sapolsky RM *et al*. Gene transfer of HSP72 protects cornu ammonis 1 region of the hippocampus neurons from global ischemia: influence of Bcl-2. *Ann Neurol* 2002; **52**: 160–167.
143. Chen JF, Sonsalla PK, Pedata F, Melani A, Domenici MR, Popoli P *et al*. Adenosine A2A receptors and brain injury: broad spectrum of neuroprotection, multifaceted actions and 'fine tuning' modulation. *Prog Neurobiol* 2007; **83**: 310–331.
144. Pierrri M, Vaudano E, Sager T, Englund U. KW-6002 protects from MPTP induced dopaminergic toxicity in the mouse. *Neuropharmacology* 2005; **48**: 517–524.
145. Hagberg H, Andersson P, Lacarewicz J, Jacobson I, Butcher S, Sandberg M 1987 Extracellular adenosine, inosine, hypoxanthine, and xanthine in relation to tissue nucleotides and purines in rat striatum during transient ischemia. *J Neurochem* 49: 227–231.
146. Kopf SR, Melani A, Pedata F, Pepeu G. Adenosine and memory storage: effect of A(1) and A(2) receptor antagonists. *Psychopharmacology (Berl)* 1999; **146**: 214–219.
147. Gianfrido M, Corsi C, Melani A, Pezzola A, Reggio R, Popoli P *et al*. Adenosine A(2A) antagonism increases striatal glutamate outflow in the quinolinic acid rat model of Huntington's disease. *Brain Res* 2003; **979**: 225–229.
148. Latini S, Pedata F. Adenosine in the central nervous system: release mechanisms and extracellular concentrations. *J Neurochem* 2001; **79**: 463–484.
149. Latini S, Bordini F, Corradetti R, Pepeu G, Pedata F. Temporal correlation between adenosine outflow and synaptic potential inhibition in rat hippocampal slices during ischemia-like conditions. *Brain Res* 1998; **794**: 325–328.
150. Matsumoto K, Graf R, Rosner G, Shimada N, Heiss WD. Flow thresholds for extracellular purine catabolite elevation in cat focal ischemia. *Brain Res* 1992; **579**: 309–314.
151. Dux E, Fastbom J, Ungerstedt U, Rudolph K, Fredholm BB. Protective effect of adenosine and a novel xanthine derivative propentofylline on the cell damage after bilateral carotid occlusion in the gerbil hippocampus. *Brain Res* 1990; **516**: 248–256.
152. Frenguelli BG, Wigmore G, Llaudet E, Dale N. Temporal and mechanistic dissociation of ATP and adenosine release during ischaemia in the mammalian hippocampus. *J Neurochem* 2007; **101**: 1400–1413.
153. Phillis JW, Perkins LM, Smith-Barbour M, O'Regan MH. Transmitter amino acid release from rat neocortex: complete versus incomplete ischemia models. *Neurochem Res* 1994; **19**: 1387–1392.
154. Phillis JW, Smith-Barbour M, O'Regan MH. Changes in extracellular amino acid neurotransmitters and purines during and following ischemias of different durations in the rat cerebral cortex. *Neurochem Int* 1996; **29**: 115–120.
155. Schwarzschild MA, Chen JF, Ascherio A. Caffeinated clues and the promise of adenosine A(2A) antagonists in PD. *Neurology* 2002; **58**: 1154–1160.
156. Ribeiro JA, Sebastiao AM, de Mendonca A 2002 Adenosine receptors in the nervous system: pathophysiological implications. *Prog Neurobiol* 68: 377–392.
157. Ongini E, Adami M, Ferri C, Bertorelli R. Adenosine A2A receptors and neuroprotection. *Ann NY Acad Sci* 1997; **825**: 30–48.
158. Fredholm BB, Cunha RA, Svenningsson P. Pharmacology of adenosine A(2A) receptors and therapeutic applications. *Curr Top Med Chem* 2003; **3**: 413–426.
159. Cunha RA. Adenosine as a neuromodulator and as a homeostatic regulator in the nervous system: different roles, different sources and different receptors. *Neurochem Int* 2001; **38**: 107–125.
160. Rudolph KA, Schubert P, Parkinson FE, Fredholm BB. Neuroprotective role of adenosine in cerebral ischemia. *Trends Pharmacol Sci* 1992; **13**: 439–445.
161. Fredholm BB. Adenosine and neuroprotection. In: Green AR, Cross AJ (eds). *Neuroprotective agents and cerebral ischemia*. Academic Press: London, 1996 pp 259–280.
162. Kitagawa H, Mori A, Shimada J, Mitsumoto Y, Kikuchi T. Intracerebral adenosine infusion improves neurological outcome after transient focal ischemia in rats. *Neural Res* 2002; **24**: 317–323.
163. Phillis JW, O'Regan MH. Deoxycoformycin antagonizes ischemia-induced neuronal degeneration. *Brain Res Bull* 1989; **22**: 537–540.
164. Phillis JW, Walter GA, Simpson RE. Brain adenosine and transmitter amino acid release from the ischemic rat cerebral cortex: effects of the adenosine deaminase inhibitor deoxycoformycin. *J Neurochem* 1991; **56**: 644–650.
165. Lin Y, Phillis JW. Deoxycoformycin and oxypurinol: protection against focal ischemic brain injury in the rat. *Brain Res* 1992; **571**: 272–280.
166. Gidday JM. Cerebral preconditioning and ischaemic tolerance. *Nat Rev Neurosci* 2006; **7**: 437–448.

167. Miller LP, Jelovich LA, Yao L, DaRe J, Ugarkar B, Foster AC. Pre- and peristroke treatment with the adenosine kinase inhibitor, 5'-deoxyiodotubercidin, significantly reduces infarct volume after temporary occlusion of the middle cerebral artery in rats. *Neurosci Lett* 1996; **220**: 73–76.
168. DeLeo J, Schubert P, Kreutzberg GW. Protection against ischemic brain damage using propentofylline in gerbils. *Stroke* 1988; **19**: 1535–1539.
169. Andine P, Rudolphi KA, Fredholm BB, Hagberg H. Effect of propentofylline (HWA 285) on extracellular purines and excitatory amino acids in CA1 of rat hippocampus during transient ischaemia. *Br J Pharmacol* 1990; **100**: 814–818.
170. Kano T, Katayama Y, Kawamata T, Hirota H, Tsubokawa T. Propentofylline administered by microdialysis attenuates ischemia-induced hippocampal damage but not excitatory amino acid release in gerbils. *Brain Res* 1994; **641**: 149–154.
171. Matsumoto K, Sakaki T, Kohmura E, Hayakawa T, Yamada K. Amelioration of ischemic brain damage by the preischemic administration of propentofylline (HWA285) in a rat focal ischemia. *Brain Res* 1996; **723**: 228–230.
172. Fritschy J-M, Johnson DJ, Mohler H, Rudolph U. Independent assembly and subcellular targeting of GABA_A-receptor subtypes demonstrated in mouse hippocampal and olfactory neurons *in vivo*. *Neurosci Lett* 1998; **249**: 99–102.
173. Parkinson FE, Xiong W, Zamzow CR. Astrocytes and neurons: different roles in regulating adenosine levels. *Neurol Res* 2005; **27**: 153–160.
174. Pignataro G, Simon RP, Boison D. Transgenic overexpression of adenosine kinase aggravates cell death in ischemia. *J Cereb Blood Flow Metab* 2007; **27**: 1–5.
175. Zetterstrom T, Fillenz M. Adenosine agonists can both inhibit and enhance *in vivo* striatal dopamine release. *Eur J Pharmacol* 1990; **180**: 137–143.
176. Corradetti R, Lo Conte G, Moroni F, Passani MB, Pepeu G. Adenosine decreases aspartate and glutamate release from rat hippocampal slices. *Eur J Pharmacol* 1984; **104**: 19–26.
177. Greene RW, Haas HL. The electrophysiology of adenosine in the mammalian central nervous system. *Prog Neurobiol* 1991; **36**: 329–341.
178. Olsson T, Cronberg T, Rytter A, Asztely F, Fredholm BB, Smith ML *et al*. Deletion of the adenosine A1 receptor gene does not alter neuronal damage following ischaemia *in vivo* or *in vitro*. *Eur J Neurosci* 2004; **20**: 1197–1204.
179. Phillis JW. The effects of selective A1 and A2a adenosine receptor antagonists on cerebral ischemic injury in the gerbil. *Brain Res* 1995; **705**: 79–84.
180. Von Lubitz DK, Lin RC, Jacobson KA. Cerebral ischemia in gerbils: effects of acute and chronic treatment with adenosine A2A receptor agonist and antagonist. *Eur J Pharmacol* 1995; **287**: 295–302.
181. Bona E, Aden U, Gilland E, Fredholm BB, Hagberg H. Neonatal cerebral hypoxia-ischemia: the effect of adenosine receptor antagonists. *Neuropharmacology* 1997; **36**: 1327–1338.
182. Monopoli A, Lozza G, Forlani A, Mattavelli A, Ongini E. Blockade of adenosine A2A receptors by SCH 58261 results in neuroprotective effects in cerebral ischaemia in rats. *Neuroreport* 1998; **9**: 3955–3959.
183. Pedata F, Gianfriddo M, Turchi D, Melani A. The protective effect of adenosine A2A receptor antagonism in cerebral ischemia. *Neurol Res* 2005; **27**: 169–174.
184. Melani A, Gianfriddo M, Vannucchi MG, Cipriani S, Baraldi PG, Giovannini MG *et al*. The selective A(2A) receptor antagonist SCH 58261 protects from neurological deficit, brain damage and activation of p38 MAPK in rat focal cerebral ischemia. *Brain Res* 2006; **1073–1074**: 470–480.
185. Chen JF, Huang Z, Ma J, Zhu J, Moratalla R, Standaert D *et al*. A(2A) adenosine receptor deficiency attenuates brain injury induced by transient focal ischemia in mice. *J Neurosci* 1999; **19**: 9192–9200.
186. Pedata F, Corsi C, Melani A, Bordoni F, Latini S. Adenosine extracellular brain concentrations and role of A2A receptors in ischemia. *Ann NY Acad Sci* 2001; **939**: 74–84.
187. Marcoli M, Raiteri L, Bonfanti A, Monopoli A, Ongini E, Raiteri M *et al*. Sensitivity to selective adenosine A1 and A2A receptor antagonists of the release of glutamate induced by ischemia in rat cerebrocortical slices. *Neuropharmacology* 2003; **45**: 201–210.
188. Aden U, Halldner L, Lagercrantz H, Dalmau I, Ledent C, Fredholm BB. Aggravated brain damage after hypoxic ischemia in immature adenosine A2A knockout mice. *Stroke* 2003; **34**: 739–744.
189. Phillis JW. Adenosine and adenine nucleotides as regulators of cerebral blood flow: roles of acidosis, cell swelling, and KATP channels. *Crit Rev Neurobiol* 2004; **16**: 237–270.
190. von Lubitz DKJE. Adenosine and cerebral ischemia: therapeutic future or death of a brave concept? *Eur J Pharmacol* 1999; **365**: 9–25.
191. Chen JF, Pedata F. Modulation of ischemic brain injury and neuroinflammation by adenosine A(2A) receptors. *Curr Pharmaceutical Design* 2008; **14**: 1490–1499.
192. Irving EA, Barone FC, Reith AD, Hadingham SJ, Parsons AA. Differential activation of MAPK/ERK and p38/SAPK in neurones and glia following focal cerebral ischaemia in the rat. *Brain Res Mol Brain Res* 2000; **77**: 65–75.
193. Wu DC, Ye W, Che XM, Yang GY. Activation of mitogen-activated protein kinases after permanent cerebral artery occlusion in mouse brain. *J Cereb Blood Flow Metab* 2000; **20**: 1320–1330.
194. Barone FC, Irving EA, Ray AM, Lee JC, Kassis S, Kumar S *et al*. Inhibition of p38 mitogen-activated protein kinase provides neuroprotection in cerebral focal ischemia. *Med Res Rev* 2001; **21**: 129–145.
195. Gao Y, Signore AP, Yin W, Cao G, Yin XM, Sun F *et al*. Neuroprotection against focal ischemic brain injury by inhibition of c-Jun N-terminal kinase and attenuation of the mitochondrial apoptosis-signaling pathway. *J Cereb Blood Flow Metab* 2005; **25**: 694–712.
196. Kawasaki H, Morooka T, Shimohama S, Kimura J, Hirano T, Gotoh Y *et al*. Activation and involvement of p38 mitogen-activated protein kinase in glutamate-induced apoptosis in rat cerebellar granule cells. *J Biol Chem* 1997; **272**: 18518–18521.