

Author's accepted manuscript (postprint)

Administration of pro re nata medications by the nurse to incapacitated patients: An ethical perspective

Vaismoradi, M., Moe, C. F., Vizcaya-Moreno, M. F. & Paal, P.

Published in: Clinical Ethics

DOI: 10.1177/14777509211034146

Available online: 23 Jul 2021

Citation:

Vaismoradi, M., Moe, C. F., Vizcaya-Moreno, M. F. & Paal, P. (2021). Administration of pro re nata medications by the nurse to incapacitated patients: An ethical perspective. *Clinical Ethics*. doi: 10.1177/14777509211034146

© 2021. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0>

This is an Accepted Manuscript of an article published by Sage in *Clinical Ethics* on 23/07/2021, available online: <https://journals.sagepub.com/doi/10.1177/14777509211034146>

Administration of PRN Medications by the Nurse to Incapacitated Patients: An Ethical Perspective

Abstract

The administration of *pro re nata* (PRN) medications is the responsibility of the nurse. However, ethical uncertainties often happen due to the inability of incapacitated patients to collaborate with the nurse in the process of decision-making for PRN medication administration. There is a lack of integrative knowledge and insufficient understanding regarding ethical considerations surrounding the administration of PRN medications to incapacitated patients. Therefore, they have been discussed in this paper and practical strategies to avoid unethical practices have been suggested. The complicated caring situation surrounding the administration of PRN medications is intertwined with ethical issues affecting the consideration of the patient's wishes and interventions that override them. The patient's right of autonomy and treatment refusal, surrogacy role, paternalism, and coercion are the main ethos of ethical PRN medication administration. Education and training can help nurses avoid legal and ethical issues in PRN medicines management and improve the quality and safety of health care. Empirical research is needed to improve our understanding of this phenomenon in the multidisciplinary environment of medicines management.

Keywords: ethics, nurse, harm, medicines management, patient safety, *pro re nata*, PRN

Introduction

A medication error is defined as “an avoidable adverse effect of healthcare, whether or not it is obvious or harmful to the patient something incorrectly done through ignorance or inadvertence; a mistake, e.g., miscalculation, judgment, speech, writing, action, or a failure to complete a planned action as intended, or the use of an incorrect plan of action to achieve a given aim” (p. 6013) ¹. Medication administration error is a high-risk situation during medicines management and with a median of 8.0% is the primary cause of medication harm. It is characterised by errors in the timing, omitted, and wrong doses of medication in various healthcare conditions and age groups ^{2, 3}. A multicentre study in 38 hospitals in the UK on the prevalence, nature, and predictors of medication administration omissions using the MedsST tool indicated that medication administration omission was frequent and was influenced by the number of administered medications to patients ⁴. Therefore, the safety of medication administration is one of the most important initiatives for the improvement of patient safety in the healthcare system. The WHO has estimated that the global cost of medication errors would be 42 billion USD annually ⁵ and the Third Global Patient Safety Challenge on ‘Medication without harm’ has discussed issues and strategies to reduce harm associated with the medication process in high-risk healthcare situations ^{5, 6}.

The application of multidisciplinary and collaborative strategies by physicians, pharmacists, and nurses can enhance the safety of medicines management ⁷. In this respect, the administration of medications is the primary role of nurses with an undeniable impact on patient safety ^{8,9}. Nurses are trained and authorized to prepare medications, calculate doses, administer, and monitor medications' side effects and adverse drug reactions ¹⁰⁻¹². Safe administration of medications requires nurses to use their theoretical knowledge and clinical skills during the medication process ^{13,14}. The 'rights of medication administration' as patient, drug, route, dose, time, documentation, and reason have been recognized as traditionally important tools in the hands of nurses to ensure the safety of medication administration ¹⁵.

PRN Medication Administration by the Nurse

“*Pro re nata*” (PRN) medication administration is a main responsibility of nurses and has been described as the administration of medications based on the patient's immediate need instead of administration at predetermined times ^{16,17}. After the prescription of PRN medications by the physician, the nurse makes a collaborative decision with participation of the patient regarding medication administration ^{17,18}. PRN medicines management has been recognized by international healthcare associations as an appropriate method for the management of physical and psychological suffering among patients with acute and chronic healthcare conditions ¹⁹⁻²¹.

Prescription and administration of PRN medications have become a main part of medicines management in various healthcare settings. For instance, 90%, of patients with mental and psychiatric health conditions and 20-86% of residents in nursing homes are prescribed at least one PRN medication ²²⁻²⁴. The exact number of medication errors due to PRN medication administration has not been reported in the international literature. PRN medication errors are characterized by not mentioning the reason behind medication administration and ambiguities in steps taken to manage the administration process, as well as to monitor their effects and side effects after administration ^{25,26}. Coercion in the administration ²⁷, and over-administration of PRN medications, when medication is not the first choice for treatment ^{18,28} are quite common. Inappropriate use of PRN medications is associated with polypharmacy and longer stays in nursing homes with a median of 2.1 years ²⁹.

The safety of medicines management is intertwined with ethical nursing practice ^{30,31}. Nurses are professionally obliged to administer medications in a manner that is consistent with ethical codes for practice as 'do the good and right thing for patients', prevent harm, and evaluate the probable outcomes of medication administration interventions ³²⁻³⁴. Insufficient knowledge of

ethical considerations influencing all types of medication administration including PRN medications has become the source of moral distress in nurses³⁵, their burn-out and job turnover³⁶.

Decision Making for Care by Incapacitated Patients

Temporary or permanent physical and psychological incapacity during each person's life is common. Over one billion people, that is about 15% of the world population, live with some form of disability and incapacity and an increasing trend is captured in this figure³⁷. Incapacitated patients often suffer from severe mental or physical conditions to the extent that they have no sufficient understanding required for making or communicating responsible healthcare decisions^{38,39}. With a capacity to make decisions, the patient should have the ability to understand the meaning of the information provided by healthcare staff, can connect them to their healthcare situation, use the information to decide, and communicate their choice and preferences regarding therapeutic measures^{40,41}.

Incapacitated patients can be older people and mentally ill that have been described as highly vulnerable patients in the healthcare literature⁴². They sometimes have no friends or family to make medical decisions as default surrogates⁴³.

According to the Convention on the Rights of Persons with Disabilities (CRPD) by the United Nations (UN), patients with any sort of disability and incapacity have the right to benefit from appropriate decisions made on their lives⁴⁴. Making appropriate decisions with regard to healthcare and therapeutic measures influence the patient's wellbeing. If the patient has no capacity to make decisions, family members and healthcare providers can be authorised to make decisions on their behalf⁴⁵.

Decision-making for incapacitated patients who cannot make decisions for themselves is considered a common concern for healthcare providers and is often intertwined with ethical issues⁴⁶, given that the patient's informed consent is a prerequisite for all types of healthcare interventions. Nevertheless, interventions that are in the interest of such incapacitated patients and when the interventions are accompanied with a minimum risk and burden are exempt from this rule⁴⁷. Having no capacity to decide imposes high levels of care burden on healthcare providers especially when medications are prescribed and administered, and in case that patient refuses to undertake healthcare procedures⁴⁸⁻⁵⁰. This situation is also accompanied by various ethical complexities affecting the society and healthcare staff individually^{51,52}.

PRN Medicines Management for Incapacitated Patients

PRN medicines management generally is the process of creating a mutual understanding and the feeling of responsibility between the nurse and the patient in order to choose the most effective medication with the lowest possibility of side effects and adverse drug reactions^{14, 17}. However, the requirements for making bilateral decisions by the nurse and the patient may not be achieved given the patient's incapacity to understand and actively take part in the medication process as a member of the healthcare team⁵³⁻⁵⁵. This situation creates barriers to ethical decision making by clinical nurses and can be the ground for the development of ethical dilemma.

Given the lack of integrative knowledge and insufficient understanding of ethical challenges surrounding the administration of PRN medications to incapacitated patients, the question is: how nurses can avoid unethical practice in PRN medicines management for these patients?

Ethics of Administration of PRN Medications to Incapacitated Patients

The nurse is obliged to play the role of the patient's advocate and put the patient's individual preferences and wishes at the centre for making the decision to administer PRN medications. However, in some situations the healthcare professional's wishes override those of the patients to benefit the patient and prevent harm. The ethical problem is raised in the preservation of the right to accept or refuse receiving medications given the patient's right of autonomy. Respect for the patient's autonomy requires to recognize his/her right to decide about own treatment and to refuse receiving treatment in case that the patient understands the action and is not forced to comply with the action⁵⁶⁻⁵⁸.

Complying with the patient's wishes

Some patients such as nonverbal trauma patients or those with severe mental and cognitive impairments may have no or very limited decision-making capacity. Therefore, they may be unable to understand the reason for PRN medication administration and collaborate in the related process. Therefore, being the patient's representative or taking the surrogacy role to predict therapeutic needs for such incapacitated patients is needed. A suggested solution is the involvement of patients' families for the prediction of their preferences during the periods of incapacity and to receive treatments that comply with their own needs and wishes^{59, 60}. Family members are appropriate sources to find the most suitable treatment and care modality⁶¹, because they can better guess the patient's preferences in similar circumstances and how their needs can be met⁶². Family members also can play the role of the patient's surrogate and the

essential partner for making decisions on the patient's behalf and can share the responsibility of decision making with nurses. This approach to decision-making can prevent moral and emotional distress in clinical nurses and enhance trust and consensus between healthcare professionals and family members ⁶³.

Making decisions to perform any sort of medical interventions including the administration of PRN medications to incapacitated patients who have no family members to take the surrogate role is also complicated and is filled by ambiguities ⁶⁴. Therefore, obtaining consent to start the medical procedure for these patients is intertwined with ethical issues and the patient's right problem ⁴³. If it is not resolved by the healthcare team, an ethical dilemma may occur given its impact on the patient's care outcome ⁶⁵. As if it is not a perfect solution, the use of clinical guidelines and a three-step approach of the physician, ethics committee, and guardian have been suggested ⁶⁴. The essential legislative core for all European member states to guarantee the protection of human rights regarding biology, medicine and healthcare is the 'Convention for the Protection of Human Rights and Dignity of the Human Being about the Application of Biology and Medicine' ⁶⁶ and its additional protocol ⁶⁷. This legislative document recommended that "Where, according to law, an adult does not have the capacity to consent to an intervention because of a mental disability, a disease or for similar reasons, the intervention may only be carried out with the authorization of his or her representative or an authority or a person or body provided for by law." ⁶⁶.

Nevertheless, ethical issues remain unresolved even if healthcare professionals including the nurse take the surrogate role as they need to identify the which extent therapeutic interventions for patient care can be performed ⁶⁸. The nurse is obliged to collect all types of data from the patient including verbal and nonverbal clues as well as consider vital signs and symptoms to find how he/she can meet the patients' physical and psychological needs before making the decision on the administration of PRN medications ⁶⁹. The patient's behaviours should be observed and interpreted to decide about the need to the involuntary administration of PRN medications to incapacitated patients. However, this could be a challenge to go beyond reported symptoms and understand their meanings in order to provide individualised care ⁷⁰. In this respect, the nurse's personal attitudes, beliefs and knowledge regarding symptoms and behavioural clues indicating the need for medication can influence his/her decision to administer medications to a great extend ⁷¹.

Overriding the patients' wishes

A source of moral concerns is to take a paternalistic role and coerce the patient to take PRN medications while the agitated patient is unwilling to cooperate and resist taking medications⁷²⁻⁷⁴. Beauchamp and Childress (2001) define paternalism as “the intentional overriding of one person’s known preferences or actions by another person, where the person who overrides justifies the action by the goal of benefitting or avoiding harm to the person whose preferences or actions are overridden” (p. 178)⁵⁶. Paternalism is the sharp end of professional beneficence⁷⁵ and has been suggested to be considered when the patient is severely incapacitated and when treatment is in the patient’s best interest⁷⁶. However, it is contradictory to the value of autonomy and is a source of tension between restraint and freedom, even if it may accompany by good intentions in patient care⁷⁷. The nurse’s effort to pursue the patient’s autonomy as an end state reflects the value given by him/her to paternalism. He/she may act paternalistically based on the assumption that his/her actions may allow the patient to have a more autonomous life in the future⁷⁸.

It is noted that even if physical force is not applied by the nurse, the persuasion of patients to take medications limits the patient’s autonomy⁷⁹. In such situations, the use of informal coercion as an ‘unpleasant good’ can be considered if it is one part of the nurse’s job description, despite inducing negative feelings in the patient and discomfort in the nurse⁸⁰⁻⁸². Although it may create feeling of safety in the patient, it is also associated with some negative consequences including the feeling of fear and disempowerment in the nurse. The nurse has the responsibility to protect the human rights of patients, and he/she should be aware of situations in which informal coercion is used and what its potential adverse psychological effects can be on therapeutic relationships^{76,83}.

As a matter of self-determination, disabled and incapacitated patients may decline invasive procedures for medication administration compared to less invasive ones such as the oral route⁸⁴. Therefore, there are some clinical situations in which covert medication administration has been suggested for those incapacitated patients who refuse taking medications but indicate a high risk of self-harm⁸⁵. Nevertheless, complex clinical and ethical issues appear when covert medication administration is used⁸⁶, as it can damage the relationship between the patient and the family member, with the nurse.

Patients have the right to refuse medications unless there is a court order obligating them to take the medication or in emergency situations and only for a limited duration. This is a change

from previous practices in which those patients who have been involuntarily hospitalized does not have decision-making capacity and is considered unable to refuse any kind of medication. In this difficult ethical situation, the nurse should face the problem of balancing their own level of control over decision making with the preferences of incapacitated patients⁸⁷. Therefore, the nurse has the responsibility to assess the situation objectively and document the medication process to avoid forthcoming legal and ethical issues^{81, 88}.

An argument in favour of paternalism emphasizes the utilitarian principle in which decisions for the administration of PRN medications may be taken because of the control of patients' behaviour to benefit other people than the patient him/herself. Roberts (2004)⁸⁹ affirms that in case of illnesses leading to severe incapacity, paternalism will be exercised by the means of interventions aiming at the alleviation of human suffering for those afflicted with that illness. From an opposing perspective, the negative consequences of forced and involuntary administration of medication may include further detachment from therapeutic measures that can lead to a violent outbreak requiring the use of restraints and seclusion^{90, 91} to gain control over the patient's threatening behaviours.

How the nurse responds to an ethical situation is influenced by his/her understanding of the illness. If mental illness is considered a 'bodily disease', medications will be seen beneficial for avoiding human suffering. On the other hand, if the mental illness is understood as a deviation from the culture's values and norms, a paternalistic medical intervention becomes a political act rather than a caring one⁸⁹.

Paternalistic culture-bound preferences should not affect patient care. Each patient must be approached impartially addressing their needs, values, and beliefs. Ethical principles and the requirement for best available care apply regardless of ethnic, cultural, and religious background, even if the patient is incapacitated. A summary of key ethical factors surrounding PRN medication administration has been presented in figure 1.




Figure 1. Ethical factors affecting PRN medication administration

Implications for clinical practice

Nurses have the required knowledge and clinical expertise to help with relieving physical and psychological suffering among patients ⁹², and the administration of PRN medications allows nurses to become more actively involved in medicines management initiatives in the multidisciplinary environment of health care ⁹³. The requirement for the safe administration of PRN medications by nurses to incapacitated patients is to comply with regulatory, professional, ethical and legal frameworks that govern it to establish a caring environment in which the human rights, dignity, values, custom, and spiritual beliefs of the patients are all respected ⁹⁴.

A summary of ethical considerations of the administration of PRN medications to incapacitated patients and clinical strategies to avoid unethical practice by the nurse has been presented (Table 1).

Table 1. PRN medication administration and strategies to avoid unethical practice

Ethical considerations	Practical strategy for the administration of PRN medications
Patient's autonomy and right of treatment refusal	Seeking partnership with the family surrogate who can help with the interpretation of the patient's behaviours and prediction of his/her preferences; Incorporation of the patient's wishes and interests into the decision for medication administration; Prediction of the impact of the decision on the patient's care outcomes; Consideration of clinical guidelines and practical directives to manage difficult situations; Approaching the physician, ethics committee, guardian or an authority endorsed by the law;
Surrogacy role	Observation and interpretation of the patient's verbal and nonverbal behavioural clues; Taking vital signs and assessment of symptoms indicating physical and psychological suffering; Self-reflection on personal attitudes and beliefs regarding the interpretation of the patient's symptoms and behaviours; Improvement of knowledge through education regarding how to identify the patient's need to medications
Paternalism and patient coercion	Use of enforcement and covert administration of PRN medications only in case that it is in the patient's best interest and can prevent further harm to the patient and others; Establishment of coercing interventions on the job description and with the consideration of legal aspects; Revision of personal beliefs justifying the use of coercion through peer discussion and seeking leadership support; Exploration of the negative consequences of coercion and covert medications for the therapeutic relationship between the nurse and family members; Prioritising the use of less invasive methods of medication administration; Seeking the legal and court order to justify the duration and possibility of medication use; Creation of balance between the patient's preferences and the nurse's control; Objective assessment and documentation of the decision-making process; Being impartial for the assessment of the patient's needs, beliefs, and values

Conclusion

The administration of PRN medications to incapacitated patients creates a complicated caring situation for the nurse and is intertwined with ethical issues affecting the consideration of the patient's wishes and interventions that override them. The patient's right of autonomy and treatment refusal, surrogacy role, paternalism, and coercion are the main ethos of ethical PRN medication administration. Education and training can help nurses avoid legal and ethical issues in PRN medicines management and improve the quality and safety of health care.

The improvement of the nurses' attitudes regarding the ethical considerations of PRN medication administration and their application to clinical practice requires discussion and reflection by nurses, multidisciplinary collaboration, and leadership support. Healthcare organisations can heed such an exchange of ideas by introducing supervision or more informal discussion rounds.

It is also suggested to incorporate these ethical aspects into clinical guidelines and directives for the safe administration of PRN medications to incapacitated patients. Also, extra efforts should be made by the nurse to evaluate the effectiveness of medication and the possibility of harm, as well as detect the possible side effects of medications and adverse drug reactions, given the inability of incapacitated patients to report them.

Empirical research is needed to explore more the interconnection between the administration of PRN medications and ethics in the multidisciplinary environment. Also, there is a need to conduct experimental studies to investigate how the ethical aspects and principles of PRN medicines management can improve the quality and safety of health care.

Conflicts of interest declaration: The authors declare that there is no conflict of interest.

Funding: This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

References

1. Aronson JK. Medication errors: definitions and classification. *Br J Clin Pharmacol* 2009; 67: 599-604. 2009/07/15. DOI: 10.1111/j.1365-2125.2009.03415.x.
2. Keers RN, Hann M, Alshehri GH, et al. Prevalence, nature and predictors of omitted medication doses in mental health hospitals: A multi-centre study. *PLoS One* 2020; 15: e0228868. 2020/02/07. DOI: 10.1371/journal.pone.0228868.
3. Keers RN, Williams SD, Cooke J, et al. Prevalence and nature of medication administration errors in health care settings: a systematic review of direct observational evidence. *Ann Pharmacother* 2013; 47: 237-256. 2013/02/07. DOI: 10.1345/aph.1R147.
4. Rostami P, Heal C, Harrison A, et al. Prevalence, nature and risk factors for medication administration omissions in English NHS hospital inpatients: a retrospective multicentre study using Medication Safety Thermometer data. *BMJ Open* 2019; 9: e028170. DOI: 10.1136/bmjopen-2018-028170.
5. World health Organization (WHO). Patient safety. WHO global patient safety challenge: medication without harm, <http://www.who.int/patientsafety/medication-safety/en/> (2017, accessed March 11, 2021).
6. World Health Organization (WHO). Medication safety in key action areas, <https://www.who.int/patientsafety/medication-safety/technical-reports/en/> (2019, March 11, 2021).
7. Manias E, Kusljic S and Wu A. Interventions to reduce medication errors in adult medical and surgical settings: a systematic review. *Therapeutic Advances in Drug Safety* 2020; 11: 2042098620968309. DOI: 10.1177/2042098620968309.
8. Hayes C, Jackson D, Davidson PM, et al. Medication errors in hospitals: a literature review of disruptions to nursing practice during medication administration. *J Clin Nurs* 2015; 24: 3063-3076. 2015/08/11. DOI: 10.1111/jocn.12944.
9. Magalhães AMM, Kreling A, Chaves EHB, et al. Medication administration - nursing workload and patient safety in clinical wards. *Rev Bras Enferm* 2019; 72: 183-189. 2019/03/28. DOI: 10.1590/0034-7167-2018-0618.
10. GuKG N. Gesundheits-und Krankenpflegegesetz 1997, BGBl.INr.108/1997i.d.F.BGBl.INr.8/2016,

https://www.ris.bka.gv.at/Dokumente/BgblAuth/BGBLA_2016_I_75/BGBLA_2016_I_75.pdf (1997, accessed March 11, 2021).

11. Health Information and Quality Authority (HIQA). Medicines Management Guidance, <https://www.hiqa.ie/sites/default/files/2017-01/Medicines-Management-Guidance.pdf> (2015, accessed March 11, 2021).
12. Pegram A and Bloomfield J. Medicines management. *Nurs Stand* 2015; 29: 36-43. 2015/04/16. DOI: 10.7748/ns.29.33.36.e9194.
13. Luokkamäki S, Härkänen M, Saano S, et al. Registered Nurses' medication administration skills: a systematic review. *Scandinavian Journal of Caring Sciences* 2021; 35: 37-54. DOI: <https://doi.org/10.1111/scs.12835>.
14. Vaismoradi M, Jordan S, Vizcaya-Moreno F, et al. PRN Medicines Optimization and Nurse Education. *Pharmacy (Basel)* 2020; 8 2020/10/30. DOI: 10.3390/pharmacy8040201.
15. Elliott M and Liu Y. The nine rights of medication administration: an overview. *British Journal of Nursing* 2010; 19: 300-305. DOI: 10.12968/bjon.2010.19.5.47064.
16. Adelaide D and Rawther F. Audit: Prescribing PRN medication. *Psychiatr Danub* 2017; 29: 568-570. 2017/09/28.
17. Vaismoradi M, Amaniyan S and Jordan S. Patient Safety and Pro Re Nata Prescription and Administration: A Systematic Review. *Pharmacy (Basel)* 2018; 6 2018/08/31. DOI: 10.3390/pharmacy6030095.
18. Hipp K, Repo-Tiihonen E, Kuosmanen L, et al. Patient participation in pro re nata medication in forensic psychiatric care: A nursing document analysis. *Journal of Psychiatric and Mental Health Nursing* 2020; n/a. DOI: <https://doi.org/10.1111/jpm.12706>.
19. Gordon DB, Dahl J, Phillips P, et al. The use of "as-needed" range orders for opioid analgesics in the management of acute pain: a consensus statement of the American Society for Pain Management Nursing and the American Pain Society. *Pain Manag Nurs* 2004; 5: 53-58. 2004/08/07. DOI: 10.1016/j.pmn.2004.04.001.
20. Gordon DB, Dahl J, Phillips P, et al. The use of 'as-needed' range orders for opioid analgesics in the management of acute pain: a consensus statement of the American Society for Pain Management Nursing and the American Pain Society. *Home Healthc Nurse* 2005; 23: 388-396; quiz 397-388. 2005/06/16. DOI: 10.1097/00004045-200506000-00012.
21. Molloy L, Field J, Beckett P, et al. PRN psychotropic medication and acute mental health nursing: reviewing the evidence. *J Psychosoc Nurs Ment Health Serv* 2012; 50: 12-15. 2012/07/18. DOI: 10.3928/02793695-20120703-03.
22. Dörks M, Allers K and Hoffmann F. Pro Re Nata Drug Use in Nursing Home Residents: A Systematic Review. *J Am Med Dir Assoc* 2019; 20: 287-293.e287. 2018/12/16. DOI: 10.1016/j.jamda.2018.10.024.
23. Griffiths AW, Surr CA, Alldred DP, et al. Pro re nata prescribing and administration for neuropsychiatric symptoms and pain in long-term care residents with dementia and memory problems: a cross-sectional study. *Int J Clin Pharm* 2019; 41: 1314-1322. 2019/07/26. DOI: 10.1007/s11096-019-00883-7.
24. Martin K, Arora V, Fischler I, et al. Descriptive analysis of pro re nata medication use at a Canadian psychiatric hospital. *Int J Ment Health Nurs* 2017; 26: 402-408. 2016/11/03. DOI: 10.1111/inm.12265.
25. Vaismoradi M, Vizcaya Moreno F, Sletvold H, et al. PRN Medicines Management for Psychotropic Medicines in Long-Term Care Settings: A Systematic Review. *Pharmacy (Basel, Switzerland)* 2019; 7: 157. DOI: 10.3390/pharmacy7040157.
26. Walsh B, Dahlke S, O'Rourke H, et al. Nurses' decision-making related to administering as needed psychotropic medication to persons with dementia: an empty systematic review. *International Journal of Older People Nursing* 2021; 16: e12350. DOI: <https://doi.org/10.1111/opn.12350>.

27. Hipp K, Kuosmanen L, Repo-Tiihonen E, et al. Patient participation in pro re nata medication in psychiatric inpatient settings: An integrative review. *Int J Ment Health Nurs* 2018; 27: 536-554. 2017/12/23. DOI: 10.1111/inm.12427.
28. Usher K, Holmes C, Lindsay D, et al. PRN psychotropic medications: The need for nursing research. *Contemporary Nurse* 2003; 14: 248-257. DOI: 10.5172/conu.14.3.248.
29. Dörks M, Schmiemann G and Hoffmann F. Pro re nata (as needed) medication in nursing homes: the longer you stay, the more you get? *Eur J Clin Pharmacol* 2016; 72: 995-1001. 2016/04/15. DOI: 10.1007/s00228-016-2059-4.
30. King CA. Clinical Ethics: Patient and Provider Safety. *AORN Journal* 2017; 106: 548-551. DOI: <https://doi.org/10.1016/j.aorn.2017.10.003>.
31. Soltanian MM, Z.; Mohammadi, E.; Sharif, F.; Rakhshan, M.,. Professional responsibility: An ethical concept extracted from practices of Iranian nurses during drug administration. *International Journal of Pharmaceutical Research* 2018; 10: 346-353.
32. Burke A. Ethical Practice: NCLEX-RN, <https://www.registerednursing.org/nclex/ethical-practice/> (2021, accessed March 11, 2021).
33. Durham B. The nurse's role in medication safety. *Nursing* 2015; 45 2015/03/19. DOI: 10.1097/01.NURSE.0000461850.24153.8b.
34. Erlen JA. Medication errors: ethical implications. *Orthop Nurs* 2001; 20: 82-85. 2002/05/25. DOI: 10.1097/00006416-200107000-00013.
35. Dodek P, Norena M, Ayas N, et al. Moral distress in intensive care unit personnel is not consistently associated with adverse medication events and other adverse events. *J Crit Care* 2019; 53: 258-263. 2019/07/14. DOI: 10.1016/j.jcrc.2019.06.023.
36. Karakachian A and Colbert A. Nurses' Moral Distress, Burnout, and Intentions to Leave: An Integrative Review. *J Forensic Nurs* 2019; 15: 133-142. 2019/08/23. DOI: 10.1097/jfn.0000000000000249.
37. World Health Organization (WHO). Disability, https://www.who.int/health-topics/disability#tab=tab_2 (2021, March 11, 2021).
38. DeMartino ES, Dudzinski DM, Doyle CK, et al. Who Decides When a Patient Can't? Statutes on Alternate Decision Makers. *N Engl J Med* 2017; 376: 1478-1482. DOI: 10.1056/NEJMms1611497.
39. Duhaime's Law Dictionary. Legal dictionary: Incapacitated Definition, <http://www.duhaime.org/LegalDictionary/I/Incapacitated.aspx#:~:text=%22Incapacitated%20person%20means%20any%20person,make%20or%20communicate%20responsible%20decisions.%22> (2021, accessed March 11, 2021).
40. Palmer BW and Harmell AL. Assessment of Healthcare Decision-making Capacity. *Arch Clin Neuropsychol* 2016; 31: 530-540. 2016/08/24. DOI: 10.1093/arclin/acw051.
41. Schröder-Bäck P, Duncan P, Sherlaw W, et al. Teaching seven principles for public health ethics: towards a curriculum for a short course on ethics in public health programmes. *BMC Med Ethics* 2014; 15: 73. 2014/10/08. DOI: 10.1186/1472-6939-15-73.
42. Courtwright A and Rubin E. Who should Decide for the Unrepresented? *Bioethics* 2016; 30: 173-180. 2015/08/27. DOI: 10.1111/bioe.12185.
43. Pope T. Unbefriended And Unrepresented: Better Medical Decision Making For Incapacitated Patients Without Healthcare Surrogates. *Georgia State University Law Review* 2017; 33.
44. United nation (UN): Department of Economic and Social Affairs Disability. Convention on the Rights of Persons with Disabilities (CRPD), <https://www.un.org/development/desa/disabilities/convention-on-the-rights-of-persons-with-disabilities.html> (2008, accessed March 11, 2021).
45. Kaplunov EH, N; () The European Proceedings of Social & Behavioural Sciences EpSBS, Volume XIX. *Incapacitated Patients' Wellbeing: Surrogate Decision Making*. In: Casati, F and Barysheva, GA and Krieger, W, (eds.). 2017.
46. Moye J, Catlin C, Kwak J, et al. Ethical Concerns and Procedural Pathways for Patients Who are Incapacitated and Alone: Implications from a Qualitative Study for Advancing Ethical Practice. *HEC Forum* 2017; 29: 171-189. 2017/01/14. DOI: 10.1007/s10730-016-9317-9.

47. World Health Organization (WHO). A declaration on the promotion of patients' rights in Europe. European consultation on the rights of patients Amsterdam 28 - 30 March 1994., https://www.who.int/genomics/public/eu_declaration1994.pdf (1994, accessed March 11, 2021).
48. Barstow C, Shahan B and Roberts M. Evaluating Medical Decision-Making Capacity in Practice. *Am Fam Physician* 2018; 98: 40-46. 2018/09/15.
49. Lee GB, Woo H, Lee SY, et al. The burden of care and the understanding of disease in Parkinson's disease. *PLoS One* 2019; 14: e0217581. 2019/06/01. DOI: 10.1371/journal.pone.0217581.
50. Rezaei H, Niksima SH and Ghanei Gheshlagh R. Burden of Care in Caregivers of Iranian patients with chronic disorders: a systematic review and meta-analysis. *Health Qual Life Outcomes* 2020; 18: 261. 2020/08/05. DOI: 10.1186/s12955-020-01503-z.
51. Kulju K, Suhonen R and Leino-Kilpi H. Ethical problems and moral sensitivity in physiotherapy: a descriptive study. *Nurs Ethics* 2013; 20: 568-577. 2013/01/19. DOI: 10.1177/0969733012468462.
52. Sequeira AL-S and Lewis A. Ethical and Legal Considerations in the Management of an Unbefriended Patient in a Vegetative State. *Neurocritical Care* 2017; 27: 173-179. DOI: 10.1007/s12028-017-0405-8.
53. Dilles T, Van Rompaey B, Van Bogaert P, et al. Resident and nurse reports of potential adverse drug reactions. *Eur J Clin Pharmacol* 2015; 71: 741-749. 2015/04/24. DOI: 10.1007/s00228-015-1848-5.
54. Murray L. The role of the registered nurse managing pro re nata (PRN) medicines in the care home (nursing): a case study of decision-making, medication management and resident involvement, <http://hdl.handle.net/2299/17989>, <https://doi.org/10.18745/th.17989> (2017, accessed March 11, 2021).
55. Virgolesi M, Pucciarelli G, Colantoni AM, et al. The effectiveness of a nursing discharge programme to improve medication adherence and patient satisfaction in the psychiatric intensive care unit. *J Clin Nurs* 2017; 26: 4456-4466. 2017/02/25. DOI: 10.1111/jocn.13776.
56. Beauchamp T and Childress J. *Principles of biomedical ethics*. 5th ed. New York: Oxford University Press, 2001.
57. Gillon R. Ethics needs principles--four can encompass the rest--and respect for autonomy should be "first among equals". *J Med Ethics* 2003; 29: 307-312. 2003/10/02. DOI: 10.1136/jme.29.5.307.
58. Sjöstrand M and Helgesson G. Coercive treatment and autonomy in psychiatry. *Bioethics* 2008; 22: 113-120. 2008/02/07. DOI: 10.1111/j.1467-8519.2007.00610.x.
59. Rid A, Wesley R, Pavlick M, et al. Patients' priorities for treatment decision making during periods of incapacity: quantitative survey. *Palliat Support Care* 2015; 13: 1165-1183. 2014/10/02. DOI: 10.1017/S1478951514001096.
60. Wendler D, Wesley B, Pavlick M, et al. A new method for making treatment decisions for incapacitated patients: what do patients think about the use of a patient preference predictor? *Journal of Medical Ethics* 2016; 42: 235-241. DOI: 10.1136/medethics-2015-103001.
61. Kelly B, Rid A and Wendler D. Systematic review: Individuals' goals for surrogate decision-making. *J Am Geriatr Soc* 2012; 60: 884-895. 2012/04/04. DOI: 10.1111/j.1532-5415.2012.03937.x.
62. Rid A and Wendler D. Can we improve treatment decision-making for incapacitated patients? *Hastings Cent Rep* 2010; 40: 36-45. 2010/10/23. DOI: 10.1353/hcr.2010.0001.
63. Sussman B. A narrative approach to the ethical dilemmas of surrogate decision making. *Progress in Palliative Care* 2015; 23: 137-141. DOI: 10.1179/1743291X14Y.0000000114.
64. Schweikart SJ. Who Makes Decisions for Incapacitated Patients Who Have No Surrogate or Advance Directive? *AMA J Ethics* 2019; 21: E587-593. 2019/07/25. DOI: 10.1001/amajethics.2019.587.
65. Haskins DR and Wick JY. Medication Refusal: Resident Rights, Administration Dilemma. *Consult Pharm* 2017; 32: 728-736. 2018/02/23. DOI: 10.4140/TCP.n.2017.728.
66. Council of Europe. Convention for the protection of Human Rights and Dignity of the Human Being with regard to the Application of Biology and Medicine: Convention on Human Rights and

Biomedicine. ETS No 164, <https://www.coe.int/en/web/conventions/full-list/-/conventions/treaty/164> (1997, accessed March 11, 2021).

67. Council of Europe. Additional Protocol to the Convention on Human Rights and Biomedicine, concerning Biomedical Research. ETS No 195, <https://www.coe.int/en/web/conventions/full-list/-/conventions/treaty/195> (2005, accessed March 11, 2021).
68. White DB, Jonsen A and Lo B. Ethical challenge: when clinicians act as surrogates for unrepresented patients. *Am J Crit Care* 2012; 21: 202-207. 2012/05/03. DOI: 10.4037/ajcc2012514.
69. Di Giulio P and Crow R. Cognitive Processes Nurses and Doctors Use in the Administration of PRN (at need) Analgesic Drugs. *Scandinavian Journal of Caring Sciences* 1997; 11: 12-19. DOI: <https://doi.org/10.1111/j.1471-6712.1997.tb00425.x>.
70. Bygstad-Landro M and Giske T. Risking existence: The experience and handling of depression. *J Clin Nurs* 2018; 27: e514-e522. 2017/09/02. DOI: 10.1111/jocn.14056.
71. Hamers JP, Abu-Saad HH, van den Hout MA, et al. Are children given insufficient pain-relieving medication postoperatively? *J Adv Nurs* 1998; 27: 37-44. 1998/03/27. DOI: 10.1046/j.1365-2648.1998.00493.x.
72. Brodtkorb K, Skisland AV, Slettebø Å, et al. Ethical challenges in care for older patients who resist help. *Nurs Ethics* 2015; 22: 631-641. 2014/08/26. DOI: 10.1177/0969733014542672.
73. Moermans VRA, Bleijlevens MHC, Verbeek H, et al. The use of involuntary treatment among older adults with cognitive impairment receiving nursing care at home: A cross-sectional study. *Int J Nurs Stud* 2018; 88: 135-142. 2018/10/09. DOI: 10.1016/j.ijnurstu.2018.09.004.
74. Wheeler AJ, Hu J, Profitt C, et al. Is higher psychotropic medication burden associated with involuntary treatment under the Mental Health Act? A four-year Australian cohort study. *BMC Psychiatry* 2020; 20: 294. DOI: 10.1186/s12888-020-02661-6.
75. Mitchell V. *Professional relationships*. In: Barker P (ed.), *Mental health ethics*. New York: Routledge, 2011.
76. Szmukler G and Appelbaum PS. Treatment pressures, leverage, coercion, and compulsion in mental health care. *Journal of Mental Health* 2008; 17: 233-244. DOI: 10.1080/09638230802052203.
77. Double D. *The psychiatrist*. In: Barker P (ed.) *Mental health ethics: the human context*. New York: Routledge, 2011.
78. Kashka MS and Keyser PK. Ethical issues in informed consent and ECT. *Perspect Psychiatr Care* 1995; 31: 15-21; quiz 22-14. 1995/04/01. DOI: 10.1111/j.1744-6163.1995.tb00461.x.
79. Ventura CAA, Austin W, Carrara BS, et al. Nursing care in mental health: Human rights and ethical issues. *Nurs Ethics* 2020; 27: 969733020952102. 2020/10/29. DOI: 10.1177/0969733020952102.
80. Andersson U, Fathollahi J and Gustin LW. Nurses' experiences of informal coercion on adult psychiatric wards. *Nurs Ethics* 2020; 27: 741-753. 2020/01/04. DOI: 10.1177/0969733019884604.
81. Lind M, Kaltiala-Heino R, Suominen T, et al. Nurses' ethical perceptions about coercion. *J Psychiatr Ment Health Nurs* 2004; 11: 379-385. 2004/07/17. DOI: 10.1111/j.1365-2850.2004.00715.x.
82. Pelto-Piri V, Kjellin L, Hylén U, et al. Different forms of informal coercion in psychiatry: a qualitative study. *BMC Research Notes* 2019; 12: 787. DOI: 10.1186/s13104-019-4823-x.
83. Hotzy F and Jaeger M. Clinical Relevance of Informal Coercion in Psychiatric Treatment-A Systematic Review. *Front Psychiatry* 2016; 7: 197-197. DOI: 10.3389/fpsy.2016.00197.
84. Shalowitz DI, Garrett-Mayer E and Wendler D. How should treatment decisions be made for incapacitated patients, and why? *PLoS Med* 2007; 4: e35. 2007/03/29. DOI: 10.1371/journal.pmed.0040035.
85. Kala AK. Covert medication; the last option: A case for taking it out of the closet and using it selectively. *Indian J Psychiatry* 2012; 54: 257-265. DOI: 10.4103/0019-5545.102427.
86. Sheldon CT. Proof in the Pudding: The Value of a Rights Based Approach to Understanding the Covert Administration of Psychotropic Medication to Adult Inpatients Determined to Be Decisionally-Incapable in Ontario's Psychiatric Settings. *The Journal of Law, Medicine & Ethics* 2017; 45: 170-181. DOI: 10.1177/1073110517720646.
87. Chiovitti RF. Theory of protective empowering for balancing patient safety and choices. *Nurs Ethics* 2011; 18: 88-101. 2011/02/03. DOI: 10.1177/0969733010386169.

88. Vaismoradi M, Jordan S, Logan PA, et al. A Systematic Review of the Legal Considerations Surrounding Medicines Management. *Medicina (Kaunas)* 2021; 57 2021/01/17. DOI: 10.3390/medicina57010065.
89. Roberts M. Psychiatric ethics; a critical introduction for mental health nurses. *J Psychiatr Ment Health Nurs* 2004; 11: 583-588. 2004/09/29. DOI: 10.1111/j.1365-2850.2004.00764.x.
90. Balevre P. Is it legal to be crazy: an ethical dilemma. *Arch Psychiatr Nurs* 2001; 15: 241-244. 2001/10/05. DOI: 10.1053/apnu.2001.27021.
91. Myers S. Seclusion: a last resort measure. *Perspect Psychiatr Care* 1990; 26: 24-28. 1990/01/01. DOI: 10.1111/j.1744-6163.1990.tb00313.x.
92. Laurant M, van der Biezen M, Wijers N, et al. Nurses as substitutes for doctors in primary care. *Cochrane Database Syst Rev* 2018; 7: Cd001271. 2018/07/17. DOI: 10.1002/14651858.CD001271.pub3.
93. Mardani A, Griffiths P and Vaismoradi M. The Role of the Nurse in the Management of Medicines During Transitional Care: A Systematic Review. *J Multidiscip Healthc* 2020; 13: 1347-1361. DOI: 10.2147/JMDH.S276061.
94. McDermott-Levy R, Leffers J and Mayaka J. Ethical Principles and Guidelines of Global Health Nursing Practice. *Nurs Outlook* 2018; 66: 473-481. 2018/09/13. DOI: 10.1016/j.outlook.2018.06.013.