

Advances in *Glomeromycota* taxonomy and classification

Fritz Oehl¹, Ewald Sieverding², Javier Palenzuela³, Kurt Ineichen⁴, and Gladstone Alves da Silva⁵

¹Agroscope Reckenholz-Tänikon Research Station (ART), Ecological Farming Systems, Reckenholzstrasse 191, CH-8046 Zürich, Switzerland; corresponding author e-mail: fritz.oehl@art.admin.ch

²Institute of Plant Production and Agroecology in the Tropics and Subtropics, University of Hohenheim, Garbenstrasse 13, D-70593 Stuttgart, Germany

³Departamento de Microbiología del Suelo y Sistemas Simbióticos, Estación Experimental del Zaidín, CSIC, Profesor Albareda 1, 18008 Granada, Spain

⁴Basel-Zürich Plant Science Center, Institute of Botany, University of Basel, Hebelstrasse 1, CH-4056 Basel, Switzerland

⁵Departamento de Micología, CCB, Universidade Federal de Pernambuco, Av. Prof. Nelson Chaves s/n, Cidade Universitaria, 50670-420, Recife, PE, Brazil

Abstract: Concomitant morphological and molecular analyses have led to major breakthroughs in the taxonomic organization of the phylum *Glomeromycota*. Fungi in this phylum are known to form arbuscular mycorrhiza, and so far three classes, five orders, 14 families and 29 genera have been described. *Sensu lato*, spore formation in 10 of the arbuscular mycorrhiza-forming genera is exclusively glomoid, one is gigasporoid, seven are scutellosporoid, four are entrophosphoroid, two are acaulosporoid, and one is pacisporoid. Spore bimorphism is found in three genera, and one genus is associated with cyanobacteria. Here we present the current classification developed in several recent publications and provide a summary to facilitate the identification of taxa from genus to class level.

Key words:

Archaeosporomycetes
endomycorrhizas
evolution
Gigasporales
Glomerales
Glomeromycetes
Paraglomeromycetes
phylogeny
VA mycorrhiza

Article info: Submitted 10 November 2011; Accepted: 16 November 2011; Published: 18 November 2011.

INTRODUCTION

Glomeromycota taxonomy was largely morphologically driven up to the end of the last millennium. All glomeromycotean fungi, except one genus, are known to form arbuscular mycorrhiza. Their identification was based on spore morphology, spore formation, and spore wall structure (e.g. Gerdemann & Trappe 1974, Walker & Sanders 1986, Morton & Benny 1990, Schenck & Pérez 1990). However, as soon as molecular phylogenetic tools became available, they were included in taxonomic analyses (e.g. Simon *et al.* 1992) and soon became the drivers of the establishment of a new taxonomy (Morton & Redecker 2001, Schüßler *et al.* 2001). In 1990, without the benefit of molecular aspects, the arbuscular mycorrhiza-forming fungi were organized in three families (*Acaulosporaceae*, *Gigasporaceae*, and *Glomeraceae*) and six genera (*Acaulospore*, *Entrophospora*, *Gigaspora*, *Glomus*, *Sclerocystis*, and *Scutellospora*) within one order, *Glomerales* (Morton & Benny 1990) of the fungal phylum *Zygomycota*. That classification was based on spore morphology and spore formation characteristics (acaulosporoid, entrophosphoroid, gigasporoid, glomoid,

radial-globose, and scutellosporoid). Differences in spore wall structure were used at the species level.

Today, we accept three classes (*Archaeosporomycetes*, *Glomeromycetes*, and *Paraglomeromycetes*), five orders (*Archaeosporales*, *Diversisporales*, *Gigasporales*, *Glomerales* and *Paraglomerales*), 14 families, 29 genera and approximately 230 species (e.g. Morton & Redecker 2001, Schüßler *et al.* 2001, Oehl & Sieverding 2004, Walker & Schüßler 2004, Sieverding & Oehl 2006, Spain *et al.* 2006, Oehl *et al.* 2008, 2011a-d, Palenzuela *et al.* 2008).

Until recently, it was unclear whether globose and gigasporoid species could be further divided into different morphological groups congruent with the major phylogenetic clades obtained by molecular analyses. A first revision of the sporogenous cell forming (gigasporoid and scutellosporoid) *Glomeromycetes* according to concomitant morphological and phylogenetic features (Oehl *et al.* 2008) was not accepted by all mycologists (Morton & Msiska 2010). However, later studies with a broader database (e.g. Goto *et al.* 2010, 2011, Oehl *et al.* 2010, 2011b) confirmed that the revised genus *Scutellospora*, as well as the new *Racocetra*, *Cetraspora*, *Dentiscutata*, and *Orbispora*, are monophyletic.

© 2011 International Mycological Association

You are free to share - to copy, distribute and transmit the work, under the following conditions:

Attribution: You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).

Non-commercial: You may not use this work for commercial purposes.

No derivative works: You may not alter, transform, or build upon this work.

For any reuse or distribution, you must make clear to others the license terms of this work, which can be found at <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>. Any of the above conditions can be waived if you get permission from the copyright holder. Nothing in this license impairs or restricts the author's moral rights.

A large group of species forms glomoid spores, and it had been believed that there were too few morphological characters of significance to differentiate them. Taxonomists have consequently started basing groupings of the glomoid species almost exclusively on molecular phylogenetic characters. A recent revision of these glomoid species has, however, shown that molecular phylogeny is actually congruent with the morphological characteristics of these fungi (Oehl et al. 2011c). Fungal species with entrophosphoroid spore formation were also revised (Oehl et al. 2011d). The objective of this paper is to present the current overall classification system of *Glomeromycota* that has emerged from these recent studies, and to summarize the major morphological features in the phylum down to genus level.

MATERIALS AND METHODS

The morphological, molecular, and phylogenetic analyses performed are presented in a series of recent publications dealing with different species groups of *Glomeromycota* (e.g. Oehl et al. 2006, 2010, 2011a, b, d, f, Sieverding & Oehl 2006, Silva et al. 2006, Spain et al. 2006, Palenzuela et al. 2008, 2010, 2011).

RESULTS AND DISCUSSION

Figure 1 is a schematic tree for *Glomeromycota* based on molecular phylogenetic analyses of the SSU, ITS region, partial LSU of the rRNA gene, and partial β -tubulin gene (e.g. Oehl et al. 2008, 2010, 2011a–d). In Table 1, the major morphological features of all higher level taxa are presented, with the taxa arranged according to their taxonomic rank down to genus. Three glomeromycetean classes, five orders, 14 families, and 29 genera have been recognized to date (Table 1). *Sensu lato*, spore formation in 10 of the arbuscular mycorrhiza-forming genera have exclusively glomoid, one has gigasporoid, seven have scutellosporoid, four have entrophosphoroid, two genera have acaulosporoid, and one has pacisporoid spore formation, while three genera show spore bimorphism, and one genus is associated with cyanobacteria (the only one not forming arbuscular mycorrhizas).

Hitherto, *Paraglomeromycetes* are monogeneric (Table 1), are characterized by mono-walled spores formed terminally on hyphae (i.e. glomoid spores *sensu lato*), and germinate directly through the spore wall. Their arbuscular mycorrhizal structures do not or only faintly stain in trypan blue. *Archaeosporomycetes* includes organisms that are exclusively bimorphic since they form either acaulosporoid or entrophosphoroid spores simultaneously with glomoid spores, or are associated with cyanobacteria. The mycorrhizal structures of *Archaeosporaceae* are similar to those of *Paraglomeraceae*, while *Ambisporaceae* form vesicular-arbuscular mycorrhizal structures staining pale blue in trypan blue. In contrast, mycorrhizal structures in *Glomeromycetes*

stain blue to dark blue in trypan blue. In *Glomeromycetes*, *Gigasporales* species do not form intraradical vesicles but auxiliary cells in soils, which clearly distinguish them from *Glomerales* and *Diversisporales*.

Gigasporales exhibit gigasporoid or scutellosporoid spore formation (Oehl et al. 2011b), i.e. spores formed terminally on sporogenous cells and with either germ warts on the inner surface of the mono-walled spore wall (gigasporoid; *Gigasporaceae*), or a discrete germination shield on the innermost (= ‘germinal wall’) of 2–4 walls (scutellosporoid). There are three families with scutellosporoid spore formation (*sensu lato*): *Dentiscutataceae*, *Racocetraceae* and *Scutellosporaceae* (Oehl et al. 2008). *Scutellosporaceae* form mono-lobed (*Orbispora*) or bi-lobed (*Scutellospora*), hyaline germination shields (Figs 2–4). *Racocetraceae* species form wavy-like, multiply lobed, hyaline germination shields and have either two (*Racocetra*) or three (*Cetraspora*) spore walls (Figs 5–8). *Dentiscutataceae* species form yellow-brown to brown germ shields that are bi-lobed (*Fuscotata*; Fig. 9) or with multiple compartments (*Dentiscutata*, triple-walled; *Quatunica* four-walled; Figs 10–11).

In *Archaeosporales* and *Diversisporales*, four genera have spore formation laterally on the neck of terminal or intercalary sporiferous saccules (= acaulosporoid *sensu lato*; Table 1): *Acaulospora*, *Otospora*, and the bi-morphic *Ambispora* and *Archaeospora*. These genera can easily be separated on spore wall number and spore wall structure (Palenzuela et al. 2008). Triple-walled *Acaulospora* species have a characteristic granular, ‘beaded’ inner wall surface (Morton & Benny 1990), which is absent in acaulo-ambisporoid spores of triple-walled *Ambispora* species (Spain et al. 2006, Palenzuela et al. 2011). The wall structure of the bi-walled *Otospora* is more complex than that of bi-walled *Archaeospora* species (Palenzuela et al. 2008).

In *Archaeosporales*, *Diversisporales*, and *Glomerales*, there are five genera with spore formation within the neck of terminal or intercalary sporiferous saccules (i.e. entrophosphoroid *sensu lato*; Table 1): *Entrophospora*, *Kuklospora*, *Sacculospora*, *Tricispore*, and bimorphic *Intraspora* (Oehl et al. 2011d). Triple-walled *Kuklospora* has the characteristic granular, ‘beaded’ inner wall surface of *Acaulosporaceae* (Sieverding & Oehl 2006), which is absent in spores of triple-walled *Sacculospora* (Oehl et al. 2011d). The wall structure of bi-walled *Entrophospora* and *Tricispore* is more complex than that of bi-walled, bimorphic *Intraspora* species (Sieverding & Oehl 2006, Oehl et al. 2011d). *Entrophospora* and *Tricispore* can be distinguished through the two cicatrices (scars) and pore structures proximal and distal to the sporiferous saccule: the proximal pore is wide in *Tricispore* and closed by a septum, while it is narrow and closed by a plug in *Entrophospora*. The distal pore and scar is absent in *Entrophospora* from the structural layer, and formed only on the overlying, hyaline, evanescent layer, while, in light microscopy, the distal pore with a distal scar is obvious in *Tricispore* (Sieverding & Oehl 2006, Palenzuela et al. 2010, Oehl et al. 2011d).





Fig. 1. Representative tree of the phylum *Glomeromycota* based on molecular (SSU, ITS region, partial LSU of the rRNA gene, and partial β -tubulin gene) and morphological analyses (spore wall structures, structures of the spore bases and subtending hyphae, germination, and germination shield structures). Adapted from (Oehl *et al.* 2008, 2011a–d). The drawings in the central columns show the spore formation types of the genera, and the typical germination shields for those genera which form persistent shields already during spore formation.


Table 1. Major morphological characters for higher level taxa of Glomeromycota from class to genus level.

| Class | Family | Genus | Spore formation | Number of spore walls | Germination; specific germination structure | Mycorrhizal structures; staining in Trypan blue |
|-----------------|-------------------|------------------------|--|-----------------------|---|---|
| Glomeromycetes | | | | | | |
| Glomerales | Glomeraceae | <i>Glomus</i> | Glomoid (terminally on hyphae) | 1 | gt through hypha | Vesicles, Hyphae |
| | | <i>Funneliformis</i> | Glomoid Funneliformoid sensu stricto | 1 | gt through hypha | V, A, H |
| | | <i>Septoglomus</i> | Glomoid Septoglooid sensu stricto | 1 | gt through hyphae | V, A, H |
| | | <i>Simiglomus</i> | Glomoid Simiglomoid sensu stricto | 1 | gt through hypha? | V, A, H |
| | Entrophosporaceae | <i>Claroideoglomus</i> | Glomoid sensu lato Claroideoglomoid sensu stricto | 1 | gt through hypha | V, A, H |
| | | <i>Albahypha</i> | Glomoid Claroideoglomoid sensu lato | 1 | gt through hypha? | V, A, H |
| | | <i>Viscospora</i> | Glomoid Claroideoglomoid sensu lato | 1 | gt through hypha | V, A, H |
| | | <i>Entrophospora</i> | Entrophosporoid (in the neck of a saccule) | 2 | gt through wall? | V, A, H |
| Diversisporales | Diversisporaceae | <i>Diversispora</i> | Glomoid Diversisporoid sensu stricto | 1 | gt through hypha | V, A, H |
| | | <i>Redeckeria</i> | Glomoid (Diversisporo-)Redeckeroid sensu stricto | 1 | gt through hypha? | V, A, H |
| | | <i>Otospora</i> | Acaulosporoid (on the neck of sporiferous saccule); otosporoid sensu stricto | 2 | Unknown? | V, A, H |
| | | <i>Tricispora</i> | Entrophosporoid Tricisporoid sensu stricto | 2 | Unknown? | V, A, H |
| | Sacculosporaceae | <i>Sacculospora</i> | Entrophosporoid Sacculosporoid sensu stricto | 3 | gt through wall; multiply lobed germ structure | V, A, H |
| | Pacioporaceae | <i>Pacispora</i> | Pacisporoid | 2 | gt through wall; mono-lobed, hyaline germ shield (=orb) | V, A, H |
| | Acaulosporaceae | <i>Kuklospora</i> | Entrophosporoid Kuklosporoid sensu stricto | 3 | gt through wall; mono-(to multiply) lobed, hyaline germ shield (=orb) | V, A, H |
| | | <i>Acaulospora</i> | Acaulosporoid | 3 | gt through wall; mono-(to multiply) lobed, hyaline germ shield (=orb) | V, A, H |

| | | | | | | | |
|---------------------------|---------------------------|-----------------------|--|------------------|---|-------------------------------|--|
| Gigasporales | Scutellosporaceae | <i>Orbispora</i> | Scutellosporoid (on sporogenous cells, and forming germ shields); <i>Orbisporoid sensu stricto</i> | 3 | gt through wall; mono-lobed, hyaline germ shield (=orb) | A, H | |
| | | <i>Scutellospora</i> | Scutellosporoid | 3 | gt through wall; bi-lobed, hyaline, violin-shaped germ shield | A, H | |
| | Dentiscutataceae | <i>Fuscata</i> | Scutellosporoid <i>Fuscataoid sensu stricto</i> | 3 | gt through wall; bi-lobed, brown, oval shield | A, H | |
| | | <i>Dentiscutata</i> | Scutellosporoid <i>Dentiscutataoid sensu stricto</i> | 3 | gt through wall; brown germ shield with multiple small compartments | A, H | |
| | | <i>Quatunica</i> | Scutellosporoid <i>Dentiscutataoid sensu stricto</i> | 4 | gt through wall; brown germ shield with multiple compartments | A, H | |
| | Racocetraeae | <i>Cetraspora</i> | Scutellosporoid <i>Racocetroid sensu stricto</i> | 3 | gt through wall; multiply lobed, hyaline germ shield | A, H | |
| | | <i>Racocetra</i> | Scutellosporoid <i>Racocetroid sensu stricto</i> | 2 | gt through wall; multiply lobed, hyaline germ shield | A, H | |
| | | <i>Gigasporaceae</i> | Gigasporoid (on sporogenous cells, and forming germ warts) | 1 | gt through wall; germ warts on inner spore wall layer | A, H | |
| | Archaeosporomyctes | | | | | | |
| | Ambisporales | <i>Ambispora</i> | Bimorph: Acaulo- & Glomo-ambisporoid | 3 (Ac) 1 (Gl) | multiply-lobed germ structure (Ac) & gt through hypha (Gl) | V, A, H | |
| | Archaeosporaceae | <i>Archaeospora</i> | Bimorph: Acaulo- & Glomo-archaeosporoid | 2 (Ac) 1 (Gl) | gt through wall; germ trunk (Ac), & gt through hypha (Gl)? | A, H | |
| | | <i>Infraspora</i> | Bimorph: Entropho- & Glomo-infrasporoid | 2 (Ac) 1 (Gl) | Unknown? | A, H | |
| | | <i>Geosiphonaceae</i> | Geosiphon | 1 | gt through hypha? | Associated with cyanobacteria | |
| Paraglomeromycetes | | | | | | | |
| Paraglomerales | Paraglomeraceae | <i>Paraglomus</i> | | | gt through wall | A, H | |



Figs 2–11. Characteristic germination shields in *Gigasporales* with germ pore (*gp*) as connection between spore cell contents and shields that are positioned on the surface of the germinal wall; germ tubes emerge from germ tube initiations (*gti*). **Fig. 2.** *Orbispora pernambucana* (isotype, ZT Myc 641) with mono-lobed, hyaline germ shield (*orb*). **Figs 3–4.** *Scutellospora calospora* (photo taken at INVAM) and *S. dipurpurescens* (holotype OSC #83343) have bi-lobed, violin-shaped, hyaline shields. **Figs 5–8.** *Racocetra coralloidea* (type, OSC #31026), *R. castanea* (ex type, ZT Myc 4377), *Cetrasporella nodosa* (isotype, DPP, Szczecin, Poland) and *C. helvetica* (isotype, ZT Myc 3038) have wavy-like, multiply lobed, hyaline shields. **Figs 9–11.** *Dentiscutataceae* shields are yellow brown to brown. **Fig. 9.** *Dentiscutata reticulata* (photo taken at INVAM) shields with multiple small compartments. **Fig. 10.** *Quatunica erythropo* (photo taken at INVAM) is assumed to be the only known species in *Glomeromycota* with four spore walls. **Fig. 11.** *Fuscotula heterogama* (ex type, ZT Myc 642) has a bi-lobed, oval to ovoid shield.



Figs 12–21. Characteristic spore bases and subtending hyphae (*sh*) in *Glomeromycetes* genera with glomoid spore formation. **Figs 12–13.** *Glomus ambisporum* (Oehl collection, from Bolivia) and *G. aureum* (type, ZT Myc 822) with two wall layers (SWL1 and SWL2), marked introverted wall thickening at *sb* and in *sh*, and a small, bridging septum (*sp*). **Fig. 14.** *Funneliformis coronatus* (ex type, Oehl collection) with funnel-shaped *sh* and conspicuous *sp*; introverted wall thickening is lacking. **Fig. 15.** *Septoglomus constrictum* (Oehl collection, from Switzerland) with conspicuous septum that sometimes resembles a plug. **Fig. 16.** *Simiglomus hoi* (Oehl collection, specimen mounted at York university) with cylindrical *sh*; *sh* wall thickened over long distances; several septae are regularly observed within the hyphae; no introverted wall thickening at *sb*, pore at *sb* generally opened. **Fig. 17.** *Claroideoglomus etunicatum* (Oehl collection, from Bolivia) with funnel/bill-shaped, white *sh*; all *Entrophosporaceae* (syn. *Claroideoglomeraceae*) with characteristic color change of structural wall layer at *sb*, if spores are pigmented. **Fig. 18.** *Albahypha drummondii* (type, DPP) with slightly funnel-shaped, white *sh*. **Fig. 19.** *Viscospora viscosa* (ex type, photo taken at INVAM) with cylindrical, white hyphae; *sp* within *sh* in some distance of *sb*; introverted wall thickening of *sh* at *sp* position, here not that obvious as usually found for this species; viscose spore surface. **Fig. 20.** *Diversispora versiformis* with short, fragile *sh* that is principally continuous with semi-persistent outermost spore wall layer (SWL1) but not with structural layer SWL2 (Oehl collection, from Tibet). **Fig. 21.** *Redeckera fulva* (Oehl ex Trappe collection) with inflating *sh* and conspicuous broad *sp* exactly at spore base.

In *Diversisporales* and *Glomerales*, 10 genera exclusively differentiate mono-walled, glomoid (9) or bi-walled pacisporoid (1) spores, all formed on subtending hyphae (Oehl & Sieverding 2004, Oehl *et al.* 2011a). The morphological differentiation of the glomoid species is mainly based on the morphology of the subtending hyphae of the spores, and spore wall structure. Spores of *Funneliformis*, *Glomus*, *Septoglomus*, and *Simiglomus* species have subtending hyphae that are concolorous or slightly

lighter in colour than the spore wall (Table 1, Figs 12–16). *Albahypha*, *Claroideoglomus*, and *Viscospora* form spores in which the structural wall layer is continuous with the subtending hyphal wall layer, but the subtending hyphae are hyaline (Figs 17–19). In contrast, *Diversispora* and *Redeckera* form spores whose structural wall layer is not obviously continuous with the hyphal wall layer (Figs 20–21); consequently, such spores appear to have included 'endospores'.

Funneliformis, *Glomus*, *Septoglomus*, and *Simiglomus* can be separated by the structure of the spore base and subtending hyphae (*sh*). *Glomus* species often have an introverted wall thickening (Oehl et al. 2011a; Figs 12–13) which is only otherwise seen in *Viscospora*. *Funneliformis* species generally have an easily visible septum in the area of the spore base, and their *sh* are regularly funnel-shaped to cylindrical (Fig. 14). *Septoglomus* species have constricted to cylindrical *sh*, and usually there is a septum at the spore base (Fig. 15). In *Simiglomus*, *sh* are cylindrical and thick-walled, and they have several septa some distance from the spore base (Fig. 16). *Claroideoglomus* has funnel- to bird-bill-shaped *sh*, with *sh* and *sh* walls that are > 2.5 times wider at the spore base than some distance from the base (Fig. 17). *Albahypha* has slightly funnel to bill-shaped *sh* and *sh* walls that are < 2.0 times wider at the spore base than at some distance from the base (Fig. 18), and *Viscospora* has cylindrical *sh* (Fig. 19) with an *sh* wall that may be thickened over large distances and may bear septa in the hyphae with introverted wall thickenings in the area of the septum. In *Diversispora*, the *sh* are usually quite fragile and hyaline, distal to the pore closure at the spore base or in the *sh* (Fig. 20). *Redeckera* species have a broad septum at the spore base (Fig. 21), and the structural wall layer does not continue more than 5–15 µm into the subtending hypha, and thus, the *sh* may inflate at this distance from the spore base.

There are three bi-morphic genera with glomoid spore formation. Glomo-ambisporoid spores have a subhyaline to ochraceous, evanescent outer wall layer continuous with the outer acaulo-ambisporoid spore wall, while the second, structural layer is hyaline and continuous with the middle wall of acaulo-ambisporoid spores (Spain et al. 2006, Palenzuela et al. 2011). Glomo-archaeosporoid and Glomo-intrasporoid spores are among the smallest within *Glomeromycota* (ca. 30 µm), and thus difficult to observe.

PERSPECTIVES

Further separations of genera and families can be expected in the near future since many species and several species groups have not yet been analyzed by molecular phylogenetic methods (e.g. *Glomus* group Ab1, *sensu* Oehl et al. 2011a). Major efforts are needed to properly describe the morphology of, in particular, small-spored *Glomus* species (Błaszkowski et al. 2009a, b, 2010a, b), and it is difficult to predict how morphological identification will develop in those fungi. Other recent progress has been made on *Acaulospora* species with pitted surface ornamentation, where several species, that superficially all resembled *A. scrobiculata*, have been separated through extensive morphological and molecular spore analyses (e.g. Oehl et al. 2006, 2011e, f). The establishment of international and national collections of arbuscular mycorrhizal fungi, such as INVAM in Morgantown (International Culture Collection of (Vesicular) Arbuscular Mycorrhizal Fungi, West Virginia State University, USA), CICG in Blumenau (International Collection of *Glomeromycota* at

FURB, Santa Catarina State, Brazil), GINCO-BEL in Louvain-La-Neuve (*Glomeromycota* In Vitro Collection at the Catholic University of Louvain, Belgium), or SAF in Zurich (Swiss Collection of Arbuscular Mycorrhizal Fungi at Agroscope ART, Switzerland) will facilitate further progresses in the taxonomy of glomeromycetean fungi that were thought to have not enough criteria to morphologically separate them unequivocally into the higher level taxa they phylogenetically belong to. Currently, several arbuscular mycorrhizal fungi are being described as new to science each year by an increasing numbers of research groups. A simple, but well justified conclusion is that, as a result of future concomitant morphological and molecular analyses, yet more higher level taxa will be proposed in this ancient fungal phylum, at all levels from class down to genus.

ACKNOWLEDGEMENTS

We acknowledge the help of many current and former students and technicians in Switzerland, Spain, and Brazil for outstanding support (especially David Schneider, Robert Bösch, Giacomo Busco, Louis Lawouin, Domingo Alvarez, Danielle Silva, Natália Sousa, and Daniele Magna). This study has been supported within the Swiss National Science Foundation (SNSF) Project 315230_130764/1, by the SNSF Programme NFP48 'Landscapes and habitats of the Alps', by the Spanish Ministry of Environment (MMA-OAPN, project 70/2005) and FAECA (Junta de Andalucía, Spain, Project 92162/11), and by the Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco (FACEPE) and the UFPE which provided grants to F. Oehl as 'visiting professor', and FACEPE which also provided financial support for G.A. Silva.

REFERENCES

- Błaszkowski J, Kovács GM, Balázs TK (2009a) *Glomus perpusillum*, a new arbuscular mycorrhizal fungus. *Mycologia* **101**: 247–255.
- Błaszkowski J, Ryszka P, Oehl F, Koegel S, Wiemken A, et al. (2009b) *Glomus achrum* and *G. bistratum*, two new species of arbuscular mycorrhizal fungi (*Glomeromycota*). *Botany* **87**: 260–271.
- Błaszkowski J, Wubet T, Harikumar VS, Ryszka P, Buscot F (2010a) *Glomus indicum*, a new arbuscular mycorrhizal fungus. *Botany* **88**: 132–143.
- Błaszkowski J, Kovács GM, Balázs TK, Orlowska E, Sadravi M, et al. (2010b) *Glomus africanum* and *G. iranicum*, two new species of arbuscular mycorrhizal fungi (*Glomeromycota*). *Mycologia* **102**: 1450–1462.
- Gerdemann JW, Trappe JM (1974) The *Endogonaceae* of the Pacific Northwest. *Mycologia Memoir* **5**: 1–76.
- Goto BT, Silva GA, Maia LC, Oehl F (2010) *Dentiscutata colliculosa*, a new species in the *Glomeromycetes* from Northeastern Brazil with colliculate spore ornamentation. *Nova Hedwigia* **90**: 383–393.
- Goto BT, Silva GA, Maia LC, Souza RG, Coyne D, et al. (2011) *Racocetra tropicana*, a new species in the *Glomeromycetes* from tropical areas. *Nova Hedwigia* **92**: 69–82.

- Morton JB, Benny GL (1990) Revised classification of arbuscular mycorrhizal fungi (*Zygomycetes*): a new order, *Glomales*, two new suborders, *Glomineae* and *Gigasporineae*, and two families, *Acaulosporaceae* and *Gigasporaceae*, with an emendation of *Glomaceae*. *Mycotaxon* **37**: 471–491.
- Morton JB, Redecker D (2001) Two new families of *Glomales*, *Archaeosporaceae* and *Paraglomaceae*, with two new genera *Archaeospora* and *Paraglomus*, based on concordant molecular and morphological characters. *Mycologia* **93**: 181–195.
- Morton JB, Msiska Z (2010) Phylogenies from genetic and morphological characters do not support a revision of *Gigasporaceae* (*Glomeromycota*) into four families and five genera. *Mycorrhiza* **20**: 483–496.
- Oehl F, Sieverding E (2004) *Pacispora*, a new vesicular-arbuscular mycorrhizal fungal genus in the *Glomeromycetes*. *Journal of Applied Botany and Food Quality* **78**: 72–82.
- Oehl F, Sýkorová Z, Redecker D, Wiemken A, Sieverding E (2006) *Acaulospora alpina*, a new arbuscular mycorrhizal fungal species characteristic for high mountainous and alpine regions of the Swiss Alps. *Mycologia* **98**: 286–294.
- Oehl F, Souza FA, Sieverding E (2008) Revision of *Scutellospora* and description of five new genera and three new families in the arbuscular mycorrhiza-forming *Glomeromycetes*. *Mycotaxon* **106**: 311–360.
- Oehl F, Jansa J, Souza FA, Silva GA (2010) *Cetraspora helvetica*, a new ornamented species in the *Glomeromycetes* from Swiss agricultural fields. *Mycotaxon* **114**: 71–84.
- Oehl F, Silva GA, Goto BT, Sieverding E (2011a) *Glomeromycetes*: three new genera and glomoid species reorganized. *Mycotaxon* **116**: 75–120.
- Oehl F, Silva DKA, Maia LC, Sousa NMF, Vieira HEE, Silva GA (2011b) *Orbispora* gen. nov., ancestral in the *Scutellosporaceae* (*Glomeromycetes*). *Mycotaxon* **116**: 161–169.
- Oehl F, Silva GA, Goto BT, Maia LC, Sieverding E (2011c) *Glomeromycota*: two new classes and a new order. *Mycotaxon* **116**: 365–379.
- Oehl F, Silva GA, Sánchez-Castro I, Goto BT, Maia LC, Vieira HEE, Barea JM, Sieverding E, Palenzuela J (2011d) Revision of *Glomeromycetes* with entrophosphoroid and glomoid spore formation with three new genera. *Mycotaxon* **117**: 297–316.
- Oehl F, Sýkorová Z, Błaszkowski J, Sánchez-Castro I, Coyne D, et al. (2011e) *Acaulospora sieverdingii*, an ecologically diverse new fungus in the *Glomeromycota*, described from lowland temperate Europe and tropical West Africa. *Journal of Applied Botany and Food Quality* **84**: 47–53.
- Oehl F, Silva GA, Palenzuela J, Sánchez-Castro I, Castillo C, Sieverding E (2011f) *Acaulospora punctata*, a new fungal species in the *Glomeromycetes* from mountainous altitudes of the Swiss Alps and Chilean Andes. *Nova Hedwigia* **93**: 353–362.
- Palenzuela J, Ferrol N, Boller T, Azcón-Aguilar C, Oehl F (2008) *Otospora bareai*, a new fungal species in the *Glomeromycetes* from a dolomitic shrub-land in the National Park of Sierra de Baza (Granada, Spain). *Mycologia* **100**: 282–291.
- Palenzuela J, Barea JM, Ferrol N, Azcón-Aguilar C, Oehl F (2010) *Entrophospora nevadensis*, a new arbuscular mycorrhizal fungus, from Sierra Nevada National Park (southeastern Spain). *Mycologia* **102**: 624–632.
- Palenzuela J, Barea JM, Ferrol N, Oehl F (2011) *Ambispora granatensis*, a new arbuscular mycorrhizal fungus, associated with *Asparagus officinalis* in Andalucía (Spain). *Mycologia* **103**: 333–340.
- Schenck NC, Pérez Y (1990) *Manual for the Identification of VA Mycorrhizal Fungi*. 3rd edn: Gainesville, FL: Synergistic Publications.
- Schüßler A, Schwarzott D, Walker C (2001) A new fungal phylum, the *Glomeromycota*: phylogeny and evolution. *Mycological Research* **105**: 1413–1421.
- Sieverding E, Oehl F (2006) Revision of *Entrophospora* and description of *Kuklospora* and *Intraspora*, two new genera in the arbuscular mycorrhizal *Glomeromycetes*. *Journal of Applied Botany and Food Quality* **80**: 69–81.
- Silva GA, Lumini E, Maia LC, Bonfante P, Bianciotto V (2006) Phylogenetic analysis of *Glomeromycota* by partial LSU rDNA sequences. *Mycorrhiza* **16**: 183–189.
- Simon L, Lalonde M, Bruns TD (1992) Specific amplification of 18S fungal ribosomal genes from vesicular-arbuscular endomycorrhizal fungi colonizing roots. *Applied and Environmental Microbiology* **58**: 291–295.
- Spain JL, Sieverding E, Oehl F (2006) *Appendicispora*, a new genus in the arbuscular mycorrhizal-forming *Glomeromycetes*, with a discussion of the genus *Archaeospora*. *Mycotaxon* **97**: 163–182.
- Walker C, Sanders FE (1986) Taxonomic concepts in the *Endogonaceae*: III. The separation of *Scutellospora* gen. nov. from *Gigaspora* Gerd. and Trappe. *Mycotaxon* **27**: 169–182.
- Walker C, Schüßler A (2004) Nomenclatural clarifications and new taxa in the *Glomeromycota*. *Mycological Research* **108**: 979–982.