



Choking recovery: threats and opportunities in an expanding fish stock

Journal:	<i>Fish and Fisheries</i>
Manuscript ID:	FaF-13-Aug-OA-130.R1
Wiley - Manuscript type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Baudron, Alan; University of Aberdeen, Zoology Fernandes, Paul; University of Aberdeen, Zoology
Key terms:	fish stock recovery, fisheries management, choke species, European hake
Abstract:	<p>Many commercial fish stocks are beginning to recover under more sustainable exploitation regimes. In this study we document the temporal and spatial changes in one remarkable example of stock recovery: northern European hake (<i>Merluccius merluccius</i>). Analysing data from several scientific surveys, we document a dramatic increase in estimates of biomass between 2004 and 2011 throughout the larger area now occupied by the stock. The largest increase occurred in the North Sea, where hake have been largely absent for over 50 years. Spatio-temporally resolved commercial landings show that high densities occur in the North Sea only between April and September, suggesting a density-dependent seasonal habitat expansion to suitable temperature and depth conditions. These changes have implications for the management of the stock which are discussed. Notably, if discards are banned as part of management revisions, the relatively low quota for hake in the North Sea will be a limiting factor (the so-called "choke" species) which may result in a premature closure of the entire demersal mixed fishery in the North Sea, jeopardising many commercial fisheries in the region. This example of the unforeseen consequences of improved stewardship, highlight the need for a more holistic, regional and responsive approach to managing our marine ecosystems.</p>

1 **Choking recovery: threats and opportunities in an expanding fish stock**

2

3 Alternative title:

4 **From a recovering stock to a 'choke' species: the example of northern hake**

5

6 Alan Ronan Baudron and Paul George Fernandes

7

8 Zoology Department, Institute of Biological and Environmental Sciences, University of

9 Aberdeen, Aberdeen AB24 2TZ, UK

10

11 Corresponding author: Alan Ronan Baudron

12 Zoology Department, Institute of Biological and Environmental Sciences, University of

13 Aberdeen, Aberdeen AB24 2TZ, UK

14 +44 (0) 1224272648

15 alan.baudron@abdn.ac.uk

16

17 Running title:

18 **Recovering fish stock and choke species**

19

20 **Abstract**

21

22 Many commercial fish stocks are beginning to recover under more sustainable exploitation
23 regimes. In this study we document the temporal and spatial changes in one remarkable
24 example of stock recovery: northern European hake (*Merluccius merluccius*). Analysing data
25 from several scientific surveys, we document a dramatic increase in estimates of biomass
26 between 2004 and 2011 throughout the larger area now occupied by the stock. The largest
27 increase occurred in the North Sea, where hake have been largely absent for over 50 years.
28 Spatio-temporally resolved commercial landings show that high densities occur in the North
29 Sea only between April and September, suggesting a density-dependent seasonal habitat
30 expansion to suitable temperature and depth conditions. These changes have implications for
31 the management of the stock which are discussed. Notably, if discards are banned as part of
32 management revisions, the relatively low quota for hake in the North Sea will be a limiting
33 factor (the so-called “choke” species) which may result in a premature closure of the entire
34 demersal mixed fishery in the North Sea, jeopardising many commercial fisheries in the
35 region. This example of the unforeseen consequences of improved stewardship, highlight the
36 need for a more holistic, regional and responsive approach to managing our marine
37 ecosystems.

38

39

40 **Keywords:** fish stock recovery, fisheries management, choke species, European hake

41

42 **Table of contents**

43

44 Introduction

45 Methods

46 Data

47 Analyses

48 Results

49 Discussion

50 Acknowledgements

51 References

52

For Review Only

53 Introduction

54

55 Fisheries of the northeast Atlantic were heavily exploited throughout the second half of the
56 20th century, and many commercial fish stocks experienced a severe decline in biomass by
57 the early 2000s. As a result, the Common Fisheries Policy (CFP) which regulates fisheries in
58 European waters undertook a major reform in 2002 (Daw and Gray 2005) and is currently
59 under further reform (EC 2013). The 2002 reform aimed to reduce fishing pressure on over-
60 exploited stocks by introducing recovery plans to allow depleted fish stocks to recover, and
61 long-term management plans to protect healthy stocks from depletion (Kraak *et al.* 2013).
62 These measures have contributed towards a substantial reduction in fishing pressure for most
63 northern European fish stocks in recent years (Cardinale *et al.* 2013) and a reversal of stock
64 decline, with prospects for recovery (Fernandes and Cook 2013). Among the stocks showing
65 signs of recovery, the northern stock of European hake (*Merluccius merluccius*, Merluccidae)
66 seems to have experienced one of the largest and fastest biomass increases over the last five
67 years. The latest assessment of this stock undertaken by the International Council for the
68 Exploration of the Sea (ICES) shows a dramatic increase in biomass since 2006, and the
69 spawning stock biomass (SSB) is now well above the recommended level (ICES 2012a).
70 Reported landings have consequently increased, especially for the northern part of the stock
71 (ICES 2012a).

72

73 European hake is a large demersal gadoid species found at depths between 70 and 200 m
74 (Kacher and Amara 2005), with a preference for depths between 70 and 100 m (Bartolino *et*
75 *al.* 2008). It is a Lusitanian species with a preferred temperature of 13.8 °C, plus or minus 2.9
76 °C (Wheeler 1969). European hake has the most extensive distribution of all gadoid species
77 in the northeast Atlantic and ranges from the tropical coast of Mauritania to the cooler waters

78 of Norway, expanding eastwards in the Mediterranean Sea, the North Sea, and the Skagerrak
79 and Kattegat (Casey and Pereriro 1995). In northeast Atlantic waters, European hake is
80 managed as two distinct stock units, a southern and northern component, separated by the
81 Cap Breton canyon in the Bay of Biscay (ICES 2012a). The northern stock ranges from the
82 southwest coast of France to Norway, covering ICES areas IIIa (Skagerrak and Kattegat), IV
83 (North Sea), VI (West of Scotland), VII (Celtic Sea) and VIIIa,b,d (Bay of Biscay) (ICES
84 2012a). The stock is, therefore, managed over an extensive area and regional assessments are
85 not carried out. Northern hake are surveyed by five scientific trawl surveys, conducted
86 annually, in the North Sea, the Celtic Sea, the West of Scotland, Irish waters, the Porcupine
87 Bank and the Bay of Biscay. These surveys provide estimates of relative abundance, as well
88 as length- and maturity-at-age estimates, used in the stock assessment. The SSB of this stock
89 peaked in 1980 at 101,917 t, but was then rapidly depleted to a historical low in 1998 at
90 24,603 t (ICES 2012a). As a result, an emergency plan was implemented in 2001 (EC 2001,
91 2002) which introduced a reduction in the Total Allowable Catch (TAC) as well as
92 stipulating a minimum mesh size (100 mm) in the cod-ends of trawl nets; a recovery plan
93 followed in 2004 (EC 2004). These multi-annual plans led to a reduction in the exploitation
94 rate and SSB has since increased to a new high of 131,075 t in 2010 (ICES 2012a).

95

96 Fish stocks experiencing an increase in abundance often show a concurrent increase in the
97 area they occupy, a mechanism known as density-dependent habitat selection (MacCall 1990;
98 Hinz *et al.* 2003; Hiddink *et al.* 2005). According to the ideal free distribution theory
99 (Shepherd and Litvak 2004), individuals expand to suitable habitats in order to avoid high
100 densities and maximise their fitness, a response which has been observed in gadoid stocks
101 (Marshall and Frank 1995). Given the large increase in biomass, an expansion of the area
102 occupied by the northern hake stock is likely to have occurred, providing that suitable

103 habitats are available in adjacent areas. Warming sea temperatures have also been linked to
104 changes in the distribution of fish stocks such as shifts towards the poles or deeper waters
105 (Hiddink and ter Hofstede 2008; Dulvy *et al.* 2008). In the northeast Atlantic several
106 examples of Lusitanian species expanding their distribution northward have been reported
107 such as European anchovy (*Engraulis encrasicolus*, Engraulidae), European pilchard
108 (*Sardina pilchardus*, Clupeidae), Atlantic horse mackerel (*Trachurus trachurus*, Carangidae)
109 and Atlantic mackerel (*Scomber scombrus*, Scombridae) and are now prevalent in new areas
110 (Beare *et al.* 2004; Petitgas *et al.* 2012). However, both a change in the area occupied and a
111 shift in distribution have yet to be documented for the northern stock of European hake.

112

113 Changes in the area occupied by a stock can result in changes to the potential catch and can
114 offer new fishing opportunities depending on the area and/or the species considered (Cheung
115 *et al.* 2012). Typical examples in the northeast Atlantic region include northward expansion
116 of the Atlantic mackerel (ICES 2013). Such changes are likely to affect fisheries regulation
117 and exploitation patterns, but also fish prices and economic performance of fishing fleets
118 (Cheung *et al.* 2012). The northern hake stock is of great economic importance (Alvarez
119 2004) especially for Spanish and French fleets which historically have accounted for 60%
120 and 25% of the landings respectively; fleets from the U.K., Denmark, Ireland, Norway,
121 Belgium, Netherlands, Germany and Sweden have contributed to the remaining 15% (ICES
122 2012a).

123

124 In the North Sea there is an additional management concern relating specifically to hake
125 which could affect the entire demersal fishery. The latest CFP reform (EC 2013) includes a
126 ban on discards due to be phased in over the next 5 years. Hake in the North Sea is caught in
127 a mixed demersal fishery including cod, haddock, whiting and other fish species. However

128 the TAC of hake in this area is very small compared to other species, simply because it was
129 not abundant when catch shares were being allocated. In the presence of a discard ban this
130 fishery will be closed once the smallest quota (hake) is taken and fishermen will not be able
131 to catch other species even though their quotas may not have been reached. This scenario is
132 commonly referred to in the USA as the “choke species” concept (Schrope 2010): a species
133 with the lowest quota in the mixed fishery “chokes” the opportunity to catch the quotas of
134 other species. In Europe this has been examined in mixed fishery models (Ulrich *et al.* 2011)
135 but hitherto the choke species in the North Sea is expected to be cod. It is, therefore,
136 important to assess the increase in regional biomass of hake and document the concomitant
137 changes in distribution which are likely to occur in order to better understand the dynamics of
138 this stock and any repercussion on fisheries. Improving our knowledge of the intra-stock
139 dynamics of northern hake is particularly important since this pan-European stock is assessed
140 over a large area.

141

142 In this study, fisheries independent data are analysed to document the regional increases in
143 the northern hake stock and investigate the potential implications for fisheries management.
144 Local abundance estimates are determined to investigate the distribution of the increasing
145 biomass and whether the area occupied by European hake has expanded amid the observed
146 increase in global biomass. Commercial landings data are used to compare seasonal changes
147 in distribution with local changes in sea temperature. Regional biomass estimates are
148 calculated to quantify the increase within ICES areas of the northern hake stock. Particular
149 attention is given to the North Sea where the TAC is much lower than in other areas of the
150 stock. Finally, the increase in biomass and expansion of spatial distribution is discussed in
151 relation to quota allocation, to highlight the opportunities and challenges presented by a
152 recovering stock in the context of the proposed management changes (CFP reform).

153

154

155 **Methods**

156

157 ***Data***

158 Data from scientific trawl surveys were obtained from ICES for the five regions covering the
159 northern stock area: North Sea, West of Scotland, Celtic Sea, Porcupine bank and Bay of
160 Biscay. Data for European hake were available in sufficient quantity over the period 1978-
161 2011 for the North Sea and West of Scotland, 1978-2008 for Ireland, 2001-2011 for the
162 Porcupine Bank and 1985-2010 for the Bay of Biscay. Additional data from Scottish (1978-
163 2007) and French (1985-2002) surveys were obtained from Marine Scotland Science (MSS)
164 and IFREMER, respectively. Available maturity-at-length data were extracted from the same
165 database. Northern hake stock estimates of total stock biomass (TSB), SSB, fishing mortality
166 and recruitment were taken from the latest stock assessment report (ICES 2012a). North Sea
167 landings of European hake from 1903 to 2010 were obtained from the Food and Agriculture
168 Organisation. Landings of European hake into Scottish ports and the associated discards
169 values for 2011 were obtained from MSS (Alastair Pout, MSS, personal communication).
170 Monthly European hake landings per ICES statistical rectangle of Scottish (from 2005 to
171 2011) and Danish (from 2000 to 2011) fleets, the two main nations targeting European hake
172 in the North Sea (ICES 2012a), were obtained from MSS (Rui Catarino, MSS, personal
173 communication) and the Danish National Institute of Aquatic Resources (Henrik Degel, DTU
174 aqua, personal communication) in order to perform spatial analyses. Monthly sea surface
175 temperature (SST) values in °C were obtained from the British Atmospheric Data Centre for
176 the northern hake stock area. SST values were available on a 1° latitude by 1° longitude grid

177 extrapolated from the Met Office Hadley Centre data (Rayner 2003). Bathymetry data for the
 178 North Sea were obtained from the National Oceanic and Atmospheric Administration.

179

180 *Analyses*

181 Data for each trawl haul included: date, longitude and latitude, depth, distance covered (a
 182 measure of the maximum length sampled by the trawl), haul duration, speed over ground,
 183 door spread (a measure of the maximum width sampled by the trawl) and numbers-at-length.
 184 For each survey, the number-at-length values were corrected to account for the differences in
 185 sampling procedures between surveys: whenever sub-sampling occurred, the values were
 186 adjusted by the corresponding raising factor. Data from each survey were then compiled in
 187 one standardised dataset for further analysis. The weight of hake in each haul was estimated
 188 by summing up the weight at each length-class as follows:

189

$$190 \quad (1) \quad w_h = (\sum_l^l (a * l^b) * n_l) / 1000$$

191

192 where w_h is the weight of hake caught in haul h in kg, l is the length-class in cm, n_l is the
 193 number of individuals in length class l in haul h , and a and b are the coefficients of the hake
 194 weight to length relationship obtained from Coull *et al.* (1989) with $a = 0.0047$ and $b = 3.099$.
 195 The relationship between depth and door spread was estimated by fitting the following
 196 equation to survey data for which both depth and door spread were recorded:

197

$$198 \quad (2) \quad ds = \frac{\alpha * dp}{\beta + dp}$$

199

200 where ds is the door spread, dp the depth and α and β are the estimated unitless coefficients.
 201 Equation 2 was then used to estimate missing door spread values. Missing distance values

202 were estimated by multiplying haul duration by speed over ground. The area sampled by each
203 haul was calculated by multiplying door spread (width) with distance (length). Using door
204 spread provides a conservative measure of density since it corresponds to the largest area
205 sampled: the estimates can, therefore, be considered as minimum values in the context of
206 whole gear selectivity which affect absolute abundance estimates. For each haul, densities in
207 weight (kg.km^{-2}) were determined by dividing the weight of fish caught by the area sampled.
208 These density estimates were plotted spatially by year and quarter from 2001 to 2011.
209 Average densities were also estimated for each survey, taking into account null observations
210 (i.e. hauls where no hake were caught).

211

212 In order to estimate the biomass in each ICES area included in the northern hake stock, the
213 time series of total biomass from each survey was estimated by multiplying the mean density
214 by the area covered by the corresponding survey. The total northern hake survey biomass
215 tsb_{TOT} was estimated by summing up the regional survey estimates tsb_{AREA} . The survey
216 catchability q was calculated as the ratio of the total northern hake survey biomass tsb_{TOT} to
217 the northern hake TSB estimates TSB from the reported stock assessments (ICES 2012a):
218 $q=tsb_{TOT}/TSB$. Regional TSB estimates TSB_{AREA} were then determined by multiplying the
219 survey biomass in each area tsb_{AREA} by the inverse of the survey catchability:
220 $TSB_{AREA}=tsb_{AREA} * q^{-1}$. A maturity ogive for the northern hake stock was fitted to the maturity-
221 at-length data available from ICES and the length at 50% maturity (L_{50}) was estimated at 31.2
222 cm. SSB was then estimated as the fraction of the TSB for which the length of individuals
223 was greater than L_{50} .

224

225 To investigate changes in the northern hake stock distribution area, a threshold was defined
226 as the maximum density determined in the North Sea prior to the implementation of the 2004

227 hake recovery plan. The percentage of ICES rectangles surveyed in which densities greater
228 than this threshold were determined was calculated for each year. This proxy of relative
229 spatial occupation differs from the positive area (the area where fish densities are strictly
230 positive) used by Woillez *et al.* (2009) to investigate fish distribution and was employed to
231 assess the magnitude of the change in the area occupied by hake. This proxy was estimated
232 for all surveys between 1985 and 2011 to ensure sufficient data in all regions with the
233 exception of Porcupine Bank where data were available from 2001 only. To assess changes in
234 regional distribution, the centre of gravity (Woillez *et al.* 2007) of the observed densities (i.e.
235 the mean location of the population) was calculated for each survey from 2001 to 2011. In
236 addition, the overall centre of gravity was calculated using only years sampled by all surveys
237 (2001-2008) and averaged both before (2001-2004), and after (2005-2008), the
238 implementation of the 2004 hake recovery plan.

239

240 To investigate changes in distribution in the North Sea throughout the year, Scottish and
241 Danish monthly hake landings per ICES statistical rectangles were aggregated over all
242 available years and plotted on monthly maps. Monthly mean SST values averaged across
243 2001-2011 as well as the 100 m depth contour were added to the maps to compare the
244 distribution of hake with these environmental variables. Generalised Additive Models
245 (GAMs) were employed to investigate statistical relationships between hake distribution and
246 the corresponding temperature and depth as follows:

247

$$248 \quad (3) \quad g(\text{landings}) = c + s(\text{SST}) + \text{Depth} + \varepsilon$$

249

250 where g is the Gaussian link function, c a constant, s a smoother, and ε a random error term.

251 Depth was set as a discrete variable with 50 m classes. GAMs were applied to the whole

252 dataset (2000-2011), and to landings recorded before (2000-2004) and after (2005-2011) the
253 2004 hake recovery plan to assess whether the increase in abundance affected the relationship
254 between distribution and environmental variables.

255

256

257 **Results**

258

259 Estimates of biomass from the northern hake stock assessment reveal that the biomass was
260 declining for most of the 1978-2010 period until a significant increase in recent years. TSB
261 (Fig. 1a) and SSB (Fig. 1b) time series exhibit a steady decline from the late 1970s to the late
262 1990s followed by a slight increase until the mid-2000s. From 2006 the biomass increased
263 dramatically, to reach a historical high in 2010 (Fig. 1a and 1b). Estimates of recruitment
264 declined steadily over the majority of the time series reaching a historical low in 2009, apart
265 from a slight increase in the mid-2000s concurrent with the start of the increase in TSB (Fig.
266 1a). Average estimates of fishing mortality show concurrent and opposite trends to SSB (Fig.
267 1b). After a rapid increase in the 1980s, F remained high until a sharp decrease occurred from
268 2005 to 2010, which was concurrent with the increase in SSB. Northern hake landings show
269 similar patterns to the biomass time series apart from a slight increase in the 1980s (Fig. 1c).
270 However, there are differences between the various areas occupied by the stock. Landings in
271 the Bay of Biscay declined by 50% between the 1980s and 2000s with a slight increase in
272 recent years, while landings from the Celtic Sea have remained constant. Landings from the
273 West of Scotland and North Sea, although much lower in comparison, have experienced a
274 proportionally larger increase in the last 5 years and are now half the size of the landings in
275 the Bay of Biscay and Celtic Sea, respectively (Fig. 1c).

276

277 Density estimates from trawl surveys showed a significant increase across the northern hake
278 stock area between 2001 and 2011, although there were differences between regions (Fig. 2).
279 From 2001 to 2004, the largest densities were observed on the western continental shelf
280 between the Bay of Biscay and the West of Scotland with the highest densities recorded on
281 the Porcupine Bank and West of Scotland (Fig. 2). Low densities were recorded in the
282 Skagerrak and Kattegat, and very few positive values were estimated in the North Sea (Fig.
283 2). In 2005, there was an increase in samples of low density in the northern North Sea around
284 Shetland, while densities in other regions remained similar (Fig. 2). From 2006 to 2011,
285 densities estimated in the northern North Sea exhibited a rapid increase reaching the largest
286 estimated values in 2010 and 2011 while densities in other regions increased steadily except
287 in Skagerrak and Kattegat (Fig. 2). Contrary to other regions, the densities observed in the
288 North Sea were much larger in quarter 3 (Q3) than in quarter 1 (Q1, Fig 2). Overall the mean
289 density quadrupled in all regions except the North Sea, where it quintupled.

290

291 The percentage of ICES rectangles with densities higher than the maximum estimated in the
292 North Sea prior to the implementation of the 2004 hake recovery plan (85.7 kg/km^2)
293 increased in all five regions demonstrating an expansion of the northern hake stock (Fig. 3a).
294 The increase in area of occupation was most pronounced from 2005 to 2011, with the largest
295 expansion observed in the North Sea, followed by West of Scotland (Fig. 3a). Summing up
296 the indicator of spatial occupation across all regions shows that the area occupied by northern
297 hake has quintupled over the last decade (Fig. 3a). Over this period, the centre of gravity of
298 the estimated densities within each survey has remained unchanged with the exception of the
299 North Sea where it shifted north-westward, from the northern tip of Denmark in 2001, to the
300 northern North Sea in 2011 (Fig. 3b). Densities in the northwest North Sea increased from
301 2005 onward, while densities in the Skagerrak and Kattegat remained low (Fig. 2), so the

302 shift in centre of gravity actually advocates for an eastward expansion into the North Sea. The
303 overall centre of gravity before and after the implementation of the 2004 hake recovery plan
304 shows a northeast displacement, supporting the hypothesis of an eastward expansion into the
305 North Sea (Fig. 3b).

306

307 In the North Sea the dichotomy between the high densities in Q3 and low densities in Q1
308 suggests that European hake are not present in the North Sea throughout the year (Fig. 2).
309 Spatially resolved landings data from the Scottish and Danish fishing fleets (the predominant
310 fleets in the area) show that from January to March (Q1) hake are mostly located west of
311 Shetland, along the 8°C temperature isotherm to the north and west of Scotland (Fig. 4). In
312 April, as sea temperature rose, landings increased in the northern North Sea around the 100 m
313 depth contour. From May to August, as landings increased, the distribution expanded
314 eastward into the North Sea along the 100 m depth contour as temperatures increased to their
315 maximum (Fig. 4). From September to December a reverse pattern was observed as the
316 landings distribution gradually retreated back to the North of Scotland as temperatures
317 declined. There was a significant relationship between the distribution of landings and
318 temperature and depth for all periods tested (Table 1). Landings increased with temperature
319 (between 5 and 17°C) and depth (until 150 m) and this pattern was observed both before and
320 after the 2004 recovery plan, although the relationship with depth was less clear for the 2000-
321 2004 period due to fewer data available (Fig. S1). The spatial monthly landings data infer that
322 hake undertake seasonal migrations as temperatures become suitable, expanding from the
323 North of Scotland into the North Sea and back out in the course of the year. This is supported
324 by the difference in survey density between Q1 and Q3. Landings were restricted mainly to
325 areas where waters were deeper than 100 m and none were from the southern North Sea

326 despite suitable temperatures, suggesting that migrations are driven by both temperature and
327 depth.

328

329 Regional SSB estimates for the period encompassing the increase in northern hake (2001-
330 2011) were calculated for the four ICES areas included in the stock for the first (Q1 and Q2)
331 and second (Q3 and Q4) half of the year. SSB increased in the second half of the 2000s in all
332 regions (Fig. 5a-d). The largest proportional increase, from 4,494 to 59,273 t, occurred in the
333 North Sea in the second half of the year (Fig. 5a). The highest SSBs were estimated in the
334 West of Scotland (84,177 t) (Fig. 5b) and Celtic Sea (86,282 t) in 2011 (Fig. 5c). A
335 dichotomy between the first and second half of the year was observed in both the North Sea,
336 where the increasing SSB in 2011 reached 59,273 t in the second half (c.f. 12,423 t in the first
337 half, Fig. 5a), and the West of Scotland, where SSB in 2011 was only 19,913 t in the second
338 half contrasting with 84,177 t estimated in the first half (Fig. 5b). This supports the
339 hypothesis that individuals migrate in and out of the North Sea in the year, and suggests that
340 much of the SSB present in the North Sea in the second half of the year comes from the West
341 of Scotland. SSB estimates suggest that the rise observed in the North Sea is unprecedented.
342 However, historical landings in the late 1940s and early 1950s ($\approx 7,800$ t) were similar to
343 those achieved nowadays (7,631 t in 2009), attesting that high biomasses previously occurred
344 in the North Sea (Fig. 5e).

345

346

347 **Discussion**

348

349 An increase in fish stock biomass can be associated with three factors, occurring
350 independently or in combination: higher recruitment success, faster body growth rate, or a

351 reduction in mortality, which in a heavily exploited stock is largely due to fishing (Hilborn
352 and Walters 1992). In the case of the northern hake stock, an increase in the recruitment
353 success resulting from improved environmental conditions between the late 1980s and mid-
354 2000s has been suggested (Goikoetxea and Irigoien 2013). However, declining recruitment
355 estimates towards historical low values in the late 2000s suggest that recruitment is unlikely
356 to be the cause of the recent increase observed in biomass. A possible increase in body
357 growth rate, albeit not explored in this study, is unlikely to generate an increase in biomass of
358 such a magnitude, particularly when associated with the changes in distribution observed
359 here, although it could have contributed. Trends in fishing mortality estimates on the other
360 hand mirror those in biomass, with the highest values corresponding to the lowest biomass
361 levels. The fact that the drop in fishing mortality following the 2004 recovery plan coincides
362 with the sudden rise in biomass strongly suggests that the reduction in fishing mortality is the
363 most probable cause behind the biomass increase currently observed. In the North Sea the
364 highest landings were recorded in the late 1940s immediately after World War II which
365 prevented commercial fishing for six years. This period of reduced harvest has been linked to
366 rapid increases in fish stock abundance with a faster response from older individuals
367 (Beverton and Holt 1957; Beare *et al.* 2010). Overfished stocks can experience rapid
368 recovery providing that a sufficient reduction in harvest rate is applied (Neubauer *et al.*
369 2013). It is likely that the recent increase in northern hake biomass resulted from the decline
370 in fishing pressure enforced by the recovery plan. The recent observed increase in the
371 proportion of stocks with rising biomass trends (and concomitant decreasing fishing mortality
372 trends) in the northeast Atlantic has been attributed to such multi-annual plans which also
373 included controls on fishing capacity (days-at-sea and vessel power) (Fernandes and Cook
374 2013).

375

376 The recent increase in northern hake biomass has also resulted in an increase of the overall
377 area occupied by the stock, with a striking expansion into the North Sea over the last ten
378 years. Both the biomass increase and the spatial expansion are proportionally higher in the
379 North Sea, showing the growing importance of northern hake in that particular area. The fact
380 that the centre of gravity of survey-based densities has remained unchanged in all areas but
381 the North Sea, advocates against a climate-induced northward distribution shift and suggests
382 instead an expansion of European hake into the North Sea. The northeast displacement of the
383 overall centre of gravity is most likely due to higher densities occurring in the North Sea.
384 These invasions are seasonal, as shown by both survey-based and commercial data, and
385 driven by both temperature and depth which is consistent with the density-dependent habitat
386 selection hypothesis and suggest that European hake expand to suitable habitats when
387 available (MacCall 1990). Northern hake landings realised in the North Sea in the late 1940s
388 and early 1950s were similar to current levels before declining throughout the 1960s and
389 1970s, probably as a consequence of a decreasing biomass resulting from overexploitation
390 (Goikoetxea and Irigoien 2013). The high values of these historical landings show that high
391 levels of biomass previously occurred in the North Sea when the fishing pressure was low
392 (Beare *et al.* 2010). This suggests that, providing that the current fishing mortality remains
393 low, the high levels of biomass experienced in the North Sea could become a permanent
394 feature.

395

396 The increase in biomass combined with an expansion of the area occupied by the stock
397 suggest that, if these two characteristics become permanent features, the northern hake stock
398 could offer new fishing opportunities. Under the current CFP, the distribution of the annual
399 TAC for the northern hake stock among ICES areas remains unchanged from year to year so
400 that each area and country is allocated the same proportion of the TAC every year, a policy

401 known as Relative Stability. This quota allocation key is based on historical catch records and
402 was set when the CFP was first adopted in 1983 (Symes 1997). At that time, hake landings in
403 the North Sea (area IV) were negligible, resulting in a strong west-east imbalance. The TAC
404 is now distributed as follows: 37% to the Bay of Biscay (area VIII) and 56% to the West of
405 Scotland and Celtic Sea (areas VI and VII respectively); while the North Sea (area IV) and
406 Skagerrak and Kattegat (area IIIa) are allocated just 4% and 3% respectively (ICES 2012a).
407 The findings from this study show that European hake undertake seasonal migrations
408 between areas VI and IV and are now present in the North Sea in large quantities during
409 summer months only. Survey-based SSB estimates for 2011 suggest that while the North Sea
410 and Skagerrak and Kattegat only contribute to 7% of the biomass in the first half of the year
411 (when West of Scotland, Celtic Sea and Bay of Biscay account for 44%, 46%, and 3%
412 respectively); they contribute to 34% of the biomass in the second half of the year when West
413 of Scotland, Celtic Sea and Bay of Biscay account for 12%, 50%, and 4% respectively. This
414 reveals a problematic aspect of the northern hake biomass increase: the quota allocation no
415 longer reflects the regional abundances. In the summer, when the fishery is active, the North
416 Sea has 34% of the entire stock SSB, but only 7% of the TAC.

417

418 The mismatch between allocated quotas and the regional abundance of commercial species
419 can result in major management challenges and unfavourable economic consequences. For
420 instance, a change in the timing of the migration in the Northeast Atlantic mackerel stock in
421 2009 led to the under-utilization of quotas worth over 100 M € as fishes were absent from
422 areas with allocated quotas (Jansen *et al.* 2012). The same mackerel stock has also been the
423 subject of a reduction in fishing pressure in the last decade (ICES 2012b). Like hake, this has
424 led to the expansion of its distribution to the west of northern Europe (ICES 2013). Mackerel
425 now appear more prominently in Icelandic & Faroese waters leading those nations to

426 unilaterally increase their allocation of catches. In 2005, Iceland caught 363 t of mackerel
427 (0.1% of the TAC): by 2011 this had risen to 155,000 t (17% of the TAC). The Faroe Islands
428 also unilaterally increased their quota from 2.4% (10,000 t) in 2005 to 13% (150,000 t) in
429 2011. These increases have since pushed the exploitation rate of mackerel beyond sustainable
430 limits and resulted in significant political disagreements between the EC (and Norway), and
431 Iceland and the Faroe Islands. In the case of European hake in the North Sea, the discrepancy
432 between available biomass and allocated quotas has led to the practice of extensive
433 discarding (Fernandes *et al.* 2011).

434

435 Discarding is a key issue for mixed-fisheries (Ulrich *et al.* 2011). In the North Sea, European
436 hake shares its habitat with other gadoid species and is caught as part of the North Sea
437 demersal mixed-fishery (Casey and Pereriro 1995). Being of similar size of other target
438 species and fished using the same gear, it is extremely difficult for fishermen to avoid
439 catching hake; especially if this species is present in large quantities as is the case today.
440 Under the current CFP it is possible for fishermen to trade quotas between ICES areas
441 (Valatin 2000) should they need it to land a species for which the quota in the area where
442 they are operating is exhausted. In 2011, a quota of 1,935 t (corresponding to 4% of the
443 northern hake TAC of 55,105 t) was allocated to the North Sea, 348 t of which were
444 distributed to the United Kingdom (ICES 2012a). By acquiring quotas from other areas,
445 Scottish fleets alone were able to land 3,035 t of hake caught in the North Sea, corresponding
446 to almost nine times the quota allocated for all British fleets. However, despite the trading of
447 quotas the large mismatch between low quotas and higher biomass still results in extensive
448 discarding occurring in the North Sea. While Scottish fleets landed 3,035 t of hake in the
449 North Sea in 2011, 4,993 t were discarded, bringing the total catches to 8,028 t which is more
450 than four times the TAC allocated to the whole North Sea and over 20 times the UK quota.

451 Such figures emphasize the difficulties created by a quota allocation scheme put in place
452 under markedly different ecological conditions.

453

454 The example of the northern hake stock shows that the management measures introduced by
455 the 2002 reform of the CFP can be successful in restoring depleted stocks' biomass providing
456 that good stewardship is applied. However, while offering new fishing opportunities a
457 recovering stock can also result in unexpected management issues, as shown in this study. In
458 the case of the northern hake stock in the North Sea, regional quotas do not reflect the
459 regional abundance of the increasing and expanding biomass, resulting in high discards. Such
460 issue challenges the relevance of the Relative Stability policy and the lack of flexibility in
461 adjusting regional quotas. The CFP is currently under reform in order to improve fish stock
462 conservation and achieve long-term economic viability of the fishing industry (EC 2013).
463 This revision includes a move towards a discard ban meaning that all fish caught at sea will
464 have to be landed: an option to ensure that this policy is realised is to close the fishery when
465 the quota of a given stock is reached. Atlantic cod (*Gadus morhua*, Gadidae) has previously
466 been identified as the "choke" species of the North Sea demersal mixed-fisheries for which
467 the quota would be exhausted first (Ulrich *et al.* 2011). However, the current level of biomass
468 and distribution of European hake in the North Sea documented here and the associated low
469 TAC suggest otherwise. If the increased levels of northern hake biomass reported here
470 persist, European hake is likely to be the "choke" species which effects a premature closure
471 of the entire demersal mixed-fishery in the North Sea.

472

473 The consideration of solutions to this problem lies beyond the scope of this paper, however it
474 is clear that a regional approach to management, which is included in the proposed reforms,
475 will help. Elements of co-management (Holmes *et al.* 2011) and new approaches to dealing

476 with a potential discard ban (Kindt-Larsen *et al.* 2011) will be needed if the potential of our
477 recovering fish stocks is not to be “choked” by inflexible management approaches which fail
478 to take into account the dynamic ecology of the oceans.

479

480

481 **Acknowledgements**

482

483 The authors wish to thank Rui Catarino, Henrik Degel, Mike Heath and Nick Bailey for their
484 contributions. This work was funded by the Seventh Framework Programme as part of the
485 European research project EcoFishMan (Grant No. FP7-265401). Paul Fernandes receives
486 funding from the MASTS pooling initiative (The Marine Alliance for Science and
487 Technology for Scotland) and their support is gratefully acknowledged. MASTS is funded by
488 the Scottish Funding Council (grant reference HR09011) and contributing institutions. The
489 authors declare no conflict of interest.

490

491

492 **References**

493

494 Alvarez, P. (2004) Distribution and abundance of European hake *Merluccius merluccius* (L.),
495 eggs and larvae in the North East Atlantic waters in 1995 and 1998 in relation to
496 hydrographic conditions. *Journal of Plankton Research* **26**, 811–826.

497 Bartolino, V., Ottavi, A., Colloca, F., Ardizzone, G.D. and Stefansson, G. (2008) Bathymetric
498 preferences of juvenile European hake (*Merluccius merluccius*). *ICES Journal of Marine*
499 *Science* **65**, 963–969.

500 Beare, D., Hölker, F., Engelhard, G.H., McKenzie, E. and Reid, D.G. (2010) An unintended
501 experiment in fisheries science: a marine area protected by war results in Mexican waves in
502 fish numbers-at-age. *Die Naturwissenschaften* **97**, 797–808.

- 503 Beare, D.J., Burns, F., Greig, A., *et al.* (2004) Long-term increases in prevalence of North
504 Sea fishes having southern biogeographic affinities. *Marine Ecology Progress Series* **284**,
505 269–278.
- 506 Beverton, R.J.H. and Holt, S.J. (1957) *On the dynamics of exploited fish populations*.
507 Fisheries Investigations, London, Ser. 2. 19:533 pp.
- 508 Cardinale, M., Dörner, H., Abella, a., *et al.* (2013) Rebuilding EU fish stocks and fisheries, a
509 process under way? *Marine Policy* **39**, 43–52.
- 510 Casey, J. and Pereriro, J. (1995) European hake (*M. merluccius*) in the Northeast Atlantic. In:
511 *Hake, Fisheries, Ecology and Markets*. (eds J. Alheit and T.J. Pitcher). Chapman and Hall,
512 London.
- 513 Cheung, W.W.L., Pinnegar, J., Merino, G., Jones, M.C. and Barange, M. (2012) Review of
514 climate change impacts on marine fisheries in the UK and Ireland. *Aquatic Conservation:
515 Marine and Freshwater Ecosystems* **22**, 368–388.
- 516 Coull, K.A., Jermyn, A.S., Newton, A.W., Henderson, G.I. and Hall, W.B. (1989) Length-
517 weight relationships for 88 species of fish encountered in the North East Atlantic. Scottish
518 Fish. Res. Rep. **43**, 1–80 pp.
- 519 Daw, T. and Gray, T. (2005) Fisheries science and sustainability in international policy: a
520 study of failure in the European Union's Common Fisheries Policy. *Marine Policy* **29**, 189–
521 197.
- 522 Dulvy, N.K., Rogers, S.I., Jennings, S., Stelzenmiller, V., Dye, S.R. and Skjoldal, H.R. (2008)
523 Climate change and deepening of the North Sea fish assemblage: a biotic indicator of
524 warming seas. *Journal of Applied Ecology* **45**, 1029–1039.
- 525 EC (2013) Amendement proposal for a regulation of the European Parliament and of the
526 Council on the European Maritime and Fisheries Fund. *Communication from the Commission
527 to the European Parliament, the Council, the European Economic and Social Committee and
528 the Committee of the Regions* Brussels, 22.4.2013 COM(2013) 245 final.
- 529 EC (2001) Commission Regulation (EC) No. 2602/2001 of 27 December 2001 establishing
530 additional technical measures for the recovery of the stock of hake in ICES subareas III, IV,
531 V, VI and VII and ICES Divisions VIIIa,b,d,e. *Official Journal of the European Communities*.
532 L 345: 49.
- 533 EC (2002) Commission Regulation (EC) No. 494/2002 of 19 March 2002 establishing
534 additional technical measures for the recovery of the stock of hake in ICES subareas III, IV,
535 V, VI and VII and ICES Divisions VIIIa,b,d,e. *Official Journal of the European Communities*.
536 L 77: 8.
- 537 EC (2004) Council Regulation (EC) No. 811/2004 of 21.4.2004 establishing measures for the
538 recovery of the Northern hake stock. *Official Journal of the European Union*. L 150:1.
- 539 Fernandes, P.G. and Cook, R. (2013) Reversal of fish stock decline in the Northeast Atlantic.
540 *Current Biology* **23**, R661–R662.

- 541 Fernandes, P.G., Coull, K., Davis, C., *et al.* (2011) Observations of discards in the Scottish
542 mixed demersal trawl fishery. *ICES Journal of Marine Science* **68**, 1734–1742.
- 543 Goikoetxea, N. and Irigoien, X. (2013) Links between the recruitment success of northern
544 European hake (*Merluccius merluccius* L.) and a regime shift on the NE Atlantic continental
545 shelf. *Fisheries Oceanography* **22**, 459–476.
- 546 Hiddink, J.G. and ter Hofstede, R. (2008) Climate induced increases in species richness of
547 marine fishes. *Global Change Biology* **14**, 453–460.
- 548 Hiddink, J.G., Jennings, S. and Kaiser, M.J. (2005) Do haddock select habitats to maximize
549 condition? *Journal of Fish Biology* **67**, 111–124.
- 550 Hilborn, R. and Walters, C.J. (1992) *Quantitative Fisheries Stock Assessment: Choice,*
551 *Dynamics and Uncertainty*. Chapman and Hall, New York.
- 552 Hinz, H., Kaiser, M.J., Bergmann, M., Rogers, S.I. and Armstrong, M.J. (2003) Ecological
553 relevance of temporal stability in regional fish catches. *Journal of Fish Biology* **63**, 1219–
554 1234.
- 555 Holmes, S.J., Bailey, N., Campbell, N., Catarino, R., Barratt, K., Gibb, A. and Fernandes,
556 P.G. (2011) Using fishery-dependent data to inform the development and operation of a co-
557 management initiative to reduce cod mortality and cut discards. *ICES Journal of Marine*
558 *Science* **68**, 1679–1688.
- 559 ICES (2013) Report of the Ad hoc Group on the Distribution and Migration of Northeast
560 Atlantic Mackerel (AGDMM). *ICES CM 2013/ACOM: 5 8*. 211 pp.
- 561 ICES (2012a) Report of the Working Group on the Assessment of Southern Shelf Stocks of
562 Hake, Monk and Megrim (WGHMM). *ICES CM 2012/ACOM: 11*. 617pp.
- 563 ICES (2012b) Report of the Working Group on Widely Distributed Stocks (WGWIDE).
564 *ICES CM 2012/ACOM: 15*. 931 pp.
- 565 Jansen, T., Campbell, A., Kelly, C., Hátún, H. and Payne, M.R. (2012) Migration and
566 fisheries of north east Atlantic mackerel (*Scomber scombrus*) in autumn and winter. *PloS one*
567 **7**. (doi:10.1371/journal.pone.0051541).
- 568 Kacher, M. and Amara, R. (2005) Distribution and growth of 0-group European hake in the
569 Bay of Biscay and Celtic Sea: a spatial and inter-annual analyses. *Fisheries Research* **71**,
570 373–378.
- 571 Kindt-Larsen, L., Kirkegaard, E. and Dalskov, J. (2011) Fully documented fishery: a tool to
572 support a catch quota management system. *ICES Journal of Marine Science* **68**, 1606–1610.
- 573 Kraak, S.B.M., Bailey, N., Cardinale, M., *et al.* (2013) Lessons for fisheries management
574 from the EU cod recovery plan. *Marine Policy* **37**, 200–213.
- 575 MacCall, A.D. (1990) *Dynamic geography of marine fish populations*. Washington Sea Grant
576 Program, Seattle.

- 577 Marshall, C.T. and Frank, K.T. (1995) Density-dependent habitat selection by juvenile
578 haddock (*Melanogrammus aeglefinus*) on the southwestern Scotian Shelf. *Canadian Journal*
579 *of Fisheries and Aquatic Sciences* **52**, 1007–1017.
- 580 Neubauer, P., Jensen, O.P., Hutchings, J.A. and Baum, J.K. (2013) Resilience and recovery
581 of overexploited marine populations. *Science (New York, N.Y.)* **340**, 347–349.
- 582 Petitgas, P., Alheit, J., Peck, M., *et al.* (2012) Anchovy population expansion in the North
583 Sea. *Marine Ecology Progress Series* **444**, 1–13.
- 584 Rayner, N. A. (2003) Global analyses of sea surface temperature, sea ice, and night marine
585 air temperature since the late nineteenth century. *Journal of Geophysical Research* **108**, 4407.
- 586 Schrope, M. (2010) What's the catch? *Nature* **465**, 540–542.
- 587 Shepherd, T.D. and Litvak, M.K. (2004) Density-dependent habitat selection and the ideal
588 free distribution in marine fish spatial dynamics: considerations and cautions. *Fish and*
589 *Fisheries* **5**, 141–152.
- 590 Symes, D. (1997) The European Community's common fisheries policy. *Ocean & Coastal*
591 *Management* **35**, 137–155.
- 592 Ulrich, C., Reeves, S.A., Vermard, Y., Holmes, S.J. and Vanhee, W. (2011) Reconciling
593 single-species TACs in the North Sea demersal fisheries using the Fcube mixed-fisheries
594 advice framework. *ICES Journal of Marine Science* **68**, 1535–1547.
- 595 Valatin, G. (2000) Quota Trading Systems in EU Fisheries. *Review of European Community*
596 *& International Environmental Law* **9**, 296–306.
- 597 Wheeler, A.J. (1969) *The Fishes of the British Isles and north-west Europe*. Michigan State
598 University Press, East Lansing, MI.
- 599 Woillez, M., Poulard, J.-C., Rivoirard, J., Petitgas, P. and Bez, N. (2007) Indices for
600 capturing spatial patterns and their evolution in time, with application to European hake
601 (*Merluccius merluccius*) in the Bay of Biscay. *ICES Journal of Marine Science* **64**, 537–550.
- 602 Woillez, M., Rivoirard, J. and Petitgas, P. (2009) Notes on survey-based spatial indicators for
603 monitoring fish populations. *Aquatic Living Resources* **22**, 155–164.
- 604
- 605

606 **Tables**

607

608 **Table 1.** Summary table of the Generalised Additive Models (GAMs) used to investigate
 609 statistical relationships between North Sea hake landings, sea surface temperature (SST) and
 610 depth (in 50 m depth classes). GAMs were performed using all available data (2000-2011),
 611 and using data recorded before (2000-2004) and after (2005-2011) the implementation of the
 612 2004 hake recovery plan (see methods) (*edf*: estimated degrees of freedom, *F*: F-ratio
 613 statistic).

614

Period	Deviance explained	<i>edf</i>		<i>F</i>		<i>P</i> -value	
		SST	Depth	SST	Depth	SST	Depth
2000-2011	18.8%	7.839	10	166.3	212.4	<0.001	<0.001
2000-2004	24.3%	5.519	9	98.24	74.15	<0.001	<0.001
2005-2011	20.9%	8.135	10	104.4	191.1	<0.001	<0.001

615

616

617 **Figure legends**

618

619 **Figure 1.** Summary of the northern hake stock assessment estimates. (a) total stock biomass
620 (TSB) together with recruitment (Rec). (b) spawning stock biomass (SSB) together with the
621 average fishing mortality (F). (c) Total landings for the northern hake stock, along with the
622 landings for each ICES area composing the stock: North Sea (area IV), West of Scotland
623 (area VI), Celtic Sea (area VII) and Bay of Biscay (area VIII).

624

625 **Figure 2.** Maps of the north east Atlantic displaying the spatial distribution of estimated
626 densities of northern hake from 2001 to 2011 showing the expansion of the stock into
627 northern areas and the North Sea. The chosen time series encompasses the increase in
628 northern hake densities observed over the last 10 years. Densities are colour coded by quarter
629 (blue: Q1, green: Q2, orange: Q3, red: Q4). Survey data were not available for the Celtic Sea
630 and Bay of Biscay in 2011.

631

632 **Figure 3.** Statistics of the spatial occupation of northern hake showing expansion and
633 changes in distribution. (a) The percentage of ICES rectangles surveyed where densities
634 greater than the maximum North Sea hake density observed prior to the 2004 recovery plan
635 (85.7 kg/km^2) were recorded. This is a proxy to assess changes in the area occupied by the
636 stock. (b) Centres of gravity of hake densities observed in each survey from 2001 to 2011,
637 labelled 1 to 11. The overall centres of gravity for the northern hake stock calculated for
638 years sampled by all regional surveys (2001-2008, see methods) are averaged prior (2001-
639 2004) and after (2005-2008) the 2004 recovery and are also represented by a black circle and
640 a black triangle respectively. The 100 m depth contour is displayed in grey.

641

642 **Figure 4.** Monthly maps of the British Isles showing Scottish and Danish hake landings per
643 ICES rectangle between 2000 and 2011 by month (data for Scottish landings were available
644 from 2005 onwards). The proportion of Scottish and Danish landings are displayed in blue
645 and red respectively. The coloured background corresponds to monthly sea surface
646 temperature (°C) averaged from 2000 to 2011. The 100 m depth contour is displayed in
647 black.

648

649 **Figure 5.** Estimates of European hake spawning stock biomass (SSB) calculated from survey
650 data for each ICES area included in the northern hake stock (a-d), and historical landings
651 from the North Sea (e). For each area, SSB estimates calculated for the first (dark grey) and
652 second (black) half of the year, according to data availability, are displayed from 2001 to
653 2011. (e) The historical North Sea landings from 1903 to 2010 (grey histogram) from FAO
654 are displayed along the North Sea SSB estimates.

Figures




Figure 1.




Figure 2.

Review Only




Figure 3.




Figure 4.

Review Only




Figure 5.

Supporting information




Figure S1. Results from Generalised Additive Models (GAMs) used to investigate statistical relationships between the distribution of North Sea hake landings and the variables temperature and depth. GAMs were performed using all available data (2000-2011, panels (a) and (b)), and using data recorded before (2000-2004, panels (c) and (d)) and after (2005-2011, panels (e) and (f)) the implementation of the 2004 hake recovery plan to assess potential impact of an increase in abundance on the relationship between hake distribution

and environmental variables. The continuous and dashed lines show the model fits and 95% confidence intervals respectively. The tick marks on the x -axis indicate the data availability.

For Review Only