

# Alcohol Interventions for Trauma Patients Treated in Emergency Departments and Hospitals

## A Cost Benefit Analysis

Larry M. Gentilello, MD,\* Beth E. Ebel, MD, MPH,†|| Thomas M. Wickizer, MPH, PhD,‡  
David S. Salkever, PhD,§ and Frederick P. Rivara, MD, MPH†||

**Objective:** To determine if brief alcohol interventions in trauma centers reduce health care costs.

**Summary Background Data:** Alcohol-use disorders are the leading cause of injury. Brief interventions in trauma patients reduce subsequent alcohol intake and injury recidivism but have not yet been widely implemented.

**Methods:** This was a cost-benefit analysis. The study population consisted of injured patients treated in an emergency department or admitted to a hospital. The analysis was restricted to direct injury-related medical costs only so that it would be most meaningful to hospitals, insurers, and government agencies responsible for health care costs. Underlying assumptions used to arrive at future benefits, including costs, injury rates, and intervention effectiveness, were derived from published nationwide databases, epidemiologic, and clinical trial data. Model parameters were examined with 1-way sensitivity analyses, and the cost-benefit ratio was calculated. Monte Carlo analysis was used to determine the strategy-selection confidence intervals.

**Results:** An estimated 27% of all injured adult patients are candidates for a brief alcohol intervention. The net cost savings of the intervention was \$89 per patient screened, or \$330 for each patient offered an intervention. The benefit in reduced health expenditures resulted in savings of \$3.81 for every \$1.00 spent on screening and intervention. This finding was robust to various assumptions regard-

ing probability of accepting an intervention, cost of screening and intervention, and risk of injury recidivism. Monte Carlo simulations found that offering a brief intervention would save health care costs in 91.5% of simulated runs. If interventions were routinely offered to eligible injured adult patients nationwide, the potential net savings could approach \$1.82 billion annually.

**Conclusions:** Screening and brief intervention for alcohol problems in trauma patients is cost-effective and should be routinely implemented.

(*Ann Surg* 2005;241: 541–550)

Alcohol intoxication is the leading risk factor for injury.<sup>1–3</sup> As a result, it offers the most promising and obvious target for injury-prevention programs. Brief alcohol interventions in trauma patients have been shown to reduce subsequent alcohol intake and injury recidivism.<sup>4–8</sup> Given accumulating evidence to support their use, a variety of expert and consensus group panels have concluded that the scientific basis for their routine provision in hospitals and emergency departments has been established, and it is time to move towards national implementation.<sup>7,9–16</sup>

New medical procedures, once confirmed as “best practice,” often become virtually mandatory for delivery and insurance coverage. However, this has not been the case for alcohol interventions. Instead, clinicians, hospital administrators, and insurance plans are more likely to require information about cost and benefits before making decisions on implementation or coverage.

Alcohol interventions in trauma patients have not yet been analyzed for cost-benefit. Prior studies have not measured financial outcomes directly because it has been difficult to obtain claims data from this patient population with multiple potential sources of insurance coverage or with no coverage. However, the assignment of dollar values can enable clinicians to make informed choices about competing available treatment options.<sup>17</sup>

From the \*Department of Surgery, University of Texas Southwestern Medical School, Dallas, Texas; †Department of Pediatrics, School of Public Health and Community Medicine, University of Washington, Seattle, Washington; ‡Department of Health Services, School of Public Health and Community Medicine, University of Washington, Seattle, Washington; §Department of Health Policy and Management, Johns Hopkins University School of Public Health, Baltimore, MD; and the ||Harborview Injury Prevention and Research Center, University of Washington, Seattle, Washington.

Supported by a grant from the Robert Wood Johnson Foundation: 042424. Reprints: Larry M. Gentilello, MD, Department of Surgery, Division of Burns, Trauma and Critical Care, Parkland Memorial Hospital, University of Texas Southwestern Medical School, 5323 Harry Hines Boulevard, MC 9158, Dallas, TX. E-mail: larry.gentilello@utsouthwestern.edu.

Copyright © 2005 by Lippincott Williams & Wilkins  
ISSN: 0003-4932/05/24104-0541  
DOI: 10.1097/01.sla.0000157133.80396.1c

This study estimates the cost savings associated with routine provision of brief alcohol interventions to trauma patients treated in hospitals and emergency departments. We chose to restrict this analysis to direct medical costs so that it would be most meaningful to hospitals, insurers, and government agencies responsible for health care costs.

## METHODS

### Determination of Eligible Subjects

We estimated the proportion of injured patients who would be candidates for a brief alcohol intervention. Patients were considered eligible if they were treated in an emergency department or admitted to a hospital after sustaining an injury, were 18 years of age or older, and had either a blood alcohol level  $\geq 100$  mg/dL or a positive result on a standard brief alcohol disorder screening questionnaire. Patients with a major concurrent psychiatric illness or severe disability precluding participation in a brief intervention were excluded.

We performed a literature search to identify studies reporting alcohol intoxication (blood alcohol concentration: BAC) or problem drinking as defined by a positive result on a standard screening questionnaire (CAGE, Michigan Alcohol Screening Test, or Alcohol Use Disorders Identification Test) in injured adults treated in emergency departments (Table 1). The screen positive rate varied, depending on the proportion of all patients screened and the methods used to classify a screen as positive. These differences prevented formal meta-analysis. To arrive at a prevalence estimate, the proportion from each study was weighted by the study sample size.<sup>18</sup> The range was used to conduct a sensitivity analysis.

Eligible patients would have to agree to counseling. Consent rates for studies enrolling patients in brief intervention trials were used to estimate acceptance rates, as no data exist on acceptance rates when offered as a routine component of trauma care.<sup>4,5,7,8,19-24</sup> The mean consent rate, representing 9116 subjects, was 76% (range 57 to 94%). This is a conservative estimate as patients are more likely to consent to routine care than to participation in a study.

### Cost-Benefit Analysis

#### Screening Costs

We determined the cost of screening all eligible adults with a BAC and a brief alcohol-disorder screening questionnaire. The direct cost of a BAC was valued at \$15 based upon the current Medicare allowable fee schedule in 2000. Screening expenses also included the cost of paper materials (\$1). Average screening costs were \$16/patient (Table 2).

#### Costs of Brief Alcohol Intervention

The cost of the brief intervention included direct costs representing professional expenses and materials. Typically, brief interventions are administered over a 20- to 30-minute period.<sup>7,25</sup> Professional costs were determined by multiplying the national average hourly wage for a psychologist by an estimated 1.4 hours of work time per intervention, which included 30 minutes for the intervention, and the remainder for follow-up and documentation. Sensitivity analysis was conducted to include salary ranges for social workers to physicians (\$15.09 to \$61.43 per hour).<sup>26</sup> The estimated cost of the intervention is \$38 per patient (Table 2).

**TABLE 1.** Prevalence of Alcohol Use in Injured Patients Treated in US Emergency Departments

| Author                              | Population   | Measurement     | Prevalence* | BAC<br>$\geq 100$ mg/dL | N    |
|-------------------------------------|--|-----------------|-------------|-------------------------|------|
| Wechsler, et al, 1969 <sup>58</sup> | ED   | Breath analysis | 23%         | 12%                     | 2989 |
| Cherpitel, 1988 <sup>59</sup>       | ED in 4 hospitals  | Breath analysis | 15%         | 8%                      | 1528 |
| Cherpitel, 1989a <sup>60</sup>      | ED, hospital   | Breath analysis | 23%         | 15%                     | 555  |
| Teplin et al, 1989 <sup>61</sup>    | ED   | Blood test      | NA          | 10%                     | 179  |
| Rivara et al, 1989 <sup>62</sup>    | ED, level 1 hospital   | Blood test      | 38%         | NA                      | 316  |
| Cherpitel, 1992 <sup>63</sup>       | ER admissions  | Breath analysis | 13%         | 6%                      | 1004 |
| Cherpitel, 1993 <sup>64</sup>       | 3 HMO EDs, 1 county hospital ED, 3 community hospital EDs            | Breath analysis | 9%          | 3%                      | 1478 |
| Cherpitel, 1994 <sup>65</sup>       | EDs in 1 county hospital, 3 community hospitals, and 3 HMO hospitals | Breath analysis | 9%          | 3%                      | 1548 |
| Cherpitel, 1994b <sup>66</sup>      | 4 EDs  | Breath analysis | 9%          | 4%                      | 1494 |
| Average (weighted by sample size)   |  |                 | 16%         | 7.4%                    |      |
| Range                               |  |                 | 9-38%       | 3-15%                   |      |

ED indicates emergency department; BAC, blood alcohol concentration.  
\*Positive for any blood alcohol.

**TABLE 2.** Determination of Model Variables

| Variable  | Baseline Value (US\$ 2000) | Range                    | Source   |
|---|----------------------------|--------------------------|--|
| Number of adults ( $\geq 18$ y) treated for injuries in US EDs, 2000  | 20,507,601                 | 18,407,275 to 22,607,928 | NCIPC <sup>28</sup>  |
| Proportion of injured ED patients who required hospitalization  | 6.00%                      | 4.61%-7.38%              | NCIPC <sup>28</sup>  |
| Proportion of injured patients with psychiatric illness prohibiting intervention  | 2.4%                       | 2.4%                     | Gentilello <sup>7</sup>  |
| Prevalence of intoxicated injured adult patients treated in ED (blood alcohol concentration $\geq 100$ mg/dL)             | 7.4%                       | 3%-12%                   | See Table 1  |
| Prevalence of nonintoxicated trauma patients with a positive substance abuse screening questionnaire                      | 19.6%                      | 11.7%-45.8%              | See references <sup>2,38,39</sup>  |
| Consent rate to participate in a brief intervention trial   | 76%                        | 57%-94%                  | See references <sup>4,5,7,8,19-24</sup>  |
| Annual proportion of injured adult patients with intoxication or alcohol problems readmitted to an ED for trauma          | 28%                        | 5%-50%                   | See Table 3  |
| Relative risk for readmission in injured patients with alcohol dependency compared to injured patients without dependency | 2.2                        | 1.4-3.5                  | Rivara <sup>27</sup>   |
| Economic annual discount rate   | 3%                         | 0%-5%                    | Weinstein et al, 1996 <sup>32</sup>  |
| Costs of screening for alcohol use  |                            |                          |  |
| Cost of blood alcohol test  | \$15                       | \$15                     | Medicare reimbursement   |
| Cost of materials (paper, photocopy)  | \$1                        | \$0-\$1                  | Estimated cost of 5 copies at \$0.20/copy                                      |
| Cost of brief alcohol intervention  |                            |                          |  |
| Provider time (30-min intervention plus administrative time)  | 84 min                     | 30-120 min               | HMC brief intervention program   |
| Provider hourly salary  | \$27.20 (psychologist)     | \$15.09-\$61.43          | Hourly salary range: social worker; psychologist; physician, BLS <sup>26</sup> |
| Health care costs   |                            |                          |  |
| Cost of ED visit for injury   | \$440                      | \$149-\$1250             | MarketScan <sup>44</sup>   |
| Cost of hospitalization   | \$16,852                   | \$4,691-\$49,672         | MarketScan <sup>44</sup>   |
| Effectiveness of brief intervention in the context of acute injury at reducing subsequent health care use                 |                            |                          |  |
| Effectiveness of intervention in reducing annual ED injury recidivism (hazard ratio)                                      | 0.53                       | 0.26-1.07                | Gentilello <sup>7</sup>  |
| Effectiveness of intervention in reducing annual injury recidivism requiring hospitalization (hazard ratio)               | 0.52                       | 0.21-1.29                | Gentilello <sup>7</sup>  |

ED indicates emergency department; NCIPC, National Center for Injury Control and Prevention, Centers for Disease Control and Prevention; HMC, Harborview Medical Center, Seattle, WA; NIAAA, National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health; BLS, Bureau of Labor Statistics.

### Emergency Department Visits and Hospitalization Rates for Problem Drinkers

To estimate the baseline proportion of screen-positive patients who will sustain an injury requiring emergency department treatment in a given year, we performed a literature review (Table 3). The proportion of patients treated for an injury in the emergency department from each study was

weighted by the study sample size.<sup>18</sup> The range of the reported fraction of alcohol-using patients was used to conduct the model sensitivity analysis. The relative risk for readmission in injured patients with an intoxicating BAC or positive screening questionnaire compared with injured patients without these characteristics was estimated to be 2.2 (range 1.4 to 3.5).<sup>27</sup> We assumed that the proportion who

**TABLE 3.** Proportion of Intoxicated or Injured Problem Drinkers With Subsequent Injury Requiring an Emergency Department Visit Within 1 Year

| Author                        | Study Site        | Proportion of Patients Revisiting Emergency Departments Within 1 y | N    | Number of ED Visits |
|-------------------------------|-------------------|--|------|---------------------|
| Fleming, 1999 <sup>21</sup>   | Primary care      | 35%  | 382  | 132                 |
| Gentilello, 1999 <sup>7</sup> | ED, trauma center | 10%  | 396  | 38                  |
| Freeborn, 2000 <sup>22</sup>  | Primary care      | 9%   | 254  | 23                  |
| Blose, 1991 <sup>41</sup>     | Primary care      | 23%  | 3729 | 858                 |
| Cryer, 1999 <sup>67</sup>     | Primary care      | 42%  | 2253 | 935                 |
| Davidson, 1997 <sup>68</sup>  | ED, trauma center | 50%  | 150  | 75                  |
| Schermer, 2001 <sup>69</sup>  | ED                | 5%   | 180  | 9                   |
| Total (range)                 |                   | 28% (5%-50%)   | 6962 | 1937                |

ED indicates emergency department.

required admission to the hospital was equivalent to national estimates in which 6% of injured emergency department patients required hospitalization.<sup>28</sup>

### Intervention Effectiveness

We conducted a literature search to identify studies reporting brief alcohol interventions for patients treated in the context of an acute injury for which subsequent utilization of health care resources was an outcome. Multiple reviews have reported that brief interventions are effective in reducing alcohol consumption, problems related to consumption, and injury.<sup>6,19,29,30</sup> Three studies conducted in the acute-injury context reported substantial decreases in reinjury rates.<sup>4,7,8</sup> However, only 1 reported the effect on injury-related emergency department utilization, thus fulfilling the search criteria. This was a randomized controlled trial in which patients who were admitted to a hospital and screened positive were offered a brief intervention.<sup>7</sup> The authors reported a 47% reduction in subsequent injuries requiring either emergency department or hospital admission (hazard ratio 0.53, 95% CI 0.26–1.07) and a 48% reduction in injuries requiring hospital admission over 3 years of follow-up (hazard ratio 0.52, 95% CI 0.21–1.29). These confidence intervals were used for sensitivity analysis.

### Costs of Emergency Visits and Hospitalization

The costs of emergency department visits and hospital admissions were derived from the 1998 MarketScan database of commercial claims (MedStat Group, Inc., Ann Arbor, MI). This database reflects reimbursed payments by commercial insurance carriers and Medicare supplemental reimbursement. Injury episodes were identified by *ICD9-CM* codes. The cost of an emergency department visit for treatment of an injury and the cost of a hospital admission for injury reflected the average reimbursement. We conducted sensitivity analyses using the 25th-percentile cost reimbursement amount as the lower limit and the average cost reimbursement plus 1

standard deviation as the upper limit (Table 2). All cost data were converted into year 2000 US dollars using the Consumer Price Index for wage data, and the Consumer Price Index for Medical Care for medical expenditures.<sup>31</sup>

To reflect the opportunity cost of future investment returns for financial or physical resources used in the present, a discount rate of 3% was employed for all future costs. Rates varying between 0% and 5% were tested in a sensitivity analysis.<sup>32</sup>

### Estimation of Cost Savings From Reduced Trauma Recidivism

The average length of stay for a hospital injury admission was estimated to be 5.1 days.<sup>33</sup> We have not included costs for postdischarge medical care required by more severely injured patients, nor have we assigned a cost to individuals who died.

### Decision Analysis

A decision-analysis model was used to determine whether the cost-benefit analysis favored the adoption of brief alcohol interventions by comparing 2 scenarios (Fig. 1). In the first, all eligible injured patients would be screened and would incur screening costs. Patients who screen positive would be asked to consent to a brief intervention, and accrue intervention costs. In the second scenario, injured patients would not be screened or offered an intervention. The decision tree compares the costs of screening and intervention to the potential cost savings from reduced trauma recidivism.

The model was constructed based on Figure 1. It incorporated an imbedded Markov model allowing for transitions between the defined states of “well,” “injured requiring emergency department care,” and “injured requiring hospital admission” in a yearly cycle. We assumed that transition probabilities and the costs of being in each state were time independent. A 1-year period was used, and the model was run for 3 periods of observation, emulating the intervention




FIGURE 1. Flow diagram illustrating potential outcomes related to screening and intervention strategies.

study in which injury recidivism was reduced over the course of 3 years.<sup>7</sup> The costs of being in each state accrued during each cycle. The analysis was performed using Data 4.0 (TreeAge Software, Inc., Williamstown, MA).

Sensitivity testing was performed for the critical variables to explore their impact on cost savings using the confidence intervals or ranges derived as discussed above (Fig. 2). A graph of these univariate sensitivity analyses called a “tornado plot” was generated to compare the relative influence of model variables on final cost estimates. Sensitivity analysis was done using Monte Carlo analyses in which key parameters were varied simultaneously, performed with Data™ 4.0 (TreeAge Software, Inc., Williamstown, MA).<sup>37</sup> Distributions were sampled 50,000 times for the final analysis (Fig. 3). Probability density functions for parameters were chosen to approximately match the high and low ranges found in the published studies cited and in our own data. Emergency department costs and hospitalization costs were modeled using a  $\gamma$  distribution.<sup>38</sup> Parameters measuring intervention effectiveness at reducing ED and hospital recidivism were modeled with a log normal distribution.<sup>38</sup> Other


probabilities were transformed using a method described by Doubilet et al.<sup>37</sup>

## RESULTS

In 2000, there were an estimated 20.5 million adult injuries requiring emergency department care.<sup>28</sup> The weighted average of studies in Table 1 indicates that approximately 7.4% (range 3% to 12%) were intoxicated. An additional 19.6% of these patients screen positive on an alcohol screening questionnaire even when not intoxicated (range 11.7% to 45.8%).<sup>2,39,40</sup>

Approximately 94% of injured adults treated in the emergency department are discharged, and 6% are admitted to the hospital.<sup>28</sup> The probability that an injured patient has a concurrent, serious psychiatric illness was 2.4%.<sup>7</sup> In total, 27% of injured adult patients treated in the emergency department screen positive and could be candidates for an intervention (5.5 million visits/y).

The proportion of problem drinkers who will return to the emergency department for treatment of a new injury within 1 year was estimated to be 28% (range 5% to 50%;



**FIGURE 2.** Sensitivity analysis demonstrating potential cost savings associated with variations in a number of primary variables. Screening and brief intervention is associated with cost savings when the bar is above zero. The estimated mean cost savings is \$89 in US dollars (year 2000). BAI, brief alcohol intervention; ED, emergency department; RR, relative risk.


Table 3). It was assumed that the proportion who will require admission to the hospital was 6.0%, reflecting the national admission rate for injured adults treated in emergency departments (Table 2).

### Cost Analysis of Universal Screening and a Brief Alcohol Intervention Policy

Using the above base-case parameter values, the analysis found that routine screening and intervention would save costs and is the preferred option (Table 4). If the brief intervention is offered, the expected cost of screening, intervention, and subsequent emergency department visits and hospital admissions over the next 3 years was \$600 per patient. In the scenario where screening and intervention are not offered, the expected cost of subsequent emergency department visits and hospital admissions was \$689 per patient over 3 years, resulting in an estimated cost savings of

\$89 per injured patient screened, or \$330 for each patient offered a brief intervention. The brief intervention resulted in \$3.81 in health care costs saved for every \$1.00 spent on screening and intervention.

Sensitivity analyses were used to determine when it was optimal to offer the intervention. The most critical variables included the costs of hospitalization, the hazard ratios for trauma recidivism following a brief intervention, and the probability of a subsequent injury resulting in emergency department care. The hazard ratio for having a subsequent injury requiring emergency department treatment was allowed to vary between 0.26 and 1.07.<sup>7</sup> The hazard ratio for hospital readmission was allowed to vary between 0.21 and 1.29.<sup>7</sup> One-way and 2-way sensitivity testing for the most critical model variables of brief intervention effectiveness showed that under most assumptions, it is preferable to offer



**FIGURE 3.** Histogram demonstrating likelihood of cost savings where variables are allowed to range simultaneously. In 91.5% of simulations, screening and intervention were associated with cost savings.

the intervention because its costs are very low compared with the costs of repeat hospital care (Fig. 2). The model was relatively invariant to the probability of accepting a brief intervention, the cost of a brief intervention, and the cost of alcohol screening, as these costs were small compared with the potential savings in medical costs.

We used Monte Carlo analysis in which each parameter was allowed to vary, using a distribution centered around the baseline value to account for the uncertainty in estimating each parameter. The results indicated that a brief alcohol intervention would result in a reduction of health care expenditures in 91.5% of the simulations (Fig. 3). In 95% of the simulations, the intervention was either cost saving or cost

less than \$24 per patient screened. Given that the benefits of reducing alcohol use greatly exceed the reduction in health care costs alone, the brief alcohol intervention is an inexpensive and likely cost-savings therapy for injured patients.

## DISCUSSION

This study indicates that routine alcohol screening and intervention in trauma patients results in net dollar savings through the effect of the intervention on reducing subsequent health care costs. Current recommendations to provide brief interventions to trauma patients are based on a number of factors. They capitalize on a “teachable moment,” wherein the health care worker can moderate the conceptual link between drinking and its consequences at a time when the consequences are obvious. Patients with alcohol problems may not seek alcohol treatment but often receive treatment of medical conditions related to their drinking. Injuries are the most common condition for which patients with alcohol problems seek medical attention.<sup>42</sup> Alcohol interventions in trauma centers may also provide an opportunity to initiate care before problems progress to a more severe stage, requiring more costly and intensive treatment and medical services.<sup>42</sup> Alcohol also plays such a major role in causing injuries that injury-prevention programs are unlikely to be successful if hazardous drinking is not addressed. This study now adds a strong financial argument to the above reasons for implementing routine alcohol screening and intervention programs in trauma centers.

Before considering the implications of this research, the potential limitations should be understood. Multiple studies have reported that brief interventions conducted in emergency departments or trauma centers reduce the rate of injury

**TABLE 4.** Cost-Savings Analysis of Brief Alcohol Intervention Policy (\$US 2000)

|   | Screening and Brief Intervention          |  |                             |                                     |
|---|---|--|-----------------------------|-------------------------------------|
|   | Patient Ineligible for Intervention (73%) | Patients Eligible for Intervention (27%) |                             | No Screening and Brief Intervention |
|   |   | Intervention Accepted (20.5%)            | Intervention Refused (6.5%) |                                     |
| Screening costs   | \$16                                      | \$16                                     | \$16                        | \$0                                 |
| Intervention costs  | \$0                                       | \$38                                     | \$0                         | \$0                                 |
| Health care costs of ED and hospital recidivism over next 3 y | \$521                                     | \$600                                    | \$1145                      | \$689                               |
| Total costs per adult trauma patient                          | \$536                                     | \$653                                    | \$1161                      | \$689                               |
| Weighted average costs  |   | \$600                                    |                             | \$689                               |
| Cost savings per patient screened                             |   | \$89                                     |                             | \$0                                 |
| Cost savings per intervention                                 |   | \$330                                    |                             | \$0                                 |
| Cost savings per dollar spent                                 | \$3.81                                    |  | \$0                         |                                     |
| Potential annual savings                                      | \$1.82 billion                            |  | \$0                         |                                     |

SBI indicates screening and brief intervention; ED, emergency department.

recidivism.<sup>4,7,9</sup> However, due to differences in method of reporting, costs could not be analyzed, and therefore, they were not included in this paper. For example, Monti et al<sup>8</sup> reported a greater than 50% reduction in reinjury rate in intervention compared with control group patients but did not report the percentage of patients who subsequently required injury-related emergency department or trauma-center care. Another study analyzed 84 different injury prevention methods and found that brief interventions for heavy drinkers were ranked towards the top in benefit-cost ratio, but outcomes other than injury recidivism were included in the analysis.<sup>43</sup>

We therefore based our model sensitivity range on a randomized study that reported the impact of trauma-center interventions on the annual rate of emergency department injury-related visits.<sup>7</sup> Despite this limitation, as shown in Figure 2, only under extremely negative assumptions regarding the intervention effect on injury recidivism rates would the net cost savings drop to zero. And, taken together, multiple recent studies support the notion that injury-related hospital visits are reduced following a brief intervention.

Another potential limitation of the study is that the results depend on the estimates used for costs of hospitalization. We used data from MarketScan, which represents a spectrum of trauma patients across Medicaid, Medicare, and private insurers.<sup>44</sup> It has been widely used in other cost-effectiveness and cost-benefit studies.<sup>45–51</sup> However, we did not include the costs of follow-up care. Data from the Medical Expenditure Panel Survey suggest that the average non-admitted injured patient treated in the emergency department generates \$832 in medical expenditures when follow-up care is included, which exceeds the \$440 cost used in this study for the ED visit alone.<sup>52</sup> Medical expenditures for follow-up care and rehabilitation services for patients who are admitted are likely to be even higher.

We also chose to provide cost-benefit information that is primarily relevant to clinicians, hospital administrators, insurance carriers, and governmental agencies that are responsible for health care costs and insurance legislation. Therefore, this study only examined the impact of interventions on direct medical costs. Many cost-benefit analyses include costs related to mental health services, property damage, lost productivity, crime, and intangible losses such as pain, suffering, and reduced quality of life. From a societal perspective, these costs are likely to be large, resulting in a much greater savings than \$3.81 for each dollar spent.

Despite these limitations, we believe that this study has important implications for trauma care in the United States. Unlike the majority of medical treatments that cost money to improve health care, offering brief interventions to injured patients saves money by reducing subsequent injuries requiring emergency department treatment or hospital admission. There are 20.5 million emergency department visits and 1 million hospitalizations of adults in the United States annu-

ally for trauma, and approximately 5.5 million of these individuals are candidates for an alcohol intervention. If a brief intervention were offered to every eligible injured person in the United States, the resulting savings from health care costs alone would be approximately \$1.82 billion annually (20.5 million adult trauma patients × \$89 health care cost savings per patient screened). To put this number in perspective, it is more than twice the aggregate charges, or total national bill in year 2000, for all forms of cholecystitis (both acute and chronic).<sup>53</sup>

Implementation of screening and intervention programs in trauma centers will require changes in health care financing. Since 1947, most states have allowed insurance companies to deny payment on a claim in which an individual was injured and alcohol use was documented.<sup>54</sup> This has been found to be a deterrent to alcohol screening in hospitals and emergency departments.<sup>55</sup> The National Association of Insurance Commissioners recently passed a model law that disallows such denials. The National Conference of Insurance Legislators has recommended that states adopt this model. Such changes, however, must be adopted by each state legislature. Several states have recently done so.<sup>56,57</sup>

Excessive alcohol use is the single most important risk factor for injury and is by far the most promising yet underutilized target for injury prevention programs. The economic value of brief alcohol interventions supports the development of a policy of routine trauma-center screening and the provision of brief interventions to those who screen positive. Current insurance statutes resulting in financial penalties for patients and providers may prevent the implementation of such programs, resulting in greater health care costs.

## ACKNOWLEDGMENTS

This study was funded by a grant from the Robert Wood Johnson Foundation. The views expressed herein are our own and do not necessarily reflect those of the foundation.

## REFERENCES

1. Soderstrom CA, Dischinger PC, Smith GS, et al. Alcoholism at the time of injury among trauma center patients: vehicular crash victims compared with other patients. *Accid Anal Prev*. 1997;29:715–721.
2. Soderstrom CA, Dischinger PC, Smith GS, et al. Psychoactive substance dependence among trauma center patients. *JAMA*. 1992;267:2756–2759.
3. Rivara FP, Jurkovich GJ, Gurney JG, et al. The magnitude of acute and chronic alcohol abuse in trauma patients. *Arch Surg*. 1993;128:907–912.
4. Longabaugh R, Woolard RE, Nirenberg TD, et al. Evaluating the effects of a brief motivational intervention for injured drinkers in the emergency department. *J Stud Alcohol*. 2001;62:806–816.
5. Hungerford DW, Pollock DA, Todd KH. Acceptability of emergency department-based screening and brief intervention for alcohol problems. *Acad Emerg Med*. 2000;7:1383–1392.
6. D'Onofrio G, Degutis LC. Preventive care in the emergency department: screening and brief intervention for alcohol problems in the emergency department: a systematic review. *Acad Emerg Med*. 2002;9:627–638.
7. Gentilello LM, Rivara FP, Donovan DM, et al. Alcohol interventions in



- a trauma center as a means of reducing the risk of injury recurrence. *Ann Surg.* 1999;230:473–480.
8. Monti PM, Colby SM, Barnett NP, et al. Brief intervention for harm reduction with alcohol-positive older adolescents in a hospital emergency department. *J Consult Clin Psychol.* 1999;67:989–994.
  9. Institute of Medicine. *Broadening the Base of Treatment for Alcohol Problems: Report of a Study by a Committee of the Institute of Medicine, Division of Mental Health and Behavioral Medicine.* Washington, DC: National Academy Press; 1990.
  10. US Department of Health and Human Services. *Healthy People 2010: Understanding and Improving Health.* Washington, DC: US Government Printing Office; 2000.
  11. US Department of Health and Human Services. *Changing the Conversation: Improving Substance Abuse Treatment: The National Treatment Plan Initiative.* Rockville, MD: US Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Substance Abuse Treatment; 2000.
  12. National Institute on Alcohol Abuse and Alcoholism. *Alcohol and Trauma. Alcohol Alert.* 1989;3.
  13. National Highway Traffic Safety Administration. *Developing Best Practices of Emergency Care of the Alcohol-Impaired Patient: Recommendations from the National Conference.* Washington, DC: US Department of Transportation; 2001. Publication DOT HS 809 281.
  14. US Department of Health and Human Services. *Model Trauma Care Plan.* Rockville, MD: US Department of Health and Human Services; 1992.
  15. American Medical Association. *Screening for Alcohol and Other Drug Use in Trauma Patients.* Chicago, IL: American Medical Association; 1991. Policy No. H-130. 956, BOT Rep. J, I-91.
  16. US Department of Health and Human Services. *Alcohol and Other Drug Screening of Hospitalized Trauma Patients: Treatment Improvement Protocol (TIP) Series 16.* Rockville, MD: US Department of Health and Human Services, Public Health Service, Substance Abuse and Mental Health Services Administration, Center for Substance Abuse Treatment; 1995. DHHS Publication No. (SMA) 95–3039.
  17. Eisenberg JM. Clinical economics: a guide to the economic analysis of clinical practices. *JAMA.* 1989;262:2879–2886.
  18. Laird NM, Mosteller F. Some statistical methods for combining experimental results. *Int J Technol Assess Health Care.* 1990;6:5–30.
  19. Babor TF, Acuda W, Campillo C, et al. A cross-national trial of brief interventions with heavy drinkers: WHO Brief Intervention Study Group. *Am J Public Health.* 1996;86:948–955.
  20. Fleming MF, Barry KL, Manwell LB, et al. Brief physician advice for problem alcohol drinkers: a randomized controlled trial in community-based primary care practices. *JAMA.* 1997;277:1039–1045.
  21. Fleming MF, Manwell LB, Barry KL, et al. Brief physician advice for alcohol problems in older adults: a randomized community-based trial. *J Fam Pract.* 1999;48:378–384.
  22. Freeborn DK, Polen MR, Hollis JF, et al. Screening and brief intervention for hazardous drinking in an HMO: effects on medical care utilization. *J Behav Health Serv Res.* 2000;27:446–453.
  23. Kristenson H, Ohlin H, Hulthen-Nosslin MB, et al. Identification and intervention of heavy drinking in middle-aged men: results and follow-up of 24–60 months of long-term study with randomized controls. *Alcohol Clin Exp Res.* 1983;7:203–209.
  24. Richmond R, Heather N, Wodak A, et al. Controlled evaluation of a general practice-based brief intervention for excessive drinking. *Addiction.* 1995;90:119–132.
  25. National Institute on Alcohol Abuse and Alcoholism. Brief intervention for alcohol problems. *Alcohol Alert.* 1999;43.
  26. Bureau of Labor Statistics. National compensation survey: occupational wages in the United States, 2001 [US Department of Labor, Bureau of Labor Statistics Web site]. Available at: <http://www.bls.gov/ncs/ocs/sp/ncbl0449.pdf>. Accessed January 31, 2003.
  27. Rivara FP, Koepsell TD, Jurkovich GJ, et al. The effects of alcohol abuse on readmission for trauma. *JAMA.* 1993;270:1962–1964.
  28. National Center for Injury Prevention and Control. WISQARS™ (Web-based Injury Statistics Query and Reporting System) [Centers for Disease Control and Prevention Web site]. Available at: <http://www.cdc.gov/ncipc/wisqars/>. Accessed December 20, 2002.
  29. US Preventive Services Task Force. *Guide to Clinical Preventive Services.* 2nd ed. Baltimore, MD: Williams & Wilkins; 1996.
  30. Sommers MS, Dyehouse JM, Howe SR. Binge drinking, sensible drinking, and abstinence after alcohol-related vehicular crashes: the role of intervention versus screening. *Annu Proc Assoc Adv Automot Med.* 2001;45:317–328.
  31. Bureau of Labor Statistics. Historical consumer price index for urban wage earners and clerical workers (CPI-W): U.S. city average, all items. Available at: <http://www.bls.gov/cpi/cpid0110.pdf>. Accessed November 27, 2001.
  32. Weinstein MC, Siegel JE, Gold MR, et al. Recommendations of the Panel on Cost-effectiveness in Health and Medicine. *JAMA.* 1996;276:1253–1258.
  33. Nathens AB, Jurkovich GJ, EJ M, et al. A resource-based assessment of trauma care in the United States. *J Trauma.* 2004;56:173–178.
  34. Deleted in proof.
  35. MacKenzie EJ, Morris JA Jr, Jurkovich GJ, et al. Return to work following injury: the role of economic, social, and job-related factors. *Am J Public Health.* 1998;88:1630–1637.
  36. Deleted in proof.
  37. Doubilet P, Begg CB, Weinstein MC, et al. Probabilistic sensitivity analysis using Monte Carlo simulation: a practical approach. *Med Decis Making.* 1985;5:157–177.
  38. Briggs A. *Handling Uncertainty in Economic Evaluation.* Oxford: Oxford University Press; 2001.
  39. Maio RF, Waller PF, Blow FC, et al. Alcohol abuse/dependence in motor vehicle crash victims presenting to the emergency department. *Acad Emerg Med.* 1997;4:256–262.
  40. Soderstrom CA, Smith GS, Dischinger PC, et al. Psychoactive substance use disorders among seriously injured trauma center patients. *JAMA.* 1997;277:1769–1774.
  41. Blose JO, Holder HD. Injury-related medical care utilization in a problem drinking population. *Am J Public Health.* 1991;81:1571–1575.
  42. National Institute on Alcohol Abuse and Alcoholism. *Fourth Special Report to the US Congress on Alcohol and Health.* Washington, DC: US Government Printing Office; 1981.
  43. Miller TR, Levy DT. Cost-outcome analysis in injury prevention and control: eighty-four recent estimates for the United States. *Med Care.* 2000;38:562–582.
  44. The MEDSTAT Group, Inc. *MarketScan Database.* Ann Arbor, MI: MEDSTAT Group; 2002.
  45. Croghan TW, Melfi CA, Crown WE, et al. Cost-effectiveness of antidepressant medications. *J Ment Health Policy Econ.* 1998;1:109–117.
  46. Goodman AC, Nishiura E, Humphreys RS. Cost and usage impacts of treatment initiation: a comparison of alcoholism and drug abuse treatments. *Alcohol Clin Exp Res.* 1997;21:931–938.
  47. Goodman AC, Nishiura E, Hankin JR. Short term drug abuse treatment costs and utilization: a multi-employer analysis. *Med Care.* 1998;36:1214–1227.
  48. Hillman AL, Schwartz JS, Willian MK, et al. The cost-effectiveness of terazosin and placebo in the treatment of moderate to severe benign prostatic hyperplasia. *Urology.* 1996;47:169–178.
  49. Mark TL, Coffey RM, King E, et al. Spending on mental health and substance abuse treatment, 1987–1997. *Health Aff (Millwood).* 2000;19:108–120.
  50. Zhao SZ, Arguelles LM, Wong JM, et al. Cost comparisons between nafarelin and leuprolide in the treatment of endometriosis. *Clin Ther.* 1998;20:592–602.
  51. Lestina DC, Miller TR, Smith GS. Creating injury episodes using medical claims data. *J Trauma.* 1998;45:565–569.
  52. Medical Expenditure Panel Survey. Agency for Health Care Research and Quality. Available at: <http://www.meps.ahrq.gov/WhatIsMEPS/Overview.htm>.
  53. Health Care Cost and Utilization Project (HCUP). Outcomes for multiple conditions ICD-9-CM diagnosis codes 575.0, 575.10, 575.11, 575.12 [Agency for Health Care Research and Quality Web site]. Available at: <http://www.ahrq.gov/data/hcup/>.
  54. Rivara FP, Tollefson S, Tesh E, et al. Screening trauma patients for alcohol problems: are insurance companies barriers? *J Trauma.* 2000;48:115–118.

55. Schermer CR, Gentilello LM, Hoyt DB, et al. National Survey of Trauma Surgeons' use of alcohol screening and intervention. *J Trauma*. 2003;45:213.
56. Uniform Individual Accident and Sickness Insurance Policy Provisions Law (#180), revisions to model 180. Kansas City, MO; 2001.
57. Resolution in Support of Amending the NAIC Uniform Accident And Sickness Policy provision Law. Albany, NY; 2001.
58. Wechsler H, Kasey EH, Thum D, et al. Alcohol level and home accidents. *Public Health Rep*. 1969;84:1043–1050.
59. Cherpitel CJ. Drinking patterns and problems associated with injury status in emergency room admissions. *Alcohol Clin Exp Res*. 1988;12:105–110.
60. Cherpitel CJ. Breath analysis and self-reports as measures of alcohol-related emergency room admissions. *J Stud Alcohol*. 1989;50:155–161.
61. Teplin LA, Abram KM, Michaels SK. Blood alcohol level among emergency room patients: a multivariate analysis. *J Stud Alcohol*. 1989;50:441–447.
62. Rivara FP, Mueller BA, Fligner CL, et al. Drug use in trauma victims. *J Trauma*. 1989;29:462–470.
63. Cherpitel CJ. Acculturation, alcohol consumption, and casualties among United States Hispanics in the emergency room. *Int J Addict*. 1992;27:1067–1077.
64. Cherpitel CJ. Alcohol consumption among emergency room patients: comparison of county/community hospitals and an HMO. *J Stud Alcohol*. 1993;54:432–440.
65. Cherpitel CJ. Injury and the role of alcohol: county-wide emergency room data. *Alcohol Clin Exp Res*. 1994;18:679–684.
66. Cherpitel CJ. Cause of casualty and drinking patterns: an emergency room study of unintentional injuries. *Drug Alcohol Depend*. 1994;35:61–67.
67. Cryer PC, Jenkins LM, Cook AC, et al. The use of acute and preventative medical services by a general population: relationship to alcohol consumption. *Addiction*. 1999;94:1523–1532.
68. Davidson P, Koziol-McLain J, Harrison L, et al. Intoxicated ED patients: a 5-year follow-up of morbidity and mortality. *Ann Emerg Med*. 1997;30:593–597.
69. Schermer CR, Qualls CR, Brown CL, et al. Intoxicated motor vehicle passengers: an overlooked at-risk population. *Arch Surg*. 2001;136:1244–1248.