

Northumbria Research Link

Citation: Newbury-Birch, Dorothy, Coulton, Simon, Bland, Martin, Cassidy, Paul, Dale, Veronica, Deluca, Paolo, Gilvarry, Eilish, Godfrey, Christine, Heather, Nick, Kaner, Eileen, McGovern, Ruth, Myles, Judy, Oyefeso, Adenekan, Parrott, Steve, Patton, Robert, Perryman, Katherine, Phillips, Tom, Shepherd, Jonathan and Drummond, Colin (2014) Alcohol Screening and Brief Interventions for Offenders in the Probation Setting (SIPS Trial): a Pragmatic Multicentre Cluster Randomized Controlled Trial. *Alcohol and Alcoholism*, 49 (5). pp. 540-548. ISSN 0735-0414

Published by: Oxford University Press

URL: <http://dx.doi.org/10.1093/alcalc/agu046> <<http://dx.doi.org/10.1093/alcalc/agu046>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/17359/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Alcohol Screening and Brief Interventions for Offenders in the Probation Setting (SIPS Trial): a Pragmatic Multicentre Cluster Randomized Controlled Trial

Dorothy Newbury-Birch^{1,*}, Simon Coulton², Martin Bland³, Paul Cassidy⁴, Veronica Dale³, Paolo Deluca⁵, Eilish Gilvarry⁶,
Christine Godfrey³, Nick Heather⁷, Eileen Kaner¹, Ruth McGovern¹, Judy Myles⁸, Adenekan Oyefeso⁸, Steve Parrott³,
Robert Patton⁵, Katherine Perryman⁹, Tom Phillips^{5,10}, Jonathan Shepherd¹¹ and Colin Drummond⁵

¹Institute of Health and Society, Newcastle University, Richardson Road, Newcastle upon Tyne, NE2 4AX, UK, ²Centre for Health Service Studies, University of Kent, The Registry, Canterbury, CT2 7NZ, UK, ³Department of Health Sciences, University of York, Seebholm Rowntree Building, Heslington, York, YO10 5DD, UK, ⁴Teams Family Practice, Watson Street, Gateshead, NE8 2PQ, UK, ⁵Addictions Department, Institute of Psychiatry, King's College London 4 Windsor Walk, London, SE5 8AF, UK, ⁶Northumberland, Tyne & Wear NHS Foundation Trust, Newcastle and North Tyneside Addictions Service, Plummer Court, Carlisle Place, Newcastle upon Tyne, NE1 6UR, UK, ⁷School of Life Sciences, Northumbria University, Northumberland Building, Newcastle upon Tyne, NE1 8ST, UK, ⁸Division of Population Health Sciences and Education, St George's, University of London, Cranmer Terrace, London, SW17 0RE, UK, ⁹Institute of Population Health, University of Manchester, Oxford Road, Manchester, M13 9PL, UK, ¹⁰Humber NHS Foundation Trust, Beverley Road, Willerby, HU10 6ED, UK and ¹¹Violence Research Group, Cardiff University, Heath Park, Cardiff, Wales, CF14 4XY, UK

*Corresponding author: Institute of Health and Society, Newcastle University, Baddiley-Clark Building, Richardson Road, Newcastle NE2 4AX, UK. Tel.: +44-191-208-8500; Fax: +44-191-222-6043; E-mail: dorothy.newbury-birch@ncl.ac.uk

(Received 10 April 2014; first review notified 2 May 2014; in revised form 26 June 2014; accepted 26 June 2014)

Abstract — **Aim:** To evaluate the effectiveness of different brief intervention strategies at reducing hazardous or harmful drinking in the probation setting. Offender managers were randomized to three interventions, each of which built on the previous one: feedback on screening outcome and a client information leaflet control group, 5 min of structured brief advice and 20 min of brief lifestyle counselling. **Methods:** A pragmatic multicentre factorial cluster randomized controlled trial. The primary outcome was self-reported hazardous or harmful drinking status measured by Alcohol Use Disorders Identification Test (AUDIT) at 6 months (negative status was a score of <8). Secondary outcomes were AUDIT status at 12 months, experience of alcohol-related problems, health utility, service utilization, readiness to change and reduction in conviction rates. **Results:** Follow-up rates were 68% at 6 months and 60% at 12 months. At both time points, there was no significant advantage of more intensive interventions compared with the control group in terms of AUDIT status. Those in the brief advice and brief lifestyle counselling intervention groups were statistically significantly less likely to reoffend (36 and 38%, respectively) than those in the client information leaflet group (50%) in the year following intervention. **Conclusion:** Brief advice or brief lifestyle counselling provided no additional benefit in reducing hazardous or harmful drinking compared with feedback on screening outcome and a client information leaflet. The impact of more intensive brief intervention on reoffending warrants further research.

INTRODUCTION

There is evidence of an association between alcohol use and offending behaviour (Boden *et al.*, 2012). Alcohol has been found to be a factor in half of all violent crimes (Flatley *et al.*, 2010). In England and Wales, alcohol-related crime is estimated to cost society £11 billion (2010–2011 costs) (Alcohol Team Home Office, 2013). However, the precise relationship is complex (Plant *et al.*, 2002), with an intricate interplay between drinking patterns, the amount of alcohol consumed and individual and contextual factors (Graham *et al.*, 2012).

Hazardous drinking is a repeated pattern of drinking that increases the risk of future health problems, whereas harmful drinking is defined by the presence of these problems (Babor *et al.*, 1994). In the UK, higher rates of alcohol misuse have been found at various stages in the criminal justice system compared with the 20–30% observed in primary care populations (Funk *et al.*, 2005), with a majority of men and women reporting problematic alcohol use. This encompasses those arrested at police stations (59%) (Brown *et al.*, 2010), within probation settings 59–63% (Newbury-Birch *et al.*, 2009b; Orr *et al.*, 2013) and those in prison 60–81% (McMurrin and Cusens, 2005; Newbury-Birch *et al.*, 2009b; MacAskill *et al.*, 2011).

Alcohol screening and brief alcohol intervention is a secondary preventative approach (Kaner *et al.*, 2009), which involves the identification via screening of hazardous and harmful drinking and the delivery of an intervention aimed at reducing consumption and concomitant problems. There is a wealth of evidence in primary care for the effectiveness of alcohol screening and brief intervention in adults who are not seeking

treatment for alcohol-related problems (Moyer *et al.*, 2002; Bertholet *et al.*, 2005; Kaner *et al.*, 2007), whereas the evidence in the other health-care settings such as Emergency Departments is conflicting (Drummond *et al.*, 2014). However, there has been very little research carried out in criminal justice settings. A recent quasi-experimental study in eight police custody settings in England found reduced alcohol consumption in those who got a brief intervention; however, those in the intervention group were statistically more likely to be arrested at 6 months post intervention (McCracken *et al.*, 2012). However, just 34% of participants were followed up at 12 months (McCracken *et al.*, 2012), and these were compared with a retrospective control group. A Scottish pilot trial in a community-based criminal justice settings of alcohol brief advice compared with delivery of an information booklet could not make any conclusions for effectiveness because of low follow-up rates of 20% at 6 months (Orr *et al.*, 2013). Nevertheless, brief alcohol intervention in a magistrate court setting resulted in lower levels of injury at 12 months compared with controls (Watt *et al.*, 2008).

Thus, the criminal justice setting represents important opportunity for alcohol intervention work and that could impact upon consumption and offending behaviour. However, more evaluative studies with robust methodological designs are required if we are to provide offender services with clear evidence of impact. We carried out a pilot study in 2007 that evaluated the most feasible criminal justice setting in which to conduct a randomized control trial (RCT) of alcohol screening and brief interventions and found that the recruitment, consent and willingness to participate rates were highest in the probation setting (Coulton *et al.*, 2012). We therefore carried out a pragmatic

cluster RCT of the effectiveness of two different brief intervention strategies compared with a control condition of feedback on screening outcome and a client information leaflet at reducing hazardous or harmful drinking in the English probation setting.

METHODS

The full trial protocol for the present study has been published (Newbury-Birch *et al.*, 2009a).

Study design

A prospective factorial pragmatic cluster RCT with randomization at the level of offender manager. Randomization was conducted by a secure remote randomization service using random permuted blocks and was stratified by geographical region. The research team was blind to the allocation schedule. Offender managers were randomly allocated initially to using one of two screening tools: Fast Alcohol Screening Test (FAST) (Hodgson *et al.*, 2002) or Modified Single Alcohol Screening Question (M-SASQ) (Canagasaby and Vinson, 2005) and then to one of the three additive conditions: feedback on screening outcome plus the delivery of an information leaflet; 5 min of brief advice; 20 min of brief lifestyle counselling delivered on a subsequent occasion. The study received ethical approval from the Multicentre Research Ethics Committee (08/HO903/2) and was conducted in accordance with the Helsinki Declaration.

Sample size

Due to the paucity of comparable research in criminal justice settings, the sample size calculation was based on outcomes reported in health-care settings, particularly primary care. Moyer *et al.* (2002) suggest that a clinically important difference in negative status on Alcohol Use Disorders Identification Test (AUDIT) between brief intervention and control is of the order 13%: 5% in the control group and 18% in the brief intervention group. To detect a difference of this magnitude at the 5% significance level with 80% power, for a two-sided test, 109 offenders in each of the three groups are required (a total of 327 offenders) (Moyer *et al.*, 2002).

Our experience with other studies of alcohol using populations (UKATT Research Team, 2005; Coulton *et al.*, 2006) and hard-to-engage populations (Gilbert *et al.*, 2012) would suggest that with assiduous follow-up regimes the loss to follow-up at 6 months should not exceed 25%, and so the sample required was inflated to 145 in each group. As the study is a cluster RCT, we adjusted the sample size for any potential design effect. We assumed that the intra-class correlation coefficient should be similar to primary care at 0.04. Assuming a cluster size of the order of five offenders per offender manager, this inflates the sample size calculation by a factor of 1.1, requiring 160 offenders in each group (a total of 480 offenders) with an expectation that at least 365 would be followed up at 6 months.

Participants

Offender managers are responsible for the high-quality assessment, robust management of contact and behaviour and provision of well-targeted interventions to offenders. They provide the

link for other organizations to work in an integrated manner (www.parliament.uk, 2011). Their work is subject to national standards (www.parliament.uk, 2011). Offender managers were recruited between May 2008 and July 2009 across three geographical regions of England: the North East, South East and London. Offender managers were considered eligible if they delivered routine probation services and were not involved in any other alcohol screening and brief intervention programme.

Individual offenders were eligible if they screened positive using the allocated screening instrument, were 18 years of age or more, were resident within 20 miles of the offender manager's office and were able to understand English sufficiently to complete the study questionnaire. Offenders were excluded if they were already receiving treatment for alcohol use, were intoxicated at assessment, were severely mentally impaired or were of no fixed abode. Offenders were provided with a £10 gift voucher for completing questionnaires at each data collection point. After discussions with senior management in the probation departments, probation offices were offered £1000 to participate in the study and recruit the required number of offenders (5 per offender manager) or alternatively individual offender managers were offered a £20 gift voucher for each offender recruited.

Screening

Offenders were screened using one of the two short validated alcohol screening tools. The M-SASQ asked 'How often do you have X or more standard drinks on one occasion?' where $X = 6$ for women and 8 for men, with monthly, weekly, daily or almost daily considered a positive screen (Canagasaby and Vinson, 2005). FAST is a four-question tool that scores between 0 and 16 with a score of 3 or more being considered a positive screen (Hodgson *et al.*, 2002). The results relating to the different screening tools will be reported elsewhere.

Interventions

Client information leaflet (control group)

Offenders received simple feedback on their screening outcome and an alcohol information leaflet from the offender manager developed by the Department of Health in England (www.sips.iop.kcl.ac.uk/pil.php). This 16-page leaflet describes the effects of alcohol on health and well-being and shows the number of units contained in popular alcoholic drinks to help the reader understand how much they were drinking. Contact details of local alcohol treatment agencies, Drinkline and websites were also provided.

Brief advice

Offender managers provided offenders with feedback on screening outcome and the client information leaflet plus 5 min of structured personalized brief advice based on the 'How much is too much?' simple structured advice tool developed as part of the UK version of the Drink-Less brief intervention programme (McAvoy *et al.*, 1997). In addition to providing specific details about the health and social consequences of hazardous and harmful drinking, offenders were shown a gender-specific graph which indicated that their drinking exceeded that of most of the population, and a list of benefits that would result from reduced drinking. Offenders were then taken through a menu of techniques to help reduce drinking and asked to consider a personal target for an achievable reduction.

Brief lifestyle counselling

Offender managers gave offenders feedback on screening outcome, the client information and structured brief advice in the initial meeting. Offenders were then asked to return for an appointment with an alcohol health worker for a 20-min session of brief lifestyle counselling. This was based on the 'How much is too much?' extended brief intervention tool developed as part of the UK version of the Drink-Less brief intervention programme (McAvoy *et al.*, 1997), which was modified in line with motivational interviewing principles outlined in the health behaviour change manual (Rollnick *et al.*, 1999). During the intervention, offenders described their typical drinking day and rated the importance of changing their drinking and their confidence about changing their drinking on a 10-point scale (where a higher number indicated greater importance or confidence and vice versa). The alcohol health worker then worked with these ratings to establish why they were at the current level and how they might be increased to a higher point, before eliciting both pros and cons of drinking and finally working through a six-step plan to help reduce drinking levels.

All interventions were coded for the presence of Behaviour Change Techniques (BCTs) by one of the authors (K.P.) who is a trained coder in the use of BCT taxonomy V1 (Michie *et al.*, 2011, 2013) (see www.sips.iop.kcl.ac.uk for further details of the screening tools, intervention materials and coding for BCTs used in this study).

Training and support

Offender managers were trained only to use the screening tool and intervention in the condition they had been allocated. All offender managers were trained by a member of the research team to implement alcohol screening and deliver the client information leaflet in accordance with the trial protocol. Offender managers allocated to brief advice received an additional 60-min training session delivered by an experienced alcohol health worker. The standardized training involved an interactive presentation with role play and feedback to develop skills and confidence in delivering the intervention. Ongoing weekly support through visits and telephone calls occurred during the trial.

Alcohol health workers delivering brief lifestyle counselling received formal training that consisted of four main elements: orientation to working in probation settings; workshops with interactive presentations and role play; recorded simulated consultations with trained actors; continuous clinical supervision. As the alcohol health workers were required to demonstrate adherence to the brief lifestyle counselling protocol and reach a required standard of competence prior to intervention delivery, simulated consultations were recorded and assessed by clinical supervisors using the Behaviour Change Counselling Index (Lane, 2002). They did not deliver interventions until they reached the correct standard. All alcohol health workers received ongoing weekly clinical supervision and support throughout the study.

Study measures

The primary outcome measure was hazardous or harmful drinking status at 6 months as measured by the AUDIT questionnaire (Saunders *et al.*, 1993). A score of ≥ 8 on this test

indicates hazardous or harmful drinking or possible dependent drinking, with a sensitivity of 92% and specificity of 94% (Saunders *et al.*, 1993). The primary outcome was negative status (≥ 8) at 6 months, which indicated a shift to lower risk drinking. Offenders also completed the Euroqol EQ-5D, which is a measure of health-related quality of life. EQ-5D has a ceiling of 1.0, which represents full quality of life, and the higher the score (max 1.0) the greater the quality of the individual's life (The EuroQol Group, 1990). Other tools completed were a short service use questionnaire to capture visits to NHS and other services and a modified Readiness Ruler that measured readiness to change drinking behaviour (Heather *et al.*, 2008).

Follow-up

At 6 and 12 months after the intervention, researchers who were blinded to the allocated condition contacted the offenders by telephone or post. Researchers administered the same instruments as at baseline plus the *Alcohol Problems Questionnaire* (Drummond, 1990). A short patient satisfaction questionnaire (Marshall and Hays, 1994) was also administered at 12 months. Police National Computer (PNC) conviction data were used as a measure of recidivism (whether offenders had been convicted of any offence) between recruitment and 12 months later.

Analysis

The primary analysis was by intention to treat, whereby offenders were analysed as members of their allocated group irrespective of intervention received. The primary outcome measure was analysed using logistic regression adjusting for known covariates: baseline AUDIT score, age and gender. In a factorial study, we tested for any interaction between screening tool and allocated intervention and, if the interaction was found to be significant, it was included in the model. To address the clustered design of the study, we employed the Huber–White sandwich estimation approach to derive robust standard errors. The impact of missing data was assessed using multiple imputation and sensitivity analysis undertaken to assess the potential influence of missing data. As no influence of missing data was observed, the results presented are those without imputation. Odds ratios and associated 95% confidence intervals were generated.

A secondary, per protocol analysis that included just those who received their allocated treatment was undertaken on the primary and dichotomous outcome measures. For continuous outcomes, we assessed distributional assumptions and where necessary undertook statistical transformations to conduct a linear regression adjusting for known covariates. Readiness to change analysis employed a similar method but used an ordinal regression modelling approach. All analysis was conducted using STATA v10.

RESULTS

Recruitment and follow-up

Of the 227 offender managers randomized, 172 (76%) were trained and of these 131 (76%) recruited at least one offender. The CONSORT diagram for the trial is shown in Fig. 1. Offender managers approached a total of 976 potential




Fig. 1. Consort statement indicating actual recruitment and intervention by screening approach and intervention.

offenders; of whom, 860 were eligible and agreed to undergo alcohol screening (88%). Of the 116 offenders who were ineligible, 52 (44.8%) were seeking treatment for alcohol problems; 13 (11.2%) were already taking part in research; 23 (19.8%) were severely unwell; 10 (8.6%) were intoxicated and 18 (15.5%) had no fixed abode. Of the 860 who were eligible to enter the trial, 574 (67%) screened positive using the M-SASQ or FAST. The numbers of individuals who screened positive were similar across intervention groups (client information leaflet: 196; brief advice: 193; brief lifestyle counselling: 185). A total of 525 offenders consented to the trial (91% of those who screened positively) with similar numbers across groups (client information leaflet: 184; brief advice: 178; brief lifestyle counselling: 163) (Fig. 1) which

was greater than the estimated sample size. All offenders received the client information leaflet; 97% of those in the brief advice group received the brief advice. In the counselling group, 99% (162) received brief advice but only 41% (67) subsequently received brief lifestyle counselling. The overall follow-up rate at 6 months was 68% (355), which was slightly lower than the sample size assumption of 75% ($n = 365$). The follow-up rates did not differ significantly across groups: client Information leaflet ($n = 120$; 65%), brief advice ($n = 123$; 69%) and brief lifestyle counselling ($n = 112$; 69%). At 12 months, 315 of the original 525 (60%) were followed up: client information leaflet group 102 (56%), brief advice group 115 (65%) and brief lifestyle counselling group 97 (60%).

Sample characteristics

The mean age of offenders was 31.0 years (SD 10.9) and was similar across groups. The majority were male (85%), white (76%) and single (67%). Less than half continued their education after 16 years of age (45%). Sixteen per cent possessed a degree or equivalent professional qualification. Almost 80% were current smokers. Mean AUDIT score at baseline was 16.07 (SD 8.57), and the mean baseline EQ-5D score was 0.85 (SD 0.22). AUDIT scores at baseline suggested that 43% of offenders consumed alcohol at harmful or possibly dependent levels (Table 1).

Primary outcome

AUDIT negative status increased in all three groups between baseline and 6 months from 17.7 to 29.1% in the client information leaflet group, 12.4 to 23.6% in the brief advice and 9.9 to 19.8% in the brief lifestyle counselling group. There were no significant differences between those in the brief advice and client information leaflet groups (OR = 0.80; 95% CI = 0.39–1.62) or brief lifestyle counselling compared with client information leaflet groups (OR = 0.73; 95% CI = 0.34–1.53) (Table 2). A per protocol analysis also showed no significant differences in this outcome between the groups.

Secondary outcomes

An analysis of AUDIT negative status at 12 months indicated no significant differences between the three groups. The odds ratio of being AUDIT negative for brief lifestyle counselling compared with client information leaflet was 0.70 (95% CI = 0.34–1.47) and for brief advice compared with client information leaflet was 1.11 (95% CI = 0.53–2.33) (Table 2).

The mean AUDIT score at 6 months was 13.96 (SE 0.78) for client information leaflet, 14.27 (SE 0.54) for brief advice and 14.22 (SE 0.77) for brief lifestyle counselling. The mean difference for brief advice compared with the client information leaflet was not significant (0.306; 95% CI = -1.60 to 2.21) as was the mean difference for brief lifestyle counselling compared with client information leaflet (0.25; 95% CI = -1.91 to 2.42). A similar pattern was observed for the AUDIT score at 12 months (Table 2).

No significant differences were observed at either 6 or 12 months for quality of life, alcohol-related problems, readiness to change or satisfaction with care (at 12 months) (Tables 3–6).

Recidivism data were obtained from 511 offenders (97% in both the client information leaflet and brief advice group and 98% in the brief lifestyle counselling group). Of the whole sample, 42% of offenders were convicted of an offence in the year following intervention. Differences were observed in the proportions in each group convicted (client information leaflet: 50%; brief advice: 36%; brief lifestyles counselling: 38%). The odds ratio of receiving a conviction was significantly lower in the brief advice (OR = 0.50; 95% CI = 0.33–0.80) and brief lifestyle counselling (OR = 0.54; 95% CI = 0.33–0.89) groups compared with the client information leaflet group (Table 7).

DISCUSSION

This is the first large-scale pragmatic RCT to evaluate alcohol screening and brief intervention in the probation setting. The

Table 1. Sample characteristics and AUDIT scores/categories at baseline

	Client information leaflet (N = 184)		Brief advice (N = 178)		Brief lifestyle counselling (N = 163)		Total (N = 525)		Intraclass correlation coefficient (SE)
	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	
Age in years	184	31.6 (10.9)	177	30.8 (10.8)	163	30.5 (11.1)	524	31.0 (10.9)	
EQ-5D	178	0.84 (0.24)	171	0.82 (0.22)	157	0.88 (0.21)	506	0.85 (0.23)	
Baseline AUDIT score	181	15.40 (8.33)	178	16.29 (8.80)	161	16.58 (8.58)	520	16.07 (8.57)	0.165 (0.048)
		%		%		%		%	
Baseline AUDIT status	32	17.7	22	12.4	16	9.9	70	13.5	0.074 (0.041)
	149	82.3	156	87.6	145	90.1	450	86.5	
AUDIT score category	32	17.7	22	12.4	16	9.9	70	13.5	0.143 (0.047)
	75	41.4	80	44.9	72	44.7	227	43.7	
	74	40.9	76	42.7	73	45.3	223	42.9	
	160/184	87.0	145/178	81.5	142/163	87.1	447/525	85.1	
Gender	140/184	76.1	138/178	77.5	122/163	74.8	400/525	76.2	
Ethnicity	125/184	67.9	125/178	70.2	103/163	63.2	353/525	67.2	
Marital status	84/182	46.2	77/178	43.3	74/163	45.4	235/523	44.9	
Education after 16 years	25/184	13.6	26/178	14.6	30/161	18.6	81/523	15.5	
Possess degree or equivalent	150/184	81.5	139/177	78.5	126/162	77.8	415/523	79.3	
Smoke tobacco	39	21.3	35	19.7	43	26.5	117	22.4	
Readiness Ruler	37	20.2	37	20.8	34	21.0	108	20.7	
	39	21.3	33	18.5	18	11.1	90	17.2	
	68	37.2	73	41.0	67	41.4	208	39.8	

Table 2. AUDIT scores and categories by condition at 6 and 12 months

	Client information leaflet		Brief advice/Client information leaflet		Brief lifestyle counselling		Brief lifestyle counselling/Client information leaflet		Intraclass correlation coefficient (SE)
	N (%)	Mean (SE)	OR (95% CI)	P-value	N (%)	Mean (SE)	OR (95% CI)	P-value	
AUDIT negative—Month 6	34/117 (29.1)	30/127 (23.6)	0.80 (0.39–1.62)	0.54	21/106 (19.8)	21/106 (19.8)	0.73 (0.34–1.53)	0.40	0.015 (0.053)
AUDIT negative—Month 12	34/102 (33.3)	37/111 (33.3)	1.11 (0.53–2.33)	0.79	25/95 (26.3)	25/95 (26.3)	0.70 (0.34–1.47)	0.35	0.031 (0.062)
	Mean (SE)	Mean (SE)	Mean difference (95% CI)	P-value	Mean (SE)	Mean (SE)	Mean difference (95% CI)	P-value	
AUDIT score—Month 6	13.96 (0.78)	14.27 (0.54)	0.306 (–1.60 to –2.21)	0.75	14.22 (0.77)	14.22 (0.77)	0.25 (–1.91 to –2.42)	0.82	0.001 (0.052)
AUDIT score—Month 12	12.18 (0.84)	13.10 (0.65)	0.91 (–1.16 to –2.99)	0.38	12.81 (0.80)	12.81 (0.80)	0.63 (–1.62 to –2.88)	0.58	0.042 (0.062)

results highlighted 67% of offenders screened positive on the M-SASQ or the FAST, which were slightly higher than those found in the probation setting in the UK previously using the AUDIT (59–63%) (Newbury-Birch *et al.*, 2009b; Orr *et al.*, 2013), but lower than in our pilot study (73%) (Coulton *et al.*, 2012). These figures may be an underestimate, as those already seeking help for alcohol problems were not eligible for the study. The prevalence of harmful or probably dependent drinkers (43%) was much higher in this population compared with 4% reported in the general population in England (Drummond *et al.*, 2004).

This study and the results of our pilot study (Coulton *et al.*, 2012) suggest that opportunistic screening and interventions are acceptable in this population, with a high proportion of offenders willing and able to participate. It is clear in this study, as with other studies in health settings (Crawford *et al.*, 2004; Kaner *et al.*, 2013), that a minority of offenders attend follow-up counselling. This was a pragmatic study; and this level of return for a subsequent session was outwith usual activity in probation and the return visit was entirely voluntary.

An increase in the proportion of AUDIT negative status at follow-up occurred in all three groups, but the difference between the groups was not statistically significant either in intention to treat or per protocol analysis on any of the outcome measures. Thus, we would not recommend the implementation of more complex brief interventions in this setting. Nevertheless, the large proportion of offenders who were drinking at the higher end of the AUDIT spectrum suggests a need for the development and evaluation of interventions that address the more severe cases in criminal justice settings, and this issue warrants further research.

We used PNC data to measure recidivism at 12 months post intervention (compared with 12 months pre-intervention) as other studies have done (Watt *et al.*, 2008). The convention for reconviction studies is often 2 years (Friendship *et al.*, 2002); however, it has been argued that the length of the follow-up period is entirely arbitrary (Maltz, 1984). It is important however, as we did, to give enough time for the PNC data to be updated (usually 3 months) before requesting the download to ensure that the data relate to a whole year. We found a significant difference in recidivism rates between the groups, with those randomized to brief advice and brief lifestyle counselling significantly less likely to be convicted in the 12 months after intervention than those randomized to the client information leaflet group. This finding is different to previous work in police settings where an increase in recidivism was found in those that received an alcohol intervention (McCracken *et al.*, 2012). This discrepancy between the two studies could be as a result of them being focused at different points in the criminal justice system. However, the reduction in recidivism in the more intensive intervention groups in the current study was found in the absence of significant differences in drinking status between the groups. This unexpected finding requires further investigation to establish whether it is a real effect (valid finding) and, if so, to explore possible explanations and/or mechanisms. Both the brief advice and brief lifestyle counselling interventions encouraged reduced consumption but also promoted awareness of risks relating to excessive alcohol use, including the risk of offending whilst under the influence of alcohol. It could be that this increased awareness resulted in a significant change in offending behaviour rather than consumption *per se* or it may be that offending is linked to

Table 3. Mean APQ (alcohol-related problems) score by treatment group

Treatment group	APQ score		Mean difference (95% CI) treatment group—client information leaflet	P-value
	n	Mean (SD)		
Month 6				
Client information leaflet	120	4.47(4.52)	—	
Brief advice	131	5.34 (5.06)	0.87 (−0.51, 2.24)	0.21
Brief lifestyle counselling	109	4.76 (5.03)	0.29 (−1.23, 1.80)	0.71
Overall test				0.41
Month 12				
Client information leaflet	103	4.38 (4.80)	—	
Brief advice	115	4.91 (5.03)	0.53 (−0.94, 2.00)	0.48
Brief lifestyle counselling	99	4.48 (5.18)	0.10 (−1.57, 1.78)	0.90
Overall test				0.72

Table 4. Results for readiness to change by condition over time

	Never think about drinking less	Sometimes think about drinking less	I have decided to drink less	Already trying to cut down	Ordered odds (95% CI) compared with client information leaflet	P-value
6 months						
Client information leaflet (n = 206)	28 (24.1%)	17 (14.7%)	26 (22.4%)	45 (38.8%)	—	—
Brief advice (n = 129)	17 (13.2%)	27 (20.9%)	27 (20.9%)	58 (45%)	1.37 (0.89–2.11)	0.15
Brief lifestyle counselling (n = 106)	19 (17.9%)	17 (16%)	21 (19.8%)	49 (46.2%)	1.38 (0.90–2.13)	0.15
12 months						
Client information leaflet (n = 101)	22 (21.8%)	16 (15.8%)	20 (19.8%)	43 (42.6%)	—	—
Brief advice (n = 112)	15 (13.4%)	30 (26.8%)	16 (14.3%)	51 (45.5%)	1.09 (0.64–1.87)	0.74
Brief lifestyle counselling (n = 307)	20 (21.3%)	16 (17%)	17 (18.1%)	41 (43.6%)	0.97 (0.53–1.78)	0.92

Table 5. EQ-5D (quality of life) scores at baseline, 6 months and 12 months

	EQ-5D baseline		EQ-5D Month 6		EQ-5D Month 12	
	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)
Client information leaflet	178	0.84 (0.24)	116	0.87 (0.21)	101	0.88 (0.20)
Brief advice	171	0.82 (0.22)	127	0.82 (0.25)	112	0.84 (0.23)
Brief lifestyle counselling	157	0.88 (0.21)	105	0.87 (0.21)	97	0.86 (0.22)
P-value ^a			0.23		0.19	
Brief advice—Client information leaflet Estimate (CI; P)			−0.040 (−0.089, 0.010; P = 0.12)		−0.035 (−0.086, 0.016; P = 0.18)	
Brief lifestyle counselling—Client information leaflet Estimate (CI; P)			−0.032 (−0.075, 0.012; P = 0.16)		−0.044 (−0.098, 0.010; P = 0.11)	

^aAdjusted for baseline EQ-5D score and using the Huber–White estimator to allow for clustering.

Table 6. Patient satisfaction at 12 months

	Score					Mean difference (95% CI), P-value					
	Client information leaflet		Brief advice		Brief lifestyle counselling		Brief advice/ Client information leaflet		Brief lifestyle counselling/ Client information leaflet		Overall P-value
	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	
General satisfaction	95	3.9 (0.63)	105	3.7 (0.65)	91	3.8 (0.80)	−0.19 (−0.38, 0.01) 0.07	−0.06 (−0.28, 0.15) 0.55		0.17	
General communication	96	4.0 (0.57)	108	3.9 (0.60)	90	3.9 (0.80)	−0.12 (−0.26, 0.03) 0.11	−0.09 (−0.28, 0.93) 0.32		0.25	
Interpersonal manner	96	4.0 (0.61)	107	3.9 (0.62)	91	3.9 (0.65)	−0.11 (−0.28, 0.05) 0.18	−0.08 (−0.26, 0.09) 0.35		0.40	

Table 7. Recidivism rates by category 12 months post screening

	Client information leaflet, <i>n</i> (%), <i>N</i> = 177	Brief advice, <i>n</i> (%), <i>N</i> = 172	Brief lifestyle counselling, <i>n</i> (%), <i>N</i> = 158	Brief advice/Client information leaflet		Brief lifestyle counselling/Client information leaflet	
				OR (95% CI)	<i>P</i> -value	OR (95% CI)	<i>P</i> -value
Reoffend	88 (49.7%)	62 (36.0%)	60 (38.0%)	0.50 (0.33–0.80)	0.005	0.54 (0.33–0.89)	0.016

Analysis adjusted for baseline AUDIT score, number of cautions in the previous year, ever been in custody, age, gender and screening method.

particular patterns of drinking (risky single occasion high intensity drinking) rather than overall consumption (such as average levels of drinking per week). More research is needed to ascertain the precise relationships between drinking level, drinking pattern, drinking context and offending behaviour.

As in other studies (Watt *et al.*, 2008; McCracken *et al.*, 2012), we included all crime rather than alcohol-related crimes such as violent offences. Watt *et al.*'s (2008) study of alcohol screening and brief intervention in the magistrates' court for alcohol-related offences found no differences between numbers of convictions for violent offences (52%) or non-violent offences (58%) amongst offenders at the 12-month follow-up. This may be because the measurement of whether an offence is alcohol related is a subjective measure made by the police and Crown Prosecution Service, and people may be convicted of offences other than those that they were originally arrested for.

Only 58% of randomized offender managers recruited at least one offender to the study, whereas 76% of those trained for the study recruited at least one offender. Although most staff were supportive of the study and saw working with offenders with alcohol issues as a legitimate part of their job, some staff were less interested. This lack of interest may have been caused by a number of contributing factors such as a perceived lack of importance of alcohol misuse in their workload, information overload, workload pressures and particularly maybe feeling aggrieved that they were told by their team leaders they were taking part in the study, which has been found in similar studies in the criminal justice system (Brown *et al.*, 2010). We operated an informal drop in/telephone support system once a week for anyone who was in the office at that time to see how they were getting on. In future work, more formal support mechanisms from both the research team and the line managers should be put in place. It was also the case that some offender managers were incentivized for participating in the study, and this raises the important issue of how interventions in criminal justice populations aimed at addressing health-related behaviours can be integrated into normal working practice.

This study involved opportunistic screening in a population not seeking alcohol treatment and so who may not be aware of their level of alcohol-related risk or harm. In light of the large field of evidence supporting the effectiveness of brief alcohol intervention (Kaner *et al.*, 2007), we did not feel it was ethically appropriate to use a no-input control group (i.e. withhold information about the screening outcome). Thus, we felt that the minimally acceptable control condition should be simple feedback following screening plus an alcohol information leaflet. There is a possibility that reductions observed across all three groups were due to regression to the mean or research participation (Hawthorn) effects, and this explanation cannot be

totally discounted in our study although the reduction in the proportion of AUDIT positives in the control arm of this study was similar to the effect sizes reported in the brief intervention arms of other trials and greater than those reported for no-treatment control conditions (Moyer *et al.*, 2002). However, it is also possible that the lack of differences between the groups was as a result of our control group including active ingredients of behaviour change such as screening, assessment, feedback and awareness raising via the written information and details of local treatment services.

Finally, it is possible that the lack of differences between interventions could have been due to the brief intervention protocols not being implemented correctly, despite the on-going support and clinical supervision during the trial, and analysis on this is currently underway. However, as a pragmatic study of effectiveness it aimed to reflect what would happen in real practice where intensive monitoring and supervision would be impractical.

The strengths of the study are as follows: this is the first large multicentre trial of screening and brief intervention in the criminal justice system; it was a pragmatic RCT in the typical probation setting; cluster randomization avoided the problems of contamination between interventions being delivered; the study used remote randomization and validated outcome measures of clinical relevance; rates of eligibility and consent were higher than in previous similar studies which add weight to the generalizability of the research. Although the follow-up rate at 6 months was 68% instead of the expected 76%, the higher than anticipated recruitment meant the numbers followed up at the primary end point were sufficient to detect the hypothesized differences stated in the power calculation. Furthermore, these were still substantially higher than the 20% follow-up rate at 6 months found in a recent study in the probation setting (Orr *et al.*, 2013).

In conclusion, this article adds to the small but growing evidence base relevant to brief interventions in the criminal justice system. However, the paucity of clear evidence on the effectiveness of alcohol screening and brief interventions in criminal justice system settings would suggest that the evidence for their implementation in this setting is currently not clear, and more research is needed around the interplay between brief interventions and AUDIT status and between brief interventions and recidivism.

Acknowledgements — We wish to thank all probation staff and offenders who have taken part in the trial.

Funding — This study was funded by the UK Department of Health. The views expressed herein do not necessarily reflect those of the Department of Health, the National Health Service, the National Offender Management Service or the Home Office.

Conflict of interest statement. None declared.

REFERENCES

- Alcohol Team Home Office. (2013) *Next Steps Following the Consultation on Delivering the Government's Alcohol Strategy*. London: Crown Copyright.
- Babor T, Campbell R, Room R *et al.* (1994) *Lexicon of Alcohol and Drug Terms*. Geneva: World Health Organization.
- Bertholet N, Daepfen J-B, Wietlisbach V *et al.* (2005) Brief alcohol intervention in primary care: systematic review and meta-analysis. *Arch Int Med* **165**:986–95.
- Boden J, Fergusson D, Horwood L. (2012) Alcohol misuse and violent behavior: findings from a 30-year longitudinal study. *Drug Alcohol Depend* **122**:135–41.
- Brown N, Newbury-Birch D, McGovern R *et al.* (2010) Alcohol screening and brief intervention in a policing context: a mixed methods feasibility study. *Drug Alcohol Rev* **29**:647–54.
- Canagasaby A, Vinson DC. (2005) Screening for hazardous or harmful drinking using one or two quantity-frequency questions. *Alcohol Alcohol* **40**:208–13.
- Coulton S, Drummond C, James D *et al.* (2006) Opportunistic screening for alcohol use disorders in primary care: comparative study. *Br Med J* **332**:511–7.
- Coulton S, Newbury-Birch D, Cassidy P *et al.* (2012) Screening for alcohol use in criminal justice settings: an exploratory study. *Alcohol Alcohol* **47**:423–7.
- Crawford MJ, Patton R, Touquet R *et al.* (2004) Screening and referral for brief intervention of alcohol-misusing patients in an emergency department: a pragmatic randomised controlled trial. *Lancet* **364**:1334–9.
- Drummond C. (1990) The relationship between alcohol dependence and alcohol-related problems in a clinical population. *Br Jnl Add* **85**:357–66.
- Drummond C, Oyefeso A, Phillips T *et al.* (2004) *Alcohol Needs Assessment Research Project (ANARP). The 2004 National Needs Assessment for England*. London: Department of Health and the National Treatment Agency.
- Drummond C, Deluca P, Coulton S *et al.* (2014) The effectiveness of alcohol screening and brief intervention in emergency departments: a multicentre pragmatic cluster randomized controlled trial. *PLoS One* In press.
- Flatley J, Kershaw C, Smith K *et al.* (2010) *Crime in England and Wales 2009/10*. London: Home Office.
- Friendship C, Beech A, Browne K. (2002) Reconviction as an outcome measure in research. *Br Jnl Crim* **42**:442–4.
- Funk M, Wutzke S, Kaner E *et al.* (2005) A multicountry controlled trial of strategies to promote dissemination and implementation of brief alcohol intervention in primary health care: findings of a World Health Organization collaborative study. *J Stud Alcohol* **66**:379–88.
- Gilbert H, Burns T, Copello A *et al.* (2012) Assertive Community Treatment for alcohol dependence (ACTAD): study protocol for a randomised controlled trial. *Trials* **13**:13–9.
- Graham L, Parkes T, McAuley A *et al.* (2012) *Alcohol Problems in the Criminal Justice System: An Opportunity for Intervention*. Denmark: World Health Organisation.
- Heather N, Smailes D, Cassidy P. (2008) Development of a Readiness Ruler for use with alcohol brief interventions. *Drug Alcohol Depend* **98**:235–40.
- Hodgson R, Alwyn T, John B *et al.* (2002) The FAST alcohol screening test. *Alcohol Alcohol* **37**:61–6.
- Kaner E, Beyer F, Dickinson H *et al.* (2007) Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database Syst Rev*: CD004148.
- Kaner E, Newbury-Birch D, Heather N. (2009) Brief interventions. In Miller P (ed). *Evidence-based Addiction Treatment*, Vol. Part 3 Treatment methods, Chapter 10. San Diego, CA: Elsevier.
- Kaner E, Bland M, Cassidy P. *et al.* (2013) Effectiveness of screening and brief alcohol intervention in primary care (SIPS trial): pragmatic cluster randomised controlled trial. *Br Med J* **346**:e8501. doi: 10.1136/bmj.e8501.
- Lane C. (2002) *The Behaviour Change Counselling Index (BECCI) Scale*. Cardiff, UK: University of Wales.
- MacAskill S, Parkes T, Brooks O *et al.* (2011) Assessment of alcohol problems using AUDIT in a prison setting: more than an 'aye or no' question. *BMC Pub Health* **11**:1–12.
- Maltz M. (1984) *Recidivism*. Orlando, FL: Academic Press.
- Marshall G, Hays R. (1994) *The Patient Satisfaction Questionnaire—Short Form (PSQ-18)*. Santa Monica, CA: Rand Corporation.
- McAvoy B, Kaner E, Haighton K *et al.* (1997) 'Drink-Less'—Marketing a brief intervention package in UK general practice. *Fam Pract* **14**:427–8.
- McCracken K, McMurrin M, Winlow S *et al.* (2012) *Evaluation of Alcohol Arrest Referral Pilot Schemes (Phase 2)*. London: Home Office. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/116267/occ102.pdf (1 April 2014, date last accessed).
- McMurrin M, Cusens B. (2005) Alcohol and violent and non-violent acquisitive offending. *Addict Res Theory* **13**:439–43.
- Michie S, Abraham C, Eccles M *et al.* (2011) Strengthening evaluation and implementation by specifying components of behaviour change interventions: a study protocol. *Imp Sci* **6**:1–8.
- Michie S, Richardson M, Johnston M *et al.* (2013) The Behavior Change Technique Taxonomy (v1) of 93 hierarchically-clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Beh Med* **46**:81–95.
- Moyer A, Finney JW, Swearingen CE *et al.* (2002) Brief interventions for alcohol problems: a meta-analytic review of controlled investigations in treatment-seeking and non-treatment-seeking populations. *Addiction* **97**:279–92.
- Newbury-Birch D, Bland M, Cassidy P *et al.* (2009a) Screening and brief interventions for hazardous and harmful alcohol use in probation services: a cluster randomised controlled trial protocol. *BMC Pub Health* **9**:418.
- Newbury-Birch D, Harrison B, Brown N *et al.* (2009b) Slashed and Sentenced: a prevalence study of alcohol use disorders amongst offenders in the North East of England. *Int J Pris Health* **5**:201–11.
- Orr K, McAuley A, Graham L *et al.* (2013) Applying and Alcohol Brief Intervention (ABI) model to the community justice setting: learning from a pilot project. *Crim Crim Jus* **0**:1–19.
- Plant M, Plant M, Thornton C. (2002) People and places: some factors in the alcohol violence link. *J Subst Abuse* **7**:207–13.
- Rollnick S, Mason P, Butler C. (1999) *Health Behaviour Change: A Guide for Practitioners*. Edinburgh: Churchill Livingstone.
- Saunders JB, Aasland OG, Babor TF *et al.* (1993) Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption. *Addiction* **88**:791–804.
- The EuroQol Group. (1990) "EuroQol—a new facility for the measurement of health-related quality of life". *Health Policy* **16**:199–208.
- UKATT Research Team. (2005) Effectiveness of treatment for alcohol problems: findings of the randomised UK alcohol treatment trial (UKATT). *Br Med J* **331**:541.
- Watt K, Shepherd J, Newcombe R. (2008) Drunk and dangerous: a randomised controlled trial of alcohol brief intervention for violent offenders. *J Exp Crim* **4**:1–19.
- www.parliament.uk. (2011) *The Role of the Probation Service—Justice Committee*. London: Parliamentary copyright, 2011.