



# ALDH2 polymorphism and alcohol-related cancers in Asians: a public health perspective

Jeffrey S. Chang<sup>1\*†</sup>, Jenn-Ren Hsiao<sup>2</sup> and Che-Hong Chen<sup>3\*†</sup>

## Abstract

The occurrence of more than 200 diseases, including cancer, can be attributed to alcohol drinking. The global cancer deaths attributed to alcohol-consumption rose from 243,000 in 1990 to 337,400 in 2010. In 2010, cancer deaths due to alcohol consumption accounted for 4.2% of all cancer deaths. Strong epidemiological evidence has established the causal role of alcohol in the development of various cancers, including esophageal cancer, head and neck cancer, liver cancer, breast cancer, and colorectal cancer. The evidence for the association between alcohol and other cancers is inconclusive. Because of the high prevalence of *ALDH2\*2* allele among East Asian populations, East Asians may be more susceptible to the carcinogenic effect of alcohol, with most evidence coming from studies of esophageal cancer and head and neck cancer, while data for other cancers are more limited. The high prevalence of *ALDH2\*2* allele in East Asian populations may have important public health implications and may be utilized to reduce the occurrence of alcohol-related cancers among East Asians, including: 1) Identification of individuals at high risk of developing alcohol-related cancers by screening for *ALDH2* polymorphism; 2) Incorporation of *ALDH2* polymorphism screening into behavioral intervention program for promoting alcohol abstinence or reducing alcohol consumption; 3) Using *ALDH2* polymorphism as a prognostic indicator for alcohol-related cancers; 4) Targeting *ALDH2* for chemoprevention; and 5) Setting guidelines for alcohol consumption among *ALDH2* deficient individuals. Future studies should evaluate whether these strategies are effective for preventing the occurrence of alcohol-related cancers.

**Keywords:** Aldehyde dehydrogenase 2, Alcohol, Cancer, Prevention, Public health

## Background

### Alcohol consumption in the world

According to the 2010 estimate published by the World Health Organization, the global average alcohol consumption for individuals aged 15 or older was 13.5 g of pure alcohol per day [1], which is approximately equivalent to 1 can of beer (12 oz. or 355 ml with 5.0% alcohol content), 1 glass of wine (5 oz. or 148 ml with 12% alcohol content), or 1 drink of distilled spirits (1.5 oz. or 44 ml with 40% alcohol content). The level of alcohol consumption varied by regions with the highest consumption found in Eastern

Europe and Russia, and the lowest consumption seen in Southeast Asia, Middle East, and North Africa [1]. Alcohol was consumed most commonly in the form of spirits (50.1%), followed by beer (34.8%) and wine (8.0%) [1]. The WHO projected that by 2025, alcohol consumption will continue to rise, particularly in the Western Pacific Region, which includes countries in the East Asia, unless public health policies are implemented to curb or reverse the trend [1].

### Alcohol-related disease burden

The occurrence of more than 200 diseases can be fully or partially attributed to alcohol drinking [2, 3]. Alcohol use was ranked as the fifth leading risk factor of disease globally in 2010, moving up from eighth place in 1990 [4]. Alcohol-related cancer, liver cirrhosis, and injury accounted for the majority of mortality attributed to alcohol consumption [3]. Together these three conditions

\* Correspondence: jeffreychang@nhri.org.tw; chehong@stanford.edu

†Equal contributors

<sup>1</sup>National Institute of Cancer Research, National Health Research Institutes, 1F, No 367, Sheng-Li Road, Tainan 70456, Taiwan

<sup>3</sup>Department of Chemical and Systems Biology, Stanford University, School of Medicine, 269 Campus Drive, CCSR Building, Rm. 3140, Stanford, CA 94305, USA

Full list of author information is available at the end of the article



caused 1,500,000 deaths worldwide, representing 2.8% of all deaths [3]. The global cancer deaths attributed to alcohol-consumption rose from 243,000 in 1990 to 337,400 in 2010 [3]. In 2010, cancer deaths due to alcohol consumption accounted for 4.2% of all cancer deaths [3]. All evidence indicates that the health impact of alcohol has been on the rise and this trend will likely to continue. In addition, the economic burden associated with alcohol consumption can be tremendous. The annual alcohol-related financial cost was estimated to be US\$249 billion for USA, CA\$14.5 billion for Canada, GB £21 billion for UK, AU\$35 billion for Australia, NT\$25.5 billion for Taiwan, and JP¥6300 billion for Japan [5–7].

#### Alcohol-related cancers: evidence from epidemiologic studies

Table 1 summarizes the association between alcohol and alcohol-related cancers

##### Head and neck cancer

The association between alcohol and head and neck cancer is one of the most studied and the results consistently showed an increased head and neck cancer risk associated with alcohol drinking. In a meta-analysis of 52 studies, Bagnardi et al. reported a positive trend in the association between alcohol drinking and head and neck cancer risk (light drinking: relative risk (RR) = 1.13, 95% confidence interval (CI): 1.00–1.26; moderate drinking: RR = 1.83, 95% CI: 1.62–2.07; heavy drinking: 5.13, 95% CI: 4.31–6.10) [8]. In Western countries, approximately 39% of head and neck cancer can be attributed to alcohol consumption (4% for alcohol drinking alone and 35% for the combined effect of alcohol and tobacco) [9].

##### Esophageal cancer

Alcohol consumption is a well-established risk factor of esophageal cancer, particularly esophageal squamous cell carcinoma. In the meta-analysis by Bagnardi et al, alcohol consumption showed a positive dose-response

relationship with esophageal squamous cell carcinoma (light drinking: RR = 1.26, 95% CI: 1.06–1.50; moderate drinking: RR = 2.23, 95% CI: 1.87–2.65; heavy drinking: RR = 4.95, 95% CI: 3.86–6.34) [8]. In a meta-analysis of 17 studies by Jarl et al. the risk of esophageal cancer was reduced to the level of never drinkers after 16.5 years of alcohol cessation, suggesting that alcohol cessation is effective in reducing the risk of esophageal cancer [10].

##### Liver cancer

Alcohol is a known risk factor of liver cancer. Chuang et al conducted a meta-analysis of 112 studies and showed an increased risk of liver cancer among ever drinkers compared to never drinkers (RR = 1.29, 95% CI: 1.16–1.42) [11]. They also reported a dose-response relationship for the positive association between alcohol consumption and liver cancer (12 g of alcohol per day: RR = 1.08, 95% CI: 1.04–1.11; 50 g of alcohol per day: RR = 1.54, 95% CI: 1.36–1.74; 75 g of alcohol per day: RR = 2.14, 95% CI: 1.74–2.62; 100 g of alcohol per day: RR = 3.21, 95% CI: 2.34–4.40; and 125 g of alcohol per day: RR = 5.20, 95% CI: 3.25–8.29) [11]. In addition, a synergistic interaction between alcohol drinking and hepatitis or diabetes on the risk of liver cancer was observed [11].

##### Breast cancer

Several meta-analyses have examined the association between alcohol consumption and breast cancer. Jayasekara et al. reported a weak non-linear positive association between alcohol drinking and breast cancer while Bagnardi et al. observed that the risk of breast cancer rose linearly with increasing level of alcohol consumption [8, 12]. Chen et al. reported that 5 g of ethanol from wine per day was associated with a reduced breast cancer risk while the risk of breast cancer began to increase with more than 10 g of ethanol from wine per day [13]. Overall, studies agreed that high level of alcohol consumption is associated with an increased breast

**Table 1** The association between alcohol and alcohol-related cancers

Cancer type	Level of association with alcohol	Summary of current literature
Head and neck cancer	Strong	With light drinking, the risk of head and neck cancer increases by 13%, while the risk increases by more than 5 times with heavy drinking [8].
Esophageal cancer	Strong	With light drinking, the risk of esophageal cancer increases by 26%, while the risk increases by 5 times with heavy drinking [8].
Liver cancer	Strong	Compared to never drinkers, the risk of liver cancer increases by 1.3 times among ever drinkers. With light drinking, the risk of liver cancer increases by 8%, while the risk increases by more than 5 times with heavy drinking [11].
Breast cancer	Moderate	Overall, studies agreed that high level of alcohol consumption is associated with an increased breast cancer risk while the risk of breast cancer at low level of alcohol consumption may require further investigation [8, 12, 13]
Colorectal cancer	Moderate	Light drinking is not associated with colorectal cancer while heavy drinking is associated with a 1.5 times increase in the risk of colorectal cancer [8, 12]

cancer risk while the risk of breast cancer at low level of alcohol consumption may require further investigation.

### **Colorectal cancer**

Studies have consistently found a positive association between alcohol consumption and colorectal cancer, although the increased risk is modest in magnitude [8, 12]. A meta-analysis by Bagnardi et al. reported a linear positive dose-response relationship between alcohol drinking and colorectal cancer (light drinking: RR = 0.99, 95% CI: 0.95–1.04; moderate drinking: RR = 1.17, 95% CI: 1.11–1.24; heavy drinking: 1.44, 95% CI: 1.25–1.65) [8]. Jayasekara et al. also reported a linear positive dose-response relationship between alcohol drinking and colorectal cancer and the highest level of alcohol consumption was associated with a 1.5 times increase in colorectal cancer risk compared to the lowest level of alcohol consumption [12].

The evidence presented so far only considers the average risk of cancer associated with alcohol consumption. Genetic background may determine an individual's susceptibility to the carcinogenic effect of alcohol. Individuals deficient in aldehyde dehydrogenase 2 (ALDH2), an enzyme that converts acetaldehyde, a carcinogenic metabolite of ethanol, to acetate, may have a higher risk of alcohol-related cancers. Based on overwhelming epidemiological evidences, the International Agency for Research on Cancer (WHO/IARC) has classified ethanol in alcoholic beverage as a Group 1 chemical, which is carcinogenic to humans [14, 15]. The IARC working group further emphasized the strong evidence of acetaldehyde derived from alcohol metabolism as the mechanistic basis in causing upper aerodigestive track (UADT) cancers in individuals with ALDH2 deficiency [14, 15].

### **The role of ALDH2 and its polymorphism**

In human, alcohol metabolism involves two major NAD-dependent enzymes, alcohol dehydrogenase (ADH) and aldehyde dehydrogenase (ALDH). Alcohol is first oxidized to acetaldehyde by ADH. Acetaldehyde is then oxidized to non-toxic acetate by ALDH for excretion. Among the different human ALDH isozymes, ALDH2, a mitochondria enzyme, is the most efficient enzyme to remove toxic acetaldehyde [16]. Greater than 90% of the consumed alcohol is detoxified by the first-pass metabolism, especially in the liver where ALDH2 is abundant, although ALDH2 is distributed and functional in many other major organs and tissues [17]. A striking genetic polymorphism that dramatically reduces ALDH2 enzyme activity and affects alcohol response is the variant *ALDH2\*2* allele. The variant *ALDH2\*2* allele is caused by a single point mutation (G to A) in exon 12, which leads to an amino acid substitution from glutamine to lysine (E487K) [18]. The normal *ALDH2\*1* allele

and the variant *ALDH2\*2* allele can be easily genotyped by the determination of single nucleotide polymorphism (SNP) rs671 from the human genome. Due to the tetrameric nature of the ALDH2 enzyme, the E487K mutation exhibits a dominant negative phenotype and affects both heterozygous and homozygous individuals who carry the variant allele. In carriers of the *ALDH2\*2/\*2* homozygous and *ALDH2\*1/\*2* heterozygous genotypes, the enzyme activity is nearly 0% and 17–38% of the normal activity, respectively [19]. Such dramatic reduction in the capacity of acetaldehyde clearance leads to accumulation of acetaldehyde in circulation even after a moderate amount of alcohol intake [20]. Hence, the *ALDH2\*2* variant causes the well-known Asian Alcohol Flushing Syndrome which is characterized by facial flushing, palpitation, tachycardia, nausea, and unpleasant feelings when alcohol is consumed by these individuals [21]. The *ALDH2\*2* variant is essentially absent among the Europeans, but is highly prevalent among East Asians [22]. Large scale genotyping and haplotype analysis of the *ALDH2* gene has traced the origin of the *ALDH2\*2* allele back to the ancient Pai-Yuei tribe about 2000–3000 years ago in Southeast China [23]. The *ALDH2\*2* allele likely dispersed from its origin toward East Asia with the highest frequencies in Southeastern coastal regions of China (e.g. Guangdong, Fujian provinces) and countries with historical Han migrations, such as Taiwan, Japan, Korea and Singapore [23, 24]. The prevalence of the *ALDH2\*2* variant varies from 28% (e.g. in Korea) to as high as 45% (e.g. in Taiwan) in populations of these regions [23] and an estimated 560 million East Asians are carriers of *ALDH2\*2* [25]. The ALDH2 deficiency is therefore one of the most common and genetically uniform enzymopathies in human carried by approximately 8% of the world population.

### **ALDH2 polymorphism and risk of alcohol-related cancers**

Many studies have investigated the role of *ALDH2* polymorphism and its interaction with alcohol consumption in the development of various cancers. The strongest and most consistent findings have been observed for head and neck cancer and esophageal cancer, while the evidence for other cancers is more limited (Table 2).

#### **Head and neck cancer**

In a meta-analysis of six case-control studies (945 cases and 2917 controls), Boccia et al. reported a reduced risk of head and neck cancer among individuals with *ALDH2\*2/\*2* genotype, likely due to the lower consumption of alcohol among these individuals [26]. In addition, *ALDH2\*1/\*2* genotype showed gene-environment interaction with the level of alcohol consumption [26]. Compared to *ALDH2\*1/\*1*, *ALDH2\*1/\*2* was associated with an increased risk of

**Table 2** The association between *ALDH2* polymorphism and alcohol-related cancers

Cancer type	Level of association with <i>ALDH2</i> polymorphism	Summary of current literature
Head and neck cancer	Strong	Most studies showed a synergistic interaction between <i>ALDH2</i> *2 allele and alcohol drinking to increase head and neck cancer risk [26–29].
Esophageal cancer	Strong	Overall, literature provides strong evidence to support the synergistic interaction between <i>ALDH2</i> *2 allele and alcohol consumption to increase the risk of esophageal cancer [31].
Liver cancer	Weak	Overall, studies to date have consistently shown that <i>ALDH2</i> *2 does not independently contribute to liver cancer risk. Whether <i>ALDH2</i> *2 can modify the association between alcohol consumption and liver cancer risk requires further investigation [32–41].
Breast cancer	Weak	Results from the only three studies published to date do not support a significant association between <i>ALDH2</i> *2 and breast cancer [42–44].
Colorectal cancer	Weak	The role of <i>ALDH2</i> *2 in the development of colorectal cancer is inconclusive and all of the studies conducted to date have a small sample size that resulted in insufficient statistical power and lack of precision [32, 45–47].

head and neck cancer among moderate drinkers (odds ratio (OR) = 1.68, 95% CI: 1.27–2.22) and heavy drinkers (OR = 3.57, 95% CI: 1.41–9.05), but not among never drinkers (OR = 0.97, 95% CI: 0.65–1.46) [26]. Three of the four studies published after the meta-analysis all showed a synergistic interaction between *ALDH2*\*2 allele and alcohol drinking to increase head and neck cancer risk [27–29] and only one study did not observe such interaction [30]. Overall, studies have indicated that *ALDH2* polymorphism can modulate the association between alcohol drinking and head and neck cancer risk.

#### Esophageal cancer

Zhao et al. conducted a meta-analysis of 31 case-control studies and found that *ALDH2*\*2/\*2 was associated with a reduced risk of esophageal cancer (OR = 0.69, 95% CI: 0.48–0.98) while *ALDH2*\*1/\*2 was associated with an increased esophageal cancer risk (OR = 2.34, 95% CI: 1.75–3.13) [31]. Individuals with *ALDH2*\*2/\*2 tend to avoid alcohol consumption due to the severe reaction after alcohol drinking and this is likely the explanation for their reduced risk of esophageal cancer. However, Zhao et al. reported that among alcohol drinkers, *ALDH2*\*2/\*2 was associated with an increased esophageal cancer risk (OR = 3.87, 95% CI: 1.67–8.96) compared to *ALDH2*\*1/\*1 [31]. In addition, the association between *ALDH2*\*1/\*2 and esophageal cancer became stronger with higher level of alcohol consumption (never drinker: OR = 1.21, 95% CI: 0.95–1.73; light drinker: OR = 3.79, 95% CI: 3.04–4.72; heavy drinker: OR = 6.50, 95% CI: 5.34–7.92) [31]. Overall, literature provides strong evidence to support the synergistic interaction between *ALDH2*\*2 allele and alcohol consumption to increase the risk of esophageal cancer.

#### Liver cancer

Among the 10 studies that examined the association between *ALDH2*\*2 and liver cancer risk, 8 found no significant association [32–39], 1 found an increased risk among individuals with at least 1 copy of the *ALDH2*\*2

allele [40], and 1 found an increased risk for heterozygous individuals only [41]. Among the six studies that examined the interaction between *ALDH2*\*2 and alcohol drinking on liver cancer risk, three found a synergistic interaction between *ALDH2*\*2 allele and alcohol consumption to increase liver cancer risk [36, 37, 40], while three found no such interaction [33, 34, 38]. Overall, studies to date have consistently shown that *ALDH2*\*2 does not independently contribute to liver cancer risk. Whether *ALDH2*\*2 can modify the association between alcohol consumption and liver cancer risk requires further investigation.

#### Breast cancer

There is limited evidence regarding the association between *ALDH2* polymorphism and risk of breast cancer. To date, only three studies have examined the role of *ALDH2*\*2 in the development of breast cancer and all found no association between *ALDH2*\*2 and risk of breast cancer [42–44]. In addition, Choi et al. and Kawase et al. examined the association between *ALDH2*\*2 and risk of breast cancer stratified by alcohol consumption status and observed no significant interaction between *ALDH2*\*2 and alcohol consumption on the risk of breast cancer [43, 44]. With only three studies published to date, it may be premature to rule out the role of *ALDH2*\*2 in the occurrence of breast cancer; however, evidence from the small number of published studies to date does not support a significant association between *ALDH2*\*2 and breast cancer.

#### Colorectal cancer

Four studies, all from Japan, have investigated the association between *ALDH2* polymorphism and colon cancer risk. Yokoyama et al. reported an increased risk of colon cancer (OR = 3.35, 95% CI: 1.51–7.45) associated with alcoholic carriers of *ALDH2*\*2 allele compared to alcoholics with homozygous wild type [32]. Murata et al. reported that alcohol showed a stronger dose-response

relationship with colon cancer among *ALDH2\*1/\*2* individuals than among *ALDH2\*1/\*1* individuals, although the results were not statistically significant; in addition, alcohol consumption was not associated with rectal cancer risk regardless of the *ALDH2* genotype [45]. Matsuo et al. reported that high level of alcohol consumption was associated with an increased rectal cancer risk but not colon cancer risk among individuals with *ALDH2\*1/\*2* [46]. Miyasaka et al. found no association between *ALDH2\*2* and colon cancer and did not assess the interaction between *ALDH2\*2* and alcohol drinking on colon cancer risk [47]. Overall, the role of *ALDH2\*2* in the development of colorectal cancer is inconclusive and all of the studies conducted to date have a small sample size that resulted in insufficient statistical power and lack of precision.

### Public health implications

Alcohol flushing and its associated unpleasant feelings due to acetaldehyde accumulation is a strong deterrent against heavy drinking and alcoholism for *ALDH2\*2* carriers [48]. Many studies have demonstrated the protective effect against alcohol addiction and abuse by *ALDH2* deficiency. For example, in 1982, Harada first showed that among Japanese alcoholics, only 2% of the subjects are *ALDH2* deficient [49]. However, influenced by social, cultural, and economic factors in the past few decades, such protection against alcohol dependence and alcohol abuse has gradually lost among the large populations of *ALDH2\*2* carriers. An alarming rise in the proportion of heavy drinkers who are carriers of *ALDH2\*1/\*2* genotype has been documented from the 1970 to 2010. In Japan, the percent of *ALDH2\*2* alcoholics was 2.5% in 1979, and has increased to 8.0% in 1986, 13.0% in 1992 [50], 13.0% in 1996–2000, 14.0% in 2001–2005, and 15.4% in 2006–2010 [51]. This rapid rise is further highlighted by a recent study in Tokyo area, showing that 26% of the heavy drinking men who consume >400 g of ethanol per week are *ALDH2\*1/\*2* heterozygotes [52]. In Taiwan, an estimation from a 1999 study indicated that 17% of the alcoholics are *ALDH2\*2* carriers [53]. It is therefore anticipated that, without intervention, the health risk and healthcare burden caused by heavy *ALDH2\*2* alcohol drinkers will become much more severe in the next few decades. The rapid and dangerous rise in alcohol consumption and dependence among *ALDH2\*2* carriers also underlines an urgent need for new public health policies and guidelines and active campaigns for public education and awareness in high *ALDH2\*2* prevalent countries. Given the strong association between *ALDH2* polymorphism and certain alcohol-related cancers, screening for *ALDH2\*2* allele may have several public health implications, including: 1) Identification of individuals at high risk of developing alcohol-related cancers; 2)

Incorporation of *ALDH2* polymorphism screening into alcohol cessation program for promoting alcohol abstinence or reducing alcohol consumption; 3) Using *ALDH2* polymorphism as a prognostic indicator for alcohol-related cancers; 4) Targeting *ALDH2* for chemoprevention; and 5) Setting guidelines for alcohol consumption among *ALDH2* deficient individuals.

### Identification of individuals at high risk of developing alcohol-related cancers

Early detection is the key to reduce cancer mortality and increase the chance for cure. However, it is not cost-effective to screen the entire population. Because cancer is a rare disease, even if a screening tool has a high sensitivity and specificity, the positive predictive value (PPV) (percent of individuals who test positive and actually have the disease) will still be low. For example, in 2012, East Asia was the region with the highest incidence of esophageal cancer with an incidence of 11 per 100,000 [54]. With this incidence, using a screening tool with 99% sensitivity and 99% specificity will generate a PPV of only 1.1%. This means that for every 100 positive cases detected by screening, only 1 case will actually have esophageal cancer. This is not cost-effective and valuable medical resources will be wasted. In addition, many individuals will undergo unnecessary medical procedures due to positive screening results. To make the screening more cost-effective, it would be important to identify the high-risk population to increase PPV. For example, if we can identify a population with an esophageal cancer incidence of 1 per 100, using a screening tool with 99% sensitivity and 99% specificity, the PPV will increase from 1.1 to 50% (1 of the 2 individuals with a positive screening result will actually have esophageal cancer), which is a substantial improvement in cost-effectiveness. Given the strong evidence for the synergistic interaction between *ALDH2\*2* allele and alcohol drinking in increasing the risk of several alcohol-related cancers, screening for the carriers of *ALDH2\*2* allele may be ideal for identifying individuals at high risk for alcohol-related cancers. For example, Yokoyama et al. have developed a health risk model for screening esophageal cancer [55, 56]. The model included information on alcohol drinking, *ALDH2* genotype or facial flushing after alcohol consumption, which is a physical symptom strongly associated with *ALDH2\*2* allele, cigarette smoking, and consumption of vegetables and fruits [55, 56]. Using this health risk model, Yokoyama et al. were able to identify individuals with higher risk of developing esophageal cancer, with an esophageal cancer detection of 2.9% in the high-risk group compared to 0.5% in the low-risk group [56]. Another study by Koyanagi et al. built a risk prediction model of UADT cancer that incorporated information on sex, age,

alcohol drinking, cigarette smoking, and *ALDH2* genotype and reported that the risk model had a good discriminative ability with an area under the curve of more than 0.8 [57]. In addition, they reported that heavy drinkers with *ALDH2*\*1/\*2 genotype have a 20% cumulative risk of developing UADT cancer by the age of 80 years, while other individuals have < 5% risk of developing UADT cancer by the age of 80 years [57]. These studies suggest that *ALDH2* genotype may be incorporated in a risk prediction model to identify individuals at high risk of developing alcohol-related cancers, particularly those cancers that are strongly influenced by gene-environment interaction between alcohol and *ALDH2*, such as head and neck cancer and esophageal cancer. For other alcohol-related cancers, including liver cancer, breast cancer, and colorectal cancer, for which the evidence for the involvement of *ALDH2* is more limited, more studies are needed to evaluate the interaction between *ALDH2* polymorphism and alcohol drinking on the risk of these alcohol-related cancers.

#### **Incorporation of *ALDH2* polymorphism screening into alcohol cessation program**

It is possible that the knowledge regarding *ALDH2* genotype and the associated disease risk may motivate individuals to reduce alcohol consumption. Hendershot et al. conducted a web-based genetic feedback intervention trial incorporating information on *ALDH2* genotype [58]. They recruited 200 northeast Asian American college students and randomized them to control and intervention groups. Individuals in the control group received a web-based feedback session that included normative information about behaviors of college students [58]. Individuals in the intervention group received different web-based feedback sessions according to their *ALDH2* genotype. Individuals with *ALDH2*\*1/\*1 received risk information on alcohol dependence while individuals with *ALDH2*\*1/\*2 received risk information on alcohol-related cancers [58]. One month after the web-based feedback session, individuals in the intervention group showed significant reduction in the frequency and the quantity of alcohol drinking while individuals in the control group showed no significant change in drinking behaviors [58]. This suggested that it may be feasible to incorporate information on *ALDH2* genotype and the associated disease risk into an alcohol cessation program to effectively reduce the consumption of alcohol. However, the use of genetic information to change health behaviors is complex and its effectiveness is still inconclusive [59]. For example, Smerecnik et al. conducted a meta-analysis of nine studies and found that genetic testing of genes associated with smoking-related diseases had only a short-term effect on risk perception and motivation to quit smoking, which did not last with longer

follow-ups [60]. More studies are needed to determine whether genetic testing of *ALDH2* will achieve a long lasting effect for reducing the consumption of alcohol. It is possible that the outward physical symptoms, including facial flushing, palpitation, tachycardia, and nausea, associated with *ALDH2*\*2 may serve as a strong reminder to maintain the motivation for reducing alcohol consumption. In addition, more investigations are needed to determine the optimal and culturally appropriate methods for delivering genetic information and the related disease risks.

#### ***ALDH2* polymorphism as a prognostic indicator for alcohol-related cancers**

*ALDH2* polymorphism may be used to predict the survival, recurrence, and development of second-primary or other alcohol-related cancers for patients with alcohol-related cancers, particularly for esophageal cancer and head and neck cancer. Kawakita et al. reported that both alcohol drinking and *ALDH2*\*2 were not independently associated with disease-free survival of head and neck cancer [61]. However, there was a significant positive dose-response between higher alcohol consumption and poorer disease-free survival of head and neck cancer among *ALDH2*\*1/\*1 individuals [61]. The exact biological mechanism underlying this relationship was not clear and the authors speculated that patients with *ALDH2*\*1/\*1 might continue to drink alcohol at a higher level after head and neck cancer treatment, resulting in their poorer outcomes [61]. Yokoyama et al. conducted a follow-up study of 100 alcoholic men diagnosed with esophageal squamous cell carcinoma and observed that during the follow-period, individuals with *ALDH2*\*1/\*2 were at higher risk of developing metachronous squamous cell carcinoma in the esophagus, oral cavity, pharynx and larynx (age-adjusted hazard ratio = 3.38, 95% CI: 1.45–7.85; alcohol-adjusted hazard ratio = 4.27, 95% CI: 1.42–12.89) compared to individuals with *ALDH2*\*1/\*1 [62]. Due to limited number of studies investigating the role of *ALDH2*\*2 in the survival or recurrence of alcohol-related cancers, it may be premature to consider its clinical application. However, once such role of *ALDH2*\*2 has been proven, the follow-up schedule and methods for alcohol-related cancer patients may be tailored according to their *ALDH2* genotypes to increase survival and improve early detection of cancer recurrence or second primary cancer.

#### **Setting guidelines for alcohol consumption among *ALDH2* deficient individuals**

If an individual does not drink alcohol, the best recommendation for alcohol consumption would be not to start. If one does drink, the Dietary Guidelines for Americans 2015–2020 recommends no more than 1

drink (14 g of pure alcohol) per day for women and 2 drinks for men [63]. The definition of low-risk alcohol consumption varies widely across different countries in the world ranging from 10 g per day to 42 g per day for women and 14 g per day to 56 g per day for men [64]. These guidelines do not consider the influence of *ALDH2* polymorphism on the association between alcohol drinking and cancer and the amount of alcohol that is considered low risk may not be suitable for individuals with *ALDH2* deficiency. For example, in a study by Lee et al., compared to never drinkers with *ALDH2*\*1/\*1, drinking < 30 g of alcohol per day was associated with a 2.2 times increase in esophageal cancer risk for those with *ALDH2*\*1/\*1, but the risk increased by 14.5 times and 17.3 times, respectively, for those with *ALDH2*\*1/\*2 and *ALDH2*\*2/\*2 [65]. This showed that even with the amount of alcohol considered as “light” or “moderate” drinking, the increase in esophageal cancer risk could be substantial for carriers of *ALDH2*\*2 allele. Therefore, when setting the level of low-risk alcohol consumption, it is important to consider the increased susceptibility to the carcinogenic effect of alcohol among carriers of *ALDH2*\*2 allele. This is particularly relevant for East Asian countries where the prevalence of *ALDH2*\*2 allele is high. More studies are needed to quantify the dose-response relationship between alcohol consumption and risk of various cancers by *ALDH2* genotypes, particularly for esophageal cancer and head and neck cancer, for both the risk is significantly increased by the synergistic interaction between alcohol consumption and *ALDH2*\*2 allele. Results from these studies will be important for experts in the field and public health officials to decide on the level of alcohol consumption that confers minimum acceptable risk.

#### **Chemoprevention by aldehyde dehydrogenase activators**

Exposure to acetaldehyde, a volatile, reactive and commonly present compound from either physiological conversion of alcoholic beverages or from the environmental sources (such as cigarette smoke [66], food items [67], automobile exhausts [68] etc.) is almost unavoidable. Considering the carcinogenicity of acetaldehyde and the significantly increased risk for head and neck cancer, esophageal cancer, and other cancers for *ALDH2*\*2 variant carriers, implementation of chemopreventive strategies may be a worthwhile effort for specific high-risk groups, such as *ALDH2*\*2 individuals who are heavy drinkers or smokers or industrial workers who have a higher acetaldehyde exposure burden. One of the chemopreventive strategies is enhancing the catalytic activity of the aldehyde dehydrogenase enzyme for faster acetaldehyde clearance. Recently, a group of novel small molecule ALDH enzyme activators have been discovered. These compounds may serve as pharmaceuticals

leads for the development of chemopreventive agents based on their enhanced capability of acetaldehyde removal. Alda-1 (N-(1,3-benzodioxol-5-ylmethyl)-2,6-dichlorobenzamide) was the first identified potent selective activator for *ALDH2* [69]. The compound has been tested and shown to be effective in several animal models of human diseases related to excessive aldehyde toxicity [70–72]. One of the unique features of Alda-1 is that the compound not only increases the catalytic activity of the wild type *ALDH2*\*1 enzyme, but can also correct the defect of the mutant *ALDH2*\*2 enzyme [69, 73]. This feature is particularly useful for the development of a chemopreventive drug specifically for the *ALDH2*\*2 carriers and the targeted high-risk human groups [74]. Another useful ALDH activator is Alda-89 (5-(2-propenyl)-1,3-benzodioxole) which was discovered as a selective *ALDH3A1* isozyme activator [75]. The *ALDH3A1* is not an enzyme that metabolizes acetaldehyde efficiently under normal physiological condition. However, with the addition of Alda-89, the catalytic activity of *ALDH3A1* toward acetaldehyde could be increased by at least 5 folds [75, 76]. In an acute alcohol intoxication animal model, it was demonstrated that Alda-89 was able to recruit *ALDH3A1* to assist the function of *ALDH2* for acetaldehyde detoxification in both wild type and *ALDH2*\*2 knock-in mice [76]. It is conceivable that ALDH activators, like Alda-1 or Alda-89, can be developed in various forms for rapid clearance of acetaldehyde regardless of its source or presence in either saliva or circulating blood. Such drugs could potentially have a great utility in reducing the risk of cancer or cancer recurrence, especially among the groups of high-risk individuals who are alcohol drinkers, cigarette smokers and carriers of *ALDH2*\*2 allele. Caution should also be given in the design of a chemopreventive program using aldehyde dehydrogenase activators, since it is not clear whether by enhancing ALDH activity, there will also be a risk of increased alcohol consumption among *ALDH2*\*2 subjects. Behavioral intervention studies and risk assessment on alcohol consumption and cessation similar to the study mentioned above by Hendershot et al. [58] should be recommended and incorporated in the design of such a chemopreventive program.

#### **Conclusions**

Given the rising consumption of alcohol around the world, the incidence of alcohol-related cancers will continue to rise, particularly in East Asian countries and regions with high concentrations of East Asian immigrants, where the percentage of *ALDH2* deficient individuals is high. Information regarding an individual's *ALDH2* genotype may help establish effective public health preventive strategies to reduce the occurrence and improve the survival of alcohol-related cancers.

Screening for carriers of *ALDH2\*2* allele may identify high-risk individuals for alcohol-related cancers. These high-risk individuals may be targeted for more frequent screening of alcohol-related cancers, health education, and alcohol cessation program. In addition, alcohol-related cancer patients carrying *ALDH2\*2* may be at higher risk for recurrence or developing second primary cancer and thus follow-up plan with more clinical visits may be required. Finally, chemo-preventive agent to restore the function of ALDH2 enzyme may be considered to prevent the occurrence or the recurrence of alcohol-related cancers among high-risk individuals.

#### Abbreviations

ADH: Alcohol dehydrogenase; ALDH: Aldehyde dehydrogenase; CI: Confidence interval; IARC: International Agency for Research on Cancer; OR: Odds ratio; PPV: Positive predictive value; RR: Relative risk; UADT: Upper aerodigestive tract.

#### Acknowledgements

Not applicable

#### Funding

Dr. Jeffrey S. Chang and Dr. Jenn-Ren Hsiao are supported by grants from the Establishment of Cancer Research System Excellence Program funded by the Ministry of Health and Welfare, Taiwan (MOHW106-TDU-B-211-144-004, MOHW105-TDU-B-212-134-013). Dr. Jeffrey S. Chang is also supported by intramural fund from the National Health Research Institutes (CA-106-SP-01). Dr. Che-Hong Chen is supported by a grant from NIH-NIAAA 11147 to Prof. Daria Mochly-Rosen, Stanford University.

#### Availability of data and materials

Not applicable

#### Authors' contributions

All authors conceived the content of the manuscript, JSC and CHC wrote the manuscript, all authors reviewed and approved the manuscript.

#### Competing interests

The authors declare that they have no competing interests.

#### Consent for publication

Not applicable

#### Ethics approval and consent to participate

Not applicable

#### Author details

<sup>1</sup>National Institute of Cancer Research, National Health Research Institutes, 1F, No 367, Sheng-Li Road, Tainan 70456, Taiwan. <sup>2</sup>Department of Otolaryngology, National Cheng Kung University Hospital, College of Medicine, National Cheng Kung University, 138 Sheng Li Road, Tainan 70456, Taiwan. <sup>3</sup>Department of Chemical and Systems Biology, Stanford University, School of Medicine, 269 Campus Drive, CCSR Building, Rm. 3140, Stanford, CA 94305, USA.

Received: 19 October 2016 Accepted: 26 February 2017

Published online: 03 March 2017

#### References

- World Health Organization. Global status report on alcohol and health 2014. Geneva: WHO Press; 2014.
- Rehm J. The risks associated with alcohol use and alcoholism. *Alcohol Res Health*. 2011;34:135–43.
- Rehm J, Shield KD. Global alcohol-attributable deaths from cancer, liver cirrhosis, and injury in 2010. *Alcohol Res*. 2013;35:174–83.
- Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, Amann M, Anderson HR, Andrews KG, Aryee M, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2224–60.
- Gilmore W, Chikritzhs T, Stockwell T, Jernigan D, Naimi T, Gilmore I. Alcohol: taking a population perspective. *Nat Rev Gastroenterol Hepatol*. 2016;13:426–34.
- Hsu WL. The economic costs attributable to alcohol consumption in Taiwan. Taipei: National Yang-Ming University; 2012. 204.
- Kanda H, Okamura T. The economic and medical costs of alcohol consumption in Japan. *West Indian Med J*. 2013;62:785–6.
- Bagnardi V, Rota M, Botteri E, Tramacere I, Islami F, Fedirko V, Scotti L, Jenab M, Turati F, Pasquali E, et al. Alcohol consumption and site-specific cancer risk: a comprehensive dose-response meta-analysis. *Br J Cancer*. 2015;112:580–93.
- Hashibe M, Brennan P, Chuang SC, Boccia S, Castellsague X, Chen C, Curado MP, Dal Maso L, Daudt AW, Fabianova E, et al. Interaction between tobacco and alcohol use and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *Cancer Epidemiol Biomarkers Prev*. 2009;18:541–50.
- Jarl J, Gerdttham UG. Time pattern of reduction in risk of oesophageal cancer following alcohol cessation—a meta-analysis. *Addiction*. 2012;107:1234–43.
- Chuang SC, Lee YC, Wu GJ, Straif K, Hashibe M. Alcohol consumption and liver cancer risk: a meta-analysis. *Cancer Causes Control*. 2015;26:1205–31.
- Jayasekara H, MacInnis RJ, Room R, English DR. Long-Term Alcohol Consumption and Breast, Upper Aero-Digestive Tract and Colorectal Cancer Risk: A Systematic Review and Meta-Analysis. *Alcohol Alcohol*. 2016;51:315–30.
- Chen JY, Zhu HC, Guo Q, Shu Z, Bao XH, Sun F, Qin Q, Yang X, Zhang C, Cheng HY, et al. Dose-Dependent Associations between Wine Drinking and Breast Cancer Risk - Meta-Analysis Findings. *Asian Pac J Cancer Prev*. 2016; 17:1221–33.
- Baan R, Straif K, Grosse Y, Secretan B, El Ghissassi F, Bouvard V, Altieri A, Coglianò V. Carcinogenicity of alcoholic beverages. *Lancet Oncol*. 2007;8:292–3.
- IARC. Alcohol consumption and ethyl carbamate. *IARC Monogr Eval Carcinog Risks Hum*. 96:3-1383.
- Klyosov AA, Rashkovetsky LG, Tahir MK, Keung WM. Possible role of liver cytosolic and mitochondrial aldehyde dehydrogenases in acetaldehyde metabolism. *Biochemistry*. 1996;35:4445–56.
- Zakhari S. Overview: how is alcohol metabolized by the body? *Alcohol Res Health*. 2006;29:245–54.
- Yoshida A, Huang IY, Ikawa M. Molecular abnormality of an inactive aldehyde dehydrogenase variant commonly found in Orientals. *Proc Natl Acad Sci U S A*. 1984;81:258–61.
- Lai CL, Yao CT, Chau GY, Yang LF, Kuo TY, Chiang CP, Yin SJ. Dominance of the inactive Asian variant over activity and protein contents of mitochondrial aldehyde dehydrogenase 2 in human liver. *Alcohol Clin Exp Res*. 2014;38:44–50.
- Chen YC, Peng GS, Tsao TP, Wang MF, Lu RB, Yin SJ. Pharmacokinetic and pharmacodynamic basis for overcoming acetaldehyde-induced adverse reaction in Asian alcoholics, heterozygous for the variant *ALDH2\*2* gene allele. *Pharmacogenet Genomics*. 2009;19:588–99.
- Harada S, Agarwal DP, Goedde HW. Aldehyde dehydrogenase deficiency as cause of facial flushing reaction to alcohol in Japanese. *Lancet*. 1981;2:982.
- Li H, Borinskaya S, Yoshimura K, Kalina N, Marusin A, Stepanov VA, Qin Z, Khaliq S, Lee MY, Yang Y, et al. Refined geographic distribution of the oriental *ALDH2\*504Lys* (nee 487Lys) variant. *Ann Hum Genet*. 2009;73:335–45.
- Luo HR, Wu GS, Pakstis AJ, Tong L, Oota H, Kidd KK, Zhang YP. Origin and dispersal of atypical aldehyde dehydrogenase *ALDH2487Lys*. *Gene*. 2009; 435:96–103.
- Tan EC, Lim L, Leong JY, Lim JY, Lee A, Yang J, Tan CH, Winslow M. Alcohol and aldehyde dehydrogenase polymorphisms in Chinese and Indian populations. *Subst Use Misuse*. 2010;45:1–14.
- Brooks PJ, Enoch MA, Goldman D, Li TK, Yokoyama A. The alcohol flushing response: an unrecognized risk factor for esophageal cancer from alcohol consumption. *PLoS Med*. 2009;6:e50.
- Boccia S, Hashibe M, Galli P, De Feo E, Asakage T, Hashimoto T, Hiraki A, Katoh T, Nomura T, Yokoyama A, et al. Aldehyde dehydrogenase 2 and head and neck cancer: a meta-analysis implementing a Mendelian randomization approach. *Cancer Epidemiol Biomarkers Prev*. 2009;18:248–54.
- Tsai ST, Wong TY, Ou CY, Fang SY, Chen KC, Hsiao JR, Huang CC, Lee WT, Lo HI, Huang JS, et al. The interplay between alcohol consumption, oral



- hygiene, ALDH2 and ADH1B in the risk of head and neck cancer. *Int J Cancer*. 2014;135:2424–36.
28. Chung CS, Lee YC, Liou JM, Wang CP, Ko JY, Lee JM, Wu MS, Wang HP. Tag single nucleotide polymorphisms of alcohol-metabolizing enzymes modify the risk of upper aerodigestive tract cancers: HapMap database analysis. *Dis Esophagus*. 2014;27:493–503.
  29. Oze I, Matsuo K, Hosono S, Ito H, Kawase T, Watanabe M, Suzuki T, Hatooka S, Yatabe Y, Hasegawa Y, et al. Comparison between self-reported facial flushing after alcohol consumption and ALDH2 Glu504Lys polymorphism for risk of upper aerodigestive tract cancer in a Japanese population. *Cancer Sci*. 2010;101:1875–80.
  30. Ji YB, Tae K, Ahn TH, Lee SH, Kim KR, Park CW, Park BL, Shin HD. ADH1B and ALDH2 polymorphisms and their associations with increased risk of squamous cell carcinoma of the head and neck in the Korean population. *Oral Oncol*. 2011;47:583–7.
  31. Zhao T, Wang C, Shen L, Gu D, Xu Z, Zhang X, Xu Y, Chen J. Clinical significance of ALDH2 rs671 polymorphism in esophageal cancer: evidence from 31 case-control studies. *Oncol Targets Ther*. 2015;8:649–59.
  32. Yokoyama A, Muramatsu T, Ohmori T, Yokoyama T, Okuyama K, Takahashi H, Hasegawa Y, Higuchi S, Maruyama K, Shirakura K, et al. Alcohol-related cancers and aldehyde dehydrogenase-2 in Japanese alcoholics. *Carcinogenesis*. 1998;19:1383–7.
  33. Koide T, Ohno T, Huang XE, Iijima Y, Sugihara K, Mizokami M, Xiang J, Tokudome S. HBV/HCV Infection, Alcohol, Tobacco and Genetic Polymorphisms for Hepatocellular Carcinoma in Nagoya, Japan. *Asian Pac J Cancer Prev*. 2000;1:237–43.
  34. Takeshita T, Yang X, Inoue Y, Sato S, Morimoto K. Relationship between alcohol drinking, ADH2 and ALDH2 genotypes, and risk for hepatocellular carcinoma in Japanese. *Cancer Lett*. 2000;149:69–76.
  35. Yu SZ, Huang XE, Koide T, Cheng G, Chen GC, Harada K, Ueno Y, Sueoka E, Oda H, Tashiro F, et al. Hepatitis B and C viruses infection, lifestyle and genetic polymorphisms as risk factors for hepatocellular carcinoma in Haimen, China. *Jpn J Cancer Res*. 2002;93:1287–92.
  36. Sakamoto T, Hara M, Higaki Y, Ichiba M, Horita M, Mizuta T, Eguchi Y, Yasutake T, Ozaki I, Yamamoto K, et al. Influence of alcohol consumption and gene polymorphisms of ADH2 and ALDH2 on hepatocellular carcinoma in a Japanese population. *Int J Cancer*. 2006;118:1501–7.
  37. Ding J, Li S, Wu J, Gao C, Zhou J, Cao H, Su PS, Liu Y, Zhou X, Chang J. Alcohol dehydrogenase-2 and aldehyde dehydrogenase-2 genotypes, alcohol drinking and the risk of primary hepatocellular carcinoma in a Chinese population. *Asian Pac J Cancer Prev*. 2008;9:31–5.
  38. Chang SC, Chang PY, Butler B, Goldstein BY, Mu L, Cai L, You NC, Baecker A, Yu SZ, Heber D, et al. Single nucleotide polymorphisms of one-carbon metabolism and cancers of the esophagus, stomach, and liver in a Chinese population. *PLoS One*. 2014;9:e109235.
  39. Liu J, Yang HJ, Lee MH, Jen CL, Hu HH, Lu SN, Wang LY, You SL, Huang YT, Chen CJ. Alcohol Drinking Mediates the Association between Polymorphisms of ADH1B and ALDH2 and Hepatitis B-Related Hepatocellular Carcinoma. *Cancer Epidemiol Biomarkers Prev*. 2016;25:693–9.
  40. Munaka M, Kohshi K, Kawamoto T, Takasawa S, Nagata N, Itoh H, Oda S, Katoh T. Genetic polymorphisms of tobacco- and alcohol-related metabolizing enzymes and the risk of hepatocellular carcinoma. *J Cancer Res Clin Oncol*. 2003;129:355–60.
  41. Tomoda T, Nouse K, Sakai A, Ouchida M, Kobayashi S, Miyahara K, Onishi H, Nakamura S, Yamamoto K, Shimizu K. Genetic risk of hepatocellular carcinoma in patients with hepatitis C virus: a case control study. *J Gastroenterol Hepatol*. 2011;27:797–804.
  42. Sangrajang S, Sato Y, Sakamoto H, Ohnami S, Khuhaprema T, Yoshida T. Genetic polymorphisms in folate and alcohol metabolism and breast cancer risk: a case-control study in Thai women. *Breast Cancer Res Treat*. 2010;123:885–93.
  43. Kawase T, Matsuo K, Hiraki A, Suzuki T, Watanabe M, Iwata H, Tanaka H, Tajima K. Interaction of the effects of alcohol drinking and polymorphisms in alcohol-metabolizing enzymes on the risk of female breast cancer in Japan. *J Epidemiol*. 2009;19:244–50.
  44. Choi JY, Abel J, Neuhaus T, Ko Y, Harth V, Hamajima N, Tajima K, Yoo KY, Park SK, Noh DY, et al. Role of alcohol and genetic polymorphisms of CYP2E1 and ALDH2 in breast cancer development. *Pharmacogenetics*. 2003;13:67–72.
  45. Murata M, Tagawa M, Watanabe S, Kimura H, Takeshita T, Morimoto K. Genotype difference of aldehyde dehydrogenase 2 gene in alcohol drinkers influences the incidence of Japanese colorectal cancer patients. *Jpn J Cancer Res*. 1999;90:711–9.
  46. Matsuo K, Hamajima N, Hirai T, Kato T, Koike K, Inoue M, Takezaki T, Tajima K. Aldehyde dehydrogenase 2 (ALDH2) genotype affects rectal cancer susceptibility due to alcohol consumption. *J Epidemiol*. 2002;12:70–6.
  47. Miyasaka K, Hosoya H, Tanaka Y, Uegaki S, Kino K, Shimokata H, Kawanami T, Funakoshi A. Association of aldehyde dehydrogenase 2 gene polymorphism with pancreatic cancer but not colon cancer. *Geriatr Gerontol Int*. 2010;10 Suppl 1:S120–6.
  48. Higuchi S, Matsushita S, Murayama M, Takagi S, Hayashida M. Alcohol and aldehyde dehydrogenase polymorphisms and the risk for alcoholism. *Am J Psychiatry*. 1995;152:1219–21.
  49. Harada S, Agarwal DP, Goedde HW, Tagaki S, Ishikawa B. Possible protective role against alcoholism for aldehyde dehydrogenase isozyme deficiency in Japan. *Lancet*. 1982;2:827.
  50. Higuchi S, Matsushita S, Imazeki H, Kinoshita T, Takagi S, Kono H. Aldehyde dehydrogenase genotypes in Japanese alcoholics. *Lancet*. 1994;343:741–2.
  51. Yokoyama A, Yokoyama T, Matsui T, Mizukami T, Kimura M, Matsushita S, Higuchi S, Maruyama K. Trends in gastrectomy and ADH1B and ALDH2 genotypes in Japanese alcoholic men and their gene-gastrectomy, gene-gene and gene-age interactions for risk of alcoholism. *Alcohol Alcohol*. 2013;48:146–52.
  52. Yokoyama A, Kato H, Yokoyama T, Tsujinaka T, Muto M, Omori T, Haneda T, Kumagai Y, Igaki H, Yokoyama M, et al. Genetic polymorphisms of alcohol and aldehyde dehydrogenases and glutathione S-transferase M1 and drinking, smoking, and diet in Japanese men with esophageal squamous cell carcinoma. *Carcinogenesis*. 2002;23:1851–9.
  53. Chen CC, Lu RB, Chen YC, Wang MF, Chang YC, Li TK, Yin SJ. Interaction between the functional polymorphisms of the alcohol-metabolism genes in protection against alcoholism. *Am J Hum Genet*. 1999;65:795–807.
  54. International Agency for Research on Cancer. Globocan 2012. <http://globocan.iarc.fr/>. Accessed on 18 Aug 2016.
  55. Yokoyama T, Yokoyama A, Kumagai Y, Omori T, Kato H, Igaki H, Tsujinaka T, Muto M, Yokoyama M, Watanabe H. Health risk appraisal models for mass screening of esophageal cancer in Japanese men. *Cancer Epidemiol Biomarkers Prev*. 2008;17:2846–54.
  56. Yokoyama A, Oda J, Iriguchi Y, Kumagai Y, Okamura Y, Matsuoka M, Mizukami T, Yokoyama T. A health-risk appraisal model and endoscopic mass screening for esophageal cancer in Japanese men. *Dis Esophagus*. 2013;26:148–53.
  57. Koyanagi YN, Ito H, Oze I, Hosono S, Tanaka H, Abe T, Shimizu Y, Hasegawa Y, Matsuo K. Development of a prediction model and estimation of cumulative risk for upper aerodigestive tract cancer on the basis of the aldehyde dehydrogenase 2 genotype and alcohol consumption in a Japanese population. *Eur J Cancer Prev*. 2017;26:38–47.
  58. Hendershot CS, Otto JM, Collins SE, Liang T, Wall TL. Evaluation of a brief web-based genetic feedback intervention for reducing alcohol-related health risks associated with ALDH2. *Ann Behav Med*. 2010;40:77–88.
  59. Austin J. The effect of genetic test-based risk information on behavioral outcomes: A critical examination of failed trials and a call to action. *Am J Med Genet A*. 2015;167A:2913–5.
  60. Smerecnik C, Grispin JE, Quaaq M. Effectiveness of testing for genetic susceptibility to smoking-related diseases on smoking cessation outcomes: a systematic review and meta-analysis. *Tob Control*. 2012;21:347–54.
  61. Kawakita D, Oze I, Hosono S, Ito H, Watanabe M, Yatabe Y, Hasegawa Y, Murakami S, Tanaka H, Matsuo K. Prognostic Value of Drinking Status and Aldehyde Dehydrogenase 2 Polymorphism in Patients With Head and Neck Squamous Cell Carcinoma. *J Epidemiol*. 2016;26:292–9.
  62. Yokoyama A, Omori T, Yokoyama T, Sato Y, Kawakubo H, Maruyama K. Risk of metachronous squamous cell carcinoma in the upper aerodigestive tract of Japanese alcoholic men with esophageal squamous cell carcinoma: a long-term endoscopic follow-up study. *Cancer Sci*. 2008;99:1164–71.
  63. US. Department of Health and Human Services and US. Department of Agriculture. 2015–2020 Dietary Guidelines for Americans. 8th Edition. December 2015. Available at <http://health.gov/dietaryguidelines/2015/guidelines/>.
  64. Kalinowski A, Humphreys K. Governmental standard drink definitions and low-risk alcohol consumption guidelines in 37 countries. *Addiction*. 2016;111:1293–8.
  65. Lee CH, Lee JM, Wu DC, Goan YG, Chou SH, Wu IC, Kao EL, Chan TF, Huang MC, Chen PS, et al. Carcinogenic impact of ADH1B and ALDH2 genes on squamous cell carcinoma risk of the esophagus with regard to the consumption of alcohol, tobacco and betel quid. *Int J Cancer*. 2008;122:1347–56.

66. Saad I, Wallach JM. Ethanal assay, using an enzyme-conductimetric method. *Anal Lett.* 1992;25:37–48.
67. Uebelacker M, Lachenmeier DW. Quantitative determination of acetaldehyde in foods using automated digestion with simulated gastric fluid followed by headspace gas chromatography. *J Autom Methods Manag Chem.* 2011;2011:907317.
68. Rodrigues MC, Guarieiro LL, Cardoso MP, Carvalho LS, da Rocha GO, de Andrade JB. Acetaldehyde and formaldehyde concentrations from sites impacted by heavy-duty diesel vehicles and their correlation with the fuel composition: Diesel and diesel/biodiesel blends. *Fuel.* 2012;92:258–63.
69. Chen CH, Budas GR, Churchill EN, Disatnik MH, Hurley TD, Mochly-Rosen D. Activation of aldehyde dehydrogenase-2 reduces ischemic damage to the heart. *Science.* 2008;321:1493–5.
70. Chen CH, Sun L, Mochly-Rosen D. Mitochondrial aldehyde dehydrogenase and cardiac diseases. *Cardiovasc Res.* 2010;88:51–7.
71. Chen CH, Ferreira JC, Gross ER, Mochly-Rosen D. Targeting aldehyde dehydrogenase 2: new therapeutic opportunities. *Physiol Rev.* 2014;94:1–34.
72. Zhang R, Wang J, Xue M, Xu F, Chen Y. ALDH2—the genetic polymorphism and enzymatic activity regulation: their epidemiologic and clinical implications. *Curr Drug Targets.* 2015. [Epub ahead of print].
73. Perez-Miller S, Younus H, Vanam R, Chen CH, Mochly-Rosen D, Hurley TD. Alda-1 is an agonist and chemical chaperone for the common human aldehyde dehydrogenase 2 variant. *Nat Struct Mol Biol.* 2010;17:159–64.
74. Gross ER, Zambelli VO, Small BA, Ferreira JC, Chen CH, Mochly-Rosen D. A personalized medicine approach for Asian Americans with the aldehyde dehydrogenase 2\*2 variant. *Annu Rev Pharmacol Toxicol.* 2015;55:107–27.
75. Banh A, Xiao N, Cao H, Chen CH, Kuo P, Krakow T, Bavan B, Khong B, Yao M, Ha C, et al. A novel aldehyde dehydrogenase-3 activator leads to adult salivary stem cell enrichment in vivo. *Clin Cancer Res.* 2011;17:7265–72.
76. Chen CH, Cruz LA, Mochly-Rosen D. Pharmacological recruitment of aldehyde dehydrogenase 3A1 (ALDH3A1) to assist ALDH2 in acetaldehyde and ethanol metabolism in vivo. *Proc Natl Acad Sci USA.* 2015;112:3074–9.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

