

Algebraic structure of the families of compatible frames of discernment

Fabio Cuzzolin

Computer Science Department, University of California at Los Angeles, 3811A Boelter Hall,
Los Angeles, CA 90095-1596, USA
E-mail: cuzzolin@cs.ucla.edu

One of the major ideas of Shafer's mathematical theory of evidence is the introduction of uncertainty descriptions on different representation domains of phenomena, called *families of compatible frames of discernment*. Here we are going to analyze these families of frames from an algebraic point of view, study the properties of *minimal refinements* of collections of domains and introduce the internal operation of *maximal coarsening* to establish the structure of *semimodular lattice*. Motivated by the search for a solution of the *conflict problem* that arises in sensor fusion applications, we will show the connection between classical *independence of frames* as Boolean subalgebras and independence of frames as elements of a locally finite Birkhoff lattice. This will eventually suggest a potential algebraic solution of the conflict problem.

Keywords: family of compatible frames, minimal refinement, Birkhoff lattices, linear independence, conflict problem

AMS subject classification (2000): 60D05, 28E10, 06C10, 68T37

1. Introduction

The *theory of evidence* [21] was introduced in the late 1970s by Glenn Shafer as a way of representing epistemic knowledge, starting from a sequence of seminal works [7–9] of Arthur Dempster. In this formalism the best representation of one's subjective belief is a *belief function* (b.f.) rather than a Bayesian mass distribution. Belief functions assign probability values to *sets* of possibilities rather than single events: their appeal rests on the fact they naturally encode evidence in favor of propositions. The theory provides a simple method for combining the evidence carried by a number of different sources (*Dempster's rule*) with no need of any *a-priori* distributions.

The other major pillar of the theory (strictly related to the nature of belief functions) is the formalization of the idea of structured collection of representations of the external world, encoded into the notion of *family of compatible frames*. As our knowledge is inherently imprecise, it cannot be exhausted by any representation on single domains. This implies, for instance, that representing ignorance in a Bayesian setup is impossible, since any non-informative distribution is not invariant when

passing from a domain to another one [21]. On the other side, acceptable belief functions admit sister functions on any other frame of a family of compatible frames, defining the class of *support* functions.

In sensor fusion applications to engineering or computer vision [5] we often face situations in which several belief functions defined on different frames of a family need to be combined to make decisions or update the knowledge state of a system. Unfortunately, their combination is not always possible: as we prove in this paper, this is true only when their domains are *independent* in a very precise way (Theorem 1). We refer to this question with the term *conflict problem*. A possible naive solution consists then in building a conflict graph whose nodes are the belief functions to combine, and recursively detecting the most coherent set of functions (see section 3.1). However, this is a rather heuristic solution which suffers a high computational cost, and no clear criteria for choosing a collection of b.f. instead of another are provided.

In this paper we try and pose the conflict problem in a rigorous mathematical context, by studying the algebraic structure of the families of compatible frames as objects obeying a small number of axioms [21]. The intuition comes from a striking resemblance between the notion of independence of frames (see Definition 9)

$$\omega_1(A_1) \cap \cdots \cap \omega_n(A_n) \neq \emptyset, \quad \forall A_i \subset \Theta_i$$

and the familiar linear independence relation for a collection of subspaces $\{V_i\}$ of a vector space V

$$v_1 + \cdots + v_n \neq 0, \quad \forall v_i \in V_i.$$

We may wonder whether this is a simple analogy, or rather the symptom of a deeper similarity of these structures at the algebraic level. Given a collection of arbitrary elements of a vector space, a well known procedure called *Gram–Schmidt algorithm* is able to generate a new collection of independent vectors, spanning the same subspace. It would be highly desirable to design an analogous procedure which, given a set of belief functions defined on arbitrary elements of a family of frames, would yield another collection of (combinable) belief functions defined on independent frames, generating an equivalent (in some formal sense) Dempster’s combination.

In this work we prepare the mathematical ground for this ambitious goal, by studying the monoidal and lattice structures of the families of compatible frames of discernment. As a matter of fact, it turns out that both vector subspaces and families of frames share the structure of *Birkhoff lattice* [25]. Birkhoff lattices admit the introduction of a linear dependence relation on their atoms. We can then reasonably conjecture (as we will see in section 7) the equivalence between the classical notion of independence of frames as Boolean subalgebras, and their independence as elements of a locally finite Birkhoff lattice. A formal proof of this equivalence would eventually prefigure a solution to the conflict problem based on a ‘pseudo Gram-Schmidt’ algorithm, transforming belief functions on arbitrary frames into ‘equivalent’ b.f. on independent frames.

1.1. Paper outline

After briefly recalling the basic notions of the theory of evidence (section 2), we give a characterization of the relationship between conflict problem and independence of frames (section 3). As Dempster's combination is guaranteed *if and only if the underlying frames are independent* (Theorem 1), it is crucial to study independence in more detail to provide practical solutions to real-world applications (section 3.1), in which different measurements are fused to make decisions or compute estimates.

This can be done in a rigorous fashion by posing the problem in an algebraic context (section 4), after finding an alternative, *constructive* set of axioms for families of compatible frames. If we assume we have a finite knowledge of the problem (a realistic assumption in practical applications) we can build the subfamily of frames of discernment generated by any given frame.

In section 5 we first focus on these finite subfamilies of frames, showing that the natural operator of *minimal refinement* \otimes induces on them a structure of commutative monoid with annihilator (Theorem 4). The collection of refinings of a finite subfamily of frames is also a monoid, as one can build an isomorphism between frames and refinings (equation (8)). More important, both frames (\mathcal{F}, \otimes) and refinings (\mathcal{R}, \otimes) of a *general family* of compatible frames of discernment also possess the algebraic structure of a commutative monoid (section 5.4), with finite subfamilies as submonoids. The situation is summarized in equation (13).

As the internal operation of a monoid induces an order relation, we are tempted to think about the problem from a lattice-theoretic point of view. As a matter of fact, in section 6 we prove that the collection of sets of a family of compatible frames is a *Birkhoff lattice* (with minimal refinement as least upper bound or *sup*, and the dual operation of *maximal coarsening* as greatest lower bound or *inf*) of locally finite length and with a smallest element. Formally, $(\mathcal{F}, \otimes, \oplus)$ is a *Birkhoff lattice bounded below*.

This is where the interplay between the conflict problem and the algebraic structure of the families of frames takes place, as Birkhoff lattices come with a notion of independence. It is then quite natural to conjecture (section 7) that the independence of frames as Boolean subalgebras (Definition 9), originally incorporated into the theory of evidence, could possibly be equivalent to the independence of frames as elements of a Birkhoff lattice (Theorem 14). This would open the way to an algebraic solution of the conflict problem, where a pseudo Gram-Schmidt orthogonalization procedure would be used to transform any given collection of b.f. into a new, combinable set by projection onto independent frames.

1.2. Previous work

Not much work has been done on the properties of the families of compatible frames. In [22], though, an analysis of the collections of partitions of a given frame in the context of the hierarchical representation of belief can be found. A wider ex-

position of the algebraic properties of the families of frames is instead presented in [14], where Chapter 7 is devoted to their lattice-theoretical interpretation and the meaning of the concept of independence, while Chapter 8 develops the consequences of the application of constraints to the structure of the family of compatible frames and deepens the properties of *Markov trees* as a useful characterization.

Some work has been done on the conflict problem, too [10]. Murphy [16], on his side, studied a related problem (the failure to balance multiple evidence), illustrated the proposed solutions and described their limitations.

2. The theory of evidence

In the theory of evidence [21], subjective degrees of belief are mathematically represented as *belief functions* (b.f.), which generalize finite probability distributions in the finite case. Belief functions encoding different pieces of evidence can be combined by means of *Dempster's rule* [7]. The concept of *family of compatible frames* is introduced to formalize the intuitive idea of different levels of description of a same phenomenon.

2.1. Belief functions

2.1.1. Axiomatic definition

The reader will of course be familiar with the classical definition of probability measure, due to Kolmogorov.

Definition 1. A *probability measure* over a σ -field $\mathbf{F} \subset 2^\Theta$ associated with a sample space Θ is a function $p : \mathbf{F} \rightarrow [0,1]$ such that

1. $p(\emptyset) = 0$;
2. $p(\Theta) = 1$;
3. if $A \cap B = \emptyset$, $A, B \in \mathbf{F}$ then $p(A \cup B) = p(A) + p(B)$ (additivity).

Now, let us restrict ourselves to finite sets, and relax the third constraint allowing the function to have the value obtained by additivity as a lower bound. We then get a new mathematical object: a *belief function*.

Definition 2. Suppose Θ is a finite set, and let 2^Θ denote the set of all the subsets of Θ . A *belief function* on Θ is a function $s : 2^\Theta \rightarrow [0,1]$ such that

1. $s(\emptyset) = 0$;
2. $s(\Theta) = 1$;

3. for every positive integer n and every collection $A_1, \dots, A_n \in 2^\Theta$

$$s(A_1 \cup \dots \cup A_n) \geq \sum_i s(A_i) - \sum_{i < j} s(A_i \cap A_j) + \dots + (-1)^{n+1} s(A_1 \cap \dots \cap A_n).$$

The third axiom is called *superadditivity*, and obviously reduces to standard additivity when we replace the inequality with an equality. Belief functions can then be seen as *generalizations* of the familiar probability distributions on finite domains. Θ can be interpreted as the set of possible answers of a decision problem, exactly one of which is the correct one. For each subset (proposition) $A \subset \Theta$ the quantity $s(A)$ measures the *degree of belief* that the true answer lies in A .

2.1.2. Basic probability assignment

Belief functions can be given an alternative but equivalent description in terms of total belief committed to a set of possibilities A . Let us call the finite set of possible outcomes for a decision problem *frame*¹ of discernment.

Definition 3. A *basic probability assignment (b.p.a.)* over a frame Θ is a function $m : 2^\Theta \rightarrow [0, 1]$ such that

$$m(\emptyset) = 0, \sum_{A \subset \Theta} m(A) = 1, m(A) \geq 0 \forall A \subset \Theta.$$

Subsets of Θ associated with non-zero values of m are called *focal elements* and their union \mathcal{C} *core*. Now suppose a b.p.a. is introduced on an arbitrary frame.

Definition 4. The belief function s associated with a basic probability assignment m is defined as (see Figure 1a):

$$s(A) = \sum_{B \subset A} m(B).$$

Conversely, a basic probability assignment m associated with a given belief function s can be uniquely recovered by means of the *Moebius inversion formula*²

$$m(A) = \sum_{B \subset A} (-1)^{|A-B|} s(B) \tag{1}$$

so that there is a 1–1 correspondence $m \leftrightarrow s$ between these two set functions. It can be proved that Definitions 4 and 2 are equivalent.

¹For a note about the intuitionistic origin of this denomination see Rosenthal, *Quantales and their applications* [20].

²See [25] for an explanation in terms of the theory of monotone functions over partially ordered sets.

2.2. Dempster's rule of combination

Belief functions representing distinct bodies of evidence can be combined by means of the so-called *Dempster's rule of combination* [7, 9].

Definition 5. The *orthogonal sum* of two belief functions s_1, s_2 is a new belief function $s_1 \oplus s_2$ whose focal elements are all the possible non-empty intersections $A_i \cap B_j$ between focal elements A_i, B_j of s_1, s_2 , respectively, and whose b.p.a. is given by

$$m(A) = \frac{\sum_{i,j:A_i \cap B_j = A} m_1(A_i) \cdot m_2(B_j)}{1 - \sum_{i,j:A_i \cap B_j = \emptyset} m_1(A_i) \cdot m_2(B_j)}, A \neq \emptyset. \tag{2}$$

The normalization constant in equation (2) measures the *weight of conflict* $Con(s_1, s_2)$ between the belief functions, and represents the amount of probability they attribute to contradictory (i.e., disjoint) subsets. If all the intersections between focal elements of s_1 and s_2 are empty ($\mathcal{C}_1 \cap \mathcal{C}_2 = \emptyset$) the two belief functions are said *non-combinable*.

The above definition can be easily extended to the combination of several belief functions (figure 1b).

2.2.1. Example

Consider as an example a gray-scale image, and imagine you need to measure the column index of the center of mass of the intensity function of the image. A naive




Figure 1. (a) Basic probability assignment and belief value. In the picture $m(B) \neq 0, B \subset A$ only for B_1 and B_2 so that $s(A) = m(B_1) + m(B_2)$. (b) Dempster's rule of combination: on the y, x axes are depicted the focal elements A_i and B_j of s_1, s_2 , respectively, where the width of columns and rows is proportional to the b.p.a. associated with each focal element (in a size 1 square). Each intersection is given the product of the corresponding masses (equal to the area of the related square). Contributions of coincident intersections are then summed. This value is finally normalized by taking into account all the empty intersections (denominator of equation (2)).




Figure 2. The frame (*middle*) associated with a partition of the column range of an image (*left*). The b.f. s_1 with focal elements A_1, A_2 and s_2 with f.e. B_1, B_2 (*middle*) can be combined through Dempster's rule yielding a new belief function $s_1 \oplus s_2$ (*right*) with f.e. X_1 and X_2 .

quantization of the column range in, say, five intervals (figure 2-left) yields a frame of discernment $\{c_1, \dots, c_5\}$ (figure 2-middle). On this frame we can define a belief function s_1 with two focal elements $A_1 = \{c_2\}$ and $A_2 = \{c_2, c_4\}$ through the basic probability assignment $m_1(A_1) = 0.7, m_1(A_2) = 0.3$. For instance, the belief values of the sets $\{c_4\}, \{c_2, c_5\}, \{c_2, c_3, c_4\}$ are then $s_1(\{c_4\}) = m_1(\{c_4\}) = 0, s_1(\{c_2, c_5\}) = m_1(\{c_2\}) + m_1(\{c_5\}) + m_1(\{c_2, c_5\}) = 0.7 + 0 + 0 = 0.7, s_1(\{c_2, c_3, c_4\}) = m_1(\{c_2\}) + m_1(\{c_2, c_4\}) = 0.7 + 0.3 = 1$, respectively. Now introduce another b.f. s_2 with b.p.a. $m_2(B_1) = m_2(\{c_2, c_3\}) = 0.6$ and $m_2(B_2) = m_2(\{c_4, c_5\}) = 0.4$. Their Dempster's combination yields a new b.f. with focal elements (figure 2-right) $X_1 = A_1 \cap B_1 = A_2 \cap B_1 = \{c_2\}$ and $X_2 = A_2 \cap B_2 = \{c_4\}$ and b.p.a. $m(X_1) = \frac{0.7 \cdot 0.6 + 0.3 \cdot 0.6}{1 - 0.7 \cdot 0.4} = 5/6, m(X_2) = \frac{0.3 \cdot 0.4}{1 - 0.7 \cdot 0.4} = 1/6$.

2.3. Random sets and belief functions

2.3.1. Multi-valued mappings

In fact, in the original approach due to Dempster belief functions are inherently associated with multi-valued maps between frames. Let us suppose we have probabilities (coming from arbitrary sources, for instance subjective judgement or objective measurements) for a question Q_1 and we want to derive a degree of belief for a related question Q_2 . For example, Q_1 could be the judgement on the reliability of a witness, and Q_2 the decision about the truth of the reported fact. Let us call X and Θ the sets (frames) of possible answers of Q_1 and Q_2 , respectively. Given a probability $P(x)$ on X we wish to derive a degree of belief $s(A)$ that $A \subset \Theta$ contains the correct answer to Q_2 .

If we call $\Gamma(x)$ the subset of responses to Q_2 compatible with x , each element $x \in X$ tells us that the answer to Q_2 is somewhere in A , whenever $\Gamma(x) \subset A$. The degree of belief $s(A)$ is then the total probability of all answers x that satisfy the above condition, namely

$$s(A) = P\{x | \Gamma(x) \subset A\}.$$

The map Γ is called a *multi-valued mapping* from X to Θ . This mapping, together with the probability measure on X , induces a belief function on Θ : $(P, \Gamma) \mapsto s$.

2.3.2. Belief functions as random sets

Having a multi-valued mapping Γ , a straightforward step is to consider the probability $P(x)$ attached to the subset $\Gamma(x) \subset \Theta$: what we obtain is a *random set* in Θ , i.e., a probability measure on a collection of subsets (see [12, 15] for the most complete introductions to the matter). The degree of belief $s(A)$ then becomes the total probability that the random set is contained in A .

This approach has been emphasized in particular by Nguyen [11, 17, 18] and resumed in [22].

Consider a multi-valued mapping $\Gamma : X \rightarrow 2^\Theta$. We can define the *lower inverse* of Γ as

$$\begin{aligned} \Gamma_* : 2^\Theta &\rightarrow 2^X \\ A &\mapsto \Gamma_*(A) \doteq \{x \in X : \Gamma(x) \subset A, \Gamma(x) \neq \emptyset\}. \end{aligned} \quad (3)$$

and in the dual way its *upper inverse* as

$$\begin{aligned} \Gamma^* : 2^\Theta &\rightarrow 2^X \\ A &\mapsto \Gamma^*(A) \doteq \{x \in X : \Gamma(x) \cap A \neq \emptyset\}. \end{aligned} \quad (4)$$

Given two σ -fields \mathcal{A}, \mathcal{B} on X, Θ respectively, Γ is said *strongly measurable* iff $\forall B \in \mathcal{B}, \Gamma^*(B) \in \mathcal{A}$. The *lower probability* measure on \mathcal{B} is defined as $P_*(B) \doteq P(\Gamma_*(B))$ for all $B \in \mathcal{B}$.

Nguyen proved that, if Γ is strongly measurable, the probability distribution \hat{P} of the random set [17] coincides with the lower probability measure

$$\hat{P}[I(B)] = P_*(B) \quad \forall B \in \mathcal{B}$$

and (in the finite case) the probability distribution of the random set Γ is precisely the basic probability assignment associated with the lower probability (belief function) P_* .

An analysis of the relations between the transferable belief model and the theory of random sets can be found in [24].

2.4. Refinings

As belief functions are inherently associated with maps between different (but related) domains, the theory of evidence quite naturally contemplates the impact of evidence on different finite descriptions of the same problem. In particular, a new amount of evidence can allow us to make decisions (by means of Dempster's rule) over more detailed domains. This argument is embodied into the notion of *refining*.

Definition 6. Given two frames Θ and Ω , a map $\omega : 2^\Theta \rightarrow 2^\Omega$ is a *refining* if it satisfies the following properties

1. $\omega(\{\theta\}) \neq \emptyset \quad \forall \theta \in \Theta$,
2. $\omega(\{\theta\}) \cap \omega(\{\theta'\}) = \emptyset$ when $\theta \neq \theta'$,
3. $\cup_{\theta \in \Theta} \omega(\{\theta\}) = \Omega$,

while its value for arbitrary subsets A of Θ is given by

$$\omega(A) \doteq \bigcup_{\theta \in A} \omega(\{\theta\}).$$

Ω is called a *refinement* of Θ , while Θ is said a *coarsening* of Ω . Both frames represent sets of possible outcomes for the same decision problem, but the finer one is a more detailed description, obtained by splitting each possible answer $\theta \in \Theta$ into a set $\omega(\{\theta\})$.

Consider for example a partition A_1, \dots, A_n of Ω . If we define $\Theta = \{A_1, \dots, A_n\}$ the map $\omega : 2^\Omega \rightarrow 2^\Theta$ with $\omega(A_i) = A_i$ is a refining.

2.4.1. Vacuous extension

Any belief function s defined on a frame Θ can be ‘projected’ onto any refinement Ω of Θ by *vacuous extension*.

Definition 7. Suppose Ω is a refinement of Θ , with refining $\omega : 2^\Omega \rightarrow 2^\Theta$. The *vacuous extension* on Ω of a belief function $s : 2^\Theta \rightarrow [0, 1]$ on Θ is the belief function $s' : 2^\Omega \rightarrow [0, 1]$ on Ω defined as

$$s'(A) = \max_{B \subset \Theta, \omega(B) \subset A} s(B) \quad \forall A \subset \Omega.$$

s' is a belief function on the frame Ω [21], whose focal elements are exactly the images of the focal elements of s ,

$$\{B_1, \dots, B_k\} \mapsto \{A_1 = \omega(B_1), \dots, A_k = \omega(B_k)\}$$

with $m_{s'}(A_i) = m_s(B_i)$.

Recall the example of section 2.2.1, and imagine to refine each column range c_1, \dots, c_5 into two subranges: $\omega(c_1) = \{c_{11}, c_{12}\}, \dots, \omega(c_5) = \{c_{51}, c_{52}\}$. The new frame $\Omega \doteq \{c_{11}, c_{12}, c_{21}, c_{22}, c_{31}, c_{32}, c_{41}, c_{42}, c_{51}, c_{52}\}$ is a refinement of $\Theta \doteq \{c_1, \dots, c_5\}$. The vacuous extension of s_2 defined above is then the belief function s'_2 on Ω with focal elements $A'_1 = \{c_{21}, c_{22}\}$ and $A'_2 = \{c_{21}, c_{22}, c_{41}, c_{42}\}$ and probability assignment $m_{s'_2}(A'_1) = 0.7, m_{s'_2}(A'_2) = 0.3$.

2.5. Families of compatible frames

Now that the basis tools (refining maps between frames) have been introduced, the intuitive idea of different descriptions of a same phenomenon can be formally expressed by the concept of *family of compatible frames* (see [21], pages 121–125).

Definition 8. A non-empty collection of finite non-empty sets \mathcal{F} is a *family of compatible frames of discernment* with refinings \mathcal{R} , where \mathcal{R} is a non-empty collection of

refinings between couples of frames in \mathcal{F} , if \mathcal{F} and \mathcal{R} meet the following requirements:

Axiom 1. *Composition of refinings:* if $\omega_1 : 2^{\Theta_1} \rightarrow 2^{\Theta_2}$ and $\omega_2 : 2^{\Theta_2} \rightarrow 2^{\Theta_3}$ are in \mathcal{R} , then $\omega_1 \circ \omega_2 \in \mathcal{R}$.

Axiom 2. *Identity of coarsenings:* if $\omega_1 : 2^{\Theta_1} \rightarrow 2^\Omega$ and $\omega_2 : 2^{\Theta_2} \rightarrow 2^\Omega$ are in \mathcal{R} and $\forall \theta_1 \in \Theta_1 \exists \theta_2 \in \Theta_2$ such that $\omega_1(\{\theta_1\}) = \omega_2(\{\theta_2\})$, then $\Theta_1 = \Theta_2$ and $\omega_1 = \omega_2$.

Axiom 3. *Identity of refinings:* if $\omega_1 : 2^\Theta \rightarrow 2^\Omega$ and $\omega_2 : 2^\Theta \rightarrow 2^\Omega$ are in \mathcal{R} , then $\omega_1 = \omega_2$.

Axiom 4. *Existence of coarsenings:* if $\Omega \in \mathcal{F}$ and A_1, \dots, A_n is a disjoint partition of Ω , then there is a coarsening Ω' of Ω in \mathcal{F} corresponding to this partition, i.e., $\forall A_i$ there exists an element of Ω' whose image under the appropriate refining is A_i .

Axiom 5. *Existence of refinings:* if $\theta \in \Theta \in \mathcal{F}$ and $n \in \mathbb{N}$ then there exists a refining $\omega : 2^\Theta \rightarrow 2^\Omega$ in \mathcal{R} and $\Omega \in \mathcal{F}$ such that $\omega(\{\theta\})$ has n elements.

Axiom 6. *Existence of common refinements:* every pair of elements in \mathcal{F} has a common refinement in \mathcal{F} .

By iterative application of Axiom 6 it follows that any finite collection of compatible frames has many common refinements. One of these is particularly simple.

Proposition 1. If $\Theta_1, \dots, \Theta_n$ are elements of a family of compatible frames \mathcal{F} then there exists a unique element $\Theta \in \mathcal{F}$ such that

1. $\forall i \exists \omega_i : 2^{\Theta_i} \rightarrow 2^\Theta$ refining;
2. $\forall \theta \in \Theta \exists \theta_i \in \Theta_i$ for $i = 1, \dots, n$ such that

$$\{\theta\} = \omega_1(\{\theta_1\}) \cap \dots \cap \omega_n(\{\theta_n\}). \quad (5)$$

This unique frame is called the *minimal refinement* $\Theta_1 \otimes \dots \otimes \Theta_n$ of the collection $\Theta_1, \dots, \Theta_n$ and is the simplest space in which you can compare propositions belonging to different compatible domains.

2.5.1. Example

In section 2.2.1 we introduced a frame Θ_1 by partitioning the column range into 5 intervals. The set of columns can be partitioned also into 10 intervals, yielding a new frame Θ_2 (remember section 2.4.1). More, the row range can also be divided in, say, 6 intervals (Θ_3). All those frames are clearly related to the location of the center

of mass: as figure 3 suggests, they all belong to a family of compatible frames, with the collection of cells $\Theta = \{e_1, \dots, e_{60}\}$ depicted in figure 3-left as common refinement. Figure 3-right shows the refinings between them where, for instance, $\omega_1(\{c_1\}) = \{c_{11}, c_{12}\}$, $\omega_2(\{c_{11}\}) = \{e_1, e_{11}, e_{21}, e_{31}, e_{41}, e_{51}\}$, $\omega_3(\{r_1\}) = \{e_1, \dots, e_{10}\}$, etcetera.

3. Independence and evidence conflict

Belief functions carrying distinct pieces of evidence over different frames of a family can be combined by means of Dempster’s rule. We just need to project each of them onto their common refinement by vacuous extension, and then combine them using Dempster’s rule. The information encoded by each belief function can then be used to evaluate propositions in this finer, common environment.

3.1. Motivations: Sensor fusion

A natural application of these attractive features of the theory of evidence can be found in *sensor fusion*, in which measurements coming from different sources have to be combined to make decisions, in order to increase the robustness of the process.

For instance, in some recent works [5, 6] we proposed an evidential framework for the solution of the *object tracking* problem. In this context, given a sequence of images of a moving articulated object you want to reconstruct the position and configuration of the body at each time instant. We represented a number of different *feature measurements* extracted from the images as belief functions and combined them to compute an estimate of the pose $\hat{q}(t) \in \tilde{Q}$, where \tilde{Q} is a sufficiently ‘good’ finite approximation of the parameter space Q of the object.

As an example, we can think of a human hand as an articulated object with 27 degrees of freedom (its position, orientation and the values of the angles between pairs of links of the fingers). As features we can choose indifferently the positions of




Figure 3. An example of family of compatible frames of discernment. Different discrete quantizations of row and column ranges of an image have a common refinement, the set of cells shown on the left. The refinings between all those frames are shown on the right.



Figure 4. The PantoMouse planar robot (*left*). Its pose is just the 2D position of its end-effector (*right*).

the fingertips in the image plane, the center of mass of the image itself, etc. In [5] we showed some experiments involving a simple planar robot (see figure 4).

These feature spaces Θ_i and the approximate parameter space \tilde{Q} form a finite subset of a family of compatible frames of discernment, where \tilde{Q} is a common refinement of the collection of feature spaces (see figure 5).

3.2. Independence and conflicting evidence

Unfortunately, the combination of evidence available on different domains is guaranteed *only for trivially interacting frames*.

Definition 9. Let $\Theta_1, \dots, \Theta_n$ be compatible frames, and $\omega_i : 2^{\Theta_i} \rightarrow 2^{\Theta_1 \otimes \dots \otimes \Theta_n}$ the corresponding refinings to their minimal refinement. They are said to be *independent*³ if

$$\omega_1(A_1) \cap \dots \cap \omega_n(A_n) \neq \emptyset \quad (6)$$

whenever $\emptyset \neq A_i \subset \Theta_i$ for $i = 1, \dots, n$.

Independence of frames and combinability of belief functions are as a matter of fact strictly connected.

Theorem 1. Let $\Theta_1, \dots, \Theta_n$ a set of compatible frames. The following conditions are equivalent:

1. all the possible collections of belief functions s_1, \dots, s_n defined respectively over $\Theta_1, \dots, \Theta_n$ are combinable over their minimal refinement $\Theta_1 \otimes \dots \otimes \Theta_n$;
2. $\Theta_1, \dots, \Theta_n$ are *independent*;

³ This definition is nothing but a formulation in the evidential language of the concept of independence of Boolean subalgebras of a Boolean algebra [23].




Figure 5. Families of feature spaces involved in the object tracking problem. The actual continuous parameter space Q of the articulated object is unknown, while its finite approximation \tilde{Q} is the common refinement where the evidence extracted from the image is combined.

- 3. there is an isomorphism $\Theta_1 \otimes \dots \otimes \Theta_n \leftrightarrow \Theta_1 \times \dots \times \Theta_n$;
- 4. $|\Theta_1 \otimes \dots \otimes \Theta_n| = \prod_{i=1}^n |\Theta_i|$.

Proof. $1 \Rightarrow 2$. We know that if s_1, \dots, s_n are combinable then s_i, s_j must be combinable $\forall i, j = 1, \dots, n$. Hence $\omega_i(C_i) \cap \omega_j(C_j) \neq \emptyset \forall i, j$ where C_i is the core of s_i , and $\omega_i : 2^{\Theta_i} \rightarrow 2^{\Theta_1 \otimes \dots \otimes \Theta_n}$ is the refining connecting Θ_i with the common refinement $\Theta_1 \otimes \dots \otimes \Theta_n$. Being s_i, s_j arbitrary, their cores C_i, C_j can be any pair of subsets of Θ_i, Θ_j , respectively, so that the condition can be rewritten as

$$\omega_i(A_i) \cap \omega_j(A_j) \neq \emptyset \quad \forall A_i \subset \Theta_i, A_j \subset \Theta_j.$$

$2 \Rightarrow 1$. Since $\Theta_1, \dots, \Theta_n$ are independent, $\omega_1(A_1) \cap \dots \cap \omega_n(A_n) \neq \emptyset$ for all non-empty $A_i \subset \Theta_i, \forall i = 1, \dots, n$. But then if we pick $A_i = C_i$ (the core of s_i) $\forall i = 1, \dots, n$ we get $\omega_1(C_1) \cap \dots \cap \omega_n(C_n) \neq \emptyset$, i.e., the vacuous extensions of s_1, \dots, s_n onto $\Theta_1 \otimes \dots \otimes \Theta_n$ are combinable (see section 2.2).

$2 \Rightarrow 3$. We note that

$$\bigcap_i \omega_i(\{\theta_i^k\}) = \bigcap_i \omega_i(\{\theta_i^l\}) \Leftrightarrow \theta_i^k = \theta_i^l \quad \forall i = 1, \dots, n.$$

In fact, if $\exists j$ s.t. $\theta_j^k \neq \theta_j^l$ then $\omega_j(\{\theta_j^k\}) \neq \omega_j(\{\theta_j^l\})$ for the properties of refinings: since they are disjoint, it must be $\bigcap_i \omega_i(\{\theta_i^k\}) \neq \bigcap_i \omega_i(\{\theta_i^l\})$. Hence their number

coincides with the number of the n -tuples of singletons $|\Theta_1| \times \cdots \times |\Theta_n|$, and since $\Theta_1 \otimes \cdots \otimes \Theta_n$ is a minimal refinement each of these subsets must correspond to a singleton.

3 \Rightarrow 2. For Proposition 1, each $\theta \in \Theta_1 \otimes \cdots \otimes \Theta_n$ corresponds to a subset of the form $\bigcap_i \omega_i(\{\theta_i^k\})$. Since the singletons are $|\Theta_1| \times \cdots \times |\Theta_n|$ the above subsets come in the same number, but then they all are non-empty, so that $\Theta_1, \dots, \Theta_n$ are independent.

3 \Rightarrow 4. Obvious.

4 \Rightarrow 3. $\Theta_1 \otimes \cdots \otimes \Theta_n = \{\bigcap_i \omega_i(\{\theta_i\}) : \theta_i \in \Theta_i\}$, so that if the intersections' number is $|\Theta_1| \times \cdots \times |\Theta_n|$ they all must be non-empty and each of them can be denoted by $(\theta_1, \dots, \theta_n)$. \square

For any collection of non-independent frames there exists at least a collection of belief functions defined on them that are not combinable. Any given set of belief functions is then characterized by a *weight of conflict* Con . When $Con = \infty$ there is no way of combining all of them by means of Dempster's rule.

3.3. The conflict problem

Going back to our sensor fusion application, since the belief functions built from the extracted features are not always compatible, we need a method for detecting the subset we are going to apply the rule of combination to. A basic property of the conflict weight is

$$Con(s_1, \dots, s_{n+1}) = Con(s_1, \dots, s_n) + Con(s_1 \oplus \dots \oplus s_n, s_{n+1}),$$

so that if $Con(s_i, s_j) = +\infty$ then $Con(s_i, s_j, s_k) = +\infty \forall s_k$. This suggests a bottom-up technique.

First the level of conflict is computed for each pair of measurement functions (s_i, s_j) , $i, j = 1, \dots, n$. Then a suitable threshold is chosen and a *conflict graph* is built, where every node represents a belief function while an edge indicates a conflict level below the threshold. Finally the subsets of combinable b.f. of size $d + 1$ are recursively computed from those of size d , eventually detecting the most coherent set of features. This approach obviously suffers a high computational cost when large groups of functions turn out to be compatible. Also, there is no clear indication about how to select a particular set of combinable belief functions.

Theorem 7 instead establishes a direct connection between conflicting evidence and independence of frames. As we have discussed in the introduction, this strongly suggests that a satisfactory, formal solution could be found in an algebraic context. In this perspective, it is necessary to understand the nature of the analogy between independence of vector spaces and independence of frames. This in turns involves a deep study of the algebraic properties of the families of compatible frames.

4. Axiom analysis

4.1. Constructive approach

To understand the algebraic properties of families of frames we necessarily need to analyze the axioms of Definition 8. As we have seen in section 3.1, in real-world applications our knowledge of a given problem is finite. We first then consider an arbitrary finite set S , and adopting a constructive approach we check the result of the application of the axioms of Definition 8.

Axiom A4 needs to be applied first, yielding the collection of all the possible partitions of S and the refinings between each of them and the basis set. By applying A4 again to the new sets all the refinings among them are achieved, while no other set is added to the collection. Axioms 2 and 3 guarantee the uniqueness of the obtained maps and sets.⁴ It is clear even at a first glance that A6 claims an existing condition but is not constructive, i.e., it does not allow to generate new sets from a given initial collection. Let us then introduce a new axiom

Axiom 7. *Existence of the minimal refinement:* for every pair of elements in \mathcal{F} there exists their minimal refinement in \mathcal{F} , i.e., a set satisfying the conditions of Proposition 1.

and consider another set of axioms by replacing A6 with A7. Let us call $A_{1..6}$ and $A_{1..5,7}$ these two formulations.

Theorem 2. $A_{1..6}$ and $A_{1..5,7}$ are equivalent formulations of the notion of family of compatible frames.

Proof. It is necessary and sufficient to prove that (i) A7 can be obtained by using the set of axioms $A1, \dots, A6$ and (ii) that A6 can be obtained from $A1, \dots, A5, A7$.

(i). See [21] or Proposition 1. (ii). Each refinement of a given pair of sets Θ_1, Θ_2 can be obtained by arbitrarily refining $\Theta_1 \otimes \Theta_2$ by means of Axiom A5. Anyway, the minimal refinement is obviously a refinement so that $A7 \Rightarrow A6$. \square

From now on we can then replace axiom A6 with axiom A7.

4.2. Finite knowledge

Again, if we assume our knowledge of the problem to be *finite and static*, Axiom A5 cannot be used. According to the established notation let us call $A_{1..4,7}$ the set of axioms corresponding to finite knowledge.

⁴ Observe that axiom A1 in this situation is redundant, for it does not add any new refinings.

Definition 10. The *subfamily generated by a collection of sets* $\Theta_1, \dots, \Theta_n$ by means of a set of axioms \mathcal{A} is the smallest collection of frames $\langle \Theta_1, \dots, \Theta_n \rangle_{\mathcal{A}}$ containing $\Theta_1, \dots, \Theta_n$ and closed under the application of the axioms in \mathcal{A} .

Given their importance in concrete applications it is natural to focus first on the properties of finite sub-families of compatible frames. We will then extend the analysis to general abstract families to get a comprehensive picture of the matter.

Lemma 1. The minimal refinement of two coarsenings Θ_1, Θ_2 of a frame S is still a coarsening of S .

Proof. From the hypothesis S is a common refinement of Θ_1 and Θ_2 , and since the minimal refinement is coarsening of every other refinement the thesis follows. \square

Immediately we conclude that

Theorem 3. The subfamily of compatible frames generated by the application of the restricted set of rules $A_{1..4,7}$ to a basis set S is the collection of all the disjoint partitions of S along with the appropriate refinings.

Note that this is not necessarily true when using Axiom A6.

5. Monoidal structure

The ‘constructive’ version of Definition 8 allows us to introduce in a family of compatible frames the internal operation

$$\begin{aligned} \otimes : \mathcal{F} \times \dots \times \mathcal{F} &\rightarrow \mathcal{F} \\ \{\Theta_1, \dots, \Theta_n\} &\mapsto \otimes_i \Theta_i \end{aligned} \quad (7)$$

associating each collection of frames with their minimal refinement. Equation (7) is well defined, for Axiom A7 ensures the existence of $\otimes_i \Theta_i$.

5.1. Finite families as commutative monoids

Let us then consider a finite subfamily of compatible frames $\mathcal{F}' \doteq \langle S \rangle_{A_{1..4,7}}$ for some $S \in \mathcal{F}$, and discuss the properties of the minimal refinement.

Theorem 4. The internal operation \otimes of minimal refinement satisfies the following properties:

1. *associativity:* $\Theta_1 \otimes (\Theta_2 \otimes \Theta_3) = (\Theta_1 \otimes \Theta_2) \otimes \Theta_3, \quad \forall \Theta_1, \Theta_2, \Theta_3 \in \mathcal{F}'$;
2. *commutativity:* $\Theta_1 \otimes \Theta_2 = \Theta_2 \otimes \Theta_1, \quad \forall \Theta_1, \Theta_2 \in \mathcal{F}'$;

3. *existence of unit*: $\exists \mathbf{1} \in \mathcal{F}'$ s.t. $\Theta \otimes \mathbf{1} = \Theta$, $\forall \Theta \in \mathcal{F}'$;
 4. *annihilator*: $\Theta \otimes S = S$, $\forall \Theta \in \mathcal{F}'$

where $\mathbf{1}$ is the unique frame of \mathcal{F}' containing a single element.

Proof. Associativity and commutativity. If we look at equation (5), associativity and commutativity follow from the analogous property of set-theoretic intersection.

Unit. Let us prove that there exists a *unique* frame $\mathbf{1}$ in \mathcal{F}' with cardinality 1. Given $\Theta \in \mathcal{F}'$ due to Axiom A4 (existence of coarsenings) there exists a coarsening $\mathbf{1}_\Theta$ of Θ and a refining $\omega_{\mathbf{1}_\Theta} : 2^{\mathbf{1}_\Theta} \rightarrow 2^\Theta$. Then by Axiom A1 there exists another refining $\mathbf{1}_\omega \in \mathcal{R}'$ such that

$$\mathbf{1}_\omega : 2^{\mathbf{1}_\Theta} \rightarrow 2^S.$$

Now, if we consider a pair of elements of \mathcal{F}' , say Θ_1, Θ_2 , the above procedure yields two pairs set-refining $(\mathbf{1}_{\Theta_i}, \mathbf{1}_{\omega_i})$ with $\mathbf{1}_{\omega_i} : 2^{\mathbf{1}_{\Theta_i}} \rightarrow 2^S$. But if we call 1_{θ_i} the unique element of $\mathbf{1}_{\Theta_i}$

$$\mathbf{1}_{\omega_i}(\{1_{\theta_i}\}) = S \quad \forall i = 1, 2$$

so that for Axiom A2 (identity of coarsenings) uniqueness follows: $\mathbf{1}_{\omega_1} = \mathbf{1}_{\omega_2}$, $\mathbf{1}_{\Theta_1} = \mathbf{1}_{\Theta_2}$.

Annihilator. If $\Theta \in \mathcal{F}'$ then obviously Θ is a coarsening of S , therefore their minimal refinement coincides with S itself. \square

In other words, a finite family of frames of discernment (\mathcal{F}', \otimes) is a *finite commutative monoid with annihilator* with respect to the internal operation of minimal refinement.

However, a family of frames is not only a collection of sets, but also the collection of refinings between these sets. Does Theorem 4 have any consequence for these refinings?

5.2. Isomorphism frames–refinings

Actually, a family of frames can be dually viewed as a set of refining maps with attached their domains and codomains (perhaps the most synthetic approach as it accounts for sets automatically, allowing to reduce the number of axioms). In the finite case we can immediately establish the following correspondence

$$\Theta \longleftrightarrow \omega_S^\Theta : 2^\Theta \rightarrow 2^S \quad (8)$$

between an element Θ of a finite subfamily of compatible frames and the unique refining from Θ to S (see Axiom A3).

Definition 11. Given $\omega_1 : 2^{\Theta_1} \rightarrow 2^S$ and $\omega_2 : 2^{\Theta_2} \rightarrow 2^S$ their composition induced by the operation of minimal refinement is the unique refining from $\Theta_1 \otimes \Theta_2$ to S (as for Axiom A3 there is only one refining connecting two frames of the family)

$$\omega_1 \otimes \omega_2 : 2^{\Theta_1 \otimes \Theta_2} \rightarrow 2^S. \quad (9)$$

Theorem 5. The subcollection of the refinings of a finite family of compatible frames $\langle S \rangle_{A1.4.7}$ with codomain S

$$\langle S \rangle_{A1.4.7}^{\omega_S^{(\cdot)}} \doteq \{\omega_S^\Theta, \Theta \in \langle S \rangle_{A1.4.7}\}$$

is a finite commutative monoid with annihilator with respect to equation (9).

Proof. Associativity and commutativity come from the analogous property of the minimal refinement operator. The unit element is $\mathbf{1}_\omega : 2^S \rightarrow 2^S$, as

$$Dom(\mathbf{1}_\omega \otimes \omega) = \mathbf{1}_S \otimes Dom(\omega) = Dom(\omega) \Rightarrow \mathbf{1}_\omega \otimes \omega = \omega$$

for Axiom A2. On the other side if we consider $\mathbf{0}_\omega : 2^S \rightarrow 2^S$ the unique refining from S onto itself (which exists by Axiom A4 with $n = 1$) we have

$$Dom(\mathbf{0}_\omega \otimes \omega) = S \otimes Dom(\omega) = S$$

so that $\mathbf{0}_\omega$ is an annihilator for $\langle S \rangle_{A1.4.7}^{\omega_S^{(\cdot)}}$. \square

It easily follows that

Proposition 2. Given a finite family of compatible frames, the bijection (8) is an isomorphism between commutative monoids.

5.2.1. Some remarks

It may be interesting to note that the existence of both unit element and annihilator in a finite family of frames are consequences of the following proposition:

Proposition 3. $(\omega_S^\Theta \circ \omega_\Theta^\Omega) \otimes \omega_S^\Theta = \omega_S^\Theta \quad \forall \Omega, \Theta.$

Proof. $Cod(\omega_S^\Theta \circ \omega_\Theta^\Omega = S)$ so the operation \otimes can be applied. By noting that $Dom(\omega_S^\Theta \circ \omega_\Theta^\Omega) = \Omega$ is a coarsening of $Dom(\omega_S^\Theta) = \Theta$ we get

$$Dom((\omega_S^\Theta \circ \omega_\Theta^\Omega) \otimes \omega_S^\Theta) = Dom(\omega_S^\Theta \circ \omega_\Theta^\Omega) \otimes Dom(\omega_S^\Theta) = Dom(\omega_S^\Theta) = \Theta$$

and from Axiom A3 the thesis follows. \square

As a matter of fact, when $\Omega = \mathbf{1}_S$ then $\omega_S^\Theta \circ \omega_\Theta^\Omega = \omega_S^\Theta \circ \omega_\Theta^{\mathbf{1}_S} = \mathbf{1}_\omega$ and we get $\mathbf{1}_\omega \otimes \omega_S^\Theta = \omega_S^\Theta$. On the other side, when $\Theta = S$ then $\omega_S^\Theta = \omega_S^S = \mathbf{0}_\omega$ and we have the annihilation property $\omega_S^\Omega \otimes \mathbf{0}_\omega = \mathbf{0}_\omega$.

5.3. Generators

Even if not strictly necessary in the context of the conflict problem, something more can be said about the dual finite monoids generated by \otimes in terms of the notion of *generator*. Given a monoid \mathcal{M} , the submonoid $\langle M \rangle$ generated by a subset M of \mathcal{M} is defined as the intersection of all the submonoids of \mathcal{M} containing M .

Definition 12. The set of *generators* of a monoid \mathcal{M} is a finite collection M of elements of \mathcal{M} such that $\langle M \rangle = \mathcal{M}$.

Theorem 6. The set of generators of the finite family of compatible frames $\langle S \rangle_{A_{1..4,7}}$ seen as finite commutative monoid with respect to the internal operation \otimes (minimal refinement) is the collections of all the binary frames.

The set of generators of the associated monoid of refinings $\langle S \rangle_{A_{1..4,7}}^{\omega_S^{(\cdot)}}$ is the collection of refinings from all the binary partitions of S to S itself:

$$\langle S \rangle_{A_{1..4,7}}^{\omega_S^{(\cdot)}} = \langle \{ \omega_{ij} : 2^{\Theta_{ij}} \rightarrow 2^S \mid |\Theta_{ij}| = 2, \omega_{ij} \in \mathcal{R} \} \rangle.$$

Proof. We need to prove that all the possible partitions of S can be obtained as minimal refinement of a number of binary partitions. Let us consider a generic partition $\Pi = \{S^1, \dots, S^n\}$ and define

$$\begin{aligned} \Pi_1 &= \{S^1, S^2 \cup \dots \cup S^n\} \doteq \{A_1, B_1\} \\ \Pi_2 &= \{S^1 \cup S^2, S^3 \cup \dots \cup S^n\} \doteq \{A_2, B_2\} \quad \dots \\ \Pi_{n-1} &= \{S^1 \cup \dots \cup S^{n-1}, S^n\} \doteq \{A_{n-1}, B_{n-1}\}. \end{aligned}$$

It is not difficult to see that every arbitrary intersection of elements of Π_1, \dots, Π_{n-1} is an element of the n -ary partition Π , in fact

$$A_i \cap B_k = \emptyset, \quad k \geq i, \quad A_i \cap A_k = A_i, \quad k \geq i, \quad B_i \cap B_k = B_k, \quad k \geq i$$

so that $\bigcap_i A_i = A_1 = S^1$, $\bigcap_i B_i = B_{n-1} = S^n$. If both A s and B s are present the resulting intersection is \emptyset whenever there exist a pair B_l, A_m with $l \geq m$. Hence the only non-empty mixed intersections in the class $\{X_1 \cap \dots \cap X_n\}$ must necessarily be of the following kind

$$B_1 \cap \dots \cap B_k \cap A_{k+1} \cap \dots \cap A_{n-1}$$

and we have

$$\begin{aligned} B_1 \cap \dots \cap B_k \cap A_{k+1} \cap \dots \cap A_{n-1} &= \\ &= (S^{k+1} \cup \dots \cup S^n) \cap (S^1 \cup \dots \cup S^{k+1}) = S^{k+1} \end{aligned}$$

where $k + 1$ goes from 2 to $n - 1$. That satisfies the fundamental condition for Π to be the minimal refinement of Π_1, \dots, Π_{n-1} (note that the choice of the binary frames is

not unique). The second part of the proof, concerning the refining maps, comes directly from the existence of the isomorphism (8). \square

The monoids associated with finite families are then really simple, their structure being completely determined by the binary frames and related refinings. This is not true for complete families of compatible frames. Even if the general case is not a copycat of the finite one, though, the analysis of finite subfamilies provides us with the necessary intuition to cope with the general case.

5.4. General families as commutative monoids

The algebraic structure of monoid is then a direct implication of the notion of minimal refinement. We may wonder whether a general family of compatible frames possess the same structure. The answer is positive.

Theorem 7. \mathcal{F} is an infinite commutative monoid with no annihilators.

Proof. The proof of Theorem 4 still holds for the first two points. *Existence of unit.* Suppose there exist two frames $\mathbf{1} = \{1\}$ and $\mathbf{1}' = \{1'\}$ of size 1. For Axiom A6 they have a common refinement Θ , with $\omega_1 : 2^{\mathbf{1}} \rightarrow 2^{\Theta}$ and $\omega_{1'} : 2^{\mathbf{1}'} \rightarrow 2^{\Theta}$ refinings. But then $\omega_1(\{1\}) = \Theta = \omega_{1'}(\{1'\})$, and by Axiom A2 $\mathbf{1} = \mathbf{1}'$. Now, for every frame $\Omega \in \mathcal{F}$ Axiom A4 ensures that there exists a partition of Ω with only one element, $\mathbf{1}_{\Omega}$. For the above argument, $\mathbf{1}_{\Omega} = \mathbf{1}$. In conclusion there is only one monic frame in \mathcal{F} , and since it is a refinement of every other frame, it is the unit element with respect to \otimes .

Annihilators. If a frame $\mathbf{0}_{\Theta}$ would exist such that $\mathbf{0}_{\Theta} \otimes \Theta = \mathbf{0}_{\Theta}$ for each Θ we would have, given a refinement Ω of $\mathbf{0}_{\Theta}$ (built by means of Axiom A5), $\mathbf{0}_{\Theta} \otimes \Omega = \Omega$ that is a contradiction. \square

The finite case also suggests the presence of a similar structure for the set of refinings \mathcal{R} . However, in a complete family of frames whatever basis set S you choose there are refinings with codomain distinct from S , so that it is impossible to write down a simple correspondence between frames and maps as in equation (8).

Nevertheless, if we note that given two maps ω_1 and ω_2 their codomains Ω_1, Ω_2 always have a common refinement Ω , we can write

$$\omega_1 \otimes \omega_2 \doteq \omega'_1 \otimes \omega'_2 \quad (10)$$

where, after calling w_1 (w_2) the refining map between Ω_1 (Ω_2) and Ω ,

$$\omega'_1 = w_1 \circ \omega_1, \quad \omega'_2 = w_2 \circ \omega_2$$

and \otimes on the right side of equation (10) stands for the composition of refinings in the finite family of compatible frames $\langle \Omega \rangle_{A_{1.4.7}}$ generated by Ω . This way the operation \otimes is again well-defined.

The composition of refinings can be rewritten in a more elegant way, proving its validity for general families too.

Definition 13. Given two maps $\omega_1, \omega_2 \in \mathcal{R}$, $\omega_1 : 2^{\Theta_1} \rightarrow 2^{\Omega_1}$ and $\omega_2 : 2^{\Theta_2} \rightarrow 2^{\Omega_2}$ we define

$$\omega_1 \otimes \omega_2 : 2^{\Theta_1 \otimes \Theta_2} \rightarrow 2^{\Omega_1 \otimes \Omega_2}. \quad (11)$$

This operation is again well-defined, for the correspondence

$$(Dom(\omega), Cod(\omega)) \leftrightarrow \omega \quad (12)$$

(guaranteed by Axiom A3) is a bijection. It obviously reduces to Definition 11 when applied to finite subfamilies.

Theorem 8. The set of refinings \mathcal{R} of a complete generated family of compatible frames is a commutative monoid with respect to the internal operation (11).

Proof. Obviously \otimes is commutative and associative for commutativity and associativity of the operation of minimal refinement of frames. To find the unit element $\mathbf{1}_\omega$ it suffices to note that $\mathbf{1}_\omega$ is in correspondence through (12) with a pair of frames (Θ, Ω) such that

$$\begin{aligned} \Theta \otimes \Theta_1 &= \Theta_1 & \forall \Theta_1 \in \mathcal{F}, \\ \Omega \otimes \Omega_1 &= \Omega_1 & \forall \Omega_1 \in \mathcal{F} \end{aligned}$$

but that means $\Theta = \Omega = \mathbf{1}$, so that $\mathbf{1}_\omega : 2^{\mathbf{1}} \rightarrow 2^{\mathbf{1}}$ and $\mathbf{1}_\omega$ turns out to be nothing but the identity map on $\mathbf{1}$. \square

Corollary 1. \mathcal{R} is a submonoid of the product monoid $(\mathcal{F}, \otimes) \times (\mathcal{F}, \otimes)$ through the map (12).

5.5. Monoidal structure of families of compatible frames

In order to summarize what we have learned about the algebraic structures induced on families of compatible frames by the operation of minimal refinement, it is worth to specify the missing relations among them. Clearly, $(\langle S \rangle_{A_{1..4,7}}^\omega, \otimes)$ (where $\langle S \rangle_{A_{1..4,7}}^\omega$ is the collection of *all* the refinings of a finite family of frames) is a monoid, too, and

Proposition 4. $(\langle S \rangle_{A_{1..4,7}}^{\omega_s^{(i)}}, \otimes)$ is a submonoid of $(\langle S \rangle_{A_{1..4,7}}^\omega, \otimes)$.

Proof. Obviously $\langle S \rangle_{A_{1.4.7}}^{\omega_s^{(,)}}$ \subset $\langle S \rangle_{A_{1.4.7}}^\omega$ in a set-theoretic sense. We only have to prove that the internal operation of the first set is inherited from the second one. If $\omega_1 : 2^{\Theta_1} \rightarrow 2^S$ and $\omega_2 : 2^{\Theta_2} \rightarrow 2^S$ then $S \otimes S = S$ and

$$\omega_1 \otimes_{\langle S \rangle^\omega} \omega_2 : 2^{\Theta_1 \otimes \Theta_2} \rightarrow 2^{S \otimes S} = 2^S \equiv \omega_1 \otimes_{\langle S \rangle^{\omega_s^{(,)}}} \omega_2 \quad \square$$

In turn,

Proposition 5. $(\langle S \rangle_{A_{1.4.7}}^\omega, \otimes)$ is a submonoid of (\mathcal{R}, \otimes) .

Proof. It suffices to prove that $\langle S \rangle_{A_{1.4.7}}^\omega$ is closed with respect to the composition operator (11). But then, given two maps ω_1, ω_2 whose domains and codomains are both coarsenings of S , $Dom(\omega_1 \otimes \omega_2) = Dom(\omega_1) \otimes Dom(\omega_2)$ and $Cod(\omega_1 \otimes \omega_2) = Cod(\omega_1) \otimes Cod(\omega_2)$ are still coarsenings of S . \square

while as far as frames are concerned,

Proposition 6. $(\langle S \rangle_{A_{1.4.7}}^\Theta, \otimes)$ is a submonoid of (\mathcal{F}, \otimes) .

Proof. Obvious, for the finite family of compatible frames $\langle S \rangle_{A_{1.4.7}}^\Theta$ is strictly included in the complete one \mathcal{F} , and $\langle S \rangle_{A_{1.4.7}}^\Theta$ is closed with respect to \otimes , i.e., if Θ_1 and Θ_2 are coarsenings of S then their minimal refinement is still a coarsening of S . \square

A comprehensive picture of the relations between the monoidal structures associated with a family of compatible frames $(\mathcal{F}, \mathcal{R})$ can then be appreciated in the following diagram:

$$\begin{array}{ccccc} (\langle S \rangle_{A_{1.4.7}}^{\omega_s^{(,)}}, \otimes) & \subset & (\langle S \rangle_{A_{1.4.7}}^\omega, \otimes) & \subset & (\mathcal{R}, \otimes) \\ \wr & & \wr & & \\ (\langle S \rangle_{A_{1.4.7}}^\Theta, \otimes) & \subset & (\langle S \rangle_{A_{1.4.7}}^\Theta, \otimes) \times (\langle S \rangle_{A_{1.4.7}}^\Theta, \otimes) & \cap & (13) \\ = & & & & \\ (\langle S \rangle_{A_{1.4.7}}^\Theta, \otimes) & \subset & (\mathcal{F}, \otimes) & \subset & (\mathcal{F}, \otimes) \times (\mathcal{F}, \otimes) \end{array}$$

where \sim denotes an isomorphism, and \subset an inclusion between monoids.

6. Lattice structure

The notion of minimal refinement is the essence of the idea of families of compatible frames of discernment. This fact is reflected in the elegant scheme above, where a hidden matrioska of encapsulated monoids is depicted. It is still unclear,

though, what is the deep root of the analogy between independence relations we discussed in the Introduction, with its potential role in the conflict problem. Let us then take a step further.

6.1. Minimal refinement as least upper bound

It is well-known (see [13], page 456) that the internal operation \cdot of a monoid \mathcal{M} induces an order relation among the elements of \mathcal{M} , namely $b \geq a \equiv \exists c \text{ s.t. } b = a \cdot c$. In our case this yields

$$\Theta_2 \geq \Theta_1 \equiv \exists \Theta_3 \text{ s.t. } \Theta_2 = \Theta_1 \otimes \Theta_3 \quad (14)$$

$\equiv \Theta_1 \otimes \Theta_2 = \Theta_2$, i.e., Θ_2 is a refinement of Θ_1 .⁵

Since both finite and general families of frames are monoids,

Proposition 7. $\langle S \rangle_{A_{1..4,7}}$ and \mathcal{F} are partially ordered set (posets) with respect to the order relation (14).

In a poset the dual notions of *least upper bound* and *greatest lower bound* of a pair of elements can be introduced.

Definition 14. Given two elements $x, y \in X$ of a poset X their *least upper bound* $\sup_X(x, y)$ is the smallest element of X that is bigger than both x and y , i.e., $\sup_X(x, y) \geq x, y$ and

$$\text{if } \exists z \text{ s.t. } z \leq \sup_X(x, y), \quad z \geq x, y \quad \text{then } z = \sup_X(x, y).$$

Definition 15. Given two elements $x, y \in X$ of a poset X their *greatest lower bound* $\inf_X(x, y)$ is the biggest element of X that is smaller than both x and y , i.e., $\inf_X(x, y) \leq x, y$ and

$$\text{if } \exists z \text{ s.t. } z \geq \inf_X(x, y), \quad z \leq x, y \quad \text{then } z = \inf_X(x, y).$$

The standard notation is $\inf(x, y) = x \wedge y$ and $\sup(x, y) = x \vee y$. By induction \sup and \inf can be defined for arbitrary finite collections too.

It must be pointed out that not any pair of elements of a poset, in general, admits \inf and/or \sup .

⁵ It is interesting to note that Proposition (3) can be expressed in terms of the order relation $\omega_2 \geq \omega_1 \Rightarrow \omega_1 \otimes \omega_2 = \omega_2$. Again, since $\omega_2 \geq \omega_1 \iff \omega_2 = \omega_1 \otimes \omega_3$ for some ω_3 , it can be rewritten as

$$\omega_1 \otimes (\omega_1 \otimes \omega_3) = (\omega_1 \otimes \omega_1) \otimes \omega_3 = \omega_1 \otimes \omega_3. \quad (15)$$

Hence $\omega \otimes \omega = \omega$ (*idempotence*) is a sufficient condition.

Definition 16. A *lattice* is a poset in which each pair of elements admits both inf and sup.

Definition 17. An infinite lattice L is said *complete* if each arbitrary collection (even not finite) of points in L admits both sup and inf.

In this case there exist $\mathbf{0} \equiv \wedge L$, $\mathbf{1} \equiv \vee L$ called respectively *initial* and *final* element of L .

The notion of lattice provides us with an alternative interpretation of the minimal refinement operation.

Theorem 9. In a family of compatible frames \mathcal{F} seen as a poset the sup of a finite collection of frames coincides with the minimal refinement,

$$\sup_{\mathcal{F}}(\Theta_1, \dots, \Theta_n) = \Theta_1 \otimes \dots \otimes \Theta_n.$$

Proof. Of course $\Theta_1 \otimes \dots \otimes \Theta_n \geq \Theta_i \forall i = 1, \dots, n$ for there exists a refining between each Θ_i and their minimal refinement. Now, if there exists another frame Ω greater than each Θ_i , then Ω is a common refinement for $\Theta_1, \dots, \Theta_n$, hence it is a refinement of the minimal refinement, i.e., $\Omega \geq \Theta_1 \otimes \dots \otimes \Theta_n$ according to the order relation (14). \square

At a first glance is not clear what $\inf\{\Theta_1, \dots, \Theta_n\}$ should represent.

6.2. Maximal coarsening as greatest lower bound

Let us then introduce a new notion.

Definition 18. A *common coarsening* of two frames Θ_1, Θ_2 is a set $\Omega \in \mathcal{F}$ such that $\exists \omega_1 : 2^\Omega \rightarrow 2^{\Theta_1}$ and $\omega_2 : 2^\Omega \rightarrow 2^{\Theta_2}$ refinings, i.e., is a coarsening of both frames.

This is well defined, as

Theorem 10. If $\Theta_1, \Theta_2 \in \mathcal{F}$ are elements of a family of compatible frames then they have a common coarsening.

Proof. From the proof of Theorem 7 they have at least $\mathbf{1}$ as common coarsening. \square

Theorem 10 is nothing but the dual of Axiom A6 (see section 2.5). This opens the way to the introduction of a new operator acting on collections of compatible frames, the dual of the minimal refinement.

Theorem 11. Given a collection $\Theta_1, \dots, \Theta_n$ of elements of a family of compatible frames \mathcal{F} there exists a unique element $\Omega \in \mathcal{F}$ such that:

1. $\forall i$ there exists a refining $\omega_i : 2^\Omega \rightarrow 2^{\Theta_i}$;
2. $\forall \theta \in \Omega \nexists A_1 \subseteq \omega_1(\{\theta\}), \dots, A_n \subseteq \omega_n(\{\theta\})$ s.t. $\eta_1(A_1) = \dots = \eta_n(A_n)$

where $\eta_i : 2^{\Theta_i} \rightarrow 2^{\Theta_1 \otimes \dots \otimes \Theta_n}$.

We first need a simple intermediate result.

Lemma 2. Suppose $\Theta_1 \otimes \dots \otimes \Theta_n$ is the minimal refinement of $\Theta_1, \dots, \Theta_n$, with refinings $\eta_i : 2^{\Theta_i} \rightarrow 2^{\Theta_1 \otimes \dots \otimes \Theta_n}$, and there exist $X_1 \subset \Theta_1, \dots, X_n \subset \Theta_n$ with $\eta_1(X_1) = \dots = \eta_n(X_n)$ such that $\nexists A_1 \subseteq X_1, \dots, A_n \subseteq X_n$ s.t. $\eta_1(A_1) = \dots = \eta_n(A_n)$. Then for every common coarsening Θ of $\Theta_1, \dots, \Theta_n$, with refinings $\omega_i : 2^\Theta \rightarrow 2^{\Theta_i}$, there exists $\theta \in \Theta$ such that $X_i \subseteq \omega_i(\{\theta\}) \forall i = 1, \dots, n$.

Proof. Let us suppose there not exists such an element θ , but X_i is covered by a subset $\bar{X} \subset \Theta$. If we consider one of these elements $\bar{\theta} \in \bar{X}$ we have

$$\eta_i(\omega_i(\bar{\theta}) \cap X_i) = \eta_i(\omega_i(\bar{\theta})) \cap \eta_i(X_i)$$

but then $\eta_1(\omega_1(\bar{\theta})) = \dots = \eta_n(\omega_n(\bar{\theta}))$ by definition of common coarsening, and $\eta_1(X_1) = \dots = \eta_n(X_n)$ for hypothesis, so that

$$\eta_1(\omega_1(\bar{\theta}) \cap X_1) = \dots = \eta_n(\omega_n(\bar{\theta}) \cap X_n)$$

with $A_i \doteq \omega_i(\bar{\theta}) \cap X_i \subseteq X_i \forall i = 1, \dots, n$, which is a contradiction. □

Now we can face the proof of Theorem 11.

Proof. Existence. The proof is constructive. Let us take an arbitrary coarsening \mathcal{L} of $\Theta_1, \dots, \Theta_n$ (which exists by Theorem 10) and check for every $l \in \mathcal{L}$ whether there exists a collection of subsets $\{A_i \subseteq \omega_i(\{l\}), i = 1, \dots, n\}$ such that $\eta_1(A_1) = \dots = \eta_n(A_n)$. If the answer is negative we have the desired frame. Otherwise we can build a new common coarsening \mathcal{L}' of $\Theta_1, \dots, \Theta_n$ by simply splitting $\{l\}$ into a pair $\{l_1, l_2\}$ where

$$\omega'_i(\{l_1\}) = A_i, \omega'_i(\{l_2\}) = B_i \quad \forall i = 1, \dots, n$$

having defined $B_i = \omega_i(\{l\}) \setminus A_i$. This can always be done, for if $\omega_i(\{l\}) \setminus A_i \neq \emptyset$ for some i then it is not void $\forall i$.

This procedure can be iterated until there are not subsets satisfying condition 2. It is also guaranteed to terminate, as the number of possible bijections of the images $\omega_i(\{l\})$ is finite. More precisely, the maximum number of steps is

$$\lceil \log_2 \max_{l \in \mathcal{L}} \min_{i=1, \dots, n} |\omega_i(\{l\})| \rceil.$$

Uniqueness. Suppose Ω' is another common coarsening satisfying condition 2, with refinings $\omega'_i : 2^{\Omega'} \rightarrow 2^{\Theta_i}$, distinct from Ω . If we define $X_i \doteq \omega_i(\{\theta\})$, $\theta \in \Omega$, by lemma 2 there exists $\theta' \in \Omega'$ such that $\omega_i(\{\theta\}) \subset \omega'_i(\{\theta'\})$. But then condition 2 implies $\omega_i(\{\theta\}) = \omega'_i(\{\theta'\})$ for every pair θ, θ' so that $\Omega = \Omega'$. \square

Definition 19. This unique frame is denoted by $\Theta_1 \oplus \dots \oplus \Theta_n$, and called *maximal coarsening* of $\Theta_1, \dots, \Theta_n$.

Theorem 12. If Ω is a common coarsening of a finite set of compatible frames $\Theta_1, \dots, \Theta_n$ then Ω is a coarsening of their maximal coarsening too, i.e., there exists a refining $\omega : 2^\Omega \rightarrow 2^{\Theta_1 \oplus \dots \oplus \Theta_n}$.

Proof. Let us consider another common coarsening Ω' of $\Theta_1, \dots, \Theta_n$, with $\alpha_i : 2^{\Omega'} \rightarrow 2^{\Theta_i}$. If it satisfies condition 2 of Theorem 11 then for the uniqueness of $\Theta_1 \oplus \dots \oplus \Theta_n$ it coincides with the maximal coarsening. Otherwise the procedure of the above proof can be used to produce such a frame. Again, uniqueness guarantees that this is the maximal coarsening and by construction it is a refinement of Ω' . \square

In other words,

Corollary 2. The maximal coarsening $\Theta_1 \oplus \dots \oplus \Theta_n$ of a collection of frames $\Theta_1, \dots, \Theta_n$ coincides with the greatest lower bound (inf) of the collection of frames as elements of the poset (\mathcal{F}, \geq) .

6.3. Birkhoff property

Proposition 1 and Theorems 9, 11 and 12 have a straightforward consequence on the algebraic structure of \mathcal{F} .

Corollary 3. The collection of sets \mathcal{F} of a family of compatible frames of discernment is a lattice, with respect to the internal operations of minimal refinement and maximal coarsening.

As we will see in this last part of the paper, the lattice-theoretic language turns out to be the right tool for discussing the algebraic nature of the conflict problem.

From a technical point of view, it must be pointed out that \mathcal{F} lacks the attribute of completeness: the axioms do not guarantee the existence of a minimal refinement for an infinite (even countable) collection of sets. A more precise characterization of $(\mathcal{F}, \otimes, \oplus)$ can be given after introducing a few other lattice-theoretic notions (a reference can be found in [25] and [26]).

Definition 20. A lattice L is modular iff if $a \geq b$, $a \wedge c = b \wedge c$, $a \vee c = b \vee c$ then $a = b$.




Figure 6. Example of partitions like Π_1 (solid line) and Π_2 (dashed line) in Theorem 13.

Definition 21. A lattice L is *Birkhoff* when $a, b \succ a \wedge b$ implies $a \vee b \succ a, b$ for all $a, b \in L$, where $x \succ y$ (x ‘covers’ y) means that $x > y$ and $\nexists z : x > z > y$.

It can be proved that the Birkhoff property is weaker than modularity [25].

Definition 22. A poset is said to have *finite length* if the length of its chains (collections of consecutive elements) is bounded.

Now we have the tools to precise the nature of the lattices of compatible frames.

Theorem 13. A finite family of compatible frames $(\langle S \rangle_{A_{1,\dots,4,7}}, \geq)$ is a *complete Birkhoff lattice of finite length*.⁶

Proof. $(\langle S \rangle_{A_{1,\dots,4,7}}, \geq)$ is *complete*. Actually each finite lattice is complete for it does not contain any infinite collection of elements. $(\langle S \rangle_{A_{1,\dots,4,7}}, \geq)$ is *Birkhoff*. Assume that Θ_1 and Θ_2 both cover their maximal coarsening. Necessarily $\Theta_1 \oplus \Theta_2$ must have rank

$$|\Theta_1 \oplus \Theta_2| = |\Theta_1| - 1 = |\Theta_2| - 1,$$

so that both refinings $\omega_1 : 2^{\Theta_1 \oplus \Theta_2} \rightarrow 2^{\Theta_1}$ and $\omega_2 : 2^{\Theta_1 \oplus \Theta_2} \rightarrow 2^{\Theta_2}$ leave unchanged each point of $\Theta_1 \oplus \Theta_2$ but one, replaced by two new elements.

Now, on the other side Θ_1 and Θ_2 represent two different partitions Π_1, Π_2 of their minimal refinement. By construction these partitions coincide but for the elements obtained by refining these two points, as shown in figure 6.

But then, their minimal refinement must be by definition the superposition of Π_1, Π_2 : analytically

$$\Pi_1 = \{\{p_1, p_2\}, p_3, \dots, p_n\}, \quad \Pi_2 = \{p_1, \dots, p_{n-2}, \{p_{n-1}, p_n\}\}$$

⁶ For an alternative proof see [26].

having denoted their minimal refinement by $\{p_1, p_2, \dots, p_{n-1}, p_n\}$. Its rank is then equal to $|\Theta_1 \otimes \Theta_2| = |\Theta_1| + 1 = |\Theta_2| + 1$. Hence it covers both frames, for there cannot exist any frame with intermediate size between Θ_1 or Θ_2 and $\Theta_1 \otimes \Theta_2$. \square

After looking at the proof of Theorem 13 we can observe how it proves the Birkhoff property for *general* families of frames, too. As a matter of fact, being Birkhoff is a *local* property involving the sublattice $[\Theta_1 \oplus \dots \oplus \Theta_n, \Theta_1 \otimes \dots \otimes \Theta_n]$.

Corollary 4. The collection of sets \mathcal{F} of a family of compatible frames is a *locally Birkhoff lattice bounded below*, i.e., is a Birkhoff lattice of locally finite length with initial element.

Proof. It just remains to point out that, for Theorem 7, each arbitrary collection of frames in \mathcal{F} admits a common coarsening $\mathbf{1}$, playing the role of initial element of the lattice. \square

We are hence finally able to discuss the difference between vector spaces and compatible frames in algebraic terms. As a matter of fact, it turns out that the collection of vector subspaces of a vector space forms a *modular* lattice [1]. Lattices of compatible frames, instead, *are not modular*: Figure 7 shows a simple counterexample. Both the pairs formed by picking one of the left-most frames and the right-most one have the same minimal refinement and maximal coarsening (the unit




Figure 7. Non-modularity of finite families of frames: a counter-example.

frame). However, they are different even though linked by a refining, violating Definition 20.

7. Independence and pseudo-Gram–Schmidt algorithm

Maybe it is time for a little summary. So far we have proved that the problem of conflicting belief functions, so important for sensor fusion applications, is inherently related to the notion of independence of frames. This in turn is strikingly similar to the notion of independence of vector subspaces. We have wondered whether this is more than a simple analogy or there is instead a deep connection, possibly providing us with a formally correct solution of the conflict problem.

We posed the question in an algebraic setup, and investigated the properties of the axioms defining a family of frames, in particular the notion of minimal refinement. We now know that both frames and refinings form monoids with respect to \otimes . This brought us to explore a possible description in terms of partially ordered sets. We learned how to define a dual operation in the families of compatible frames of discernment, called maximal coarsening \oplus . These families form in fact a lattice with respect to \oplus, \otimes .

The bottom line is that this particular class of lattices, called ‘Birkhoff,’ also includes the lattice of the vector subspaces of a vector space. Actually, the latter is something more (a ‘modular’ lattice). As we are going to see, Birkhoff lattices are among the algebraic structures where the concept of *independence* can be introduced. In this last Section we will discuss the possible outcomes of our algebraic analysis, and conjecture a possible formal solution of the conflict problem.

7.1. Linear independence and Birkhoff lattices

The following definition is due to Szasz [26].

Definition 23. Consider the collection $\mathcal{P}(M)$ of all the finite subsets of a given set M and define in the product $M \times \mathcal{P}(M)$ a relation Λ . Λ is said to be a *linear dependence* on the set M if it satisfies the following conditions:

1. $p_j \Lambda \{p_1, \dots, p_m\}, j = 1, \dots, m$;
2. if $p \Lambda \{p_1, \dots, p_m\}$ and $\forall j p_j \Lambda \{q_1, \dots, q_n\}$, then $p \Lambda \{q_1, \dots, q_n\}$
3. if $p \Lambda \{p_1, \dots, p_m, q\}$ but $p \not\Lambda \{p_1, \dots, p_m\}$, then $q \Lambda \{p_1, \dots, p_m, p\}$

where Λ' means ‘is not in relation with.’

A set provided with a linear (in)dependence relation is often called *matroid* [19]. Definition 23 is rather cryptic, so maybe it may be useful to give of it an alternative

interpretation. More intuitively, a collection \mathcal{I} of elements of a set M (matroid) is called the *independent sets* of M if it meets the independence augmentation axiom: if I_1 and I_2 are in \mathcal{I} and $|I_1| < |I_2|$ then there exists an element e of $I_2 - I_1$ such that $I_1 \cup e \in \mathcal{I}$. Try and think at the usual vector space case, and you will realize it does indeed satisfy this condition.

Some elements of a Birkhoff lattice do admit a linear dependence relation. Let us call *rank* $h(a)$ of an element a of a finite length semimodular lattice bounded below L , the length of the interval $I[0, a]$ (i.e., the sublattice $\{x \in L : 0 \leq x \leq a\}$). In a lattice bounded below, the elements ‘covering’ 0 (i.e., such that there are no other elements in the chains containing them) are called *atoms*. It can be proved that (see [4], Chapter 6).

Theorem 14. If L is a Birkhoff lattice bounded below, the equivalence relation $\mathcal{LD} \subset L \times \mathcal{P}(L)$ defined as

$$p \mathcal{LD} \{p_1, \dots, p_n\} \equiv h(p \vee p_1 \vee \dots \vee p_n) < h(p_1 \vee \dots \vee p_n) + h(p) \quad (16)$$

where $p, p_1, \dots, p_n \in A$ (the set of atoms) and h is the rank of the elements of the lattice is a *linear dependence* relation.

In other words, the atoms of a Birkhoff lattice form a matroid. Actually, the notion of linear independence of atoms of a semimodular lattice is due to G. Birkhoff and H. Whitney [27], and is well known since the first half of the last century (consult his seminal works [2], or [3]).

7.2. Equivalence of independence relations

As a family of compatible frames \mathcal{F} is a Birkhoff lattice, it admits the lattice-theoretic *dependence* relation (16) on its atoms (the binary frames). On the other side, \mathcal{F} is also endowed with the evidence-theoretic independence relation (6). Are they related?

A connection between these two different notions of independence becomes clear when we notice that the linear *independence* relation associated with (16) is

$$\begin{aligned} p \mathcal{LI} \{p_1, \dots, p_n\} &\equiv p \overline{\mathcal{LD}} \{p_1, \dots, p_n\} \equiv h(p \vee p_1 \vee \dots \vee p_n) = \\ &= h(p_1 \vee \dots \vee p_n) + h(p) = h(p_1) + \dots + h(p_n) + h(p) \end{aligned} \quad (17)$$

where for a frame lattice $\Theta \vee \Omega = \Theta \otimes \Omega$.

But now, condition 4 of Theorem 1 can be written as

$$f(\Theta_1 \otimes \dots \otimes \Theta_n) = f(\Theta_1) + \dots + f(\Theta_n)$$

where $f = \lg(|\Theta|) = \lg(r(\Theta))$ and $r = |\Theta|$ is the rank of a frame Θ as element of the lattice $(\mathcal{F}, \otimes, \oplus)$. We can then conjecture that

Conjecture 1. The extension of the linear-independence relation (17) to *arbitrary elements* of the frame lattice and the evidence-theoretic independence of frames (6) are equivalent.

The necessary steps to achieve an equivalence proof become obvious:

1. proving that equation (17) defines a linear independence relation for arbitrary elements of a Birkhoff lattice, possibly enriched with additional properties;
2. showing that this can be also true for at least a suitable class of monotone functions of the rank (like logarithm).

To our knowledge none has yet proposed to give a matroidal structure to whole Birkhoff lattices. Maybe a naive, straightforward implementation of (17) is not feasible: the problem is likely to deserve a paper of its own.

7.3. Towards an algebraic solution of the conflict problem

Anyway, the above discussion suggests a way to solve the conflict among measurement belief functions without resorting to the conflict graph mentioned in section 3 [5, 6].

We have pointed out in the Introduction the striking similarity between independence of frames and vectors. Their formal expressions in terms of vectors (top) and subsets (bottom), respectively, confirm this impression

$$\begin{array}{lcl}
 \sum_i v_i \neq 0 & \iff & v_1 + \dots + v_n \neq 0, \forall v_i \in V_i \\
 & & \updownarrow \\
 \bigcap_i V_i = 0 & \iff & \text{span}\{V_1, \dots, V_n\} = \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_n} \\
 & & \updownarrow \\
 \bigoplus_i \Theta_i = \mathbf{0}_{\mathcal{F}} & \iff & \Theta_1 \otimes \dots \otimes \Theta_n = \Theta_1 \times \dots \times \Theta_n \\
 & & \updownarrow \\
 \bigcap A_t \neq \bigwedge & \iff & \omega_1(A_1) \cap \dots \cap \omega_n(A_n) \neq \emptyset
 \end{array}$$

where $\bigcap A_t \neq \bigwedge$ (with \bigwedge the initial element of a Boolean algebra) is the independence condition for Boolean sub-algebras [23].

This analogy has a formal root in Corollary 4. $L(V)$ and \mathcal{F} are both Birkhoff lattices, admitting the notion of independent elements. Moreover, the elements of these lattices (subspaces and frames) are *commutative monoids*. A vector space, indeed, is a group with respect to the operation ‘+’ and a frame is a Boolean algebra, i.e., a complete distributive lattice, hence they are both commutative monoids ([13]). If

Conjecture 1 is true, independence of frames and vector spaces may be nothing but different expressions of the matroidal structure of Birkhoff lattices.

We remember that, given a collection of arbitrary elements of a vector space, the well known Gram-Schmidt algorithm generates another collection of independent vectors, spanning the same subspace. Its basis elements are the independence condition and the *projection* of vectors onto other subspaces.

Conjecture 1 suggests the concrete possibility of designing a ‘pseudo Gram-Schmidt’ method resting on the shared algebraic structure of Birkhoff lattice of commutative monoids, and the associated independence relation. This algorithm, starting from a set of belief functions defined over a finite collection of discrete feature spaces, would build another collection of *independent* frames of the same family

$$\begin{array}{ccc} \Theta_1, \dots, \Theta_n \in \mathcal{F} & \longrightarrow & \Theta'_1, \dots, \Theta'_m \in \mathcal{F} \\ s_1, \dots, s_n & \mapsto & s'_1, \dots, s'_m \end{array}$$

with $m \neq n$ in general, and the same minimal refinement

$$\Theta_1 \otimes \dots \otimes \Theta_n = \Theta'_1 \otimes \dots \otimes \Theta'_m$$

Once projected the n b.f. s_1, \dots, s_n onto the new set of frames we would achieve a set of *surely combinable* belief functions s'_1, \dots, s'_m *equivalent*, in some sense, to the previous one. A formal definition of the equivalence of possibly non-combinable collections of belief functions needs of course to be introduced, even if it is reasonable to say it has to involve Dempster’s combination.

7.4. Perspectives

In this work, in conclusion, we have shown how an algebraic description of the families of frames could open the way to a solution of the problem of conflicting evidence by means of algebraic tools.

However, although we think the proposed method is the most elegant and rigorous approach to cope with groups of conflicting belief functions, the fundamental issues of Conjecture 1 and the pseudo Gram-Schmidt algorithm still need to be addressed prior to a practical application to information fusion problems (like the pose estimation task we discussed in section 3). For instance, a precise definition of the projection operator in the context of Birkhoff lattices of commutative monoids has to be formulated. Furthermore, power sets are not groups, so we cannot simply subtract the projected elements like in the original G-S algorithm. Difficult questions concern the meaning of the obtained collection of independent frames (in the object tracking context, we could ask whether they represent actual features), or the order in which the input data is ‘orthogonalized’.

The prize of this challenging pursuit would be an elegant and definitive arrangement of the conflict problem, and by this means a rigorous approach to sensor fusion tasks.

Of course the present algebraic analysis's applications are far more extended, as many concepts of the theory of evidence are inherently connected to the structure of the underlying domains. For example, the notion of *support function* depends on the idea of refining and could quite likely be reformulated using the algebraic language. Its analysis in the light of the lattice structure of \mathcal{F} can lead eventually to an alternative solution of the *canonical decomposition* problem.

For a more technical point of view, other properties of the frame lattices still need to be investigated: the presence and nature of complements and semi-complements, intervals and the relations of filters and ideals with finite and general families of frames.

Acknowledgements

This work has been partially supported by the Italian Space Agency (ASI) at the Autonomous Navigation and Computer Vision Laboratory of the University of Padova, Italy, led by Ruggero Frezza.

References

- [1] A. Beutelspacher and U. Rosenbaum, *Projective Geometry* (Cambridge University Press, Cambridge, 1998).
- [2] G. Birkhoff, Abstract linear dependence and lattices, *American Journal of Mathematics* 57 (1935) 800–804.
- [3] G. Birkhoff, *Lattice Theory* (3rd ed.) Vol. 25 (Amer. Math. Soc. Colloquium Publications, Providence, Rhode Island, 1967).
- [4] F. Cuzzolin, Visions of a generalized probability theory. PhD dissertation, Università di Padova, Dipartimento di Elettronica e Informatica (2001).
- [5] F. Cuzzolin and R. Frezza, An evidential reasoning frame-work for object tracking, in: *SPIE – Photonics East 99*, Vol. 3840 (19–22 September 1999) pp. 13–24.
- [6] F. Cuzzolin and R. Frezza, Integrating feature spaces for object tracking, in: *Proc. of MTNS2000* (21–25 June 2000).
- [7] A. Dempster, Upper and lower probabilities generated by a random closed interval, *Annals of Mathematical Statistics* 39 (1968) 957–966.
- [8] A. Dempster, Upper and lower probabilities inferences for families of hypothesis with monotone density ratios, *Annals of Mathematical Statistics* 40 (1969) 953–969.
- [9] A.P. Dempster, Upper and lower probability inferences based on a sample from a finite univariate population, *Biometrika* 54 (1967) 515–528.
- [10] M. Deutsch-McLeish, A study of probabilities and belief functions under conflicting evidence: Comparisons and new method, in: *Proceedings of IPMU'90* (Paris, France, 2–6 July 1990) pp. 41–49.
- [11] I.R. Goodman and H.T. Nguyen, *Uncertainty Models for Knowledge-Based Systems* (North Holland, New York, 1985).
- [12] J. Goutsias, R.P. Mahler and H.T. Nguyen, Random sets: theory and applications, in: *IMA Volumes in Mathematics and Its Applications, Vol. 97* (Springer, Berlin Heidelberg New York, December 1997).

- [13] N. Jacobson, *Basic Algebra I* (Freeman, New York, 1985).
- [14] J. Kohlas and Monney P.-A., *A Mathematical Theory of Hints – An Approach to the Dempster–Shafer Theory of Evidence*. Lecture Notes in Economics and Mathematical Systems, (Springer, 1995).
- [15] G. Matheron, *Random Sets and Integral Geometry*, Wiley Series in Probability and Mathematical Statistics.
- [16] C.K. Murphy, Combining belief functions when evidence conflicts, *Decision Support Systems* 29 (2000) 1–9.
- [17] H. Nguyen, On random sets and belief functions, *J. Mathematical Analysis and Applications* 65 (1978) 531–542.
- [18] H. Nguyen and T. Wang, Belief functions and random sets, in: *Applications and Theory of Random Sets, The IMA Volumes in Mathematics and its Applications*, Vol. 97 (Springer, 1997) pp. 243–255.
- [19] J.G. Oxley, *Matroid Theory* (Oxford University Press, Great Clarendon Street, Oxford, UK, 1992).
- [20] K.I. Rosenthal, *Quantales and Their Applications* (Longman scientific and technical, Longman house, Burnt Mill, Harlow, Essex, UK, 1990).
- [21] G. Shafer, *A Mathematical Theory of Evidence* (Princeton University Press, 1976).
- [22] G. Shafer, P.P. Shenoy and K. Mellouli, Propagating belief functions in qualitative Markov trees, *International Journal of Approximate Reasoning* 1(4) (1987) 349–400.
- [23] R. Sikorski, *Boolean Algebras* (Springer, Berlin Heidelberg New York, 1964).
- [24] P. Smets, The transferable belief model and random sets, *International Journal of Intelligent Systems* 7 (1992) 37–46.
- [25] M. Stern, *Semimodular Lattices* (Cambridge University Press, 1999).
- [26] G. Szasz, *Introduction to Lattice Theory* (Academic, New York, 1963).
- [27] H. Whitney, On the abstract properties of linear dependence, *American Journal of Mathematics* 57 (1935) 509–533.