

Mediterranean Marine Science

Vol 21, No 3 (2020)

Vol 21, n3



Alien and cryptogenic Foraminifera in the Mediterranean Sea: A revision of taxa as part of the EU 2020 Marine Strategy Framework Directive

RUTA STULPINAITE, ORIT HYAMS-KAPHZAN,
MARTIN R. LANGER

doi: [10.12681/mms.24673](https://doi.org/10.12681/mms.24673)

To cite this article:

STULPINAITE, R., HYAMS-KAPHZAN, O., & LANGER, M. R. (2020). Alien and cryptogenic Foraminifera in the Mediterranean Sea: A revision of taxa as part of the EU 2020 Marine Strategy Framework Directive. *Mediterranean Marine Science*, 21(3), 719–758. <https://doi.org/10.12681/mms.24673>

Alien and cryptogenic Foraminifera in the Mediterranean Sea: A revision of taxa as part of the EU 2020 Marine Strategy Framework Directive

Ruta STULPINAITE¹, Orit HYAMS-KAPHZAN² and Martin R. LANGER¹

¹Institut für Geowissenschaften, Paläontologie, Nussallee 8, 53115 Bonn, Germany

²Geological Survey of Israel, 32 Yeshu'ayahu Leibowitz, Jerusalem 9692100, Israel

Corresponding author: martin.langer@uni-bonn.de

Handling Editor: Argyro ZENETOS

Received: 11 September 2020; Accepted: 19 October 2020; Published online: 20 November 2020

Abstract

The human-mediated translocation of marine alien species beyond their natural ranges started as early as people began navigating the sea and is of growing concern to nature conservation. The Mediterranean Sea is among the most severely affected areas by biological invasions, a phenomenon that has been fostered by the opening and recent extension of the Suez Canal, the transport and release of ballast water, aquaculture and aquarium trade, ichthyochory and other active or passive dispersal mechanisms. The increase of marine invasions has stimulated considerable research, but for some important groups, in particular microorganisms, data are still limited. In this paper we have reviewed the current status of marine alien foraminifera in the Mediterranean Sea. Our survey includes a comprehensive taxonomic revision of previously recognized alien taxa, and new information obtained from the fossil record and from molecular studies. Our survey and reexamination of alien benthic foraminifera yielded a total of 44 validly recognized species and two species of cryptogenic taxa and reduces the number of previous recordings. The revised list includes both larger symbiont-bearing and smaller benthic foraminifera, including 16 hyaline-perforate, 3 agglutinated and 25 porcelaneous taxa. The vast majority of alien foraminifera recorded so far have become established in the Eastern and Central Mediterranean Sea, indicative for translocation and introduction via the Suez Canal pathway. Only one species, *Amphistegina lobifera*, causes significant ecological impacts and fulfills the criteria to be considered as an invasive alien. This species is a prolific carbonate producer, and displays extreme forms of ecosystem invasibility with capabilities to reduce native diversity and species richness. The proliferation and rates of recently observed range extensions, track contemporary sea surface temperature increases, provide strong support for previous species distribution models, and corroborate findings that rising water temperatures, global climate change and the extension of climate belts are major drivers fueling the latitudinal range expansion of larger symbiont-bearing and smaller epiphytic foraminifera. Intensified efforts to study alien foraminifera on a molecular level, in dated cores and in ballast water are required to trace their source of origin, to identify vectors of introduction and to verify their status as true aliens.

Keywords: Alien foraminifera; invasive species; taxonomy; biogeography; Mediterranean Sea.

Introduction

Alien or non-indigenous species (NIS), as defined in the European *Union's* Regulation 1143/2014 (EU, 2014a), are organisms introduced outside their natural range and dispersal potential. While most of them have little impact and are not a cause of concern, others can become invasive and pose a serious threat to biodiversity, ecosystem services, and coastal economies. More than 1400 marine species are currently recognized to be alien, cryptogenic or questionable to European waters, among which the Mediterranean Sea is the most impacted (Tsimis *et al.*, 2019, 2020; Galil *et al.*, 2015; Zenetos & Galanidi, 2020; Zenetos, 2017). The number of marine alien species introductions continues to grow as global trade,

shipping, fishing, travel, commercial mariculture, and aquarium trade allow species to be transported over large distances and to reach regions outside their native range not previously accessible to them.

The transfer of species started as early as shipping trade, and ballast water has been recognized as a major vector for the translocation of aquatic species across biogeographic boundaries. A substantial number of alien species are also translocated as hitchhikers by using floating objects for attachment and dispersal [e.g. ship hulls, marine plastic litter, trash, mollusk veliger shells (Gollasch, 2002; Nessbit, 2005; Rech *et al.*, 2016)]. Today, more than 80% of global trade is transported by commercial shipping (UNCTAD, 2019) and maritime transport will remain the most important transport mode in the

foreseeable future. International shipping is responsible for transporting and discharging ~ 3-12 billion tons of ballast water and 100 million tons of sediment each year (UNCTAD, 2019), and Cohen (1998) estimated that up to 10,000 aquatic alien species are transported in ships around the world per day. The rise of alien species around the globe is a consequence of an increasingly connected world and the result of human population growth (Pyšek *et al.*, 2020).

Biological invasions are now recognized as a major driver of biodiversity change across the globe (Vitousek *et al.*, 1996; Ojaveer *et al.*, 2015). The capacity to mitigate potential risks posed by alien species depends on precise, up-to-date and easily accessible information. To increase awareness and to facilitate effective prevention and management activities, comprehensive databases were established through large-scale international collaborations and advances in analytical bioinformatic tools. Comprehensive accounts are now available for plants, bryophytes, terrestrial snails, ants, spiders, amphibians, birds and mammals (Pyšek *et al.*, 2020). For some important groups of organisms, in particular microorganisms, data are still limited. Here we provide a comprehensive review of alien and invasive benthic foraminifera from the Mediterranean Sea. Foraminifera are a highly diverse group of single-celled eukaryotes, characterized by granular pseudopodia and a test that can be organic, agglutinated or calcareous. They are among the most abundant testate protists in the world's ocean, have an excellent fossil record and their shells represent a major and globally significant sink for calcium carbonate (Langer, 2008). To assess the current state, we carefully reviewed the presence of NIS benthic foraminifera in the Mediterranean Sea.

The Mediterranean is a semi-enclosed sea at the crossroads between Europe, Africa and Asia and a global hotspot of marine alien species (Costello *et al.*, 2010; Molnar *et al.*; 2008; Edelist *et al.*, 2013; Katsanevakis *et al.*, 2014a). During the 10th meeting of the Conference of the Parties at Nagoya in 2010, a New Strategic Plan for Biodiversity 2011-2020 was adopted in the Convention on Biological Diversity (CBD, 2012, 2014). This included *Aichi Target 9* on alien species, with the requirement that “*By 2020, invasive alien species and pathways are identified and prioritized, priority species are controlled and eradicated and measures are in place to manage pathways to prevent their introduction and establishment*”. The Marine Strategy Framework Directive (MSFD) of the European Union (Directive 2008/56/EC), initiated to achieve Good Environmental Status (GES) in the EU's marine waters by 2020, included the assessment of Non-Indigenous Species (NIS) as key Descriptor D2 (EU, 2008). In 2014, the EU Regulation 1143/2014 on Invasive Alien Species (IAS) has been implemented to oversee the detection, main pathways and early control or eradication until 2020 (EU, 2014a). To satisfy the need and in partial fulfillment to achieve the goals set by the EU Directive and Aichi Target 9, this survey provides a comprehensive reexamination, status-report and overview of NIS benthic foraminifera morphospecies and molecular types currently recognized in Mediterranean waters.

Material and Methods

The European Alien Species Information Network (EASIN) was launched with the aim to enable easy access to data and information on alien species in Europe and to assist scientists and policy makers in their efforts to prevent, control and manage alien invasions (Katsanevakis *et al.*, 2012). EASIN is an initiative of the Joint Research Centre (JRC) of the European Commission and has been established as a platform upon the recognition of the increasing serious threat posed by alien species in Europe. The EASIN information system facilitates the implementation of the EU Regulation 1143/2014 on Invasive Alien Species (EU, 2014a), and entered into force to fulfill Action 16 of Target 5 of the EU 2020 Biodiversity Strategy, as well as Aichi Target 9 of the Strategic Plan for Biodiversity 2011-2020 under the Convention of Biological Diversity.

EASIN (2020) currently contains the most complete register of foraminifera considered to be alien to the Mediterranean Sea. Primary data inserted into the EASIN database originate from published resources and contain the scientific name of the species, coordinates where the species was found, notes on the location, the date of observation, and a reference to a scientific publication/data source to allow proper citation and to link data to its original source. To respect high standards, quality assurance of the EASIN catalogue is provided by the EASIN Editorial Board (EB), a group of taxonomic experts responsible for specific groups of species. In an effort to update the existing account on invasive alien foraminifera in the Mediterranean Sea, we have conducted a large-scale survey on the existing literature. This includes a careful revision of the current taxonomic status of species, records and observations from new core material, and recent findings from molecular genetic analysis. The bulk of data have been carefully extracted from literature records published by generations of micropaleontologists. Literature data have been carefully surveyed and with a few exceptions include only adequately illustrated studies to assure taxonomic consistency. Because species-level taxonomy may vary from author to author and affects biogeography, all literature records have been critically reevaluated by the authors. The uniformity of nomenclature thus provides a framework of recognizable species that minimizes human-created taxonomic and identification weaknesses.

According to the Convention on Biological Diversity (CBD, 2012), alien species “*refers to a ‘species, subspecies or lower taxon, introduced outside its natural past or present distribution; including any part, gametes, seeds, eggs, or propagules of such species that might survive and subsequently reproduce*”. Because foraminifera are extremely abundant and provide an excellent fossil record, past occurrences can be traced from dated fossil material, a feature potentially masked by recent invasions through the opening of the Suez Canal in 1869, an artificial waterway connecting the Indian Ocean and the Mediterranean Sea. Thus, our survey includes fossil records from dated cores (0.3-120 m) using OSL, ¹⁴C and ²¹⁰Pb (Toueg, 1996; Avital, 2002; Tapiero, 2002; Tapiero *et al.*,

2003; Avnaim-Katav, 2010; Hyams-Kaphzan, unpublished data) with a focus on sites from the shallow Mediterranean coast of Israel: Achziv, Haifa, Dor, Beit-Yanai, Caesarea, Herzliya, Palmachim and Ashkelon. Benthic foraminifera obtained from material of other cores in the Mediterranean Sea apart from Israel (e.g. Koukousioura *et al.*, 2012; Milker & Schmiedl, 2012; Melis *et al.*, 2015) were also examined to verify the presence and identity of species pre-dating the opening of the Suez Canal in 1869. In the following we discuss the foraminifera species listed in the EASIN database together with new additions recently reported in the literature. For each species we provide the first record and locality, original citation, references, a synonymy, occurrence records in Mediterranean subregions, and discuss their status classified as alien, invasive, native, cryptogenic, or absent in respect to

previously published studies on alien foraminifera (Table 1, Table 2, Fig. 1). Potential pathways for the introduction (Table 1) are inferred from habitat preferences, live modes (epiphytic, epifauna, infauna), field and laboratory studies (Finger 2017; Langer, 1993; Jorissen, 1987; Hyams *et al.*, 2002; Alve & Goldstein, 2003; Langer *et al.*, 2009; Guy-Haim *et al.*, 2017; Weinmann *et al.*, 2019; McGann *et al.* 2020). Additional information is provided when a species was recorded in fossil or core material and when new molecular genetic data were available. The taxonomy adopted here generally follows Cimerman & Langer (1991), Hottinger *et al.* (1993), and Langer & Schmidt-Sinns (2006), with additional modern revisions provided by Parker (2009), Förderer & Langer (2018, 2019), WORMS (2020), and Fajemila *et al.* (2020).




Fig. 1: Mediterranean marine subregions according to the MSFD (Directive 2008/56/EC) as delineated by Article 4 of the MSFD (Jensen *et al.*, 2017; modified after Servello *et al.*, 2019). The border between ADRIA and CMED is defined by a line that runs from Capo Santa Maria di Leuca (Italy) (39.8° N, 18.36666° E) to the west coast of Corfu (Greece) (39.75194° N, 19.62777° E); the border between CMED and WMED is defined by a line that joins Capo Bon (37.08333° N, 11.05° E) (Tunisia) with Capo Lilibeo (37.8° N, 12.43333° E) (Sicily, Italy); between Sicily and the mainland of Italy, the border of WMED is a line that connects Capo Peloro (North-East Sicily, Italy) (38.26666° N, 15.65° E) to Capo Paci (Calabria, Italy) (38.25° N, 15.7° E) on the mainland. The border between the Central Mediterranean Sea and Eastern Mediterranean Sea (=Aegean-Levantine Sea) subregions is based on the reporting of Greece of its marine waters in 2012 under MSFD Article 8, 9 and 1020 and runs from the Greek mainland to the Island of Kithira (Greece), to the western coast of Crete and to Ras Al Hilal (Libya).

Table 1. Alphabetical list of currently recognized alien foraminifera in the Mediterranean Sea.

Abbreviations: Year=Year of first detection in the Mediterranean Sea as reported for the collection of the material in the publication; REF=References for the first year of detection in the Mediterranean Sea; NDR= Native distribution (Oceanic region from where the holotype was described); RS=Red Sea, IO=Indian Ocean, IP=Indo-Pacific, PTW=Pacific Tropical West, PTE=Pacific Tropical East, PT=Pacific Tropical, PO=Pacific Ocean, PW=Pacific West, PN=Pacific North, PS=Pacific South, PSE=Pacific Southeast, PSW=Pacific Southwest, PNE=Pacific Northeast, PNW=Pacific Northwest, ASE=Atlantic Southeast, ASW=Atlantic Southwest, ANE=Atlantic Northeast, ANW=Atlantic Northwest, ATW=Atlantic Tropical West, ATE=Atlantic Tropical East, AT=Atlantic Tropical, AO=Atlantic Ocean, AW=Atlantic West, AN=Atlantic North, ArO=Arctic Ocean, CoC=Cosmopolitan Circumtropical, CoT=Cosmopolitan Temperate, UNK=Unknown; Success=population success, CAS=casual, EST=established, Inv=invasive, UNK=unkown; PP=Potential Pathway; COR=Transport via the Suez Canal Corridor; TC/AN=Transport Contaminant: Contaminant on animals (species transported by host/vector); TS=TRANSPORT STOWAWAY: (Ship/boat ballast water, Ship/boat hull fouling); TS/AFS=TRANSPORT STOWAWAY: Attached to floating substrates (wood, seagrass leaves, rhizomes, algae) including plastic waste; Occurrence records of alien foraminifera in MSFD subregions of the Mediterranean Sea (for details see Servello *et al.*, 2019): MSFD subregions: the Adriatic (ADRIA), the central Mediterranean (CMED), the western Mediterranean (WMED), and the eastern Mediterranean (EMED).

<i>Species Name</i>	Year	REF (* not illustrated or unpublished record)	NDR	MSFD subregions	Success	PP
<i>Agglutinella compressa</i> El-Nakhal, 1983	1998	Hyams-Kaphzan <i>et al.</i> , 2008	RS	EMED	CAS	COR, TS
<i>Amphistegina lessonii</i> d'Orbigny, 1826	1974	Hollaus & Hottinger, 1997	IO	EMED, CMED, WMED	EST	COR, TS, TC/AN, TS/AFS
<i>Amphistegina lobifera</i> Larsen, 1976	1955-1964	Blanc-Vernet, 1969	RS	EMED, ADRIA, CMED, WMED	INV	COR, TS, TC/AN, TS/AFS
<i>Amphistegina</i> cf. <i>A. papillosa</i> Said, 1949	2005	Caruso & Cosentino, 2014	RS?/IO?	CMED, WMED	EST	COR, TS, TC/AN, TS/AFS
<i>Borelis schlumbergeri</i> (Reichel, 1937)	1999	Hyams (2000)	IO	EMED	EST	COR, TS, TS/AFS
<i>Brizalina simpsoni</i> (Heron-Allen & Earland, 1915)	2007	Meriç <i>et al.</i> , 2010*, Meriç <i>et al.</i> , 2014	IO	EMED	CAS	COR, TS, TS/AFS
<i>Cibicides mabahethi</i> Said, 1949	1998	Hyams (2000)	RS	EMED	CAS	COR, TS, TS/AFS
<i>Clavulina</i> cf. <i>C. multicamerata</i> Chapman, 1907	1972-1974	Blanc-Vernet <i>et al.</i> , 1979	PSW/IO	EMED, CMED	EST	COR, TS
<i>Cycloforina</i> sp.	2008	Langer, 2008	RS	EMED	CAS	COR, TS, TS/AFS
<i>Cyclorbiculina compressa</i> (d'Orbigny, 1839)	2002	Meriç <i>et al.</i> , 2008a	ATW	EMED	EST	TS, TC/AN, TS/AFS
<i>Elphidium striatopunctatum</i> (Fichtel & Moll, 1798)	1998	Hyams, 2000	RS	EMED	EST	COR, TS, TS/AFS
<i>Epistomaroides punctulata</i> (Said, 1949)	2005	Almogi-Labin & Hyams-Kaphzan, 2012	RS	EMED	EST	COR, TS, TS/AFS
<i>Haddonina</i> sp.	1996	Hyams-Kaphzan, unpubl.	RS	EMED	CAS	COR, TS
<i>Hauerina diversa</i> Cushman, 1946	1996	Hyams, 2000	PT	EMED	EST	COR, TS, TS/AFS
<i>Heterostegina depressa</i> d'Orbigny, 1826	1965	Moncharmont Zei, 1968	ASE	EMED	EST	COR, TS, TS/AFS
<i>Loxostomina</i> cf. <i>L. africana</i> (Smitter, 1955)	1972-74	Alavi, 1980	IO	EMED	EST	COR, TS, TS/AFS
<i>Mimosina affinis</i> Millett, 1900	2012	Mouanga, 2017	PTW	ADRIA	CAS	COR, TS, TS/AFS
<i>Neoconorbina clara</i> (Cushman, 1934)	1997	Hyams-Kaphzan <i>et al.</i> , 2008	PTW	EMED	CAS	COR, TS, TS/AFS

continued

Table 1 continued

<i>Species Name</i>	Year	REF (* not illustrated or unpublished record)	NDR	MSFD subregions	Success	PP
<i>Nodophthalmidium antillarum</i> (Cushman, 1922)	1968	Moncharmont Zei, 1968	ATW	EMED	EST	COR, TS, TS/AFS
<i>Operculina ammonoides</i> (Gronovius, 1781)	2015	Merkado, 2016	IO	EMED	EST	COR, TS, TS/AFS
<i>Pararotalia</i> cf. <i>P. socorroensis</i> (McCulloch, 1977)	1997	Hyams-Kaphzan <i>et al.</i> , 2008	PTE	EMED	CAS	COR, TS, TS/AFS
<i>Parasorites orbitolitoides</i> (Hofker, 1930)	2016	Nadal Nebot, 2017	PTW	WMED	CAS	COR, TS, TS/AFS
<i>Paratrochammina madeirae</i> Brönnimann, 1979	1997	Hyams, 2000	ASW	EMED	CAS	TS, TS/AFS
<i>Pegidia lacunata</i> McCulloch, 1977	1994	Langer, 2008	PTW	EMED	CAS	COR, TS, TS/AFS
<i>Planispirinella exigua</i> (Brady, 1879)	1988-1991	Oflaz, 2006	PTW	EMED, ADRIA?	EST	COR, TS, TS/AFS
<i>Planogypsina acervalis</i> (Brady, 1884)	1909	Sidebottom, 1909	PTW	EMED, ADRIA?	EST	COR, TS, TS/AFS
<i>Procerolagena oceanica</i> (Albani, 1974)	2003	Hyams, 2006	PW	EMED	CAS	COR, TS, TS/AFS
<i>Pseudohauerinella dissidens</i> (McCulloch, 1977)	1998	Hyams, 2000	PTE	EMED	CAS	COR, TS, TS/AFS
<i>Pseudomassilina australis</i> (Cushman, 1932)	1988-1991	Oflaz, 2006	PT	EMED	CAS	COR, TS, TS/AFS
<i>Pseudomassilina reticulata</i> (Heron-Allen & Earland, 1915)	1998	Hyams, 2000	IO	EMED	EST	COR, TS, TS/AFS
<i>Pyrgo denticulata</i> (Brady, 1884)	1994, 1996	Hyams, 2000, Samir <i>et al.</i> , 2003	PTE	EMED	EST	COR, TS, TS/AFS
<i>Quinqueloculina</i> cf. <i>Q. mosharrafai</i> Said, 1949	2002	Meriç <i>et al.</i> , 2008a	RS	EMED	EST	COR, TS, TS/AFS
<i>Quinqueloculina</i> cf. <i>Q. multimarginata</i> Said, 1949	1996	Hyams, 2000	RS	EMED	CAS	COR, TS, TS/AFS
<i>Schlumbergerina alveoliniformis</i> (Brady, 1879)	2008	Meriç <i>et al.</i> , 2008b	PTW	EMED	EST	COR, TS
<i>Sigmamiliolinella australis</i> (Parr, 1932)	1977	Alavi, 1980	IO	EMED, CMED, ADRIA	EST	COR, TS, TS/AFS
<i>Siphonaperta distorquata</i> (Cushman, 1954)	1977	Alavi, 1980	PTW	EMED	EST	COR, TS, TS/AFS
<i>Sorites variabilis</i> Lacroix, 1941	1996	Langer (unpublished)	RS	EMED, WMED	EST	COR, TS, TS/AFS
<i>Spiroloculina angulata</i> Cushman, 1917	1968	Moncharmont Zei, 1968	PN	EMED	EST	COR, TS, TS/AFS
<i>Spiroloculina antillarum</i> d'Orbigny, 1826	1913	Wiesner, 1913	PN	CMED, ADRIA	EST	COR, TS, TS/AFS
<i>Spiroloculina attenuata</i> Cushman & Todd, 1944	1997	Hyams, 2000	PTW	EMED	CAS	COR, TS, TS/AFS
<i>Spiroloculina</i> aff. <i>S. communis</i> Cushman & Todd, 1944	1996	Hyams-Kaphzan <i>et al.</i> , 2008	PTW	EMED	EST	COR, TS, TS/AFS
<i>Spiroloculina nummiformis</i> Said, 1949	1997	Hyams, 2000	RS	EMED	CAS	COR, TS, TS/AFS
<i>Triloculina</i> cf. <i>T. fichteliana</i> d'Orbigny, 1839	2002	Meriç <i>et al.</i> , 2008a	ATW	EMED, CMED	EST	TS, TS/AFS
<i>Varidentella</i> cf. <i>V. neostriatula</i> (Thalman, 1950)	1996	Hyams, 2000	PTW	EMED	CAS	COR, TS, TS/AFS

Table 2. Alphabetical list of currently recognized cryptogenic foraminifera in the Mediterranean Sea (for abbreviations see captions in in Table 1).

<i>Species Name</i>	Year	REF (* not illustrated)	NDR	MSFD subregions	Suc- cess	PP
<i>Cymbaloporetta</i> sp. 1	1968	Moncharmont Zei, 1968	PTE	EMED, CMED, ADRIA, WMED	CAS	ST/SH- BAL, ST/AFS, UNAI
<i>Euthymonacha polita</i> (Chapman, 1900)	2007	Meriç <i>et al.</i> , 2010	PW	EMED, ADRIA, CMED	EST	ST/SH- BAL, ST/AFS, UNAI

Results

Acervulina inhaerens Schulze, 1854

- 1854 *Acervulina inhaerens* Schulze – Schulze, p. 68, pl. 6, fig. 12.
 2015 *Acervulina inhaerens* Schulze – Delliou *et al.*, p. 570, pl. 1, fig. 5.

Acervulina inhaerens was first described by Schulze (1854) from the Adriatic Sea and found to be alive on phytal substrates, on sand and along rocky shores off Ancona (Italy). The species was reported as alien by Delliou *et al.* (2015) from coastal habitats in the Gulf of Kavala (NE Aegea Sea, Greece). The record by Schulze (1854) shows the species to be present in the Mediterranean prior to the opening of the Suez Canal and the taxon must therefore be considered to be native.

Adelosina carinata-striata (Wiesner, 1912)

- 1912 *Miliolina milletti* var. *carinata-striata* Wiesner – Wiesner, p. 221
 1923 *Adelosina milletti* var. *carinata-striata* Wiesner – Wiesner, p. 77, figs. 190-191
 1970 *Quinqueloculina milletti* var. *carinata-striata* (Wiesner) – von Daniels, p. 74, pl. 2, figs. 17a-c, textfig. 49
 1987 *Quinqueloculina poeyana carinata* Albani – Baccaert, p. 97, pl. 47, figs. 5a-c
 1991 *Adelosina carinata-striata* (Wiesner) – Cimerman & Langer, p. 28, pl. 20, figs. 1-4
 1993 *Cycloforina? carinata* (Albani) – Hottinger *et al.*, p. 49, pl. 32, figs. 1-9
 2005 *Adelosina carinata striata* (Wiesner) – Debenay *et al.*, pl. 1, figs. 14-15
 2007 *Quinqueloculina carinatastriata* (Wiesner) – Bouchet *et al.*, pl. 1
 2012 *Adelosina carinatastriata* (Wiesner) – Koukousioura *et al.*, pl. 10, fig. 18
 2013 *Adelosina carinata-striata* (Wiesner) – Bassler-Veit *et al.*, pl. 5, figs. 1a-b
 2014 *Adelosina carinata-striata* (Wiesner) – Meriç *et al.*, pl. 9, figs. 8-9
 2017 *Adelosina carinata-striata* (Wiesner) – Mouanga, pl. 3, fig. 11

Adelosina carinata-striata was first described by Wiesner (1912) from the Adriatic Sea. The species is present off the European Atlantic coast (Bouchet *et al.*, 2007) and in the Red Sea (Hottinger *et al.*, 1993). Illustrated Mediterranean Sea records include specimens from the Adriatic Sea (Wiesner, 1923; von Daniels, 1970; Cimerman & Langer, 1991; Mouanga, 2017), Greece (Debenay *et al.*, 2005) and Turkey (Bassler-Veit, 2013; Meriç *et al.*, 2014). Subfossil specimens from the Aegean Sea were recorded by Koukousioura *et al.* (2012). *Adelosina carinata-striata* is present in the lower part of a borehole from Alykes Kitros, and the illustrated specimen was dated at 7460±40yBP. The species *Adelosina carinata-striata* is therefore native to the Mediterranean Sea. The specimen recorded by Triantaphyllou *et al.* (2005) under the name *Adelosina milletti* var. *carinata* differs from *Adelosina carinata-striata* (Wiesner) and belongs to a different species.

Adelosina milletti (Wiesner, 1911)

- non 1839 *Quinqueloculina bosciiana* – d’Orbigny, p. 191, pl. 11, figs. 22-24
 1898 *Miliolina bosciiana* d’Orbigny – Millett, p. 267, pl. 6, fig. 1
 1911b *Miliolina milletti* Wiesner – Wiesner, p. 506
 1912 *Miliolina milletti* Wiesner – Wiesner, p. 220
 1923 *Adelosina milletti* Wiesner – Wiesner, p. 76
 non 1980 *Quinqueloculina milletti* (Wiesner) – Alavi, p. 20 (appendix), pl. 17, fig. 5, pl. 18, fig. 6
 non 1990 *Massilina milletti* (Wiesner) – Dermitzakis & Triantafyllou, pl. 2, fig. 8
 non 1993 *Quinqueloculina milletti* (Wiesner) – Sgarrella & Moncharmont Zei, p. 172, pl. 7, fig. 7
 non 2000 *Quinqueloculina milletti* (Wiesner) – Hyams, pl. 7, fig. 8
 non 2006 *Edentostomina milletti* (Wiesner) – Oflaz, p. 145, pl. 2, figs. 4-6
 non 2016 *Quinqueloculina milletti* (Wiesner) – Hyams-Kaphzan, pl. 2, fig. 10
 non 2016a *Quinqueloculina milletti* (Wiesner) – Dimiza *et al.*, pl. 2, fig. 2

Adelosina milletti was first reported by Wiesner (1911b, as *Miliolina milletti*) from the Adriatic Sea (not illustrated) and a morphological description was provided by Wiesner in 1912. The author (1923) considered the

species to be identical with *Miliolina bosciiana* of Millett (1898) from the Malay Archipelago and to represent the type species. However, Wiesner (1923) pointed out that *Miliolina bosciiana* Millett is not present in the Adriatic Sea and that the species is characterized by rounded chambers, a smooth surface, a short, curved neck and therefore differs from *Miliolina bosciiana* d'Orbigny (1839). The species was illustrated by Wiesner in his monograph on miliolids from the eastern Adriatic Sea (1923). Mediterranean records by Alavi (1980), Dermitzakis & Triantafyllou (1990), Oflaz (2006) and Dimiza *et al.* (2016a) provide illustrated specimens but they all differ from the original of Millett (1898). The specimens of *Quinqueloculina milletti* (see Sgarrella & Moncharmont Zei, 1993; Hyams, 2000; Hyams-Kaphzan, 2016), a species also considered to be alien to the Mediterranean in EASIN (2020), also differ substantially from Millett's *Miliolina bosciiana* and require further study. To date, there is no reliable record of *Adelosina milletti* in the Mediterranean Sea, and we therefore consider the species to be absent from Mediterranean waters.

***Agglutinella compressa* El-Nakhal, 1983**

- 1983 *Agglutinella compressa* El-Nakhal – El-Nakhal, p. 129, pl. 1, figs. 1-3, pl. 2, figs. 10-11
- 1993 *Agglutinella compressa* El-Nakhal – Hottinger *et al.*, p. 48, pl. 29, figs. 5-10
- non 2003 *Agglutinella compressa* El-Nakhal – Samir *et al.*, pl. 1, figs. 5, 9-10
- non 2012 *Agglutinella compressa* El-Nakhal – Aloulou *et al.*, pl. 1, fig. 5

Agglutinella compressa was first described by El-Nakhal (1983) from the Gulf of Suez. The specimens illustrated are identical with the specimens reported by Hottinger *et al.* (1993) from the Gulf of Aqaba. Rare specimens reported from the Mediterranean Sea (Samir *et al.*, 2003; Aloulou *et al.*, 2012) have more inflated chambers, are less elongate and compressed and have an elliptical aperture. The species was collected off the Mediterranean coast of Israel between the years 1998-1999 (Hyams-Kaphzan *et al.*, 2008). The data currently available indicate that *Agglutinella compressa* is alien to the Mediterranean Sea.

***Agglutinella robusta* El-Nakhal, 1983**

- 1983 *Agglutinella robusta* El-Nakhal – El-Nakhal, p. 130, pl. 1, figs. 4-6, pl. 2, figs. 12-15
- 1993 *Agglutinella robusta* El-Nakhal – Hottinger *et al.*, p. 48, pl. 29, figs. 11-14, pl. 2, figs. 12-15
- non 2003 *Agglutinella robusta* El-Nakhal – Samir *et al.*, pl. 1, figs. 6-8
- non 2019 *Agglutinella robusta* El-Nakhal – Badr-ElDin *et al.*, pl. 1, figs. 5a-c

The holotype of *Agglutinella robusta* was described and illustrated by El-Nakhal (1983) from the Red Sea. Mediterranean records of this species were provided by Samir *et al.* (2003), but their findings differ from the holotype in having a more elongated shape, less inflated chambers and a coarser agglutination pattern. Specimens illustrated by Badr-ElDin *et al.* (2019) differ substantially from the holotype. Yokeş & Meriç (2009) have reported the species to be present in the Mediterranean Sea, but do not provide an illustration. We therefore consider *Agglutinella robusta* to be absent from the Mediterranean Sea.

***Amphisorus hemprichii* Ehrenberg, 1839**

- 1839 *Amphisorus hemprichii* Ehrenberg – Ehrenberg, p. 130, pl. 3, fig. 3
- 1993 *Amphisorus hemprichii* Ehrenberg – Hottinger *et al.*, p. 71, pl. 81, pl. 82
- 2013 *Amphisorus hemprichii* Ehrenberg – Merkado *et al.*, fig. 7.3
- 2014 *Amphisorus hemprichii* Ehrenberg – Meriç *et al.*, pl. 40, figs. 3-6, pl. 41, figs. 1-2
- 2015 *Amphisorus hemprichii* Ehrenberg – Corsini-Foka *et al.*, p. 356, fig. 2

The holotype of *Amphisorus hemprichii* has been described by Ehrenberg (1839) from modern sediments off Alexandria, Egypt. Present-day Mediterranean occurrences were reported by Meriç *et al.* (2014) from shallow water sites off Turkey. Living specimens reported as *Amphisorus hemprichii* by Gruber (2007) from Shikmona (Haifa, Israel) were reexamined by us and belong to *Sorites orbiculus*. Only a lateral view is illustrated by Corsini-Foka *et al.* (2015) for a specimen from Rhodes Island (Greece), but the species is described to possess two separate layers of chamberlets and a double row of apertures, and as such fulfills the criteria for *Amphisorus*. No living *A. hemprichii* are currently present along the Mediterranean coast off Israel (Merkado *et al.*, 2013; Hyams-Kaphzan, unpublished data). The lower temperature limit for symbiont-bearing species of *Amphisorus hemprichii* is at around 18°C in the Mediterranean Sea (Langer, unpubl. data). *Amphisorus hemprichii* is widely present in all oceans (Langer & Hottinger, 2000) and shows a circumtropical distribution. The species is native to the Mediterranean Sea.

***Amphistegina lessonii* d'Orbigny, 1826**

- 1826 *Amphistegina lessonii* – d'Orbigny, p. 304
- 1993 *Amphistegina lessonii* d'Orbigny – Hottinger *et al.*, p. 132, pl. 184, fig. 1-11; pl. 185, fig. 1-7
- 1997 *Amphistegina lessonii* d'Orbigny – Hollaus & Hottinger, p. 592
- 2000 *Amphistegina lessonii* d'Orbigny – Hyams, pl. 28, figs. 1-4
- 2002 *Amphistegina lessonii* d'Orbigny – Hyams *et al.*, p. 174, pl. 1, fig. 1
- 2006 *Amphistegina lessonii* d'Orbigny – Langer

- & Lipps, p. 349
- 2009 *Amphistegina lessonii* d'Orbigny – Parker, p. 498, fig. 355a-d
- 2011 *Amphistegina lessonii* d'Orbigny – Makled & Langer, p. 248, fig. 9: 30, 31
- 2015 *Amphistegina lessonii* d'Orbigny – Fajemila *et al.*, Moorea, fig. 2: 12
- 2016 *Amphistegina lessonii* d'Orbigny – Langer *et al.*, p. 76, pl. 2, fig. 28-31
- 2018 *Amphistegina lessonii* d'Orbigny – Förderer & Langer, p. 124, pl. 51, figs. 4-6
- 2019 *Amphistegina lessonii* d'Orbigny – Guastella *et al.*, p. 5, figs. 2.7-2.10
- 2020 *Amphistegina lessonii* d'Orbigny – Fajemila *et al.*, Moorea, p. 79, pl. 29, fig. 24-29.

Amphistegina lessonii has first been described by d'Orbigny (1826) from Mauritius (Indian Ocean). The species has a circumtropical distribution and has also been reported from the Pacific and the Atlantic Ocean, and the Red and the Mediterranean Sea (see Förderer and Langer, 2018 and Fajemila *et al.*, 2020 for biogeographic details). The biogeographic range of *A. lessonii* is strongly controlled by water temperature and in the Mediterranean Sea (Hollaus & Hottinger, 1997; material collected in 1974) the species has recently reached the coast of Sicily (Guastella *et al.*, 2019). *Amphistegina lessonii* has established self-maintaining populations, and reaches high abundances in some areas (e.g. Central and eastern Mediterranean; Triantaphyllou *et al.*, 2005; Guastella *et al.*, 2019; Hyams-Kaphzan, unpublished). Whether the species has indeed the capability to cause a negative impact on the diversity of native biotas or to modify local habitats requires further study. The species is considered to be alien to the Mediterranean Sea, but molecular genetic studies are needed to confirm that the species has been introduced from the Atlantic via the Strait of Gibraltar and/or the Red Sea via the Suez Canal.

Amphistegina lobifera Larsen, 1976

- 1968 *Amphistegina madagascariensis* d'Orbigny – Moncharmont Zei, p. 29, pl. 5, figs. 2a-b
- 1969 *Amphistegina madagascariensis* d'Orbigny – Blanc-Vernet, p. 214, pl. 10, 2, nr. 30
- 1976 *Amphistegina lobifera* Larsen – Larsen, p. 4, pl. 3, figs. 1-5, pl. 7, fig. 3, pl. 8, fig. 3
- 1980 *Amphistegina lobifera* Larsen – Alavi, p. 67 (appendix), pl. 34, fig. 3, pl. 35, figs. 1, 3
- 1993 *Amphistegina lobifera* Larsen – Hottinger *et al.*, p. 133, pl. 186, figs. 1-11, pl. 187, figs. 1-7, pl. 188, figs. 1-6
- 1998 *Amphistegina lobifera* Larsen – Yanko *et al.*, p. 193, pl. 3, figs. 14-16
- 2000 *Amphistegina lobifera* Larsen – Langer & Hottinger, p. 112
- 2000 *Amphistegina lobifera* Larsen – Hyams, pl. 28, figs. 5-6
- 2002 *Amphistegina lobifera* Larsen – Hyams *et al.*, p. 174, pl. 1, figs. 2-4

- 2005 *Amphistegina lessonii* d'Orbigny - Triantaphyllou *et al.*, p. 283, pl. 1, fig. 5
- 2008 *Amphistegina lobifera* Larsen – Langer, p. 404, fig. 3
- 2009 *Amphistegina lobifera* Larsen – Triantaphyllou *et al.*, p. 79, pl. 1, figs. 1-5
- 2010 *Amphistegina lobifera* Larsen – Koukousioura *et al.*, p. 159, pl. 1, figs. 1-2
- 2010 *Amphistegina lobifera* Larsen – Avnaim-Katav, p. 232, pl. 8, figs. 11-12
- 2013 *Amphistegina lobifera* Larsen – Triantaphyllou & Dimiza in Siokou *et al.*, p. 242, fig. 7
- 2016 *Amphistegina lobifera* Larsen – Langer & Mouanga, fig. 4
- 2020 *Amphistegina lobifera* Larsen – Manda *et al.*, pl. 3c

Amphistegina lobifera has first been described by Larsen (1976) from the Red Sea. Its biogeographic range is strongly controlled by water temperature and the distributional range is currently delimited by the 13.7°C winter isotherm (Zmiri *et al.*, 1974; Hollaus & Hottinger, 1997; Langer & Hottinger, 2000). The species is widely present in the eastern Mediterranean and the Ionian Sea (Blanc-Vernet, 1969; Langer, 2008; Guy-Haim *et al.*, 2017; Manda *et al.*, 2020; Triantaphyllou *et al.*, 2005; Triantaphyllou & Dimiza in Siokou *et al.*, 2013), has recently invaded the Adriatic Sea (Langer & Mouanga, 2016), the coast of Sicily (Guastella *et al.*, 2019) and is present along the coast of Libya (Langer unpubl. data) and Tunisia (Blanc-Vernet *et al.*, 1979; Glacon, 1962; Langer, 2008; El Kateb *et al.*, 2018; Damak *et al.*, 2019; Langer unpubl. data). The species is currently among the 26 most invasive taxa recorded in the Mediterranean (Servello *et al.*, 2019; Tsiamis *et al.*, 2020). Core material recovered from Haifa Bay (Israel) shows *Amphistegina lobifera* to be present at a depth of 1.95 m (Avnaim-Katav, 2010) but dating of this core is not clear cut and requires further studies. A sediment core date prior to the opening of the Suez Canal, would support previous hypothesis (Langer & Hottinger, 2000; Langer, 2008) that Mediterranean populations were restocked from the Atlantic via the Strait of Gibraltar.

Amphistegina cf. *A. papillosa* Said, 1949

- 2014 *Amphistegina* cf. *A. papillosa* Said – Caruso & Cosentino fig. 5, 8

Amphistegina cf. *A. papillosa*, an ecomorphotype of *Amphistegina papillosa* Said, was recently reported by Caruso & Cosentino (2014) from shallow waters off the Pelagian Islands, Central Mediterranean Sea. The ecomorphotype is similar to *A. papillosa*, which is commonly found attached to green algae in the Red Sea at water depth of 112 and 135 m (Hottinger *et al.*, 1993), and is one of the Amphisteginids that occupies the deepest waters (Murray, 2006 and reference therein). The ecomorphotype has not been formally described but was considered to be a putative exotic form (Caruso & Cosentino,

2014) and was listed as an alien species by Servello *et al.* (2019). The holotype of *A. papillosa* was described by Said (1949) from the Red Sea, but neither *A. papillosa* nor a related “eco”-morphotype has been reported prior the finding of Caruso & Cosentino (2014) from the Mediterranean Sea. *Amphistegina* cf. *A. papillosa* and *Amphistegina papillosa* Said are not known from the Mediterranean coast of Israel, and the occurrence at the Pelagian Islands is puzzling. For the time being, *Amphistegina* cf. *A. papillosa* is here considered to be alien to the Mediterranean Sea, but molecular genetic studies are needed to confirm that the species is identical with the Red Sea species.

Archaias angulatus (Fichtel & Moll, 1798)

- 1798 *Nautilus angulatus* Fichtel & Moll – Fichtel & Moll, p. 113, pl. 22, figs. a-e
 1928 *Archaias angulatus* Fichtel & Moll – Cushman, p. 220, pl. 31, fig. 9

The holotype of *Archaias angulatus* has been described by Fichtel & Moll (1798) from the Red Sea but was not found in present-day assemblages recorded by Said (1949) and Hottinger *et al.* (1993). *Archaias angulatus* has been recorded from the Cape Verde Islands (Rocha & Mateu, 1971). An obscure finding of a single, small, megalospheric test of *A. angulatus* was reported from around the Kornati Islands, Adriatic Sea (Langer & Hottinger, 2000). To date, this represents the only record in the Mediterranean Sea. Cushman (1931) and Smout & Eames (1958) consider the species to be absent from the Mediterranean Sea. Other Mediterranean occurrences of *A. angulatus* have never been reported and we therefore consider the species to be absent from the Mediterranean Sea.

Articulina alticostata Cushman, 1944

- 1944 *Articulina alticostata* Cushman – Cushman, p. 16, pl. 4, figs. 10-13
 non 2004 *Articulina alticostata* Cushman – Meriç *et al.*, p. 109, pl. 16, fig. 10
 non 2006 *Articulina alticostata* Cushman – Oflaz, p. 180, pl. 4, fig. 6
 non 2009 *Articulina alticostata* Cushman – Yokeş & Meriç, pl. 2, figs. 10-11

Meriç *et al.* (2004) consider *Articulina alticostata* to be present in the Mediterranean Sea but the initial part of the specimen illustrated is lacking. Therefore, the species can not be assigned with certainty to the genus *Articulina*. Specimens illustrated by Yokeş & Meriç (2009) show chambers that are less inflated than in the holotype and are similar to *Nodophthalmidium antillarum*. The specimen identified by Oflaz (2006) as *Articulina alticostata* is an aberrant juvenile variety of *Articulina pacifica* (for further information see remarks on *Articulina pacifica*). To date, convincing evidence for the presence of *Articulina alticostata* in the Mediterranean Sea is lacking and we therefore consider the species to be absent.

Articulina mayori Cushman, 1922

- 1922 *Articulina mayori* Cushman – Cushman, p. 71, pl. 13, fig. 5
 non 2006 *Articulina mayori* Cushman – Oflaz, pl. 4, fig. 9

Articulina mayori is a tropical species that was first described from the Tortugas (Florida) by Cushman (1922). The illustration provided by Oflaz (2006) has a very faint striation pattern and is not identical with the distinctly ornamented specimen described and illustrated by Cushman (1922). The specimen depicted by Oflaz (2006) may belong to the genus *Nodophthalmidium* but its true identification requires further study. For the time being, we consider *Articulina mayori* to be absent from the Mediterranean Sea.

Articulina pacifica Cushman, 1944

- 1904 *Articulina sulcata* Reuss – Sidebottom, p. 16, pl. 4, figs. 16-17
 1944 *Articulina pacifica* Cushman – Cushman, p. 17, pl. 4, figs. 14-18
 1993 *Articulina pacifica* Cushman – Hottinger *et al.*, p. 48, pl. 31, figs. 5-9
 2006 *Articulina alticostata* Cushman – Oflaz, p. 180, pl. 4, fig. 6

The holotype of *Articulina pacifica* was described by Cushman (1944) from Fiji. The species is present in the Red Sea (Hottinger *et al.*, 1993). Mediterranean records include the specimens illustrated by Sidebottom (1904, as *Articulina sulcata* Reuss), Oflaz (2006) and Hyams-Kaphzan *et al.* (2008). As the oldest record of *Articulina pacifica* was provided by Sidebottom (1904) from the island of Delos (Greece), and the species is considered to be native to the Mediterranean Sea.

Astacolus insolitus (Schwager, 1866)

- 1866 *Cristellaria insolita* Schwager – Schwager, p. 242, pl. 6, fig. 85
 1980 *Astacolus insolitus* (Schwager) – Srinivasan & Sharma, p. 28, pl. 5, figs. 12-13
 non 2009 *Astacolus insolitus* (Schwager) – Yokeş & Meriç, pl. 4, fig. 1
 non 2009 *Astacolus insolitus* (Schwager) – Mojtahid *et al.*, pl. 2, fig. 26
 non 2014 *Astacolus insolitus* (Schwager) – Meriç *et al.*, pl. 46, figs. 11a-b

Astacolus insolitus has been described from fossil deposits of Car Nicobar, north-eastern Indian Ocean (Schwager, 1866). A neotype has been illustrated by Srinivasan & Sharma (1980). Records from the Mediterranean Sea were illustrated by Yokeş & Meriç (2009), Mojtahid *et al.* (2009) and Meriç *et al.* (2014) but differ substantially from Schwager’s illustrations and the neotype and belong to a different species, most probably to *Astacolus*

crepidulus (Fichtel & Moll, 1798). To date, there is no record for *Astacolus insolitus* in the Mediterranean Sea and we therefore consider the species to be absent.

***Brizalina simpsoni* (Heron-Allen & Earland, 1915)**

- 1915 *Bolivina simpsoni* Heron-Allen & Earland – Heron-Allen & Earland, p. 648, pl. 49, figs. 18-35
- 1993 *Brizalina simpsoni* (Heron-Allen & Earland) – Hottinger *et al.*, p. 92, pl. 111, figs. 8-13, pl. 112, figs. 1-2
- 1997 *Brizalina simpsoni* (Heron-Allen & Earland) – Haunold *et al.*, fig. 5
- 2014 *Brizalina simpsoni* (Heron-Allen & Earland) – Meriç *et al.*, pl. 50 fig. 1
- 2017 *Brizalina simpsoni* (Heron-Allen & Earland) – Thissen & Langer, pl. 13, figs. 6-8

The holotype of *Brizalina simpsoni* has been described by Heron-Allen & Earland (1915) from the Quirimba Islands off northeastern Mozambique. The species has also been reported from the Red Sea (Hottinger *et al.*, 1993; Haunold *et al.*, 1997) and Zanzibar (Thissen & Langer, 2017). The first Mediterranean record was given by Meriç *et al.* (2010) from the Pamučak Cove (northwest Kuşadasi, Turkey). The species has later been illustrated by Meriç *et al.* (2014) from the Gulf of Kuşadasi, Aegean Sea. For the generic assignment, we follow the concept of Revets (1996). The available records show that the species is alien to the Mediterranean Sea.

***Borelis schlumbergeri* (Reichel, 1937)**

- 1973 *Neoalveolina pygmaea schlumbergeri* Reichel – Reichel, p. 110, pl. 10, figs. 1-3, pl. 11, fig. 6b
- 1993 *Borelis schlumbergeri* (Reichel) – Hottinger *et al.*, p. 68, pl. 75, figs. 1-17
- ? 2000 *Borelis* sp. – Hyams, pl. 14, figs. 9-10
- ? 2002 *Borelis* sp. – Hyams *et al.*, pl. 1, figs. 5-6
- 2008 *Borelis* sp. – Langer, fig. 5
- ? 2017 *Borelis* sp. – Makled *et al.*, pl. 13, figs. 19-20
- 2019 *Borelis schlumbergeri* (Reichel) – Bassi *et al.*, p. 14, figs. 12-13

The genus *Borelis* has a circumtropical distribution and is represented by two extant species (Langer & Hottinger, 2000): an elongate *Borelis schlumbergeri* and a spherical *Borelis pulchra*. Fossil occurrences of other species of *Borelis* spp. from the Mediterranean region were reported by Reiss & Gvirtzman (1966), Adams (1976), Jones *et al.* (2006), Makled *et al.* (2017) and Bassi *et al.* (2019). Unpublished records of modern *B. schlumbergeri* from northern Israel (Hyams-Kaphzan) show the species to be present in the year 1999 at Akhziv. Later, more *Borelis* were found at Akhziv by Lazar (2007), showing the typical elongate test shape (see also Langer, 2008). Specimens reported by Hyams (2000) and Hyams *et al.* (2002) represent subspherical forms that may represent juvenile stages of *Borelis schlumbergeri* or *Borelis pulchra*. Sub-spherical forms were also reported from fossil sediments in the Nile Delta (Makled *et al.*, 2017). For the time being, we consider the elongate *Borelis schlumbergeri* to be alien to the Mediterranean Sea.

Cibicides mabahethi Said, 1949

1949 *Cibicides mabahethi* Said – Said, p. 42, pl. 4, figs. 20a-c

1993 *Cibicides mabahethi* Said – Hottinger *et al.*, p. 115, pl. 151, figs. 6-12

2000 *Cibicides mabahethi* Said – Hyams, pl. 22, figs. 1-2

***Cibicides mabahethi* Said, 1949**

- 1949 *Cibicides mabahethi* Said – Said, p. 42, pl. 4, figs. 20a-c
- 1993 *Cibicides mabahethi* Said – Hottinger *et al.*, p. 115, pl. 151, figs. 6-12
- 2000 *Cibicides mabahethi* Said – Hyams, pl. 22, figs. 1-2

Cibicides mabahethi was first described from the Red Sea by Said (1949). Recent material from the eastern Mediterranean Sea has been recorded by Hyams (2000), Hyams *et al.* (2002) and Hyams-Kaphzan *et al.* (2008). This species is considered to be alien to the Mediterranean Sea.

***Clavulina angularis* d'Orbigny, 1826**

- 1826 *Clavulina angularis* d'Orbigny – d'Orbigny, p. 268, pl. 12, fig. 7
- non 1979 *Clavulina angularis* d'Orbigny – Blanc-Vernet *et al.*, pl. 21, fig. 3
- non 1980 *Clavulina angularis* d'Orbigny – Alavi, p. 11 (appendix), pl. 5, fig. 12, pl. 7, fig. 1
- 1993 *Clavulina angularis* d'Orbigny – Hottinger *et al.*, p. 41, pl. 21, figs. 1-13
- non 2000 *Clavulina angularis* d'Orbigny – Hyams, pl. 2, fig. 8
- non 2008a *Clavulina angularis* d'Orbigny – Meriç *et al.*, pl. 1, figs. 7a-b
- non 2009 *Clavulina angularis* d'Orbigny – Yokeş & Meriç, pl. 1, figs. 6-7
- non 2014 *Clavulina angularis* d'Orbigny – Meriç *et al.*, pl. 7, figs. 3-4
- non 2016 *Clavulina angularis* d'Orbigny – Ayadi *et al.*, pl. 2, fig. 6

Clavulina angularis has first been reported by d'Orbigny (1826) from Corsica in the western Mediterranean Sea. Jones and Parker (1860) reported it from the Gulf of La Spezia (Italy) and Crete (as *Valvulina angularis*, not illustrated). The specimens reported by Alavi (1980), Hyams (2000), Yokeş & Meriç (2009), Meriç *et al.* (2008a; 2014) and Ayadi *et al.* (2016) show rounded last chambers that are absent in the holotype and belong to *Clavulina* cf. *C. multicamerata*. The first record of *Clavulina angularis* shows the species to be present before the opening of the Suez Canal and therefore native to the Mediterranean Sea.

***Clavulina* cf. *C. multicamerata* Chapman, 1907**

- 1907 cf. *Clavulina parisiensis* d'Orbigny var.

- multicamerata* Chapman – Chapman, p. 127, pl. 9, fig. 5
- 1979 *Clavulina angularis* d'Orbigny – Blanc-Vernet *et al.*, pl. 21, fig. 3
- 1980 *Clavulina angularis* d'Orbigny – Alavi, p. 11 (appendix), pl. 5, fig. 12, pl. 7, fig. 1
- 1993 *Clavulina* cf. *C. multicamerata* Chapman – Hottinger *et al.*, p. 42, pl. 22, figs. 1-6
- 2000 *Clavulina angularis* d'Orbigny – Hyams, pl. 2, fig. 8
- 2008a *Clavulina angularis* d'Orbigny – Meriç *et al.*, pl. 1, fig. 7
- 2008a *Clavulina* cf. *C. multicamerata* Chapman – Meriç *et al.*, pl. 1, fig. 8
- 2009 *Clavulina angularis* d'Orbigny – Yokeş & Meriç, pl. 1, figs. 6-7
- 2009 *Clavulina* cf. *C. multicamerata* Chapman – Yokeş & Meriç, pl. 1, figs. 8-9
- 2014 *Clavulina angularis* d'Orbigny – Meriç *et al.*, pl. 7, figs. 3-4
- 2014 *Clavulina* cf. *C. multicamerata* Chapman – Meriç *et al.*, pl. 7, figs. 5-6
- 2014b *Clavulina* cf. *C. multicamerata* Chapman – Katsanevakis *et al.*, p. 688, fig. 20
- 2008b *Peneroplis arietinus* (Batsch) – Meriç *et al.*, pl. 4, figs. 17-19, pl. 5, figs. 1-3, 5-6
- 2010 *Coscinospira hemprichii* Ehrenberg – Koukousioura *et al.*, p. 171, pl. 1, fig. 3

Coscinospira hemprichii was originally described from the Red Sea by Ehrenberg (1839). The specimens illustrated by Batsch (1791) are believed to originate from recent sediments in the Adriatic Sea (Cushman, 1931) and were assigned to *Spirolina hemprichii* (Ehrenberg) or *Spirolina cylindracea* (see Cushman, 1931, p. 72). Modern Mediterranean records include material from the Adriatic Sea, Greece, Israel and Turkey (Cimerman & Langer, 1991; Hollaus & Hottinger, 1997; Hyams *et al.*, 2002; Langer, 2008; Meriç *et al.*, 2008b; Koukousioura *et al.*, 2010). Fossil material of *Coscinospira hemprichii* from the Mediterranean was recently reported by Melis *et al.* (2015), Yümün *et al.* (2016) and Hyams-Kaphzan (unpublished) from the Adriatic Sea, Turkey and Israel, respectively. Both the fossil record and the material from Batsch show that *Coscinospira hemprichii* was present in the Mediterranean prior to the opening of the Suez Canal and the species must therefore be considered to be native.

Cribromiliolinella milletti (Cushman, 1954)

- 1954 *Hauerina milletti* Cushman – Cushman *et al.*, p. 337, pl. 84, fig. 23

Cribromiliolinella milletti has been reported to be alien to the Mediterranean Sea (EASIN, 2020). Cushman *et al.* (1954) described the holotype, *Hauerina milletti*, from the Marshall Islands. No published records exist for this species from the Mediterranean Sea and the taxon is therefore considered to be absent.

Cushmanina striatopunctata (Parker & Jones, 1865)

Coscinospira hemprichii Ehrenberg, 1839

- ? 1791 *Nautilus (Lituus) arietinus* Batsch – Batsch, p. 15, figs. 15d-f
- 1839 *Coscinospira hemprichii* Ehrenberg – Ehrenberg, p. 143, pl. 2, fig. 2
- ? 1987 *Peneroplis pertusus* (Forskål) *arietinus* (Batsch) – Baccaert, p. 60, pl. 19, figs. 3a-b, pl. 20, figs. 1-2
- 1991 *Coscinospira hemprichii* Ehrenberg – Cimerman & Langer, p. 49, pl. 47, figs. 8-11
- 1993 *Coscinospira hemprichii* Ehrenberg – Hottinger *et al.*, p. 69, pl. 76, figs. 1-12, pl. 77, figs. 1-8
- 1997 *Coscinospira hemprichii* Ehrenberg – Hollaus & Hottinger, p. 593
- 2002 *Coscinospira hemprichii* Ehrenberg – Hyams *et al.*, pl. 1, figs. 7-8
- 2008 *Coscinospira hemprichii* Ehrenberg – Langer, fig. 4
- 1865 *Lagena sulcata* Walker and Jacob, var. *striatopunctata* Parker & Jones – Parker & Jones, p. 350, pl. 13, figs. 25-27
- non 1910 *Lagena striatopunctata* Parker & Jones – Sidebottom, p. 17, pl. 2, figs. 5a-b
- non 1940 *Lagena striatopunctata* Parker & Jones – Buchner, p. 444, pl. 7, fig. 100
- 1987 *Cushmanina striatopunctata* (Parker & Jones) – Patterson & Richardson, p. 217, pl. 1, figs. 2-6
- non 2009 *Cushmanina striatopunctata* (Parker & Jones) – Yokeş & Meriç, pl. 4, fig. 5
- non 2014 *Cushmanina striatopunctata* (Parker & Jones) – Meriç *et al.*, pl. 48, fig. 7

Yokeş & Meriç (2009) consider *C. striatopunctata* (Parker & Jones) to be alien to the Mediterranean Sea. However, the illustrated specimen (see also Meriç *et al.*, 2014) differs significantly from the original by Parker & Jones (1865, number of costae and ornamental pattern on the neck) and belongs to a different species. The specimens reported as *Lagena striatopunctata* by Sidebottom

(1910) and Buchner (1940) also differ from the type specimen. To date, there is no reliable record for *Cushmanina striatopunctata* in the Mediterranean Sea and the species is therefore not alien to the Mediterranean Sea. The status and the taxonomic identification of the species illustrated by Yokeş & Meriç (2009) and Meriç *et al.* (2014) require further study.

***Cycloforina quinquecarinata* (Collins, 1958)**

- 1958 *Quinqueloculina quinquecarinata* Collins – Collins, p. 360, pl. 2, figs. 8a-c
- 1993 *Cycloforina quinquecarinata* (Collins, 1958) – Hottinger *et al.*, p. 49, pl. 33, figs. 7-15
- 2016 *Cycloforina quinquecarinata* (Collins, 1958) – Hyams-Kaphzan, pl. 2, figs. 6-7
- non 2016 *Cycloforina quinquecarinata* (Collins, 1958) – Martins *et al.*, pl. 2, figs. 3a-b

Cycloforina quinquecarinata has an Indopacific origin (Collins, 1958) and is also present in the Red Sea (Hottinger *et al.*, 1993). The species has been recorded along the Mediterranean coast of Israel by Hyams-Kaphzan *et al.* (2008; 2009) and Hyams-Kaphzan (2016). Martins *et al.* (2016) reported the species from off Bizerte (Tunisia) but their specimen lacks the typical short neck, has somewhat inflated chambers and the surface shows microstriae that are not present in the original material. New findings from core records along the coast of Israel (Hyams-Kaphzan, unpublished) show that the species has been present in the Mediterranean Sea for at least 250 years (core depth: 80 cm). The species is therefore native to the Mediterranean Sea.

***Cycloforina* sp.**

- 1993 *Cycloforina* sp. C – Hottinger *et al.*, p. 50, pl. 35, figs. 7-13
- 2008 *Cycloforina* sp. – Langer, p. 407, fig. 8

Cycloforina sp. has been reported by Hottinger *et al.* (1993) from the Red Sea (as *Cycloforina* sp. C). The species is characterized by its reticulated test surface ornamentation, produced by intersecting costae (Langer, 2008). It resembles *Quinqueloculina pseudoreticulata* but is provided with a neck and a rounded aperture. Rare specimens were found off Haifa, Israel (Langer, 2008). *Cycloforina* sp. is here considered to be alien to the Mediterranean Sea.

***Cyclorbiculina compressa* (d’Orbigny, 1839)**

- 1839 *Orbiculina compressa* d’Orbigny – d’Orbigny, p. 66, pl. 8, figs. 4-6
- 1884 *Cyclorbiculina compressa* (d’Orbigny) – Brady, pl. 14, figs. 7-9
- 2008a *Cyclorbiculina compressa* (d’Orbigny) – Meriç *et al.*, pl. 11, fig. 12 (non fig. 11)
- 2008b *Cyclorbiculina compressa* (d’Orbigny) – Meriç *et al.*, p. 314, pl. 5, fig. 10 (non figs.

8, 9)

- 2009 *Cyclorbiculina compressa* (d’Orbigny) – Yokeş & Meriç, pl. 3, fig. 2 (non fig. 1)
- 2014 *Cyclorbiculina compressa* (d’Orbigny) – Meriç *et al.*, pl. 40, fig. 2 (non fig. 1)

The holotype of *Cyclorbiculina compressa* has been described by d’Orbigny from Cuba (1839). Smout & Eames (1958) consider the biogeographic range of *Cyclorbiculina compressa* to be restricted to the Caribbean, the Gulf of Mexico and other parts of the tropical western Atlantic. The species has not yet been recorded from the Red Sea (Said, 1949; Hottinger *et al.*, 1993). Some specimens illustrated by Meriç *et al.* (2008a; 2008b; 2014) and Yokeş & Meriç (2009) from Turkey show similarity to the holotype of d’Orbigny, although no profile views were shown; other specimens closely resemble *Sorites*. For the time being, we consider *Cyclorbiculina compressa* to be alien to the Mediterranean Sea. The species was probably introduced into the Mediterranean Sea via passage through the Strait of Gibraltar. Records by Brady (1884) from the Cape Verde Islands support this conclusion.

***Cymbaloporetta* sp. 1**

- ? 1924 *Tretomphalus bulloides* var. *plana* Cushman – Cushman, p. 36, pl. 10, fig. 8
- ? 1968 *Tretomphalus planus* (Cushman) – Moncharmont Zei, pl. 4, figs. 8a-b
- ? 1979 *Pseudotretomphalus planus* (Cushman) – Hofker, p. 23, figs. 12-13
- ? 1987 *Cymbaloporetta plana* (Cushman) – Loeblich & Tappan, pl. 649, figs. 1-3
- ? 1991 *Cymbaloporetta* sp. 1 – Cimerman & Langer, pl. 80, figs. 1-5
- ? 1993 *Cymbaloporetta* sp. A – Hottinger *et al.*, p. 120, pl. 160, figs. 1-8
- ? 1994 *Cymbaloporetta plana* (Cushman) – Jones, pl. 102, figs. 7-8, 12
- ? 2000 *Cymbaloporetta* sp. 1 – Hyams, pl. 23, figs. 6-8
- ? 2008b *Cymbaloporetta plana* (Cushman) – Meriç *et al.*, pl. 8, figs. 2-7
- 2010 *Cymbaloporetta plana* (Cushman) – Koukousioura *et al.*, pl. 2, figs. 1-2
- 2014 *Cymbaloporetta plana* (Cushman) – Meriç *et al.*, pl. 62, figs. 12-13
- ? 2014 *Cymbaloporetta plana* (Cushman) – Meriç *et al.*, pl. 62, figs. 14-16

The holotype of *Cymbaloporetta plana* Cushman (1924) was described from Samoa (Pacific Ocean). Although the quality of the original illustration is poor, photographs of the holotype provided by the Smithsonian Museum of Natural History (2020) appear to be identical with most specimens illustrated from the Mediterranean Sea. *Cymbaloporetta* sp. 1 has been frequently reported from the Mediterranean Sea (as *C.* sp. 1 and *C. plana*; see synonymie list). The species has been recorded by Hottinger *et al.* (1993) from the Red Sea (as *C.* sp. A). Hofker (1979) recorded the species (as *Pseudotretomphalus planus*) from the Caribbean. Some authors (Rückert-Hilbig,

1983) consider *C. plana* to be a synonym of *C. bulloides* (d'Orbigny, 1839). The complex taxonomic status of the species (e.g. *C. plana*, *C. bulloides*, *C. sp.*) has not yet been resolved and requires further studies. The species is widespread and present in most oceans (Pacific, Atlantic, Red Sea, Mediterranean Sea). Until molecular genetic data are available for the specimens from the type locality and other regions, and because of its complex taxonomic status, we consider the status of *Cymbaloporetta* sp. 1 from the Mediterranean Sea as cryptogenic.

***Edentostomina cultrata* (Brady, 1881)**

- 1881 *Miliolina cultrata* Brady – Brady, p. 45
- 1884 *Miliolina cultrata* Brady – Brady, p. 161, pl. 5, figs. 1-2
- 1987 *Edentostomina cultrata* (Brady) – Loeblich & Tappan, pl. 334, figs. 6-8
- 1980 *Edentostomina cultrata* (Brady) – Alavi, p. 14 (appendix), pl. 10, fig. 11
- non 2004 *Edentostomina cultrata* (Brady) – Meriç *et al.*, pl. 5, fig. 1
- 2006 *Edentostomina cultrata* (Brady) – Oflaz, pl. 2, fig. 3
- 2006 *Edentostomina milletti* (Wiesner) – Oflaz, pl. 2, figs. 4-6
- non 2008b *Edentostomina cultrata* (Brady) – Meriç *et al.*, pl. 2, figs. 1-5

Edentostomina cultrata was first described by Brady (1884) from Papua New Guinea. Mediterranean records from Turkey were given by Alavi (1980) and Oflaz (2006). The specimens identified as *E. cultrata* by Meriç *et al.* (2004; 2008b) do not show the typical thick, everted apertural rim, lack a pronounced neck and therefore belong to a different species of *Edentostomina*. The species *Edentostomina milletti*, originally described by Cushman (1917, as *Biloculina milletti*), looks similar but has a short bifid tooth. Records of *Edentostomina cultrata* in recent surface samples along the Mediterranean coast of Israel as well as findings of this species in core material deposited prior to the opening of the Suez Canal (Hyams-Kaphzan, unpublished) indicate that this species is native to the Mediterranean Sea.

***Elphidium charlottense* (Vella, 1957)**

- 1957 *Elphidiononion charlottensis* Vella – Vella, p. 38, pl. 9, figs. 187-188
- 1958 *Elphidium advenum* (Cushman) – Parker, pl. 4, figs. 3-4
- 1980 *Elphidium advenum* (Cushman) – Alavi, pl. 32, fig. 12
- 1991 *Elphidium* sp. 1 – Cimerman & Langer, p. 79, pl. 89, fig. 8
- 1997 *Elphidium charlottense* (Vella) – Hayward *et al.*, p. 72, pl. 6, figs. 15-16
- 2004 *Elphidium advenum* (Cushman) – Meriç *et al.*, pl. 32, figs. 9-10
- 2006 *Elphidium* sp. 1 – Langer & Schmidt-Sinns,

- pl. 15, fig. 5
- 2008a *Elphidium charlottense* (Vella) – Meriç *et al.*, pl. 16, figs. 1-3
- ? 2012 *Elphidium advenum* (Cushman) – Debenay, p. 218
- 2017 *Elphidium namibium/advenum* – Mouanga, pl. 16, fig. 17

The holotype of *Elphidium charlottense* has been described by Vella (1957) from Rellings Bay, Queen Charlotte Sound, Marlborough (New Zealand). The species is characterized by its rounded keel, translucent wall and non-inflated chambers resulting in a comparatively smooth test surface. The species differs from *Elphidium advenum* (Cushman), a species that has more inflated chambers and deeper interocular septae and spaces. *Elphidium charlottense* has not been recorded from the Red Sea (Hottinger *et al.*, 1993) but is present in shallow waters off the coast of Namibia (Langer, unpublished). *Elphidium charlottense* has been found in core material collected off Greece (Parker, 1958, as *Elphidium advenum*) and is therefore considered to be native to the Mediterranean Sea.

***Elphidium striatopunctatum* (Fichtel & Moll, 1798)**

- 1798 *Nautilus striatopunctatus* Fichtel & Moll – Fichtel & Moll, p. 61, pl. 9, figs. a-c
- non 1909 *Polystomella striatopunctata* (Fichtel & Moll) – Sidebottom, pl. 4, fig. 10, pl. 5, figs. 1-2
- 1993 *Elphidium striatopunctatum* (Fichtel & Moll) – Hottinger *et al.*, p. 149, pl. 213, figs. 1-8, pl. 214, figs. 1-6
- 2000 *Elphidium striatopunctatum* (Fichtel & Moll) – Hyams, pl. 27, figs. 7-8
- 2001 *Elphidium striato-punctatum* (Fichtel & Moll) – Avşar *et al.*, pl. 4, figs. 8-11
- 2009 *Elphidium striatopunctatum* (Fichtel & Moll) – Yokeş & Meriç, pl. 5, figs. 8-10
- 2014 *Elphidium striatopunctatum* (Fichtel & Moll) – Meriç *et al.*, pl. 83, figs. 16-17

Elphidium striatopunctatum has been described from the Red Sea by Fichtel & Moll (1798). Modern Mediterranean records include specimens from Israel (Hyams, 2000) and Turkey (Avşar *et al.*, 2001; Yokeş & Meriç, 2009; Meriç *et al.*, 2014). Not illustrated records include Jones and Parker (1860), Schaudinn (1911) and Wiesner (1911a; 1911b). The illustrations given by Sidebottom (1909) show a different species. *Elphidium striatopunctatum* is therefore considered to be alien to the Mediterranean Sea.

***Epistomaroides punctulata* (Said, 1949)**

- 1949 *Epistomaria punctata* Said – Said, p. 37, pl. 4, figs. 23a-c
- 1980 *Anomalina punctulata* d'Orbigny – Hansen & Rögl, pl. 1, figs. 4-8

- 1993 *Epistomaroides punctatus* (Said) – Hottinger *et al.*, p. 131, pl. 180, figs. 1-11, pl. 181, figs. 1-6
- 2012 *Epistomaroides punctatus* (Said) – Almogi-Labin & Hyams-Kaphzan, fig. 2
- 2017 *Epistomaroides punctulatus* (d'Orbigny) – Thissen & Langer, pl. 19, figs. 1-5

The holotype of *Epistomaroides punctulata* was described by Said (1949) from the Red Sea. According to Hottinger *et al.* (1993), the specimens recorded by d'Orbigny (1826) as *Anomalina punctulata* are identical to the specimens described by Hansen & Rögl (1980). The material of d'Orbigny (1826) was collected around the Island of Mauritius, Indian Ocean. The first Mediterranean record of this species along the northern coast of Israel was provided by Almogi-Labin & Hyams-Kaphzan (2012). Today, the species is very abundant and widely distributed along the Mediterranean rocky coast of northern Israel (Hyams-Kaphzan, unpublished). We consider this species to be alien to the Mediterranean Sea.

Euthymonacha polita (Chapman, 1900)

- 1900 *Peneroplis (Monalysidium) polita* Chapman – Chapman, p. 4, pl. 1, fig. 5
- 1994 *Monalysidium politum* (Chapman) – Jones, p. 29, pl. 13, figs. 24-25
- 2010 *Euthymonacha polita* (Chapman) – Meriç *et al.*, p. 192, fig. 2
- 2013a *Euthymonacha polita* (Chapman) – Langer *et al.*, fig. 7, 23-24
- 2014 *Euthymonacha polita* (Chapman) – Meriç *et al.*, pl. 35, figs. 8-15
- 2015 *Euthymonacha polita* (Chapman) – Delliou *et al.*, pl. 1, fig. 3
- 2016 *Euthymonacha polita* (Chapman) – Huth, p. 31, pl. 1, figs. 10-12
- 2017 *Euthymonacha polita* (Chapman) – Mouanga, pl. 1, figs. 1a-d
- 2018 *Euthymonacha polita* (Chapman) – Emter, pl. 21, fig. 6

Euthymonacha polita was first described by Chapman (1900) from Funafuti (Pacific Ocean). Mediterranean records include findings from the Gulf of Kuşadası, Ilica Bay and the Karaburun Peninsula in Turkey (Meriç *et al.*, 2010; 2011), from the Kavala Gulf in Greece (Delliou *et al.*, 2015), from off Otranto, Italy (Huth, 2016), Libya (Emter, 2018) and the Balearic Islands off Spain (Alvira Romero, 2019). Brady (1884) recorded it from the Cape Verde Islands (Atlantic Ocean) as *Peneroplis pertusus* var. *e* (see Jones, 1994), and Langer *et al.* (2013a) recorded it from the Bazaruto Archipelago (Mozambique). So far, the species has not been recorded from the Red Sea or the Mediterranean coast of Israel (Said, 1949; Hottinger *et al.*, 1993; Hyams-Kaphzan *et al.*, 2008). The circum-tropical distribution and the widespread occurrence in the Mediterranean Sea suggest that this species may possibly be native to the Mediterranean. However, no record prior

to the opening of the Suez Canal exists to date. For the time being we consider the species to be cryptogenic and further evidence is needed to resolve its status.

Euvigerina sp.

- 1982 *Uvigerina peregrina* Cushman – Foraminiferi Padani, pl. 34, figs. 1-1bis
- 2004 *Euvigerina* sp. – Meriç *et al.*, pl. 22, figs. 2-3

Euvigerina sp. was reported by Meriç *et al.* (2004) from Turkey and claimed to be alien to the Mediterranean Sea. The illustration provided is identical to specimens of *Uvigerina peregrina* reported in Foraminiferi Padani (1982), which, according to the authors, has been present in the Mediterranean Sea since the Pliocene. However, the holotype of *Uvigerina peregrina* Cushman differs from *Euvigerina* sp. of Meriç *et al.* (2004). The taxonomic status of *Euvigerina* sp. requires further study, but the species is certainly present in fossil material and therefore native to the Mediterranean Sea.

Guttulina? sp.

- 2008a *Entosigmomorphina* sp. – Meriç *et al.*, pl. 13, figs. 8a-b
- 2014 *Entosigmomorphina* sp. – Meriç *et al.*, pl. 49, figs. 8a-b
- non 2014 *Entosigmomorphina* sp. – Caruso & Cosentino, pl. 4, fig. 12

Meriç *et al.* (2008a) provide the first illustrated record of this species from the Mediterranean Sea. The chambers of the illustrated specimen are rapidly increasing in size and height and successive chambers appear to be 144° apart. The sutures are strongly depressed and the aperture is radiate. This would place the specimen in the genus *Guttulina* d'Orbigny, 1839. As this represents the first record of this taxon, it is considered to be native to the Mediterranean Sea. The specimen illustrated by Caruso & Cosentino (2014) does not belong to the genus *Entosigmomorphina*, differs from Meriç *et al.* (2008a, 2014), and requires further study.

Haddonia sp.

- 1993 *Haddonia?* sp. C – Hottinger *et al.*, p. 31, pl. 4, figs. 5-9
- 2008a *Haddonia* sp. – Meriç *et al.*, pl. 1, figs. 1-4
- 2008b *Haddonia* sp. – Meriç *et al.*, pl. 1, figs. 1-14
- 2009 *Haddonia* sp. – Yokeş & Meriç, pl. 1, figs. 1-3
- 2014 *Haddonia* sp. – Meriç *et al.*, pl. 2, figs. 7-17

Haddonia is a genus described from coral reefs from the Torres Strait (off northern Australia) by Chapman (1898). Hottinger *et al.* (1993) recorded a total of four different species in the Red Sea. *Haddonia* sp. was reported from Turkey by Meriç *et al.* (2007, see also Meriç *et al.*, 2008b; Yokeş & Meriç, 2009; and Meriç *et al.*,

2014). Poorly preserved specimens of *Haddonina* were also found in surface samples from rocky reefs off the coast of Israel (Hyams-Kaphzan, unpublished; material collected in 1996). The species *Haddonina* sp. is here considered to be alien to the Mediterranean Sea.

***Hauerina diversa* Cushman, 1946**

- 1932 *Hauerina bradyi* Cushman – Cushman, p. 44, pl. 10, figs. 12-15
 1946 *Hauerina diversa* Cushman – Cushman, p. 11, pl. 2, figs. 16-19
 1993 *Hauerina diversa* Cushman – Hottinger *et al.*, p. 50, pl. 36, figs. 1-7
 1998 *Hauerina diversa* Cushman – Yanko *et al.*, p. 195, pl. 5, figs. 1-3
 2000 *Hauerina diversa* Cushman – Hyams, pl. 6, figs. 6-7
 2006 *Sigmoihauerina bradyi* (Cushman) – Oflaz, p. 170, pl. 3, fig. 9
 2008a *Hauerina diversa* Cushman – Meriç *et al.*, pl. 4, figs. 9-12
 2008b *Hauerina diversa* Cushman – Meriç *et al.*, pl. 3, figs. 12-14
 2010 *Hauerina diversa* Cushman – Avnaim-Katav, pl. 2, fig. 12
 2014 *Hauerina diversa* Cushman – Meriç *et al.*, pl. 16, figs. 11-15
 2017 *Hauerina diversa* Cushman – Guy-Haim *et al.*, pl. 4 fig. 16, pl. 5, figs. 10-11

Hauerina diversa was first described by Cushman (1946) from Hereheretue, French Polynesia. Mediterranean records of *Hauerina diversa* include material from the Turkish Aegean coast (Meriç *et al.*, 2008a; 2008b) and from Israel (Yanko *et al.*, 1998; Hyams, 2000; Hyams-Kaphzan *et al.*, 2008; Avnaim-Katav, 2010 core-top). Non-illustrated records from Turkey also include Avşar (1997 and Avşar *et al.* (2001)). The species has recently been found in large numbers along the rocky coast of northern Israel (Hyams-Kaphzan, unpublished). The Mediterranean and Red Sea forms do not show internal septulae and are placed into the genus *Hauerina*. We consider *Hauerina diversa* to be alien to the Mediterranean Sea. The species has been recovered from fish guts collected in the the Suez Canal and the Mediterranean Sea (Guy-Haim *et al.*, 2017), indicating that ichthyochory is a potential vector for long-distance transport and species introduction.

***Heterocyclus tuberculata* (Möbius, 1880)**

- 1880 *Heterostegina tuberculata* Möbius – Möbius, p. 107, pl. 12, figs. 3-7
 1977 *Heterocyclus tuberculata* (Möbius) – Reiss *et al.*, p. 104, figs. 35 A-C, 36 A
 1993 *Heterocyclus tuberculata* (Möbius) – Hottinger *et al.*, p. 156, pl. 226, figs. 1-7, pl. 227, figs. 1-8, pl. 230, fig. 10

Heterocyclus tuberculata has been described by Möbius (1880) from the western Indian Ocean. It has been recorded from the Turkish Mediterranean coast by Avşar *et al.* (2001; 2008) and from Greece (Debenay *et al.*, 2005) but the specimens are not illustrated. Langer & Hottinger (2000) consider *Heterocyclus tuberculata* to be “restricted to the Indian Ocean and the Red Sea”. Illustrations of this species from the Mediterranean Sea were not published and we therefore consider the species to be absent from the Mediterranean Sea.

***Heterostegina depressa* d’Orbigny, 1826**

- 1826 *Heterostegina depressa* d’Orbigny – d’Orbigny, p. 305, pl. 17, figs. 5-7
 non 1865 *Heterostegina depressa* d’Orbigny – Parker *et al.*, p. 34, pl. 3, fig. 100
 non 1907 *Heterostegina depressa* d’Orbigny – Silvestri, p. 43, pl. 2, fig. 5
 1968 *Heterostegina antillarum* d’Orbigny – Moncharmont Zei, p. 27, pl. 5, figs. 3-7
 1980 *Heterostegina antillarum* d’Orbigny – Alavi, p. 27, pl. 35, figs. 3-7
 1993 *Heterostegina depressa* d’Orbigny – Hottinger *et al.*, p. 157, pl. 228, figs. 1-11, pl. 229, figs. 1-8, pl. 230, fig. 9
 2000 *Heterostegina depressa* d’Orbigny – Langer & Hottinger., p. 110, text-fig. 4A
 2002 *Heterostegina depressa* d’Orbigny – Hyams *et al.*, pl. 1, fig. 9
 2008 *Heterostegina depressa* d’Orbigny – Langer, p. 403, fig. 2
 2008b *Heterostegina depressa* d’Orbigny – Meriç *et al.*, p. 323, pl. 10, figs. 1-12
 2010 *Heterostegina depressa* d’Orbigny – Avnaim-Katav, p. 236, pl. 10, fig. 20

Heterostegina depressa has been described by d’Orbigny (1826) from the island of Saint Helena in the South Atlantic. The species is a true circumtropical foraminifer and its distribution encompasses the Caribbean, the North and South Atlantic, the entire Indian and Pacific Oceans and the Red and Arabian Sea (Langer & Hottinger, 2000; Langer, 2008 and references therein). Mediterranean records include specimens from Lebanon (Moncharmont Zei, 1968, as *H. antillarum*), Israel (Yanko *et al.*, 1998, not illustrated; Hyams *et al.*, 2002; Avnaim-Katav, 2010 core-top) and Turkey (Alavi, 1980; Meriç *et al.*, 2008b). Fossil occurrences of this species were reported by Parker *et al.* (1865) from “Middle Tertiary limestones of Malta and Vienna” and by Silvestri (1907) from a cave near Turin, Italy. However, the quality of the illustrations is too poor to assign the species to *H. depressa* and the specimens probably belong to a different species (Hottinger, pers. comm.). *Heterostegina depressa* is therefore considered to be alien to the Mediterranean Sea.

***Loxostomina* cf. *L. africana* (Smitten, 1955)**

- 1955 cf. *Loxostomum africanum* Smitten –

- Smutter, p. 118, fig. 40 (o)
- 1975 *Brizalina (Parabrizalina)* cf. *B. (P.) africana* (Smutter) – Zweig-Strykowski & Reiss, p. 100, pl. 3, figs. 1-8
- 1980 *Bolivina africana* (Smutter) – Alavi, p. 42 (appendix), pl. 26, figs. 5, 9-10
- 1993 *Loxostomina* cf. *L. africana* (Smutter) – Hottinger *et al.*, p. 97, pl. 119, figs. 10-15
- 2000 *Loxostomina* cf. *L. africana* (Smutter) – Hyams, pl. 18, fig. 1
- non 2017 *Brizalina africana* (Smutter) – Makled *et al.*, p. 479, pl. 16, fig. 18

Loxostomina africana has been described by Smutter (1955). The material was collected from sediments off Mozambique and are “probably Upper Miocene” in age. The holotype has a smooth surface texture. Hottinger *et al.* (1993) illustrated a similar species. However, the Red Sea specimens have distinct longitudinal striae across the entire test surface and were therefore referred to as *Loxostomina* cf. *L. africana*. Striated representatives of this species were also recorded from Turkey by Alavi (1980) and from Israel by Hyams (2000), Hyams (2006), and Hyams-Kaphzan *et al.* (2008). So far, the species has not been recorded in fossil material. The first record of *Loxostomina* cf. *L. africana* from the Mediterranean Sea was provided by Alavi (1980). As the oldest record for this species originates from the Red Sea (Zweig-Strykowski & Reiss, 1975) the species is considered to be alien to the Mediterranean Sea.

Loxostomina costulata (Cushman, 1922)

- 1922 *Bolivina limbata* var. *costulata* Cushman – Cushman, p. 26, pl. 3, fig. 8
- 1993 *Loxostomina? limbata* (Brady) *costulata* (Cushman) – Hottinger *et al.*, p. 97, pl. 120, figs. 8-13
- ? 2000 *Loxostomina? limbata* (Brady) *costulata* (Cushman) – Hyams, pl. 18, fig. 2
- 2016 *Loxostomina limbata* (Brady) *costulata* (Cushman) – Huth, p. 32, pl. 2, figs. 1-2
- 2017 *Loxostomina limbata* (Brady) *costulata* (Cushman) – Mouanga, pl. 12, figs. 10a-b

Loxostomina costulata has been described by Cushman (1922) from the Tortugas (Florida). Hottinger *et al.* (1993) recorded specimens from the Gulf of Aqaba, Red Sea. The species was also found in the Adriatic Sea off Otranto, Italy (Huth, 2016). Specimens from the coast of Israel provided by Hyams (2000) do not show the strong costate surface as illustrated by Hottinger *et al.* (1993) but may represent a morphological variety of this species. Fossil specimens of *Loxostomina costulata* with faint costate surfaces, were also found in core material from Israel (Hyams-Kaphzan, unpublished). The species is therefore considered to be native to the Mediterranean Sea.

Mimosina affinis Millett, 1900

- 1900 *Mimosina affinis* Millett – Millett, p. 548, pl. 4, fig. 11
- 1993 *Mimosina affinis* Millett – Hottinger *et al.*, p. 104, pl. 133, figs. 9-12, pl. 134, figs. 1-3
- 2017 *Mimosina affinis* Millett – Mouanga, pl. 13, figs. 1a-b

Mimosina affinis has been first described by Millett (1900) from the Malay Archipelago. A single Mediterranean record of this species was provided by Mouanga (2017) from the Bay of Vlore, Albania. *Mimosina affinis* is considered to be alien to the Mediterranean Sea.

Monalysidium acicularis (Batsch, 1791)

- 1791 *Nautilus lituus acicularis* Batsch – Batsch, p. 3, pl. 6, figs. 16a-b
- 1993 *Monalysidium acicularis* (Batsch) – Hottinger *et al.*, p. 70, pl. 78, figs. 1-14
- 2011 *Coscinospira acicularis* (Batsch) – Meriç *et al.*, p. 2, fig. 3
- 2019 *Monalysidium acicularis* (Batsch) – Förderer & Langer, pl. 2, figs. 32-35

Monalysidium acicularis was first described by Batsch (1791) but the type locality was not provided. Cushman (1931) considers the specimens to originate “from Rimini or at least the Adriatic Sea”. The species is abundant in the Red Sea (Hottinger *et al.*, 1993). Records of *Spirolina acicularis* from the Mediterranean include specimens from Turkey (Alavi, 1980), Israel (Yanko *et al.*, 1998) and Italy (Aiello *et al.*, 2006), but all lack illustrations. A single illustrated record of *M. acicularis* from the Mediterranean Sea was provided by Meriç *et al.* (2011, as *Coscinospira acicularis*). The information provided by Cushman on the type locality from the Adriatic Sea suggests that *Monalysidium acicularis* is native to the Mediterranean Sea.

Neoconorbina clara (Cushman, 1934)

- 1934 *Tretomphalus clarus* Cushman – Cushman, p. 99, pl. 11, figs. 6 a-c, pl. 12, figs. 16-17
- 1993 *Tretomphaloides clara* (Cushman) – Hottinger *et al.*, p. 112, pl. 145, figs. 6-11
- 2008 *Tretomphaloides clara* (Cushman) – Hyams-Kaphzan *et al.*, p. 344

The holotype of *Neoconorbina clara* has been described by Cushman (1934) from Guam Anchorage, Ladrone Islands (western Pacific Ocean). Hottinger *et al.* (1993) recorded this species from the Gulf of Aqaba, Red Sea. The species was reported from the Mediterranean coast of Israel (Hyams-Kaphzan *et al.*, 2008; Hyams-Kaphzan, unpublished). *Neoconorbina clara* is here considered to be alien to the Mediterranean Sea.

Nodophthalmidium antillarum (Cushman, 1922)

- 1922 *Articulina antillarum* Cushman – Cushman, p. 71, pl. 12, fig. 5
- 1929 *Nodophthalmidium antillarum* (Cushman) – Cushman, p. 52, pl. 12, fig. 4
- 1949 *Nodophthalmidium antillarum* (Cushman) – Said, p. 20, pl. 2, fig. 3
- 1968 *Nodophthalmidium antillarum* (Cushman) – Moncharmont Zei, pl. 3, fig. 3
- 1980 *Nodophthalmidium antillarum* (Cushman) – Alavi, pl. 20, fig. 9
- 1993 *Nodophthalmidium antillarum* (Cushman) – Hottinger *et al.*, p. 44, textfig. 2, pl. 23, figs. 4-7
- 2000 *Nodophthalmidium antillarum* (Cushman) – Hyams, pl. 3, fig. 4
- non 2008b *Nodophthalmidium antillarum* (Cushman) – Meriç *et al.*, pl. 2, fig. 10 (non figs. 11-14)
- non 2009 *Nodophthalmidium antillarum* (Cushman) – Yokeş & Meriç, pl. 1, figs. 10-11
- 2013a *Articulina antillarum* Cushman – Langer *et al.*, pl. 7, figs. 6-7
- 2014 *Nodophthalmidium antillarum* (Cushman) – Meriç *et al.*, pl. 8, figs. 11-12
- non 2017 *Nodophthalmidium antillarum* (Cushman) – Mouanga, pl. 3, fig. 6

Nodophthalmidium antillarum was first described by Cushman (1922) from Tortugas, Florida (Cushman, 1922). The species was also recorded from the Red Sea (Said, 1949; Hottinger *et al.*, 1993) and the Bazaruto Archipelago, Mozambique (Langer *et al.*, 2013a). The first Mediterranean record was provided by Moncharmont Zei (1968) from material collected at Ras Muker Ben and Ras Minet el Hosn (Lebanon) at depth between 75 m and 246 m. Other Mediterranean records include specimens from Turkey (Alavi, 1980; Meriç *et al.*, 2008b) and Israel (Hyams, 2000; Yanko *et al.*, 1998, not illustrated). Specimens illustrated by Meriç *et al.* (2008b) and Yokeş & Meriç (2009) differ substantially from the holotype (chamber shape and striation pattern) and belong to a different species. The juvenile specimen illustrated and recorded by Mouanga (2017) from the coast of Albania is also a different species. *Nodophthalmidium antillarum* is considered to be alien to the Mediterranean Sea.

Operculina ammonoides (Gronovius, 1781)

- 1781 *Nautilus ammonoides* Gronovius – Gronovius, p. 282 (no. 1220), pl. 19, figs. 5-6
- 1993 *Assilina ammonoides* (Gronovius) – Hottinger *et al.*, p. 154, pl. 222, figs. 1-8, pl. 223, figs. 1-14, pl. 224, figs. 1-8, pl. 225, figs. 1-9
- 2016 *Operculina ammonoides* (Gronovius) – Merkado, fig. 2 A, fig. 21, fig. 24, fig. 26

The holotype of *Operculina ammonoides* has been described by Gronovius (1781) from the Bay of Bengal (In-

dian Ocean). Hottinger *et al.* (1993) recorded this species from the Gulf of Aqaba, Red Sea. The species also has been reported from the Adriatic Sea by Wiesner (1913) and Silvestri (1950) and by Yokeş & Meriç (2009) from Turkey. The latter records, however, lack illustrations. *Operculina ammonoides* was recently reported from the Mediterranean coast of Israel (Merkado, 2016). It is currently also present in rocky reefs of Shikmona, Haifa (Hyams-Kaphzan, unpublished). Populations along the coast of Israel are genetically closely related to specimens from the Red Sea and Japan (Merkado, 2016). The data currently available indicate that *Operculina ammonoides* is alien to the Mediterranean Sea.

Pararotalia calcariformata McCulloch, 1977

- 1977 *Pararotalia calcariformata* McCulloch – McCulloch, p. 428, pl. 177, figs. 10-11
- 1980 *Pararotalia* aff. *P. bisaculeata* (d'Orbigny) – Alavi, pl. 30, fig. 8
- 1994 *Eponides repandus* (Fichtel & Moll) – Yanko *et al.*, pl. 2, figs. 1-9
- 1994 *Pararotalia spinigera* (Le Calvez) – Reinhardt *et al.*, pl. 2, figs. 11-12
- 1998 *Pararotalia spinigera* (Le Calvez) – Yanko *et al.*, pl. 1, figs. 6-9
- 2000 *Pararotalia spinigera* Le Calvez – Hyams, pl. 24, figs. 5-7
- 2010 *Pararotalia spinigera* Le Calvez – Avnaim-Katav, pl. 9, figs. 7-10
- 2013 *Pararotalia calcariformata* McCulloch – Meriç *et al.*, figs. 2-3
- 2015 *Pararotalia calcariformata* McCulloch – Schmidt *et al.*, fig. 1
- 2016 *Pararotalia calcariformata* McCulloch – Titelboim *et al.*, fig. 5, 3-4
- 2020 *Pararotalia calcariformata* McCulloch – Avnaim-Katav *et al.*, fig. 4, 18a, b
- 2020 *Pararotalia calcariformata* McCulloch – Manda *et al.*, fig. 3C

The holotype of *Pararotalia calcariformata* was collected at Colombo Bay, Ceylon, Indian Ocean (McCulloch, 1977). This species has been considered to be alien to the Mediterranean Sea by Meriç *et al.* (2013) and Schmidt *et al.* (2015). *Pararotalia calcariformata* has been frequently reported as *Pararotalia spinigera* (Reinhardt *et al.*, 1994; Yanko *et al.*, 1998; Hyams, 2000; Hyams-Kaphzan *et al.*, 2008). It is currently widely present and increasing in abundance off the coast of Israel from very shallow water depths down to 40 m (Hyams-Kaphzan *et al.*, 2014; Titelboim *et al.*, 2016; Avnaim-Katav *et al.*, 2020; Manda *et al.*, 2020). It has not yet been found in the Red Sea but is present around Bazaruto (Mozambique; Langer *et al.*, 2013a). However, fossil evidence shows *P. calcariformata* (reported as *P. spinigera*) to be present at Caesarea Maritima (Israel, Mediterranean Sea) by 171BC-78AD (Reinhardt *et al.*, 1994) as well as in Haifa Bay (HB30, 5 m), Israel (Avnaim-Katav, 2010). We therefore consider the species to be native to the Mediterranean Sea.

***Pararotalia cf. P. socorroensis* (McCulloch, 1977)**

- 1977 cf. *Praeglobotruncana? socorroensis*
McCulloch – McCulloch, p. 424, pl. 178,
fig. 4, pl. 179, figs. 1, 3
- 1993 *Pararotalia cf. P. socorroensis* (McCulloch)
– Hottinger *et al.*, p. 141, pl. 200, figs. 1-11
- 2008 *Pararotalia cf. P. socorroensis* (McCulloch)
– Hyams-Kaphzan *et al.*, p. 344

The holotype of *Pararotalia socorroensis* is from Socorro Island, eastern Pacific Ocean (McCulloch, 1977). The species has been recorded in the Red Sea by Hottinger *et al.* (1993) and from the eastern Mediterranean Sea by Hyams-Kaphzan *et al.* (2008) and Hyams-Kaphzan (unpublished). *Pararotalia cf. P. socorroensis* is considered to be alien to the Mediterranean Sea.

***Pararotalia spinigera* Loeblich & Tappan, 1957 (ex. Le Calvez, 1949)**

- 1987 *Pararotalia spinigera* (Le Calvez) –
Loeblich & Tappan, p. 18, pl. 4, figs. 1a-3
- non 1994 *Pararotalia spinigera* (Le Calvez) –
Reinhardt *et al.*, pl. 2, figs. 11-12
- non 2000 *Pararotalia spinigera* Le Calvez – Hyams,
pl. 24, figs. 5-7

Pararotalia spinigera was considered to be alien to the Mediterranean Sea (Zenetos *et al.*, 2012). All illustrated Mediterranean records are now considered to be identical with *Pararotalia calcariformata*, a species that is native to the Mediterranean Sea (for further details, see *Pararotalia calcariformata*).

***Parasorites orbitolitoideis* (Hofker, 1930)**

- 1930 *Praesorites orbitolitoideis* Hofker – Hofker,
p. 149, pl. 55, figs. 8, 10, 11, pl. 58, figs.
1-5, pl. 59, figs. 3, 14
- non 1977 *Sorites orbitolitoideis* (Hofker) – Hottinger
et al., figs. 11B, 13, 30A-C, 32A
- 1994 *Sorites orbitolitoideis* (Hofker) –
Gudmundsson, figs. 55-58, pl. 7, figs. 1-3
- 2017 *Parasorites/Broeckina orbitolitoideis*
(Hofker) – Nadal Nebot, pl. 1, fig. *
- 2018 *Parasorites* sp. – Mateu-Vicens *et al.*, 2018.
- 2018 *Parasorites orbitolitoideis* (Hofker) –
Ferragut Perelló, pl. 1, fig. 6, textfig. 6
- 2019 *Parasorites orbitolitoideis* (Hofker) – Alvira
Romero, fig. 1

Parasorites orbitolitoideis has been first described by Hofker (1930) from Indonesia. The species has not been recorded in the Red Sea (Hottinger *et al.*, 1993). Reliable records for *Parasorites orbitolitoideis* from the Mediterranean Sea are known from the Balearic Islands, Spain (Nadal Nebot, 2017; Ferragut Perelló, 2018; Mateu-Vicens *et al.*, 2018 (not illustrated, but re-examined); Alvira Romero, 2019). Populations of Mediterranean *P. orbitolitoideis*

show a genetic similarity of 98% with the Indo-Pacific population analyzed from Guam (Alvira Romero, 2019). For the time being, we consider this species to represent a true alien in the Mediterranean Sea.

***Paratrochammina madeirae* Brönnimann, 1979**

- 1979 *Paratrochammina madeirae* Brönnimann
– Brönnimann, p. 7, pl. 7, figs. a-c, f, h, pl.
10, figs. b, e
- 1993 *Paratrochammina madeirae* Brönnimann –
Hottinger *et al.*, p. 32, pl. 7, figs. 11-15
- 2000 *Paratrochammina* sp. 1 – Hyams, pl. 1,
figs. 9-10
- non 2004 *Paratrochammina madeirae* Brönnimann –
Diz *et al.*, pl. 1, figs. 4a-b
- 2008 *Paratrochammina madeirae* Brönnimann
– Hyams-Kaphzan *et al.*, p. 336 (not
illustrated)

Paratrochammina madeirae has first been recorded from Brazil (Brönnimann, 1979) and later from the Red Sea (Hottinger *et al.*, 1993). The species has also been found along the coast of Israel (Hyams, 2000; Hyams-Kaphzan *et al.*, 2008) and is considered to be alien to the Mediterranean Sea.

***Pegidia lacunata* McCulloch, 1977**

- 1977 *Pegidia lacunata* McCulloch – McCulloch,
p. 347, pl. 154, fig. 2
- 1993 *Pegidia lacunata* McCulloch – Hottinger *et al.*,
p. 108, pl. 139, figs. 7-9, pl. 140, figs. 1-5
- 2008 *Pegidia lacunata* McCulloch – Langer, p.
407, fig. 9

Pegidia lacunata has been described by McCulloch (1977) from the Philippine Islands, western Pacific. Hottinger *et al.* (1993) have reported the species from the Gulf of Aqaba at depths between 20 and 190 meters. The species has been rarely observed at a single sample site off Haifa, Israel (Langer, 2008, material collected in 1994) and is here considered to be alien to the Mediterranean Sea.

***Peneroplis cf. P. antillarum* (d'Orbigny, 1839)**

- 1826 cf. *Dendritina antillarum* d'Orbigny –
d'Orbigny, p. 285
- non 1839 *Dendritina antillarum* d'Orbigny –
d'Orbigny, p. 58, pl. 7, figs. 3-6
- non 1994 *Peneroplis antillarum* (d'Orbigny) –
Gudmundsson, p. 111, textfigs. 19, 20, pl. 3,
fig. 4, pl. 4, fig. 4
- non 2000 *Peneroplis antillarum* (d'Orbigny) –
Hohenegger *et al.*, text-fig. 6
- 2002 *Peneroplis antillarum* (d'Orbigny) – Hyams
et al., pl. 1, figs. 10-11
- non 2019 *Peneroplis antillarum* (d'Orbigny) –
Förderer and Langer, pl. 2, figs. 16-18

Peneroplis antillarum has been described from the Antilles, Caribbean Sea (d'Orbigny, 1826). Specimens recorded by Hyams *et al.* (2002) from the Mediterranean coast off Israel differ from the specimens illustrated by d'Orbigny (1826, 1839) and Hohenegger *et al.* (2000). The test surface of the Mediterranean specimen is ornamented by costae, has curved sutures and is more rounded in outline. The aperture is characterized by irregular, bilobate, rimmed openings. We consider the Mediterranean species to be a separate taxon, that is tentatively regarded as *P. cf. P. antillarum*. As this represents the first record of this species, it is considered to be native to the Mediterranean Sea.

***Peneroplis arietinus* (Batsch, 1791)**

- 1791 *Nautilus arietinus* Batsch – Batsch, pl. 6, fig. 15c
 non 1791 *Nautilus arietinus* Batsch – Batsch, pl. 6, figs. 15a-b, 15d-f
 ? 1884 *Peneroplis arietinus* (Batsch) – Brady, p. 204, pl. 13, figs. 18-19
 non 1884 *Peneroplis arietinus* (Batsch) – Brady, p. 204, pl. 13, fig. 22
 1923 *Peneroplis arietinus* (Batsch) – Wiesner, p. 96, fig. 289
 1979 *Spirolina arietinus* (Batsch) – Blanc-Vernet *et al.*, pl. 24, fig. Sp a
 1994 *Peneroplis arietinus* (Batsch) – Gudmundsson, pl. 2, fig. 3, pl. 3, fig. 2
 non 2008a *Peneroplis arietinus* (Batsch) – Meriç *et al.*, pl. 9, figs. 14-16, pl. 10, figs. 1-5
 non 2017 *Spirolina arietina* (Batsch) – Lamourou *et al.*, fig. 7g
 2019 *Peneroplis arietinus* (Batsch) – Förderer & Langer, p. 13, pl. 2, figs. 6-12

Peneroplis arietinus has been described by Batsch (1791). The type locality is not known, but Cushman (1931) states that the collection site of the Batsch species is “from Rimini or at least the Adriatic Sea” based on the faunal content. The species resembles *Peneroplis planatus* and mainly differs from the latter in having fused pit rows that form deep grooves in between the ribs (see also Förderer & Langer, 2019). *Peneroplis arietinus* also resembles *Coscinospira hemprichii*, a species that is native to the Mediterranean Sea (discussed above). Several not illustrated records of *P. arietinus* (Sidebottom, 1904; Wiesner, 1911a; 1911b; 1913; Alavi, 1980; Davaud & Septfontaine, 1995) from the Mediterranean Sea may belong to *C. hemprichii* but the diagnostic features that separate both taxa require further study. Because of lower temperatures in the Mediterranean Sea, *C. hemprichii* masks its prominent morphological features, has lower calcification rates and resembles the sister genus *Peneroplis*. The test ornamentation and the pit row pattern of specimens illustrated by Gudmundsson (1994) resemble *C. hemprichii* as illustrated by Hottinger *et al.* (1993) and Förderer & Langer (2019). The illustrations provided by Meriç *et al.* (2008a) reveal apertural openings centered in

the terminal face and distinct pore pits between the longitudinal ribs. Both features are not present in true *Peneroplis arietinus*. The figures provided by Wiesner (1923) and Blanc-Vernet *et al.* (1979) do not provide sufficient resolution to identify the taxon as true *P. arietinus*. *Coscinospira arietina* has previously been reported to be alien to the Mediterranean Sea (Servello *et al.*, 2019), but the latter is in fact *Coscinospira hemprichii*, a species that is native to the Mediterranean (see above). As outlined above, the type locality of *P. arietinus* is believed to be in the Adriatic Sea (Cushman, 1931). We therefore consider *P. arietinus* to be native to the Mediterranean, but further studies are required.

***Planispirinella exigua* (Brady, 1879)**

- 1884 *Hauerina exigua* Brady – Brady, p. 196, pl. 12, figs. 1-4
 1954 *Planispirina exigua* (Brady) – Cushman *et al.*, p. 341, pl. 85, fig. 28
 1994 *Planispirinella exigua* (Brady) – Loeblich & Tappan, p. 38, pl. 57, figs. 7-8
 2006 *Planispirinella exigua* (Brady) – Oflaz, p. 141, pl. 1, figs. 10-11, pl. 11, fig. 3

The first description of *Planispirinella exigua* was provided by Brady (1879) from the Admiralty Islands and New Guinea. Not illustrated records from the Adriatic Sea include Wiesner (1911a) and Ćosović *et al.* (2011). Oflaz (2006) illustrates *P. exigua* from Turkey. This species is also present off the coast of Israel (Hyams-Kaphzan, unpublished, material collected in 1997). We therefore consider *P. exigua* to be alien to the Mediterranean Sea.

***Planogypsina acervalis* (Brady, 1884)**

- 1884 *Planorbulina acervalis* Brady – Brady, p. 657, pl. 92, fig. 4
 1909 *Planorbulina acervalis* Brady – Sidebottom, p. 2, pl. 1, fig. 4
 1949 *Planorbulina mediterraneensis* d'Orbigny – Said, p. 44, pl. 4, fig. 25
 non 1949 *Planorbulina acervalis* Brady – Said, p. 43, pl. 4, fig. 28
 1993 *Planogypsina acervalis* (Brady) – Hottinger *et al.*, p. 125, pl. 169, figs. 1-9, 170, figs. 1-8
 non 2006 *Planorbulina acervalis* (Brady) – Oflaz, pl. 8, fig. 16
 2008b *Planogypsina acervalis* (Brady) – Meriç *et al.*, pl. 8, fig. 9 (non figs. 4-5, 10-11)
 non 2010 *Planogypsina acervalis* (Brady) – Koukousioura *et al.*, p. 163, pl. 2, figs. 4-5
 2014 *Planogypsina acervalis* (Brady) – Meriç *et al.*, pl. 63, figs. 4a-b (non figs. 5-6)

Planogypsina acervalis has first been described by Brady (1884) from Booby Island, located in the Pacific Ocean (~ 14 m water depth). Said (1949) and Hottinger *et al.* (1993) reported the species from the northern Red

Sea. The first record from the Mediterranean Sea was provided by Sidebottom (1909, Island of Delos, Greece). Records of *P. acervalis* were given by Wiesner (1923) from Rimini (Adria) and by Moncharmont Zei (1968) from off Lebanon, but they all lack illustrations. Other Mediterranean records include Meriç *et al.* (2008b; 2014) and Hyams-Kaphzan (unpublished, material collected in 1998). The currently available evidence indicates that this species is alien to the Mediterranean Sea.

Planogypsina squamiformis (Chapman, 1901)

- 1901 *Gypsina vesicularis* (Parker & Jones) var. *squamiformis* Chapman – Chapman, p. 200, pl. 19, fig. 15
- 1901 *Gypsina vesicularis* (Parker & Jones) var. *monticulus* Chapman – Chapman, p. 200, pl. 19, fig. 14
- 1949 *Planorbulina acervalis* Brady – Said p. 43, pl. 4, fig. 28
- 1979 *Planorbulina* aff. *P. acervalis* Brady – Pereira, p. 288, pl. 41, figs. N-Q
- 1993 *Plangypsina squamiformis* (Chapman) – Hottinger *et al.*, p. 126, pl. 171, figs. 1-9
- non 2008a *Plangypsina squamiformis* (Chapman) – Meriç *et al.*, pl. 15, figs. 1-4
- non 2008b *Planogypsina squamiformis* (Chapman) – Meriç *et al.* pl. 8, figs. 12-15
- non 2009 *Planogypsina squamiformis* (Chapman) – Yokeş & Meriç pl. 4, figs. 14-15
- non 2014 *Planogypsina squamiformis* (Chapman) – Meriç *et al.* pl. 63, figs. 7-10

Planogypsina squamiformis was described by Chapman (1901) from Funafuti, Pacific Ocean. The species has also been recorded by Said (1949, Red Sea), Pereira (1979, Kenya) and Hottinger *et al.* (1993, Red Sea). The specimens illustrated by Meriç *et al.* (2008a; 2008b; 2014) and Yokeş & Meriç (2009) differ from the specimens recorded from the Pacific and the Indian Oceans and belong to *Planorbulina mediterraneensis*. To date, there is no reliable record for *Planogypsina squamiformis* in the Mediterranean Sea, and we therefore consider the species to be absent.

Procerolagena oceanica (Albani, 1974)

- 1974 *Lagena oceanica* Albani – Albani, p. 37, pl. 1, figs. 7, 10-11
- 1993 *Lagena oceanica* Albani – Hottinger *et al.*, p. 78, pl. 90, figs. 9-11

Procerolagena oceanica has first been described from Australia by Albani (1974). Mediterranean records are limited to specimens from the Mediterranean coast off Israel (*Lagena oceanica*: Hyams, 2006; Hyams-Kaphzan, 2016; not illustrated but reexamined). We consider *Procerolagena oceanica* to represent a species that is alien to the Mediterranean Sea.

Pseudohauerinella dissidens (McCulloch, 1977)

- 1977 *Pseudohauerina dissidens* McCulloch – McCulloch, p. 237, pl. 102, fig. 7
- 1993 *Pseudohauerinella dissidens* (McCulloch) – Hottinger *et al.*, p. 67, pl. 74, figs. 1-8
- 1998 *Pseudohauerinella dissidens* (McCulloch) – Piller & Haunold, pl. 6, fig. 15
- 2000 *Pseudohauerinella dissidens* (McCulloch) – Hyams, pl. 14, fig. 8
- 2017 *Pseudohauerinella dissidens* (McCulloch) – Thissen & Langer, pl. 9, figs. 15-18

The type locality of *Pseudohauerinella dissidens* is Sulphur Bay, Clarion Island (eastern Pacific, 31m depth; McCulloch, 1977). *Pseudohauerinella dissidens* has been recorded from the Red Sea by Hottinger *et al.* (1993) and Piller & Haunold (1998) and from Zanzibar by Thissen & Langer (2017). The first Mediterranean record was provided by Hyams (2000) from the coast off Israel. The Mediterranean representatives of *P. dissidens* are not as strongly plicated as their tropical counterparts. A reduction of test ornamental features is typical for several alien foraminifera that have migrated from warm tropical waters to the colder waters of the Mediterranean Sea. We consider this species to be alien to the Mediterranean Sea.

Pseudolachlanella slitella Langer, 1992

- 1980 *Quinqueloculina* “*laevigata*” d’Orbigny – Alavi, p. 19 (appendix), pl. 10, fig. 5, pl. 11, fig. 5
- 1992 *Pseudolachlanella slitella* Langer – Langer, p. 90, pl. 2, figs. 4-6
- 1993 “*Quinqueloculina*” *eburnea* d’Orbigny – Hottinger *et al.*, p. 59, pl. 53, figs. 9-11, pl. 54, figs. 1-5
- 2003 *Pseudolachlanella slitella* Langer – Samir *et al.*, pl. 3, figs. 3-5
- 2008 *Pseudolachlanella slitella* Langer – Langer, p. 406, fig. 7

Pseudolachlanella slitella has first been described by Langer (1992) from the Lagoon at Madang, Papua New Guinea. The species has also been recorded by Loeblich & Tappan (1994) from the Timor Sea, by Debenay (2012) from New Caledonia, and Hottinger *et al.* (1993) from the Red Sea (as “*Quinqueloculina*” *eburnea*). Mediterranean records include the studies by Alavi (1980, Turkey, as *Quinqueloculina* “*laevigata*”), Samir *et al.* (2003, Egypt) and Langer (2008, Gulf of Gabes). The species is rarely found along the coast off Israel (Hyams-Kaphzan, unpublished). *Pseudolachlanella slitella* was considered to be alien to the Mediterranean Sea by Langer (2008), but the first documented occurrence by Alavi (1980) shows that it was present in the Mediterranean Sea before its first record from Papua New Guinea, New Caledonia, and the Timor Sea. *Pseudolachlanella slitella* is therefore considered to be native to the Mediterranean Sea.

Pseudomassilina australis (Cushman, 1932)

- 1932 *Massilina australis* Cushman – Cushman, p. 32, pl. 8, figs. 2 a-b
non 1980 *Pseudomassilina* cf. *P. australis* (Cushman) – Alavi, pl. 19 (appendix), fig. 8
1993 *Pseudomassilina australis* (Cushman) – Hottinger *et al.*, p. 53, pl. 41, figs. 3-11
non 2000 *Pseudomassilina* cf. *P. australis* (Cushman) – Hyams, pl. 11, fig. 1
2006 *Pseudomassilina australis* (Cushman) – Oflaz, p. 171, pl. 3, fig. 10

The holotype of *Pseudomassilina australis* was described by Cushman (1932) from Rarotonga, Cook Islands. Hottinger *et al.* (1993) recorded *P. australis* from the Gulf of Aqaba, Red Sea. The specimen illustrated by Oflaz (2006) from Turkey shares all morphological features with the holotype but appears to be a semi-adult individual. Similar specimens were recorded by Alavi (1980) from the Cilicia Basin, Turkey, and Hyams (2000) from the coast off Israel, but they show strong transverse ribs and were therefore assigned to *Pseudomassilina* cf. *P. australis*. For the time being we consider the specimen illustrated by Oflaz (2006) to be identical with the holotype and *Pseudomassilina australis* to be alien to the Mediterranean Sea.

Pseudomassilina reticulata (Heron-Allen & Earland, 1915)

- 1915 *Miliolina* (*Massilina*) *secans*, var. *reticulata* Heron-Allen & Earland – Heron-Allen & Earland, p. 582, pl. 45, figs. 1-4
1987 *Pseudomassilina australis* (Cushman) subsp. *reticulata* (Heron-Allen & Earland) – Baccaert, p. 111, pl. 51, fig. 2
1993 *Pseudomassilina reticulata* (Heron-Allen & Earland) – Hottinger *et al.*, p. 54, pl. 42, figs. 5-8, pl. 43, figs. 1-8
2000 *Pseudomassilina reticulata* (Heron-Allen & Earland) – Hyams, pl. 11, figs. 2-3
2008a *Pseudomassilina reticulata* (Heron-Allen & Earland) – Meriç *et al.*, pl. 7, fig. 4
2008b *Pseudomassilina reticulata* (Heron-Allen & Earland) – Meriç *et al.*, pl. 4, fig. 6
2009 *Pseudomassilina reticulata* (Heron-Allen & Earland) – Yokeş & Meriç, pl. 2, fig. 6
2014 *Pseudomassilina reticulata* (Heron-Allen & Earland) – Meriç *et al.*, pl. 27, fig. 16

Heron-Allen & Earland (1915) first described *Pseudomassilina reticulata* from several localities in the Quirimbas Archipelago off northern Mozambique. Their illustrations (figs. 1-2) show a distinct reticulate surface pattern as a characteristic feature of this taxon. The species has been reported from Zanzibar (Thissen & Langer, 2017) and the Red Sea by Hottinger *et al.* (1993). Mediterranean records include Hyams (2000), Avsar *et al.*, 2001 (not illustrated), Meriç *et al.* (2008a; 2008b), Yokeş & Meriç (2009) and

Meriç *et al.* (2014). In the Mediterranean Sea, the test reticulation appears to be less pronounced, a morphological feature that is typical for species that have migrated from warm tropical waters to the colder Mediterranean. Hottinger *et al.* (1993) consider test reticulation to be variable. We therefore consider the specimens illustrated by Hyams (2000) and Meriç (2008a) to be identical with the holotype of *Pseudomassilina reticulata* and the species to be alien to the Mediterranean Sea.

Pseudoschlumbergerina ovata (Sidebottom, 1904)

- 1904 *Sigmoilina ovata* Sidebottom – Sidebottom, p. 6, fig. 1
1968 *Sigmoilina ovata* Sidebottom – Moncharmont Zei, p. 27, pl. 2, fig. 11
1980 “*Sigmoilina*” *ovata* Sidebottom – Alavi, p. 25 (appendix), pl. 11, fig. 12, pl. 17, fig. 11
1990 *Septloculina rotunda* El-Nakhal – El-Nakhal, p. 91, pl. 1, figs. 8-11, pl. 2, fig. 13
1993 *Pseudoschlumbergerina ovata* (Sidebottom) – Hottinger *et al.*, p. 55, pl. 46, figs. 1-6
2000 *Pseudoschlumbergerina ovata* (Sidebottom) – Hyams, pl. 11, figs. 5-6
2006 *Septloculina rotunda* El-Nakhal – Oflaz, p. 164, pl. 3, fig. 5
2012 *Pseudoschlumbergerina ovata* (Sidebottom) – Milker & Schmiedl, p. 68, pl. 17, figs. 29-30
2017 *Pseudoschlumbergerina ovata* (Sidebottom) – Guy-Haim *et al.*, pl. 4, fig. 19, pl. 2, fig. 15

Septloculina rotunda is listed as alien to the Mediterranean in EASIN (2020). The species was described by El-Nakhal (1990) from the Red Sea but is a junior synonym of *Pseudoschlumbergerina ovata* (Sidebottom, 1904), a species that was first recorded from the Mediterranean. *Septloculina rotunda* has been considered to be alien to the Mediterranean Sea by Oflaz (2006). It has been recorded by Hyams-Kaphzan *et al.* (2008, 2014) as *Pseudoschlumbergerina ovata*. Marriner *et al.* (2005), Milker & Schmiedl (2012) and Hyams-Kaphzan (unpublished) recorded fossil occurrences from the Mediterranean Sea. *Pseudoschlumbergerina ovata* is therefore native to the Mediterranean Sea. Guy-Haim *et al.* (2017) found living specimens of *Pseudoschlumbergerina ovata* in fecal pellets of herbivorous rabbitfish collected in the Suez Canal, indicating that present-day species introductions via ichthyochory vectors are ongoing.

Pseudotriloculina subgranulata (Cushman, 1918)

- 1918 *Triloculina subgranulata* Cushman – Cushman, p. 290, pl. 96, figs. 4a-c
1993 *Pseudotriloculina subgranulata* (Cushman) – Hottinger *et al.*, p. 56, pl. 47, figs. 8-13, pl. 48, figs. 1-8
? 1994 *Triloculina subgranulata* Cushman – Reinhardt *et al.*, pl. 1, figs. 12-13
1998 *Pseudotriloculina subgranulata* (Cushman)

- Yanko *et al.*, pl. 6, figs. 9-11
 2000 *Pseudotriloculina subgranulata* (Cushman)
 – Hyams, pl. 12, fig. 6
 non 2017 *Pseudotriloculina subgranulata* (Cushman)
 – Makled *et al.*, pl. 12, fig. 12

Pseudotriloculina subgranulata has been described by Cushman (1918, as *Triloculina subgranulata*) from the Tortugas (Florida). The species is characterized by its rough test surface, resulting from distinct microstriae. *Pseudotriloculina subgranulata* was also reported from the coast of Israel by Yanko *et al.* (1998) and Hyams (2000). Other records from the Mediterranean Sea include Moncharmont Zei (1968) and Hyams-Kaphzan *et al.* (2008), but lack illustrations. The fossil specimen reported by Reinhardt *et al.* (1994) requires further study. The species was recently recorded in core material from the Mediterranean coast off Israel (Hyams-Kaphzan, unpublished). *Pseudotriloculina subgranulata* is therefore native to the Mediterranean Sea.

***Pyrgo denticulata* (Brady, 1884)**

- 1884 *Biloculina ringens* var. *denticulata* Brady –
 Brady, p. 143, pl. 3, figs. 4-5
 1950 *Pyrgo denticulata* (Brady) – Said, p. 7, pl.
 1, fig. 15
 1993 *Pyrgo denticulata* (Brady) – Hottinger *et*
al., p. 56, pl. 49, figs. 8-12
 2000 *Pyrgo denticulata* (Brady) – Hyams, pl. 12,
 fig. 2
 2008a *Pyrgo denticulata* (Brady) – Meriç *et al.*, pl.
 7, fig. 10
 2010 *Pyrgo denticulata* (Brady) – Avnaim-Katav,
 p. 224, pl. 4, fig. 7

Pyrgo denticulata has been reported from Honolulu and Tongatapu (Tonga) by Brady (1884), from the Malay Archipelago (Millett, 1898), Funafuti (Chapman, 1901), the Kerimba Archipelago (Heron-Allen & Earland, 1915), Jamaica (Cushman, 1929) and the Red Sea (Said, 1950; Hottinger *et al.*, 1993). The first occurrence in the Mediterranean Sea was provided by Hyams (2000) from off Israel. A single figure is provided by Meriç *et al.* (2007; 2008a; 2008b; 2014) and Yokeş & Meriç (2009), a figure that is identical in all four publications. Avnaim-Katav (2010) recorded fossil occurrences from the Mediterranean Sea, Haifa Bay, Israel (HB30, 1.95 m, Israel). However, dating of the core material is not clear-cut and requires further study. For the time being we consider *Pyrgo denticulata* to be alien to the Mediterranean Sea.

***Quinqueloculina* cf. *Q. mosharrafai* Said, 1949**

- 1949 cf. *Quinqueloculina mosharrafai* Said –
 Said, p. 10, pl. 1, fig. 23
 1993 *Quinqueloculina* cf. *Q. mosharrafai* Said –
 Hottinger *et al.*, p. 59, pl. 54, figs. 6-9, pl.
 55, figs. 1-6

- 2008a *Quinqueloculina* cf. *Q. mosharrafai* Said –
 Meriç *et al.*, pl. 6, fig. 12
 2009 *Quinqueloculina* cf. *Q. mosharrafai* Said –
 Yokeş & Meriç, pl. 2, fig. 4
 2014 *Quinqueloculina* cf. *Q. mosharrafai* Said –
 Meriç *et al.*, pl. 24, fig. 8
 2014 *Quinqueloculina* cf. *Q. mosharrafai* Said –
 Hyams-Kaphzan *et al.*, p. 13

Quinqueloculina mosharrafai was first described from the Red Sea by Said (1949). Hottinger *et al.* (1993) illustrate a similar species (*Quinqueloculina* cf. *Q. mosharrafai*) from the Red Sea that differs from the holotype in having a bifid tooth and a partially agglutinated test surface. The holotype illustrated by the Smithsonian National Museum of Natural History (2020) shows agglutinated particles on the test surface but lacks the bifid tooth. The specimen described by Meriç *et al.* (2008a; 2014) and Yokeş & Meriç (2009) from Turkey is identical to *Quinqueloculina* cf. *Q. mosharrafai* from the Red Sea (Hottinger *et al.*, 1993). *Quinqueloculina* cf. *Q. mosharrafai* has also been reported from the coast of Israel (Hyams-Kaphzan *et al.*, 2014, material collected in 2013) and is here considered to be alien to the Mediterranean Sea.

***Quinqueloculina* cf. *Q. multimarginata* Said, 1949**

- 1949 cf. *Quinqueloculina multimarginata* Said –
 Said, p. 10, pl. 1, fig. 34
 1993 *Quinqueloculina* cf. *Q. multimarginata* Said –
 Hottinger *et al.*, p. 59, pl. 55, figs. 7-10
 2000 *Quinqueloculina* cf. *Q. multimarginata* Said –
 Hyams, pl. 7, fig. 9
 2018 *Quinqueloculina* cf. *Q. multimarginata* Said –
 Förderer & Langer, pl. 23, figs. 28-30

The holotype of *Quinqueloculina multimarginata* described by Said (1949) from the Red Sea has thin keels and irregular costae on parts of the test surface. *Quinqueloculina* cf. *Q. multimarginata* described by Hottinger *et al.* (1993) differs from Said's illustration by having an angular periphery and a rounded carina. In addition, the test surface is covered with distinct anastomosing microstriations. The Mediterranean specimen illustrated by Hyams (2000) represents a morphological variety of *Q.* cf. *Q. multimarginata* and has a slightly smoother surface than the specimens illustrated by Hottinger *et al.* (1993, Gulf of Aqaba) and Förderer & Langer (2018, Indonesia). For the time being, we consider *Quinqueloculina* cf. *Q. multimarginata* to be alien to the Mediterranean Sea.

***Schlumbergerina alveoliniformis* (Brady, 1879)**

- 1879 *Miliolina alveoliniformis* Brady – Brady, p. 268
 1884 *Miliolina alveoliniformis* Brady – Brady, p.
 181, pl. 8, figs. 15-20
 1987 *Schlumbergerina alveoliniformis* (Brady)
 – Baccaert, p. 150, pl. 65, figs. 4-5, pl. 66,
 figs. 1a-c

- 1993 *Schlumbergerina alveoliniformis* (Brady) – Hottinger *et al.*, p. 61, pl. 58, figs. 11-14, pl. 59, figs. 1-9
- 2008b *Schlumbergerina alveoliniformis* (Brady) – Meriç *et al.*, p. 331, pl. 3, figs. 8-11
- 2009 *Schlumbergerina alveoliniformis* (Brady) – Yokeş & Meriç, pl. 1, figs. 16-17
- 2014 *Schlumbergerina alveoliniformis* (Brady) – Meriç *et al.*, pl. 24, fig. 8
- non 1990 *Miliolinella subrotunda* (Montagu) – Dermitzakis & Triantafyllou, p. 150, pl. 3, fig. 5
- 2005 *Miliolinella subrotunda* (Montagu) – Triantaphyllou *et al.*, pl. 1, fig. 3
- 2006 *Miliolinella subrotunda* (Montagu) – Langer & Schmidt-Sinns, pl. 8, figs. 8-9
- 2010 *Sigmamiliolinella australis* (Parr) – Avnaim-Katav, pl. 4, fig. 18
- 2011 *Miliolinella subrotunda* (Montagu) – Koukousioura *et al.*, pl. 1, fig. 7
- 2013a *Miliolinella australis* (Parr) – Langer *et al.*, pl. 5, figs. 40-41
- non 2014 *Miliolinella australis* (Parr) – Meriç *et al.*, pl. 25, fig. 21
- 2017 *Miliolinella australis* (Parr) – Fajemila and Langer, figs. 6, 30, 31
- 2017 *Sigmamiliolinella australis* (Parr) – Mouanga, pl. 8, figs. 15a-b
- 2018 *Sigmamiliolinella australis* (Parr) – Förderer & Langer, p. 91, pl. 18, figs. 25-30

Schlumbergerina alveoliniformis has been described by Brady (1879) from the Admiralty Islands in the Pacific Ocean with occurrences in Honolulu and Tongatabu. Mediterranean specimens illustrated by Meriç *et al.* (2008b; 2014) and Yokeş & Meriç (2009) represent the only Mediterranean records. We consider the species to be alien to the Mediterranean Sea.

***Septloculina angulata* El-Nakhal, 1990**

- 1904 *Sigmoilina ovata* Sidebottom – Sidebottom, p. 6 (pars), text-fig. 1 (non pl. 2, figs. 12-13)
- 1990 *Septloculina angulata* El-Nakhal – El-Nakhal, p. 91, pl. 1, figs. 1-7
- non 2006 *Septloculina angulata* El-Nakhal – Oflaz, p. 164, pl. 3, fig. 4

The species *Septloculina angulata* was described by El-Nakhal (1990) from the littoral zone off Tartus Town, Syria (Mediterranean Sea). *Septloculina angulata* is a species with seven, externally visible chambers, a subangular thickened periphery, a rough surface, and a semicircular terminal aperture provided with a bifid tooth. The species illustrated by Oflaz has less chambers, lacks the thickened chamber periphery and probably belongs to a different genus. The species was first described from the eastern Mediterranean is therefore native to the Mediterranean Sea.

***Septloculina tortuosa* El-Nakhal, 1990**

- 1990 *Septloculina tortuosa* El-Nakhal – El-Nakhal, p. 91, pl. 2, fig. 4-9
- non 2006 *Septloculina tortuosa* El-Nakhal – Oflaz, p. 165, pl. 3, fig. 6

The species *Septloculina tortuosa* has been described by El-Nakhal from Alexandria (1990, Egypt, Mediterranean Sea) and is therefore native to the Mediterranean Sea. The specimen reported by Oflaz (2006) from Iskenderun Bay, Turkey shows a quinqueloculine chamber arrangement and belongs to a different genus.

***Sigmamiliolinella australis* (Parr, 1932)**

- 1932 *Quinqueloculina australis* Parr – Parr, pl. 1, figs. 8a-c
- 1987 *Miliolinella australis* (Parr) – Baccaert, p. 138, pl. 60, figs. 6-8

The type locality for *Sigmamiliolinella australis* is east of Cape Pillar, Tasmania (Parr, 1932). Baccaert (1987) noted that the species is typical for tropical and temperate waters of “Indopacific shallow-water localities exclusively”. *Sigmamiliolinella australis* is not known from the northern Red Sea (Said, 1949; Hottinger *et al.*, 1993; Piller & Haunold, 1998). A typical feature of *S. australis* is the partially coarse test surface at the proximal part of the shell. Some specimens of *S. australis* from the Mediterranean have been assigned to *Miliolinella subrotunda*. The first Mediterranean reference of *Miliolinella australis* was provided by Alavi (1980), p. 90. The first illustrated record of *Sigmamiliolinella australis* in the Mediterranean was given by Triantaphyllou *et al.* (2005, identified as *Miliolinella subrotunda*). Other records include material from Greece (Koukousioura *et al.*, 2011), Elba Island, Italy (Langer & Schmidt-Sinns, 2006), Dhermi, Albania (Mouanga, 2017), Corfu (Langer, unpublished; material collected in 2014) and Israel (Avnaim-Katav, 2010 core-top, material collected in 2005; Hyams-Kaphzan *et al.*, 2014, material collected in 2013). The specimen illustrated in Dermitzakis & Triantafyllou (1990) has inflated chambers, a smooth surface, lacks the typical thickened apertural rim and thus differs from *Sigmamiliolinella australis* (Parr). Recently re-illustrated material from the same material (Triantaphyllou, unpublished) has a similar test surface but shows a specimen with an oblique middle chamber and small aperture that also lacks the typical thickened apertural rim. To date, the species has not yet been found in core material. *Sigmamiliolinella australis* is currently abundant on macroalgae at sites off Haifa (Hyams-Kaphzan, unpublished) and appears to be a recent arrival in the southeastern Mediterranean Sea. We consider *Sigmamiliolinella australis* to be alien to the Mediterranean Sea.

***Sigmoihauerina bradyi* (Cushman, 1917)**

- 1884 *Hauerina compressa* d'Orbigny – Brady, pl. 11, figs. 12-13 (*non H. compressa* d'Orbigny)
- non 1904 *Hauerina compressa* d'Orbigny – Sidebottom, p. 19, no. 20
- 1917 *Hauerina bradyi* Cushman – Cushman, p. 62, pl. 23, figs. 2a-b
- 1949 *Hauerina bradyi* Cushman – Said, p. 17, pl. 2, fig. 5
- non 1968 *Hauerina bradyi* Cushman – Moncharmont Zei, p. 14, pl. 3, fig. 7
- 1993 *Hauerina bradyi* Cushman – Hottinger *et al.*, p. 62, pl. 60, figs. 1-12
- non 2006 *Sigmoihauerina bradyi* (Cushman) – Oflaz, p. 170, pl. 3, fig. 9

Brady (1884) provided the original illustrations for *Sigmoihauerina bradyi* (Cushman) from material collected in the Torres Strait and at Booby Island (Pacific Ocean). The species is characterized by its compressed test and thus differs from the more inflated species *Hauerina compressa* of d'Orbigny (1846) from Miocene deposits of the Vienna Basin. The Mediterranean records provided by Sidebottom (1904, as *Hauerina compressa*), Moncharmont Zei (1968), and Oflaz (2006) show the inflated form and that does not match the holotype or the specimens illustrated by Hottinger *et al.* (1993). Specimens with the typical large and compressed test are not yet known from the Mediterranean Sea and we therefore consider *Sigmoihauerina bradyi* to be absent from this region.

***Siphonaperta cf. S. pittensis* (Albani, 1974)**

- non 1957 *Quinqueloculina parvagggluta* Vella – Vella, p. 27, pl. 4, figs. 71-73
- 1974 *cf. Quinqueloculina pittensis* Albani – Albani, p. 33, pl. 1, figs. 1-3
- non 1993 *Siphonaperta pittensis* (Albani) – Hottinger *et al.*, p. 63, pl. 64, figs. 1-6
- 2000 *Siphonaperta cf. S. pittensis* (Albani) – Hyams, pl. 9, fig. 6
- non 2010 *Quinqueloculina parvagggluta* Vella – Hayward *et al.*, p. 154, pl. 5, figs. 1-2
- non 2012 *Quinqueloculina parvagggluta* Vella – Debenay, p. 125
- non 2012 *Quinqueloculina pittensis* (Albani) – Debenay, p. 125
- non 2013a *Siphonaperta pittensis* (Albani) – Langer *et al.*, pl. 4, figs. 33-34

Siphonaperta pittensis was considered to be a Lessepsian invader by Hyams-Kaphzan *et al.* (2008). The Mediterranean records are now considered to constitute a separate species. *Siphonaperta cf. S. pittensis* is more elongated than *S. pittensis* and its chambers are less inflated (see also Hyams, 2000). *Siphonaperta cf. S. pittensis* also resembles *Quinqueloculina parvagggluta* as described by Vella (1957). The relationship between

Siphonaperta pittensis and *Quinqueloculina parvagggluta* remains uncertain. Specimens of *Siphonaperta cf. S. pittensis*, as illustrated by Hyams (2000), were recorded from the Mediterranean coast off Israel. As this constitutes the only record, we consider the species to be native to the Mediterranean Sea.

***Siphonaperta distortaqueata* (Cushman, 1954)**

- 1954 *Quinqueloculina distortaqueata* Cushman – Cushman *et al.*, p. 333, pl. 83, fig. 27
- 1980 *Quinqueloculina berthelotiana* d'Orbigny – Alavi, pl. 15, fig. 2
- 1993 *Siphonaperta distortaqueata* (Cushman) – Hottinger *et al.*, p. 63, pl. 62, figs. 4-9, pl. 63, figs. 1-6
- 2000 *Siphonaperta distortaqueata* (Cushman) – Hyams, pl. 9, figs. 2-3
- 2009 *Quinqueloculina distortaqueata* Cushman – Parker, p. 195, figs. 138-139
- 2011 *Siphonaperta distortaqueata* (Cushman) – Makled & Langer, fig. 7, 7-9
- 2017 *Siphonaperta distortaqueata* (Cushman) – Thissen & Langer, pl. 7, figs. 24-26
- 2017 *Quinqueloculina distortaqueata* Cushman – Fajemila & Langer, fig. 4, 30-32

Siphonaperta distortaqueata has been described from the Marshall Islands by Cushman *et al.* (1954). The species has been recorded from the Red Sea by Hottinger *et al.* (1993), from around Australia by Parker (2009, and references therein), from the Chuuk Atoll (western Pacific) by Makled & Langer (2011), from the Atlantic Ocean (Fajemila & Langer, 2017) and from Zanzibar by Thissen & Langer (2017). Mediterranean records include specimens from Turkey (Alavi, 1980, as *Quinqueloculina berthelotiana*) and from shallow waters off Israel (Hyams, 2000; Hyams-Kaphzan *et al.*, 2008; 2014). The species is here considered to be alien to the Mediterranean Sea.

***Sorites variabilis* Lacroix, 1941**

- 1941 *Sorites variabilis* Lacroix – Lacroix. (pars), p. 14
- 1941 *Sorites variabilis* Lacroix – Lacroix., p. 11 (pars), figs. 12, 18
- ? 1979 *Sorites variabilis* Lacroix – Blanc-Vernet *et al.*, pl. 21, fig. 14, pl. 24, fig. S
- ? 1993 *Sorites variabilis* Lacroix – Hottinger *et al.*, p. 73, pl. 84, figs. 1-15
- ? 2008a *Sorites variabilis* Lacroix – Meriç *et al.*, pl. 12, figs. 12-15, pl. 13, figs. 1-4
- ? 2008b *Sorites variabilis* Lacroix – Meriç *et al.*, pl. 7, figs. 1-8
- 2013 *Sorites variabilis* Lacroix – Merkado *et al.*, pl. 7, figs. 1-2
- ? 2016 *Sorites variabilis* Lacroix – Ayadi *et al.*, pl. 1, fig. 9, pl. 3, figs. 2-3
- non 2017 *Sorites variabilis* Lacroix – Lamourou *et al.*, fig. 7(h)

Sorites variabilis has been first described by Lacroix (1941) from the Gulf of Aqaba, Red Sea. Lacroix described three types of soritid foraminifera. Only type II has been designated to represent *S. variabilis* (see also Hottinger *et al.*, 1993). The Mediterranean records from Turkey (Meriç *et al.*, 2008a; 2008b) do not provide illustrations of the apertural features and their true identity requires further study. The specimens from Israel (Hyams-Kaphzan *et al.*, 2008, not illustrated) and from Villefranche (Langer, unpublished; material collected in 1996) were reexamined by us, show the typical morphological features of *Sorites variabilis* (particularly thin-shelled, single row of apertures). All *Sorites* specimens described or illustrated from Naxos (Greece, Cherif, 1970) or from along the coast of Tunisia (Glacon, 1962; Blanc-Vernet *et al.*, 1979; Langer, unpublished) are thick-shelled, show the typical features of *Sorites orbiculus*, and do not belong to *Sorites variabilis* (see also Baccaert, 1987). For the time being, this species is considered to be alien to the Mediterranean, but further molecular studies are required.

Spiroloculina angulata Cushman, 1917

- 1917 *Spiroloculina grata* Terquem var. *angulata* Cushman – Cushman, p. 36, pl. 7, fig. 5;
 1968 *Spiroloculina grata* Terquem var. *angulata* Cushman – Moncharmont Zei, pl. 3, fig. 10
 non 2008 *Spiroloculina angulata* Cushman – Yalçın *et al.*, pl. 1, fig. 8
 non 2013 *Spiroloculina angulata* Cushman – Cosentino *et al.*, p. 8786, pl. 1, fig. 18

The type species of *Spiroloculina angulata* has been described by Cushman (1917) from the northern Pacific. The species is characterized by its truncated periphery and a characteristic median carina. A single illustrated record of this species exists from the coast of Lebanon (Moncharmont Zei, 1968). Other records of *Spiroloculina angulata* include Alavi (1980), Morhange *et al.* (2000), Basso & Spezzaferri (2000), Samir & El-Din (2001), Avşar *et al.* (2001), Meriç *et al.* (2004), Zaïbi *et al.* (2016), Altınışlı *et al.* (2017, Sea of Marmara), but they all lack illustrations. For the time being, *Spiroloculina angulata* is considered to be alien to the Mediterranean Sea.

Spiroloculina antillarum d'Orbigny, 1839

- 1839 *Spiroloculina antillarum* d'Orbigny – d'Orbigny, p. 166, pl. 9, figs. 3-4
 1920 *Spiroloculina antillarum* d'Orbigny – Martinotti, p. 261, figs. 16-17
 1923 *Spiroloculina antillarum* d'Orbigny – Wiesner, p. 33, pl. 4, fig. 20
 non 1970 *Spiroloculina antillarum* d'Orbigny – Cherif, p. 37, pl. 3, fig. 3
 1977 *Spiroloculina antillarum* d'Orbigny – Le Calvez, p. 91, pl. 17, figs. 1-6
 non 1979 *Spiroloculina antillarum* d'Orbigny –

- Blanc-Vernet *et al.*, pl. 24, fig. Sa
 non 1993 *Spiroloculina antillarum* d'Orbigny – Hottinger *et al.*, p. 45, pl. 24, figs. 15-17, pl. 25, figs. 1-2
 non 2008a *Spiroloculina antillarum* d'Orbigny – Meriç *et al.*, pl. 3, figs. 5-11
 non 2008b *Spiroloculina antillarum* d'Orbigny – Meriç *et al.*, pl. 2, fig. 19, pl. 3, figs. 1-4, 6-7 (non fig. 5)
 non 2010 *Spiroloculina antillarum* d'Orbigny – Avnaim-Katav, p. 220, pl. 2, fig. 2
 non 2017 *Spiroloculina antillarum* d'Orbigny – Mouanga, p. 175, pl. 1, figs. 3a-b
 non 2017 *Spiroloculina antillarum* d'Orbigny – Thissen & Langer, pl. 3, figs. 7-10
 non 2018 *Spiroloculina antillarum* d'Orbigny – Emter, figs. 19, 21-22

Spiroloculina antillarum was first recorded from Cuba by d'Orbigny (1839), is elongate fusiform in lateral view and possess a densely costate test surface and long neck (see Le Calvez, 1977). The first record from the Mediterranean was provided by Wiesner (1913, not illustrated). The specimen was later illustrated by Martinotti (1920, Libya), Wiesner (1923, Adriatic Sea). Records without illustrations include Moncharmont-Zei (1968, Lebanon), Samir *et al.* (2003, Egypt), Mkawar *et al.* (2007, Tunisia), and Serandrei-Barbero *et al.* (2011, Adriatic Sea). We consider *Spiroloculina antillarum* to be alien in the Mediterranean Sea. The specimens illustrated by Hottinger *et al.* (1993), Meriç *et al.* (2008 a,b), Avnaim-Katav (2010), Mouanga (2017), Thissen & Langer (2017) and Emter (2018) are coarsely costate, broadly fusiform, and have a short neck and thus differ from the original of d'Orbigny and represent a separate species. The Red Sea/Indian Ocean type (e.g. Hottinger *et al.*, 1993) was also reported from fossil sediments by Mateu (1972), Avnaim-Katav (2010), Melis *et al.* (2015), Yümün *et al.* (2016) and Hyams-Kaphzan (unpublished) and is native to the Mediterranean Sea.

Spiroloculina attenuata Cushman & Todd, 1944

- 1944 *Spiroloculina attenuata* Cushman & Todd – Cushman & Todd, p. 54, pl. 9, figs. 23-25
 1987 *Spiroloculina communis* Cushman & Todd subsp. *attenuata* – Baccaert, p. 118, pl. 53, figs. 4-5
 1993 *Spiroloculina attenuata* Cushman & Todd – Hottinger *et al.*, p. 45, pl. 25, figs. 3-9
 2000 *Spiroloculina attenuata* Cushman & Todd – Hyams, pl. 4, fig. 2

Spiroloculina attenuata was described by Cushman and Todd (1944) with the type locality near Pago Pago Harbor, Samoa (Pacific Ocean). The WoRMS Editorial Board (2020) has recently placed the taxon into *Naxotia* (Al-Zamel & Cherif, 1997), a new genus that was introduced for evolute quinqueloculine forms, with five chambers visible externally, tests with a greasy luster and a necked

circular aperture with two teeth. *Spiroloculina attenuata* does not fulfil the range of features required to be placed in *Naxotia*. Hyams (2000) recorded this species from the Mediterranean coast of Israel. We consider *Spiroloculina attenuata* to be alien to the Mediterranean Sea.

Spiroloculina* aff. *S. communis Cushman & Todd, 1944

1944 *Spiroloculina communis* Cushman & Todd – Cushman & Todd, p. 63, pl. 9, figs. 4-5, 7-8

? 1949 *Spiroloculina communis* Cushman & Todd – Said, p. 14, pl. 1, fig. 37

1993 *Spiroloculina* aff. *S. communis* Cushman & Todd – Hottinger *et al.*, p. 45, pl. 25, figs. 10-15

The holotype of *Spiroloculina communis* recorded by Cushman & Todd (1944) originates from the San Andres Island in the Philippines. *Spiroloculina communis* was later illustrated by Said (1949) from the Red Sea. However, the quality of the illustration is poor. *Spiroloculina* aff. *S. communis* was described from the Red Sea by Hottinger *et al.* (1993). This species resembles *S. communis* but possesses only one bifid tooth. Recent material from the Mediterranean Sea for *Spiroloculina* aff. *S. communis* has been recorded by Hyams-Kaphzan *et al.* (2008). We therefore consider the species to be alien to the Mediterranean Sea.

Spiroloculina nummiformis Said, 1949

1949 *Spiroloculina nummiformis* Said – Said, p. 16, pl. 1, fig. 39

1993 *Spiroloculina nummiformis* Said – Hottinger *et al.*, p. 46, pl. 27, figs. 1-9

2000 *Spiroloculina nummiformis* Said – Hyams, pl. 4, fig. 6

non 2011 *Spiroloculina nummiformis* Said – Elshanawany *et al.*, pl. 7, fig. 8

Spiroloculina nummiformis has been described from the Red Sea by Said (1949). Mediterranean occurrences include two records from the Levantine coast off Israel (Hyams, 2000; Hyams-Kaphzan *et al.*, 2008). The specimen illustrated by Elshanawany *et al.* (2011) from Abu-Qir Bay off Alexandria (Egypt) differs substantially from the holotype and the specimens illustrated by Hottinger *et al.* (1993) from the Red Sea. *Spiroloculina nummiformis* is considered to be alien to the Mediterranean Sea.

Textularia agglutinans d’Orbigny, 1839

1839 *Textularia agglutinans* d’Orbigny – d’Orbigny, pl. 1, figs. 17, 18, 32–34

1884 *Textularia agglutinans* d’Orbigny – Brady, pl. 43 figs. 1–2

1977 *Textularia agglutinans* d’Orbigny – Le Calvez, p. 13–14, fig. 1

1981 *Textularia agglutinans* d’Orbigny – Banner

& Pereira, pl. 1, figs. 6–7, pl. 2, fig. 1

Textularia agglutinans d’Orbigny – Dermitzakis & Triantafyllou, p. 161, pl. 3, fig. 6

1991 *Textularia agglutinans* d’Orbigny – Cimerman & Langer, pl. 10, figs. 1–2

1993 *Textularia agglutinans* d’Orbigny – Hottinger *et al.*, pl. 13, figs. 1-9

2012 *Textularia agglutinans* d’Orbigny – Milker & Schmiedl, fig. 10, 15-16

2015 *Textularia agglutinans* d’Orbigny – Merkado *et al.*, fig. 4, 1-4; fig. 5, 1-9

2020 *Textularia agglutinans* d’Orbigny – Manda *et al.*, fig. 3c

Textularia agglutinans d’Orbigny is a cosmopolitan agglutinated foraminiferal species, with an elongated biserial test in its adult stage and a low arched aperture. It was first described by d’Orbigny in 1839 from sandy beaches around Cuba and has been reported from numerous locations worldwide: Atlantic Ocean (Culver & Buzas, 1980), Red Sea (Hottinger *et al.*, 1993), Timor Sea (Loeblich & Tappan, 1994), Pacific Ocean (Debenay, 2012), Indian Ocean (Langer *et al.*, 2013a), western and eastern Mediterranean (Cimerman & Langer, 1991; Hyams-Kaphzan *et al.*, 2008, p. 336; Milker & Schmiedl, 2012), Adriatic Sea (Jorissen, 1987), Tyrrhenian Sea (Sgarrella & Moncharmont Zei, 1993) and Marmara Sea (Armynot du Châtelet *et al.*, 2013). Recent molecular analyses (Merkado *et al.*, 2015) indicate that the northern Red Sea and the eastern Mediterranean hard-bottom populations of *T. agglutinans* off Israel belong to the same genetic population, regardless of their large morphological variability. Further molecular studies are required to validate if the Mediterranean species are indeed genetically identical to the material from the Caribbean type locality. Fossil *T. agglutinans* material was recorded from Holocene deposits by Milker & Schmiedl (2012) from the western Mediterranean Sea (off Mallorca, Alboran Platform and Oran Bight) and from the Gulf of Corinth (Dermitzakis & Triantafyllou, 1990). The species is also abundant in core material (Hyams-Kaphzan, unpublished) collected off Israel. Fossil evidence and the global biogeographic distribution makes it difficult to consider *T. agglutinans* as alien to the Mediterranean Sea. The species is therefore considered to be native.

Triloculina asymmetrica Said, 1949

1949 *Triloculina asymmetrica* Said – Said, p. 18, pl. 2, fig. 11

1993 *Triloculina asymmetrica* Said – Hottinger *et al.*, p. 64, pl. 66, figs. 4-9

non 2017 *Triloculina asymmetrica* Said – Mouanga, pl. 9, figs. 14-15

The holotype of *Triloculina asymmetrica* was described by Said (1949) from the Red Sea. Hottinger *et al.* (1993) recorded it from the Gulf of Aqaba, Red Sea. The record by Oflaz (2006) lacks an illustration. The pho-

tographs provided by Mouanga (2017) from Albania do not show the typical asymmetrical test shape. *Triloculina asymmetrica* has not yet been found in the Mediterranean Sea, is therefore absent and not alien.

***Triloculina* cf. *T. fichteliana* d'Orbigny, 1839**

- non 1839 *Triloculina fichteliana* d'Orbigny – d'Orbigny, p. 171, pl. 9, figs. 8, 10
- non 1929 *Triloculina fichteliana* d'Orbigny – Cushman, p. 63, pl. 17, figs. 1a-c
- non 1977 *Triloculina fichteliana* d'Orbigny – Le Calvez, p. 106, fig. 1
- 1993 *Triloculina fichteliana* d'Orbigny – Hottinger *et al.*, p. 65, pl. 66, figs. 10-15
- 2008a *Triloculina* cf. *T. fichteliana* d'Orbigny – Meriç *et al.*, pl. 7, figs. 14-16, pl. 8, figs. 1-2
- 2008b *Triloculina* cf. *T. fichteliana* d'Orbigny – Meriç *et al.*, pl. 4, figs. 9-12
- 2010 *Triloculina fichteliana* d'Orbigny – Koukousioura *et al.*, p. 164, pl. 2, fig. 3
- 2015 *Miliolinella fichteliana* (d'Orbigny) – Delliou *et al.*, pl. 1, fig. 1
- 2017 *Triloculina fichteliana* d'Orbigny – Thissen & Langer, pl. 8, figs. 10-12

Triloculina fichteliana was originally described by d'Orbigny (1839) from Cuba (see also Le Calvez, 1977). The specimens recorded from the Mediterranean and the Red Sea share a common morphology, that differs from the material described from Caribbean localities (Cushman, 1929). The main differences concern the test ornamentation and the shape of the tooth. We therefore consider specimens from the Red Sea, the Mediterranean Sea (Hottinger *et al.*, 1993; Meriç *et al.*, 2008a; 2008b) and the Indian Ocean (Thissen & Langer, 2017) to represent a species that differs slightly from d'Orbigny (1839), herein referred to as *Triloculina* cf. *T. fichteliana*. As such, we consider it to represent a species alien to the Mediterranean Sea. The specimen illustrated by Cushman (1929) differs from the holotype of d'Orbigny in having a lip instead of a tooth (see also Le Calvez, 1977), belongs to a different genus, but does not have taxonomic priority over *Triloculina fichteliana* of d'Orbigny (1839).

***Vaginulinopsis sublegumen* Parr, 1950**

- 1950 *Vaginulinopsis sublegumen* Parr – Parr, p. 325, pl. 11, figs. 18a-b
- non 2004 *Astacolus sublegumen* (Parr) – Meriç *et al.*, p. 126, pl. 19, fig. 9

Astacolus sublegumen was reported to be present in Turkey by Meriç *et al.* (2004). The holotype of this species originates from the Antarctic Ocean and is currently referred to as *Vaginulinopsis sublegumen* (see Jones, 1994; and WoRMS Editorial Board, 2020). The specimen illustrated by Meriç *et al.* (2004) significantly differs from the holotype. The same illustration was used in

Yokeş & Meriç (2009) and Meriç *et al.* (2014). To date, there is no valid record of *Vaginulinopsis sublegumen* in the Mediterranean Sea and the species is therefore absent and not alien.

***Varidentella* cf. *V. neostriatula* (Thalman, 1950)**

- 1932 cf. *Quinqueloculina striatula* Cushman – Cushman, p. 27, pl. 7, figs. 3-4
- 1950 *Quinqueloculina striatula* Thalman – Said, p. 5, pl. 1, fig. 9
- 1987 *Quinqueloculina neostriatula* Thalman – Baccaert, p. 91, pl. 43, figs. 1-6
- 1993 *Varidentella* cf. *V. neostriatula* (Thalman) – Hottinger *et al.*, p. 66, pl. 70, figs. 5-11, pl. 71, figs. 1-7
- 1998 *Varidentella neostriatula* (Thalman) – Piller & Haunold, pl. 6, fig. 9
- 2000 *Varidentella* cf. *V. neostriatula* (Thalman) – Hyams, pl. 14, fig. 4

Quinqueloculina striatula was described by Cushman (1932) from Mojaoukar Anchorage, Fiji (Melanesia). The holotype is characterized by five chambers, a subacute chamber periphery and a test wall that is well ornamented by numerous costae. *Quinqueloculina neostriatula* recorded by Said (1950) and Piller & Haunold (1998) from the northern Red Sea resembles the specimens illustrated by Hottinger *et al.* (1993). However, Said (1950) provides only a single side view illustration. Baccaert (1987) illustrated specimens from Australia with variable morphologies, some of which resemble specimens from the northern Red Sea. A single record of *Varidentella* cf. *V. neostriatula* from the Mediterranean Sea was illustrated by Hyams (2000). It is identical to some of the specimens illustrated by Hottinger *et al.* (1993). The morphotypes of this species are highly variable and further studies are required. For the time being, we consider *Varidentella* cf. *V. neostriatula* to be alien to the Mediterranean Sea.

Discussion

Our survey and reexamination of alien benthic foraminifera currently present in the Mediterranean Sea yielded a total of 44 validly recognized species and two crypto-genic taxa (Table 1). This reduces the number of previous recordings of alien foraminifera (Hyams-Kaphzan *et al.*, 2008; Zenetos *et al.*, 2012; EASIN, 2020), and is mainly due to new occurrence records, new findings in the fossil record, erroneous identifications, and evidence predating the presence of alien species prior to the opening of the Suez Canal. The revised list of accepted alien species includes both larger symbiont-bearing and smaller benthic foraminifera, including 16 hyaline-perforate, 3 agglutinated and 25 porcelaneous taxa.

The vast majority of alien foraminifera recorded so far have become established in the Eastern Mediterranean Sea (41 species = 93.1 %; Tbl. 1, Fig. 1), primarily along the Levantine coasts (Langer & Hottinger, 2000; Hyams *et al.*, 2002; Hyams-Kaphzan *et al.*, 2008; Langer, 2008;

Almogi-Labin & Hyams-Kaphzan, 2012; Langer *et al.*, 2012; Mouanga & Langer, 2014; Langer & Mouanga, 2016; Guastella *et al.*, 2019; Servello *et al.*, 2019; Katsanevakis *et al.*, 2020). The number of alien species decreases rapidly from east to west. Eleven species of alien benthic foraminifera were recorded in the central Mediterranean (27.2 %), 8 species in the Adriatic (20.4 %), and only 5 (11.6 %) in the western Mediterranean Sea. The preferential establishment of NIS of benthic foraminifera within the warmer eastern Mediterranean is indicative for a point source from a tropical location, in particular the Red Sea/Indian Ocean with a pathway through the Suez Canal. In fact, recent molecular phylogenetic analyses have provided evidence that modern populations of a few selected species belong to the same genetic population and were therefore considered Lessepsian migrants (Merkado *et al.*, 2013; 2015; Schmidt *et al.*, 2015). However, the fossil record provides evidence that all of them (*Textularia agglutinans*, *Sorites orbiculus*, *Pararotalia calcariformata*) were present in the Mediterranean prior to the opening of the Suez Canal (see Results). They are therefore native to the Mediterranean Sea and, per definition, can neither be considered alien nor invasive aliens. In addition, fully conclusive evidence to trace the origin of an alien species can only be provided when molecular data are available from the type locality. To trace the origin of a putative alien species via molecular genetic analysis of Mediterranean and Red Sea/Indian/Pacific Ocean material remains therefore inconclusive, if the type locality is located in a different geographic zone (e.g. the Caribbean Sea for *Textularia agglutinans*).

Being located at the crossroads between the Atlantic and the Indo-Pacific Oceans, alien species enter the Mediterranean Sea mainly via the Suez Canal but also via the Strait of Gibraltar and the Dardanelles. Foraminifera entering the Mediterranean via the Suez Canal pathway, generally follow the counterclockwise current pattern and first settle along the eastern shores of the Levantine coast (Langer, 2008). Biological invasions through the Suez Canal have been facilitated as a consequence of expanded connectivity including the extension, broadening and enlargement of the Suez Canal (Galil *et al.*, 2017; Zenetos, 2017). This has raised concern over rising propagule pressure and introductions of new NIS with potential consequences on native biotas (Galil *et al.*, 2017). The Red-to-Med invasion of tropical taxa has also been promoted by rising sea surface temperatures (Raitsos *et al.*, 2010; Manda *et al.*, 2020) and the Nile damming (Skirris & Lascaratos, 2004). While rising sea surface temperatures increase thermal habitat suitability (Marras *et al.*, 2015), damming of the Nile has caused drastic modifications in sedimentation patterns, circulation and salinity, diminished pre-existing environmental barriers and provided an additional stimulus for the invasion and establishment of alien species.

The distribution of tropical larger benthic foraminifera (LBF) is strongly constrained by water temperature (Langer & Hottinger, 2000). As they invade the temperate waters of the eastern Mediterranean, they meet a physical border that limits their rapid distribution. The tempera-

ture requirements of tropical foraminifera restrict their distributional capacity and migration towards the colder western Mediterranean, which is influenced by cold water currents from the Atlantic Ocean. Some of the larger symbiont-bearing foraminifera have reached Sicily and Tunisia (Langer & Hottinger, 2000, Langer *et al.*, 2012; El Kateb *et al.*, 2018; Damak *et al.*, 2019; Guastella *et al.*, 2019), entered the Adriatic Sea (Langer & Mouanga, 2016), and continue to their range expansion towards the northwestern Mediterranean with more than 10km/yr (Langer *et al.*, 2012; Guastella *et al.*, 2019).

To project future species distributions, species distribution modelling based on ecological niche constraints of current distributions were applied to foraminifera by Langer *et al.* (2012; 2013b) and Weinmann *et al.* (2013). The projection onto the RCP4.5 scenario (Collins *et al.*, 2013) suggests that the overall habitat suitability will increase with potential range expansions into the western area of the Mediterranean along the coast of northern Africa, into the Alboran and Balearic Sea, along the Tyrrhenian coast in Italy, and deep into the Adriatic Sea, following a continuous northwestward dispersal. The range expansions are fueled in response to temperature increases and the extension of climate belts (Tittensor *et al.*, 2010, Langer *et al.* 2012; Guastella *et al.*, 2019). Minimum winter sea surface temperatures (SST) has previously been invoked to be among the main agents controlling the latitudinal distribution of LBF (Zmiri *et al.*, 1974; Langer & Hottinger, 2000) and the observed range extension of thermophilic LBF and endosymbiotic foraminifera were shown to track contemporary SST increase (Langer *et al.*, 2012; Weinmann *et al.*, 2013; Langer & Mouanga, 2016; Schmidt *et al.*, 2015; 2018; Guastella *et al.*, 2019; Manda *et al.*, 2020). In addition, the proliferation and recent range expansion rates of LBF (Langer & Mouanga, 2016; Guastella *et al.*, 2019; Damak *et al.*, 2019) provide strong support for previous species distribution models projecting the northward migration and range shifts and corroborate findings that rising water temperatures and warm currents are the most likely agents controlling the latitudinal extension of larger symbiont-bearing foraminifera. Compared to alien LBF, hardly anything is known about range expansions rates of smaller benthic foraminifera. Previous studies have documented that populations of putative Lessepsian migrants (*Textularia agglutinans* and *Pararotalia calcariformata*) became very dominant along the Israeli Mediterranean coast (Hyams-Kaphzan *et al.*, 2014; Merkado *et al.*, 2015; Schmidt *et al.*, 2015) possibly as a result of rising water temperatures (e.g. Manda *et al.*, 2020).

Almost all alien foraminifera present in the Mediterranean Sea appear to be innocuous and only one fulfills the criteria to be considered as an invasive alien: *Amphistegina lobifera* (Langer *et al.*, 2012; Guastella *et al.*, 2019; Servello *et al.*, 2019). This species is a prolific carbonate producer (Hallock, 1981; Langer *et al.*, 1997; Langer, 2008) and displays extreme forms of ecosystem invasibility (Langer *et al.*, 2012; Mouanga & Langer, 2014; Langer & Mouanga, 2016). This includes mass occurrences, hyperabundances and the appearance in mono-

cultures, with capabilities to reduce native diversity and species richness (Mouanga & Langer, 2014), to transform the composition, grain size and chemistry of sediments from predominantly siliceous to carbonate deposits (Hyams *et al.*, 2002; Samir *et al.*, 2003; Gruber, 2007; Yokeş & Meriç, 2009; Abu Tair & Langer, 2010; Langer *et al.*, 2012; Weinmann *et al.*, 2013; Caruso & Cosentino, 2014; Mouanga & Langer, 2014; Guastella *et al.*, 2019; Meriç *et al.*, 2020), and to trigger changes in ecosystem functioning (Langer *et al.*, 2012). Triantaphyllou *et al.* (2005), Koukousioura *et al.* (2011), and Dimiza *et al.* (2016b) have suggested that their presence and numerical abundance may be used as proxy to assess the degree of environmental disturbance. At some sites extreme abundances of amphisteginid invaders were shown to result in the formation of large amounts of “living sands” (Meriç, 2008; Meriç *et al.*, 2008a; Yokeş & Meriç 2009; Abu Tair & Langer 2010; Langer *et al.*, 2012). The invasion and prolific occurrences indicate that the amphisteginid invaders successfully fill an open niche that is obviously providing ideal conditions. While the immediate impact of *Amphistegina lobifera* appears to be obvious, the resilience of native biotas to key invaders remains yet to be determined.

Amphisteginid invasions selectively affect taxa that share the same microhabitat and suggests that competitive exclusion is a major driving force regulating species richness in invaded communities. The displacement of native species may be of local nature and natives may persist in nearby uninvaded areas. Whether *Amphistegina lobifera* displaces any other organisms than foraminifera and what role they play in the food web of metacommunities, is currently not known. Mouanga & Langer (2014) have demonstrated that massive invasions of amphisteginids locally result in the homogenization of the foraminiferal faunas with a clear correlation between the diversity of foraminiferal biotas and percent abundances of amphisteginid invaders (Langer & Mouanga, 2016). *Amphistegina lobifera* was recently included in the list of the top 26 high-priority species that display invasive traits and for which the performance of risk assessments is encouraged (Tsiamis *et al.*, 2019) under Descriptor 2 of the European Union Marine Strategy Framework Directive (EU, 2008) and the Biodiversity Strategy (EU, 2014b).

Means of dispersal of benthic foraminifera are multifold and involve natural and human-mediated sources. Natural means involve transportation by ocean currents, attachment to bodies of birds and fish, the incidental consumption and translocation by macroorganisms, and rafting on algae and other objects (Lipps, 1983; Langer, 1993; Langer *et al.*, 1998; Goldbeck *et al.*, 2005; Guy-Haim *et al.*, 2017; Finger, 2017), where the ranges of native species are broadly constrained by limitations on their capacity for dispersal. Knowledge of the processes and mechanisms that govern dispersal and colonization in benthic organisms is crucial for understanding biodiversity patterns and historical development of biogeography. However, the rate at which humans translocate species has substantially intensified with increasing globalization and does not show any signs of saturation (Seebens *et al.*, 2017). Human-me-

diated introductions of NIS include ballast water, hull fouling and sea chests (water-intake recesses in the hull), intentional and accidental releases of aquaculture species, aquarium discards and aquarium trade.

Ballast water is generally invoked to be among of the major pathways for the introduction of nonindigenous marine species (Ruiz *et al.*, 1997; Carlton *et al.*, 1999; Molnar *et al.*, 2008; Seebens *et al.*, 2016), but hardly anything is known about the potential role of transoceanic ship deballasting as a vector for the introduction of alien foraminifera in the Mediterranean Sea (Galil & Hülsmann 1997; McGann *et al.*, 2020). The release of ballast water and the transport stowaway pathway related to shipping traffic (biofoulers and hitchhikers) have been widely suspected as vectors for the introduction of non-native foraminifera (Hayward, 1997; Hayward *et al.*, 1999; 2004; McGann *et al.*, 2000; Calvo-Marcilese & Langer, 2010; Polovodova Asteman & Schönfeld, 2016; Eichler *et al.*, 2018; Deldicq *et al.*, 2019), has led to a breakdown of classical biogeographic regions and is considered a major vector for Red-to-Med alien species invasions (Zenetos *et al.*, 2012).

Cargo ships and tankers take up and release large volumes of ballast water. The ballast water is used for stability and maneuverability and may also be added to add weight to pass under bridges and other structures. Ballasting and deballasting most frequently occurs in shallow water, in ports and in estuaries. Ballast water was shown to contain abundant and diverse biotas, is moved across oceans and constitutes “a conveyor belt of marine organisms wrapping around the world”. Together with the ballast water, ships also pump substantial amount of sediment (including foraminifera) into their ballast tanks (Carlton & Geller 1993; Chu *et al.*, 1997; Galil & Hülsmann 1997; Gollasch *et al.*, 1998; Macdonald, 1998; Lavoie *et al.*, 1999; Smith *et al.*, 1999; McGann *et al.*, 2000; 2020; Drake *et al.*, 2005; Johengen *et al.*, 2005). Ballast water may contain both benthic and planktic foraminifera, including all life stages ranging from propagules to adult life-stages. The invasion success of ballast water aliens is highly probabilistic and a game of ecological roulette, where the outcome depends on a myriad of dynamic factors (Ricciardi, 2016), including species traits, available niche space, interactions with native biotas, trade routes, climate change and water temperature (among many others).

Predicting foraminiferal invasions is a challenging task and there is general international consensus that multi-vector pathways-based management is a priority in minimizing marine alien species (Ojaveer *et al.*, 2018). To date, there is no evidence that foraminifera are toxic (Langer & Bell, 1995), they are of no concern to human health, and by virtue of their substantial carbonate production, they may be even beneficial to coastal protection (Hallock, 2002; Hohenegger, 2006; Langer, 2008, Langer *et al.*, 2013b; Doo *et al.*, 2014). Management of invasive alien species has been among the main priorities of the IAS Regulation (Article 4, EU, 2014a), but measures to mitigate the impact of alien foraminifera, or to efficiently control or even eradicate are neither feasible

nor costeffective (Tsiamis *et al.*, 2020), and may be even counterproductive. Because of their abundance, high reproduction rates, and ubiquity in virtually all marine environments, however, foraminifera are excellent predictors of rates of global change.

The revision and revised list of alien foraminifera presented here is a dynamic resource and will continue to evolve as the number of alien species is likely to increase. Regular updates, more accurate information and the application of molecular techniques will assist our understanding of climate driven range shifts and the magnitude of alien species impacts in the Mediterranean Sea and in global oceans. In addition, recent molecular genetic studies provide accumulating evidence that traditional morphospecies taxonomy vastly underestimates true diversity in foraminifera and routinely report a higher number of previously unrecognized species (e.g. Pawlowski *et al.*, 2014; Prazeres *et al.*, 2020). On a molecular level, alien foraminifera present in the Mediterranean Sea are currently underresearched and require intensified efforts to trace their source. Rigorous morpho- and molecular species identifications, studies on ballast water, species distribution modeling and new evidence from the fossil record will thus continue to shed new light on the status (native/alien/cryptogenic), pathway, vector and origin of species (Atlantic/Red Sea).

Summary

A large-scale survey on alien benthic foraminifera in the Mediterranean Sea has been conducted and resulted in a revised list of 44 validly recognized alien and two cryptogenic species, including both larger symbiont-bearing and smaller benthic taxa. The reexamination of previous alien species records is based on a critical and careful revision of the current taxonomic status of each species, new findings in the fossil record, and recently published data obtained through molecular genetic analysis. The vast majority of alien foraminifera recorded so far have established self-sustaining populations in the Eastern and Central Mediterranean Sea. It is anticipated that rising sea surface temperatures and the expansion of climate belts will convey the range expansion and northwestward migration of alien species. To date, only *Amphistegina lobifera* meets the full set of criteria to be considered as an invasive alien. Measures to mitigate its impact on ecosystems and native biotas, to efficiently control, manage or even eradicate are neither feasible nor costeffective and may even be counterproductive. The number of alien foraminifera is likely to evolve as travel, shipping trade, rising temperatures and increasing globalization promote the spread of alien taxa in the Mediterranean Sea. Because of their abundance, ubiquity, and high reproduction rates, foraminifera are excellent proxies for rates of global climate change. Monitoring future range expansions of key alien taxa (e.g. *Amphistegina* spp.) will thus provide a baseline to assess the magnitude, rate and impact of predicted global change.

Acknowledgements

We are grateful to the reviewers and the editor (Argyro Zenetos) for their constructive comments and suggestions. Field and laboratory assistance were provided by Gloria Mouanga, Jens Thissen, Anna Weinmann, Laura Huth, Christian Lott, Miriam Weber, Boris Unger, Michael Freise, Nisan Sariaslan, Leon Paiano and Nasir Abu Tair. We also thank Nicholas-Jason Xentidis and Gianna Servello for allowing us to use the illustration for the Mediterranean subregions, Baki Yokes for information on *Cyclorbiculina*, and Ahuva-Almogi-Labin from the Geological Survey of Israel (GSI) for sharing core-material and for fruitful discussions. Yael Edelman-Furstenberg (GSI) is appreciated for sharing her core-material (Herzliya, Israel) and Michael Kitin (GSI) is thanked for preparing sample material. Material from Israel studied by OHK was supported by the Israel Science Foundation (Grant No. 145/02-13.0 to Ahuva Almogi-Labin, Barak Herut and Chaim Benjamini) and by the Ministry of National Infrastructures (grant no. 98-17-017 to Ahuva Almogi-Labin and Sivan Dorit; grant no. 20-17-20 to Chaim Benjamini). Collection and study of the material by MRL was supported by the German Science Foundation (LA 884/10-1, LA 884/14-1).

References

- Abu Tair, N., Langer, M.R., 2010. Foraminiferal invasions: the effect of Lessepsian migration on the diversity and composition of benthic foraminiferal assemblage around Cyprus (Mediterranean Sea). p. 42. In: *FORAMS 2010, International Symposium on Foraminifera – Abstracts Volume with Program*. Langer, M.R. (Ed.). Bayleydruck GmbH, University of Bonn.
- Adams, C.G., 1976. Larger foraminifera and the late Cenozoic history of the Mediterranean region. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 3 (20), 47-66.
- Aiello, G., Barra, D., Coppa, M.G., Valente, A., Zeni, F., 2006. Recent infralittoral Foraminiferida and Ostracoda from the Porto Cesareo Lagoon (Ionian Sea, Mediterranean). *Bollettino della Società Paleontologica Italiana*, 45 (1), 1-14.
- Alavi, S.N., 1980. *Micropalaeontological studies of recent sediments from the Cilicia Basin (NE Mediterranean)*. PhD Dissertation. University of London, London, 90 pp.
- Albani, A.D., 1974. New benthonic foraminiferida from Australian waters. *Journal of Foraminiferal Research*, 4 (1), 33-39.
- Almogi-Labin, A., Hyams-Kaphzan, O., 2012. *Epistomaroides punctatus* (Said, 1949) - a new alien foraminifera found at Akhziv-Rosh HaNikra, northern Israel, eastern Mediterranean Sea. *Mediterranean Marine Science*, 13 (2), 294-296.
- Aloulou, F., EllEuch, B., Kallel, M., 2012. Benthic foraminiferal assemblages as pollution proxies in the northern coast of Gabes Gulf, Tunisia. *Environmental Monitoring and Assessment*, 184 (2), 777-795.
- Altınışağı, S., Yardımcı, C.H., Altınışağı, S., Paçal, F.P., 2017. The species list belonging to some benthic invertebrate groups in a coastal lagoon: Kamil Abduş Lagoon (Istanbul,

- Turkey). *Journal of Entomology and Zoology Studies*, 5, 307-313.
- Alve, E., Goldstein, S.T., 2003. Propagule transport as a key method of dispersal in benthic foraminifera (Protista). *Limnology and Oceanography*, 48, 2163-2170.
- Alvira Romero, B., 2019. *Primera presencia y distribución de las especies exóticas Parasorites orbitoloides (Hofker, 1930) y Euthymonacha polita (Chapman, 1900): efectos sobre la comunidad de macroforaminíferos bentónicos del Mediterráneo Occidental*. MSc Thesis. Máster Universitario en Ecología Marina de la Universitat de les Illes Balears, 78 pp.
- Armynot du Châtelet, E., Bout-Roumazeilles, V., Coccioni, R., Frontalini, F., Guillot, F. *et al.*, 2013. Environmental control on shell structure and composition of agglutinated foraminifera along a proximal–distal transect in the Marmara Sea. *Marine Geology*, 335, 114-128.
- Avital, A., 2002. *Geological history of the Plio-Pleistocene-Holocene offshore based on cores off Ashqelon, southern Israel*. MSc. Thesis, Ben-Gurion University of the Negev, Beer Sheva, Israel, 180 pp. (In Hebrew, English abstract).
- Avnaim-Katav, S., 2010. *Chronostratigraphy and Paleo-environments: Reconstruction of Quaternary subsurface successions, Southern Haifa Bay*. PhD Dissertation. Department of Maritime Civilizations, University of Haifa, Haifa, Israel, 242 pp (In Hebrew, English abstract).
- Avnaim-Katav, S., Almogi-Labin, A., Kanari, M., Herut, B., 2020. Living benthic foraminifera of southeastern Mediterranean ultra-oligotrophic shelf habitats: Implications for ecological studies. *Estuarine, Coastal and Shelf Science*, 234, 106633.
- Avşar, N., 1997. Doğu Akdeniz kıyı bölgesi bentik foraminiferleri. *Ç.Ü. Yerbilimleri*, 31, 67-81.
- Avşar, N., Meriç, E., Ergin, M., 2001. Foraminiferal content of the benthogenic sediments in the İskenderun Bay. *Yerbilimleri*, 24, 97-112.
- Avşar, N., Meriç, E., Alramazanoğlu, A., Dinçer, F., 2008. Recent benthic foraminiferal assemblages in the continental shelf of the Gulf of Antalya (SW Turkey). *Yerbilimleri*, 29 (3), 111-136.
- Ayadi, N., Zghal, I., Aloulou, F., Bouzid, J., 2016. Impacts of several pollutants on the distribution of recent benthic foraminifera: the southern coast coast of Gulf of Gabes, Tunisia. *Environmental Science and Pollution Research*, 23 (7), 6414-6429.
- Baccaert, J., 1987. *Distribution patterns and taxonomy of benthic foraminifera in the Lizard Island Complex, northern Great Barrier Reef, Australia*. PhD Thesis. Université de Liège – CAPS Laboratoire de Biosédimentologie, Liège, 298 pp.
- Badr-ElDin, A.M., Charle, C.M., El-Sabrouti, M.A.F., 2019. Response of benthic foraminifera to coastal protection of the western coast of Alexandria, Egypt. *Egyptian Journal of Aquatic Research*, 45 (1), 1-9.
- Bassi, D., Braga, J.C., Domenico, G.D., Pignatti, J., Abramovich, S. *et al.*, 2019. Palaeobiogeography and evolutionary patterns of the larger foraminifer *Borelis* de Montfort (Borelidae). *Papers in Palaeontology*, 1-27.
- Bassler-Veit, B., Barut, İ.F., Meriç, E., Avşar, N., Nazik, A., *et al.*, 2013. Distribution of microflora, meiofauna, and macrofauna assemblages in the hypersaline environment of northeastern Aegean Sea coasts. *Journal of Coastal Research*, 29 (4), 883-898.
- Basso, D., Spezzaferri, S., 2000. The distribution of living (stained) benthic foraminifera in Iskenderun Bay (Eastern Turkey): A statistical approach. *Bolletino della Società Paleontologica Italiana*, 39 (3), 359-379.
- Batsch, A.J.G.K., 1791. *Sechs Kupfertafeln mit Conchylien des Seesandes*. In Commission der academischen Buchhandlung, Jena, 15 pp.
- Blanc-Vernet, L., 1969. Contributions à l'étude des foraminifères de Méditerranée. *Recueil des Travaux de la Station Marine d'Endoume*, 48, 315.
- Blanc-Vernet, L., Clairefond, P., Orsolini, P., Templeton, R.S.M., Desprat, R., 1979. B. - Les Foraminifères/ Foraminifera. *Géologie Méditerranéenne*, 6 (1), 171-209.
- Bouchet, V.M.P., Debenay, J.-P., Sauriau, P.G., 2007. First report of *Quinqueloculina carinatastriata* (Wiesner 1923) (Foraminifera) along the French Atlantic coast (Marennes-Oléron Bay and Ile de Ré). *Journal of Foraminiferal Research*, 37 (3), 204-212.
- Brady, H.B., 1879. Notes on some of the Reticularian Rhizopoda of the "Challenger" Expedition. *Quarterly Journal of Microscopical Science. New Series*, 19, 20-62.
- Brady, H.B., 1881. Notes on some of the Reticularian Rhizopoda of the "Challenger" Expedition: Part III. *Quarterly Journal of Microscopical Science. New Series*, 21 (81), 31-71.
- Brady, H.B., 1884. Notes on the foraminifera dredged by H.M.S. Challenger during the years 1873-1876. *Report of the Scientific Results of the Voyage of H.M.S. Challenger, Zoology*, 9, 1-814.
- Brönnimann, P., 1979. Recent benthonic foraminifera from Brasil. Morphology and ecology. Part IV: Trochamminids from the Campos shelf with description of *Paratrochammina* n.gen.. *Paläontologische Zeitschrift*, 53 (1-2), 5-25.
- Buchner, P., 1940. Die Lagenen des Golfes von Neapel und der marinen Ablagerungen auf Ischia. *Nova Acta Leopoldina*, 9 (62), 363-560.
- Calvo-Marcilese, L., Langer, M.R., 2010. Breaching biogeographic barriers: The invasion of *Haynesina germanica* (Foraminifera, Protista) in the Bahia Blanca estuary, Argentina. *Biological Invasions*, 12, 3299-3306.
- Carlton, J.T., 1999. The scale and ecological consequences of biological invasions in the world's oceans. p. 195-212. In: *Invasive Species and Biodiversity Management*. Sandlund, O.T., Schei, P.J., Viken, A. (Eds.). Kluwer, Dordrecht, Netherlands.
- Carlton, J.T., Geller, J.B., 1993. Ecological roulette: the global transport of nonindigenous marine organisms. *Science*, 261 (5117), 78-82.
- Caruso, A., Cosentino, C., 2014. The first colonization of the Genus *Amphistegina* and other exotic benthic foraminifera of the Pelagian Islands and south-eastern Sicily (central Mediterranean Sea). *Marine Micropaleontology*, 111, 38-52.
- CBD (Convention on Biological Diversity), 2012. *Aichi Biodiversity Targets*. <https://www.cbd.int/sp/targets/> (Accessed 2 April 2020)
- CBD (Convention on Biological Diversity), 2014. Pathways of introduction of invasive species, their prioritization and

- management. UNEP/CBD/SBSTTA/18/9/Add. 1. Secretariat of the Convention on Biological Diversity, Montreal.
- Chapman, F., 1898. On *Haddonia*, a new Genus of the Foraminifera, from Torres Straits. *Journal of the Linnean Society of London, Zoology*, 26 (169), 452-456.
- Chapman, F., 1901. On some new and interesting Foraminifera from the Funafuti Atoll, Ellice Islands / Foraminifera from the Lagoon at Funafuti. *Journal of the Linnean Society (Zoology)*, 28 (179), 161-210.
- Chapman, F., 1907. Recent Foraminifera of Victoria: Some Littoral Gatherings. *Journal of the Quekett Microscopical Club*, 10, 117-146.
- Cherif, O.H., 1970. *Die Miliolacea der Westküste von Naxos (Griechenland) und ihre Lebensbereiche*. PhD Thesis. Technische Universität Clausthal, Fakultät Natur- und Geisteswissenschaften, Clausthal, 176 pp.
- Chu, K.H., Tam, P.F., Fung, C.H., Chen, Q.C., 1997. A biological survey of ballast water in container ships entering Hong Kong. *Hydrobiologia*, 352, 201-206.
- Cimernan, F., Langer, M.R., 1991. *Mediterranean Foraminifera*. Slovenska Akademija Znanosti in Umetnosti, Ljubljana, 117 pp.
- Çinar, M.E., Bilecenoğlu, M., Ozturk, B., Katagan, T., Yokeş, M.B. *et al.*, 2011. An updated review of alien species on the coasts of Turkey. *Mediterranean Marine Science*, 12 (2), 257-315.
- Cohen, A.N., 1998. *Ship's Ballast Water and the Introduction of Exotic Organisms into the San Francisco Estuary. Current Status of the Problem and Options for Management*. San Francisco Estuary Institute, Richmond CA, 84 pp.
- Collins, A.C., 1958. Foraminifera. p. 335-437. In: *Scientific reports / Great Barrier Reef expedition 1928-29*. Eyre and Spottiswoode Limited, London.
- Collins, M., Knutti, R., Arblaster, J., Dufresne, J.L., Fichetef, T. *et al.*, 2013. Chapter 12 – Long-term climate change: Projections, commitments and irreversibility. p. 1029-1136. In: *Climate Change 2013 – The Physical Science Basis: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Stocker, T.F., Qin, D., Plattner, G.K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.). Cambridge University Press, New York.
- Corsini-Foka, M., Zenetos, A., Crocetta, F., Çinar, M.E., Koçak, F. *et al.*, 2015. Inventory of alien and cryptogenic species of the Dodecanese (Aegean Sea, Greece): collaboration through COST action training school. *Management of Biological Invasions*, 6, 351–366.
- Cosentino, C., Pepe, F., Scopelliti, G., Calabrò, M., Caruso, A., 2013. Benthic foraminiferal response to trace element pollution - the case study of the Gulf of Milazzo, NE Sicily (Central Mediterranean Sea). *Environmental Monitoring and Assessment*, 185 (10), 8777-8802.
- Costello, M.J., Coll, M., Danovaro, R., Halpin, P., Ojaveer, H. *et al.*, 2010. A Census of Marine Biodiversity Knowledge, Resources, and Future Challenges. *PLoS ONE*, 5 (8), e12110.
- Ćosović, V., Zavodnik, D., Borčić, A., Vidović, J., Deak, S. *et al.*, 2011. A checklist of Foraminifera of the Eastern Shelf of the Adriatic Sea. *Zootaxa*, 3035 (1), 1-56.
- Culver, S.J., Buzas, M.A., 1980. Distribution of recent benthic foraminifera off the North American Atlantic coast. *Smithsonian Contributions to the Marine Sciences*, 6, 1-512.
- Cushman, J.A., 1917. A monograph of the Foraminifera of the north Pacific Ocean. Part VI. Miliolidae. *Bulletin of the United States National Museum*, 71, 1-108.
- Cushman, J.A., 1918. Foraminifera from Murray Island, Australia. *Papers from the Department of Marine Biology. Carnegie Institute of Washington*, 9, 289-291.
- Cushman, J.A., 1922. *Shallow-water Foraminifera of the Tortugas region*. Department of Marine Biology of the Carnegie Institution Publication No. 311, Washington DC, 85 pp.
- Cushman, J.A., 1923. The Foraminifera of the Atlantic Ocean. Part 4: Lagenidae. *Bulletin of the United States National History Museum*, 104, 1-228.
- Cushman, J.A., 1924. Samoan foraminifera. *Tortugas Laboratory. Carnegie Institution of Washington*, 21.
- Cushman, J.A., 1928. Foraminifera. Their classification and economic use. *Cushman Laboratory for Foraminiferal Research Special Publication*, 1, 1-401.
- Cushman, J.A., 1929. The Foraminifera of the Atlantic Ocean. Part 6: Miliolidae, Ophthalmitidae and Fischerinidae. *Bulletin of the United States National Museum*, 104, 1-129.
- Cushman, J.A., 1931. Notes on the Foraminifera described by Batsch in 1791. *Contributions from the Cushman Laboratory for Foraminiferal Research*, 7 (3), 62-72.
- Cushman, J.A., 1932. The foraminifera of the tropical Pacific collections of the Albatross, 1899-1900. Part 1: Astrothizidae to Trochamminidae. *Bulletin of the United States National Museum*, 161, 1-85.
- Cushman, J.A., 1933. The foraminifera of the tropical Pacific collections of the Albatross, 1899-1900. Part 2: Lagenidae to Alveolinellidae. *Bulletin of the United States National Museum*, 161.
- Cushman, J.A., 1934. Notes on the genus *Tretomphalus*, with descriptions of some new species and a new genus, *Pyropilus*. *Contributions from the Cushman Laboratory for Foraminiferal Research*, 10 (4), 79-101.
- Cushman, J.A., 1944. The genus *Articulina* and its species. *Cushman Laboratory for Foraminiferal Research Special Publication no. 10*, 1-21.
- Cushman, J.A., 1946. The genus *Hauerina* and its species. *Contributions from the Cushman Laboratory for Foraminiferal Research*, 22 (1), 2-14.
- Cushman, J.A., Todd, R., 1944. The genus *Spiroloculina* and its species. *Cushman Laboratory for Foraminiferal Research Special Publication no. 11*, 1-55.
- Cushman, J.A., Todd, R., Post, R.J., 1954. Recent Foraminifera of the Marshall Islands: Bikini and nearby atolls: part 2: Oceanography (biologic). *U. S. Geological Survey Professional Paper*, 260, 319-394.
- Damak, M., Frontalini, F., Elleuch, B., Kallel, M., 2019. Benthic foraminiferal assemblages as pollution proxies along the coastal fringe of the Monastir Bay (Tunisia). *Journal of African Earth Sciences*, 150, 379-388.
- Daniels, C.H. von, 1970. Quantitative ökologische Analyse der zeitlichen und räumlichen Verteilung rezenter Foraminiferen im Limski-kanal bei Rovinj (nördliche Adria). *Göttinger Arbeiten zur Geologie und Paläontologie*, 8, 1-109.
- Davaud, E., Septfontaine, M., 1995. Post-mortem onshore transportation of epiphytic foraminifera; recent example

- from the Tunisian coastline. *Journal of Sedimentary Research*, 65 (1a), 136-142.
- Debenay, J.-P., 2012. *A Guide to 1,000 Foraminifera from Southwestern Pacific: New Caledonia*. IRD Editions Marseille/Publication Scientifiques du Muséum, Paris, 378 pp.
- Debenay, J.-P., Millet, B., Angelidis, M.O., 2005. Relationships between foraminiferal assemblages and hydrodynamics in the Gulf of Kalloni, Greece. *Journal of Foraminiferal Research*, 35 (4), 327-343.
- Deldicq, N., Alve, E., Schweizer, M., Polovodova Asteman, I., Hess, S. *et al.*, 2019. History of the introduction of a species resembling the benthic foraminifera *Nonionella stella* in the Oslofjord (Norway): Morphological, molecular and paleo-ecological evidences. *Aquatic Invasions*, 14, 182-205.
- Delliou, A., Antoniadou, C., Almpnakis, K., Tsoukala, E., Chintiroglou, C., 2015. Biodiversity of sublittoral foraminiferal assemblages in the northeast Aegean Sea: preliminary results. *Proceedings of the 11th Panhellenic Symposium on Oceanography and Fisheries*, 569-572.
- Dermitzakis, M.D., Triantafyllou, M.V., 1990. Ecostratigraphical observations at the eastern part of Corinthiakos Gulf. *Annales Geologiques pays Helleniques*, 34 (2), 127-161.
- Dimiza, M.D., Koukousioura, O., Triantaphyllou, M.V., Dermitzakis, M.D., 2016a. Live and dead benthic foraminiferal assemblages from coastal environments of the Aegean Sea (Greece): Distribution and diversity. *Revue de Micropaleontologie*, 59, 19-32.
- Dimiza, M.D., Triantaphyllou, M.V., Koukousioura, O., Hallock, P., Simboura, N., *et al.*, 2016b. The Forum Stress Index: A new tool for environmental assessment of soft-bottom environments using benthic Foraminifera. A case study from the Saronikos Gulf, Greece, Eastern Mediterranean. *Ecological Indicators*, 60, 611-621.
- Diz, P. Francés, G., Costas, S., Souto, C., Alejo, I., 2004. Distribution of benthic foraminifera in coarse sediments, Ría de Vigo, NW Iberian margin. *Journal of Foraminiferal Research*, 34 (4), 258-275.
- Doo, S.S., Fujita, K., Byrne, M., Uthicke, S., 2014. Fate of calcifying tropical symbiont-bearing large benthic foraminifera: Living sands in a changing ocean. *The Biological Bulletin*, 226 (3), 169-186.
- Dorst, S., Schönfeld, J., 2013. Diversity of benthic foraminifera on the shelf and slope of the NE Atlantic: Analysis of datasets. *Journal of Foraminiferal Research*, 43 (3), 238-254.
- Drake, L.A., Jenkins, P.T., Dobbs, F.C., 2005. Domestic and international arrivals of NOBOB (no ballast on board) vessels to lower Chesapeake Bay. *Marine Pollution Bulletin*, 50, 560-565.
- EASIN, 2020. *European Alien Species Information Network*. <https://easin.jrc.ec.europa.eu/easin> (accessed 25 February 2020)
- Edelist, D., Rilov, G., Golani, D., Carlton, J.T., Spanier, E., 2013. Restructuring the Sea: Profound shifts in the world's most invaded marine ecosystem. *Diversity and Distributions*, 19, 69-77.
- Ehrenberg, C.G., 1839. Über die Bildung der Kreidefelsen und des Kreidemergels durch unsichtbare Organismen. *Physikalische Abhandlungen der Königlichen Akademie der Wissenschaften zu Berlin*, 59-147.
- Eichler, P.P.B., McGann, M., Rodrigues, A.R., Mendonça, A., Amorim, A. *et al.*, 2018. The occurrence of the invasive foraminifera *Trochammina hadai* Uchio in Flamengo Inlet, Ubatuba, São Paulo State, Brazil. *Micropaleontology*, 64, 391-402.
- El Kateb, A., Stalder, C., Neururer, C., Fentimen, R., Spangenberg, J.E. *et al.*, 2018. Distribution of benthic foraminiferal assemblages in the transitional environment of the Djerba lagoon (Tunisia). *Swiss Journal of Geosciences*, 111 (3), 589-606.
- El-Nakhal, H.A., 1983. *Agglutinella*, a new miliolid genus (Foraminiferida). *Journal of Foraminiferal Research*, 13 (2), 129-133.
- El-Nakhal, H.A., 1990. *Septoculina*, a new genus in Hauerininae (Foraminiferida). *Micropaleontology*, 36, 88-95.
- Elshanawany, R., Ibrahim, M.I., Milker, Y., Schmiedl, G., Badr, N. *et al.*, 2011. Anthropogenic impact on benthic foraminifera, Abu-Qir Bay, Alexandria, Egypt. *Journal of Foraminiferal Research*, 41 (4), 326-348.
- Emter, A., 2018. *Flachwasser-Foraminiferen als Indikatoren für Habitat-Typologien und Strömungsmuster von Libyen*. MSc Thesis. Steinmann Institut für Geologie, Paläontologie und Mineralogie der Rheinischen Friedrich-Wilhelms-Universität, Bonn, 76 pp (in German, English abstract).
- EU, 2008. Directive 2008//56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive). *Official Journal of the European Union*, L164, 19-40.
- EU, 2014a. Regulation (EU) No 1143/2014 of the European Parliament and of the Council of 22 October 2014 on the prevention and management of the introduction and spread of invasive alien species. *Official Journal of the European Union*, L317, 35.
- EU, 2014b. Report from the Commission to the European Parliament and the Council. The mid-term review of the EU Biodiversity Strategy to 2020. *European Commission Report*. Brussels, Belgium.
- Fajemila O.T., Langer, M.R., Lipps, J.H. 2015. Spatial Patterns in the Distribution, Diversity and Abundance of Benthic Foraminifera around Moorea (Society Archipelago, French Polynesia). *PLoS ONE* 10(12): e0145752.
- Fajemila, O.T., Langer, M.R., 2017. Spatial distribution and biogeographic significance of foraminiferal assemblages from São Tomé and Príncipe, Gulf of Guinea, West Africa. *Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen*, 285 (3), 337-360.
- Fajemila, O.T., Langer, M.R., Lipps, J.H. (2020): Atlas of Shallow-Water Tropical Benthic Foraminifera from Moorea (Society Islands, French Polynesia). *Cushman Foundation for Foraminiferal Research, Special Publication*, 48, 1-107.
- Ferragut Perelló, F., 2018. *Detecció de la dispersió de foraminífers bentònics mitjançant l'anàlisi del contingut estomacal de peixos demersals*. Facultat de Ciències, Universitat de les Illes Balears, Palma, 25 pp.
- Fichtel, L., Moll, J.P.C., 1798. *Testacea microscopica aliaque minuta ex generibus argonauta et nautilus ad naturam picta et descripta: Mikroskopische und andere kleine Schalthiere aus den Geschlechtern Argonaute und Schiffer; nach der Natur gemahlet und beschrieben*. Anton Pichler, Wien.
- Finger, K.L., 2017. Tsunami-generated rafting of foraminifera

- across the North Pacific Ocean. *Aquatic Invasions*, 13 (1), 17-30.
- Foraminiferi Padani (Terziario e Quaternario): Atlante iconografico e distribuzione stratigrafica*, 1982. Agip Mineraria, S.p.A., Presso la Tipo-Litografia Saccardo Carlo & Figli Snc, Ornavasso (Novara).
- Förderer, M., Langer, M.R., 2018. Atlas of benthic foraminifera from coral reefs of the Raja Ampat Archipelago (Irian Jaya, Indonesia). *Micropaleontology*, 64 (1-2), 1-170.
- Förderer, M., Langer, M.R., 2019. Exceptionally species-rich assemblages of modern larger benthic foraminifera from nearshore reefs in northern Palawan (Philippines). *Revue de Micropaléontologie*, 65, 100387.
- Galil, B.S., Hülsmann, N., 1997. Protist transport via ballast water – biological classification of ballast tanks by food web interactions. *European Journal of Protistology*, 33, 244-253.
- Galil, B.S., Boero, F., Campbell, M.L., Carlton, J.T., Cook, E. et al., 2015. ‘Double trouble’: The expansion of the Suez Canal and marine bioinvasions in the Mediterranean Sea. *Biological Invasions*, 17 (4), 973-976.
- Galil, B.S., Marchini, A., Occhipinti-Ambrogi, A., Ojaveer, H., 2017. The enlargement of the Suez Canal – Erythraean introductions and management challenges. *Management of Biological Invasions*, 8, 141-152.
- Glacon, G. 1962. Foraminifères des dépôts actuels des côtes de Tunisie Sud Orientale. Thèse de Doctorat Université de Montpellier, France, 270 pp.
- Goldbeck, E., Langer, M.R., 2009. Biogeographic provinces and patterns of diversity in selected Upper Cretaceous (Santonian-Maastrichtian) larger foraminifera. In: Demchuk, T. and A.C. Gary (eds.); Geological Problem Solving with Microfossils: A Volume in Honor of Garry D. Jones. *SEPM Special Publication*, 93, 187-232.
- Gollasch, S., 2002. The importance of ship hull fouling as a vector of species introductions into the North Sea. *Biofouling*, 18, 1051-1121.
- Gollasch, S., Dammer, M., Lenz, J., Andres, H. G., 1998. Non-indigenous organisms introduced via ships into German waters. p. 50-64. In: Ballast Water: Ecological and Fisheries Implications: ICES Conference (83rd Statutory Meeting), 21–29 September 1995, Aalborg, Denmark. Cooperative Research Report No. 224. Carlton, J. T. (Ed.); International Council for the Exploration of the Sea, Copenhagen. Gronovius, L.T., 1781. *Zoophylacii Gronoviani*. Theodorus Haak et Soc., Leyden.
- Gruber, L., 2007. *Population Dynamics of Two Larger Symbiont-Bearing Foraminifera as Environmental Indicators of the Carmel Coast*. MSc. Thesis. University of Haifa Leon Recanati Institute for Maritime Studies, Haifa, Israel, 90 pp (In Hebrew, English abstract).
- Guastella, R., Marchini, A., Caruso, A., Cosentino, C., Evans, J. et al., 2019. “Hidden invaders” conquer the Sicily Channel and knock on the door of the Western Mediterranean Sea. *Estuarine, Coastal and Shelf Science*, 225, 1-12.
- Gudmundsson, G., 1994. Phylogeny, ontogeny and systematics of recent Soritacea Ehrenberg 1839 (Foraminiferida). *Micropaleontology*, 40, 101-155.
- Guy-Haim, T., Hyams-Kaphzan, O., Yeruham, E., Almogi-Labin, A., Carlton, J.T. 2017. A novel marine bioinvasion vector: Ichthyochory, live passage through fish. *Limnology and Oceanography Letters*, 2 (3), 81-90.
- Hallock, P., 1981. Production of carbonate sediments by selected large benthic foraminifera on two Pacific coral reefs. *Journal of Sedimentary Research*, 51 (2), 467-474.
- Hallock, P., 2002. Larger foraminifera as contributors to carbonate beach sands. p. 97-98. In: *Carbonate Beaches 2000*. Robbins, L.L., Magoon, O.T., Ewing, L. (Eds.). ASCE American Society of Civil Engineers Conference Proceedings, Reston, VA.
- Hansen, H.J., Rögl, F., 1980. On *Anomalina punctulata* d’Orbigny, 1826. *Journal of Foraminiferal Research*, 10 (2), 153-155.
- Haunold, T.G., Baal, C., Piller, W.E., 1997. Benthic foraminiferal associations in the northern Bay of Safaga, Red Sea, Egypt. *Marine Micropaleontology*, 29 (3-4), 185-210.
- Hayward, B.W., 1997. Introduced marine organisms in New Zealand and their impact in the Waitemata Harbour, Auckland. *Tane*, 36, 197-223.
- Hayward, B.W., Hollis, C.J., Grenfell, H.R., 1999. *Recent Elphidiidae (Foraminiferida) of the south-west Pacific and fossil Elphidiidae of New Zealand*. Institute of Geological and Nuclear Sciences, Lower Hutt.
- Hayward, B.W., Holzmann, M., Grenfell, H.R., Pawlowski, J., Triggs, C.M., 2004. Morphological distinction of molecular types in *Ammonia* – towards a taxonomic revision of the world’s most commonly misidentified foraminifera. *Marine Micropaleontology*, 50, 237-271.
- Hayward, B.W., Grenfell, H.R., Sabaa, A.T., Neil, H., Buzas, M.A., 2010. *Recent New Zealand deep-water benthic foraminifera: taxonomy, ecologic distribution, biogeography, and use in paleoenvironmental assessment*. GNS Science monograph, Lower Hutt, 363 pp.
- Heron-Allen, E., Earland, A., 1915. The Foraminifera of the Kerimba Archipelago (Portuguese East Africa). - Part II. *Transactions of the Zoological Society of London*, 20, 543-798.
- Hofker, J., 1930. *Foraminifera of the Siboga Expedition; Part 2; Families Astrorhizidae, Rhizamidae, Rheopacidae, Anomalinidae, Peneroplidae*. Siboga Expeditie, Monographie IVa. Brill, Leiden, 170 pp.
- Hofker, J., 1979. Rare and remarkable foraminifera of the Caribbean Sea. *Studies on the Fauna of Curaçao and other Caribbean Islands*, 58 (1), 1-43.
- Hohenegger, J., 2000. Coenoclines of larger foraminifera. *Micropaleontology*, 46 (1), 127-151.
- Hohenegger, J., 2006. The importance of symbiont-bearing foraminifera for West Pacific carbonate beach environments. *Marine Micropalaeontology*, 61, 4-39.
- Hollaus, S.S., Hottinger, L., 1997. Temperature dependence of endosymbiotic relationships? Evidence from the depth range of Mediterranean *Amphistegina lessonii* (Foraminiferida) truncated by the thermocline. *Eclogae geologicae Helveticae*, 90 (3), 591-598.
- Hottinger, L., 1977. Distribution of larger Peneroplidae, *Borelis* and Nummulitidae in the Gulf of Elat, Red Sea. *Utrecht Micropaleontological Bulletin*, 15, 35-110.
- Hottinger, L., Halicz, E., Reiss, Z., 1993. *Recent foraminiferida from the Gulf of Aqaba, Red Sea*. Slovenska Akademija Znanosti in Umetnosti, Ljubljana, 179 pp.

- Huth, L., 2016. *Status of non-indigenous aquatic species in the Harbor of Otranto (Mediterranean Sea): A survey on the diversity and composition of foraminiferal assemblages*. BSc Thesis. Steinmann Institut für Geologie, Paläontologie und Mineralogie der Rheinischen Friedrich-Wilhelms-Universität, Bonn, 41 pp.
- Hyams, O., 2000. *Benthic foraminifera from the Mediterranean inner shelf (to 40 m) of Israel*. MSc. Thesis. Ben-Gurion University of the Negev, Israel, 92 pp. (In Hebrew, English abstract).
- Hyams, O., 2006. *Eutrophication affecting benthic foraminifera on the oligotrophic eastern Mediterranean shallow shelf*. Ph.D. Dissertation, Ben-Gurion University of the Negev, Beer Sheva, Israel, 157 pp.
- Hyams, O., Almogi-Labin, A., Benjaminia, C., 2002. Larger foraminifera of the southeastern Mediterranean shallow continental shelf off Israel. *Israel Journal of Earth Sciences*, 51, 169-179.
- Hyams-Kaphzan, O., 2016. *Eutrophication Affecting Benthic Foraminifera on the Oligotrophic Eastern Mediterranean Shallow Shelf*. Geological Survey of Israel Report, GSI/07/2016, 176 pp.
- Hyams-Kaphzan, O., Almogi-Labin, A., Sivan, D., Benjamini, C., 2008. Benthic foraminifera assemblage change along the southeastern Mediterranean inner shelf due to fall-off of Nile-derived siliciclastics. *Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen*, 248 (3), 315-344.
- Hyams-Kaphzan, O., Almogi-Labin, A., Benjamini, C., Herut, B., 2009. Natural oligotrophy vs. pollution-induced eutrophy on the SE Mediterranean shallow shelf (Israel): environmental parameters and benthic foraminifera. *Marine Pollution Bulletin*, 58 (12), 1888-1902.
- Hyams-Kaphzan, O., Perelis Grossowicz, L., Almogi-Labin, A., 2014. *Characteristics of benthic foraminifera inhabiting rocky reefs in northern Israeli Mediterranean shelf*. Geological Survey of Israel Report, GSI/36/2014, 31 pp.
- Jensen, H.M., Panagiotidis, P., Reker, J., 2017. *Delineation of the MSFD Article 4 Marine Regions and Subregions. Version 1.0*. Technical document, ICES & EEA, 21 pp.
- Johengen, T., Reid, D., Fahnenstiel, G., MacIsaac, H., Dobbs, F. et al., 2005. *Assessment of transoceanic NOBOB vessels and low salinity ballast water as vectors for nonindigenous species introductions to the Great Lakes*. Cooperative Institute for Limnology and Ecosystems Research, School of Natural Resources and Environment, University of Michigan, 287 pp.
- Jones, R.W. 1994. *The Challenger Foraminifera*. Oxford University Press, 149 pp.
- Jones, R.W., Simmons, M.D., Whittaker, J.E., 2006. On the stratigraphical and palaeobiogeographical significance of *Borelis melo melo* (Fichtel & Moll, 1978) and *B. melo curdica* (Reichel, 1937) (Foraminifera, Miliolida, Alveolinidae). *Journal of Micropalaeontology*, 25 (2), 175-185.
- Jones, T.R., Parker, W.K., 1860. On the rhizopodal fauna of the Mediterranean, compared with that of the Italian and some other Tertiary deposits. *Quarterly Journal of the Geological Society*, 16 (1-2), 292-307.
- Jorissen, F.J., 1987. The distribution of benthic foraminifera in the Adriatic Sea. *Marine Micropaleontology*, 12, 21-48.
- Katsanevakis, S., Bogucarskis, K., Gatto, F., Vandekerckhove, J., Deriu, I. et al., 2012. Building the European Alien Species Information Network (EASIN): a novel approach for the exploration of distributed alien species data. *BioInvasions Records*, 1 (4), 235-245.
- Katsanevakis, S., Coll, M., Piroddi, C., Steenbeek, J., Ben Rais Lasram, F. et al., 2014a. Invading the Mediterranean Sea: biodiversity patterns shaped by human activities. *Frontiers in Marine Science*, 1, 32.
- Katsanevakis, S., Acar, Ü., Ammar, I., Balci, B.A., Bekas, P. et al., 2014b. New Mediterranean Biodiversity Records (October, 2014). *Mediterranean Marine Science*, 15 (3), 675-695.
- Katsanevakis, S., Zenetos, A., Corsini-Foka, M., Tsiamis, K., 2020. Biological Invasions in the Aegean Sea: Temporal Trends, Pathways, and Impacts. In: *The Handbook of Environmental Chemistry*. Springer, Berlin, Heidelberg.
- Koukousioura, O., Dimiza, M.D., Triantaphyllou, M.V., 2010. Alien foraminifera from Greek coastal areas (Aegean Sea, Eastern Mediterranean). *Mediterranean Marine Science*, 11 (1), 155-172.
- Koukousioura, O., Dimiza, M.D., Triantaphyllou, M.V., Hallock, P., 2011. Living benthic foraminifera as an environmental proxy in coastal ecosystems: A case study from the Aegean Sea (Greece, NE Mediterranean). *Journal of Marine Systems*, 88 (4), 489-501.
- Koukousioura, O., Triantaphyllou, M.V., Dimiza, M.D., Pavlopoulos, K., Syrides, G. et al., 2012. Benthic foraminiferal evidence and paleoenvironmental evolution of Holocene coastal plains in the Aegean Sea (Greece). *Quaternary International*, 261, 105-117.
- Lacroix, E., 1932. Textularidae du plateau continental méditerranéen entre Saint-Raphael et Monaco. *Bulletin de l'Institut Océanographique Monaco*, 591, 1-28.
- Lacroix, E., 1941. Les Orbitolites du Golfe d'Akaba. *Bulletin de l'Institut Océanographique, Monaco*, 794, 792-799.
- Lamourou, A., Touir, J., Fagel, N., 2017. Reconstructing the Holocene depositional environments along the northern coast of Sfax (Tunisia): mineralogical and sedimentological approaches. *Journal of African Earth Sciences*, 129, 713-727.
- Langer, M.R., 1992. New recent foraminiferal genera and species from the lagoon at Madang, Papua New Guinea. *Journal of Micropaleontology*, 11 (1), 85-93.
- Langer, M.R., 1993. Epiphytic Foraminifera. *Marine Micropaleontology*, 2, 235-265.
- Langer, M.R., 2008. Foraminifera from the Mediterranean and the Red Sea. p. 399-417. In: *Aqaba-Eilat, the Improbable Gulf: Environment, Biodiversity and Preservation*. Por, F.D. (Ed). Magnes Press, Jerusalem.
- Langer, M.R., Bell, C., 1995. Toxic Foraminifera: Innocent Until Proven Guilty. *Marine Micropaleontology*, 24, 205-214.
- Langer, M.R., Lipps, J.H., 2006. Assembly and persistence of foraminifera in introduced mangroves on Moorea, French Polynesia. *Micropaleontology*, 52, 343-355.
- Langer, M.R., Hottinger, L., 2000. Biogeography of selected "larger" foraminifera. *Micropaleontology*, 46 (1), 105-126.
- Langer, M.R., Mouanga, G.H., 2016. Invasion of amphisteginid foraminifera in the Adriatic Sea. *Biological Invasions* 18 (1335).
- Langer, M.R., Schmidt-Sinns, J., 2006. *The 100 most common*

- Foraminifera from the Bay of Fetovaia, Elba Island (Mediterranean Sea)*. Steinmann Institut für Geologie, Paläontologie und Mineralogie der Rheinischen Friedrich-Wilhelms-Universität, Bonn, 37 pp.
- Langer, M.R., Silk, M.T., Lipps, J.H., 1997. Global ocean carbonate and carbon dioxide production: The role of reef foraminifera. *Journal of Foraminiferal Research*, 27, 271-277.
- Langer, M.R., Frick, H., Silk, M.T., 1998. Photophile and sciaphile foraminifera from Lavezzi Island, Corsica, *Revue de Paléobiologie*, 17(2), 525-530.
- Langer, M.R., Makled, W., Pietsch, S.J., Weinmann, A.E., 2009. Asynchronous calcification in embryonic megalospheres from Chuuk Island (Micronesia): An ontogenetic window into the life cycle and polymorphism of the foraminifer *Peneroplis* sp. *Journal of Foraminiferal Research*, 39 (1), 8-14.
- Langer, M.R., Weinmann, A.E., Lötters, S., Rödder, D., 2012. "Strangers" in paradise: Modeling the biogeographic range expansion of the foraminifera *Amphistegina* in the Mediterranean Sea. *Journal of Foraminiferal Research*, 42 (3), 234-244.
- Langer, M.R., Thissen, J.M., Makled, W.A., Weinmann, A.E., 2013a. The foraminifera from the Bazaruto Archipelago (Mozambique). *Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen*, 267 (2), 155-170.
- Langer, M.R., Weinmann, A.E., Lötters, S., Bernhard, J.M., Rödder, D., 2013b. Climate-Driven Range Extension of *Amphistegina* (Protista, Foraminiferida): Models of Current and Predicted Future Ranges. *PloS One*, 8 (2), e54443.
- Larsen, A.R., 1976. Studies of recent *Amphistegina*, taxonomy and some ecological aspects. *Israel Journal of Earth-Sciences*, 25, 1-26.
- Larsen, A.R., 1977. A neotype of *Amphistegina lessonii* d'Orbigny, 1826. *The Journal of Foraminiferal Research*, 7 (4), 273-277.
- Lavoie, D.M., Smith, L.D., Ruiz, G.M., 1999. The potential for introcoastal transfer of nonindigenous species in the ballast water of ships. *Estuarine Coastal and Shelf Science*, 48, 551-564.
- Lazar, S., 2007. *Recent and late Pleistocene carbonate-rich sediments in the Mediterranean shelf of Israel: Sedimentary, biogenic and genetic analysis*. Geological Survey of Israel Report GSI/08/2007.
- Le Calvez, J., Le Calvez, Y., 1958. Répartition des foraminifères dans la Baie de Villefrance I. Miliolidae. *Annales de l'Institut Océanographique Monaco*, 35, 159-234.
- Le Calvez, Y., 1977. *Révision des foraminifères de la collection d'Orbigny: Foraminifères de L'Île de Cuba*. I. Muséum national d'Histoire naturelle, Paris, 127 pp.
- Lipps, J.H., 1983. Biotic Interactions in Benthic Foraminifera. In: Tevesz M.J.S., McCall P.L. (eds.) *Biotic Interactions in Recent and Fossil Benthic Communities*. *Topics in Geobiology*, 3. Springer, Boston, MA.
- Lipps, J.H., Langer, M.R., 1999. Benthic Foraminifera from the meromitic, Mecherchar Jellyfish Lake, Palau (western Pacific). *Micropaleontology*, 45, 278-284.
- Loeblich, A.R., Tappan, H., 1955. Revision of some recent foraminiferal genera. *Smithsonian Miscellaneous Collections*, 128.
- Loeblich, A.R., Tappan, H., 1957. Morphology and taxonomy of the foraminiferal genus *Pararotalia* Le Calvez, 1949. *Smithsonian Miscellaneous Collections*, 135.
- Loeblich, A.R., Tappan, H., 1964. *Treatise on Invertebrate Paleontology: Part C, Protista 2, Sarcodina, Chiefly Thecamoebians and Foraminiferida*. Geological Society of America, Incorporated and University of Kansas Press.
- Loeblich, A.R., Tappan, H., 1987. *Foraminiferal genera and their classification*. Nostrand Reinhold Company, New York, 970 pp.
- Loeblich, A.R., Tappan, H., 1994. Foraminifera of the Sahul Shelf and Timor Sea. *Cushman Foundation of Foraminiferal Research Special Publication*, 31, 1-661.
- Macdonald, E.M., 1998. Dinoflagellate resting cysts and ballast water discharge in Scottish ports. p. 24-35. In: *Ballast Water: Ecological and Fisheries Implications: ICES Conference (83rd Statutory Meeting), 21-29 September 1995, Aalborg, Denmark. Cooperative Research Report No. 224*. Carlton, J. T. (Ed.): International Council for the Exploration of the Sea, Copenhagen.
- Makled, W.A., Langer, M.R., 2011. Benthic Foraminifera from the Chuuk Lagoon Atoll System (Caroline Islands, Pacific Ocean). *Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen*, 259 (2), 231-249.
- Makled, W.A., Soliman, S.I., Langer, M.R., 2017. Manual of Neogene foraminifera from off-shore Nile Delta (Egypt) and their biostratigraphic significance. *Marine and Petroleum Geology*, 80, 450-491.
- Manda, S., Titelboim, D., Ashckenazi-Polivoda, S., Almogi-Labin, A., Herut, B. *et al.*, 2020. Epiphytic benthic foraminiferal preferences for macroalgal habitats: Implications for coastal warming. *Marine Environmental Research*, 161, 105084.
- Marras, S., Cucco, A., Antognarelli, F., Azzurro, E., Milazzo, M. *et al.*, 2015. Predicting future thermal habitat suitability of competing native and invasive fish species: From metabolic scope to oceanographic modelling. *Conservation Physiology*, 3, 1-14.
- Marriner, N., Morhange, C., Boudagher-Fadel, M., Bourcier, M., Carbonel, P. *et al.*, 2005. Geoarchaeology of Tyre's ancient northern harbour, Phoenicia. *Journal of Archaeological Science*, 32 (9), 1302-1327.
- Martinotti, A., 1920. Foraminiferi della Spiaggia di Tripoli. *Atti della Società italiana di scienze naturali e del Museo civico di storia naturale di Milano*, 59, 249.
- Martins, M.V.A., Helali, M.A., Zaaboub, N., Boukef-BenOmrane, I., Frontalini, F. *et al.*, 2016. Organic matter quantity and quality, metals availability and foraminiferal assemblages as environmental proxy applied to the Bizerte Lagoon (Tunisia). *Marine Pollution Bulletin*, 105 (1), 161-179.
- Mateu, G., 1972. Les foraminifères du Tyrrhénien de la baie de Palma de Majorque et les conditions bioécologiques de l'ancienne mer des Baléares. *Rapports et Procès-Verbaux des Réunions Conseil Internationale pour l'Exploration Scientifique de la Mer Méditerranée*, 20 (4), 539-540.
- Mateu-Vicens, G., Holzmann, M., Nadal, M.V., Khokhlova, A., Ferriol, P. *et al.*, 2018. First occurrence of the large benthic foraminifer *Parasorites* sp. in the Mediterranean. FORAMS 2018, Program with Abstracts, Edinburgh 17-22 June.
- McCulloch, I.A., 1977. *Qualitative observations on recent for-*

- aminiferal tests with emphasis on the Eastern Pacific*. University of southern California, Los Angeles.
- McGann, M., Sloan, D., Cohen, A.N., 2000. Invasion by a Japanese marine microorganism in western North America. *Hydrobiologia*, 421, 25-30.
- McGann, M., Ruiz, G.M., Hines, A.H., Smith, G., 2020. A ship's ballasting history as an indicator of foraminifer invasion potential – An example from Prince William Sound, Alaska, USA. *Journal of Foraminiferal Research*, 49 (4), 434-455.
- Melis, R., Bernasconi, M.P., Colizza, E., Di Rita, F., Schneider, E.E. *et al.*, 2015. Late Holocene palaeoenvironmental evolution of the northern harbour at the Elaiussa Sebaste archaeological site (south-eastern Turkey): Evidence from core ELA6. *Turkish Journal of Earth Sciences*, 24 (6), 566-584.
- Meriç, E., Avşar, N., Bergin, F., 2004. *Benthic foraminifera of eastern Aegean Sea (Turkey): Systematics and autoecology*. Turkish Marine Research Foundation, Istanbul, 306 pp.
- Meriç, E., Avşar, N., Yokeş, M.B., 2007. Alien foraminifers along the Aegean and southwestern coasts of Turkey. *Rapports et procès-verbaux des réunions - Commission internationale pour l'exploration scientifique de la Mer Méditerranée*, 38, 540.
- Meriç, E., Avşar, N., Nazik, A., Yokeş, M. B., Dinçer, F., 2008a. A review of benthic foraminifers and ostracodes of the Antalya coast. *Micropaleontology*, 54, 199-240.
- Meriç, E., Avşar, N., Yokeş, M.B., 2008b. Some Alien Foraminifers along the Aegean and Southwestern Coasts of Turkey. *Micropaleontology*, 54 (3-4), 307-349.
- Meriç, E., Yokeş, M.B., Avşar, N., Bircan, C., 2010. An oasis for alien benthic Foraminifera in the Aegean Sea. *Aquatic Invasions*, 5 (2), 191-195.
- Meriç, E., Yokeş, M.B., Avşar, N., 2011. A new guest in Ilica Bay (Çeşme-Izmir-Turkey), *Coscinospira acicularis*. *Marine Biodiversity Records*, 4, 1-5.
- Meriç, E., Yokeş, M.B., Avşar, N., Kirçi-Elmas, E., Dinçer, F. *et al.*, 2013. First report of *Pararotalia calcariformata* from the Hatay coastline (Turkey-north-eastern Mediterranean). *Marine Biodiversity Records*, 6.
- Meriç, E., Niyazi, A., Yokeş, M.B., Dinçer, F., 2014. Atlas of Recent Benthic foraminifera from Turkey. *Micropaleontology*, 60, 211-398.
- Meriç, E., Avşar, N., Yokeş, M.B., Dinçer, F., Bircan, C. *et al.*, 2020. Ecology of benthic foraminifera of the eastern Aegean Sea coasts. *International Journal of Environment and Geoinformatics (IJECEO)*, 7 (1), 6-22.
- Merkado, G., 2016. *Breaking Biogeographic Barriers: Molecular and Morphologic Evidences for the Lessepsian Invasion of Foraminifera to the Mediterranean*. PhD Thesis. Ben-Gurion University of the Negev, Faculty of Natural Sciences, Department of Geological and Environmental Sciences, Israel, 98 pp.
- Merkado, G., Holzmann, M., Apothéloz-Perret-Gentil, L., Pawlowski, J., Abdu, U. *et al.*, 2013. Molecular evidence for Lessepsian invasion of Soritids (larger symbiont bearing benthic foraminifera). *PLoS ONE*, 8 (10), 1-12.
- Merkado, G., Titelboim, D., Hyams-Kaphzan, O., Holzmann, M., Pawlowski, J. *et al.*, 2015. Molecular Phylogeny and Ecology of *Textularia agglutinans* d'Orbigny from the Mediterranean Coast of Israel: A Case of a Successful New Incumbent. *PLoS ONE*, 10 (11).
- Milker, Y., Schmiedl, G., 2012. A taxonomic guide to modern benthic shelf foraminifera of the western Mediterranean Sea. *Palaeontologia Electronica*, 15 (2), 1-134.
- Millett, F.W., 1898. Report on the recent foraminifera of the Malay Archipelago collected by Mr. A. Durrand, F.R.M.S. – Part II. *Journal of the Royal Microscopical Society*, 18 (5), 499-513.
- Millett, F.W., 1900. Report on the recent foraminifera of the Malay Archipelago collected by Mr. A. Durrand, F.R.M.S. – Part IX. *Journal of the Royal Microscopical Society*, 20 (5), 539-549.
- Mkawar, S., Azri, C., Kamoun, F., Montacer, M., 2007. Impact sur les biophases marines des rejets anthropiques, notamment des métaux lourds rejetés sur le littoral nord de la ville de Sfax (Tunisie). *Techniques Sciences Méthodes*, 10, 71-85.
- Möbius, K.A., 1880. *Beiträge zur Meeresfauna der Insel Mauritius und der Seychellen*. Verlag der Gutmann'schen Buchhandlung (Otto Enslin), Berlin, 352 pp.
- Mojtahid, M., Jorissen, F., Lansard, B., Fontanier, C., Bombléd, B. *et al.*, 2009. Spatial distribution of live benthic foraminifera in the Rhône prodelta: Faunal response to a continental-marine organic matter gradient. *Marine Micropaleontology*, 70 (3-4), 177-200.
- Molnar, J.L., Gamboa, R.L., Revenga, C., Spalding, M.D., 2008. Assessing the global threat of invasive species to marine biodiversity. *Frontiers in Ecology and the Environment*, 6, 485-492.
- Moncharmont Zei, M., 1968. I foraminiferi di alcuni campioni di fondo prelevati lungo la costa di Beirut (Libano). *Bollettino della Società dei Naturalisti in Napoli*, 77, 3-34.
- Montfort, D. de, 1808. *Conchyliologie systématique: Coquilles univalves cloisonnées*. F. Schoell, Paris, 408 pp.
- Morhange, C., Goiran, J.P., Bourcier, M., Carbonel, P., Le Campion, J. *et al.*, 2000. Recent Holocene paleo-environmental evolution and coastline changes of Kition, Larnaca, Cyprus, Mediterranean Sea. *Marine Geology*, 170 (1-2), 205-230.
- Mouanga, G.H., 2017. *Impact and Range Extension of Invasive Foraminifera in the NW Mediterranean Sea: Implications for Diversity and Ecosystem Functioning*, PhD Thesis. Bonn: Steinmann-Institut für Geologie, Mineralogie und Paläontologie.
- Mouanga, G.H., Langer, M.R., 2014. At the front of expanding ranges: Shifting community structures at amphisteginid species range margins in the Mediterranean Sea. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 271 (2), 141-150.
- Nadal Nebot, V., 2017. *L'alga verda Halimeda incrassata com a possible vector d'introducció d'epibionts exòtics en la Badia de Palma*. Facultat de Ciències, Universitat de les Illes Balears, Palma, 30 pp.
- Nesbitt, E.A., 2005. A novel trophic relationship between caudo gastropods and mysticete whale carcasses. *Lethaia*, 38, 17-25.
- Oflaz, S.A., 2006. *Taxonomy and distribution of the benthic foraminifera in the Gulf of Iskenderun, Eastern Mediterranean*. MSc Thesis, Graduate School of Natural and Applied Sciences Middle East Technical University, Ankara, 306 pp (In Turkish, English abstract).

- Ojaveer, H., Galil, B.S., Campbell, M.L., Carlton, J.T., Canning-Clode, J. *et al.*, 2015. Classification of non-indigenous species based on their impacts: Considerations for application in marine management. *PLoS Biology*, 13, e1002130.
- Ojaveer, H., Galil, B.S., Carlton, J.T., Allevay, H., Gouletquer, P., Lehtiniemi, M. *et al.*, 2018. Historical baselines in marine bioinvasions: Implications for policy and management. *PLoS One*, 13, e0202383.
- Orbigny, A.D.d', 1826. Tableau Méthodique de la classe des Céphalopodes. *Annales des Sciences Naturelles. Series*, 1 (7), 96-314.
- Orbigny, A.D.d', 1839. Foraminifères. p. 1-224. In: *Histoire physique, politique et naturelle de L'Ile de Cuba*. Sagra, M.R. de la (Ed). Arthus Bertrand, Paris.
- Orbigny, A.D.d', 1846. *Foraminifères fossiles du Bassin tertiaire de Vienne (Autriche)*. Von Gide et comp., Paris, 312 pp.
- Parker, F.L., 1958. Eastern Mediterranean foraminifera. *Reports of the Swedish Deep-sea Expedition*, 8 (4), 219-285. Elanders boktr, Goteborg.
- Parker, J.H., 2009. Taxonomy of foraminifera from Ningaloo Reef, Western Australia. *Memoirs of the Association of Australasian Palaeontologists*, 36.
- Parker, K.W., Jones, T.R., 1865. On some Foraminifera from the North Atlantic and Arctic Oceans, including Davis Straits and Baffin's Bay. *Philosophical Transactions of the Royal Society of London*, 155, 325-441.
- Parker, K.W., Jones, T.R., Brady, H.B., 1865. On the nomenclature of the Foraminifera. Part X. The species enumerated by d'Orbigny in the "Annales des Sciences Naturelles". *The Annals and magazine of natural history; Zoology, Botany, and Geology. Series 3*, 16, 15-41.
- Parr, W.J., 1932. Victorian and South Australian shallow-water foraminifera. Part I. *Proceedings of the Royal Society of Victoria*, 44, 1-15.
- Parr, W.J., 1950. Foraminifera. *B. A. N. Z. Antarctic Research Expedition 1929-1931 under the command of Sir Douglas Mawson. Reports - Series B (Zoology and Botany)*, 5, 233-392.
- Patterson, R.T., Richardson, R.H., 1987. A taxonomic revision of the unilocular foraminifera. *Journal of Foraminiferal Research*, 17 (3), 212-226.
- Pawlowski, J., Lejzerowicz, F., Esling, P., 2014. Next-generation environmental diversity surveys of foraminifera: Preparing the future. *Biological Bulletin*, 227, 93-106.
- Pereira, C.P.G., 1979. *Foraminiferal distribution and ecology in the fringing reef complex of the coast, near Mombasa, Kenya*. PhD Thesis. University College of Swansea, University of Wales.
- Piller, W.E., Haunold, T.G., 1998. The Northern Bay of Safaga (Red Sea, Egypt): An actuopalaeontological approach V. Foraminifera. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft*, 548.
- Polovodova Asteman, I., Schönfeld, J., 2016. Recent invasion of the foraminifer *Nonionella stella* Cushman and Moyer, 1930 in northern European waters: Evidence from the Skagerrak and its fjords. *Journal of Micropalaeontology*, 35, 20-25.
- Prazeres, M., Martínez-Colón, M., Hallock, P., 2020. Foraminifera as bioindicators of water quality: The FoRAM Index revisited. *Environmental Pollution*, 257, 113612.
- Pyšek P., Hulme, P.E., Simberloff, D., Bacher, S., Blackburn, T.M. *et al*, 2020. Scientists' warning on invasive alien species. *Biological Reviews*.
- Raitsos, D.E., Beaugrand, G., Georgopoulos, D., Zenetos, A., Pancucci-Papadopoulou, M.A., Theocharis, A., Papatthanassiou, E., 2010. Global climate change amplifies the entry of tropical species into the eastern Mediterranean Sea. *Limnology and Oceanography*, 55, 1478-1484.
- Rech, S., Borrell, Y., García-Vazquez, E., 2016. Marine litter as a vector for non-native species: What we need to know. *Marine Pollution Bulletin*, 113, 40-43.
- Reichel, M., 1937. Étude sur les Alvéolines, II. *Schweizerische Paläontologische Abhandlung*, 59 (3), 95-147.
- Reinhardt, E.G., Patterson, R.T., Schröder-Adams, C.J., 1994. Geoarchaeology of the ancient harbor site of Caesarea Maritima, Israel: Evidence from sedimentology and paleoecology of benthic foraminifera. *Journal of Foraminiferal Research*, 24 (1), 37-48.
- Reiss, Z., Gvirtzman, G., 1966. *Borelis* from Israel. *Eclogae Geologicae Helvetiae*, 59, 437-447.
- Reiss Z, Leutenegger, S, Hottinger, L, Fermont, W.J.J., Meulenkamp, J.E., Thomas, E., Hansen, H.J., Buchardt, B., Larsen, A.R., Drooger, C.W., 1977. Depth-relations of recent larger foraminifera in the Gulf of Aqaba-Elat. *Utrecht Micropaleontological Bulletin*, 15, 1-244.
- Revs, S.A., 1996. The generic revision of five families of rotaliine foraminifera: Part I. The Bolivinitidae. *Cushman Foundation for Foraminiferal Research Special Publication*, 34, 1-55.
- Ricciardi, A., 2016. Tracking marine alien species by ship movements. *PNAS*, 113(20), 5470-5471.
- Rocha, A.T., Mateu, G., 1971. *Contribuição para o conhecimento dos foraminíferos actuais da ilha de Maio (Arquipélago de Cabo Verde)*. Instituto de investigação científica de Angola, Angola, 108 pp.
- Rückert-Hilbig, A., 1983. Megalospheric gamonts of *Rosalina globularis*, *Cymbaloporeta bulloides* and *Cymbaloporeta millesti* (Foraminifera) with differently constructed swimming-apparatus. *Tübinger Mikropaläontologische Mitteilungen*, 1, 1-69.
- Ruiz, G.M., Carlton, J.T., Grosholz, E.D., Hines, A.H., 1997. Global invasions of marine and estuarine habitats by non-indigenous species: Mechanisms, extent, and consequences. *American Zoologist*, 37 (6), 621-632.
- Said, R., 1949. Foraminifera of the Northern Red Sea. *Cushman Laboratory for Foraminiferal Research Special Publication*, 26, 1-44.
- Said, R., 1950. Additional foraminifera from the Northern Red Sea. *Contributions from the Cushman Foundation for Foraminiferal Research*, 1, 4-9.
- Samir, A.M., El-Din, A.B., 2001. Benthic foraminiferal assemblages and morphological abnormalities as pollution proxies in two Egyptian bays. *Marine Micropaleontology*, 41 (3-4), 193-227.
- Samir, A.M., Abdou, H.F., Zazou, S.M., El-Menhawey, W.H., 2003. Cluster analysis of recent benthic foraminifera from the northwestern Mediterranean coast of Egypt. *Revue de Micropaléontologie*, 46 (2), 111-130.
- Schaudinn, F., 1911. Notizen über die Fauna bei Rovigno: I: Verzeichnis von Foraminiferen aus Rovigno. *Zoologischer*

- Anzeiger*, 37, 254-256.
- Schmidt, C., Morard, R., Almogi-Labin, A., Weinmann, A.E., Titelboim, D. *et al.*, 2015. Recent invasion of the symbiont-bearing foraminifera into the Eastern Mediterranean facilitated by the ongoing warming trend. *PLoS ONE*, 10 (8), e0132917.
- Schmidt, C., Morard, R., Romero, O., Kucera, M., 2018. Diverse internal symbiont community in the endosymbiotic Foraminifera *Pararotalia calcariformata*: implications for symbiont shuffling under thermal stress. *Frontiers in Microbiology*, 9.
- Schwager, C., 1866. Fossile Foraminiferen von Kar Nikobar. *Reise der Österreichischen Fregatte Novara um die Erde in den Jahren 1857, 1858, 1859 unter den Befehlen des Commodore B. von Wüllerstorff-Urbair. Geologischer Theil (Zweite Abtheilung, Paläontologische Mittheilungen)*, 2, 187-268.
- Seebens, H., Schwartz, N., Schupp, P.J., Blasius, B., 2016. Predicting the spread of marine species introduced by global shipping. *Proceedings of the National Academy of Sciences*, 113 (20), 5646-5651.
- Seebens, H., Blackburn, T., Dyer, E. E., Genovesi, P., Hulme, P.E., *et al.*, 2017. No saturation in the accumulation of alien species worldwide. *Nature Communications*, 8, 14435.
- Serandrei-Barbero, R., Donnici, S., Madricardo, F., 2011. Supratidal foraminifera as ecological indicators in anthropically modified wetlands (Lagoon of Venice, Italy). *Ecological Engineering*, 37 (8), 1140-1148.
- Servello, G., Andaloro, F., Azzurro, E., Castriota, L., Catra, M. *et al.*, 2019. Marine alien species in Italy: A contribution to the implementation of descriptor D2 of the marine strategy framework directive. *Mediterranean Marine Science*, 20 (1), 1-48.
- Sgarrella, F., Moncharmont Zei, M., 1993. Benthic Foraminifera of the Gulf of Naples (Italy): systematics and autoecology. *Bollettino della Società Paleontologica Italiana*, 32 (2), 145-264.
- Sidebottom, H., 1904. Report on the recent foraminifera from the coast of the Island of Delos (Grecian Archipelago), Part 1. *Memoirs and Proceedings of the Manchester Literary and Philosophical Society*, 48 (5), 1-26.
- Sidebottom, H., 1909. Report on the recent foraminifera from the coast of the Island of Delos (Grecian Archipelago), Part 6. *Memoirs and Proceedings of the Manchester Literary and Philosophical Society*, 53 (21), 1-32.
- Sidebottom, H., 1910. Report on the recent foraminifera from the Bay of Palermo, Sicily, 14--20 fms. (Off the harbour). *Memoirs and Proceedings of the Manchester Literary and Philosophical Society*, 54 (16), 1-28.
- Silvestri, A., 1907. Considerazioni paleontologiche e morfologiche sui generi *Operculina*, *Heterostegina*, *Cycloclypeus*. *Bollettino della Società Geologica Italiana*, 26 (1), 29-62.
- Silvestri, A., 1950. I Foraminiferi della Laguna Veneta. *Bollettino di Pesca, Piscicoltura e di Idrobiologia*, 5, 22-98.
- Skliris, N., Lascaratos, A., 2004. Impacts of the Nile River damming on the thermohaline circulation and water mass characteristics of the Mediterranean Sea. *Journal of Marine Systems*, 52 (1-4), 121-143.
- Siokou, I., Ates, A.S., Ayas, D., Ben Souissi, J., Chatterjee, T., *et al.*, 2013. New Mediterranean Marine biodiversity records. *Mediterranean Marine Science* 14 (1), 238-249.
- Smith, L.D., Wonham, M.J., McCann, L.D., Ruiz, G.M., Hines, A.H. *et al.*, 1999. Invasion pressure to a ballast flooded estuary and an assessment of inoculant survival. *Biological Invasions*, 1, 67-87.
- Smithsonian National Museum of Natural History Paleobiology Collections, 2020. <https://collections.nmnh.si.edu/search/paleo/> (Accessed 4 April 2020)
- Smitter, Y.H., 1955. A foraminiferal fauna from the Tertiary sediments of southern Moçambique. *Palaeontologica Africana*, 3, 109-118.
- Smout, A.H., Eames, F.E., 1958. The genus *Archaias* (foraminifera) and its stratigraphical distribution. *Palaeontology*, 1 (3), 207-225.
- Srinivasan, M.S., Sharma, V., 1980. *Schwager's Car Nicobar Foraminifera in the Reports of the Novara Expedition: a revision*. Today and Tomorrow's Pringers and Publishers, New Delhi.
- Tapiero, I., 2002. *High-resolution paleoecologic and paleoclimatic changes of the Holocene, based on benthic foraminifera and sediments from the Mediterranean inner shelf, Israel*. MSc Thesis. Ben-Gurion University of the Negev, 130 pp. (In Hebrew, English abstract).
- Tapiero, I., Almogi-Labin, A., Benjamini, C., 2003. Holocene climate and environmental variability based on benthic foraminifera and sediments from the inner shelf of the southeastern Mediterranean Sea. EGS - AGU - EUG Joint Assembly, Abstracts from the meeting held in Nice, France, 6 - 11 April 2003, abstract id. 9499.
- Thalman, H.E., 1950. New names and homonyms in Foraminifera. *Contributions from the Cushman Foundation for Foraminiferal Research*, 1 (3-4), 41-45.
- Tittensor, D.P., Mora, C., Jetz, W., Lotze, H.K., Ricard, D., Vanden Berghe, E., Worm, B., 2010. Global patterns and predictors of marine biodiversity across taxa. *Nature*, 466, 1098-1101.
- Thissen, J.M., Langer, M.R., 2017. Spatial patterns and structural composition of foraminiferal assemblages from the Zanzibar Archipelago (Tanzania). *Palaeontographica A (Palaeozoology, Stratigraphy)*, 308, 1-67.
- Titelboim, D., Almogi-Labin, A., Herut, B., Kucera, M., Schmidt, C. *et al.*, 2016. Selective responses of benthic foraminifera to thermal pollution. *Marine Pollution Bulletin*, 105 (1), 324-336.
- Toueg, R., 1996. *History of the Inner Harbor in Caesarea*. M.A. Thesis. Department of Maritime Civilizations, University of Haifa, Haifa, Israel, 139 pp. (In Hebrew, English abstract).
- Triantaphyllou, M.V., Tsourou, T., Koukousioura, O., Dermitzakis, M.D., 2005. Foraminiferal and ostracod ecological patterns in coastal environments of SE Andros Island (Middle Aegean Sea, Greece). *Revue de Micropaléontologie*, 48 (4), 279-302.
- Triantaphyllou, M.V., Koukousioura, O., Dimiza, M.D., 2009. The presence of Indo-Pacific symbiont bearing foraminifer *Amphistegina lobifera* in Greek coastal ecosystems (Aegean Sea, Eastern Mediterranean). *Mediterranean Marine Science*, 10 (2), 73-85.
- Triantaphyllou, M.V., Dimizak, M.D., Koukousioura, O., Hallock, P., 2012. Observations on the life cycle of the symbi-

- ont-bearing foraminifer *Amphistegina lobifera* Larsen, an invasive species in coastal ecosystems of the Aegean Sea (Greece, Mediterranean). *Journal of Foraminiferal Research*, 42 (2), 143-150.
- Tsiamis, K., Palialexis, A., Stefanova, K., Ninčević Gladan, Ž. *et al.*, 2019. Non-indigenous species refined national baseline inventories: A synthesis in the context of the European Union's Marine Strategy Framework Directive. *Marine Pollution Bulletin*, 145, 429-435.
- Tsiamis, K., Azzurro, E., Bariche, M., Çinar, M.E., Crocetta, F. *et al.*, 2020. Prioritizing marine invasive alien species in the European Union through horizon scanning. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 1-52.
- UNCTAD (United Nations Conference on Trade and Development), 2019. *Review of Maritime Transport*. United Nations Publications, Sales No. E.19.II.D.20, New York and Geneva, 132 pp.
- Vella, P., 1957. Studies in New Zealand foraminifera. Part I. Foraminifera from Cook Strait. *New Zealand Geological Survey*, 28, 1-43.
- Vitousek, P.M., D'Antonio, C.M., Loope, L.L., Westbrooks, R., 1996. Biological invasions as global environmental change. *American Scientist*, 84, 468-478.
- Weinmann, A.E., Rödder, D., Lötters, S., Langer, M.R., 2013. Traveling through time: The past, present and future biogeographic range of the invasive foraminifera *Amphistegina* spp. in the Mediterranean Sea. *Marine Micropaleontology*, 105, 30-39.
- Weinmann, A.E., Goldstein, S.T., Triantaphyllou, M.V., Langer, M.R., 2019. Effects of sampling site, season, and substrate on foraminiferal assemblages grown from propagule banks from lagoon sediments of Corfu Island (Greece, Ionian Sea). *PloS One* 14 (6), e0219015.
- Wiesner, H., 1911a. Notizen über die Fauna der Adria bei Rovigno. VI. Foraminifera von dem Sandgrunde der Bucht S. Pelagio bei Rovigno, in 3 m Tiefe. *Zoologischer Anzeiger*, 37, 478-480.
- Wiesner, H., 1911b. Notizen über die Fauna der Adria bei Rovigno. VI. Schalentragende Foraminiferen von der Westküste Istriens. *Zoologischer Anzeiger*, 38, 505-510.
- Wiesner, H., 1912. Zur Systematik adriatischer Nubecularien, Spiroloculinen, Miliolinen und Biloculinen. *Archiv für Protistenkunde*, 25 (2), 201-239.
- Wiesner, H., 1913. Notizen über die Fauna der Adria bei Rovigno: Die Foraminiferen aus den im Jahre 1911 gehobenen Grundproben. *Zoologischer Anzeiger*, 41, 521-528.
- Wiesner, H., 1923. *Die Miliolideen der östlichen Adria*. Prag-Bubeneč, 113 pp.
- WoRMS Editorial Board, 2020. *World Register of Marine Species*. <https://www.marinespecies.org> (Accessed 7 February 2020).
- Yalçın, H., Meriç, E., Avşar, N., Tetiker, S., Barut, İ. F. *et al.*, 2008. Mineralogical and geochemical features of colored benthic foraminifers from Aegean and southwestern coasts of Turkey. *Micropaleontology*, 54 (3), 351-370.
- Yanko, V., Kronfeld, J., Flexer, A., 1994. Response of benthic foraminifera to various pollution sources: implications for pollution monitoring. *Journal of Foraminiferal Research*, 24 (1), 1-17.
- Yanko, V., Ahmad, M., Kaminski, M., 1998. Morphological deformities of benthic foraminiferal tests in response to pollution by heavy metals: Implications for pollution monitoring. *Journal of Foraminiferal Research*, 28 (3), 177-200.
- Yokeş, M.B., Meriç, E., 2009. Drowning in the Sand: Invasion by Foraminifera. p. 7-20. In: *Invasive Species: Detection, Impact and Control*. Wilcox, C.P., Turpin, R.P. (Eds). Nova Science Publishers, Inc., New York.
- Yümün, Z.Ü., Meriç, E., Avşar, N., Nazik, A., Barut, İ.F. *et al.*, 2016. Meiofauna, microflora and geochemical properties of the late quaternary (Holocene) sediments in the Gulf of Izmir (Eastern Aegean Sea, Turkey). *Journal of African Earth Sciences*, 124, 383-408.
- Zaïbi, C., Kamoun, F., Viehberg, F., Carbonel, P., Jedoui, Y. *et al.*, 2016. Impact of relative sea level and extreme climate events on the Southern Skhira coastline (Gulf of Gabes, Tunisia) during Holocene times: Ostracodes and foraminifera associations' response. *Journal of African Earth Sciences*, 118, 120-136.
- Zenetos, A., 2017. Progress in Mediterranean bioinvasions two years after the Suez Canal enlargement. *Acta Adriatica*, 58 (2), 347-358.
- Zenetos, A., Meriç, E., Verlaque, M., Galli, P., Boudouresque, C.F. *et al.*, 2008. Additions to the annotated list of marine alien biota in the Mediterranean with special emphasis on Foraminifera and Parasites. *Mediterranean Marine Science*, 9 (1), 119-166.
- Zenetos, A., Gofas, S., Verlaque, M., Çinar, M.E., Garcia Raso, J.E. *et al.*, 2010. Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science*, 11 (2), 381.
- Zenetos, A., Gofas, S., Morri, C., Rosso, A., Violanti, D. *et al.*, 2012. Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. *Mediterranean Marine Science*, 13 (2), 328-352.
- Zenetos, A., Galanidi, M. 2020. Mediterranean non indigenous species at the start of the 2020s: recent changes. *Marine Biodiversity Records* 13 (10).
- Zmiri, A., Kahan, D., Hochstein, S., Reiss, Z., 1974. Phototaxis and thermotaxis in some species of *Amphistegina* (Foraminifera). *The Journal of Protozoology*, 21 (1), 133-138.
- Zweig-Strykowski, M., Reiss, Z., 1975. Bolivinitidae from the Gulf of Elat. *Israel Journal of Earth-Sciences*, 24, 97-111.