

ALL MAPS OF TYPE 2^∞ ARE BOUNDARY MAPS

VÍCTOR JIMÉNEZ LÓPEZ AND L'UBOMÍR SNOHA

(Communicated by Mary Rees)

ABSTRACT. Let f be a continuous map of an interval into itself having periodic points of period 2^n for all $n \geq 0$ and no other periods. It is shown that every neighborhood of f contains a map g such that the set of periods of the periodic points of g is finite. This answers a question posed by L. S. Block and W. A. Coppel.

1. INTRODUCTION

Let I be a real compact interval and $C(I, I)$ be the metric space of continuous maps of I into itself with the distance between two elements f, g defined by $\varrho(f, g) = \max\{|f(x) - g(x)| : x \in I\}$. Let \mathbb{N} be the set of positive integers. A point $p \in I$ is a *periodic point* of a map f if $f^n(p) = p$ for some $n \in \mathbb{N}$. The *period* of p is the least such integer n , and the *orbit* of p under f is the set $\{f^k(p) : k = 0, 1, \dots, n-1\}$. We refer to such an orbit as a *periodic orbit* of f of period n .

A map $f \in C(I, I)$ is *piecewise monotone* if there are points $\min I = a_0 < a_1 < \dots < a_n = \max I$ such that for every $k \in \{1, 2, \dots, n\}$, the restriction of f to the interval $[a_{k-1}, a_k]$ is (not necessarily strictly) monotone. When speaking of a piecewise monotone map we can always take n as the minimal positive integer with this property and call the points a_1, \dots, a_{n-1} *turning points* of f (though they still are not uniquely determined by f).

Consider the Sharkovskii ordering of the set $\mathbb{N} \cup \{2^\infty\}$:

$$3 \succ 5 \succ 7 \succ \dots \succ 2 \cdot 3 \succ 2 \cdot 5 \succ 2 \cdot 7 \succ \dots \succ 4 \cdot 3 \succ 4 \cdot 5 \succ 4 \cdot 7 \succ \dots \succ \dots \\ \succ 2^n \cdot 3 \succ 2^n \cdot 5 \succ 2^n \cdot 7 \succ \dots \succ \dots \succ 2^\infty \succ \dots \succ 2^n \succ \dots \succ 4 \succ 2 \succ 1.$$

We will also use the symbol \succeq in the natural way. For $t \in \mathbb{N} \cup \{2^\infty\}$ we denote by $S(t)$ the set $\{k \in \mathbb{N} : t \succeq k\}$ ($S(2^\infty)$ stands for the set $\{1, 2, 4, \dots, 2^k, \dots\}$). Let $f \in C(I, I)$ and $\text{Per}(f)$ be the set of periods of its periodic points.

Sharkovskii's Theorem ([Sh1],[Sh2]). *For every $f \in C(I, I)$ there exists a $t \in \mathbb{N} \cup \{2^\infty\}$ with $\text{Per}(f) = S(t)$. On the other hand, for every $t \in \mathbb{N} \cup \{2^\infty\}$ there exists an $f \in C(I, I)$ with $\text{Per}(f) = S(t)$.*

Received by the editors February 27, 1995 and, in revised form, May 2, 1995.

1991 *Mathematics Subject Classification*. Primary 26A18; Secondary 58F08, 54H20.

Key words and phrases. Map of type 2^∞ , periodic point, solenoid.

Most of the work on this paper was done during the stay of the first author at the Matej Bel University. The invitation and the support of this institution is gratefully acknowledged. The first author has been partially supported by the DGICYT PB91-0575 and the second author by the Slovak grant agency, grant number 1/1470/94.

If $\text{Per}(f) = S(t)$, then f is said to be of *type* t . When speaking of types we consider them to be ordered by the Sharkovskii ordering. So if a map f is of type 2^∞ or greater than 2^∞ or less than 2^∞ , then, respectively, $\text{Per}(f) = \{1, 2, \dots, 2^k, \dots\}$ or f has a periodic point with period not a power of 2 or $\text{Per}(f) = \{1, 2, \dots, 2^n\}$ for some \mathbb{N} . The set $\text{Per}(f)$ is finite if and only if f is of type less than 2^∞ . The topological entropy of f is positive if and only if f is of type greater than 2^∞ (see [BF] for the “if” part and [Mi] for the “only if” part or [ALM, Theorem 4.4.19]).

Recall that it is very easy to see that any neighborhood of any map f contains maps of types greater than 2^∞ (even maps of type 3) (see [Kl]). Contrary to the maps of types greater than 2^∞ , the maps of types less than 2^∞ do not form a dense set in $C(I, I)$. In fact, this set is nowhere dense in $C(I, I)$. To see this use the following Block’s Theorem.

Block’s Theorem ([Bl]). *Let $f \in C(I, I)$ and let $n \in \text{Per}(f)$. Then there exists a neighborhood $U(f, n)$ of f such that for all $g \in U(f, n)$ we have $\text{Per}(g) \supset S(n) \setminus \{n\}$.*

So, if f is of type greater than 2^∞ , then there is a neighborhood of f containing no map of type less than (or equal to) 2^∞ . L. S. Block and W. A. Coppel (see [BC], the end of chapter II.4) posed the question of whether any neighborhood of any map of type 2^∞ contains a map of type less than 2^∞ . We answer this question in the affirmative by proving the following

Theorem. *Let $f \in C(I, I)$ be of type 2^∞ . Then every neighborhood of the map f contains a piecewise monotone map of type less than 2^∞ .*

We prove this theorem in two steps. First we prove it under the additional assumption that f is piecewise monotone. Then we prove that in any neighborhood of a map of type 2^∞ there is a piecewise monotone map of type at most 2^∞ . But before going to the proof we wish to mention some aspects of this theorem.

Denote by G or E or L , respectively, the set of all maps of types greater than or equal to or less than 2^∞ . For any set A in the metric space $C(I, I)$ let $\text{Bd } A$ denote the boundary of A . Our theorem and Block’s Theorem show that $\text{Bd } G = \text{Bd } L = E \cup L$. This is what is meant by the title of this paper.

In the Sharkovskii ordering the smallest element is 1 and the largest one is 3. For any $n \in \mathbb{N} \setminus \{1\}$ denote by $\nu(n)$ the predecessor of n , i.e., the maximum (in the Sharkovskii ordering) of the set $S(n) \setminus \{n\}$. If f is of type $n \in \mathbb{N} \setminus \{1\}$ then, by Block’s Theorem, there is a neighborhood U of f such that for every $g \in U$, the type of g is at least $\nu(n)$. It is also known (and not difficult to show) that for any $n \in \mathbb{N}$ there exist a map f_n of type n and a neighborhood U of f_n such that for every $g \in U$, the type of g is at least n . Our theorem shows that this is not true for $n = 2^\infty$. In other words, if f is of type 2^∞ , and if for every $n \in \mathbb{N}$ the neighborhood U_n of f satisfies that every map in U_n is at least of type 2^n (the existence of such neighborhoods follows from Block’s Theorem), then $\bigcap_{n=0}^\infty U_n = \{f\}$.

The Sharkovskii Theorem holds also for the set $C(\mathbb{R}, \mathbb{R})$ of continuous maps from the real line \mathbb{R} into itself (in this case we have the additional possibility $\text{Per}(f) = \emptyset$) (see, e.g., [ALM, Corollary 2.1.2]). It is known that Block’s Theorem works also for maps from $C(\mathbb{R}, \mathbb{R})$ (now $\varrho(f, g) = \sup\{|f(x) - g(x)| : x \in \mathbb{R}\}$ may be infinite) (see [ALM], Remark 2.8.5). So, it is natural to ask whether at least the weaker form of the theorem without the words “piecewise monotone” is true in $C(\mathbb{R}, \mathbb{R})$. The answer is negative. In fact, for any $n \in \mathbb{N}$ take a map $\varphi_n \in C([0, 1], [0, 1])$ of type 2^n with $\varphi_n([0, 1]) \subset [1/4, 3/4]$ and $0 < \varepsilon_n < 1/4$ such that the ball $B(\varphi_n, \varepsilon_n)$

contains only maps with types at least 2^{n-1} . Then there are $a_n > 0$ and a map $\alpha_n \in C([0, a_n], [0, a_n])$ of type 2^n with $\alpha_n([0, a_n]) \subset [1, a_n - 1]$ such that the ball $B(\alpha_n, 1)$ contains only maps with types at least 2^{n-1} (take $a_n = 1/\varepsilon_n$ and $\alpha_n = h_n \circ \varphi_n \circ h_n^{-1}$ where h_n is the increasing affine map of $[0, 1]$ onto $[0, a_n]$). Further, put $b_0 = 0$, $b_1 = 1$ and $b_{2n} = n + \sum_{i=1}^n a_i$, $b_{2n+1} = b_{2n} + 1$ for $n = 1, 2, \dots$ and let $\beta_n \in C([b_{2n-1}, b_{2n}], [b_{2n-1}, b_{2n}])$ be defined by $\beta_n = k_n \circ \alpha_n \circ k_n^{-1}$ where k_n is the increasing affine map from $[0, a_n]$ onto $[b_{2n-1}, b_{2n}]$. Then $f \in C(\mathbb{R}, \mathbb{R})$ defined by

$$f(x) = \begin{cases} x, & \text{if } x < 0, \\ \beta_n(x), & \text{if } x \in [b_{2n-1}, b_{2n}], \\ \text{affine,} & \text{if } x \in [b_{2n-2}, b_{2n-1}] \end{cases}$$

is a map from $C(\mathbb{R}, \mathbb{R})$ of type 2^∞ such that all maps from the ball $B(f, 1)$ have types at least 2^∞ (use that $g([b_{2n-1}, b_{2n}]) \subset [b_{2n-1}, b_{2n}]$ whenever $g \in B(f, 1)$).

2. PROOF OF THE THEOREM

If $A \subset I$, then $\text{int } A$ or $\text{diam } A$ will denote the interior or the diameter of A , respectively. If $J_1, J_2 \subset I$ are intervals, then $J_1 < J_2$ will mean that $\sup J_1 < \inf J_2$. If f is a map and A is a set, then $f|_A$ is the restriction of f to A . Let $f \in C(I, I)$. We say that $x \in I$ is *eventually periodic* if $f^n(x)$ is periodic for some n .

Given $f \in C(I, I)$, a closed subinterval J of I is periodic of period n if $f^n(J) = J$ and $f^k(J) \cap f^l(J) = \emptyset$ for any $0 \leq k < l < n$. Further, we say that $S \subset I$ is a (*simple*) *solenoid* of f if $S = \bigcap_{n=0}^\infty \bigcup_{k=0}^{2^n-1} f^k(I^n)$ where for any n , I^n is a periodic interval of period 2^n such that $I^n \supset I^{n+1}$. The equality $S = \bigcap_{n=0}^\infty \bigcup_{k=0}^{2^n-1} f^k(I^n)$ will be said to be a *standard representation* of the solenoid S . Clearly, S is a compact set, $f(S) = S$ and S cannot contain any eventually periodic point of f .

Lemma 1. *Let $f \in C(I, I)$ and let R, S be different solenoids of f . Then $R \cap S = \emptyset$.*

Proof. This follows immediately from the proofs of Proposition 2.2 (7) and Lemma 2.15 in [Pr] (these proofs work without the assumption of piecewise monotonicity). □

Lemma 2. *Let $f \in C(I, I)$ be piecewise monotone and S be a solenoid of f . Then S must contain a turning point. In particular the number of solenoids of f is finite.*

Proof. If $S = \bigcap_{n=0}^\infty \bigcup_{k=0}^{2^n-1} f^k(I^n)$ is a standard representation of a solenoid S of f , then, taking any n , f cannot be monotone on all intervals $f^k(I^n)$, $k = 0, 1, \dots, 2^n - 1$. Consequently, there is at least one of the finitely many turning points of f belonging to S . The rest of the lemma follows from Lemma 1. □

The set of all limit points of the trajectory $(f^n(x))_{n=0}^\infty$ of a point x is called the *ω -limit set* of x under f and is denoted by $\omega_f(x)$. A standard well-known result says that every finite ω -limit set is a periodic orbit.

A well-known result implicit in several of Sharkovskii's papers and proved in [Sm] (see also [FS]) states that every infinite ω -limit set of a map of type 2^∞ is contained in a solenoid. In [Ge] it is proved that if a map f of type 2^∞ is additionally piecewise monotone, then any infinite ω -limit set of f is Cantor-like. Finally, we will substantially use a result from [JS] saying that every piecewise monotone map of type 2^∞ must have an infinite (Cantor-like) ω -limit set. Combining these facts we get

Lemma 3. *Every piecewise monotone map $f \in C(I, I)$ of type 2^∞ has infinite ω -limit sets, each of them being Cantor-like and contained in a solenoid.*

Now we are able to prove our main result under the additional assumption that f is piecewise monotone.

Lemma 4. *Let $f \in C(I, I)$ be of type 2^∞ and piecewise monotone. Then every neighborhood of f contains a piecewise monotone map of type less than 2^∞ .*

Proof. Let $\varepsilon > 0$. We wish to find a map $g \in C(I, I)$ of type less than 2^∞ with $\varrho(f, g) < \varepsilon$. Take $\delta > 0$ such that $\text{diam } f(J) < \varepsilon$ whenever $\text{diam } J < \delta$.

Special case. Assume that the map f has only one solenoid S (see Lemma 3 and Lemma 2).

In the standard representation $S = \bigcap_{n=0}^{\infty} \bigcup_{k=0}^{2^n-1} I_k^n$, $I_k^n = f^k(I^n)$, we may assume that $I_0^n \supset I_0^{n+1}$ for every n . Then $I_k^{n+1}, I_{k+2^n}^{n+1} \subset I_k^n$ for every n and $k = 0, 1, \dots, 2^n - 1$. Denote by K_k^n the (maximal) closed interval lying between $\text{int } I_k^{n+1}$ and $\text{int } I_{k+2^n}^{n+1}$. Further, for every n denote $I(n) = \{I_k^n : k = 0, 1, \dots, 2^n - 1\}$, $K(n) = \{K_k^n : k = 0, 1, \dots, 2^n - 1\}$ and $\bigcup I(n) = \bigcup_{k=0}^{2^n-1} I_k^n$, $\bigcup K(n) = \bigcup_{k=0}^{2^n-1} K_k^n$.

Realize the following:

(i) There is a (sufficiently large) l_1 such that $\bigcup I(l_1)$ contains only those turning points of f which belong to S . Then, for any $n \geq l_1$, $\text{int}(\bigcup K(n))$ does not contain any turning point of f .

(ii) There is an l_2 such that $I(l_2)$ contains an interval with diameter less than δ .

Take $l = \max\{l_1, l_2\}$. Then $I(l)$ contains an interval, say I_0^l , with diameter less than δ . Further, f is monotone on each of the intervals belonging to $K(l)$.

We have $P = I_0^{l+1} \cup K_0^l \cup I_{2^l}^{l+1} \subset I_0^l$ where we may assume that $I_0^{l+1} < I_{2^l}^{l+1}$. Define $g \in C(I, I)$ by

$$g(x) = \begin{cases} f(x), & \text{for } x \in I \setminus (K_0^l \cup I_{2^l}^{l+1}), \\ f(\max I_{2^l}^{l+1}), & \text{for } x \in I_{2^l}^{l+1}, \\ \text{affine on } K_0^l. \end{cases}$$

The map g is piecewise monotone and since $\text{diam } f(P) < \varepsilon$, $g(P) \subset f(P)$ and f coincides with g on $I \setminus P$, we have $\varrho(f, g) < \varepsilon$. To finish the proof of the Special case, it is sufficient to show that g is of type less than 2^∞ .

Take any $x \in I$. If the trajectory $(g^n(x))_{n=0}^{\infty}$ does not visit $K_0^l \cup I_{2^l}^{l+1}$, then it coincides with the trajectory $(f^n(x))_{n=0}^{\infty}$, whence $\omega_g(x) = \omega_f(x)$. Since the set $\omega_g(x) = \omega_f(x)$ is not a subset of S (otherwise the trajectory $(g^n(x))_{n=0}^{\infty}$ would intersect $I_{2^l}^{l+1}$), it is a finite set, a periodic orbit of f (as well as of g) of period a power of 2.

Now suppose that $(g^n(x))_{n=0}^{\infty}$ visits the set $I_{2^l}^{l+1}$. Then the point x is eventually periodic with period 2^{l+1} and so $\omega_g(x)$ is finite.

Finally suppose that $(g^n(x))_{n=0}^{\infty}$ does not visit $I_{2^l}^{l+1}$ and, for some n_0 , $g^{n_0}(x) \in K_0^l$. Then $(g^n(x))_{n=n_0}^{\infty}$ lies in $\bigcup K(l)$. Due to the fact that g is monotone on each of the intervals belonging to $K(l)$, for any $k \in \{0, 1, \dots, 2^l - 1\}$ the set $M_k^l = \{y \in K_k^l : (g^n(y))_{n=0}^{\infty} \text{ lies in } \bigcup K(l)\}$ is an interval (possibly degenerate), the map $h = g^{2^l}$ maps this interval into itself and is monotone on it. Hence, for any $y \in M_k^l$ the cardinality of $\omega_h(y)$ is at most 2 (see [Co] or [Sh2]). Therefore, for our point x we get that $\omega_g(x)$ has cardinality 2^l or 2^{l+1} .

We have proved that for every $x \in I$, $\omega_g(x)$ is finite and, moreover, has cardinality a power of 2. So g is of type at most 2^∞ . It cannot be of type 2^∞ since otherwise, being piecewise monotone, by [JS] (cf. Lemma 3) it would have an infinite ω -limit set. Therefore g is of type less than 2^∞ .

General case. The map f has the solenoids S_1, S_2, \dots, S_r (and no other ones) for some $r \in \mathbb{N}$.

Since the solenoids S_i , $i = 1, 2, \dots, r$, are compact and mutually disjoint, each of them has a positive distance from the others. So, if $S_i = \bigcap_{n=0}^\infty \bigcup_{k=0}^{2^n-1} (I_i)_k^n$ is a standard representation of S_i , there exists an m such that the sets $\bigcup_{k=0}^{2^m-1} (I_i)_k^m$, $i = 1, 2, \dots, r$, are pairwise disjoint. Now the same procedure which was applied in the Special case to a map having one maximal solenoid S can be applied r -times to our map f having r solenoids S_1, S_2, \dots, S_r . As a result we get a piecewise monotone map g of type less than 2^∞ with $\varrho(f, g) < \varepsilon$. \square

In what follows, $h(f)$ denotes the topological entropy of f (see [AKM] or [ALM] for the definition and properties). Here we just recall that $h(f) \in [0, +\infty]$.

Lemma 5. *Let X be a compact metric space, and let $T : X \rightarrow X$ be a continuous map. Suppose that $(U_n)_{n \geq 1}$ is a sequence of open subsets of X , and let $S : X \rightarrow X$ be a continuous map such that $S(x) = T(x)$ for all $x \in X \setminus \bigcup_{n=1}^\infty U_n$ and $S|_{U_n}$ is constant for all $n \geq 1$. Then $h(S) \leq h(T)$.*

Proof. Set $V := \bigcup_{k=0}^\infty \bigcup_{n=1}^\infty S^{-k}(U_n)$ and $C := X \setminus V$. Obviously $S(C) \subset C$ and $S|_C = T|_C$. An easy calculation shows that $\Omega(S) \cap V$ equals an at most countable union of periodic orbits ($\Omega(S) \cap V$ may be empty), where $\Omega(S)$ denotes the nonwandering set of S (see [W] for the definition and properties). The variational principle (see [W, Theorem 8.6 and Corollary 8.6.1] and [W, Theorem 6.15]) give

$$h(S) = \begin{cases} h(S|_C), & \text{if } C \neq \emptyset, \\ 0, & \text{otherwise.} \end{cases}$$

In the first case $h(S|_C) = h(T|_C) \leq h(T)$, and in the second case $h(S) \leq h(T)$ is trivial. \square

Remark. An easy consequence of Lemma 5 is the following fact. If $f, g \in C(I, I)$ coincide outside an open set G , and if g is constant on every connected component of G , then $h(g) \leq h(f)$.

Lemma 6. *Let $f \in C(I, I)$ and let $[u, v] \subset I$ be a closed interval with $\max f([u, v]) \in \{f(u), f(v)\}$. Then there is a map $g \in C(I, I)$ with the following properties:*

- (1) $g(x) = f(x)$ whenever $x \notin (u, v)$,
- (2) $g|_{[u, v]}$ is (not necessarily strictly) monotone, and
- (3) $h(g) \leq h(f)$.

Proof. Assume that $\max f([u, v]) = f(v)$ (if the maximum is attained at the point u , the proof is analogous). Define the map $g|_{[u, v]}$ as the so-called rising sun function corresponding to the map $f|_{[u, v]}$ when a rising sun is on the x -axis at $-\infty$ (equivalently, pour water into the graph of $f|_{[u, v]}$ until you get all “holes in the ground” full of water). More precisely, put

$$g(x) = \begin{cases} f(x), & \text{if } x \in I \setminus [u, v], \\ \max\{f(t) : u \leq t \leq x\}, & \text{if } x \in [u, v]. \end{cases}$$

Then (1) and (2) hold trivially, the continuity of g follows from the assumption that $\max f([u, v]) = f(v)$. Finally, the set $G = \{x \in [u, v] : f(x) < g(x)\}$ is open, g equals f on $I \setminus G$ and g is constant on every connected component of G . So, Lemma 5 gives (3). \square

Lemma 7. *Let $f \in C(I, I)$. Then in every neighborhood of f there is a piecewise monotone map g with $h(g) \leq h(f)$.*

Proof. Let $\varepsilon > 0$. Take $\delta > 0$ such that $\text{diam } f(J) < \varepsilon$ whenever $\text{diam } J < \delta$. Now take points $\min I = z_1 < z_2 < \dots < z_k = \max I$ with $|z_i - z_{i+1}| < \delta$ for $i = 1, 2, \dots, k-1$. In each interval $[z_i, z_{i+1}]$ take a point s_i such that $f(s_i) = \max f([z_i, z_{i+1}])$. Let

$$\{z_i : i = 1, 2, \dots, k\} \cup \{s_i : i = 1, 2, \dots, k-1\} = \{x_1, x_2, \dots, x_n\}$$

with $\min I = x_1 < x_2 < \dots < x_n = \max I$. Since each of the intervals $[x_i, x_{i+1}]$, $i = 1, 2, \dots, n-1$, can be viewed as the interval $[u, v]$ in Lemma 6, we can use the lemma $n-1$ times to get a piecewise monotone map g with $h(g) \leq h(f)$ and $\varrho(f, g) < \varepsilon$. \square

Proof of Theorem. Since for $\varphi \in C(I, I)$ we have $h(\varphi) = 0$ if and only if φ is of type at most 2^∞ , it suffices to use Lemma 7 and Lemma 4. \square

REFERENCES

- [AKM] R. L. Adler, A. G. Konheim and M. H. McAndrew, *Topological entropy*, Trans. Amer. Math. Soc. **114** (1965), 309–319. MR **30**:5291
- [ALM] L. Alsedà, J. Llibre and M. Misiurewicz, *Combinatorial Dynamics and Entropy in Dimension One*, World Scientific Publ., Singapore, 1993. MR **95j**:58042
- [Bl] L. Block, *Stability of periodic orbits in the theorem of Sarkovskii*, Proc. Amer. Math. Soc. **81** (1981), 333–336. MR **82b**:58071
- [BC] L. S. Block and W. A. Coppel, *Dynamics in One Dimension*, Lecture Notes in Math., vol. 1513, Springer, Berlin, 1992. MR **93g**:58091
- [BF] R. Bowen and J. Franks, *The periodic points of maps of the disk and the interval*, Topology **15** (1976), 337–342. MR **55**:4283
- [Co] W. A. Coppel, *The solutions of equations by iteration*, Proc. Camb. Phil. Soc. **51** (1955), 41–43. MR **16**:577a
- [FS] V. V. Fedorenko and J. Smital, *Maps of the interval Ljapunov stable on the set of nonwandering points*, Acta Math. Univ. Comenian. (N. S.) **60** (1991), 11–14. MR **92g**:58063
- [Ge] T. Gedeon, *Stable and non-stable non-chaotic maps of the interval*, Math. Slovaca **41** (1991), 379–391. MR **92k**:58074
- [JS] V. Jiménez López and L'. Snoha, *There are no piecewise linear maps of type 2^∞* , Trans. Amer. Math. Soc. (to appear). CMP 96:12
- [Kl] P. E. Kloeden, *Chaotic difference equations are dense*, Bull. Austral. Math. Soc. **15** (1976), 371–379. MR **55**:5809
- [Mi] M. Misiurewicz, *Horseshoes for mappings of an interval*, Bull. Acad. Pol. Sci., Sér. Sci. Math. **27** (1979), 167–169. MR **81b**:58033
- [Pr] C. Preston, *Iterates of Piecewise Monotone Mappings on an Interval*, Lecture Notes in Mathematics, vol. 1347, Springer, Berlin, 1988. MR **89m**:58109
- [Sh1] A. N. Sharkovskii, *Coexistence of cycles of a continuous mapping of the line into itself*, Ukrain. Math. Zh. **16** (1964), 61–71 (Russian). MR **96j**:58058
- [Sh2] A. N. Sharkovskii, *On cycles and the structure of a continuous mapping*, Ukrain. Math. Zh. **17** (1965), 104–111 (Russian).

- [Sm] J. Smítal, *Chaotic functions with zero topological entropy*, Trans. Amer. Mat. Soc. **297** (1986), 269–282. MR **87m**:58107
- [W] P. Walters, *An Introduction to Ergodic Theory*, Springer, New York, 1982. MR **84e**:28017

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE MURCIA, CAMPUS DE ESPINARDO, APTDO.
DE CORREOS 4021, 30100 MURCIA, SPAIN
E-mail address: `vjimenez@fcu.um.es`

DEPARTMENT OF MATHEMATICS, FACULTY OF NATURAL SCIENCES, MATEJ BEL UNIVERSITY,
TAJOVSKÉHO 40, 974 01 BANSKÁ BYSTRICA, SLOVAKIA
E-mail address: `snoha@bb.sanet.sk`