

Allelopathic effect of invasive Canadian goldenrod (*Solidago canadensis* L.) on early growth of red clover (*Trifolium pratense* L.)

Peiman ZANDI^{1,2}, Beata BARABASZ-KRASNY³, Alina STACHURSKA-SWAKOŃ⁴, Joanna PUŁA⁵, Katarzyna MOŹDŹEN^{3*}

¹Yibin University, International Faculty of Applied Technology, Yibin, Sichuan, 644600, China; z_rice_b@yahoo.com

²Institute of Environment and Sustainable Development in Agriculture, Chinese Academy of Agricultural Science, Beijing, China;

³Pedagogical University of Krakow, Institute of Biology, Podchorążych 2 St., 30-084 Kraków, Poland; beata.barabasz-krasny@up.krakow.pl; katarzyna.mozdzen@up.krakow.pl (*corresponding author)

⁴Jagiellonian University, Institute of Botany, Gronostajowa St. 3, 30-387 Kraków, Poland; alina.stachurska-swakon@uj.edu.pl

⁵University of Agriculture, Faculty of Agriculture and Economics, Al. Mickiewicza 21 Ave, 31-120 Kraków, Poland; joanna.pula@urk.edu.pl

Abstract

Solidago canadensis belongs to highly invasive species in Europe, it is established in ruderal, semi-natural and natural communities. To test the traits related to invasiveness, the experiment with the potential of allelopathic compounds produced by the species was conducted. The aqueous extract from the goldenrod leaves was used to examine the germination and early stages of development of *Trifolium pratense*, the species as an example of fodder plant common in meadows and pastures. Three types of aqueous extract were used: decoction, infusion, and macerate. All extracts of Canadian goldenrod had a negative influence on the germination process of red clover, however, the effect changed in time and type of extract. The most inhibiting influence was documented for the macerate type of aqueous extract. Similarly, the negative influence of macerate was the highest for the length and weight of the seedlings, the content of chlorophyll *a* and *b*, and electrolyte leakage. As the procedure of obtaining the macerate is the most like the process of natural extraction of allelopathic substances in nature, there is the practical conclusion to remove the moved biomass of the goldenrod during the restoration process of areas colonised by the species.

Keywords: allelopathy; chlorophyll; electrolyte leakage; germination; invasive species; leaves exudates

Introduction

Invasions of alien species is a worldwide serious ecological problem leading to changes in biodiversity and ecosystem functioning (Pejchar and Mooney, 2002; Pyšek and Richardson, 2010; Pyšek *et al.*, 2010). Invasions also influence a local economy (Pimentel, 2011) and they are an important task for nature conservation (Sołtys-Lelek and Barabasz-Krasny, 2010). The invasive success of non-native plants depends on their biological traits and the vulnerability of the ecosystem (Rejmánek *et al.*, 2005; Bartoszek and Stachurska-Swakoń, 2016). Natural disturbances and anthropogenic changes in environments or in management alter the possibility of invasiveness (Hobbs, 2000; Trzcińska-Tacik and Stachurska-Swakoń, 2011). Allelopathic

compounds produced by plants are considered as an additional element facilitating competition for invasive species during settlements (Abhilasha *et al.*, 2008; Yuan *et al.*, 2013).

Solidago canadensis L. s.l. (Canadian goldenrod) is one of the successful invaders of North America origin that colonised large areas across Europe, Asia, Australia, New Zealand (Semple and Cook, 2006; Lu *et al.*, 2007; Abhilasha *et al.*, 2008; Yuan *et al.*, 2013). The species occurs in ruderal, semi-natural and natural communities, especially in moist habitats (Guzikowa and Maycock, 1986; Zając and Zając, 2015; Towpasz and Stachurska-Swakoń, 2018). It forms single-species aggregation or species-poor communities that lead to landscape homogenisation (Chen *et al.*, 2005; Yuan *et al.*, 2013), reducing diversity of native plant species (Szymura and Wolski, 2006; Szymura *et al.*, 2018) and animals, mainly insects and birds (Moron *et al.*, 2009; Skórka *et al.*, 2010; Masło and Najberek, 2014).

Comprehension of the invasion mechanism of *S. canadensis* is an important issue in the management and recultivation of the areas colonised with the goldenrod. The species propagates vegetatively by rhizomes forming large clonal colonies and produces a huge number of small seeds easily dispersed by wind (Voser-Huber, 1983; Rosef *et al.*, 2019). The rhizomes of the species produce substances that suppress soil pathogens (Zhang *et al.*, 2009). There are few studies testing the allelopathic potential of the Canadian goldenrod (Butcko and Jensen, 2002; Sun *et al.*, 2006; Abhilasha *et al.*, 2008; Yuan *et al.*, 2013; Domaradzki *et al.*, 2017). The aim of the current experiment was to check if the allelopathic compounds from decaying biomass of goldenrod could inhibit the native species. For this experiment, meadow red clover (*Trifolium pratense* L. cv. 'Rozeta') from the Fabaceae family was selected as a fodder plant species of economic importance, due to its feed qualities. The aim of the study was to examine the effect of three types of *S. canadensis* leaf extracts in the form of decoction, infusion and macerate on seed germination and early growth of *T. pratense*. The following parameters were determined: germination indexes (1), the seedlings length (2), the fresh and dry weight and water content (3) the chlorophyll content (4) and the degree of cell membrane destabilisation by electrolyte leakage method (5).

Materials and Methods

Plant material

Trifolium pratense L. cv. 'Rozeta' (fodder, diploid cultivar, characterised by quite tall plants, prone to lying on ground) seeds were obtained from the Sadowniczy store (Poland). The *Solidago canadensis* leaves were collected in the south-eastern part of Poland - Suchoraba 49°58'37"N 20°11'49"E.

Extracts preparation

The aqueous extracts of goldenrod leave in the form of infusion (crushed dry leaves flooded with hot water), decoction (crushed dry leaves boiled in water) and macerate (crushed dry leaves flooded with water and left to stand) were prepared according to the method used by Czerwińska *et al.* (2015). The infusion was prepared from 5 g of dry *S. canadensis* leaves, which were poured over with 250 ml of boiling distilled water and left covered for 30 min. After cooling, the extract was filtered through filter papers. The decoction was prepared by weighing out 8.75 g of dry plant material, which was poured with a 1 litre of distilled water. The solution was mixed thoroughly and left for 24 h in the dark, at room temperature (20-25 °C). After one day, the aqueous extract was boiled for 15 min and filtered in the same way as the infusion. Extract in the form of a macerate was prepared by flooding 5 g of dry leaves of 100 ml cold distilled water and left in the dark, at room temperature for 24 h. After one day, the extract was filtered the same form like as infusion and decoction. The extracts were stored during the experiment at 8 °C.

Germination conditions

The Petri dishes experiment was performed as follows: clover seeds were rinsed in a 1% acetone solution, followed by distilled water and put into sterile dishes (Ø 9 cm) with three layers of filter paper (50 seeds per dish). Each Petri dish was soaked in 6 ml of the appropriate extract and watered with 3 ml of extract every other day. The control consisted of Petri dishes with seeds watered with distilled water in the same quantities as dishes with extracts. Seeds were stored in a growth chamber (Angelantoni Industrie, Italy) at 25 °C/20 °C temperature (day/night), with 200 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ light intensity and relative humidity RH% 60-70%, during 8 days. The experiment was carried out in 3 repetitions in 2 independent series.

Germination indexes

Germination rate (GR), germination speed (GS), allelopathic effect response index (RI) (Gao *et al.*, 2009) and seedling vigour index (SVI) (Islam *et al.*, 2009).

$$\text{GR} = (\text{number of germinated seeds} / \text{total number of seeds}) \times 100$$

$$\text{GS} = ((\text{GT} \times \text{D}) / (\text{GC} \times \text{D})) \times 100$$

where: GT is the number of germinated seeds daily in the treatment, GC is the number of germinated seeds daily in the control, and D is the number of corresponding days

$$\text{RI} = 1 - \text{C} / \text{T} \text{ (when } \text{T} \geq \text{C})$$

where: C is the control germination speed and T is the treatment germination speed

$$\text{SVI} = (\text{seedling length (cm)} \times \text{percentage of germinated seeds}) / 100.$$

Biometric analysis

The length of *T. pratense* (root and hypocotyl) seedlings treated the aqueous extracts from *S. canadensis* was measured using a calliper (Topex 31C615, Poland), to the nearest of 0.1 cm, after the 8 days from start of the experiment. The inhibition of percentage growth (IP), expressed as a percentage of control seedlings index, was determined according to formula used by Mominul Islam and Kato-Noguchi (2012).

Fresh and dry weight and tissue water content

Seedlings were weighted (fresh weight - FW) (Radwag WPS120, Poland) and dried (dry weight - DW) at 105 °C for 48 h in oven (Wamed SUP 100, Poland). Tissue water content (TWC) was studied according to Black and Pritchard (2002) with some modifications.

Chlorophyll content

The content of chlorophyll *a*, *b*, *a + b* was determined according to Barnes *et al.* (1992). The aboveground parts of the seedlings were extracted in dimethyl sulfoxide (SIGMA-Aldrich) at 65 °C for 12 hours. The absorbance of chlorophyll *a* and *b* was determined at the wavelength: $\lambda = 665$ and 648 nm, using a spectrophotometer Aquarius 9500 (Cecil Instruments, Cambridge, United Kingdom).

Electrolyte leakage

The degree of cell membranes destabilisation by the electrolyte leakage was carried out on clover seedlings according to the method described by Redmann *et al.* (1986).

Statistical analysis

The importance of variability between objects by Anova was checked. The analysis of the differences between groups was examined by Duncan test for homogeneous groups, at the level of $p \leq 0.05$; values marked with different letters (a, b, c) differ significantly. The calculations were performed using Statistica for Windows 13.1.

Results

Germination indexes

Compared to the control, in 8 days, the highest germination rate (GR) values were found in seeds grown on the *Solidago canadensis* decoction. The lowest values of germinated *Trifolium pratense* cv. Rozeta seeds was observed in Petri dishes with macerate. Germination speed (GS), expressed as a percentage of control, was significantly highest for seeds watered with decoction, and the lowest for seeds on the macerate. Allelopathic effect response index (RI) showed that decoction has a positive influence on seed germination. In turn, the infusion and macerate inhibited clover seed germination. Compared to the control, the highest inhibitory effect was observed for seeds watered with macerate. The seedling vigour index (SVI) was significantly highest for the control in relation to the extracts used. Regardless of the examined seedlings organ, the infusion and macerate resulted in a significant reduction the SVI (Figure 1, Table 1).




Figure 1. Germination rate (GR) of *Trifolium pratense* L. cv. 'Rozeta' seeds, treated with different types of *Solidago canadensis* L. aqueous extracts from dry leaves
Mean values (\pm SD) marked in different letters (a, b, c) differ significantly according to Duncan's test $p \leq 0.05$

Table 1. Germination speed (GS) expressed as a percentage of control, allelopathic effect response (RI) and seedling vigour index (SVI) of *Trifolium pratense* L. cv. 'Rozeta' seeds treatment different types of *Solidago canadensis* L. aqueous extracts from dry leaves

| Parameters | Control | Decoction | Infusion | Macerate |
|---------------------------|-------------------|--------------------|--------------------|--------------------|
| GS (%) | 100 | 70.29 \pm 0.82 a | 45.06 \pm 0.80 b | 43.41 \pm 0.84 b |
| RI (a.u.) | 0 \pm 0.00 b | 0.40 \pm 0.02 a | -1.08 \pm 0.06 c | -1.50 \pm 0.11 d |
| SVI whole seedling (a.u.) | 2.95 \pm 0.81 a | 2.29 \pm 0.70 a | 0.28 \pm 0.05 b | 0.19 \pm 0.07 b |
| SVI root (a.u.) | 1.25 \pm 0.33 a | 1.15 \pm 0.42 a | 0.14 \pm 0.05 b | 0.08 \pm 0.04 b |
| SVI hypocotyl (a.u.) | 1.70 \pm 0.54 a | 1.14 \pm 0.38 a | 0.14 \pm 0.04 b | 0.11 \pm 0.03 b |

Note: mean values (\pm SD) marked in different letters (a, b, c) in row differ significantly according to Duncan's test $p \leq 0.05$

Biometric analysis

Biometric analysis of *T. pratense* seedlings revealed a statistically inhibitory effect of aqueous extracts from *S. canadensis* leaves (Figure 2). On the macerate, a significant growth inhibition for the above- and underground organs of seedlings was observed. Regardless of the type of extracts, the growth of the aboveground organs was inhibited, compared to the control. Analysis of the length of whole clover seedlings

showed a negative impact of each of the extracts. Compared to the control, no significant differences in length seedling growth were observed for *T. pratense* germinated on the decoction.




Figure 2. Length of seedlings (cm) (A) and inhibition percentage index IP (B), of *Trifolium pratense* L. cv. Rozeta seedlings, which were treated with different types of *Solidago canadensis* L. aqueous extracts from dry leaves
Mean values (\pm SD) marked different letters (a, b, c) differ significantly according to Duncan's test $p \leq 0.05$; (B) - a negative (-) value indicates stimulation of growth, and a positive (+) value indicates growth inhibition

Fresh and dry weight and tissue water content

The fresh weight values of *T. pratense* seedlings revealed a statistically significant negative effect of extracts on biomass increase, regardless of the type of extract used. Contrary to the dry weight, for which the values increased significantly for each extract compared to the control. Water content in seedlings was similar in each treatment, compared to control (Table 2).


Chlorophyll content

In clover seedlings grown on the infusion and macerate, significant differences in the chlorophyll *a* and *b* concentration were found. In relation to the control, the smallest chlorophyll content was found in seedlings watered with macerate. Only in the case of seedlings treated with decoction no statistically significant changes in the content of the analysed pigments were observed (Figure 3).

Table 2. Fresh, dry weight and tissue water content in *Trifolium pratense* L. cv. 'Rozeta' seedlings treatment different types of *Solidago canadensis* L. aqueous extracts from dry leaves

| Type of extracts | Fresh weight (g) | Dry weight (g) | Tissue water content (%) |
|------------------|------------------|------------------|--------------------------|
| Control | 0.0156 ± 0.01 a | 0.00018 ± 0.01 b | 99.81 ± 0.52 a |
| Decoction | 0.0094 ± 0.01 b | 0.00045 ± 0.01 a | 94.69 ± 6.14 a |
| Infusion | 0.0097 ± 0.01 b | 0.00033 ± 0.01 a | 96.36 ± 3.78 a |
| Macerate | 0.0094 ± 0.01 b | 0.00032 ± 0.01 a | 96.90 ± 3.35 a |


Note: mean values (±SD) marked with different letters (a, b, c) in column differ significantly according to Duncan's test $p \leq 0.05$

**Figure 3.** Content of chlorophyll a, b, a + b in *Trifolium pratense* L. cv. 'Rozeta' seedlings treated with different types of *Solidago canadensis* L. aqueous extracts from dry leaves

Mean values (±SD) marked with different letters (a, b, c) differ significantly according to Duncan's test $p \leq 0.05$

Electrolyte leakage

An increase of electrolyte leakage from *T. pratense* seedlings was observed in each of the *S. canadensis* extracts used. Compared to the control, the smallest changes in water and ion management were found in seedlings watered with decoction and infusion, and the largest in seedlings germinated on macerate extracts (Figure 4).

**Figure 4.** Electrolyte leakage in *Trifolium pratense* L. cv. Rozeta seedlings treated with different types of *Solidago canadensis* L. aqueous extracts from dry leaves

Mean values (±SD) marked with different letters (a, b, c) differ significantly according to Duncan's test $p \leq 0.05$

Discussion

Alien invasive species, similarly as local weeds, form communities with an additive system of components in which individuals compete for limited habitat resources. The effect of the competition is weakening the ability of native plants to adapt to dynamic environmental conditions (Richardson and Pyšek, 2012). This is usually manifested by inhibiting seed germination, reducing the amount of biomass produced, reducing their size, disorganising of metabolic processes in plants, etc. With a strong allelopathic interaction, even individuals die from a given type of community (Hierro *et al.*, 2003; Zandi *et al.*, 2018, 2019; Puła *et al.*, 2020).

In this experiment, aqueous extracts of *Solidago canadensis* leaves showed a pronounced inhibitory effect on the germination and growth of *Trifolium pratense* cv. 'Rozeta', in the early phase stages of development (Figure 1-2, Table 1). This confirms the known thesis that alien invasive plants release substances that cause allelopathic effects, influencing seed germination and the growth of native species (Tokarska-Guzik *et al.*, 2012; Yuan *et al.*, 2013). For example, of the 11 native species found naturally occurring in southern China, most showed sensitivity to alcoholic extracts from *S. canadensis* roots and rhizomes during germination and growth (Chen *et al.*, 2005; Yang *et al.*, 2007). However, the allelopathic effect was not observed for all species. It has been reported that *T. pratense* and *Medicago lupulina* L. react by stimulating of germination process, and during early stage of growth of red clover the stimulating of root elongation was documented. In this case, it can be concluded that root and rhizome extracts have less inhibitory properties than the various forms of leaf extracts used in the current experiment. The leaves of *S. canadensis*, as previously demonstrated, contain a wide range of compounds that may be responsible for their biological activity, among them are phenols, saponins, tannins and flavonoids (Zhao *et al.*, 2005; Zhang *et al.*, 2011). The leaves and other aboveground parts of various plants are the organs that produce the largest amount of this type of compound. The effect of allelopathic compounds on other plants may be negative in higher concentrations or positive in lower ones, as has been stated many times before (Inderjit and Duke, 2003; Bing-Yao *et al.*, 2006; Inderjit *et al.*, 2006; Sun and He, 2010; Yuan *et al.*, 2013), and the example cited above and the current experiment clearly confirm this.

Allelopathic compounds are released into the environment from plants in the form of volatile substances, by secretion from the root system into the soil, they are also washed out or released from dead parts of plants (Barabasz-Krasny *et al.*, 2017; Szafraniec *et al.*, 2019; Zandi *et al.*, 2019; Możdżeń *et al.*, 2020; Puła *et al.*, 2020). In this experiment, the negative effects of extracts varied depending on the type of extract used. Compared to the control, the highest inhibition of growth in length and weight of seedlings was found in the case of macerate, and the lowest in the presence of decoction (Figure 2, Table 2). Macerate is a kind of solution that most resembles the natural extraction of allelopathic substances from dead plant debris, occurring in nature. As the experiment showed, it contains a higher concentration of allelopathic compounds than the other extracts and significantly inhibits the development of plants. This is a very important conclusion for the management of areas colonised by Canadian goldenrod. Currently, according to recommendations, the most common way to eradicate this invasive species on large abandoned agricultural land is mowing (Kabuce and Priede, 2010). In view of the above research results, a very important here seems to be the removal of mown organic matter, which is a source of allelopathic compounds, inhibiting germination and development of native species returning to former habitats.

One of the first effects of allelopathic compounds at the cell level is membrane depolarization. It causes disorders in the transport of anions and cations, which is associated with increased permeability of these structures. Damage to membranous cell structures depends, among others on the concentration and solubility of allelopathic substances and the pH of the environment (Shann and Blum, 1987; Einhellig, 2004). Low pH values may promote the activity of allelopathic compounds released by *S. canadensis* (Wang *et al.*, 2016). In addition, the environmental stress factors increase allelochemical production and thus increase potential toxicity (An, 2005). The conducted studies showed an increase in the destabilisation of cell membranes in the presence of extracts from *S. canadensis*. Depending on the type of extract and the substances it contains, their

actions resulted in a lower or higher electrolytes leakage, indicating the stress of plants treated with extracts (Figure 4). Disturbances of water-ion economy have a consequent effect on the course of other life processes of plants. For example, they play an important role in the production of chlorophyll, which is responsible for photochemical reactions. In this experiment, clover seedlings were characterised by different content of chlorophylls; the most negative effects were observed with macerate and infusion (Figure 3). Usually, when the concentration of allelopathic compounds increases in extracts, there is a significant reduction in the chlorophyll content (Puła *et al.*, 2020).

The sensitivity and reactive ability on the changes in the environment are an important mechanism of ecological invasion. An explanation in terms of physiological and ecological adaptation of alien invasive plants to the environment would certainly help to introduce effective and synthetic schemes to destruction of them. It would also facilitate forecasting potential distribution areas and estimating threats to other native flora species. To the control of invasive plants such as *S. canadensis*, attention should be paid to their scattered populations, monitored so that they are not a secondary source for seed dispersion (Hua *et al.*, 2007). It is also worth look for among native species of plants that are relatively resistant to allelopathic compounds and can pave the way for re-colonisation of areas colonised by alien invasive plants. In the light of other experiments, legumes are relatively good for this (Chen *et al.*, 2005; Yang *et al.*, 2007).

Conclusions

(1) The results presented in this study confirm allelopathic properties of various types of the aqueous extracts from *Solidago canadensis* dry leaves; chemical compounds contained in goldenrod leaves had a negative effect on *Trifolium pratense* cv. 'Rozeta' germination indexes. (2) Biometric analysis of clover seedlings revealed significant growth inhibition in the presence of infusion and macerate on *T. pratense* seedlings. (3) Each of the extracts used had a negative effect on fresh weight; the highest decrease in the value of these parameters was recorded for seedlings germinated on macerate. Dry weight was increased in all extracts, compared to the control values. (4) The aqueous extracts reduced chlorophyll content in clover seedlings; the most negative effects were caused by macerate and infusion. (5) The negative effect of aqueous extracts from *S. canadensis* leaves was confirmed by electrolyte leakage; water-ion disturbances were weaker in seedlings germinated on the decoction and infusion than in those watered with the macerate. (6) Macerate as a type of solution, is the closest to the natural extraction from dead organic debris, and its allelopathic effect is in this case the largest, so when performing treatments eliminating goldenrod, which is an important source of allelochemical compounds that inhibit germination and growth of other plants.

Authors' Contributions

Conceptualization: PZ, BBK, KM; Formal analysis: KM, BBK, PZ, AS-S; Funding acquisition BB-K; Investigation: BB-K, KM, PZ, JP; Methodology: KM, BB-K, PZ; Project administration: PZ, KM, BB-K, JP; Resources: KM, BB-K, PZ, AS-S; Software: BB-K, PZ, AS-S, JP; Visualization: PZ, KM, BB-K, AS-S; Writing - original draft: KM, BB-K, PZ, AS-S; Writing - review and editing: PZ, KM, BB-K.

All authors read and approved the final manuscript.

Acknowledgements

This study was financially supported by the Pedagogical University of Krakow Grant: Ecophysiological properties of invasive plants in natural and anthropogenic habitats: the importance of the phenomenon of allelopathy - part I. (signature: WPBU/2020/05/00172; registration number: BN.610-169/PBU/2020).

Conflict of Interests

The authors declare that there are no conflicts of interest related to this article.

References

- Abhilasha D, Quintana N, Vivanco J, Joshi J (2008). Do allelopathic compounds in invasive *Solidago canadensis* s.l. restrain the native European flora? *Journal of Ecology* 96(5):993-1001. <https://doi.org/10.1111/j.1365-2745.2008.01413.x>
- An M (2005). Mathematical modelling of dose-response relationship (hormesis) in allelopathy and its application. *Nonlinearity in Biology, Toxicology and Medicine* 3(2):153-172. <https://dx.doi.org/10.2201/2Fnonlin.003.02.001>
- Barnes JD, Balaguer L, Manrique E, Elvira, S, Davison AW (1992). A reappraisal of the use of DMSO for the extraction and determination of chlorophylls a and b in lichens and higher plants. *Environmental and Experimental Botany* 32(2):85-100. [https://doi.org/10.1016/0098-8472\(92\)90034-Y](https://doi.org/10.1016/0098-8472(92)90034-Y)
- Bartoszek W, Stachurska-Swakoń A (2016). *Gypsophila perfoliata* (Caryophyllaceae) in Poland. *Polish Botanical Journal* 61(2):257-262. <http://dx.doi.org/10.1515/pbj-2016-0030>
- Bing-Yao S, Jian-Zhong T, Zhi-Gang T, Fu-Gen G, Ming-De Z (2006). Allelopathic effects of extracts from *Solidago canadensis* L. against seed germination and seedling growth of some plants. *Journal of Environmental Sciences* 18(2):304-309.
- Black M, Pritchard HW (2002). *Desiccation and survival in plants: Drying without dying*. CAB International Wallingford, United Kingdom.
- Butcko VM, Jensen RJ (2002). Evidence of tissue-specific allelopathic activity in *Euthamia graminifolia* and *Solidago canadensis* (Asteraceae). *The American Midland Naturalist* 148(2):253-262. [https://doi.org/10.1674/0003-0031\(2002\)148\[0253:EOTSAA\]2.0.CO;2](https://doi.org/10.1674/0003-0031(2002)148[0253:EOTSAA]2.0.CO;2)
- Chen X, Mei LX, Tang JJ (2005). Allelopathic effects of invasive *Solidago canadensis* on germination and root growth of native Chinese plants. The Regional Institute - online publishing; Retrieved 2020 September 09 from http://www.regional.org.au/au/allelopathy/2005/2/1/2503_chena.html
- Czerwińska E, Szparaga A, Deszcz E (2015). Estimation of effect of dressing in plant extracts on germination capacity of yellow lupine and field pea seed. *Zeszyty Naukowe Uniwersytetu Przyrodniczego we Wrocławiu, Rolnictwo* 113(612):7-19.
- Domaradzki K, Sekutowski TR, Jezierska-Domaradzka A, Matkowski A, Stochmal A (2017). Aktywność herbicydowa wyciągów wodnych z dwóch gatunków z rodzaju *Solidago* w stosunku do *Thlaspi arvense* i *Stellaria media* (Comparison the herbicidal activity of water extracts from two *Solidago* species against *Thlaspi arvense* and *Stellaria media*). *Polish Journal of Agronomy* 31:11-15.
- Einhellig FA (2004). Mode of allelochemical action of phenolic compounds. In: Macias FA, Galindo JCG, Jose M, Molinillo G (Eds). *Allelopathy: Chemistry and mode of action of allelochemicals*. CRC Press, London pp 217-374.
- Gao X, Li M, Gao Z, Li C, Sun Z (2009). Allelopathic effects of *Hemistepta lyrata* on the germination and growth of wheat, sorghum, cucumber, rape, and radish seeds. *Weed Biology and Management* 9(3):243-249. <https://doi.org/10.1111/j.1445-6664.2009.00345.x>
- Guzikowa M, Maycock PF (1986). The invasion and expansion of three North American species of goldenrod *Solidago canadensis* L. *sensu lato*, *S. gigantea* Ait. and *S. graminifolia* (L.) Salisb. in Poland. *Acta Societatis Botanicorum Poloniae* 55(3):367-384. <https://doi.org/10.5586/asbp.1986.034>

- Hierro JL, Callaway RM (2003). Allelopathy and exotic plant invasion. *Plant and Soil* 256(1):29-39. <https://doi.org/10.1023/A:1026208327014>
- Hobbs RJ (2002). Land use changes and invasions. In: Mooney HA, Hobbs RJ (Eds). *Invasive species in a changing world*. Washington, DC: Island Press.
- Hua H, Shuiliang G, Guoqi C (2007). Reproductive biology in an invasive plant *Solidago canadensis*. *Frontiers of Biology in China* 2(2):196-204. <https://doi.org/10.1007/s11515-007-0030-6>
- Inderjit, Callaway RM, Vivanco JM (2006). Can plant biochemistry contribute to understanding of invasion ecology? *Trends Plant Science*.11(2):574-580. <https://doi.org/10.1016/j.tplants.2006.10.004>
- Inderjit, Duke SO (2003). Ecophysiological aspects of allelopathy. *Planta* 217(4):529-539. <https://doi.org/10.2307/23388096>
- Islam AKMA, Anuar N, Yaakob Z (2009). Effect of genotypes and pre-sowing treatments on seed germination behaviour of *Jatropha*. *Asian Journal of Plant Sciences* 8(6):433. <http://dx.doi.org/10.3923/ajps.2009.433.439>
- Kabuce N, Priede N (2010). Invasive alien species fast sweet - *Solidago canadensis*. Online database of the European network on invasive alien species - NOBANIS, www.nobanis.org.
- Lu JZ, Weng ES, Wu XW, Weber E, Zhao B, Li B (2007). Potential distribution of *Solidago canadensis* in China. *Acta Phytotaxonomica Sinica* 45:670-674. <http://dx.doi.org/10.1360/asp06200>
- Masło D, Najberek K (2014) Amerykańskie nawłocie kontra polskie motyle dzienne [American goldenrod versus Polish daytime butterflies]. In: Mirek Z, Nikel A (Eds). *Ochrona przyrody w Polsce wobec współczesnych wyzwań cywilizacyjnych* [Nature protection in Poland in the face of contemporary civilization challenges]. Kraków: Komitet Ochrony Przyrody PAN pp 189-195.
- Mominul Islam AKM, Kato-Noguchi H (2012). Allelopathic potentiality of medicinal plant *Leucas aspera*. *International Journal of Sustainable Agriculture* 4(1):1-7.
- Moroń D, Lenda M, Skórka P, Szentgyörgyi H, Settele J, Woyciechowski M (2009). Wild pollinator communities are negatively affected by invasion of alien goldenrods in grassland landscapes. *Biological Conservation* 142(7):1322-1332. <https://doi.org/10.1016/j.biocon.2008.12.036>
- Możdżeń K, Barabasz-Krasny B, Zandi P, Kliszcz A, Puła J (2020). Effect of aqueous extracts from *Solidago canadensis* L. leaves on germination and early growth stages of three cultivars of *Raphanus sativus* L. var. *radicula* Pers. *Plants* 9:1549. <https://doi.org/10.3390/plants9111549>
- Pejchar L, Mooney H (2002). Invasive species, ecosystem services and human well-being. *Trends in Ecology and Evolution* 24(9):497-504. <http://dx.doi.org/10.1016/j.tree.2009.03.016>
- Pimentel D (2011). *Biological Invasion: economic and environmental costs of alien plant, animal and microbe species*. New York, CRC Press, Taylor and Francis Group, Boca Raton-London.
- Puła J, Zandi P, Stachurska-Swakoń A, Barabasz-Krasny B, Możdżeń K, Wang Y (2020). Influence of alcoholic extracts from *Helianthus annuus* L. roots on the photosynthetic activity of *Sinapis alba* L. cv. *Barka* plants. *Acta Agricultura Scandinavica, Section B, Soil and Plant Science* 70(1):8-13. <https://doi.org/10.1080/09064710.2019.1661509>
- Pyšek P, Jarošík V, Hulme P, Kühn I, Wild J, Arianoutsou M, ... Winter M (2010). Disentangling the role of environmental and human pressures on biological invasions across Europe. *PNAS* 107(27):12157-12162. <https://doi.org/10.1073/pnas.1002314107>
- Pyšek P, Richardson DM (2010). Invasive species, environmental change and management, and health. *Annual Review of Environment and Resources* 100(1):45-52. <https://doi.org/10.1146/annurev-environ-033009-095548>
- Redmann RE, Haraldson J, Gusta LV (1986). Leakage of UV-absorbing substances as a measure of salt injury in leaf tissue of woody species. *Physiologia Plantarum* 67(1):87-91. <https://doi.org/10.1111/j.1399-3054.1986.tb01267.x>
- Rejmánek M, Richardson D, Higgins S, Pitcairn MJ, Grotkopp E (2005). Ecology of invasive plants: State of the art. In: Harold A, Mooney (Eds). *Invasive alien species: Searching for solutions*. DC: Island Press, Washington.
- Richardson DM, Pyšek P (2012). Naturalization of introduced plants: ecological drivers of biogeographical patterns. *New Phytologist* 196(2):383-396. <https://doi.org/10.1111/j.1469-8137.2012.04292.x>
- Rosef L, Ingebrigtsen HH, Heegaard E (2019). Vegetative propagation of *Solidago canadensis*- do fragment size and burial depth matter? *Weed Research* 60(2):132-141. <https://doi.org/10.1111/wre.12395>
- Semple JC, Cook RE (2006). *Flora of North America. Magnoliophyta: Asteridae, Part 7: Asteraceae*. Oxford University Press, New York, and Oxford.
- Shann JR, Blum U (1987). The uptake of ferulic acid and p-hydroxybenzoic acids by *Cucumis sativus*. *Phytochemistry* 26(11):2959-2964. [https://doi.org/10.1016/S0031-9422\(00\)84571-9](https://doi.org/10.1016/S0031-9422(00)84571-9)

- Skórka P, Lenda M, Tryjanowski P (2010). Invasive alien goldenrods negatively affect grassland bird communities in Eastern Europe. *Biological Conservation* 143(4):856-861. <https://doi.org/10.1016/j.biocon.2009.12.030>
- Sołtys-Lelek A, Barabasz-Krasny B (2010). Ekspansja wybranych gatunków obcego pochodzenia we florze i szacie roślinnej Ojcowskiego Parku Narodowego (Południowa Polska) (Expansion of selected species of foreign origin in the flora and vegetation of the Ojców National Park (Southern Poland)). *Prądnik. Prace Muzealne im. Władysława Szafera* 20:333-376.
- Sun BJ, Tan JZ, Wan ZG, Gu FG, Zhu MD (2006). Allelopathic effects of extracts from *Solidago canadensis* L. against seed germination and seedling growth of some plants. *Journal of Environmental Sciences* 18(2):304-309.
- Sun ZK, He WM (2010). Evidence for enhanced mutualism hypothesis: *Solidago canadensis* plants from regular soils perform better. *PLoS ONE* 5:e15418. <https://doi.org/10.1371/journal.pone.0015418>
- Szymura M, Wolski K (2006). Zmiany krajobrazu pod wpływem ekspansywnych bylin północnoamerykańskich z rodzaju *Solidago* L. [Landscape changes under the influence of expansive North American perennials of the genus *Solidago* L.] *Problemy Ekologii Krajobrazu* 16:451-460.
- Szymura TH, Szymura M, Zając M, Zając A (2018). Effect of anthropogenic factors, landscape structure, land relief, soil and climate on risk of alien plant invasion at regional scale. *Science of the Total Environment* 626:1373-1381. <https://doi.org/10.1016/j.scitotenv.2018.01.13>
- Tokarska-Guzik B, Dajdok Z, Zając M, Zając A, Urbisz A, Danielewicz W, Hołdyński C (2012). Rośliny obcego pochodzenia w Polsce ze szczególnym uwzględnieniem gatunków inwazyjnych [Plants of foreign origin in Poland, with particular emphasis on invasive species]. Warszawa, Generalna Dyrekcja Ochrony Środowiska pp 196.
- Towpasz K, Stachurska-Swakoń A (2018). Occurrence of alien species in the agriculture landscape: a case of Proszowice Plateau (Southern Poland). *Annales Universitatis Paedagogicae Cracoviensis Studia Naturae* 3:7-21. <https://doi.org/10.24917/25438832.3.1>
- Trzcińska-Tacik H, Stachurska-Swakoń A (2011). *Alopecurus myosuroides* (Poaceae) as the permanent segetal weed of the Skala vicinity in the Krakow-Częstochowa upland. *Fragmenta Floristica Geobotanica Seria Polonica* 18(2):221-229.
- Voser-Huber ML (1983). Studien an eingebürgerten Arten der Gattung *Solidago* L. (Studies on naturalized species of the genus *Solidago* L.) *Dissertationes Botanicae* 68:1-97.
- Wang C, Xiao H, Zhao L, Liu J, Wang L, Zhang F, ... Du D (2016). The allelopathic effects of invasive plant *Solidago canadensis* on seed germination and growth of *Lactuca sativa* enhanced by different types of acid deposition. *Ecotoxicology* 25(3):555-562. <https://doi.org/10.1007/s10646-016-1614-1>
- Yang RY, Mei LX, Tang JJ, Chen X (2007). Allelopathic effects of invasive *Solidago canadensis* L. on germination and growth of native Chinese plant species. *Allelopathy Journal* 19(1):241-248.
- Yuan YG, Wang B, Zhang SS, Tang JJ, Tu C, Hu SJ, ... Chen X (2013). Enhanced allelopathy and competitive ability of invasive plant *Solidago canadensis* in its introduced range. *Journal of Plant Ecology* 6(3):253-263. <https://doi.org/10.1093/jpe/rts033>
- Zając A, Zając M (2015). Distribution of kenophytes in the polish Carpathians and their foreland. Kraków: Nakładem Instytutu Botaniki Uniwersytetu Jagiellońskiego.
- Zandi P, Barabasz-Krasny B, Stachurska-Swakoń A, Puła J, Możdżeń K (2018). Allelopathic effects of *Stellaria media* (L.) Vill. on germination and early stages of growth of *Raphanus sativus* var. *radicula*. *Annales Universitatis Paedagogicae Cracoviensis Studia Naturae* 3:90-99. <https://doi.org/10.24917/25438832.3.7>
- Zhang S, Jin Y, Tang J, Chen X (2009). The invasive plant *Solidago canadensis* L. suppresses local soil pathogens through allelopathy. *Applied Soil Ecology* 41(2):215-222. <https://doi.org/10.1016/j.apsoil.2008.11.002>
- Zhang SS, Zhu W, Wang B, Tang JJ, Chen X (2011). Secondary metabolites from the invasive *Solidago canadensis* L. accumulation in soil and contribution to inhibition of soil pathogen *Pythium ultimum*. *Applied Soil Ecology* 48(3):280-286. <https://doi.org/10.1016/j.apsoil.2011.04.011>
- Zhao Y, Padilla-Zakour O, Yang G (2005). Polyphenols, antioxidant and antimicrobial activities of leaf and bark extracts of *Solidago canadensis* L. *Industrial Crops Products* 74:803-809. <https://doi.org/10.1016/j.indcrop.2015.06.014>



The journal offers free, immediate, and unrestricted access to peer-reviewed research and scholarly work. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.



License - Articles published in *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* are Open-Access, distributed under the terms and conditions of the Creative Commons Attribution (CC BY 4.0) License.

© Articles by the authors; UASVM, Cluj-Napoca, Romania. The journal allows the author(s) to hold the copyright/to retain publishing rights without restriction.