

Alleviation of Salt Stress in Seedlings of Black Glutinous Rice by Seed Priming with Spermidine and Gibberellic Acid

Sumitahnun CHUNTHABUREE¹, Jirawat SANITCHON²,
Wattana PATTANAGUL¹, Piyada THEERAKULPISUT^{1*}

¹Department of Biology, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand; piythe@gmail.com (*corresponding author)

²Department of Plant Science and Agricultural Resources, Faculty of Agriculture, Khon Kaen University, Khon Kaen 40002, Thailand

Abstract

This study was carried out to elucidate the spermidine (Spd) and gibberellic acid (GA₃) priming-induced physiological and biochemical changes responsible for induction of salinity tolerance in two rice (*Oryza sativa* L.) cultivars, namely 'Niewdam Gs. no. 00621' (salt tolerant) and 'KKU-LLR-039' (salt sensitive). The seeds of the two cultivars were primed separately with distilled water, 1 mM Spd or 0.43 mM GA₃. Primed seeds were germinated and the resultant seedlings were hydroponically grown for 14 days before being exposed to salinity stress (150 mM NaCl) for 10 days. Seed priming with Spd or GA₃ slightly improved salt-induced reductions in growth, anthocyanin and chlorophyll contents of the seedlings. Salt stress induced pronounced increases in Na⁺/K⁺ ratio, proline and H₂O₂ contents, particularly in the sensitive cultivar. The levels of these salt-sensitivity physiological indicators tended to be mitigated by priming with Spd and GA₃. Salt-stressed seedlings grown from seeds primed with these growth regulators also possessed higher phenolic contents and greater antioxidant capacity than the control seedlings. Based on all growth and physiological data, Spd tended to be more effective than GA₃ in improving salt tolerance in both rice cultivars.

Keywords: antioxidant capacity, Na⁺/K⁺, polyamine, seed pretreatment

Introduction

Soil salinity is a major environmental stress that drastically affects plants growth and productivity of several crops by creating low osmotic potential outside the seeds, thereby inhibiting or slowing down seed germination and later retarding seedling growth because of both osmotic and ion toxicity effects (Khajeh-Hosseini *et al.*, 2003). Numerous attempts have been made to improve the salinity tolerance of a variety of crops by traditional breeding programs (Duan *et al.*, 2008), but the progress to develop such salt-resistant plants is very slow and commercial success is limited. Therefore, the use of some other simple, cost-effective methods, such as seed priming could be an attractive approach to overcome growth retardation of young plants in the salt-affected soils. Seed quality, seed germination rate and seedling vigor are important factors for sustainable crop production particularly under adverse environmental conditions (Koorneef *et al.*, 2002; Sun *et al.*, 2007). Priming is a process by which seeds are exposed to restricted water availability under controlled conditions, which allows some pre-germination metabolic activities (physiological and chemical) to proceed, before germination is completed, followed by re-drying for short-term storage before sowing (Bradford, 1986; Farooq *et al.*, 2010). The

effectiveness of different priming agents, however, varies under different stresses as well as in different crop species. Plant hormones are active members of the signal cascade involved in the induction of plant stress responses (Pedranzani *et al.*, 2003). Several chemicals have been used for seed priming (Afzal *et al.*, 2012). Of various priming agents employed for seed priming, plant growth regulators have gained much attention from researchers all over the world because of their consistent effects on seed germination as well as growth of a variety of plant species (Farooq *et al.*, 2007; Iqbal and Ashraf, 2013).

Polyamines (PAs), including the triamine spermidine (Spd³⁺), tetramine spermine (Spm⁴⁺) and their obligate precursor, the diamine putrescine (Put²⁺) are now regarded as plant growth regulators and secondary messenger in signaling pathways (Kusano *et al.*, 2008). Because of their cationic nature at physiological pH, PAs are able to interact with proteins, nucleic acids, membrane phospholipids and cell wall constituents, thereby stabilizing these molecules (Bouchereau *et al.*, 1999). Apart from their implication in growth and development, PAs have been reported to be involved in defense response to biotic and abiotic stresses (Alcázar *et al.*, 2010). Exogenously applied PAs have been reported to substantially enhance salt tolerance in rice plants (Chattopadhyay *et al.*, 2002). Moreover, Khan *et al.* (2012)

reported that seed priming with Put, Spd and Spm improved seed germination as well as promoted early seedling growth in hot pepper.

Gibberellins (GAs) play an important role in many essential plant growth and development processes, including seed germination, stem elongation, leaf expansion, flower and fruit development, and floral transition (Razem *et al.*, 2006). The mechanisms by which Spd and GA₃ priming could induce salt tolerance in plants have been widely proposed (Khan *et al.*, 2012; Iqbal and Ashraf, 2013). However, studies concerning Spd and GA₃ priming-induced physiological and biochemical changes in rice in saline conditions are scarce and little is known about the effect of GAs on physiological and biochemical changes in induction of salt tolerance in rice plants. Thus, investigating the possibility of exogenously applied Spd and GA₃ in induction of salt tolerance in rice is of great importance. The overall objective of this study was to elucidate the Spd and GA₃ priming-induced physiological and biochemical changes responsible for induction of salinity tolerance in two rice (*Oryza sativa* L.) cultivars, namely, 'Niewdam Gs. no. 00621' and 'KKU-LLR-039'.

Materials and methods

Plant materials and priming treatments

Two black, glutinous rice (*O. sativa* L.) cultivars, 'Niewdam Gs. no. 00621' (salt-tolerant) and 'KKU-LLR-039' (salt-sensitive) were used in the present study. The seeds were obtained from the Department of Plant Science and Agricultural Resources, Faculty of Agriculture, Khon Kaen University, Thailand. Solutions of 1 mM Spd and 0.43 mM (150 mg L⁻¹) GA₃ were utilized for seed priming. Distilled water was used for hydro-priming. Prior to priming treatments, rice seeds were surface-sterilized in 0.5% sodium hypochlorite for 10 min and rinsed thrice with distilled water. The rice seeds were primed separately, in the dark, in 5 ml solutions of GA₃, Spd and distilled water for 24 hours at ambient temperature. Following pre-soaking treatments, the seeds were rinsed with distilled water and air-dried for 48 hours at ambient temperature to their original moisture content (about 9-10%). The primed and non-primed seeds were sown for 3 days to produce seedlings. The obtained seedlings were transferred to continuously aerated Yoshida's solution in a greenhouse with twenty seedlings per treatment. The solution pH was adjusted to 5.0 and was renewed at 4-days intervals. Following a 14-day planting period, the seedlings were transferred to Yoshida's solution supplemented with 150 mM NaCl for 10 days. The seedlings that were cultivated in the absence of additional NaCl were set as controls. Seedlings were harvested after 10 days, separated into aerial and radical parts for growth analyses, including shoot length (SL), root length (RL), fresh weight (FW) and dry weight (DW) of shoot and root. To measure the seedling DW, the samples were dried in a hot-air oven at 70 °C for 3 days. Leaf samples were collected and stored at -80 °C for further use in physiological and biochemical analyses.

Measurement of ion concentration

Following 3 days of oven-drying of seedling shoots at 60 °C, about 0.5 g of each dried powdered sample was successively digested with 10 ml of nitric acid at 300 °C, 5 ml of perchloric

acid at 200 °C and 20 ml of 6 M hydrochloric acid at 200 °C, for 30 min. The concentrations of Na⁺ and K⁺ were analyzed using atomic absorption spectroscopy.

Analysis of pigment accumulation

Chlorophyll content was determined following the method described by Arnon (1949) with minor modifications. About 0.5 g of leaf samples were extracted for chlorophyll by soaking the leaf samples in 10 ml of 80% acetone overnight in the dark. The extract was centrifuged at 10000 × g for 5 min and the absorbance of the supernatant was read at 645 and 663 nm using a spectrophotometer and chlorophyll concentrations were then calculated and expressed as mg g⁻¹ FW, using the following equations:

$$\text{Total-Chl} = [20.2(A_{645}) + 8.02(A_{663})] \times (V/1000W),$$

$$\text{Chl-}a = [12.7(A_{663}) - 2.69(A_{645})] \times (V/1000W),$$

$$\text{Chl-}b = [22.9(A_{645}) - 4.69(A_{663})] \times (V/1000W).$$

where A₆₄₅ and A₆₆₃ represent absorbance of chlorophyll concentrations extract at 645 and 663 nm respectively, V is the total extract volume and W is the leaf fresh weight.

Total anthocyanin content (TAC) was estimated according to the method described by Abdel-Aal and Hucl (1999). Approximately 0.1 g of leaf tissues were soaked for 72 hour in 10 ml of acidified ethanol (ethanol : 1N HCl, 85:15 v/v). The suspension was filtered through Whatman No.1 filter paper and absorbance was measured at 535 nm.

Analysis of proline content

Proline content was quantified according to the method described by Bates *et al.* (1973) with minor modifications. About 0.1 g fresh weight of leaf tissues were homogenized and extracted utilizing 5 ml of 3% aqueous sulfosalicylic acid. Then 2 ml of extract were reacted with 2 ml of acid ninhydrin and 2 ml of glacial acetic acid and boiled at 100 °C for 1 hour. The reaction was quenched by placing the tubes rapidly on ice. The resulting solutions were extracted with toluene, and the absorbance of the toluene fraction was monitored at 520 nm. Proline content was estimated with reference to a calibration curve and expressed as μg g⁻¹ tissue FW.

Analysis of H₂O₂

For determination of H₂O₂ concentration, 0.1 g fresh weight of leaf tissues were extracted with 3 ml of 0.1 %, w/v trichloroacetic acid (TCA) in an ice bath and centrifuged at 12,000 × g for 15 min and 0.5 ml of the supernatant was added to 0.5 ml of 10 mM potassium phosphate buffer (pH 7.0) and 1 ml of 1 M potassium iodide. The absorbance of supernatant was read at 390 nm. The content of H₂O₂ was estimated based on a standard curve (Velikova *et al.*, 2000).

Extraction procedure to determine the antioxidant properties

The method of extraction for determination of antioxidant properties of rice leaf was adapted from Sultana *et al.* (2009). Rice leaf samples (0.1 g) were extracted with 10 ml of 80% aqueous ethanol for 48 hours at room temperature. After centrifugation at 10,000 × g for 10 min, the suspension was filtered through Whatman No.1 filter paper and used to determine total phenolic content (TPC) and antioxidant capacity and the measurements were replicated four times.

Analysis of TPC

Total phenolic content was assayed by Folin-Ciocalteu's reagent method (Singleton *et al.*, 1999). Briefly, 120 μ l of leaf extract was placed into test tubes and then 600 μ l of freshly diluted 10-fold Folin-Ciocalteu's reagent was added. After 2 min, 480 μ l of sodium carbonate solution (75 g L⁻¹) was added. The mixtures were vigorously shaken and allowed to stand for 40 min. The absorbance of the resulting blue color was measured at 760 nm. Gallic acid was used as standard and TPC and was expressed as mg gallic acid (GAE) equivalent per 1 g leaf FW (mg GAE g⁻¹).

2,2'-diphenyl-1-picrylhydrazyl (DPPH) radical-scavenging assay

The DPPH radical-scavenging ability of rice extracts was estimated according to the method described by Brand-Williams *et al.* (1995). The reaction mixture contained 1.5 ml DPPH working solution (4.73 mg of DPPH in 100 ml ethanol HPLC-grade) and 100 μ l rice leaf extract. The mixture was shaken and incubated for 1 hour in the dark at room temperature. The absorbance was read at 515 nm relative to the control using a spectrophotometer. The percentage of radical scavenging was calculated according to the following equation:

$$\% \text{ DPPH scavenging activity (\%)} = \frac{[(\text{Abs}_{\text{control}} - \text{Abs}_{\text{sample}})]}{(\text{Abs}_{\text{control}})} \times 100,$$

where Abs_{control} is the absorbance of DPPH + ethanol;

Abs_{sample} is the absorbance of DPPH radicals + sample extract.

2,2'-azino-bis (3-ethylbenzothiazoline 6-sulfonic acid) (ABTS⁺) scavenging assay

Radical cation ABTS⁺ scavenging assay was performed following the method described by Re *et al.* (1999) with minor modifications. A stable stock solution of ABTS radicals was prepared by reacting 7 mM aqueous solution of ABTS with potassium persulfate in the dark at room temperature for 12-16 hours before use. Rice leaf extract (120 μ l) was allowed to react with 1.5 ml of a diluted ABTS radical solution (absorbance of 0.70 \pm 0.02 at 734 nm). The absorbance at 734 nm of the mixture was measured after a 30-min reaction period. The percentage of radical scavenging was calculated according to the below-given equation:

$$\% \text{ ABTS scavenging activity} = \frac{[(\text{Abs}_{\text{control}} - \text{Abs}_{\text{sample}})]}{(\text{Abs}_{\text{control}})} \times 100,$$

where Abs_{control} is the absorbance of ABTS⁺ radical + ethanol;

Abs_{sample} is the absorbance of ABTS⁺ radical + sample extract.

Ferric reducing antioxidant power (FRAP) assay

The method adapted from Benzie and Strain (1996) was used for the FRAP assay. A 200 μ l volume of rice leaf extract was mixed with 1.3 ml of FRAP reagent and the mixture was incubated at 37 °C for 30 min. Following incubation, the absorbance of the mixture was read at 595 nm. The FRAP reagent was freshly prepared as required and consisted of 0.3 M acetate buffer (pH 3.6), 10 mM 2, 4, 6-tripyridyl-s-triazine (TPTZ) in 40 mM HCl and 20 mM FeCl₃ (10:1:1, in v/v/v). The antioxidant activity was expressed as mmol of Trolox equivalents (TE) per 100 g FW.

Statistical analysis

The experiments were arranged as a factorial experiment based on a completely randomized design with 4 replications. Treatments included two rice cultivars, four priming treatments (unprimed, hydropriming, priming with Spd and with GA₃) and two levels of salinity (0 and 150 mM NaCl). The data was subjected to an analysis of one-way and two-way ANOVA and all means were separated at the $p < 0.05$ level using the Duncan's multiple range test. All calculations and data analyses were performed using the SPSS 16.0 for Windows software package. Pearson's correlation was calculated for the relationship between seedling growth, physiological parameters and also antioxidant capacity of seedlings grown from primed and non-primed seeds under saline conditions.

Results and discussions*Spd and GA₃ priming modulates growth parameters*

The imposition of salt stress caused a significant reduction in seedling growth of both rice cultivars (Fig. 1). A significant increase in shoot length was observed in 'KKU-LLR-039' when Spd was used as a priming agent. Analysis of variance (Tab. 3) showed that all six growth traits were significantly affected by salt stress while only SL, SFW and SDW were significantly affected by different priming treatments and there were no cultivars and priming treatments (C \times P) interaction for all of the growth traits. Priming with Spd and GA₃ improved root length of both cultivars although not statistically significantly. Gibberellins are generally involved in growth and development; they control seed germination, leaf expansion, stem elongation and flowering (Magome *et al.*, 2004). Gibberellic acid has been reported to be helpful in enhancing growth under saline conditions in rice (Wen *et al.*, 2010), corn (Ghodrat and Roust, 2012), wheat (Iqbal and Ashraf, 2013) and cucumber (Radhakrishnan and Lee, 2014). Of the two priming agents applied, Spd was more effective in enhancing salt tolerance in both cultivars. Furthermore, exogenous Spd also has been reported to be effective in enhancing rice growth under saline conditions (Saleethong *et al.*, 2011; Roychoudhury *et al.*, 2011).

Spd and GA₃ priming alters Na⁺/K⁺ ratios

Salt stress caused a significant increase in Na⁺/K⁺ ratios in seedlings of both cultivars grown from unprimed seeds; by 146.65% for the salt-tolerant cultivar 'Niewdam Gs. no. 00621' and by 188.98% for the salt-sensitive cultivar 'KKU-LLR-039' (Tab. 1). Priming with Spd and GA₃ significantly reduced the seedling Na⁺/K⁺ ratios in both cultivars. However, the interaction between cultivars and priming treatments (C \times P) was not significant (Tab. 3). Considering consistency and effectiveness of the priming treatments, Spd treatments were more effective in reducing the seedling Na⁺/K⁺ ratios in both cultivars in saline conditions. Rice plants with salt-resistant characteristics commonly possess superior ion homeostasis strategies, particularly a high K⁺/Na⁺ ratios, through exclusion, compartmentation and partitioning of Na⁺ in shoots or roots (Blumwald, 2000). Under saline condition, the Na⁺/K⁺ ratios were positively correlated with proline and H₂O₂ but negatively correlated with SL, Chl-*b*, TAC, TPC and antioxidant capacities (Tab. 4). Based on our study, pre-treatment of seeds

with Spd and GA₃ was noted to alleviate the inhibitory effect of salt stress in both cultivars under saline conditions. The findings in our study were well-supported by an earlier study (Iqbal and Ashraf, 2013) that seed priming with GA₃ decreased Na⁺ concentrations in the roots and shoots of two wheat cultivars, namely MH-97 (salt-intolerant) and Inqlab-91 (salt-tolerant), when grown under saline conditions. Zhu *et al.* (2006) also reported that the exogenous application of PAs, specifically Spd inhibits Na⁺ transport in barley seedlings under

salinity conditions. Moreover, Salethong *et al.* (2011) observed that exogenously applied Spd contributed to the improvement of K⁺/Na⁺ homeostasis in rice seedlings exposed to salt stress. Even though the mechanisms underlying the alleviation of the inhibitory effect of salinity by PAs have not yet been fully understood, several previous studies have suggested that exogenous application of PAs contributed to the reduced accumulation of toxic Na⁺ in rice (Chattopadhyay *et al.*, 2002; Quinet *et al.*, 2010).




Fig. 1. Growth parameters of rice seedlings in saline conditions as affected by priming seeds with H₂O, Spd and GA₃. Results are expressed as means ± standard errors (SE). Bars with different letters are significantly different according to Duncan's multiple range tests (p < 0.05).

Tab. 1. Effects of seed priming with H₂O, Spd and GA₃ on some physiological traits of rice seedlings grown under salt stress (NaCl 150 mM) and non-stress conditions (Control)

| Cultivars | Treatment | Na ⁺ /K ⁺ | | | Proline (µg g ⁻¹ FW) | | | TAC (mg g ⁻¹ FW) | | |
|-----------------------|-------------------------|-----------------------------------|--------------------|--------|---------------------------------|--------------------|--------|-------------------------------|--------------------|-------|
| | | Control | Salt | % (+) | Control | Salt | % (+) | Control | Salt | % (-) |
| Niewdam Gs. no. 00621 | Unprimed | 0.48 ^a | 1.19 ^d | 146.65 | 183 ^a | 1.77 ^b | 873.70 | 2.15 ^e | 0.95 ^b | 55.74 |
| | Hydropriming | 0.47 ^a | 1.14 ^d | 140.65 | 170 ^a | 1.69 ^b | 897.65 | 2.28 ^f | 1.13 ^b | 50.26 |
| | Spd 1 mM | 0.45 ^a | 1.01 ^b | 126.79 | 181 ^a | 1.51 ^b | 737.12 | 2.39 ^f | 1.22 ^b | 48.96 |
| | GA ₃ 0.43 mM | 0.45 ^a | 1.03 ^b | 128.60 | 187 ^a | 1.55 ^b | 731.55 | 2.32 ^f | 1.18 ^b | 49.07 |
| KKU-LLR-039 | Unprimed | 0.49 ^a | 1.42 ^f | 188.98 | 194 ^a | 2.04 ^b | 951.80 | 1.58 ^c | 0.64 ^c | 59.55 |
| | Hydropriming | 0.46 ^a | 1.35 ^f | 193.91 | 184 ^a | 1.96 ^b | 972.48 | 1.65 ^{cd} | 0.76 ^c | 53.93 |
| | Spd 1 mM | 0.45 ^a | 1.19 ^d | 164.40 | 186 ^a | 1.68 ^b | 804.30 | 1.88 ^{bc} | 0.90 ^b | 52.02 |
| | GA ₃ 0.43 mM | 0.46 ^a | 1.26 ^e | 176.86 | 190 ^a | 1.72 ^b | 809.23 | 1.90 ^{bc} | 0.91 ^b | 51.99 |
| Cultivars | Treatment | Total-chl (mg g ⁻¹ FW) | | | Chl-a (mg g ⁻¹ FW) | | | Chl-b (mg g ⁻¹ FW) | | |
| | | Control | Salt | % (-) | Control | Salt | % (-) | Control | Salt | % (-) |
| Niewdam Gs. no. 00621 | Unprimed | 4.21 ^{bd} | 2.92 ^a | 30.69 | 3.07 ^{cd} | 1.98 ^a | 35.71 | 1.12 ^{bc} | 0.93 ^b | 16.83 |
| | Hydropriming | 4.50 ^{cd} | 3.41 ^{ab} | 24.19 | 3.25 ^{cd} | 2.34 ^{ab} | 28.18 | 1.23 ^{bc} | 1.06 ^{bc} | 13.60 |
| | Spd 1 mM | 4.57 ^{cd} | 3.53 ^{ab} | 22.78 | 3.34 ^{cd} | 2.41 ^{ab} | 27.79 | 1.21 ^{bc} | 1.11 ^{bc} | 8.90 |
| | GA ₃ 0.43 mM | 4.63 ^{cd} | 3.60 ^{ab} | 22.41 | 3.38 ^{cd} | 2.43 ^{ab} | 28.28 | 1.23 ^{bc} | 1.16 ^{bc} | 6.25 |
| KKU-LLR-039 | Unprimed | 4.82 ^{cd} | 3.63 ^{ab} | 24.55 | 3.59 ^{cd} | 2.68 ^{cd} | 25.49 | 1.21 ^{bc} | 0.95 ^b | 21.71 |
| | Hydropriming | 4.83 ^{cd} | 3.66 ^{ab} | 24.21 | 3.60 ^{cd} | 2.78 ^{cd} | 22.56 | 1.21 ^{bc} | 0.86 ^c | 29.11 |
| | Spd 1 mM | 5.04 ^d | 4.00 ^{bc} | 20.56 | 3.72 ^d | 2.79 ^{cd} | 24.81 | 1.30 ^c | 1.19 ^{bc} | 8.41 |
| | GA ₃ 0.43 mM | 5.03 ^d | 4.03 ^{bc} | 19.87 | 3.67 ^d | 2.92 ^{bc} | 20.39 | 1.35 ^c | 1.10 ^{bc} | 18.43 |

Means in columns followed by same superscript letters are not significantly different according to Duncan's multiple range test (p < 0.05). TAC = Total anthocyanin content, Total-chl = Total chlorophyll, Chl-a = Chlorophyll a, Chl-b = Chlorophyll b

Tab. 2. Effects of seed priming with H₂O, Spd and GA₃ on H₂O₂, TPC and antioxidant capacity (DPPH, ABTS and FRAP assays) of rice seedlings grown under salt stress (NaCl 150 mM) and non-stress conditions (Control)

| Cultivars | Treatment | H ₂ O ₂ | | | TPC (mg g ⁻¹ FW) | | | DPPH (% radical scavenging) | | | ABTS (% radical scavenging) | | | FRAP (mmol TE g ⁻¹ 100 FW) | | |
|-----------------------------|-------------------------|-------------------------------|--------------------|--------|-----------------------------|---------------------|-------|-----------------------------|---------------------|-------|-----------------------------|---------------------|-------|---------------------------------------|--------------------|-------|
| | | Control | Salt | % (+) | Control | Salt | % (-) | Control | Salt | % (-) | Control | Salt | % (-) | Control | Salt | % (-) |
| Niewdam Gs. no. 00621 | Unprimed | 5.26 ^c | 25.50 ^b | 384.47 | 1.16 ^d | 0.84 ^{ab} | 27.75 | 81.09 ⁶ | 67.51 ^d | 16.75 | 62.16 ^c | 45.54 ^{bc} | 26.74 | 0.42 ^{ab} | 0.29 ^{cd} | 30.22 |
| | Hydropriming | 5.29 ^a | 24.35 ^b | 360.45 | 1.20 ^{bc} | 0.86 ^{abc} | 28.20 | 81.64 ⁶ | 69.42 ^{bc} | 14.97 | 64.19 ^c | 48.18 ^{bc} | 24.94 | 0.44 ^{ab} | 0.30 ^{cd} | 30.11 |
| | Spd 1 mM | 5.07 ^a | 21.90 ^b | 331.75 | 1.35 ^c | 1.02 ^{ab} | 24.87 | 88.38 ⁶ | 76.58 ^d | 13.35 | 68.51 ^c | 53.14 ^{cd} | 22.44 | 0.48 ^a | 0.35 ^{cd} | 27.14 |
| | GA ₃ 0.43 mM | 5.36 ^c | 22.72 ^b | 323.77 | 1.34 ^c | 0.99 ^{bc} | 25.94 | 88.15 ⁶ | 74.75 ^{cd} | 15.21 | 68.15 ^c | 52.75 ^{cd} | 22.61 | 0.48 ^a | 0.34 ^{cd} | 27.46 |
| KKU- LLR-039 | Unprimed | 6.44 ^d | 26.01 ^b | 303.73 | 0.95 ^{cd} | 0.71 ^a | 25.75 | 56.39 ^c | 29.68 ^a | 47.37 | 48.40 ^c | 33.59 ^a | 30.58 | 0.31 ^{cd} | 0.13 ^a | 58.51 |
| | Hydropriming | 6.63 ^c | 24.76 ^b | 273.19 | 0.94 ^{cd} | 0.74 ^a | 21.29 | 56.74 ^c | 38.43 ^b | 32.27 | 46.83 ^c | 38.62 ^b | 17.52 | 0.31 ^{cd} | 0.18 ^a | 41.55 |
| | Spd 1 mM | 5.99 ^b | 20.91 ^b | 249.40 | 1.06 ^{cd} | 0.86 ^{abc} | 19.00 | 71.23 ^{bc} | 49.52 ^c | 30.47 | 52.75 ^{cd} | 43.72 ^{bc} | 17.11 | 0.39 ^{cd} | 0.24 ^b | 38.32 |
| | GA ₃ 0.43 mM | 6.08 ^b | 22.31 ^b | 266.80 | 1.04 ^{cd} | 0.87 ^{abc} | 16.77 | 70.85 ^{bc} | 50.40 ^c | 28.86 | 51.89 ^c | 44.54 ^{bc} | 14.16 | 0.37 ^{cd} | 0.25 ^b | 33.13 |

Means in columns followed by same superscript letters are not significantly different according to Duncan's multiple range test (p < 0.05). TPC = Total phenolic content. DPPH = 2,2'-diphenyl-1-picrylhydrazyl, ABTS = 2,2'-azinobis (3-ethylbenzothiazoline 6-sulfonic acid), FRAP = Ferric reducing antioxidant power

Tab. 3. Mean squares from analysis of variance for growth traits, physiological characteristics and antioxidant capacity in two cultivars of rice seedlings grown under salt stress from unprimed seeds or seeds primed with H₂O, Spd and GA₃

| Source of variation | df | SL (cm) | RL (cm) | SFW (g/plant) | RFW (g/plant) | SDW (g/plant) | RDW (g/plant) | Na ⁺ /K ⁺ | Total-chl (mg g ⁻¹ FW) | Chl-a (mg g ⁻¹ FW) | Chl-b (mg g ⁻¹ FW) |
|---------------------|----|----------|---------|---------------|---------------|---------------|---------------|---------------------------------|-----------------------------------|-------------------------------|-------------------------------|
| Cultivars (C) | 1 | 2.93ns | 13.868* | 0.0705ns | 0.00233ns | 0.01136* | 0.00077* | 0.08925* | 3.3658* | 3.1301* | 0.0033ns |
| Priming (P) | 3 | 93.06* | 0.537ns | 0.1151* | 0.00256ns | 0.00512* | 0.00014ns | 0.02376* | 0.6280ns | 0.2336ns | 0.1032* |
| Salt (S) | 1 | 8298.97* | 133.00* | 31.7327* | 2.02126* | 0.32880* | 0.3236* | 4.29978* | 19.5687* | 13.31* | 0.5732* |
| C x P | 3 | 1.03ns | 0.063ns | 0.0280ns | 0.00991ns | 0.0016ns | 0.00006ns | 0.00069ns | 0.0980ns | 0.0497ns | 0.0303ns |
| P x S | 3 | 0.38ns | 5.204ns | 0.0301ns | 0.02467* | 0.00163ns | 0.00010ns | 0.01121* | 0.0394ns | 0.199ns | 0.1801ns |
| C x S | 1 | 51.56* | 10.360* | 0.0778ns | 0.00001ns | 0.00247ns | 0.00080* | 0.09138* | 0.0013ns | 0.0633ns | 0.4673ns |
| C x P x S | 4 | 8.08ns | 6.119* | 0.0306ns | 0.01611* | 0.00130ns | 0.00018ns | 0.01801* | 0.00493ns | 0.02193ns | 0.01731ns |
| CV (%) | | 6.72 | 12.58 | 14.78 | 24.20 | 15.56 | 23.33 | 4.75 | 12.18 | 13.67 | 16.02 |

| Source of variation | df | TAC (mg g ⁻¹ FW) | Proline (ug g ⁻¹ FW) | H ₂ O ₂ | TPC (mg g ⁻¹ FW) | DPPH (% inhibition) | ABTS (%inhibition) | FRAP (mmol TE g ⁻¹ 100 FW) |
|---------------------|----|-----------------------------|---------------------------------|-------------------------------|-----------------------------|---------------------|--------------------|---------------------------------------|
| Cultivars (C) | 1 | 2.9096* | 205889* | 3.38ns | 0.6376* | 10432.1* | 2615.42* | 0.2134* |
| Priming (P) | 3 | 0.2475* | 89613.4* | 18.06ns | 0.1096* | 682.6* | 202.57* | 0.0223* |
| Salt (S) | 1 | 17.88* | 3.893E+07* | 5064.58* | 1.17208* | 4773.4* | 2641.02* | 0.30523* |
| C x P | 3 | 0.170* | 3822.90ns | 0.88ns | 0.0023ns | 96.7* | 1.08ns | 0.0013ns |
| P x S | 3 | 0.0027ns | 91071.9* | 13.03ns | 0.00034ns | 16.9ns | 15.10ns | 0.00068ns |
| C x S | 1 | 0.1843* | 1777031* | 0.88ns | 0.06850* | 327.0* | 144.45ns | 0.00106ns |
| C x P x S | 4 | 0.04812ns | 46277* | 1.554ns | 0.01831ns | 89.3* | 42.39ns | 0.00089ns |
| CV (%) | | 13.32 | 9.16 | 24.25 | 10.10 | 7.96 | 11.90 | 11.57 |

SL = Shoot length, RL = Root length, SFW = Shoot fresh weight, RFW = Root fresh weight, SDW = Shoot dry weight, RDW = Root dry weight, Total-chl = Total chlorophyll content, Chl-a = Chlorophyll a, Chl-b = Chlorophyll b, TAC = Total anthocyanin content, TPC = Total phenolic content, DPPH = 2,2'-diphenyl-1-picrylhydrazyl, ABTS = 2,2'-azinobis (3-ethylbenzothiazoline 6-sulfonic acid), FRAP = Ferric reducing antioxidant power.

Tab. 4. Pearson's correlation coefficient of growth traits, physiological characteristics and antioxidant capacity in two rice cultivars under saline conditions

| Physiological traits | SL | RL | SFW | SDW | RFW | RDW | Na ⁺ /K ⁺ | Total-chl | Chl-a | Chl-b | TAC | Proline | H ₂ O ₂ | TPC | DPPH | ABTS | FRAP |
|---------------------------------|----|------|--------------------|--------------------|--------------------|--------------------|---------------------------------|-------------------|--------------------|---------------------|---------------------|---------------------|-------------------------------|---------------------|---------------------|---------------------|---------------------|
| SL | 1 | .440 | .893 ^{**} | .790 ^{**} | .723 [*] | .724 [*] | -.802 ^{**} | .290 | .028 | .756 [*] | .734 [*] | -.885 ^{**} | -.911 ^{**} | .847 ^{**} | .630 [*] | .785 [*] | .732 [*] |
| RL | | 1 | .506 | .669 [*] | .700 [*] | .817 ^{**} | -.205 | .489 | .450 | .259 | .331 | -.344 | -.461 | .342 | .164 | .336 | .264 |
| SFW | | | 1 | .878 ^{**} | .917 ^{**} | .851 ^{**} | -.585 | .521 | .276 | .786 [*] | .515 | -.753 ^{**} | -.955 ^{**} | .684 [*] | .414 | .610 | .550 |
| SDW | | | | 1 | .894 ^{**} | .885 ^{**} | -.546 | .334 | .146 | .585 | .530 | -.715 ^{**} | -.780 ^{**} | .689 [*] | .474 | .625 [*] | .596 |
| RFW | | | | | 1 | .875 ^{**} | -.292 | .688 [*] | .537 | .603 | .261 | -.511 | -.860 ^{**} | .459 | .136 | .361 | .291 |
| RDW | | | | | | 1 | -.537 | .503 | .297 | .690 [*] | .582 | -.711 ^{**} | -.779 ^{**} | .667 [*] | .449 | .636 [*] | .573 |
| Na ⁺ /K ⁺ | | | | | | | 1 | .197 | .479 | -.661 ^{**} | -.974 ^{**} | .964 ^{**} | .574 | -.960 ^{**} | -.951 ^{**} | -.982 ^{**} | -.973 ^{**} |
| Total-chl | | | | | | | | 1 | .942 ^{**} | .479 | -.221 | -.019 | -.622 [*] | -.019 | -.435 | -.183 | -.296 |
| Chl-a | | | | | | | | | 1 | .155 | -.482 | .288 | -.362 | -.303 | -.674 [*] | -.451 | -.555 |
| Chl-b | | | | | | | | | | 1 | .600 | -.796 ^{**} | -.864 ^{**} | .730 [*] | .473 | .628 [*] | .567 |
| TAC | | | | | | | | | | | 1 | -.933 ^{**} | -.493 | .927 ^{**} | .964 ^{**} | .985 ^{**} | .976 ^{**} |
| Proline | | | | | | | | | | | | 1 | .741 [*] | -.977 ^{**} | -.880 ^{**} | -.962 ^{**} | -.938 ^{**} |
| H ₂ O ₂ | | | | | | | | | | | | | 1 | -.679 [*] | -.346 | -.560 | -.482 |
| TPC | | | | | | | | | | | | | | 1 | .876 [*] | .960 ^{**} | .932 [*] |
| DPPH | | | | | | | | | | | | | | | 1 | .961 ^{**} | .986 ^{**} |
| ABTS | | | | | | | | | | | | | | | | 1 | .991 ^{**} |

SL = Shoot length, RL = Root length, SFW = Shoot fresh weight, RFW = Root fresh weight, SDW = Shoot dry weight, RDW = Root dry weight, Total-chl = Total chlorophyll content, Chl-a = Chlorophyll a, Chl-b = Chlorophyll b, TAC = Total anthocyanin content, TPC = Total phenolic content, DPPH = 2,2'-diphenyl-1-picrylhydrazyl, ABTS = 2,2'-azinobis (3-ethylbenzothiazoline 6-sulfonic acid), FRAP = Ferric reducing antioxidant power.

** Correlation is significant at the p < 0.01; * Correlation is significant at the p < 0.05

Spd and GA₃ priming modulates pigment accumulation

The imposition of salinity caused a significant reduction in chlorophyll accumulation in both cultivars (Tab. 1). Salt stress generally causes the inhibition of pigment accumulation in plants owing to the instability of protein complex and destruction of chlorophyll by inducing the activity of chlorophyllase, chlorophyll-degrading enzyme (Reddy and Vora, 1986). Protective roles of Spd on reduction of chlorophyll loss was previously reported by Roychoudhury *et al.* (2011) and Salethong *et al.* (2011). Salt stress also had a negative effect on TAC content. In seedlings grown from unprimed seeds, TAC contents in salt-stressed seedlings were reduced by 55.74% in 'Niewdam Gs. no. 00621' and 59.55% in 'KKU-LLR-039'. The findings in our study were in good agreement with an earlier study (Kachout *et al.*, 2013) demonstrating that high salt concentrations (180-260 mM) resulted in a significant reduction in anthocyanin contents in the leaves of *Atriplex hortensis*. Similarly, Trivellini *et al.* (2014) demonstrated that salt stress adversely affected the content of anthocyanins in *Hibiscus rosasinensis* L. (cv. Porto), resulting in a visually perceptible loss of color. By contrast, some previous studies have reported the contribution of salt stress to an increase in anthocyanin contents in plants (Eryilmaz, 2006; Daiponmak *et al.*, 2010). In general, changes in anthocyanin contents in plants depend largely on the level of salinity plants are exposed to, plant species and various environmental conditions (Chalker-Scott, 1999). The adverse effects of salt stress on TAC contents were slightly alleviated in seedlings raised from all three priming treatments. Our data demonstrated that seedlings of both rice cultivars exposed to salinity showed decreased pigment accumulation, but those raised from seeds pre-treated with Spd and GA₃ displayed improved accumulation of Chl-*a*, Chl-*b* and total-Chl, as well as TAC under saline conditions.

Analysis of variance (Tab. 3) revealed that the cultivar main effect was significant for TAC, total-Chl and Chl-*a*, but was not significant for Chl-*b*. The interaction between cultivars and priming treatments (C × P) was not significant for chlorophyll contents but was significant for TAC. For both cultivars, all three priming treatments had positive effects on maintaining the chlorophyll contents under salt stress (except the effects of hydropriming on Chl-*b* content of 'KKU-LLR-039'). However, priming with Spd and GA₃ were more effective than hydropriming. There was positive correlation of pigment accumulation with some growth traits (SL, SFW and RDW) under saline condition (Tab. 4). These findings in our study were well-supported by earlier reports (Ali *et al.*, 2012; Ratnakar and Rai, 2014) that exogenous application of GA₃ assisted in the restoration of altered pigment concentrations in saline conditions. Moreover, Salethong *et al.* (2011) reported that improved accumulation of total-Chl exerted by exogenous Spd was observed in rice seedlings under saline conditions. Roychoudhury *et al.* (2011) disclosed that salt-induced accumulation of anthocyanins in seedlings of three rice cultivars was substantially enhanced by exogenously applied Spd.

Spd and GA₃ priming modulates proline content

Proline accumulation is a well-known mechanism numerous plants have evolved to counteract salt stress. In addition to its crucial roles in protecting the subcellular

structures and mediating osmotic adjustment in stressed conditions (Parvaiz and Satyawati, 2008; Rao *et al.*, 2013a), proline is known to play diverse adaptive roles including protection of cellular functions by scavenging ROS (Smirnoff and Cumbes, 1989). Our data showed that seedlings under saline conditions accumulated higher levels of proline when compared with those under non-saline conditions. Salinity caused a dramatic increase (873% in 'Niewdam Gs. no. 00621' and 951% in 'KKU-LLR-039') in proline content in both cultivars (Tab. 1). However, seedlings raised from seeds pre-soaked in Spd and GA₃ was noted to exhibit decreased accumulation of proline in both rice cultivars in saline conditions (Tab. 1). Strong positive correlations were noted between proline content and the two salt-damaged parameters i.e. Na⁺/K⁺ and H₂O₂ while significant negative correlations were found between proline and seedling growth, TAC, TPC and antioxidant activity under salt conditions (Tab. 4). The cultivar main effect, salt and priming treatments were significant for proline. Nevertheless, there were no cultivars × priming treatments (C × P) interactions for proline (Tab. 3). The findings in our study were consistent with an earlier study (Salethong *et al.*, 2011) that exogenous Spd tangibly decreased the level of proline in rice seedlings under saline conditions. Moreover, exogenous application Spd was found to reduce the level of salt-induced proline, associated with improved leaf yield, chlorophyll and photosynthetic efficiency, in mulberry (Das *et al.*, 2002). However, Duan *et al.* (2008) reported that exogenously applied Spd substantially increased the level of proline in cucumber roots in saline conditions.

Spd and GA₃ priming alters H₂O₂

In general, the imposition of salinity causes damage to plant tissues as a result of excessive ROS like H₂O₂ produced at a high rate due to ion imbalance and hyperosmotic stresses. ROS accumulation leads to lipid oxidation and has a negative effect on cellular metabolism and physiology, thus detrimentally affecting the membrane integrity (Munns *et al.*, 2006). The imposition of salinity significantly increased H₂O₂ content in both cultivars. Under the control (non-stressed conditions), H₂O₂ content tended to be higher in 'KKU-LLR-039' than 'Niewdam Gs. no. 00621'. On the other hand, the lessened production of H₂O₂ was observed in seedlings of both cultivars raised from seeds pre-soaked in Spd and GA₃ (Tab. 2). Previous research has indicated that exogenously applied PAs resulted in decreased production of H₂O₂ under salinity (Verma and Mishra, 2005). Farooq *et al.* (2009) also found that under drought stress the curtailed production of H₂O₂ was observed in rice seedlings raised from seeds pre-treated with Spd. However, analysis of variance showed that the main effect of cultivars, priming treatments and their interactions were not significant for H₂O₂ (Tab. 3).

Spd and GA₃ priming modulates phenolic compound accumulation and antioxidant capacity

Phenolic compounds exhibit antioxidant capacity inactivating lipid free radicals or by preventing the decomposition of hydroperoxides into free radicals (Pokorný *et al.*, 2001). In our study, salinity caused a marked reduction in TPC in both rice cultivars (Tab. 2). The findings in our study were consistent with one previous study (Rao *et al.*, 2013b) that the accumulation of phenolic compounds in wheat plants was reduced under salt stress. Moreover, salinity stress was

reported to cause an inhibitory effect on the production of bound phenolic compounds in wheat and bean cultivars (Radi *et al.*, 2013). On the other hand, Valifard *et al.* (2014) elucidated that the accumulation of phenolic compounds was induced under moderate salinity whilst high salinity caused a reduction in phenolic contents, as observed for *Salvia mirzayanii*. Phenolic compounds may also be associated with antioxidant potential, due to their ability to directly protect the damage in plant cell (Wahid and Ghazanfar, 2006; Wahid, 2007). Seed priming with H₂O, Spd and GA₃ tended to improve the amount of TPC in both cultivars (Tab. 2). The significantly improved accumulation of TPC was noted in 'Niewdam Gs. no. 00621' in Spd treatments when compared with the unprimed treatment, but significant improvement in the accumulation of TPC was not observed in GA₃ treatments. The findings in our study corroborate one previous study (Farooq *et al.*, 2009) that exogenous Spd substantially increased the production of phenolic compounds in rice under drought stress. The activation of several enzymatic and non-enzymatic antioxidants can detoxify ROS in stressed cells (Blokchina *et al.*, 2003). In general, plant genotypes with stress-resistant traits possess superior ability of ROS scavenging through the production of higher levels of antioxidants. The positive roles of phenolic compounds on plant growth were evidenced by its strong positive correlations with shoot length, seedling fresh and dry weight, chl-*b*, TAC and its negative correlations with the negative physiological indicators for salt sensitivity, the Na⁺/K⁺ and proline content.

The antioxidant activity observed for both cultivars was significantly reduced under saline conditions (Tab. 2). Pre-treatment with Spd and GA₃, however, contributed to improved antioxidant performance. Seedlings of the salt-tolerant cultivar ('Niewdam Gs. no. 0062') raised from seeds primed with Spd displayed the significantly enhanced activity of DPPH, whereas significant improvement in antioxidant activity was not observed in seedlings raised from seeds pre-treated with H₂O and GA₃ in saline conditions. Seedlings of the salt-sensitive cultivar ('KKU-LLR-039') were raised from seeds pre-soaked in Spd and GA₃ showed a significant increase in the activities of DPPH, ABTS and FRAP under both saline and non-saline conditions. The findings in our study were consistent with an earlier study that exogenously applied Spd contributed to the elevated activity of antioxidant enzymes in rice (Roychoudhury *et al.*, 2011). Moreover, Choudhary *et al.* (2012) elucidated that exogenous application of Spd was noted to enhance the total antioxidant system in radish under Cu stress, as suggested by increased levels of DPPH and FRAP. Variations in antioxidant capacity were observed (Tab. 3). The main effect of cultivars and priming treatments were significant for TPC, DPPH, ABTS and FRAP. The interactions between cultivars and priming treatment were significant for only DPPH. Simultaneously phenolic compounds were positively correlated with the antioxidant capacity (DPPH, ABTS and FRAP) under saline condition (Tab. 4). Although the mechanisms underlying the functions of PAs in salt stress are not fully understood, recent studies have elucidated that exogenous application of PAs was observed to reduce salt-induced oxidative damage by activating antioxidant enzymes in Virginia pine (Tang and Newton, 2005). In addition, our results are supported by the findings reported by Verma and Mishra (2005) who found that PAs could enhance activities of several antioxidant enzymes and non-enzymatic antioxidants,

leading to less stress damage in *Brassica juncea* seedlings.

Although the mechanisms of actions is still unclear, our results suggested, in accordance with earlier reports, that exogenous application of Spd and GA₃ as seed priming is an effective approach for mitigating deleterious effects of salinity in rice seedlings. Seed priming with GA₃ helps accelerate metabolic reactions before germination process by modulating ion uptake and partitioning, enhancing ABA catabolism and modulating hormone homeostasis, thus allowing enhanced seed germination and seedling growth under salinity stress conditions (Gonai *et al.*, 2004; Iqbal and Ashraf, 2013). Exogenous Spd has been found to alleviate salt-stress damage by protecting membrane phospholipids, scavenging free radicals, activating antioxidant enzymes and prevented NaCl-induced K⁺ efflux by inhibition of non-selective cation channel (Shabala *et al.*, 2007; Groppa and Benavides, 2008). Moreover, Li *et al.* (2014) revealed that beneficial effects of Spd as the seed priming agent were associated with its effects on improving starch metabolism by enhancing amylase activities which is the well-known effect of GAs. Although Spd and GA₃ belong to different groups of hormones, their beneficial effects as seed-priming agents are strikingly similar. It would be extremely interesting to further elucidate the mechanisms of cross-talking between GA₃ and Spd to increase our understanding of integrating hormonal actions on alleviation of damaging effects of salinity stress.

Conclusion

The imposition of salinity caused a significant reduction in growth parameters in both rice cultivars. Pre-treatment of seeds with Spd and GA₃ improved growth of seedlings of both rice cultivars, with the salt-sensitive cultivars exhibiting apparent improvement. Salt stress was also observed to cause adverse physiological and biochemical changes in seedlings of both cultivars. Pre-soaking of seeds in Spd and GA₃ contributed to improved ion homeostasis as well as delaying loss of pigment accumulation. Moreover, Spd and GA₃ priming treatments reduced the production of H₂O₂, but enhanced the antioxidant systems in rice plants exposed to salt stress. Based on our results, Spd was more effective than GA₃ in improving salt tolerance in seedlings of both rice cultivars. It was noted that the promotive effects of Spd and GA₃ on seedling growth in this study were quantitatively small which may limit the practical application of these compounds. Nevertheless, the wide array of physiological parameters investigated, particularly the antioxidant capacities, provided a new set of useful data with respect to the beneficial actions of hormonal priming.

Acknowledgments

This research was supported by a grant from the Food and Functional Food Cluster under the National Research University Project of Thailand's Office of the Higher Education Commission and Salt-tolerant Rice Research Group, Khon Kaen University. The authors are grateful to Dr. Poramet Buntaung of the Faculty of Agriculture of Khon Kaen University for kindly providing the rice seed material used in this study.

References

- Abdel-Aal EM, Hucl PA (1999). Rapid method for quantifying total anthocyanins in blue aleurone and purple pericarp wheats. *Cereal Chem* 76:350-354.
- Afzal I, Butt A, Rehman HU, Basra SMA, Afzal A (2012). Alleviation of salt stress in fine aromatic rice by seed priming. *Aust J Crop Sci* 6(10): 1401-1407.
- Alcázar R, Altabella T, Marco F, Bortolotti C, Reymond M, Koncz C, Carrasco P, Tiburcio AF (2010). Polyamines: molecules with regulatory functions in plant abiotic stress tolerance. *Planta* 231:1237-1249.
- Ali HM, Siddiqui MH, Basalah MO, Al-Whaibi MH, Sakran AM, Al-Amri A (2012). Effects of gibberellic acid on growth and photosynthetic pigments of *Hibiscus sabdariffa* L. under salt stress. *Afr J Biotechnol* 11:800-804.
- Arnon DI (1949). Copper enzymes in isolated chloroplasts. Polyphenyloxidase in *Beta vulgaris*. *Plant Physiol* 24:1-15.
- Bates IS, Waldern RP, Teare ID (1973). Rapid determination of free proline for water stress studies. *Plant Soil* 39:205-207.
- Bradford KJ (1986). Manipulation of seed water relations via osmotic priming to improve germination under stress conditions. *Hort Sci* 21:1105-1111.
- Brand-Williams W, Cuvelier ME, Berset C (1995). Use of free radical method to evaluate antioxidant activity. *LWT - Food Sci Technol* 28:25-30.
- Benzie IFF, Strain JJ (1996). The ferric reducing ability of plasma (FRAP) as a measure of 'antioxidant power': The FRAP assay. *Anal Biochem* 239:70-76.
- Blokchina O, Virolainen E, Fagerstedt KV (2003). Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Ann Bot* 91:179-194.
- Blumwald E (2000). Sodium transport and salt tolerance in plants. *Curr Opin Cell Biol* 12:431-434.
- Bouchereau A, Aziz A, Larher F, Martin-Tanguy J (1999). Polyamines and environmental challenges: recent development. *Plant Sci* 140:103-125.
- Chalker-Scott L (1999). Environmental significance of anthocyanins in plant stress responses. *Photochem Photobiol* 70(1):1-9.
- Chattopadhyay MK, Tiwari BS, Chattopadhyay G, Bose A, Sengupta DN, Ghosh B (2002). Protective role of exogenous polyamines on salinity-stressed rice (*Oryza sativa*) plants. *Physiol Plant* 116:192-199.
- Choudhary SP, Oral HV, Bhardwaj R, Yu JQ, Tran LS (2012). Interaction of brassinosteroids and polyamines enhances copper stress tolerance in raphanus sativus. *J Exp Bot* 63:5659-5675.
- Daiponmak W, Theerakulpisut P, Thanonkao P, Vanavichit A, Prathepha P (2010). Changes of anthocyanin cyanidin-3-glucoside content and antioxidant activity in Thai rice varieties under salinity stress. *Sci Asia* 36:286-291.
- Das C, Sengupta T, Chattopadhyay S, Setua M, Das NK, Saratchandra B (2002). Involvement of kinetin and spermidine in controlling salinity stress in mulberry (*Morus alba* L. cv. S1). *Acta Physiol Plant* 24:53-57.
- Duan J, Li J, Guo S, Kang Y (2008). Exogenous spermidine affects polyamine metabolism in salinity-stressed *Cucumis sativus* roots and enhances short-term salinity tolerance. *J Plant Physiol* 165:1620-1635.
- Eryilmaz F (2006). The relationships between salt stress and anthocyanin content in higher plants. *Biotechnol Biotech Eq* 20(1):47-52.
- Farooq M, Basra SMA, Wahid A, Khaliq A, Kobayashi N (2010). Rice seed invigoration: a review, p. 137-175. In: *Organic farming, pest control and remediation of soil pollutants*. Springer, Netherlands.
- Farooq M, Basra SMA, Hussain M, Rehman H, Saleem BA (2007). Incorporation of polyamines in the priming media enhances the germination and early seedling growth in hybrid sunflower (*Helianthus annuus* L.). *Int J Agr Biol* 9:868-872.
- Farooq M, Wahid A, Lee D-J (2009). Exogenously applied polyamines increase drought tolerance of rice by improving leaf water status, photosynthesis and membrane properties. *Acta Physiol Plant* 31:937-945.
- Gonai T, Kawahara S, Tougou M, Satoh S, Hashiba T, Hirai N, Kawaide H, Kamaiya Y, Yoshioka T (2004). Abscisic acid in the thermoinhibition of lettuce seed germination and enhancement of its catabolism by gibberellin. *J Exp Bot* 55:111-118.
- Ghodrat V, Rousta MJ (2012). Effect of priming with gibberellic acid (GA₃) on germination and growth of Corn (*Zea mays* L.) under saline conditions. *Int J Agr Crop Sci* 4(13):882-885.
- Groppa MD, Benavides MP (2008). Polyamines and abiotic stress: recent advances. *Amino Acid* 34:35-45.
- Iqbal M, Ashraf M (2013). Gibberellic acid mediated induction of salt tolerance in wheat plants: Growth, ionic partitioning, photosynthesis, yield and hormonal homeostasis. *Environ Exp Bot* 86:76-85.
- Kachout SS, Hamza KJ, Karray N, Bouraoui NK, Leclerc JC, Ouergui Z (2013). Salt-induced changes in antioxidative enzyme activities in shoot tissues of two Atriplex varieties. *Not Bot Horti Agrobo* 41(1):115-121.
- Khajeh-Hosseini M, Powell AA, Bingham IJ (2003). The interaction between salinity stress and seed vigor during germination of soybean seeds. *Seed Sci Technol* 31:715-725.
- Khan HA, Ziaf K, Amjad M, Iqbal Q (2012). Exogenous application of polyamines improves germination and early seedling growth of hot pepper. *Chil J Agr Res* 72:429-433.
- Koornneef M, Bentsink L, Hilhorst H (2002). Seed dormancy and germination. *Curr Opin Plant Biol* 5:33-36.
- Kusano T, Berberich T, Tateda C, Takahashi Y (2008). Polyamines: essential factors for growth and survival. *Planta* 228:367-381.
- Li Z, Peng Y, Zhang X-Q, Ma X, Huang L-K, Yan Y-H (2014). Exogenous spermidine improves seed germination of white clover under water stress via involvement in starch metabolism, antioxidant defenses and relevant gene expression. *Molecules* 19:18003-18024.
- Magome H, Yamaguchi S, Hanada A, Kamiya Y, Oda K (2004).

- Dwarf* and *delayed-flowering 1*, a novel *Arabidopsis* mutant deficient in gibberellin biosynthesis because of overexpression of a putative AP2 transcription factor. *Plant J* 37:720-729.
- Munns R, James RA, Lauchli A (2006). Approaches to increasing the salt tolerance of wheat and other cereals. *J Exp Bot* 57:1025-1043.
- Parvaiz A, Satyawati S (2008). Salt stress and phyto-biochemical responses of plants – a review. *Plant Soil Environ* 54:89-99.
- Pedranzani H, Racagni G, Alemano S, Miersch O, Ramírez I, Peña-Cortés H, Taleisnik E, Machado-Domenech E, Abdala G (2003). Salt tolerant tomato plants show increased levels of jasmonic acid. *Plant Growth Regul* 41:149-158.
- Pokorny J, Yanishlieva N, Gordon M (2001). Antioxidants in food, practical applications. Woodhead Publishing Limited, Cambridge, 1-3 p.
- Quinet M, Ndayiragije A, Lefevre I, Lambillotte B, Dupont-Gillain CC, Lutts S (2010). Putrescine differently influences the effect of salt stress on polyamine metabolism and ethylene synthesis in rice cultivars differing in salt resistance. *J Exp Bot* 61:2719-2733.
- Rao PS, Mishra B, Gupta SR (2013a). Effect of salinity and alkalinity on grain quality of tolerant, semi-tolerant and sensitive rice genotypes. *Rice Sci* 20:284-291.
- Rao A, Ahmad SD, Sabir SM, Shahid Awan S, Shah AH, Khan MF, Khan SA, Shafique S, Arif S, Abbas SR, Gohar M (2013b). Antioxidant activity and lipid peroxidation of selected wheat cultivars under salt stress. *J Med Plants Res* 7(4):155-164.
- Radhakrishnan R, Lee IJ (2014). Effect of low dose of spermidine on physiological changes in salt-stressed cucumber plants. *Russian J Plant Physiol* 61:90-96.
- Radi AA, Farghaly FA, Hamada AM (2013). Physiological and biochemical responses of salt-tolerant and salt-sensitive wheat and bean cultivars to salinity. *J Biol Earth Sci* 3(1):72-73.
- Ratnakar A, Rai A (2014). Improvement of salt tolerance in *Trigonella foenum-graecum* L. var. PEB by plant growth regulators. *J Stress Physiol Biochem* 10(2):135-143.
- Razem FA, Baron K, Hill RD (2006). Turning on gibberellin and abscisic acid signaling. *Curr Opin Plant Biol* 9:454-459.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 26:1231-1237.
- Reddy MP, Vora AB (1986). Changes in pigment composition. Hill reaction activity and saccharides metabolism in bajra (*Penisetum typhoides* S & H) leaves under NaCl salinity. *Photosynthetica* 20:50-55.
- Roychoudhury A, Basu S, Sengupta DN (2011). Amelioration of salinity stress by exogenously applied spermidine or spermine in three varieties of indica rice differing in their level of salt tolerance. *J Plant Physiol* 168:317-328.
- Saleethong P, Sanitchon J, Kong-ngern K, Theerakulpisut P (2011). Pretreatment with spermidine reverses inhibitory effects of salt stress in two rice (*Oryza sativa* L.) cultivars differing in salinity tolerance. *Asian J Plant Sci* 10:245-254.
- Shabala S, Cui TA, Pottosin I (2007). Polyamines prevent NaCl-induced K⁺ efflux from pea mesophyll by blocking non-selective cation channels. *FEBS Lett* 581:1993-1999.
- Singleton VL, Orthofer R, Lamuela-Raventos RM (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin–Ciocalteu reagent. *Meth Enzymol* 299:152-178.
- Smirnov N, Cumbes QJ (1989). Hydroxyl radical scavenging activity of compatible solutes. *Phytochemistry* 28:1057-1060.
- Sultana B, Anwar F, Ashraf M (2009). Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. *Molecules* 14:2167-2180.
- Sun Q, Wang JH, Sun BQ (2007). Advances on seed vigor physiological and genetic mechanisms. *Agric Sci China* 6:1060-1066.
- Tang W, Newton RJ (2005). Polyamines reduce salt-induced oxidative damage by increasing the activities of antioxidant enzymes and decreasing lipid peroxidation in *Virginia pine*. *Plant Growth Regul* 46:31-43.
- Trivellini A, Gordillo B, Rodríguez-Pulido FJ, Borghesi E, Ferrante A, Vernieri P, Quijada-Morín N, González-Miret ML, Heredia FJ (2014). Effect of salt stress in the regulation of anthocyanins and color of *Hibiscus* flowers by digital image analysis. *J Agric Food Chem* 62(29):6966-6974.
- Valifarda M, Mohsenzadeh S, Kholdebarin B, Rowshan V (2014). Effects of salt stress on volatile compounds, total phenolic content and antioxidant activities of *Salvia mirzayanii*. *S Afr J Bot* 93:92-97.
- Velikova V, Yordanov I, Edreva A (2000). Oxidative stress and some antioxidant systems in acid rain-treated bean plants: protective role of exogenous polyamines. *Plant Sci* 151:59-66.
- Verma S, Mishra SN (2005). Putrescine alleviation of growth in salt stressed *Brassica juncea* by inducing antioxidative defense system. *J Plant Physiol* 162:669-677.
- Wahid A (2007). Physiological implications of metabolites biosynthesis in net assimilation and heat stress tolerance of sugarcane (*Saccharum officinarum*) sprouts. *J Plant Res* 120:219-228.
- Wahid A, Ghazanfar A (2006). Possible involvement of some secondary metabolites in salt tolerance of sugarcane. *J Plant Physiol* 163:723-730.
- Wen F-P, Zhang Z-H, Bai T, Xu Q, Pan Y-H (2010). Proteomics reveals the effects of gibberellic acid (GA₃) on salt-stressed rice (*Oryza sativa* L.) shoots. *Plant Sci* 178:170-175.
- Zhu H, Ding GH, Fang K, Zhao FG, Qin P (2006). New perspective on the mechanism of alleviating salt stress by spermidine in barley seedlings. *Plant Growth Regul* 49:147-156.