

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

ALMOST PERIODIC BEHAVIOR OF SOLUTIONS OF
A NONLINEAR VOLTERRA SYSTEM *

by

R. K. Miller

Center for Dynamical Systems
Division of Applied Mathematics
Brown University
Providence, Rhode Island

FACILITY FORM 602	N69-39012	
	45	1
	CR-106093	19

* This research was supported in part by the National Aeronautics and Space Administration under Grant No. NGL 40-002-015 and in part by the Air Force Office Scientific Research under Grant No. AF-AFOSR 67-0693A.

ALMOST PERIODIC BEHAVIOR OF SOLUTIONS OF
A NONLINEAR VOLTEIRA SYSTEM

R. K. Miller

1. Introduction

The purpose of this paper is to study the behavior as $t \rightarrow \infty$ of solutions of a system of two nonlinear equations of the form

$$(1.1) \quad \begin{aligned} x_1(t) &= f_1(t) - \int_0^t a_1(t-s)g_1(s, x_1(s))ds \\ &\quad - \int_0^t a_2(t,s)g_2(s, x_2(s))ds, \\ x_2(t) &= f_2(t) - \int_0^t a_2(t-s)g_1(s, x_1(s))ds \\ &\quad - \int_0^t a_1(t-s)g_2(s, x_2(s))ds \end{aligned}$$

where $f_1(t)$ and $f_2(t)$ are asymptotically almost periodic and both $g_1(t, x)$ and $g_2(t, x)$ are almost periodic in t uniformly for x on compact sets. We seek conditions which guarantee that the solutions $x_1(t)$ and $x_2(t)$ of (1.1) exist for all $t \geq 0$ and are asymptotically almost periodic.

System (1.1) arises in a natural way from the partial differential equation

$$(1.2) \quad u_t = u_{xx} \quad (t > 0, 0 < x < L)$$

with initial conditions

$$(1.3) \quad u(0, x) = F(x) \quad (0 < x < L)$$

and nonlinear boundary conditions of the form

$$(1.4) \quad u_x(t, 0) = g_1(t, u(t, 0)), \quad u_x(t, L) = -g_2(t, u(t, 0)),$$

for all $t > 0$. Indeed, if $A_1(t) = u_x(t, 0)$ and $A_2(t) = u_x(t, L)$ are assumed to be known functions and if $A_j \in C[0, \infty) \cap C^1(0, \infty)$ with $A_j(t)$ absolutely continuous in a neighborhood of $t = 0$, then well-known elementary methods imply that

$$(1.5) \quad u(t, x) = F_0/2 + \sum_{n=1}^{\infty} F_n \exp(-(n\pi/L)^2 t) \cos(n\pi x/L) \\ + L^{-1} \int_0^t \{1 + 2 \sum_{n=1}^{\infty} \exp\{-(n\pi/L)^2(t-s)\} \cos(n\pi x/L)\} A_1(s) ds \\ - L^{-1} \int_0^t \{1 + 2 \sum_{n=1}^{\infty} (-1)^n \exp\{-(n\pi/L)^2(t-s)\} \cos(n\pi x/L)\} A_2(s) ds$$

where

$$(1.6) \quad F_n = (2/L) \int_0^L F(x) \cos(n\pi x/L) dx \quad (n = 0, 1, 2, \dots)$$

is the sequence of Fourier cosine coefficients of F . Setting $x = 0$ and then $x = L$ in (1.5) and using (1.4) one obtains the

integral equations

$$(1.7a) \quad u(t,0) = F_0/2 + \sum_{n=1}^{\infty} F_n \exp \{-(n\pi/L)^2 t\} \\ -L^{-1} \int_0^t \{1+2 \sum_{n=1}^{\infty} \exp \{-(n\pi/L)^2 (t-s)\} g_1(s, u(s,0))\} ds \\ -L^{-1} \int_0^t \{1+2 \sum_{n=1}^{\infty} (-1)^n \exp \{-(n\pi/L)^2 (t-s)\} g_2(s, u(s,L))\} ds,$$

and

$$(1.7b) \quad u(t,L) = F_0/2 + \sum_{n=1}^{\infty} F_n (-1)^n \exp \{-(n\pi/L)^2 t\} \\ -L^{-1} \int_0^t \{1+2 \sum_{n=1}^{\infty} (-1)^n \exp \{-(n\pi/L)^2 (t-s)\} g_1(s, x_1(s))\} ds \\ -L^{-1} \int_0^t \{1+2 \sum_{n=1}^{\infty} \exp \{-(n\pi/L)^2 (t-s)\} g_2(s, x_2(s))\} ds.$$

Equations (1.7) clearly have the form (1.1) with $x_1(t) = u(t,0)$ and $x_2(t) = u(t,L)$. On the other hand if $u(t,0)$ and $u(t,L)$ are the known unique solutions of (1.7), then $u(t,x)$ may be obtained using (1.4) and then (1.5). This formal equivalence of (1.2-4) and (1.7) will be made precise in section II below.

Equations (1.2-4) and also our assumption of almost periodicity may be physically motivated using C. C. Lin's theory of superfluidity of helium, c.f. [1]. In three dimensional space with coordinates (x,y,z) let the planes $x = 0$ and $x = L$ represent two infinite plates. Suppose the region $0 < x < L$ between these

plates is filled with liquid helium initially at rest. If the boundary plates $x = 0$ and $x = L$ are both given sinusoidal oscillations in the y -direction, then a one-dimensional flow will be set up in the liquid. Let $u(t,x)$ be the velocity profile at time $t > 0$ of any point (x,y,z) with first coordinate x . Then $u(t,x)$ satisfies (1.2) and (1.3), $F(x) \equiv 0$. Lin's theory implies boundary conditions of the form (1.4), indeed

$$(1.8) \quad g_i(t,u) = B(u - C \sin(k_i t))^3$$

for $i = 1, 2$ where B and C are positive constants. For this problem we prove the following:

Theorem 1. Suppose $F \in C^2[0,L]$. Let $g_i(t,u)$ be given by (1.8) where $B > 0$, $C \neq 0$ and $k_i \neq 0$. Then (1.7) has unique continuous solutions $x_1(t) = u(t,0)$ and $x_2(t) = u(t,L)$ defined for all $t \geq 0$. Moreover, there exist two almost periodic functions $X_i(t)$ with Fourier series of the form

$$(1.9) \quad X_j(t) \sim \sum_{m,n=-\infty}^{\infty} X_{mn} \exp(i(mk_1 + nk_2)t)$$

such that

$$\lim \{x_i(t) - X_i(t)\} = 0 \quad \underline{\text{as}} \quad t \rightarrow \infty.$$

This result follows as a special case of more general theorems which will be proved below. These more general theorems concern a two-dimensional system of the form

$$(E) \quad x(t) = f(t) - \int_0^t A(t-s)G(s, x(s))ds,$$

where $A(t)$ is a matrix of the form

$$A(t) = \begin{pmatrix} a_1(t) & a_2(t) \\ a_2(t) & a_1(t) \end{pmatrix}.$$

In section 3 we use the special form of $A(t)$ to show that system (E) may be transformed into an equivalent system of the form

$$(E_N) \quad y(t) = \int_0^t R_N(t-s)\{y(s) - G_N(s, y(s))\}ds$$

where $R_N(t)$ is a positive definite, diagonal matrix of class $L^1(0, \infty)$. Subsequent work deals with equations of the form (E_N) rather than directly with (E).

Section 4 contains results concerning the global existence and boundedness of solutions of (E_N) . In section 5 we study the existence and uniqueness of almost periodic solutions of a related equation of the form

$$(1.10) \quad Y(t) = \int_{-\infty}^t R_N(t-s)(Y(s) - G_N(s, Y(s)))ds.$$

In the last section we show that the solutions $y(t)$ and $Y(t)$ of (E_N) and (1.10) are asymptotic, that is

$$\lim \{y(t) - Y(t)\} = 0 \quad \text{as} \quad t \rightarrow \infty.$$

Transforming (E_N) back to (E) then yields Theorem 1 as a corollary. Section 6 also contains results concerning the mean values of the solution $x(t)$ of (E) . This information on mean values is important in any study of the behavior of the nonlinear problem (1.2-4).

If $L = +\infty$ and if the second boundary condition in (1.4) is dropped, then (1.2-4) and (1.8) model the limiting case of a one-dimensional flow in a half space. This problem has been studied by Levinson [2]. Some of Levinson's results have been generalized in papers of Friedman [3, 4] and Miller [5]. A similar problem involving heat flow has been extensively studied by Mann and Wolf [6] and others [7, 8, 9]. The methods used in this paper are extensions of the methods used in [5]. The main tools in our analysis will be the "variation of constants" equation (E_N) and invariance results similar to those used in [5, section V].

2. Equivalence of the Problems

Let R^2 denote real Euclidean two-dimensional space of column vectors $x = \text{col}(x_1, x_2)$. Throughout the remainder of this paper the norm $|x|$ in R^2 will always mean

$$|x| = \max(|x_1|, |x_2|).$$

Many of our results are explicitly dependent on the use of this norm rather than some other equivalent norm.

Define

$$(2.1a) \quad f_1(t) = F_0/2 + \sum_{n=1}^{\infty} F_n \exp\{-(n\pi/L)^2 t\},$$

and

$$(2.1b) \quad f_2(t) = F_0/2 + \sum_{n=1}^{\infty} F_n (-1)^n \exp\{-(n\pi/L)^2 t\}$$

where F_n is defined by (1.6). Define

$$(2.2a) \quad a_1(t) = (1+2 \sum_{n=1}^{\infty} \exp\{-(n\pi/L)^2 t\})/L$$

$$(2.2b) \quad a_2(t) = (1+2 \sum_{n=1}^{\infty} (-1)^n \exp\{-(n\pi/L)^2 t\})/L$$

and

$$(2.3) \quad x(t) = \begin{pmatrix} x_1(t) \\ x_2(t) \end{pmatrix}, \quad f(t) = \begin{pmatrix} f_1(t) \\ f_2(t) \end{pmatrix}.$$

Let $A(t)$ be the matrix

$$(2.4) \quad A(t) = \begin{pmatrix} a_1(t) & a_2(t) \\ a_2(t) & a_1(t) \end{pmatrix}.$$

Then equation (1.7) has the form (E) where

$$(2.5) \quad G(t, x) = \begin{pmatrix} g_1(t, x_1) \\ g_2(t, x_2) \end{pmatrix}.$$

Theorem 2. Suppose $u(t, x)$ is a function which satisfies the following conditions:

- (i) $u(t, x)$ is continuous on $\{0 \leq t < \infty, 0 \leq x \leq L\}$.
- (ii) u_t and u_{xx} exist and are continuous for all
 (t, x) in the set $\{0 < t < \infty, 0 < x < L\}$.
- (iii) $u(t, x)$ satisfies (1.2), (1.3) and also (1.4) in the
sense that

$$\lim_{x \rightarrow 0^+} u_x(t, x) = g_1(t, u(t, 0))$$

and

$$\lim_{x \rightarrow L^-} u_x(t, x) = -g_2(t, u(t, L)). \quad (t > 0)$$

(iv) The functions $A_1(t) = g(t, u(t, 0))$ and $A_2(t) = -g_2(t, u(t, L))$ are of class $C[0, \infty) \cap C^1(0, \infty)$ and are absolutely continuous in a neighborhood of
 $t = 0$.

If $F \in C^2[0, L]$ and if $g_1, g_2 \in C^1$ for all (t, ∞) , then the functions

$$x_1(t) = u(t, 0), \quad x_2(t) = u(t, L)$$

satisfy (1.1) for all $t \geq 0$.

Proof. Define functions $\alpha(x) = x^2/2L$, $K(t, x) = \alpha(x)A_2(t) - \alpha(L-x)A_1(t)$ and $v(t, x) = u(t, x) - K(t, x)$. Then

$$\begin{aligned} v_t - v_{xx} &= Q(t, x) \equiv \alpha(x)A_2'(t) - \alpha(L-x)A_1'(t) \\ &\quad + \{A_2(t) - A_1(t)\}/L, \\ v_x(t, 0) &= v_x(t, L) = 0, \end{aligned}$$

and

$$v(0, x) = H(x) \equiv F(x) - \alpha(x)A_2(0) + \alpha(L-x)A_1(0).$$

The functions H and Q are sufficiently smooth in order to uniquely solve for $v(t,x)$ in the usual way, c.f. [10, Theorems 1 and 2]. Therefore

$$\begin{aligned}
 (2.6) \quad u(t,x) &= K(t,x) + v(t,x) \\
 &= K(t,x) + H_0/2 + \sum_{n=1}^{\infty} H_n \exp \{-(n\pi/L)^2 t\} \cos (n\pi x/L) \\
 &\quad + \int_0^t \int_0^L \{L^{-1} + \sum_{n=1}^{\infty} \exp \{-(n\pi/L)^2 (t-s)\} \cos (n\pi y/L) \\
 &\quad \quad \quad \cos (n\pi x/L)\} Q(s,y) dy ds.
 \end{aligned}$$

Here H_n is the sequence of Fourier cosine coefficients of H . By the definition of $\alpha(x)$ it follows that

$$\alpha(x) = L/6 + 2 \sum_{n=1}^{\infty} (-1)^n L^2 (n\pi)^{-2} \cos (n\pi x/L)$$

and

$$\alpha(L-x) = L/6 + 2 \sum_{n=1}^{\infty} L^2 (n\pi)^{-2} \cos (n\pi x/L)$$

when $0 < x < L$. Therefore, the definitions of K, Q and H together with integration by parts suffice to put the above expression for $u(t,x)$ into the form (1.5). Since $u(t,x)$ is continuous, then setting $x = 0$ and $x = L$ in (1.5) yields (1.7). Q.E.D.

Theorem 3. Suppose (2.1-5) are true, $F \in C^2[0,1]$ and the functions

$g_1(t,u)$ and $g_2(t,u)$ are of class C^1 for all (t,u) . If the solution $x(t)$ of equation (E) exists for all $t \geq 0$, then $u(t,0) = x_1(t)$ and $u(t,L) = x_2(t)$ are the boundary values of a function $u(t,x)$ which satisfies conditions (i) - (iv) of Theorem 2.

Proof. The conditions $F \in C^2[0,L]$ and (2.1) are sufficient to insure that $f \in C[0,\infty) \cap C^1(0,\infty)$ and that f' is locally of class L^1 on $0 \leq t < \infty$. Since g_1 and $g_2 \in C^1$, then it follows from results in [11] that $x_1(t)$ and $x_2(t)$ have these same smoothness properties, that is $x(t) \in C[0,\infty) \cap C^1(0,\infty)$ and $x'(t) \in L^1$ near $t = 0$.

Define $A_1(t) = g_1(t, x_1(t))$, $A_2(t) = -g_2(t, x_2(t))$ and define $u(t,x)$ by line (1.5). Condition (i) of Theorem 2 can easily be verified directly using (1.5). Since $A_1(t)$ and $A_2(t)$ are smooth, the steps in the proof of Theorem 2 can be reversed to obtain (2.6). Therefore, the results in [10] imply (ii), (1.2), (1.3) and the boundary conditions

$$\lim_{x \rightarrow 0} u_x(t,x) = A_1(t), \quad \lim_{x \rightarrow L} u_x(t,x) = A_2(t).$$

Setting $x = 0$ in (1.5) and using the present definitions of A_1 and A_2 it follows that

$$u(t,0) = f_1(t) - \int_0^t a_1(t-s)g_1(s, x_1(s))ds - \int_0^t a_2(t-s)g_2(s, x_2(s))ds.$$

(There is a similar formula for $u(t,L)$.) Comparing this with (1.1a) one sees that $u(t,0) = x_1(t)$ for all $t \geq 0$. Similarly $u(t,L) = x_2(t)$. Q.E.D.

3. Preliminary Transformations

Given any matrix $A(t)$ the resolvent $R(t)$ of $A(t)$ is defined to be the solution of the linear equation

$$(3.1) \quad R(t) = A(t) - \int_0^t A(t-s)R(s)ds.$$

If the entries of $A(t)$ are locally of class L^1 on $0 \leq t < \infty$ then it is known (c.f. [12, Chapter IV]) that $R(t)$ exists a.e., is locally L^1 on $0 \leq t < \infty$, and $R(t)$ also satisfies the equation

$$(3.1') \quad R(t) = A(t) - \int_0^t R(t-s)A(s)ds$$

a.e. on $0 < t < \infty$.

Let Q denote the symmetric, unitary matrix

$$(3.2) \quad Q = 2^{-1/2} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

Then clearly Q diagonalizes any matrix of the form (2.4), that is $QA(t)Q$ is diagonal.

Lemma 1. Suppose $A(t)$ is any matrix of the form (2.4) where $a_1(t)$ and $a_2(t)$ are locally L^1 on $0 \leq t < \infty$. For any $N > 0$ define

$$A_N(t) = NQA(t)Q$$

and let $R_N(t)$ be the resolvent of $A_N(t)$. Then the following statements are true:

- (i) $A_N(t) = N$ diagonal $(a_1(t) + a_2(t), a_1(t) - a_2(t))$.
- (ii) $R_N(t) =$ diagonal $(\lambda_{1N}(t), \lambda_{2N}(t))$.
- (iii) If $a_1(t)$ and $a_2(t)$ are defined by (2.2) then $\lambda_{1N}(t)$ and $\lambda_{2N}(t)$ are positive and continuous on $0 < t < \infty$ and

$$\int_0^{\infty} \lambda_{1N}(t) dt = 1, \quad \int_0^{\infty} \lambda_{2N}(t) dt < 1.$$

Proof. The first two parts follow immediately from (3.2) and equation (3.1) for the resolvent. Indeed, $\lambda_{1N}(t)$ is the resolvent of the scalar function

$$W_1(t) = N\{a_1(t) + a_2(t)\}$$

and $\lambda_{2N}(t)$ is the resolvent of the function

$$W_2(t) = N\{a_1(t) - a_2(t)\}.$$

If (2.2) is true, then

$$W_1(t) = N\{2 + 4 \sum_{n \text{ even}} \exp\{-(n\pi/L)^2 t\}\}$$

$$W_2(t) = N \left(4 \sum_{n \text{ odd}} \exp \{ -(n\pi/L)^2 t \} \right).$$

These formulas show that W_1 and W_2 are nonconstant, locally integrable, and completely monic on $0 < t < \infty$, that is

$$(-1)^j (W_k)^{(j)}(t) > 0 \text{ for } 0 < t < \infty, j = 0, 1, 2, \dots \text{ and } k = 1, 2.$$

It follows from a theorem of Reuter [13] that $\lambda_{kN}(t)$ is completely monic on $0 < t < \infty$. The results in section II of [5] immediately give the two integral estimates in (iii). Q.E.D.

Lemma 2. Suppose (E) satisfies (2.3-4), Q is defined by (3.2) and both $a_1(t)$ and $a_2(t)$ are functions which are locally L^1 on $0 \leq t < \infty$. For any fixed $N > 0$ let R_N be the resolvent of the matrix valued function $A_N(t) = NQA(t)Q$. Then the transformation

$$y = Q\{x-f(t)\} \quad (\text{or } x = Qy + f(t))$$

may be used to transform (E) into the equivalent system

$$(E_N) \quad y(t) = \int_0^t R_N(t-s) \{y(s) - G_N(s, y(s))\} ds$$

where

$$(3.3) \quad G_N(t, y) = QG(t, Qy+f(t))/N.$$

Proof. Define $\delta(t) = \text{diagonal } (\delta_d(t), \delta_d(t))$ where $\delta_d(t)$ is the

Dirac delta function. Let $*$ denote the convolution operation.

Then the resolvent equation

$$R_N(t) = A_N(t) - \int_0^t A_N(t-s)R_N(s)ds$$

may be written in the symbolic form

$$R_N = A_N - A_N * R_N$$

or, equivalently

$$(3.4) \quad (\delta - R_N) * (\delta + A_N) = \delta.$$

Equation (E) has the form $x = f - A * G(x)$. If $y = Q(x-f)$ then (E) becomes

$$\begin{aligned} y &= -(QA) * G(Qy+f) \\ &= -N(QAQ) * (QG(Qy+f)/N) \\ &= -A_N * G_N(y). \end{aligned}$$

Adding $A_N * y$ to both sides of this equation yields

$$y + A_N * y = (\delta + A_N) * y = A_N * \{y - G_N(y)\}.$$

Applying $\delta - R_N$ to both sides and using (3.4) one obtains

$$y = \delta * y = (\delta - R_N) * (\delta + A_N) * y = (\delta - R_N) * A_N * \{y - G_N(y)\}$$

or

$$y = R_N * \{y - G_N(y)\}.$$

This is equation (E_N) . The calculation is completely reversible so that (E_N) also implies (E) . Q.E.D.

4. Existence of Bounded Solutions

Assume the functions f, G and A of equation (E) satisfy the following conditions:

- (A1) f, A and G satisfy (2.2-4).
- (A2) $f \in C[0, \infty)$ and $f(t)$ is bounded on $[0, \infty)$.
- (A3) $G(t, x) \in C(\mathbb{R}^3)$ and $G(t, 0) \equiv 0$ for all $t \geq 0$.
- (A4) There exist positive numbers N and K such that if $|y| \leq K$ then $|y - G_N(t, y)| \leq K$ uniformly in $t \in \mathbb{R}^1$.

Note that more generally one could assume the existence of a vector valued function $r(t)$ such that $G(t, r(t)) \equiv 0$ for all $t \geq 0$. (This is the situation in Theorem 1 above.) However, the transformation $X = x - r(t)$ puts (E) in the form

$$X(t) = \{f(t) - r(t)\} - \int_0^t A(t-s)G(s, r(s) + X(s))ds.$$

If $r(t)$ is continuous and if $|f(t) - r(t)|$ is bounded, then the new equation satisfies (A3).

Theorem 4. Suppose (A1-4), (3.2) and (3.3) are all true. Then there exists a solution $x(t)$ of (E) such that $|x(t)| \leq K$ for all $t \geq 0$.

Proof. Let $C = C([0, \infty), \mathbb{R}^2)$ be the space of all continuous functions $\varphi: [0, \infty) \rightarrow \mathbb{R}^2$. Let C have the topology of uniform convergence on

compact subsets of the interval $0 \leq t < \infty$. Define

$$S = \{\varphi \in C: |\varphi(t)| \leq K \text{ for all } t \geq 0\}.$$

For any $\varphi \in S$ define

$$(M\varphi)(t) = \int_0^t R_N(t-s) \{\varphi(s) - G_N(s, \varphi(s))\} ds.$$

Clearly, $M: S \rightarrow C$ and M is completely continuous. Since the norm $|z| = |(z_1, z_2)|$ is defined by $|z| = \max\{|z_1|, |z_2|\}$, then (A4), Lemma 1 parts (ii) and (iii) and the definitions of S and M easily imply that $|(M\varphi)(t)| \leq K$ for all $t \geq 0$. This means that $M\varphi \in S$ if $\varphi \in S$. By the Schauder fixed point theorem the operator M has at least one fixed point $x(t)$. This fixed point solves (E_N) on $0 \leq t < \infty$ and thus also solves (E) . Q.E.D.

It can be shown that if G is defined by (1.8) then (A4) is true. More generally assume:

$$(A4') \quad G(t, x_1, x_2) = \text{col} (g(t, x_1), g(t, x_2)) \text{ for all } (t, x_1, x_2) \in R^3. \text{ Moreover, } g(t, z) \text{ is an odd, nondecreasing function of } z \text{ and is bounded in } t \in R^1 \text{ uniformly for } z \text{ on compact subsets of } R^1.$$

Lemma 3. Suppose G satisfies (A3) and (A4'). Let $B = \sup\{|f(t)|: t \geq 0\}$. Then for any $M > \sqrt{2}B$ and for any ϵ in the

range: $0 < \varepsilon < B$ there exists $N > 0$ such that (A_h) is true with
 $K = M + \varepsilon$.

Proof. Fix any such values of M and ε . Pick $N > 0$ so large
 that $2|g(t, z)| < \varepsilon$ uniformly in $t \geq 0$ and $|z| \leq 5M$. The map
 $w = u - G_N(t, u)$ may be written in the form

$$(4.1) \quad w_1 - u_1 = -\{g(t, (u_1+u_2)/\sqrt{2} + f_1(t)) \\
 + g(t, (u_1-u_2)/\sqrt{2} + f_2(t))\}/(\sqrt{2}N)$$

and

$$(4.2) \quad w_2 - u_2 = -\{g(t, (u_1+u_2)/\sqrt{2} + f_1(t)) - \\
 g(t, (u_1-u_2)/\sqrt{2} + f_2(t))\}/(\sqrt{2}N)$$

for all $t \geq 0$.

If $|u| = \max\{|u_1|, |u_2|\} \leq M$ then for any t one has

$$|(u_1+u_2)/\sqrt{2} + F_1(t)|, |(u_1-u_2)/\sqrt{2} + F_2(t)| \leq \sqrt{2}M + M \leq 5M.$$

Thus (4.1) and the choice of N imply

$$|w_1| < |u_1| + (\varepsilon/\sqrt{2} + \varepsilon/\sqrt{2})/\sqrt{2} \leq M + \varepsilon.$$

Similarly, $|w_2| < M + \varepsilon$.

Now consider the region

$$D = \{u \in \mathbb{R}^2: M < |u| \leq M + \varepsilon\}.$$

We must show that if $u \in D$ then $|w_1|, |w_2| < M + \varepsilon$. For $j = 0, 1, 2, \dots, 7$ define

$$S_j = \{(u_1, u_2) \in \mathbb{R}^2: u_1 + iu_2 = re^{i\theta} \text{ for some } r > 0 \text{ and} \\ \text{some } \theta \text{ in } (j-1)(\pi/2) \leq \theta \leq j\pi/2\}$$

and define $D_j = D \cap S_j$. Since $g(t, x)$ is an odd, nondecreasing function of x with $g(t, 0) = 0$, then the map $G(t, z_1, z_2) = \text{col}(g(t, x_1), g(t, x_2))$ maps each region S_j into itself. Also recall that

$$G_N(t, z) = QG(t, Qz + f(t))/N.$$

If $u \in D_1$, then (3.2) implies that $v = Qu \in D_1$. Since $B < M/\sqrt{2}$ and $|f(t)| \leq B$, then $y = v + f(t)$ lies in S_1, S_2 or S_8 . Again Q maps $S_1 \rightarrow S_1, S_2 \rightarrow S_8$ and $S_8 \rightarrow S_2$ so that

$$Z = Qx = QG(t, Qu + f(t))/N = G_N(t, u)$$

is in S_1, S_2 or S_8 . Finally, (4.1-2) show that

$$Z = G_N(t, u) = w - u.$$

If $Z = G_N(t, u)$ is in S_1 or S_2 , then the right hand sides of (4.1) and (4.2) both lie in the range $(-\epsilon, 0)$. Since $u \in D_1$, then $M < u_1 \leq M + \epsilon$ and $0 \leq u_2 \leq M + \epsilon$. Therefore,

$$w_1 = u_1 + z_1$$

lies in the range $0 < M - \epsilon < w_1 < M + \epsilon$ and

$$w_2 = u_1 + z_2$$

lies in the range $-\epsilon < w_2 < M + \epsilon$. Therefore, $|w| \leq M + \epsilon$.

Now suppose $u \in D_1$ and $z = G_N(t, u) \in S_8$. Since $|f(t)| \leq B < M/\sqrt{2}$ one must have $M < u_1 \leq M + \epsilon$ and $0 \leq u_2 \leq \sqrt{2}B < M$ in order that $Z \in S_8$. Therefore, the right hand side of (4.1) is in the range $(-\epsilon, 0)$ and the right hand side of (4.2) in the range $(0, \epsilon)$. This and $u_1 \in D_1$ mean that $M - \epsilon < w_1 < M + \epsilon$ and $\epsilon < w_2 \leq \sqrt{2}B + \epsilon < M + \epsilon$.

The analysis of the other seven regions S_2, S_3, \dots, S_8 follow in a similar manner. The various maps involved in the analysis are illustrated in Figure 1. Q.E.D.

(Printers: Put Figure 1 near here.)

Corollary 1. Suppose (A1-3), (A4'), (3.2) and (3.3) are true. If G is sufficiently smooth to insure the uniqueness of the solution x(t) of (E) then x(t) exists for all t ≥ 0 and satisfies

$$|Q\{x(t) - f(t)\}| \leq \sqrt{2}B \quad (0 \leq t < \infty)$$

where $B = \sup \{|f(t)| : 0 \leq t < \infty\}$.

Proof. By Lemma 3 and Theorem 4 above the solution $x(t)$ satisfies

$$\sup \{|Q\{x(t) - f(t)\}| : t \geq 0\} \leq M + \varepsilon$$

for each $\varepsilon > 0$ and each $M > \sqrt{2}B$. Q.E.D.

5. Almost Periodic Solutions

The purpose of this section is to study the existence and uniqueness of almost periodic solutions of equations of the form (1.10). First, we give appropriate definitions and background information concerning almost periodic functions. The first result in this section (Theorem 5) asserts that if $Y(t)$ is an almost periodic solution of (1.10) for some fixed $N_0 > 0$ then it is also a solution of (1.10) for all other $N > 0$. This result will be important since one value of N will be needed to prove existence of almost periodic solutions of (1.10) and a second value of N will be needed to obtain uniqueness and prove the asymptotic relationships between solutions of (E_N) and (1.10).

The rest of section is devoted to the existence and uniqueness of almost periodic solutions of (1.10). Lemma 4 is an invariance theorem for bounded solutions of (1.10). Lemma 5 asserts the uniqueness of bounded solutions of (1.10). The last result of the section asserts that the unique bounded solution of (1.10) is almost periodic.

Definition. A continuous function $S(t,x)$ defined for all $(t,x) \in \mathbb{R}^{n+1}$ is called almost periodic in t (uniformly for x on compact sets) if and only if given any sequence $\{t_n\}$ of real numbers there exists a subsequence $\{t_{nk}\}$ and a function $S^*(t,x)$ such that

$$\lim_{k \rightarrow \infty} S(t+t_{nk}, x) = S^*(t,x)$$

with convergence uniform in (t,x) for all $t \in R^1$ and x on compact subsets of R^n . In this case we write $S \in AP$.

The set of all functions S^* which may be obtained in this way is called the closed hull of S , written $CH(S)$.

As general references on almost periodic functions see the books of Favard [14] and Besicovitch [15] or the original papers of Bohr [16]. The results listed below are well-known results in this field.

Given a function $S(t,x)$ which is almost periodic in t uniformly for x on compact sets define $FM(S)$ to be the set of all A.P. functions $f(t)$ with range in the same space as S and satisfying the following condition:

If $\{t_n\}$ is any real sequence such that $\{S(t+t_n, x)\}$ is a Cauchy sequence uniformly in $t \in R^1$ and x on compact subsets of R^n , then $\{f(t+t_n)\}$ is a Cauchy sequence uniformly in $t \in R^1$.

The set $FM(S)$ is called the function module of S .

Given $S \in AP$ there exists a countable set of Fourier exponents $\{\lambda_n\} \subset R^1$ and a set $\{S_n(x)\}$ of continuous nontrivial functions such that S has Fourier series

$$S(t,x) \sim \sum_{n=1}^{\infty} S_n(x) \exp(i\lambda_n t).$$

If S is independent of x , then so are the functions S_n . The module of S , written $M(S)$, is the additive group of real numbers

generated by the sequence $\{\lambda_n\}$ of Fourier exponents. In other words $M(S)$ is the smallest additive subgroup of R^1 containing the set $\{\lambda_n\}$. An almost periodic function f is in the function module $FM(S)$ if and only if the Fourier exponents of f are contained in the module $M(S)$.

Let the functions f , A and G satisfy (A1-4) and in addition some or all of the following conditions:

(A5) There exist almost periodic functions $p(t)$ and $h_j(t,x)$ such that

$$\lim_{t \rightarrow \infty} \{f(t) - p(t)\} = 0, \quad \lim_{t \rightarrow \infty} \{g_j(t,x) - h_j(t,x)\} = 0$$

with the last limit uniform in x on compact sets of R^1 .

(A6) For each $t \in R^1$ and for $j = 1,2$ the function $h_j(t,x)$ is nondecreasing in x .

(A7) The functions $h_1(t,x)$ and $h_2(t,x)$ are locally Lipschitz continuous in x with Lipschitz constants independent of $t \in R^1$.

In Theorem 1 above $g_1(t,x) = g_2(t,x) = h_1(t,x) = h_2(t,x) = BX^3$. Moreover, (2.1) implies that

$$f_j(t) + C \sin(k_j(t)) \rightarrow F_0/2 + C \sin(k_j t)$$

as $t \rightarrow \infty$. Thus (A5-7) are all true for this special case. Note that (A5) implies that $G(t, x_1, x_2)$ has the special form (2.5).

Under the above assumptions the invariance theorem in [17, Theorem 1] implies that the equation

$$(E_N) \quad y(t) = \int_0^t R_N(t-s) \{y(s) - G_N(s, y(s))\} ds$$

has the limiting form

$$(5.1) \quad Y(t) = \int_{-\infty}^t R_N(t-s) \{Y(s) - H_N(s, Y(s))\} ds$$

where $H(t, y) = H(t, y_1, y_2) = \text{col} (h_1(t, y_1), h_2(t, y_2))$ and

$$(5.2) \quad H_N(t, y) = QH(t, Qy + p(t))/N.$$

Theorem 5. Suppose (A1-5), (3.2) and (5.2) are true. Suppose $Y(t)$ is any almost periodic solution of (5.1) for some fixed N . If $Y \in \text{FM}(H, p)$ then Y is also a solution of (5.1) for all other values of $N > 0$.

Proof. Pick any $M > 0$ with $M \neq N$. Let S_N be the resolvent of $NA(t)$ and S_M the resolvent of $MA(t)$. Write (5.1) in the form

$$(5.3) \quad Y(t) = h(t) + \int_0^t R_N(t-s) \{Y(s) - H_N(s, Y(s))\} ds$$

where

$$\begin{aligned} h(t) &= \int_{-\infty}^0 R_N(t-s) \{Y(s) - H_N(s, Y(s))\} ds \\ &= \int_t^{\infty} R_N(s) \{Y(t-s) - H_N(t-s, Y(t-s))\} ds \rightarrow 0 \end{aligned}$$

as $t \rightarrow \infty$. Let $\delta_d(t)$ be the Dirac delta function and let $\delta(t) =$ diagonal. $(\delta_d(t), \delta_d(t))$. If $*$ denotes convolution then (5.3) may be written in the form

$$Y = h + R_N * \{Y - QH(QY + p)/N\}.$$

Since $QR_NQ = S_N =$ resolvent of $NA(t)$, then the transformation $Z = QY$ puts the equation in the form

$$Z = Qh + S_N * \{Z - H(Z + p)/N\},$$

or

$$(\delta - S_N) * Z = Qh - S_N * H(Z + p)/N.$$

Applying $(\delta + NA)$ to both sides one obtains

$$\begin{aligned} Z &= (\delta + NA) * Qh - A * H(Z + p) \\ &= (\delta + NA) * Qh - (MA) * H(Z + p)/M. \end{aligned}$$

Add $(MA)*Z$ to both sides and apply $(\delta - S_M)$:

$$Z = (\delta - S_M)*(\delta + NA)*Qh + S_M*[Z - H(Z+p)/M].$$

Letting $Y = QZ$ one obtains

$$Y = Q(\delta - S_M)*(\delta + NA)*Qh + R_M*[Y - QH(Qy+p)/M].$$

Note that

$$\begin{aligned} Q(\delta - S_M)*(\delta + NA)Q &= Q\{\delta - S_M + NA - (N/M)(MA - S_M)\}Q \\ &= \delta + (1 - N/M)QS_MQ = \delta + (1 - N/M)R_M. \end{aligned}$$

Therefore

$$Y = h + (1 - N/M)R_M*h + R_M*[Y - H_M(Y)].$$

Writing this equation in the usual form one has

$$(5.4) \quad \begin{aligned} Y(t) &= h(t) + \int_0^t (1 - N/M)R_M(t-s)h(s)ds \\ &\quad + \int_0^t R_M(t-s)\{Y(s) - H_M(s, Y(s))\}ds \end{aligned}$$

for $t \geq 0$.

Let $t_n \rightarrow \infty$ be an increasing sequence such that $p(t+t_n) \rightarrow p(t)$ and $H(t+t_n, y) \rightarrow H(t, y)$ as $n \rightarrow \infty$. Since $Y \in FM(H, p)$, then

$Y(t+t_n) \rightarrow Y(t)$ as $n \rightarrow \infty$. Note that $h(t) \rightarrow 0$ as $t \rightarrow \infty$ and $R_N \in L^1(0, \infty)$ implies that $h(t) + (1-N/M)R_M^*h(t) \rightarrow 0$ as $t \rightarrow \infty$. Replacing t by $t + t_n$ in (5.4) yields

$$Y(t+t_n) = h(t+t_n) + (1-N/M)R_M^*h(t+t_n) + \int_{-t_n}^t R_M(t-s)\{Y(s+t_n) - H_M(s+t_n, Y(s+t_n))\}ds.$$

Taking the limit as $n \rightarrow \infty$ gives (5.1) with N replaced by M . Q.E.D.

We now turn to the existence-uniqueness problem. The following lemma will be needed.

Lemma 4. Suppose (5.2), (A1-3) and (A5) are true. Suppose (5.1) has a bounded solution $y(t)$ on $-\infty < t < \infty$. Then given any sequence $\{t_n\}$ of real numbers there exists a subsequence $\{t_{nk}\}$, a function $(H_N)^* \in CH(H_N)$ and a function $y^*(t)$ such that

$$y(t+t_{nk}) \rightarrow y^*(t), \quad H_N(t+t_{nk}, y) \rightarrow H_N^*(t, y)$$

and

$$(5.5) \quad y^*(t) = \int_{-\infty}^t R_N(t-s)\{y^*(s) - H_N^*(s, y^*(s))\}ds \quad (-\infty < t < \infty).$$

Proof. If $\{t_n\}$ contains a subsequence which tends to a finite limit point τ , then the result is trivial with $y^*(t) = y(t+\tau)$ and

$H_N^*(t, y) = H_N(t + \tau, y)$. Therefore, assume $t_n \rightarrow +\infty$. Since $p(t)$ and $H(t, y)$ are almost periodic in t , then there is a subsequence (which we shall also index by n) and functions $p^* \in CH(p)$, $H^* \in CH(H)$ such that $p(t + t_n) \rightarrow p^*(t)$ and $H(t + t_n, y) \rightarrow H^*(t, y)$. Then

$$\begin{aligned} H_N(t + t_n, y) &= QH(t + t_n, Qy + p(t + t_n)) / N \\ &\rightarrow QH^*(t, Qy + p^*(t)) / N = H_N^*(t, y). \end{aligned}$$

Since $y(t)$ is bounded and H is AP, then $|y(t) - H_N(t, y(t))|$ is bounded on $-\infty < t < \infty$. The convolution of a function in $L^1(-\infty, \infty)$ and a function of class $L^\infty(-\infty, \infty)$ results in a bounded uniformly continuous function. Since $y(t)$ solves (5.1), then $y(t)$ must be uniformly continuous. This in turn means that the sequence $\{y(t + t_n)\}$ is a uniformly bounded, equicontinuous family of functions on each finite subinterval of R^1 . By possibly taking a subsequence we may assume that $y(t + t_n) \rightarrow y^*(t)$ as $n \rightarrow \infty$ for some function y^* . Replacing t by $t + t_n$ in (5.1) one obtains

$$y(t + t_n) = \int_{-\infty}^t R_N(t-s) \{y(s + t_n) - H_N(s + t_n, y(s + t_n))\} ds$$

taking the limit as $n \rightarrow \infty$ gives (5.5). Q.E.D.

Lemma 5. Assume the hypotheses of Lemma 4. Assume (A6-7) are also

Thm. If $H > 0$ is sufficiently large, depending only on a bound for $y(t)$ and the Lipschitz constant of H , then $y(t)$ is the unique bounded solution of (5.1).

Proof. Suppose there exist two distinct solutions $y(t)$ and $z(t)$ of (5.1). Pick a sequence t_n such that

$$|y(t_n) - z(t_n)| \rightarrow L = \sup \{|y(t) - z(t)| : -\infty < t < \infty\}.$$

By possibly taking a subsequence we may assume that $y(t_n) - z(t_n) \rightarrow u_0$ as $n \rightarrow \infty$ where u_0 is some point on the boundary of the square $\{u : |u| \leq 1\}$. By possibly taking another subsequence Lemma 8 insures that $H_N(t+t_n, y) \rightarrow H_N^*(t, y)$, $y(t+t_n) \rightarrow y^*(t)$ and $z(t+t_n) \rightarrow z^*(t)$ where $H_N^* \in CH(H_N)$ and y^* and z^* solve (5.5). Clearly H_N^* satisfies the same hypotheses as H_N . Moreover, $u_0 = y^*(0) - z^*(0) = \lim \{y(t_n) - z(t_n)\}$ as $n \rightarrow \infty$. Thus we have reduced the problem to the case where $|y(0) - z(0)| = L = \sup \{|y(t) - z(t)| : -\infty < t < \infty\}$.

The two components of $b(t, y) = y - H_N(t, y)$ have the form

$$(5.7a) \quad b_1(t, y) = y_1 - \{h_1(t, (y_1+y_2)/\sqrt{2} + p_1(t)) + h_2(t, (y_1-y_2)/\sqrt{2} + p_2(t))\}/(\sqrt{2} N)$$

and

$$(5.7b) \quad b_1(t, y) = y_2 - \{h_1(t, (y_1 + y_2)/\sqrt{2} + p_1(t)) - h_2(t, (y_1 - y_2)/\sqrt{2} + p_2(t))\} / (\sqrt{2}N).$$

Set $u(t) = y(t) - z(t)$ and define

$$(5.8a) \quad m_1(t) = \{h_1(t, (y_1(t) + y_2(t))/\sqrt{2} + p_1(t)) - h_1(t, (z_1(t) + z_2(t))/\sqrt{2} + p_1(t))\} \{(u_1(t) + u_2(t))/\sqrt{2}\}^{-1}$$

if $u_1(t) + u_2(t) \neq 0$ and $m_1(t) = 0$ if $u_1(t) + u_2(t) = 0$. Since $u(t)$ is bounded and (A7) is true, then $m_1(t) \in L^\infty(-\infty, \infty)$. Moreover, $m_1(t) \geq 0$ by (A6). Similarly define

$$(5.8b) \quad m_2(t) = \{h_2(t, (y_1(t) - y_2(t))/\sqrt{2} + p_2(t)) - h_2(t, (z_1(t) - z_2(t))/\sqrt{2} + p_2(t))\} \{(u_1(t) - u_2(t))/\sqrt{2}\}^{-1}$$

if $u_1(t) \neq u_2(t)$ and $m_2(t) = 0$ otherwise. Since $u(t) = y(t) - z(t)$, then (5.1), (5.7) and (5.8) imply that

$$(5.9a) \quad u_1(t) = \int_{-\infty}^t \lambda_{1N}(t-s) \{[1 - (m_1(s) + m_2(s))/N]u_1(s) + [(m_2(s) - m_1(s))/N]u_2(s)\} ds$$

and

$$(5.9b) \quad u_2(t) = \int_{-\infty}^t \lambda_{2N}(t-s) \{[(m_2(s) - m_1(s))/N]u_1(s) + [1 - (m_1(s) + m_2(s))/N]u_2(s)\} ds$$

for all t in \mathbb{R}^1 . In system form (5.9) becomes

$$(5.9') \quad u(t) = \int_{-\infty}^t R_N(t-s)(I-M(s)/N)u(s)ds$$

where $M(s)$ is the appropriate matrix.

Pick $N > 0$ so large that $0 \leq m_1(t), m_2(t) \leq N/3$ a.e. on $-\infty < t < \infty$. For any fixed s the map $\hat{u} = (I-M(s)/N)u$ maps the square

$$S = \{u: |u_1|, |u_2| \leq L\}$$

into the region

$$S' = \{u: |u_1|, |u_2| \leq L \max \{1-2m_1(s)/N, 1-2m_2(s)\}\}.$$

Since $0 \leq 2m_1(s)/N \leq 2/3$ (by the choice of N), then $S' \subset S$.

Using the properties of λ_{2N} obtained in Lemma 2, (5.6) and (5.9b)

it follows that

$$|u_2(t)| \leq \int_{-\infty}^t \lambda_{2N}(t-s)Lds = L \int_0^{\infty} \lambda_{2N}(s)ds = L_0 < L.$$

Letting $t = 0$ we see that $|u_2(0)| < L$. Therefore, $u(t)$ is in the set

$$S_0 = \{u: |u_1| \leq L, |u_2| \leq L_0\}.$$

Since $\hat{u} = (I-M(s)/L)$ u maps S_0 strictly inside of the square S , say $|\hat{u}| \leq \beta L (0 < \beta < 1)$, then for any t line (5.9a) implies that

$$|u_1(t)| \leq \int_{-\infty}^t \lambda_{1N}(t-s)\beta L ds = \left(\int_0^{\infty} \lambda_{1N}(s) ds \right) (\beta L) = \beta L < L.$$

Therefore, $|u_1(0)| < L$ and $|u_2(0)| < L$ which contradicts $|u(0)| = \max \{|u_1(0)|, |u_2(0)|\} = L$. Q.E.D.

Theorem 6. Suppose the hypotheses of Lemma 4 are true. Then
 $y(t) \in FM(H,p)$ so that in particular $y \in AP$. Moreover, $y(t)$ solves
each equation of the (5.1) for all $N > 0$ and for N sufficiently
large is the unique bounded solution of (5.1).

Proof. Fix $N > 0$ and large. Let $\{t_n\}$ be any real sequence such that $\{p(t+t_n)\}$ and $\{H(t+t_n, y)\}$ are Cauchy sequences, $p(t+t_n) \rightarrow p^*(t) \in CH(p)$ and $H(t+t_n, y) \rightarrow H^*(t, y) \in CH(H)$. It must be shown that $\{y(t+t_n)\}$ is a Cauchy sequence uniformly in $t \in R^1$. Suppose this is not true. Then there exists $\epsilon > 0$, subsequences n_k and m_k , and a sequence τ_k such that $n_k > m_k \geq k$ and $|y(\tau_k + t_{n_k}) - y(\tau_k + t_{m_k})| \geq \epsilon$. Define

$$T_k = \tau_k + t_{n_k}, \quad S_k = \tau_k + t_{m_k} \quad (k = 1, 2, 3, \dots).$$

By possibly taking a subsequence of the k 's it follows that

$$H^*(t+\tau_k, y) \rightarrow H_0(t, y), \quad p^*(t+\tau_k) \rightarrow p_0(t)$$

for some functions $H_0 \in CH(H^*)$ and $p_0 \in CH(p^*)$. Then $p(t+T_k) = p(t+\tau_k+t_{nk}) \rightarrow p_0(t)$ and

$$H(t+T_k, y) = H(t+\tau_k+t_{nk}) \rightarrow H_0(t, y)$$

as $k \rightarrow \infty$. Similarly $p(t+S_k) \rightarrow p_0(t)$ and $H(t+S_k, y) \rightarrow H_0(t, y)$.

By Lemma 4 there exists a subsequence (which will again be indexed by k) and functions $y_1(t)$ and $y_2(t)$ such that $y(t+S_k) \rightarrow y_1(t)$, $y(t+T_k) \rightarrow y_2(t)$ and

$$y_j(t) = \int_{-\infty}^t R_N(t-s) \{y_j(s) - (H_0)_N(s, y_j(s))\} ds$$

for $j = 1, 2$ and $t \in R^1$. Since

$$|y_1(0) - y_2(0)| = \lim_{k \rightarrow \infty} |y(S_k) - y(T_k)| \geq \epsilon,$$

then $y_1(t) \neq y_2(t)$. But this violates the uniqueness asserted in Lemma 5. This contradiction shows that $y \in FM(H, p)$. Theorem 5 shows that y solves (5.1) for all $N > 0$. The uniqueness of y is Lemma 5. Q.E.D.

6. Proof of Theorem 1 and Generalizations.

Lemma 6. Suppose (A1-7) and (3.2) are true. Then for any $\eta > 0$ equation (5.1) has at least one bounded solution. In particular, then (5.1) has a solution $Y \in FM(H,p)$.

Proof. By Theorem 4 equation (E) has a bounded solution $x(t)$. Since (E) is equivalent to (E_N) for all $N > 0$, then each (E_N) has the same bounded solution. The results in [17] imply the existence of at least one bounded solution of (5.1) for any $N > 0$. Now apply Theorem 6. Q.E.D.

Theorem 7. Suppose (A1-7) and (3.2) are true. Then there exists a unique function $X \in FM(H,p)$ such that if $x(t)$ is any bounded solution of (E) then $x(t) - X(t) \rightarrow 0$ as $t \rightarrow \infty$.

Proof. Let $Y \in FM(H,p)$ be the function given by Lemma 6. Define $X(t) = QY(t) + p(t)$ and $y(t) = Q(x(t) - f(t))$. We must show that $x(t) - X(t) \rightarrow 0$ as $t \rightarrow \infty$ or equivalently that $y(t) - Y(t) \rightarrow 0$. If this is not true, then there exists an $\epsilon > 0$ and a sequence $t_n \rightarrow \infty$ such that $|y(t_n) - Y(t_n)| \geq \epsilon$. By Lemma 3 we may assume that $p(t+t_n) \rightarrow p^*(t)$, $H(t+t_n, y) \rightarrow H^*(t, y)$ and $Y(t+t_n) \rightarrow Y^*(t)$ as $n \rightarrow \infty$ where $p^* \in CH(p)$, $H^* \in CH(H)$ and Y^* solves (5.5). Write (E_N) in the form

$$y(t) = E(t) + \int_0^t R_N(t-s) \{y(s) - H_N(s, y(s))\} ds,$$

where

$$E(t) = \int_0^t R_N(t-s) \{H_N(s, y(s)) - G_N(s, y(s))\} ds.$$

Assumption (A5), (3.3) and (5.2) insure that $H_N(t, y(t)) - G_N(t, y(t)) \rightarrow 0$ as $t \rightarrow \infty$. Since $R_N \in L^1(0, \infty)$, then $E(t) \rightarrow 0$ as $t \rightarrow \infty$.

Therefore, Theorem 1 of [17] implies that by possibly taking a subsequence of t_n one has $y(t+t_n) \rightarrow y^*(t)$ where $y^*(t)$ solves (5.5) for the same value of N . By uniqueness of solutions of (5.5) for large N , $y^*(t) \equiv Y^*(t)$. On the other hand

$$|y^*(0) - Y^*(0)| = \lim_n |y(t_n) - Y(t_n)| \geq \epsilon > 0.$$

This contradiction proves the theorem. Q.E.D.

Theorem 8. Under the hypotheses of Theorem 7 the function $H(t, X(t))$ has mean value zero. Moreover, if the mean values of $p_1(t)$ and $p_2(t)$ are equal then the two components of $X(t)$ have equal mean values.

Proof. Recall that for any $\varphi(t) \in AP$ the mean value of φ is defined to be

$$m(\varphi) = \lim_{T \rightarrow \infty} T^{-1} \int_0^T \varphi(s) ds.$$

Since $Y(t) = Q(X(t) - p(t))$, then Y solves (5.1) on $-\infty < t < \infty$.

Taking mean values of both sides of (5.1) one obtains

$$(6.1) \quad m(Y) = R_N^*(0) (m(Y) - m(H_N)),$$

where $R_N^*(\omega)$ is the Fourier transform of R_N . By Lemma 1 above

$$R_N^*(0) = \text{diagonal } (1, N\beta(1+N\beta)^{-1})$$

where

$$\beta = \int_0^{\infty} \{a_1(t) - a_2(t)\} dt.$$

Write $Y(t) = \text{col } (Y_1(t), Y_2(t))$. Then the equation in the first component of (6.1) is

$$m(Y_1) = m(Y_1) - m(H_{N1}).$$

Therefore, $m(H_{N1}) = 0$, that is the first component of

$QH(t, QY(t) + p(t))/N = QH(t, X(t))/N$ has mean value zero. The second component of (6.1) is

$$m(Y_2) = N\beta(1+N\beta)^{-1} m(Y_2) - m(H_{N2}),$$

or

$$(6.2) \quad m(H_{N2}) = -N(1+N^2)^{-1} m(Y_2) \quad (0 < N < \infty).$$

Since the left hand side of (6.2) is independent of N , then (6.2) can be true for all $N > 0$ only if $m(Y_2) = m(H_{N2}) = 0$.

We have shown that the mean value of

$$\begin{aligned} H_N(t, Y(t)) &= QH(t, QY(t) + p(t))/N \\ &= QH(t, X(t))/N \end{aligned}$$

is the zero vector. Since Q is not singular and $N \neq 0$, then $H(t, X(t))$ also have mean value zero. We have also shown that

$$m(Y_2) = m(X_1 - X_2 - p_1 + p_2) / \sqrt{2} = 0,$$

that is $m(X_1 - X_2) = m(p_1 - p_2)$. Since $m(p_1 - p_2) = 0$, then $m(X_1) = m(X_2)$. Q.E.D.

Theorem 1 follows as a special case of the results in this section. In this special case $g(x) = Bx^3$ is smooth so that the solution of (E) is unique. Assumptions (A1-7) are easily verified with $p(t) = \text{col. } (p_1(t), p_2(t))$ having components $p_j(t) = F_0/2 + c \sin(k_j t)$. If k_1 and k_2 are linearly independent over the integers then $p(t)$ is quasiperiodic with fundamental frequencies (k_1, k_2) . If k_1 and k_2 are linearly dependent then there exist

integers M_1 and M_2 such that $k_3 = M_1 k_1 + M_2 k_2$ and $k_3/2\pi$ is the least common period of $p(t)$. In this case the functions X_j in (1.9) will have the form

$$X_j(t) = \sum_{n=-\infty}^{\infty} X_{jn} \exp \{in(M_1 k_1 + M_2 k_2)t\}.$$

If $u(t,x)$ is the function defined by (1.5) and X_j the functions defined by (1.9) then

$$(6.3a) \quad u(t,0) = X_1(t) + E_1(t)$$

and

$$(6.3b) \quad u(t,L) = X_2(t) + E_2(t)$$

where X_1 and X_2 both have the same mean value, $E_j(t) \in C[0,\infty) \cap C^1(0,\infty)$, $d/dt E_j(t)$ is L^1 near $t=0$ and $E_j(t) \rightarrow 0$ as $t \rightarrow \infty$.

Therefore, one would conjecture that

$$\lim_{t \rightarrow \infty} u(t,x) = U(t,x)$$

where U solves the problem

$$(6.4) \quad \begin{aligned} U_t &= U_{xx} & (-\infty < t < \infty, 0 < x < L) \\ U(t,0) &= X_1(t), \quad U(t,L) = X_2(t) & (-\infty < t < \infty). \end{aligned}$$

Similarly $u(t,x)$ satisfies boundary conditions of the form

$$(6.5a) \quad \partial u / \partial x(t, 0) = B\{X_1(t) - C \sin(k_1 t)\}^3 + E_1(t)$$

and

$$(6.5b) \quad \partial u / \partial x(t, L) = -B\{X_2(t) - C \sin(k_2 t)\}^3 + E_2(t),$$

for all $t \geq 0$. Here E_1 and E_2 have the same properties as the corresponding terms in (6.3) and the two functions

$B\{X_j(t) - C \sin(k_j t)\}^3$ have mean value zero. If it is true that $u(t, x)$ tends to a solution U of (6.4) then $U(t, x)$ should also satisfy the boundary conditions

$$\partial U / \partial x(t, 0) = B\{X_1(t) - C \sin(k_1 t)\}^3$$

and

$$\partial U / \partial x(t, L) = -B\{X_2(t) - C \sin(k_2 t)\}^3,$$

for $-\infty < t < \infty$.




FIGURE 1

References

1. C. C. Lin, Hydrodynamics of liquid helium II, Phys. Rev. Letters 2(1959), 245-246.
2. N. Levinson, A nonlinear Volterra equation arising in the theory of superfluidity, J. Math. Anal. Appl. 1(1960), 1-11.
3. A. Friedman, On integral equations of Volterra type, J. d'Analyse Math. 11(1963), 381-413.
4. _____, Periodic behavior of solutions of Volterra integral equations, J. d'Analyse Math. 15(1965), 287-303.
5. R. K. Miller, On Volterra integral equations with non-negative integrable resolvents, J. Math. Anal. Appl. 22(1968), 319-340.
6. W. R. Mann and F. Wolf, Heat transfer between solids and gases under nonlinear boundary conditions, Quart. Appl. Math. 9(1951), 163-184.
7. J. H. Roberts and W. R. Mann, On a certain nonlinear integral equation of the Volterra type, Pacific J. Math. 1(1951), 431-445.
8. K. Padmavally, On a nonlinear integral equation, J. Math. Mech. 7(1958), 533-555.
9. A. Friedman, Partial Differential Equations of Parabolic Type, Prentice-Hall, Englewood Cliffs, N. J., 1964.

10. S. Ito, Fundamental solutions of parabolic differential equations and boundary value problems, Jap. J. Math. 27(1957), 55-102.
11. R. K. Miller and A. Feldstein, Numerical solution of weakly singular integral equations, to appear.
12. R. K. Miller and J. A. Nohel, Nonlinear Volterra Integral Equations, W. A. Benjamin, to appear.
13. G. E. H. Reuter, Über eine Volterresche Integralgleichungen mit Totalmonotonem Kern, Arch. der Math. 7(1956), 59-66.
14. J. Favard, Lecons sur les Fonctions Presque-Periodiques, Gauthier-Villars, Paris, 1933.
15. A. S. Besicovitch, Almost Periodic Functions, Cambridge University Press, 1932.
16. H. Bohr, Zur Theorie der fastperiodischen Funktionen, I, II, III, Acta Math. 45(1924), 29-127; 46(1925), 101-237-281.
17. R. K. Miller, Asymptotic behavior of solutions of nonlinear Volterra equations, Bull. Amer. Math. Soc. 72(1966), 153-156.