

Viviane Gabriela Nascimento
Simon^I

José Maria Pacheco de Souza^{II}

Sonia Buongiorno de Souza^{III}

Breastfeeding, complementary feeding, overweight and obesity in pre-school children

ABSTRACT

OBJECTIVE: To analyze the association of overweight and obesity with breastfeeding and complementary feeding in pre-school children.

METHODS: Cross-sectional study with 566 children, enrolled in private schools of the city of São Paulo, Southeastern Brazil, in 2004-2005. The dependent variable was overweight and obesity. Body Mass Index percentile curves were employed to classify children's nutritional status, considering values $\geq P85$ and $< P95$ as overweight, and values $\geq P95$ as obesity. The following explanatory variables were analyzed: child and family socio-demographic characteristics; birth weight; parents' nutritional status; breastfeeding; complementary feeding; and current feeding. Analysis of association between explanatory variables and outcome was performed with simple logistic regression and multiple logistic regression with hierarchical model.

RESULTS: Prevalence of overweight and obesity in the population studied was 34.4%. The following were protective factors against overweight and obesity: exclusive breastfeeding for six months or more (95% CI [0.38;0.86]; OR=0.57; $p=0.02$) and breastfeeding for more than 24 months (95% CI [0.05;0.37]; OR=0.13; $p=0.00$).

CONCLUSIONS: Results suggest that breastfeeding can protect children against overweight and obesity, thus representing yet another advantage of maternal milk.

DESCRIPTORS: Child, Preschool. Breast Feeding. Feeding. Overweight. Obesity. Cross Sectional Studies.

^I Programa de Pós-Doutorado em Saúde Pública. Departamento de Saúde Materno-Infantil. Faculdade de Saúde Pública. Universidade de São Paulo. São Paulo, SP, Brasil

^{II} Departamento de Epidemiologia. Faculdade de Saúde Pública. Universidade de São Paulo. São Paulo, SP, Brasil

^{III} Departamento de Nutrição. Faculdade de Saúde Pública. Universidade de São Paulo. São Paulo, SP, Brasil

Correspondence:

Sonia Buongiorno de Souza
Faculdade de Saúde Pública
Universidade de São Paulo
Av. Dr. Arnaldo, 715
01246-904 São Paulo, SP, Brasil
E-mail: buonger@usp.br

Received: 10/9/2007

Revised: 5/29/2008

Approved: 7/2/2008

INTRODUCTION

The term "metabolic imprinting" describes a phenomenon by which an early nutritional experience, acting during a critical and specific developmental stage, can lead to permanent effects throughout an individual's life, predisposing them to certain diseases.²⁰

Breastfeeding represents one of the newborn's earliest nutritional experiences. Thus, maternal milk's unique composition could be involved with the metabolic imprinting process, changing adipocytes' number and/or size, or inducing the phenomenon of metabolic differentiation.¹

It is possible that breastfed infants develop mechanisms capable of regulating their energy intake and bottle feeding. These mechanisms could, for example, promote the development of overweight by stimulating an excessive milk intake and/or by hindering the development of self-regulatory mechanisms.⁵

The hypothesis that breastfeeding has a protective effect against obesity is not recent. However, controversial results have been found, and the issue continues to this day, especially because of the important increase in prevalence of obesity.²

In a cross-sectional study⁵ with 2,565 American children aged between three and five years, those who had been breastfed showed lower prevalence of "risk of overweight", compared to the others who had never been breastfed. However, the authors did not observe a protective effect against overweight, defined as Body Mass Index (BMI) equal to or above the 95 percentile.

In Brazil, there have been few studies that verified the relationship of breastfeeding to child overweight and obesity. As excess weight in children has become a relevant nutritional problem, this study aimed to analyze the association of overweight and obesity with breastfeeding and complementary feeding in pre-school children.

METHODS

A cross-sectional study with children, aged between two and six years, living in the city of São Paulo, Southeast Brazil, was performed in 2004-2005.

Sample was calculated using the EpiInfo 6.04 statistical package, Statcalc module, with the following parameters: level of significance of the α test=5%; power of test (1- β)=80%, odds ratio=2.2; 3:1 ratio between non-obese and obese individuals; % of exposed among non-obese individuals at 11%; resulting in 544 children.

Seven private schools, located in the district of Santana, in the city of São Paulo, were contacted by telephone to enable research feasibility and better performance. All the schools agreed to participate in the study. There were 809 children, aged between two and six years, enrolled in these schools.

A questionnaire to collect information about demographic and socioeconomic aspects of the child and its family, breastfeeding, complementary feeding and current feeding was designed. Data were collected between August and November 2004, and between March and May 2005. The questionnaire, with instructions to be self-applied, was distributed by the schools to be filled out by the mothers themselves or those responsible for the children and subsequently returned to the school. Of all the 809 formularies distributed, 30% were not returned. Thus, the sample studied was comprised of 566 children.

As regards breastfeeding, it was asked until what age, in months, the child was exclusively breastfed (exclusive breastfeeding) and breastfed (breastfeeding).

In terms of complementary feeding, it was asked at what age (in months) the following foods were introduced in the child's diet: water and/or tea, non-maternal milk, chocolate drink powders, sugar and/or honey, thickeners, fruits, leafy vegetables, cereals and tubers, beans, beef, chicken, fish, egg yolk, whole eggs, cold cuts, yogurt, cookies, and candies (drops and/or lollipops, chocolate and other processed foods). In regard to current feeding, the mother was asked to report what the child usually consumed, listing daily meal foods exclusively, without the need to quantify them. Each food mentioned was considered as a serving consumed by the child, according to the following food groups: bread and cereals; vegetables; fruits; legumes; meat, offal and eggs; milk and dairy; sugar and candies; oils and fats. To assess current feeding, the U.S. Department of Agriculture Center for Nutrition Policy and Promotion (USDA) Food Pyramid^a for two to six-year-old children was used as reference.

In addition to the questionnaire, anthropometric data were collected in the school itself, after consent was obtained from parents or those responsible, according to Lohman et al's⁸ method (1988). Child weight was measured with a Tanita 1632 Solar scale, and height measured with a Seca Bodymeter 208 stadiometer.

All formularies were analyzed; when information was incomplete or unclear, the mother or those responsible were contacted by telephone again to obtain this information, so that there were no losses at this stage of research.

The variable-response considered was child nutritional status. To classify it, BMI (weight [kg]/ height² [m]) percentile curves per age were used, in accordance with the National Center for Health Statistics (CDC, 2000):⁶ low weight values <5 percentile; normal weight values \geq 5 percentile and <85 percentile; overweight values \geq 85 percentile and <95 percentile; and obesity values \geq 95 percentile. For the statistical analysis, the variable-response was categorized into low weight + normal weight (<85 percentile) and overweight + obesity (\geq 85 percentile).

Explanatory variables were included, using a hierarchical approach on three levels: child age and sex, birth weight, and father's age and level of education at the distal level; age, mother's level of education and working conditions; number of siblings; and parents' nutritional status at the intermediate level; exclusive breastfeeding, breastfeeding, complementary feeding and current feeding of child at the proximal level. The Figure shows the complete hierarchical model.

Median values were considered to categorize family income and parents' age variables; and two groups were considered (\leq 4 years and >4 years) for child age, due to possible differences in feeding patterns.

^a Department of Agriculture Center for Nutrition Policy and Promotion. Food guide pyramid for young children. 1999 [citado 2007 fev 14]. Disponível em: URL:<http://www.cnpp.usda.gov>

In regard to child complementary feeding, foods that could be associated with child nutritional status were analyzed. Water and/or tea, fruits and non-maternal milk were considered as foods responsible for the interruption of exclusive breastfeeding. Early introduction of sugar and others (chocolate drink powder, sugar and/or honey, thickeners, yogurt, cookies and candies) can lead to obesity. Cold cuts were included in the analysis as they are a source of fat.

Food groups that are more related to overweight and obesity were selected for child current feeding, when consumed in great amounts: bread and cereals; milk and dairy; oils and fats; sugar and candies; and meat, offal and eggs.

Analysis of association of exclusive breastfeeding, breastfeeding, complementary feeding and current feeding with current child nutritional status was performed with simple logistic regression and multiple logistic regression.


The value of $p < 0.25$, for inclusion of variables in the multiple logistic regression analysis, was used in the simple or bivariate analyses, thus enabling a greater number of variables in the model. According to the hierarchical proposal, once included in the model, variables remained until the end, regardless of p -value

in the subsequent stages. In the final model, associations where variables showed $p < 0.10$ were considered to be of greater interest.

The reference category used was always the first category of each variable, except for the family income variable, whose reference category was $\leq R\$6,450.00$. This is because the first category was “family income not informed”, included in the analysis so that there were no losses of sample size.

Two multiple logistic regression models with the same variable-response (overweight + obesity) and same independent variables were performed. The main independent variable included in one of the models was exclusive breastfeeding, and in the other model, breastfeeding. Possible interactions between variables were tested with an α significance level=5%.

Statistical analysis was performed using the Stata statistical package, version 9.2. This study was approved by the *Comitê de Ética em Pesquisa* of *Faculdade de Saúde Pública* (School of Public Health Research Ethics Committee) at the Universidade de São Paulo. Informed consent forms were distributed by the school, along with research formularies. These were subsequently returned to the school itself, duly signed by the mother or those responsible.



* Distal level
 ** Intermediate level
 *** Proximal level

Figure. Hierarchical model for explanatory variables

RESULTS

Median exclusive breastfeeding was four months and median breastfeeding was seven months. In regard to complementary feeding, early introduction of all foods researched, among children aged between zero and six months of age, was found. For the majority of children (72.1%), water and/or tea was the earliest food introduced, followed by fruits (66.4%) and non-maternal milk (53.2%). Other foods were also introduced in this same period, such as: sugar and/or honey (15.2%); thickeners (10.2%); chocolate drink powders (3.0%); yogurts (4.6%); cereals and tubers (25.6%); beef, chicken or fish (54.1%); eggs (14.1%); leafy vegetables (40.3%) and beans (12.0%).

Fruit consumption in the current feeding was in accordance with recommendations. Children consumed a lower number of servings than what is recommended for the bread and cereals and vegetables groups, and almost 50% more than what is recommended for the milk and dairy group.

As regards child nutritional status, 5.3% were found to be low weight; 60.2% were normal weight; 17.8% were overweight and 16.6% were obese, resulting in a prevalence of 34.4% for overweight and obesity.

Table 1 shows the result of bivariate analysis.

Variables with an association of $p < 0.25$ in the bivariate analysis were selected for multiple analysis (Table 2): distal level – child age, birth weight, father's age and family income; intermediate level – mother's working conditions, number of siblings, parents' nutritional status; and proximal level – exclusive breastfeeding, breastfeeding, introduction of cold cuts, introduction of sugar and others, and number of servings of the meat, offal and eggs group in current feeding. There was no interaction between type of breastfeeding and child age ($p = 0.47$).

Tables 3 and 4 show results of multiple analysis for both models: exclusive breastfeeding model and breastfeeding model.

Hierarchical multiple analysis of the exclusive breastfeeding model shows that exclusive breastfeeding was a protective factor against overweight and obesity. Risk factors were the following: age > 4 years; birth weight $> 3.500\text{g}$; early introduction of sugar and others in the diet; and having an obese father.

Hierarchical multiple analysis of breastfeeding model shows that the longer breastfeeding lasts, the better protected against overweight and obesity the child is. In this model, risk factors were the following: age > 4 years; birth weight $> 3.500\text{g}$; mother working out of the home; paternal nutritional status.

DISCUSSION

Prevalence of excess weight found was 34.4% and could be considered high for the age group of children studied.

Sample was obtained from private schools in the city of São Paulo, based on the hypothesis that these children would have better socioeconomic level, raising the possibility of finding higher prevalence of overweight and obesity, and thus obtaining an adequate sample size for statistical analysis. According to the literature, overweight and obesity are more prevalent in children with higher socioeconomic level. As early as 1986, Campino⁴ pointed to income as an important socioeconomic factor to determine nutritional status. More recently, Saldiva et al¹⁴ (2004) found that the higher the per capita income, the higher the prevalence of obesity in children younger than five years of age, living in five cities of the state of São Paulo.

Leão et al⁷ (2003) studied 387 children aged between five and ten years, in public and private schools of the city of Salvador, state of Bahia, using the BMI to determine nutritional status. These authors found prevalence of 30% for obesity in private schools, and only 8% in public schools, showing that obesity is indeed more present in the population with higher socioeconomic level.

One possible limitation to this study is the occurrence of memory bias, when collecting information about duration of exclusive breastfeeding and age of introduction of complementary feeding, as this is obtained in a retrospective manner.

Median duration of exclusive breastfeeding (four months) was high, if compared to other studies,^{9,15} and so was the prevalence of exclusive breastfeeding in children older than six months (21.4%), a period when the child should already be given other foods.

Even though exclusive breastfeeding until the age of six months is admittedly important, complementary feeding continues to be introduced early on, as observed by this study and others.^{3,11,13,17,19}

Final models of analyses showed that exclusive breastfeeding for six months or more and breastfeeding continuing for more than 24 months of age are protective factors against overweight and obesity.

As regards breastfeeding, when both variables (exclusive breastfeeding and breastfeeding) were included in the model, the relationship between explanatory variables and the response-variable decreased. To avoid this "weakening", the choice was to perform modeling with each of them separately to better understand the phenomenon.

The protective relationship of exclusive breastfeeding to overweight and obesity remained in all stages of analysis, regardless of variables that entered it and

Table 1. Prevalence of overweight and obesity, OR and p-values, according to sociodemographic variables. São Paulo, Southeast Brazil, 2004-2005.

Variable	n	Overweight and obesity (%)	OR	p	p*
Sex					
Male	285	34.4	1	-	
Female	281	34.5	1.00	0.97	
Child age (in years)					
≤4	219	27.8	1	-	
>4	347	38.6	1.63	0.01	
Birth weight					
<2,500 g	62	25.8	1	-	
2,500 — 3,500g	380	32.9	1.41	0.28	0.01
≥3,500 g	124	43.5	2.22	0.02	
Father's age (in years)					
≤38	299	31.8	1	-	
>38	267	37.4	1.28	0.16	
Father's level of education					
Elementary school or high-school	110	39.1	1	-	
University	456	33.3	0.78	0.25	
Family income					
Not informed	107	44.9	-	-	
≤R\$ 6,450.00	230	28.3	1	-	
>R\$ 6,450.00	229	35.8	1.42	0.08	
Mother's age (in years)					
≤35	299	35.1	1	-	
>35	267	33.7	0.94	0.72	
Mother's level of education					
Elementary school or high-school	87	35.6	1	-	
University	479	34.2	0.94	0.80	
Mother's working condition					
Does not work	150	29.3	1	-	
Works	416	36.3	1.37	0.12	
Number of siblings					
0	368	36.1	1	-	
1 or more	198	31.3	0.80	0.24	
Maternal nutritional status					
Low or normal weight	417	32.6	1	-	
Pre-obesity	123	39.0	1.32	0.19	0.12
Obesity – levels I, II and III	26	42.3	1.51	0.31	
Paternal nutritional status					
Low or normal weight	149	29.5	1	-	
Pre-obesity	307	32.9	1.17	0.47	0.01
Obesity – levels I, II and III	110	45.4	1.99	0.01	

* linear trend p

could possibly interfere with this relationship. In regard to breastfeeding protection, for longer duration (more than 24 months), it is possible to raise the hypothesis

that the greater the amount of maternal milk an infant is fed in the beginning of life, the greater the protection in relation to overweight and obesity.

Until now, there have been few studies on the relationship of current overweight and obesity to history of breastfeeding, in Brazil. Methodological differences among studies hinder comparison between them, in a way.

By analyzing the association between exposure to breastfeeding in childhood and obesity in school-aged children from Brazilian families of high socioeconomic level, Siqueira & Monteiro¹⁶ (2007) observed that risk of obesity in children who had never been breastfed were two times higher than the risk in other children.

Dose-response effect in the association between duration of breastfeeding and obesity in school-aged children was not observed.

Balaban et al¹ (2004) performed a cross-sectional study with 409 children from daycare centers of the city of Recife government to investigate if breastfeeding had a protective effect against overweight in pre-school children (two to six years). Authors considered BMI ≥ 85 percentile as overweight. It was concluded that children who had been breastfed for a period shorter

Table 2. Prevalence of overweight and obesity, OR and p-values, according to breastfeeding, complementary feeding and current feeding. São Paulo, Southeast Brazil, 2004-2005.

Variable	n	Overweight and obesity (%)	OR	p	p*
Exclusive breastfeeding (in months)					
<6	445	36.4	1	-	
≥ 6	121	27.3	0.65	0.06	
Breastfeeding (in months)					
0 — 6	191	35.6	1	-	
6 — 12	239	35.6	1.04	0.88	
12 — 18	84	39.3	1.18	0.64	
18 — 24	21	28.6	0.62	0.41	
≥ 24	31	9.7	0.16	0.02	
Complementary feeding					
Water and/or tea were introduced (in months)					
0 — 6	408	35.3	1	-	
≥ 6	158	32.3	0.87	0.50	
Age when fruits were introduced (in months)					
0 — 6	376	35.1	1	-	
≥ 6	190	33.2	0.92	0.64	
Age when non-maternal milk was introduced (in months)					
0 — 6	301	32.9	1	-	
6 — 12	197	36.5	1.17	0.40	
12 — 18	51	37.2	1.21	0.54	
≥ 18	17	29.4	0.85	0.77	
Age when cold cuts were introduced (in months)					
0 — 12	21	23.8	1	-	
12 — 18	135	36.3	1.82	0.27	
18 — 24	78	25.6	1.10	0.86	
≥ 24	332	36.4	1.83	0.24	
Age when sugar and others were introduced (in months)					
0 — 6	144	31.9	1	-	
6 — 12	322	33.5	1.07	0.73	0.17
≥ 12	100	41.0	1.48	0.15	
Current feeding					
Meat, offal and eggs (servings)					
0	11	45.4	1	-	
1	94	36.2	0.81	0.24	
2	442	34.4			
3	19	21.1			

* linear trend p

Table 3. Multiple logistic regression analysis with hierarchical model for exclusive breastfeeding and overweight and obesity. São Paulo, Southeast Brazil, 2004-2005.

Variable	OR	p	90% CI
Child age (in years)			
≤4	1	-	-
>4	1.65	0.00	1.20;2.27
Birth weight			
<2500g	1	-	-
2500 — 3500g	1.36	0.34	0.80;2.30
≥3500g	2.12	0.03	1.18;3.80
Family income			
≤R\$ 6.450,00	1	-	-
>R\$ 6.450,00	1.36	0.15	0.96;1.93
Mother works out of the home			
No	1	-	-
Yes	1.38	0.15	0.95;1.99
Duration of exclusive breastfeeding (in months)			
<6	1	-	-
≥6	0.57	0.02	0.38;0.86
Age when sugar and others were introduced (in months)			
0 — 6	1	-	-
6 — 12	1.24	0.34	0.85;1.81
≥12	1.65	0.08	1.03;2.67
Paternal nutritional status			
Low or normal weight	1	-	-
Pre-obesity	1.20	0.40	0.83;1.74
Obesity – levels I, II and III	1.99	0.01	1.27;3.11

than four months showed greater prevalence of overweight than those who had been breastfed for four months or more.

Study by Victora et al¹⁸ (2003) showed that prevalence of obesity in male adolescents aged 18 years was three times lower among those who had been breastfed from three to five months of age, compared to the remaining categories.

An international systematic review¹² showed the protective effect of breastfeeding against overweight and obesity. Despite methodological differences among studies, the importance of breastfeeding to prevent overweight and obesity during childhood and adolescence was evident.

In the present study, among the remaining explanatory variables, child age over four years was a risk factor, perhaps because, at this age, they already choose and ask for foods they want to eat, giving preference to non-nutritious, high-energy foods. As the children studied belonged to families of high socioeconomic level, they would probably have more access to the media, which stimulates consumption of foods such as

cookies, candies, and salty snack foods. Another factor associated with risk at this age could be sedentarism, resulting from the daily hours spent on television and video-games, thus decreasing physical activity.

Birth weight >3500g was found to be a risk factor to develop overweight and obesity. Monteiro et al¹⁰ (2003), while studying infants at birth, at 20 months and 40 months, and at 14 and 16 years of age, concluded that birth weight has a positive linear association with overweight.

Consumption of sugar and other foods that contain sugar or need to be prepared with sugar only seems to be risk factors in the exclusive breastfeeding model and for children with introduction of foods at 12 months or older. Introduction of sugar in the diet at this age, when children are developing their food habits, can lead to greater consumption of these foods. Preparations such as milk with chocolate drink powder, porridge and cookies, which have an appealing taste, please children, leading to consumption in great amounts, thus increasing the diet's total energy intake and also the risk of overweight and obesity.

Table 4. Multiple logistic regression analysis with hierarchical model for breastfeeding and overweight and obesity. São Paulo, Southeast Brazil, 2004-2005.

Variable	OR	p	IC 90%
Child age (in years)			
≤4	1	-	-
>4	1,71	0,01	1,24;2,37
Birth weight			
<2500g	1	-	-
2500 — 3500g	1,42	0,28	0,83;2,42
≥3500g	2,29	0,02	1,27;4,14
Family income			
≤R\$ 6.450,00	1	-	-
>R\$ 6.450,00	1,39	0,13	0,97;1,98
Mother works out of the home			
No	1	-	-
Yes	1,47	0,09	1,01;2,14
Duration of breastfeeding (in months)			
0 — 6	1	-	-
6 — 12	0,80	0,29	0,56;1,14
12 — 18	1,07	0,80	0,67;1,72
18 — 24	0,48	0,17	0,20;1,16
≥24	0,13	0,00	0,05;0,37
Age when sugar and others were introduced (in months)			
0 — 6	1	-	-
6 — 12	1,15	0,53	0,79;1,68
≥12	1,50	0,16	0,93;2,41
Paternal nutritional status			
Low or normal weight	1	-	-
Pre-obesity	1,21	0,41	0,83;1,75
Obesity – levels I, II and III	2,08	0,00	1,32;3,26

As children whose mothers do not work seem to be better protected, mother's work becomes a possible risk factor. Mothers who work out of the home all day long probably leave their children in school full-time, where they have their meals. When leaving school at the end of the day, these children have already had dinner, unlike their parents. This can lead them to eat again with the family at home, thus increasing their diet's total energy intake. Another factor is that the mother who works tends to please the child with high-energy foods, such as cookies, chocolate, salty snack foods, drops and lollipops, in an attempt to make up for her absence due to work.

As regards the association of child overweight and obesity with paternal obesity, it could be supposed

that, in addition to the genetic factor, these fathers are not concerned about healthy eating and maintaining a normal weight. This image is passed onto the children, apart from usually eating fast-food meals or offering non-nutritious, high-energy foods to them. There were no studies in the literature that analyzed the association between child nutritional status and paternal nutritional status, thus making the comparison impossible.

In conclusion, in addition to all the breastfeeding benefits that have already been studied and acknowledged, protection against overweight and obesity during childhood becomes yet another benefit, regardless of child age, family income, nutritional status and parents' level of education.

REFERENCES

1. Balaban G, Silva GAP, Dias MLCM, Dias MCM, Fortaleza GTM, Morotó FMM, et al. O aleitamento materno previne o sobrepeso na infância? *Rev Bras Saude Matern Infant*. 2004;4(3):263-8. DOI: 10.1590/S1519-38292004000300006
2. Balaban G, Silva GAP. Efeito protetor do leite materno contra a obesidade infantil. *J Pediatr (Rio J)*. 2004;80(1):7-16. DOI: 10.1590/S0021-75572004000100004
3. Bruncker GS, Silva SM, França GV, Escuder MM, Venâncio SI. Fatores associados à interrupção precoce do aleitamento materno exclusivo e à introdução tardia da alimentação complementar no centro-oeste brasileiro. *J Pediatr (Rio J)*. 2006;82(6):445-51. DOI: 10.2223/JPED.1569
4. Campino AAC. Aspectos socioeconômicos da desnutrição no Brasil. *Rev Saude Publica*. 1986;20(1):83-101. DOI: 10.1590/S0034-89101986000100007
5. Hediger ML, Overpeck MD, Kuczmarski RJ, Ruan J. Association between infant breastfeeding and overweight in young children. *JAMA*. 2001;285(19):2453-60. DOI: 10.1001/jama.285.19.2453
6. Kuczmarski RJ, Ogden CL, Grummer SLM, et al. CDC Growth charts: United States – methods and development. Hyattsville, MD: National Center for Health Statistics; 2000.
7. Leão LSCS, Araújo LMB, Moraes LTLP, Assis AM. Prevalência de obesidade em escolares de Salvador, Bahia. *Arq Bras Endocrinol Metab*. 2003;47(2):151-7.
8. Lohman TG, Roche AF, Matorell R. Antropometric standardization reference manual. Illinois: Human Kinetics; 1988.
9. Mascarenhas MLW, Albernaz EP, Silva MB, Silveira RB. Prevalência de aleitamento materno exclusivo nos 3 primeiros meses de vida e seus determinantes no Sul do Brasil. *J Pediatr (Rio J)*. 2006;82(4):289-94. DOI: 10.2223/JPED.1506
10. Monteiro PO, Victora CG, Barros FC, Monteiro LM. Birth size, early childhood growth, and adolescent obesity in a Brazilian birth cohort. *Int J Obes Relat Metab Disord*. 2003;27(10):1274-82. DOI: 10.1038/sj.ijo.0802409
11. Oliveira LPM, Assis AMO, Pinheiro SMC, Prado MS, Barreto ML. Alimentação complementar nos primeiros dois anos de vida. *Rev Nutr*. 2005;18(4):459-69. DOI: 10.1590/S1415-52732005000400002
12. S Arenz, Ruckerl R, Koletzko B, Von Kries R. Breast-feeding and childhood obesity- a systematic review *Int J Obes Relat Metab Disord*. 2004;28(10):1247-56. DOI: 10.1038/sj.ijo.0802758
13. Saldiva SRDM, Escuder MM, Mondini L, Levy RB, Venâncio SI. Práticas alimentares de crianças de 6 a 12 meses e fatores maternos associados. *J Pediatr (Rio J)*. 2007;83(1):53-8. DOI: 10.2223/JPED.1588
14. Saldiva SRDM, Escuder MML, Venâncio SI, Benicio D' Aquino MH. Prevalence of obesity in preschool children from five towns in São Paulo state, Brazil. *Cad Saude Publica*. 2004;20(6):1627-32. DOI: 10.1590/S0102-311X2004000600021
15. Silveira FJF, Lamounier JA. Prevalência do aleitamento materno e práticas de alimentação complementar em crianças com até 24 meses de idade na região do Alto Jequitinhonha, Minas Gerais. *Rev Nutr*. 2004;17(4):437-47. DOI: 10.1590/S1415-52732004000400004
16. Siqueira RS, Monteiro CA. Amamentação na infância e obesidade na idade escolar em famílias de alto nível socioeconômico. *Rev Saude Publica*. 2007;41(1):5-12. DOI: 10.1590/S0034-89102007000100002
17. Simon VGN, Souza JMP, Souza SB. Introdução de alimentos complementares e sua relação com variáveis demográficas e socioeconômicas, em crianças no primeiro ano de vida, nascidas em Hospital Universitário no município de São Paulo. *Rev Bras Epidemiol*. 2003;6(1):29-38. DOI: 10.1590/S1415-790X2003000100005
18. Victora CG, Barros FC, Lima RC, Horta BL, Wells J. Anthropometry and body composition of 18 year old men according to duration of breast feeding: birth cohort study from Brazil. *BMJ*. 2003;327(7420):901. DOI: 10.1136/bmj.327.7420.901
19. Vieira GO, Silva LR, Vieira TO, Almeida JAG, Cabral VA. Hábitos alimentares de crianças menores de 1 ano amamentadas e não-amamentadas. *J Pediatr (Rio J)*. 2004;80(5):411-6. DOI: 10.1590/S0021-75572004000600013
20. Waterland RA, Garza C. Potencial mechanisms of metabolic imprinting that lead to chronic disease. *Am J Clin Nutr*. 1999;69(2):179-97.

Article based on Doctoral thesis by Simon VGN, presented to Faculdade de Saúde Pública at Universidade de São Paulo, in 2007.