

An Acid-Free Anionic Oxoborane Isoelectronic with Carbonyl: Facile Access and Transfer of a Terminal B=O Double Bond

Ying Kai Loh, Kieran Porteous, M. Ángeles Fuentes, Dinh Cao Huan Do, Jamie Hicks, Simon Aldridge*

Inorganic Chemistry Laboratory, Department of Chemistry, University of Oxford, South Parks Road, Oxford, OX1 3QR UK


Supporting Information Placeholder

ABSTRACT: We disclose the synthesis and structural characterization of the first acid-free anionic oxoborane, [K(2.2.2-crypt)][(HCDippN)₂BO] (**1**), which is isoelectronic with classical carbonyl compounds. **1** can readily be accessed from its borinic acid by a simple deprotonation/sequestration sequence. Crystallographic and DFT analyses support the presence of a polarized terminal B=O double bond. Subsequent π bond metathesis converts the B=O bond to a heavier B=S containing system, affording the first anionic thioxoborane [K(2.2.2-crypt)][(HCDippN)₂BS] (**2**), isoelectronic with thiocarbonyls. Facile B=O bond cleavage can also be achieved to access B–H and B–Cl bonds, and via a remarkable oxide (O^{2-}) ion abstraction to generate a borenium cation [(HCDippN)₂B(NC₅H₅)]⁺[OTf]⁻ (**4**). By extension, **1** can act as an oxide transfer agent to organic substrates, a synthetic role traditionally associated with transition metal compounds. Hence we show that B–O linkages, which are often considered to be thermodynamic sinks, can be activated under mild conditions towards bond cleavage and transfer, by exploiting the higher reactivity inherent in the B=O double bond.

Boron has great affinity for oxygen, forming thermodynamically strong B–O bonds (809 kJ mol⁻¹),¹ naturally occurring boron is therefore typically found in the form of borate minerals.² In synthetic chemistry, the highly oxophilic nature of boron is exploited to drive important transformations such as Suzuki coupling and carbonyl hydroboration.³ While these processes are increasingly being adopted in pharmaceutical production, the chemical recycling of B–O bonds requires harsh conditions and/or reagents – as in the commercial production of BCl₃ from B₂O₃, carbon and Cl₂ at 500 °C.

Given the higher reactivity typically found for E=O multiple bonds (e.g. relatively inert C–O single bonds in ethers versus more reactive C=O double bonds in carbonyls), we sought to explore new modes of reactivity

Scheme 1. Examples of acid-stabilized oxoboranes (top), acid- and base-free main group E=O doubly bonded species (middle); Previous work with the NHBO ligand and present work (bottom). (Dipp = 2,6-^tPr₂C₆H₃).




for the BO fragment by developing the chemistry of simple isolable B=O double bonds.

Oxoboranes (R–B=O) represent one such class of organoboron species (albeit transiently stable) that contain a B=O double bond.⁴ The quest for isolable oxoboranes is hampered by the high polarity of the BO fragment and the Lewis acidic nature of the boron atom, making such species prone to head-to-tail oligomerization. Thus, monomeric species are highly elusive and observable only in the gas phase, in a low-temperature matrix, or by means of chemical trapping.⁵ In 2005, Cowley *et al.* exploited an additional neutral donor at boron and Lewis acid coordination at oxygen, to generate a monomeric oxoborane featuring a short B=O distance (1.304(2) Å;

Scheme 1).^{6a} Subsequently, the concept of Lewis/Brønsted acid stabilization has at O been successfully employed to generate other protected oxoboranes featuring similar B=O distances.⁶ Another strategy for the isolation of systems containing BO multiple bonds is by trapping within the coordination sphere of a transition metal, as exemplified by Braunschweig^{7a-d} and by Yamashita.^{7e}

Aside from boron, significant progress has been made recently in the isolation of compounds featuring acid- and base-free main group element E=O double bonds, which are isoelectronic with the C=O double bonds of classical carbonyl compounds. In 2012, Tamao and co-workers documented a stable monomeric germanone [R₂Ge=O], a heavier ketone analogue possessing a Ge=O double bond (Scheme 1).^{8a} Another landmark achievement was the fulfillment of “Kipping’s dream” by the groups of Kato and Inoue, through the isolation of silanones, R₂Si=O, featuring Si=O double bonds.^{8b-d} In addition, Dielmann and co-workers have described a Lewis base free oxophosphonium cation, [R₂P=O]⁺, containing a P=O double bond.^{8c} Inspired by these reports, we wondered whether an isoelectronic acid-free anionic oxoborane of the type [R₂B=O]⁻ could be accessed.

Scheme 2. Synthesis of [K(2.2.2-crypt)][(HCDippN)₂BO] (1), and its subsequent chalcogenation to give [K(2.2.2-crypt)][(HCDippN)₂BS] (2) and {K[(HCDippN)₂BS]}₄ (3).



Very recently, we introduced a new class of N-heterocyclic boryloxy (NHBO) ligand, (HCDippN)₂BO⁻, and demonstrated its ability to stabilize heavier group 14 dioxycarbene analogues (Scheme 1).⁹ We speculated whether the NHBO framework, possessing a 6π electron diazaborolyl scaffold flanked by sterically demanding Dipp groups, would possess the necessary attributes for the isolation of a monomeric, acid-free oxoborane. With this in mind, we exploited the potassiated ligand {K[(HCDippN)₂BO]}₂ (which is dimeric in the solid state) as the starting point of our investigation. To sequester the potassium ion, [2.2.2]cryptand was em-

ployed (Scheme 2), resulting in an immediate colour change from yellow to red. The resulting ¹¹B{¹H} NMR spectrum displays a broad singlet at 20.7 ppm, which is upfield shifted by *ca.* 2 ppm, suggesting slightly increased electron density at boron. Single crystals obtained by slow evaporation from a benzene solution were subjected to X-ray diffraction analysis.




Figure 1. Solid-state structure of **1**, Kohn-Sham HOMO-2, WBI bond order (red) and NPA charges (blue). For clarity, [K(2.2.2-crypt)]⁺ and hydrogen atoms are omitted. Thermal ellipsoids set at 50% probability. Key bond lengths (Å) and angles (°): B1–O1 1.273(8), B1–N1 1.484(7), B1–N2 1.489(7), N1–B1–N2 100.6(4).

The solid-state structure reveals successful sequestration of the potassium ion and the generation of a monomeric anionic oxoborane (**1**, as the [K(2.2.2-crypt)]⁺ salt) (Figure 1). The oxygen atom assumes a terminal position, with the closest contacts being an O1–K1 distance of 5.919(6) Å, and O–H interactions involving the C–H bonds of the flanking ⁱPr groups or the methylene units of the cryptand (*ca.* 2.5 Å). As such, **1** represents the first example of a simple (*i.e.* non-acid stabilized) oxoborane. The central boron atom adopts a trigonal planar geometry, and the most striking feature is the very short B–O bond distance of 1.273(8) Å, which is towards the shorter end of the range of reported distances for Lewis/Brønsted acid stabilized B=O double bonds (1.287(4)–1.329(6) Å).⁶ In comparison with the parent borinic acid (HCDippN)₂BOH (1.373(3) Å),¹⁰ shortening of the B–O distance by 0.1 Å (*ca.* 8%) suggests delocalization of additional electron density by O-to-B π donation, leading to significant B=O double bond character. Consistently, the adjacent B–N bonds are lengthened markedly from (HCDippN)₂BOH (mean 1.429(3) Å) to **1** (mean 1.487(7) Å).

To probe the electronic structure of **1**, we exploited Density Functional Theory (DFT) using a PBE1PBE hybrid exchange-correlation functional and TZVP basis set. B=O π bond character can be located in the HOMO-2 (Figure 1); the Wiberg bond index (WBI) of 1.40, and NPA charges on B (+0.99) and O (-1.03) support a description as a B=O double bond that is polarized towards oxygen. For comparison, analogous calculations on the isoelectronic carbonyl compound (the cyclic urea

(HCDippN)₂CO) reveal a WBI of 1.61 for the C=O fragment and NPA charges of +0.72(C)/−0.64(O).

In contrast to its lighter congener, monomeric thioxoboranes (R−B=S) have been isolated in the absence of acid protection, as demonstrated by reports from the groups of Cui,^{11a} Singh^{11b} and Braunschweig.^{11c} While a cationic thioxoborane has subsequently been reported by Inoue and co-workers,^{11d} the anionic counterpart [R₂B=S][−] (which is isoelectronic with thiocarbonyl compounds) remains unknown. With this in mind, the unquenched B=O motif in **1** was thought to be an ideal candidate to explore carbonyl-like metathesis of the terminal oxygen.

Treatment of **1** with carbon disulfide, CS₂ at room temperature leads to an upfield shift of the ¹¹B{¹H} NMR signal by *ca.* 12 ppm to 33.1 ppm, which is close to that of Inoue's cationic thioxoborane (33.9 ppm) (Scheme 2).^{11d} In addition, a signal at 153.4 ppm was detected in the ¹³C{¹H} NMR spectrum due to carbonyl sulfide, COS, hinting at successful B=O / B=S metathesis. X-ray crystallography unambiguously revealed the identity of the product as [K(2.2.2-crypt)][(HCDippN)₂BS] (**2**). **2** is isostructural with **1**, with sulfur in a terminal position (Figure 2), and therefore represents the first example of a monomeric thioxoborane anion. Closer inspection reveals the B−S distance to be 1.774(1) Å, which is marginally longer than those found in previously reported neutral (1.739(2)–1.752(5) Å)^{11a–c} and cationic (1.710(5) Å)^{11d} systems, indicating a stepwise decrease in bond order with accumulation of negative charge on the BS fragment. As with **1**, we turned to DFT calculations for further insight. The B=S π bond can be located in the HOMO−2 (Figure 2); the WBI of 1.56 and NPA charges on B (+0.61) and S (−0.71) support a description as a B=S double bond that is polarized towards the chalcogen, although to a lesser extent than in **1**.

2 can also be synthesized from dimeric {K[(HCDippN)₂BO]}₂ via an analogous reaction with CS₂. This reaction initially generates the tetrameric inter-




Figure 2. Solid-state structure of **2**, Kohn-Sham HOMO−2, WBI bond order (red) and NPA charges (blue). For clarity, [K(2.2.2-crypt)]⁺ and hydrogen atoms are omitted. Thermal ellipsoids set at 50% probability. Key bond lengths (Å) and angles (°): B1−S1 1.774(1), B1−N1 1.457(2), B1−N2 1.461(2), N1−B1−N2 101.4(9).





Figure 3. Solid-state structure of **3**. For clarity, hydrogen atoms and *i*-Pr groups are omitted, Dipp groups are simplified as wireframes. Thermal ellipsoids set at 50% probability. Key bond lengths (Å) and angles (°): B1−S1 1.789(4), S1−K1 3.112(1), S1−K2 3.084(1), S1−K3 3.185(1).

-mediate {K[(HCDippN)₂BS]}₄ (**3**; Scheme 2 and Figure 3); subsequent treatment with [2.2.2]cryptand leads to sequestration of the K⁺ cations to give monomeric **2**.

The facile access of a B=S bond via π bond metathesis from its lighter congener resembles the chalcogenation of carbonyl compounds. With this in mind, we wanted to explore the applicability of other classical carbonyl transformations to the B=O bond. A key fundamental reaction of the carbonyl function is hydrogenation, typically accomplished by highly reactive metal hydride reagents. In the case of **1**, hydrogenation can be achieved with dimethylamineborane, Me₂NH·BH₃ as the hydrogen transfer source to afford (HCDippN)₂BH (Scheme 3),¹⁰ by analogy with hydrogen transfer from H₃N·BH₃ to a cationic thioxoborane reported by Inoue and co-workers.^{11d}

While B−H containing compounds are valuable reagents in their own right, we wondered whether broader scope could be demonstrated for the conversion of the B=O double bond in **1** to species of the type R₂BX. A conventional protocol to activate carbonyl groups towards nucleophilic attack is by chlorination. To probe this analogy, we exposed **1** to POCl₃ at room temperature, leading to an immediate colour change from red to yellow (Scheme 3). ¹H NMR monitoring reveals clean

Scheme 3. Carbonyl-like reactions of 1 to give (HCDippN)₂BH, (HCDippN)₂BCl, [(HCDippN)₂B(−NC₅H₅)] [OTf][−] (4**).**




conversion to a new species, which can be shown to be the chloroborane $(\text{HCDippN})_2\text{BCl}$.¹⁰ Due to its labile nature, B–Cl containing compounds are versatile building blocks for key organoboron species, such as boronic acids for Suzuki coupling, or even boryl nucleophiles boron for umpolung B–C bond formation.¹⁰

Carbonyl compounds are also known to undergo rearrangements in the presence of trifluoromethanesulfonic anhydride, TF_2O , and transient cations are often proposed as reactive intermediates.¹² With this in mind we wondered whether we could generate a boron cation in analogous fashion, in effect by removal of oxide ion (O^{2-}) from **1**. Accordingly, the reaction of **1** with TF_2O in the presence of pyridine generates the triflate anion, and a cationic species which can be shown by X-ray crystallography to feature a pyridine-ligated borenium cation (*i.e.* $[(\text{HCDippN})_2\text{B}(\text{NC}_5\text{H}_5)][\text{OTf}]$ (**4**), Scheme 3).¹³

In a landmark report, Bertrand and co-workers demonstrated in 2012 that a main group compound could mimic a transition metal complex in acting as a nitrogen atom transfer agent (Scheme 4). Reaction of an isolable phosphinonitrene with isonitrile initially forms an adduct, which can subsequently be cleaved with an alkyltriflate to afford a mixture of cyanamide and carbodiimide products.¹⁴ The isoelectronic BO^-/PN relation

Scheme 4. Bertrand's phosphinonitrene as a nitrogen atom transfer agent (top); an analogous synthetic cycle with **1 as an oxide ion transfer agent (middle); solid-state structure of **5** (bottom).**



-ship between **1** and this phosphinonitrene prompted us to investigate the possibility of using **1** as an oxide ion transfer agent, a task typically accomplished by transition metal systems.¹⁵

As a proof of concept, we treated **1** with 1,3-di-*p*-tolylcarbodiimide. X-ray diffraction analysis of the product reveals not simple adduct formation, but 1,2-addition of the B=O bond across the C=N double bond to afford iso-ureate **5**. Although unsymmetrical binding of the $\text{NC}(\text{O})\text{N}$ fragment at boron is implied by crystallography, the ^1H NMR of **5** shows a single *p*-Tol group environment in solution, hinting at the facile fluxional exchange. Formally, this transformation can be considered as insertion of a carbodiimide into the B=O double bond. To release the functionalized substrate, oxalyl chloride was employed to generate the cyclic urea derivative $(\text{OC}(p\text{-Tol})\text{N})_2\text{CO}$ and $(\text{HCDippN})_2\text{BCl}$.¹⁰ While this stepwise procedure closely resembles the N-transfer reported by Bertrand, we could also achieve the same transformation in a one-pot sequence without isolation of intermediate **5**. Moreover, to close the oxide ion transfer cycle, $(\text{HCDippN})_2\text{BCl}$ can be converted to borinic acid $(\text{HCDippN})_2\text{BOH}$ using water in the presence of pyridine and subsequently back to **1** via simple deprotonation and cation sequestration.⁹

In conclusion, we disclose the first acid-free anionic oxoborane **1**, which exhibits classical carbonyl-like reactivity, and also mimics the activity of transition metal systems by acting as an oxide ion transfer agent. Hence, boron-oxygen linkages which are often considered to be thermodynamic sinks, can be shown to be activated towards facile bond cleavage and transfer, by exploiting the higher reactivity intrinsic in the B=O double bond.

ASSOCIATED CONTENT

Supporting Information. Full synthetic and characterizing data for new compounds, representative NMR spectra, details of crystallographic and computational studies (PDF). CIFs for X-ray crystal structures (CCDC: 1906845-1906849). The Supporting Information is available free of charge on the ACS Publications website.

AUTHOR INFORMATION

Corresponding Author

Prof Simon Aldridge, simon.aldridge@chem.ox.ac.uk.

Funding Sources

A*STAR (scholarship for YKL), the Jardine-Oxford Scholarship (DDCH) and EPSRC (MAF, EP/K014714/1).

REFERENCES

- (1) For bond dissociation energies: Haynes, W. M. *CRC Handbook of Chemistry and Physics, Vol. 96*; CRC Press: Boca Raton, 2015.
- (2) For minerals: Klein, C.; Hurlbut, C. S. *Manual of Mineralogy*; Wiley: New York, 1985.
- (3) (a) Miyaura, N.; Suzuki, A. Palladium-catalyzed cross-coupling reactions of organoboron compounds. *Chem. Rev.* **1995**, *95*, 2457-2483. (b) Brown, H. C. Hydroboration – a powerful synthetic tool. *Tetrahedron* **1961**, *12*, 117-138.

(4) See for example: (a) Franz, D.; Inoue, S. Advances in the development of complexes that contain a group 13 element chalcogen multiple bond. *Dalton Trans.* **2016**, *45*, 9385-9397. (b) Fischer, R. C.; Power, P. P. π -Bonding and the lone pair effect in multiple bonds involving heavier main group elements: developments in the new millennium. *Chem. Rev.* **2010**, *110*, 3877-3923.

(5) For gas phase studies: (a) Hildenbrand, D. L.; Theard, L. P.; Saul, A. M. Transpiration and mass spectrometric studies of equilibria involving BOF(g) and (BOF)₂(g). *J. Chem. Phys.* **1963**, *39*, 1973-1978. (b) Bock, H.; Cederbaum, L.; Niessen, W. v.; Paetzold, P.; Rosmus, R.; Solouki, B. Methylboron oxide, H₃CBO. *Angew. Chem., Int. Ed.* **1989**, *28*, 88-90. For matrix isolation: (c) Burkholder, T. R.; Andrews, L. Reactions of boron atoms with molecular oxygen. Infrared spectra of BO, BO₂, B₂O₂, B₂O₃, and BO₂⁻ in solid argon. *J. Chem. Phys.* **1991**, *95*, 8697-8709. (d) Lanzisera, D. V.; Andrews, L. Reactions of laser-ablated boron atoms with methanol. Infrared spectra and *ab initio* calculations of CH₃BO, CH₂BOH, and CH₂BO in solid argon. *J. Phys. Chem. A.* **1997**, *101*, 1482-1487. (e) Bettinger, H. F. Reversible formation of organyl(oxo)boranes (RBO) (R = C₆H₅ or CH₃) from boroxins ((RBO)₃): a matrix isolation study. *Organometallics* **2007**, *26*, 6263-6267. For chemical trapping: (f) Ito, M.; Tokitoh, N.; Okazaki, R. A novel approach to an oxoborane and its Lewis base complex. *Tetrahedron Lett.* **1997**, *38*, 4451-4454. (g) Ito, M.; Tokitoh, N.; Okazaki, R. Formation of oxoborane and thioxoborane from a dithiastannaboretane derivative. *Phosphorus, Sulfur Silicon Relat. Elem.* **1997**, *124*, 533-536. (h) Paetzold, P.; Neyses, S.; Geret, L. Oxo(trisilyl)boran (Me₃Si)₃CBO als Zwischenstufe. *Z. Anorg. Allg. Chem.* **1995**, *621*, 732-736. (i) Groteklaes, M.; Paetzold, P. Oxo(tert-butylphenyl)boran ArBO als Zwischenstufe. *Chem. Ber.* **1988**, *121*, 809-810. (j) Pachchaly, B.; West, R. Synthesis of a 1,3-dioxo-2,4-diboretane, an oxoborane precursor. *J. Am. Chem. Soc.* **1985**, *107*, 2987-2988. (k) Hanecker, R.; Noth, H.; Wietelmann, U. Beiträge zur chemie des bors, 171. Kristall- und molekülstruktur von 2,4-bis(2,2,6,6-tetramethylpiperidino)-1,3,2,4-dichalcogendiboretanen. *Chem. Ber.* **1986**, *119*, 1904-1910.

(6) For acid-stabilized monomeric oxoboranes: (a) Vidovic, D.; Moore, J. A.; Jones, J. N.; Cowley, A. H. Synthesis and characterization of a coordinated oxoborane: Lewis acid stabilization of a boron-oxygen double bond. *J. Am. Chem. Soc.* **2005**, *127*, 4566-4567. (b) Wang, Y.; Hu, H.; Zhang, J.; Cui, C. Comparison of anionic and Lewis acid stabilized N-heterocyclic oxoboranes: their facile synthesis from a borinic acid. *Angew. Chem., Int. Ed.* **2011**, *50*, 2816-2819. (c) Loh, Y. K.; Chong, C. C.; Ganguly, R.; Li, Y.; Vidovic, D.; Kinjo, R. 1,2,4,3-Triazaborole-based neutral oxoborane stabilized by a Lewis acid. *Chem. Commun.* **2014**, *50*, 8561-8564. (d) Swarnakar, A. K.; Hering-Junghans, C.; Ferguson, M. J.; McDonald, R.; Rivard, E. Oxoborane (RBO) complexation and concomitant electrophilic bond activation processes. *Chem. Eur. J.* **2017**, *23*, 8628-8631.

(7) For transition metal oxoboranes: (a) Braunschweig, H.; Radacki, K.; Schneider, A. Oxoboryl complexes: boron-oxygen triple bonds stabilized in the coordination sphere of platinum. *Science* **2010**, *328*, 345-347. (b) Westcott, S. A. BO chemistry comes full circle. *Angew. Chem., Int. Ed.* **2010**, *49*, 9045-9046. (c) Braunschweig, H.; Radacki, K.; Schneider, A. Reactivity of an oxoboryl complex toward fluorinated aryl boron reagents. *Chem. Commun.* **2010**, *46*, 6473-6475. (d) Bertsch, S.; Brand, J.; Braunschweig, H.; Hupp, F.; Radacki, K. Platinum oxoboryl complexes as substrates for the formation of 1:1, 1:2, and 2:1 Lewis acid-base adducts and 1,2-dipolar additions. *Chem. Eur. J.* **2015**, *21*, 6278-6285. (e) Miyada, T.; Yamashita, M. Oxygenation of a ruthenium complex bearing a PBP-pincer ligand inducing the

formation of a boronate ligand with a weak Ru-O bond. *Organometallics* **2013**, *32*, 5281-5284.

(8) For acid/base free main group E=O bonds: (a) Li, L.; Fukawa, T.; Matsuo, T.; Hashizume, D.; Fueno, H.; Tanaka, K.; Tamao, K. A stable germanone as the first isolated heavy ketone with a terminal oxygen atom. *Nat. Chem.* **2012**, *4*, 361-365. (b) Alvarado-Beltran, I.; Rosas-Sanchez, A.; Baceiredo, A.; Saffon-Merceron, N.; Branchadell, V.; Kato, T. A fairly stable crystalline silanone. *Angew. Chem., Int. Ed.* **2017**, *56*, 10481-10485. (c) Wendel, D.; Reiter, D.; Porzelt, A.; Altmann, J.; Inoue, S.; Rieger, B. Silicon and oxygen's bond of affection: an acyclic three-coordinate silanone and its transformation to an iminosiloxysilylene. *J. Am. Chem. Soc.* **2017**, *139*, 17193-17198. (d) Rosas-Sanchez, A.; Alvarado-Beltran, I.; Baceiredo, A.; Saffon-Merceron, N.; Massou, S.; Hashizume, D.; Branchadell, V.; Kato, T. Cyclic (amino)(phosphonium bora-ylide)silanone: a remarkable room-temperature-persistent silanone. *Angew. Chem., Int. Ed.* **2017**, *56*, 15916-15920. (e) Wünsche, M. A.; Witteler, T.; Dielmann, F. Lewis base free oxophosphonium ions: tunable, trigonal-planar Lewis acids. *Angew. Chem., Int. Ed.* **2018**, *57*, 7234-7239.

(9) Loh, Y. K.; Ying, L.; Fuentes, M. A.; Do, D. C. H.; Aldridge, S. An N-heterocyclic boryloxy ligand isoelectronic with N-heterocyclic imines: access to an acyclic dioxysilylene and its heavier congeners. *Angew. Chem., Int. Ed.* **2019**, *58*, 4847-4851.

(10) Segawa, Y.; Suzuki, Y.; Yamashita, M.; Nozaki, N. Chemistry of boryllithium: synthesis, structure and reactivity. *J. Am. Chem. Soc.* **2008**, *130*, 16069-16079.

(11) For monomeric thioxoboranes: (a) Wang, H.; Zhang, J.; Hu, H.; Cui, C. Access to B=S and B=Se double bonds via sulfur and selenium insertion into a B-H bond and hydrogen migration. *J. Am. Chem. Soc.* **2010**, *132*, 10998-10999. (b) Jaiswal, K.; Prashanth, B.; Ravi, S.; Shasundar, K. R.; Singh, S. Reactivity of a dihydroboron species: synthesis of a hydroborenum complex and an expedient entry into stable thioxo- and selenoxo-boranes. *Dalton Trans.* **2015**, *44*, 15779-15785. (c) Liu, S.; Legare, M.; Hofmann, A.; Braunschweig, H. A boradiselenirane and a boraditellurirane: isolable heavy analogs of dioxiranes and dithiiranes. *J. Am. Chem. Soc.* **2018**, *140*, 11223-11226. (d) Franz, D.; Irran, E.; Inoue, S. Isolation of a three-coordinate boron cation with a boron-sulfur double bond. *Angew. Chem., Int. Ed.* **2014**, *53*, 14264-14268.

(12) Barznenok, I. L.; Nenajdenko, V. G.; Balenkova, E. S. Chemical transformations induced by triflic anhydride. *Tetrahedron* **2000**, *56*, 3077-3119.

(13) See for example: Franz, D.; Inoue, S. Cationic complexes of boron and aluminum: an early 21st century viewpoint. *Chem. Eur. J.* **2019**, *25*, 2898-2926.

(14) (a) Dielmann, F.; Back, O.; Henry-Ellinger, M.; Jerabek, P.; Frenking, G.; Bertrand, G. A crystalline singlet phosphinonitrene: a nitrogen atom-transfer agent. *Science* **2012**, *337*, 1526-1528. (b) Schulz, A.; Villinger, A. Stabilized transient R₂PN species. *Angew. Chem., Int. Ed.* **2013**, *52*, 3068-3070. (c) Dielmann, F.; Moore, C. E.; Rheingold, A. L.; Bertrand, G. Crystalline, Lewis base-free, cationic phosphoranimes (iminophosphonium salts). *J. Am. Chem. Soc.* **2013**, *135*, 14071-14073. (d) Dielmann, F.; Andrada, D. M.; Frenking, G.; Bertrand, G. Isolation of bridging and terminal coinage metal-nitrene complexes. *J. Am. Chem. Soc.* **2014**, *136*, 3800-3802. (e) Dielmann, F.; Bertrand, G. Reactivity of a stable phosphinonitrene towards small molecules. *Chem. Eur. J.* **2015**, *21*, 191-198.

(15) For oxide ion transfer by transition metals, see for example: Lee, G. R.; Cooper, N. J. Transfer of an oxide ion from an anionic carbon dioxide complex to a cationic carbonyl complex. *Organometallics* **1985**, *4*, 1467-1468.

