

University of Montana

ScholarWorks at University of Montana

Numerical Terradynamic Simulation Group
Publications

Numerical Terradynamic Simulation Group

2012

An assessment of the carbon balance of arctic tundra: comparisons among observations, process models, and atmospheric inversions

A. D. McGuire

T. R. Christensen

D. J. Hayes

A. Herault

E. Euskirchen

See next page for additional authors

Follow this and additional works at: https://scholarworks.umt.edu/ntsg_pubs

Let us know how access to this document benefits you.

Recommended Citation

McGuire, A. D., Christensen, T. R., Hayes, D., Herault, A., Euskirchen, E., Kimball, J. S., Koven, C., Lafleur, P., Miller, P. A., Oechel, W., Peylin, P., Williams, M., and Yi, Y.: An assessment of the carbon balance of Arctic tundra: comparisons among observations, process models, and atmospheric inversions, *Biogeosciences*, 9, 3185-3204, doi:10.5194/bg-9-3185-2012, 2012.

This Article is brought to you for free and open access by the Numerical Terradynamic Simulation Group at ScholarWorks at University of Montana. It has been accepted for inclusion in Numerical Terradynamic Simulation Group Publications by an authorized administrator of ScholarWorks at University of Montana. For more information, please contact scholarworks@mso.umt.edu.

Authors

A. D. McGuire, T. R. Christensen, D. J. Hayes, A. Heroult, E. Euskirchen, Y. Yi, John S. Kimball, C. Koven, P. Lafleur, P. A. Miller, Walt Oechel, P. Peylin, and M. Williams

**An assessment of
the carbon balance of
arctic tundra**

A. D. McGuire et al.

An assessment of the carbon balance of arctic tundra: comparisons among observations, process models, and atmospheric inversions

A. D. McGuire¹, T. R. Christensen^{2,3}, D. Hayes⁴, A. Heroult², E. Euskirchen⁵, Y. Yi⁶, J. S. Kimball⁶, C. Koven⁷, P. Lafleur⁸, P. A. Miller², W. Oechel⁹, P. Peylin¹⁰, and M. Williams¹¹

¹U.S. Geological Survey, Alaska Cooperative Fish and Wildlife Research Unit, University of Alaska Fairbanks, Fairbanks, AK, USA

²Department of Earth and Ecosystem Science, Lund University, Lund, Sweden

³Greenland Climate Research Centre, Nuuk, Greenland

⁴Oak Ridge National Laboratory, Oak Ridge, TN, USA

⁵Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK, USA

⁶Flathead lake Biological Station, Division of Biological Sciences, University of Montana, Missoula, MT, USA

⁷Lawrence Berkeley National Laboratory, Berkeley, CA, USA

⁸Department of Geography, Trent University, Peterborough, ON, Canada

⁹Department of Biology, San Diego State University, San Diego, CA, USA

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



¹⁰Laboratoire Sciences Climat Environnement, Institut Pierre Simon Laplace, Paris, France

¹¹School of GeoScience, University of Edinburgh, Edinburgh, UK

Received: 23 January 2012 – Accepted: 23 March 2012 – Published: 17 April 2012

Correspondence to: A. D. McGuire (adm McGuire@alaska.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

9, 4543–4594, 2012

**An assessment of
the carbon balance of
arctic tundra**

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Although arctic tundra has been estimated to cover only 8% of the global land surface, the large and potentially labile carbon pools currently stored in tundra soils have the potential for large emissions of carbon (C) under a warming climate. These emissions as radiatively active greenhouse gases in the form of both CO₂ and CH₄ could amplify global warming. Given the potential sensitivity of these ecosystems to climate change and the expectation that the Arctic will experience appreciable warming over the next century, it is important to assess whether responses of C exchange in tundra regions are likely to enhance or mitigate warming. In this study we compared analyses of C exchange of Arctic tundra between 1990–1999 and 2000–2006 among observations, regional and global applications of process-based terrestrial biosphere models, and atmospheric inversion models. Syntheses of the compilation of flux observations and of inversion model results indicate that the annual exchange of CO₂ between arctic tundra and the atmosphere has large uncertainties that cannot be distinguished from neutral balance. The mean estimate from an ensemble of process-based model simulations suggests that arctic tundra acted as a sink for atmospheric CO₂ in recent decades, but based on the uncertainty estimates it cannot be determined with confidence whether these ecosystems represent a weak or a strong sink. Tundra was 0.6°C warmer in the 2000s compared to the 1990s. The central estimates of the observations, process-based models, and inversion models each identify stronger sinks in the 2000s compared with the 1990s. Similarly, the observations and the applications of regional process-based models suggest that CH₄ emissions from arctic tundra have increased from the 1990s to 2000s. Based on our analyses of the estimates from observations, process-based models, and inversion models, we estimate that arctic tundra was a sink for atmospheric CO₂ of 110 Tg C yr⁻¹ (uncertainty between a sink of 291 Tg C yr⁻¹ and a source of 80 Tg C yr⁻¹) and a source of CH₄ to the atmosphere of 19 Tg C yr⁻¹ (uncertainty between sources of 8 and 29 Tg C yr⁻¹). The suite of analyses conducted in this study indicate that it is clearly important to reduce uncertainties in the

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



observations, process-based models, and inversions in order to better understand the degree to which Arctic tundra is influencing atmospheric CO₂ and CH₄ concentrations. The reduction of uncertainties can be accomplished through (1) the strategic placement of more CO₂ and CH₄ monitoring stations to reduce uncertainties in inversions, (2) improved observation networks of ground-based measurements of CO₂ and CH₄ exchange to understand exchange in response to disturbance and across gradients of hydrological variability, and (3) the effective transfer of information from enhanced observation networks into process-based models to improve the simulation of CO₂ and CH₄ exchange from arctic tundra to the atmosphere.

1 Introduction

The distribution of the arctic tundra biome is generally north of the boreal forest tree-line and covers approximately 8% of the global land surface (McGuire et al., 1997). The exact location of the biome's southern border is subjective, with the transition between closed boreal forest and treeless tundra up to several hundred kilometers wide in regions of low topographic relief (Vlassova, 2002; Callaghan et al., 2005). Climate in the Arctic is harsh, characterized by cold winters and cool summers, with mean July temperatures below ~12°C (Callaghan et al., 2005), and annual mean temperatures typically below -10°C (New et al., 1999). Consequently, plant growth is restricted to a relatively short growing season on the order of three months or less during the boreal summer. The tundra biome is home to approximately 1800 species of vascular plants and has less species diversity than more temperate biomes (Callaghan et al., 2005). The stature of vascular plant species is limited by environmental conditions, with trees almost entirely absent and woody plant species restricted largely to shrubs and dwarf shrubs. In addition to vascular plant species, non-vascular mosses and lichens play a very important role in the structure and function of tundra ecosystems. Frozen soils are prevalent in northern high latitudes and there is a gradient of continuous to discontinuous permafrost from north to south. Most of the tundra biome is underlain

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



by continuous permafrost. The spatial and temporal dynamics of permafrost and periodic disturbance are crucial in shaping the arctic landscape and its heterogeneity with important consequences for the areal extent of wetlands and the exchange of carbon dioxide (CO₂) and methane (CH₄).

5 Future climate warming is predicted to be very pronounced over the Arctic, especially during winter and spring. The arctic autumn-winter is expected to warm between 3 and 6 °C by 2080 (SWIPA Assessment Executive Summary, 2011), which is expected to lead longer growing seasons, thawing of permafrost, warming and deepening of the soil active layer, and large changes in hydrology. These changes are likely to substantially
10 affect tundra ecosystem structure and function. In fact, there is increasing evidence that physical and ecological changes are already occurring throughout the tundra biome (Serreze et al., 2003; Hinzman et al., 2005; McDonald et al., 2004; Piao et al., 2008, 2011; Post et al., 2009; Rawlins et al., 2010; Rowland et al., 2010; Beck and Goetz, 2011; Kim et al., 2011).

15 The large and potentially labile carbon (C) pools currently stored in arctic soils (Ping et al., 2008; Tarnocai et al., 2009) have the potential to be emitted as radiatively active greenhouse gases in the form of both CO₂ and CH₄ under warmer conditions (Schuur et al., 2008, 2011; Chapin et al., 2008; McGuire et al., 2009; Schaefer et al., 2011; Koven et al., 2011). Whether the emissions of CO₂ from tundra soils tends to amplify
20 or mitigate global warming depends on the degree to which C accumulation in tundra plants responds to warming (Sitch et al., 2007). This balance determines whether the tundra is a source or sink of CO₂. Changes in the emissions of CH₄ may also affect the degree to which tundra amplifies or mitigates global warming. While CH₄ has only a small role on the mass balance of C between the atmosphere and tundra, it is a highly
25 potent greenhouse gas. Changes in CH₄ emissions are likely to be strongly linked to changes in hydrology. Current emissions of CH₄ are difficult to quantify because of substantial variability in time and space due to variations in the environment associated with topography, hydrology, and soil chemistry.

**An assessment of
the carbon balance of
arctic tundra**

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**An assessment of
the carbon balance of
arctic tundra**A. D. McGuire et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Because of the substantial changes that are already affecting the structure and function of arctic tundra, it is important to assess how C exchange of arctic tundra has been changing in recent decades. The response of C dynamics of arctic tundra to environmental change can be evaluated through a synthesis of (1) observations of C exchange with the atmosphere, (2) the application of process-based models, and (3) the analysis of atmospheric inversions of C exchange with the atmosphere. Each of these scaling approaches has its strengths, weaknesses, and limitations in assessing the carbon dynamics of arctic tundra. In this paper we compare analyses of C dynamics of arctic tundra in the two most recent decades among the three scaling approaches to gain insight on how C exchange of arctic tundra may be responding to ongoing environmental changes. The analysis in this paper represents the arctic tundra contribution to the Global Carbon Project's REgional Carbon Cycle and Assessment Processes (RECCAP) synthesis (Canadell et al., 2011).

2 Methods

2.1 Estimates from flux observations

Methods for ground-based observation of the exchange of C between land and atmosphere face great challenges in arctic environments. The challenges include (1) comprehensive spatial coverage in the face of a heterogeneous landscape mosaic that is often characterized by “hot spots”; (2) continuous sampling to achieve full year-round estimates of carbon dynamics; (3) high temporal resolution to sample episodic exchanges of CO₂ and CH₄; and (4) collection of C exchange data without line power in remote conditions.

A single technique is not available that meets all of these challenges. Currently, manual chambers, automatic chambers, and eddy covariance towers are the primary techniques being used to measure C exchange between tundra with the atmosphere. For CO₂ flux measurements, these sampling techniques are linked to infra-red gas

analyzers that measure CO₂ concentrations. For CH₄, the field technology is less developed, and has relied on gas sample collection in the field, with laboratory estimates of CH₄ concentrations using gas chromatographs. Measurement systems have recently been developed that allow direct CH₄ concentration estimates in the field. In Table 1 we compare the relative performance of these methods for a number of requirements and considerations. Because typical tundra areas are heterogeneous, it is often necessary to employ chamber methods of measurement to differentiate the C exchange for the individual components (e.g., soil, moss, and vascular plants) and to better understand the underlying processes of land-atmosphere C exchange. Chamber-based measurements complement tower-based measurements that more effectively integrate across a heterogeneous landscape. In Appendix A in the Supplement we have compiled approximately 250 estimates from 120 published papers of the mean exchange of CO₂ and CH₄ between Arctic tundra and the atmosphere at growing season, winter season, and annual time scales based on published observational studies. The exchange of CO₂ with the atmosphere is reported as net ecosystem exchange (NEE), i.e., net land-atmosphere CO₂ flux, in which a positive NEE represents a loss of CO₂ from tundra to the atmosphere. Similarly the exchange of CH₄ is reported as a positive flux when the net exchange is to the atmosphere and as a negative flux when the net exchange is into the ecosystem. Both CO₂ and CH₄ estimates are reported in units of C. In this study we used only estimates of CO₂ and CH₄ exchange from Appendix A in the Supplement for the time period between 1990 and 2009 unless stated otherwise.

2.2 Estimates from process-based models

The spatial domain of arctic tundra we considered in this study (Fig. 1) was defined by the Regional Carbon Cycle Assessment and Processes (RECCAP) Activity. It is important to note that the spatial domain of arctic tundra was defined from an atmospheric perspective as a region that could potentially be resolved by the applications of inversion models. The region does extend into boreal forest in some areas (for example in western North America). In this study we compare the carbon dynamics of Arctic tundra

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



between 1990 and 2006 estimated by regional applications of three models that have focused on representing processes in ecosystems underlain by permafrost: LPJ-Guess WHyMe (Smith et al., 2001; Wania et al., 2009a, b, 2010; Hickler et al., 2012), Orchidee (Koven et al., 2009, 2011), and version 6 of the Terrestrial Ecosystem Model (TEM6; McGuire et al., 2010; Hayes et al., 2011). For evaluating the production estimates of the three regional process-based models, we have also included the Terrestrial Carbon Flux (TCF) model (Kimball et al., 2009) in the regional process-based model analysis. The general features of the models are compared in Table 2 (see Appendix B in the Supplement for more details).

The TCF is unique among the models in that it is partially driven by satellite-based vegetation gross primary production (GPP) estimates from the NASA Moderate Resolution Imaging Spectroradiometer (MODIS) for the period between 2000 and 2009 (Zhao and Running, 2010). The MODIS (MOD17) GPP estimates are used to derive vegetation net primary production (NPP), while heterotrophic respiration (RH) is determined from a simulated surface (<10 cm depth) soil organic carbon pool and dynamic soil moisture and temperature constraints to litter decomposition. The TCF does not account for other carbon emission sources, including fire disturbance, so NEE is derived as the residual difference between NPP and RH, and assumed equivalent to net ecosystem production (NEP). The TCF calculations assume dynamic steady-state conditions between NPP and RH so that estimated NEE/NEP has no trend over the decade from 2000 to 2009. The other three models were driven in a prognostic fashion by atmospheric CO₂ and climate data of their own choosing over the simulation period. Because of each model's choice of climate data, both LPJ-Guess WHyMe and TEM6 produced estimates for only the 1990–2006 time period. Therefore, we only compare 2000–2006 among the models in the most recent decade. See Appendix B in the Supplement for more details on the application of the regional models in this study.

All models make estimates of GPP, NPP, RH, and NEP. Both LPJ-Guess and TEM6 make estimates of losses of carbon to the atmosphere associated with fire and biogenic CH₄ emissions. In addition, TEM6 also calculates ecosystem losses of carbon

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



associated with the export of harvested products and dissolved organic carbon. For each of the models we also calculate the net ecosystem carbon balance (NCB) as the sum of NPP and the atmospheric and export losses. The sign conventions for NEP and NCB are positive for a net flux of carbon into the ecosystem. We also report net CO₂-C exchange with a sign convention of positive representing a source to the atmosphere and negative representing a sink into tundra ecosystems. In this study, we compare the mean C budget estimated from 1990–1999 and 2000–2006 among simulations of LPJ-Guess WHyMe, Orchidee, and TEM6. We also report net CO₂-C exchange for each of the models as previously defined. We also provide mean GPP and NPP estimates simulated by TCF for 2000–2006. To explore issues involving the mean seasonal cycle of CO₂ exchange, we compare the mean monthly flux estimates for GPP, NPP, RH, and NEP for 1990–2006 simulated by LPJ-Guess, Orchidee, and TEM6, and for 2000–2009 simulated by TCF. We also compare interannual variability for estimates of GPP, NPP, RH, NEP, other atmospheric losses, export losses, and NCB among the models. To explore the importance of changing climate on the regional applications of the process-based models, we conducted additional simulations that were driven by constant climate keeping all other drivers unchanged. The constant climate for the three models was based on the 1901–1930 climate used to drive their transient simulations. We estimate the effect of a changing climate on NPP, RH, and NEP between 1990 and 2006 by subtracting the estimates of the constant climate simulation of each model from that of the corresponding transient climate simulation.

The RECCAP activity is also comparing the mean C budgets for the 1990s and 2000s estimated by a suite of global applications of dynamic global vegetation models (DGVMs). These DGVM applications were conducted as part of the Trendy project (<http://dgvm.ceh.ac.uk>) to examine trends in the net land C exchange over the period 1980–2009. In this study we compare the mean C budget of these global applications for the Arctic tundra region of this study for the time periods 1990–1999 and 2000–2006. The DGVM applications compared in this study include contributions from CLM4C (Lawrence et al., 2010), CLM4CN (Thornton et al., 2007, 2009; Bonan and

BGD

9, 4543–4594, 2012

**An assessment of
the carbon balance of
arctic tundra**

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Levis, 2010; Lawrence et al., 2010), Hyland (Levy et al., 2004), LPJ (Sitch et al., 2003), LPJ-Guess (Smith et al., 2001), O-CN (v0.74; Zaehle and Friend, 2010), SDGVM (Woodward et al., 1995; Woodward and Lomas, 2004), and TRIFFID (Cox, 2001). The models used a common protocol (<http://dgvm.ceh.ac.uk>) applying CRU-NCEP climatology over the period 1901–2009. Note that the global application of LPJ-Guess is quite different from the regional application of LPJ-Guess WHyMe, which represents processes relevant to arctic tundra function and structure including (1) soil water freezing; (2) arctic shrub and open ground plant functional types (e.g., *Sphagnum* mosses and tundra graminoids; (3) peatland hydrology, decomposition, and plant functional types; (4) a methane module for peatlands; and (5) root exudates (see Appendix B in the Supplement for details).

2.3 Estimates from inversion-based models

We also analyzed the mean land-atmosphere CO₂-C exchange for the arctic tundra domain from a set of ten inversion models that were applied in support of REC-CAP analyses (Gurney et al., 2012). The inversion models include C13_CCAM_law, C13_MATCH_rayner, CTRACKER_EU, CTRACKER_US, JENA_s96_v3.3, JMA_2010, LSCE_an_v2.1, LSCE_var_v1.0, NICAM_niwa_woaia, and rigc_patra. Among these inversion models, the applications span the time period from 1985–2009. However, the period of application is highly variable among the models. We report the mean NEE estimate for arctic tundra from 1990–1999 and 2000–2006 for eight of the applications of these inversion models; we do not report the results from CTRACKER_EU or CTRACKER_US as these models did not start making estimates until 2001. We report the mean season cycle of CO₂-C exchange estimates based on the time period of application of each of the ten models. Similarly, we report the interannual variability of NEE anomalies across the time period of application of each of the ten models.

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results

3.1 Estimates based on flux observations

3.1.1 CO₂ exchange

Most direct observational studies of the exchange of CO₂ between tundra and the atmosphere have been conducted during the summer growing season. These studies generally indicate that arctic tundra has been a sink for atmospheric CO₂ during the summer in all subregions of the Arctic (i.e., NEE is largely negative; Fig. 2) and that there has not been a substantial change in the sink strength between the 1990s and 2000s (Table 3). While it appears from Table 3 that the summer sink strength in Eurasia has more than doubled since 2000, the mean estimates are not significantly different as the confidence limits since 2000 completely overlap the confidence limits for the 1990s. The existing observations suggest that wet (lowland) tundra is strong sink for CO₂ during the growing season, while dry/mesic (upland) tundra tends to be a growing season source of CO₂ to the atmosphere with a confidence interval that overlaps neutral balance (Table 4).

Only a handful of studies have estimated the exchange of CO₂ in winter, as there are considerable challenges in maintaining accurate flux measurements outside the growing season. The available estimates indicate that tundra ecosystems are sources of CO₂ to the atmosphere (Fig. 2, Table 3). Given that few studies have been conducted, it does not appear that the strength of sources differs among subregions (Table 3). Also, the scarcity of winter exchange data does not allow us to evaluate if there are differences in source strength between the 1990s and 2000s. The existing observations suggest that there is little difference in the source strength of CO₂ during the winter between wet and dry/mesic tundra (Table 4).

There are a growing number of observationally based studies that are attempting to estimate annual CO₂ exchange between tundra and the atmosphere. In general, the range of variability among estimates is scattered around neutral annual CO₂ exchange

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 in all subregions (Fig. 2). In North America, the data suggest that tundra ecosystems tended to be annual sources of CO₂ to the atmosphere prior to 2000, but have been approximately neutral or weak sinks since 2000 (Table 3). The existing observations suggest that wet tundra is a strong sink for CO₂ annually, while dry/mesic tundra tends to be an annual source of CO₂ to the atmosphere with a confidence interval the overlaps neutral balance (Table 4).

10 Based on the mean and range of NEE observations reported in Table 3 for the different geographical regions of the Arctic, we developed first order estimates of NEE and ranges in uncertainty in those estimates for Arctic tundra between 1990 and 2009 (see Appendix C in the Supplement for details). This analysis suggests that tundra was source of 138 Tg C yr⁻¹ as CO₂ to the atmosphere in the 1990s with a range of uncertainty between a 102 Tg C yr⁻¹ sink and a 378 Tg C yr⁻¹ source. In contrast, the analysis suggests that tundra was a 202 Tg C yr⁻¹ sink in the 2000s with an uncertainty range between a 628 Tg C yr⁻¹ sink and 224 Tg C yr⁻¹ source. Across the two
15 decades, we estimate that tundra was a sink of 103 Tg C yr⁻¹ with an uncertainty between a 297 Tg C yr⁻¹ sink and 89 Tg C yr⁻¹ source.

3.1.2 CH₄ exchange

20 Similar to data on CO₂ exchange, most of the studies of the exchange of CH₄ between tundra and the atmosphere have been conducted during the summer growing season. These studies generally indicate that arctic tundra is a substantial source of CH₄ to the atmosphere during the summer (Fig. 2) and that there has not been a substantial change in the strength of the source between the 1990s and 2000s (Table 3). However, the existing observations suggest that there are differences among different tundra types as mean summer emissions of CH₄ for wet tundra are 9.2 g C m⁻² compared with 0.8 g C m⁻² for dry/mesic tundra with no overlap in the confidence intervals
25 (Table 4). There are only two studies that have estimated the exchange of CH₄ in winter, and these studies indicate that tundra ecosystems are a weak source of around

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.0 (range 0.1 to 6.0) g CH₄-C m⁻² winter⁻¹ to the atmosphere. The inference from the studies in North America during the 2000s suggest that there are substantial emissions of CH₄ during the winter, but the studies in northern Europe in the 2000s suggest that CH₄ emissions in winter are negligible. The inconsistency is likely to be due to bias associated with different representations of wet and dry/mesic tundra in the summer vs. annual estimates, as the summer and annual fluxes are consistent with each other when broken down by landscape position/wetness (Table 4).

Based on the mean and range of CH₄ observations reported in Table 3 for the different geographical regions of the Arctic, we developed estimates of CH₄ emissions and ranges in uncertainty in those estimates for Arctic tundra before and since 2000 (see Appendix C in the Supplement for details). This analysis suggests that tundra emitted 10 Tg C yr⁻¹ as CH₄ to the atmosphere in the 1990s with a range of uncertainty between -1 and 22 Tg C yr⁻¹. The analysis suggests that tundra was stronger emitter of CH₄ during the 2000s (20 Tg C yr⁻¹), but the uncertainties since 2000 are much larger than in the 1990s (between a sink of 11 and a source of 51 Tg C yr⁻¹). Across the two decades, our analysis indicates that tundra emitted 11 Tg C yr⁻¹ as CH₄ with a range from 0 to 22 Tg C yr⁻¹.

3.2 Process-based model estimates

3.2.1 Mean C budgets for 1990–1999 and 2000–2006

GPP estimated by the regional applications of process-based models over the arctic tundra region from 1990 through 1999 varies from 1755 Tg C yr⁻¹ (191 g C m⁻² yr⁻¹) for LPJ-Guess WHyMe to 5295 Tg C yr⁻¹ (577 g C m⁻² yr⁻¹) for Orchidee (Table 5). NPP is estimated to be approximately 65 %, 61 %, and 40 % of GPP by LPJ-Guess WHyMe, Orchidee, and TEM6, respectively. TEM6 estimates a higher proportion of GPP allocated to autotrophic respiration because the temperature sensitivity of autotrophic respiration in the model increases with decreasing mean annual temperature

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(McGuire et al., 1992). RH estimates vary from 875 TgC yr⁻¹ (95 gC m⁻² yr⁻¹) for TEM6 to 2954 TgC yr⁻¹ (322 gC m⁻² yr⁻¹) for Orchidee, and RH is estimated to be less than NPP by each model. NEP estimates vary from 85 TgC yr⁻¹ (10 gC m⁻² yr⁻¹) for TEM6 to 255 TgC yr⁻¹ (28 gC m⁻² yr⁻¹) for Orchidee. TEM estimates 30 TgC yr⁻¹ in fire emissions, which is 20 TgC yr⁻¹ more than is estimated by LPJ-Guess WHyMe. After accounting for fire emissions of CO₂, the estimates of the net exchange of CO₂ vary from 55 TgC yr⁻¹ taken up from the atmosphere by TEM6 to 255 TgC yr⁻¹ taken up by Orchidee, which is a higher range of uptake than is estimated by the global applications of process models (from null balance estimated by Hyland to an uptake of 188 TgC yr⁻¹ by LPJ-Guess). Most of the NEP estimated by TEM6 is lost to fire emissions, biogenic CH₄ emissions, and the export of harvested products and DOC. Approximately 20 % of NEP estimated by LPJ-Guess WHyMe is lost to fire and CH₄ emissions. Thus, NCB estimated by the models varies from approximately 20 TgC yr⁻¹ (TEM6) to 255 TgC yr⁻¹ (Orchidee).

In comparison to the 1990s, GPP estimated by the regional applications of process-based models over the Arctic tundra region from 2000 through 2006 is higher (from an increase of 9 gC m⁻² yr⁻¹ by TEM to an increase of 38 gC m⁻² yr⁻¹ by Orchidee; compare Table 6 to Table 5). The satellite-based estimate of GPP by TCF from 2000–2006 is 307 gC m⁻² yr⁻¹, which is 47 % and 14 % higher than the estimates by LPJ-Guess WHyMe and TEM6, respectively, and 50 % of the estimate by Orchidee. Similar to GPP, both NPP and RH estimates of the regional applications are higher in the 2000s compared to the 1990s. Although NEP estimates increase by 1 to 6 gC m⁻² yr⁻¹ between the two decades across the models, the increase from 1990 to 2006 is significant only for the LPJ-Guess WHyMe simulation (0.57 gC m⁻² yr⁻¹; *P* = 0.001). The TEM6 simulation estimates that fire emissions doubled in the 2000s compared to the 1990s. Estimates of the net uptake of CO₂ increase (lower or more negative net CO₂-C exchange) for both LPJ-Guess WHyMe and Orchidee in the 2000s compared to the 1990s, but decrease for TEM6. Between 1990 and 2006 both Orchidee and LPJ-Guess WHyMe estimate substantially greater net uptake of CO₂ (31 and 4 gC m⁻² yr⁻¹, respectively;

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 7) than TEM6 ($4 \text{ g C m}^{-2} \text{ yr}^{-1}$). Among the global models, only LPJ and SDGVM estimate less net uptake of CO_2 (less negative net $\text{CO}_2\text{-C}$ exchange) in the 2000s (Table 6) compared to the 1990s (Table 5). The greatest uptake between 1990 and 2006 among the global models is simulated by LPJ-Guess ($23 \text{ g C m}^{-2} \text{ yr}^{-1}$), which is similar to the uptake estimated by LPJ-Guess WHyMe (Table 7).

To explore the importance of changing climate on the regional applications of the process-based models, we conducted additional simulations that were driven by constant climate. We estimated the effect of a changing climate on NPP, RH, and NEP between 1990 and 2006 by subtracting the estimates of the constant climate simulation of each model from that of the corresponding transient climate simulation. This analysis indicated that climate change between 1990 and 2006 caused NPP and RH of all three models to increase (Table 8). In comparison to TEM6, NPP was 69 % and 106 % more sensitive in the LPJ-Guess WHyMe and Orchidee simulations, respectively. In contrast, the RH sensitivity of LPJ-Guess WHyMe was similar to that of TEM6, while the RH sensitivity of Orchidee was 146 % more sensitive than TEM6. The different sensitivities of NPP and RH caused quite different sensitivities in NEP. Climate change between 1990 and 2006 caused LPJ-Guess WHyMe NEP to increase by $8 \text{ g C m}^{-2} \text{ yr}^{-1}$, TEM6 NEP to increase by $1 \text{ g C m}^{-2} \text{ yr}^{-1}$, and Orchidee NEP to decrease by $4 \text{ g C m}^{-2} \text{ yr}^{-1}$.

3.2.2 Seasonal cycle and changes in the seasonal cycle

The shape of the seasonal cycle of NPP, RH, and NEP is similar among the regional applications of the process-based models (Fig. 3). In general, all models indicate that the growing season lasts from June through September with three of the models indicating that NPP starts to increase above winter values in May (LPJ-Guess WHyMe, Orchidee, TCF) and one of the models indicating a similar level of May production in October (Orchidee). In contrast to the other models, TCF is the only model that predicts positive, albeit very small, fluxes of NPP throughout the winter. All models estimate that the month of maximum production is July, followed closely by August. RH is generally

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



estimated to be small from November through April, to increase in magnitude from May through July/August, and to decrease in magnitude during September and October. All the models estimate that the month of maximum RH is July, except for LPJ-Guess WHyMe, for which August is characterized by the greatest RH. All models indicate that the maximum NEP occurs in July, but the number of months with positive NEP varies among the models between two (TCF) and four (LPJ-Guess WHyMe and Orchidee).

Both LPJ-Guess WHyMe and Orchidee have the same pattern of differences in monthly NEP between the 2000s and 1990s (Fig. 4), with the largest increases in July. LPJ-Guess WHyMe has relatively larger increases in May, while Orchidee has relatively larger increases in August. The summer increases of LPJ-Guess WHyMe and Orchidee are driven by increases in NPP that are greater than increases in RH. In contrast, TEM6 has the largest increases in September, followed by August as NPP increases in August and September are greater than increases in RH; in June and July there are similar increases in both NPP and RH. All three models indicate substantially lower NEP in October because of increases in RH when NPP is close to zero in both decades.

3.2.3 Interannual variability

Among the regional applications of the process-based models, Orchidee stands out as having the highest range of interannual variability in GPP, NPP, RH, and NEP (Fig. 5). Correlations are high among the models for interannual variability in the anomalies of GPP ($R = 0.73$ to 0.88 ; Fig. 5a) and RH ($R = 0.81$ to 0.97 ; Fig. 5c). The correlations for NPP anomalies (Fig. 5b) are slightly lower ($R = 0.66$ to 0.80), except for a low correlation between TCF and TEM6 ($R = 0.23$). In contrast, correlations among the anomalies of NEP (Fig. 5d) are poor and range from negative correlations ($R = -0.64$ between Orchidee and TCF) to low positive correlations ($R = 0.30$ between Orchidee and TEM6).

TEM6 estimates of fire emissions are characterized by substantial interannual variability in comparison to LPJ-Guess WHyMe, which has little interannual variability

(Fig. 6a); the variability is uncorrelated between the models ($R = 0.07$). LPJ-Guess estimates of biogenic CH_4 emissions are correlated with those of TEM6 ($R = 0.69$; Fig. 6b), but are characterized by more interannual variability than those of TEM6. The other flux anomalies estimated by TEM6 have less interannual variability ($\sim 0.25 \text{ g C m}^{-2}$; Fig. 6c) than fire emissions ($\sim 16 \text{ g C m}^{-2}$; Fig. 6a) and biogenic CH_4 emissions ($\sim 0.5 \text{ g C m}^{-2}$; Fig. 6b). In general, the correlations among the models for interannual variability in NCB (Fig. 6d) are similar to those for NEP (Fig. 6d), except that all of the correlations are weaker between TEM6 and the other models. This suggests that fire emissions, biogenic CH_4 emissions, and other export fluxes are important to consider in evaluating interannual variability in carbon storage of Arctic tundra.

3.3 Atmospheric inversion estimates

We analyzed the net exchange of CO_2 (i.e., NEE) between arctic tundra and the atmosphere estimated by inversions for 1990–1999 and 2000–2006 (Table 9). Among the three models that made estimates between 1990 and 1999, the mean annual exchange ranged from a source of 140 Tg C yr^{-1} ($15 \text{ g C m}^{-2} \text{ yr}^{-1}$) to a sink of 321 Tg C yr^{-1} ($35 \text{ g C m}^{-2} \text{ yr}^{-1}$). In comparison to the 1990–1999 time period, the range among the eight models that made estimates for 2000–2006 is wider and ranges from a source of 206 Tg C yr^{-1} ($22 \text{ g C m}^{-2} \text{ yr}^{-1}$) to a sink of 439 Tg C yr^{-1} ($48 \text{ g C m}^{-2} \text{ yr}^{-1}$).

The shape of the mean seasonal cycle of NEE between 2000 and 2006 is generally similar among the inversions (Fig. 7). All models indicate that the maximum NEE occurs in July, but the number of months with negative NEE varies among the models between two (C13_MATCH_rayner) and four (LSCE_an_v2.1). Among the inversion models, the NEE estimates of individual months are generally within 10 g C m^{-2} except for LSCE_an_v2.1, which estimates higher releases of CO_2 to the atmosphere than the other models in April and May and higher uptake of CO_2 in July, August, and September.

Among the inversion models, interannual variability is smallest for LSCE_an_v2.1 (standard deviation of NEE anomalies = $2.1 \text{ g C m}^{-2} \text{ yr}^{-1}$) and largest for rigc_patra

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(standard deviation = $13.1 \text{ g C m}^{-2} \text{ yr}^{-1}$) (Fig. 8). Similar to the correlations of interannual variability in NEP anomalies among the process-based models, the correlations of interannual variability in NEE anomalies among the inversion models is poor with a mean correlation of 0.03; correlations range between -0.38 (between CTRACKER_EU and C13_CCAM_low) to +0.99 (between CTRACKER_US and LSCE_an.v2.1).

4 Discussion

The changing C balance of Arctic tundra has been an issue of concern for several decades (Billings et al., 1983; Oechel et al., 1993; McGuire et al., 2000, 2009, 2010; Chapin et al., 2000; Sitch et al., 2007; Hayes et al., 2011). It has been hypothesized that tundra will become a source of C to the atmosphere because of C emissions associated with the warming of soil organic matter in the active layer as well as the exposure of previously frozen C to decomposition as the active layer deepens. Some recent model applications that consider soil C stocks at depth in high latitudes and the exposure of those stocks to decomposition upon permafrost thaw indicate that northern terrestrial ecosystems will release soil C to the atmosphere (Koven et al., 2011; Schaefer et al., 2011; Schneider von Deimling et al., 2012). It has also been hypothesized that tundra could become a sink for atmospheric CO_2 if N-limited plants in tundra regions take up a substantial proportion of N that is released by enhanced decomposition (Shaver et al., 1992). Some coupled climate-carbon model simulations predict that the northern high latitudes will serve as a substantial land carbon sink during the 21st century because both climate warming and elevated global $[\text{CO}_2]$ favor increased productivity and CO_2 uptake in the region (Friedlingstein et al., 2006; Qian et al., 2010; Sitch et al., 2008). Whether tundra becomes a source or a sink of atmospheric CO_2 in response to warming is an important scientific issue to resolve, as substantial source activity could compromise efforts to mitigate the increase of greenhouse gases in the atmosphere. Changes in CH_4 emissions are also important, because of the high global warming potential of CH_4 . In terms of climate forcing, increasing CH_4 emissions could

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



offset the effects of a CO₂ sink, or enhance the effects of a CO₂ source. In this study, we attempt to shed some light on these issues by analyzing the C balance of Arctic tundra through a synthetic comparison among estimates of CO₂ and CH₄ fluxes based on observations, regional and global applications of process-based models, and inversion models. We focused our comparison on the mean CO₂ and CH₄ budgets for the time periods 1990–1999 and 2000–2006, on aspects of the seasonal cycle of CO₂ exchange, and on interannual variability of CO₂ exchange.

4.1 Mean C budgets for the 1990s and 2000s

Table 10 compares the mean net exchanges of CO₂-C and CH₄-C from Arctic tundra to the atmosphere among observations, process-based models, and inversion models (see Appendix C in the Supplement for documentation of the estimates reported in Table 10). Syntheses of the compilation of flux observations and of inversion model results for Arctic tundra indicate that the annual exchange of CO₂ between Arctic tundra and the atmosphere has large uncertainties that cannot be distinguished from neutral balance. The synthesis of process-based model simulations indicate that Arctic has been acting as a sink for atmospheric CO₂, but based on the uncertainty estimates it is not clear if Arctic tundra acted as a weak or a strong sink. In comparison to the global process-based models, the regional process-based models indicate that Arctic tundra acted as a stronger sink.

Analysis of the CRU-NCEP data sets indicates that the region was 0.6 °C warmer in the 2000s compared to the 1990s. Most of the warming was in the autumn and winter (1.1 °C warmer) followed by summer (0.3 °C warmer), and little difference in the spring. The pattern of warmer autumns, winters, and summers in the 2000s occurred in all of the subregions except North America in which the mean summer temperature was not different between the two decades. It is notable that the central estimates of the observations, process-based models, and inversion models each identify stronger CO₂ sinks in the warmer 2000s compared with the 1990s. A stronger sink in the 2000s compared to the 1990s suggests that the efficiency of the tundra CO₂ sink is not currently

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



weakening; a common response of process-based models to warming in this region is that NPP increases faster in response to warming than RH (Sitch et al., 2007).

The largest changes in central estimates between the 1990s and 2000s are for those of the observations and the inversion models, and there is more convergence among the central estimates in the 2000s than in the 1990s. The large changes between decades for the observations and inversion models might reflect biases in the 1990s since the diversity of flux observations, and the number of CO₂ concentration measuring stations and inversion model applications were fewer than in the 2000s. The only mean source activity is that suggested by the observations in the 1990s. In particular, the source strength of the observations in North America in the 1990s is driving the overall source estimate for Arctic tundra in the 1990s. It is possible that tundra in North America was responding differently than tundra in other regions of the Arctic as northwestern North America warmed more strongly than other regions of the Arctic at the end of the last century (Serreze et al., 2000). It is also possible that sampling of observations in the 1990s was biased toward dry tundra, which tends to act as a source for C in the observations in both the 1990s and 2000s.

In general, the regional process-based model applications predict stronger sinks than the global applications. This is primarily due to the response of Orchidee, which has the highest NPP and NEE among the 11 models. It is notable that NPP and the sink strength of O-CN is much less than that of Orchidee, which might reflect the role of N in limiting productivity in O-CN. Although the sink strength of the central estimates of both regional and global process-based model applications increase from the 1990s and 2000s, 5 of the 11 process-based models show either no change or a weaker sink between decades (TEM6, CLM4CN, Hyland, LPJ, and SDGVM). Detailed analysis of one of the model applications (TEM6) suggests that tundra became a weaker sink from the 1970s through the 2000s because of the effects of climate on net ecosystem carbon balance (McGuire et al., 2010). The constant climate experiments we conducted with the regional model applications suggests that warming increases both NPP and RH in all three models, but that the relative responses of NPP and RH to warming

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are different among the models. This analysis indicates that the relative responses of NPP and RH to climate change are major sources of uncertainty in the application of process-based models to assess whether Arctic tundra will act as a positive or negative feedback to climate change.

5 Our analysis of CH₄ responses between decades is limited to comparison between the observations and the regional model applications (Table 10). There is substantial uncertainty in the 2000s CH₄ flux observation-based estimate. Because of big differences in estimates of CH₄ fluxes between wet and dry/mesic tundra (Table 4), uncer-
10 tainties could be reduced by adequately sampling CH₄ exchange between these two types of tundra within each of the subregions that we considered in scaling the CH₄ observations. The synthesis of observations produces central estimates in the 1990s (10 Tg CH₄-C yr⁻¹) and 2000s (20 Tg CH₄-C yr⁻¹) that are consistent with the range of uncertainty (23 to 75 Tg CH₄-C yr⁻¹) among observation-based and process-model es-
15 timates (McGuire et al., 2009) for northern high latitude terrestrial regions that include boreal forest in addition to tundra. The central estimates in Table 10 are also consistent with the range of uncertainty (11 to 38 Tg CH₄-C yr⁻¹) among CH₄ inversion models for northern high latitudes (McGuire et al., 2009). Similarly, the range of uncertainty of CH₄ exchange estimated by LPJ-Guess WHyMe and TEM6 are within the ranges of uncer-
20 tainty for both the bottom-up (based on observations and process-based models) and top-down (based on inversion models) reported by McGuire et al. (2009). The central estimates of the observations suggest that CH₄ emissions have increased more than 100 % from the 1990s to the 2000s, while the applications of LPJ-Guess WHyMe and TEM6 suggest that CH₄ emissions have only increased ~10–20 % from the 1990s to the 2000s. Previous analyses with TEM6 suggest that increasing temperature plays an
25 important role in this response (Zhuang et al., 2007; McGuire et al., 2010).

4.2 Seasonal cycle and changes in the seasonal cycle

The sink strength of Arctic tundra for CO₂ could increase between decades because of more net CO₂ uptake during the middle of the growing season or because of greater

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



net uptake of CO₂ at either the start or end of the growing season. In general, the pattern of mean seasonal cycle NEP of the regional model applications is consistent with the pattern of mean seasonal cycle NEE of the inversion models. All applications identify that the greatest CO₂ uptake occurs in July, but there are differences among both the regional model applications and inversions about the length of the net uptake period. Our analysis of the seasonal response of the regional model applications indicates that NPP of LPJ-Guess WHYMe, Orchidee, and TEM6 increased during the growing season in the 2000s compared to the 1990s. This is consistent with a number of remote sensing studies that have concluded that the Arctic tundra has become more productive during the last several decades (Nemani et al., 2003; Zhang et al., 2008; Beck and Goetz, 2011; Goetz et al., 2011) in association with warmer summers (Piao et al., 2011). However, the models show different patterns in the response of NEP between decades. Both LPJ-Guess and Orchidee show stronger patterns of early and mid-growing season NEP increases than TEM6, which has stronger NEP increases late in the growing season. A number of studies have concluded that the growing season in northern high latitude regions has increased in recent decades (McDonald et al., 2004; Euskirchen et al., 2006; Parmesan, 2007; Karlsen et al., 2009; Piao et al., 2011), although the increase varies both spatially and temporally. The TEM6 late season NEP response is consistent with a recent analysis for boreal Eurasia that suggests that the springtime extension of the growing season has stalled from 1997–2006 while the fall season has continued to lengthen and warm (Piao et al., 2011). The pattern of lower October NEP between decades among all three models is consistent with the analysis of Piao et al. (2008), which concludes that warmer and longer falls lead to greater CO₂ release in northern terrestrial ecosystems.

4.3 Interannual variability

Our analysis of inter-annual NEP anomalies among the regional model applications and of inter-annual NEE anomalies among the inversion models indicates that there is little agreement among the models on the pattern of inter-annual exchanges of CO₂

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



between Arctic tundra and the atmosphere. Although different inversions generally agree on the pattern inter-annual variability of regional NEE (Gurney et al., 2008), this is not the case for the Arctic tundra region and suggests that the inversions are not well constrained in this region. The regional applications of the process-models generally agree on patterns of inter-annual variability for GPP, NPP, and RH. The interannual patterns of GPP for LPJ-Guess WhyMe, Orchidee, and TEM6 are consistent with the satellite-based inter-annual variability of TCF, but the inter-annual variability of TEM6 NPP is not well correlated with that of TCF. Clearly, differences between GPP/NPP and RH need to be better constrained for the models to improve estimates of inter-annual variability.

4.4 Best estimates of carbon balance from 1990–2009

For the arctic tundra region, the use of observations, process-based models, and inversion models each have shortcomings with respect to estimating the net exchanges of CO₂ and CH₄ with the atmosphere in the 1990s and 2000s. Problems with observations include small sample size in comparison to the area being considered, biases in tundra types sampled in both space and time, different sampling technologies among the samples, and changes in sampling technology through time. Process-models have uncertainties with respect to conceptualization, formulation, and parameterization issues. Inversion models are not well constrained for the tundra regions. Given the shortcomings of these approaches, we decided to weight them equally in making estimates of net CO₂ and CH₄ exchange with the atmosphere for the time period from 1990–2009. For estimating CO₂ exchange, we first averaged the regional and global process-based model central estimates and high and low estimates of uncertainty for the 1990–2006 period in Table 10 and then average those estimates with the corresponding estimates for the observations and inversion models. This procedure results in an estimate of the net CO₂-C exchange of CO₂ between the atmosphere and tundra ecosystems of a 110TgCyr⁻¹ sink with an uncertainty range between a sink of 291TgCyr⁻¹ and a source of 80TgCyr⁻¹. For estimating CH₄ exchange, we averaged the central and

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the high and low estimates of uncertainty for the 1990–2006 period in Table 10 between the observations and the regional process-based models. This procedure results in an estimate of net CH₄-C exchange between the atmosphere and tundra ecosystems of a source of 19 Tg C yr⁻¹ with an uncertainty range between sources of 8 and 29 Tg C yr⁻¹.

5 Conclusions

The syntheses of the compilation of flux observations and of inversion model results for Arctic tundra in this study indicate that the annual exchange of CO₂ between Arctic tundra and the atmosphere has large uncertainties that cannot be distinguished from neutral balance in the 1990s and 2000s. In contrast, the synthesis of process-based model simulations indicate that Arctic tundra acted as a sink for atmospheric CO₂ in recent decades, but based on the uncertainty estimates it is not clear if Arctic tundra acted as a weak or a strong sink. Our analyses do suggest that both the CO₂ sink strength and the CH₄ emissions of Arctic tundra are increasing in recent decades. However, the analyses we conducted in this study indicate that responses of the seasonal exchange of CO₂ between decades and the interannual variability in CO₂ exchange of process-based models are not consistent. Although the regional models generally agree on patterns of inter-annual variability in production and decomposition, the constant climate experiments we conducted with the regional models indicates that the relative sensitivity of production and decomposition to climate change are different among the models. Thus, it is clearly important to reduce uncertainties in the observations, process-based models, and inversions in order to better understand the degree to which Arctic tundra is influencing atmospheric CO₂ and CH₄ concentrations. As inversion models are currently poorly constrained for making estimates of CO₂ exchange for Arctic tundra, there is a need to identify and place additional atmospheric CO₂ monitoring stations in a strategic fashion for better constraining inversion models. The availability and technology of ground-based observations in Arctic tundra are

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



improving, particularly through the implementation of the Arctic Observing Network that has been ongoing since the beginning of the International Polar Year in 2007 (Sorlin and Danell, 2008). However, it is important to improve the network so that observations can be effectively stratified into those for dry/mesic tundra vs. wet tundra so that regional estimates based on ground-based observations can be improved and uncertainties reduced. More importantly, observation networks need to be designed so that the observations can ultimately be synthesized to understand how and why the net annual and seasonal exchanges of CO₂ and CH₄ are changing in response to climate variability and change in different tundra types that span hydrological variability. Also, the effects of disturbances such as fire and thermokarst on the exchange of CO₂ and CH₄ are not well represented in observation networks. Information from enhanced observation networks needs to be effectively transferred into process-based models to improve the simulation of CO₂ and CH₄ exchange so that process-based models can more reliably assess whether Arctic tundra will act as to amplify or mitigate global climate change.

Supplementary material related to this article is available online at:

<http://www.biogeosciences-discuss.net/9/4543/2012/bgd-9-4543-2012-supplement.pdf>

Acknowledgements. Support for this study was provided by the National Science Foundation, the US Geological Survey, the US Fish and Wildlife Service, the US Department of Energy Office of Science (Biological and Environmental Research), the US National Aeronautics and Space Administration Terrestrial Ecology program, the European Union RTN GREENCYCLES II network, the Swedish Research Council VR and FORMAS, the Greenland Climate Research Center, the Mistra Swedish Research Programme for Climate, Impacts and Adaptation, the Lund University research programme Modelling the Regional and Global Earth System, the European Union 6th Framework CARBO-North project, and the Lund University Centre for Studies of Carbon Cycle and Climate Interactions. The data for the global process-model sim-

BGD

9, 4543–4594, 2012

**An assessment of
the carbon balance of
arctic tundra**

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ulations was provided by the Trendy multi-model evaluation project (<http://dgvm.ceh.ac.uk>), as part of the Global Carbon Project.

References

- 5 Beck, P. S. A and Goetz, S. J.: Satellite observations of high northern latitude vegetation productivity changes between 1982 and 2008: ecological variability and regional differences, *Environ. Res. Lett.*, 6, 045501, doi:10.1088/1748-9326/6/4/045501, 2011.
- Billings, W. D., Loken, J. O., Mortensen, D. A., and Peterson, K. M.: Increasing atmospheric carbon dioxide: possible effects on arctic tundra, *Oecologia*, 58, 286–289, 1983.
- 10 Bonan, G. B. and Levis, S.: Quantifying carbon-nitrogen feedbacks in the Community Land Model (CLM4), *Geophys. Res. Lett.*, 37, L07401, doi:10.1029/2010GL042430, 2010.
- Callaghan, T. V., Björn, L. O., Chapin III, F. S., Chernov, Y., Christensen, T. R., Huntley, B., Ims, R. A., Johansson, M., Jolly, D., Jonasson, S., Matveyeva, N., Oechel, W. C., Panikov, N., Shaver, G. R., Elster, J., Henttonen, H., Jónsdóttir, I. S., Laine, K., Schaphoff, S., Sitch, S., Taulavuori, E., Taulavuori, K., and Zöckler, C.: Arctic tundra and polar desert ecosystems, in: Arctic climate impact assessment (ACIA), Cambridge University Press, Cambridge, UK, 243–352, 2005.
- 15 Canadell, J. G., Ciais, P., Gurney, K., Le Quere, C., Piao, S., Raupach, M. R., and Sabine, C. L.: An international effort to quantify regional carbon fluxes, *Eos*, 92, 81–82, 2011.
- Chapin III, F. S., McGuire, A. D., Randerson, J., Pielke Sr., R., Baldocchi, D., Hobbie, S. E., Roulet, N., Eugster, W., Kasischke, E., Rastetter, E. B., Zimov, S. A., Oechel, W. C., and Running, S. W.: Feedbacks from arctic and boreal ecosystems to climate, *Glob. Change Biol.*, 6, S211–S223, 2000.
- 20 Chapin III, F. S., Woodwell, G. M., Randerson, J. T., Lovett, G. M., Rastetter, E. B., Baldocchi, D. D., Clark, D. A., Harmon, M. E., Schimel, D. S., Valentini, R., Wirth, C., Aber, J. D., Cole, J. J., Goulden, M. L., Harden, J. W., Heimann, M., Howarth, R. W., Matson, P. A., McGuire, A. D., Melillo, J. M., Mooney, H. A., Neff, J. C., Houghton, R. A., Pace, M. L., Ryan, M. G., Running, S. W., Sala, O. E., Schlesinger, W. H., and Schulze, E.-D.: Reconciling carbon-cycle concepts, terminology, and methods, *Ecosystems*, 9, 1041–1050, 2006.
- 25

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chapin III, F. S., Randerson, J. T., McGuire, A. D., Foley, J. A., and Field, C. B.: Changing feedbacks in the climate-biosphere system, *Front. Ecol. Environ.*, 6, 313–320, doi:10.1890/080005, 2008.

Cox, P. M.: Description of the “TRIFFID” dynamic global vegetation model, Hadley Centre Technical Note 24, 2001.

Euskirchen, E. S., McGuire, A. D., Kicklighter, D. W., Zhuang, Q., Clein, J. S., Dargaville, R. J., Dye, D. G., Kimball, J. S., McDonald, K. C., Melillo, J. M., Romanovsky, V. E., and Smith, N. V.: Importance of recent shifts in soil thermal dynamics on growing season length, productivity, and carbon sequestration in terrestrial high-latitude ecosystems, *Glob. Change Biol.*, 12, 731–750, 2006.

Farquhar, G. D., von Caemmerer, S., and Berry, J. A.: A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species, *Planta*, 149, 78–90, 1980.

Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate-carbon cycle feedback analysis: Results from the C4MIP model intercomparison, *J. Climate*, 19, 3337–3353, 2006.

Goetz, S. J., Epstein, H. E., Alcaraz, D., Beck, P., Bhatt, U., Bunn, A. G., Comiso, J., Jia, G. J., Kaplan, J. O., Lischke, H., Lloyd, A. H., Yu, Q., and Walker, D. A.: Recent changes in arctic vegetation: satellite observations and simulation model predictions, in: *Eurasian Arctic Land Cover and Land Use in a Changing Climate*, edited by: Gutman, G. and Reissell, A., Springer, ISBN 978-90-481-9117-8, 2011.

Gurney, K. R.: Global atmospheric carbon budget, *Biogeosciences Discuss.*, in preparation, 2012.

Gurney, K. R., Baker, D., Rayner, P., Denning, S., Law, R., Bousquet, P., Bruhwiler, L., Chen, Y. H., Ciais, P., Fung, I., Heimann, M., John, J., Maki, T., Maksyutov, S., Peylin, P., Prather, M., Pak, B., and Taguchi, S.: Interannual variations in continental-scale net carbon exchange and sensitivity to observing networks estimated from atmospheric CO₂ inversions for the period 1980–2005, *Global Biogeochem. Cy.*, 22, GB3025, doi:10.1029/2007GB003082, 2008.

Haxeltine, A. and Prentice, I. C.: A general model for the light-use efficiency of primary production, *Funct. Ecol.*, 10, 551–561, 2006.

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hayes, D. J., McGuire, A. D., Kicklighter, D. W., Gurney, K. R., Burnside, T. J., and Melillo, J. M.: Is the northern high latitude land-based CO₂ sink weakening?, *Global Biogeochem. Cy.*, 25, GB3018, doi:10.1029/2010GB003813, 2011.

Hickler, T., Vohland, K., Feehan, J., Miller, P. A., Smith, B., Costa, L., Giesecke, T., Fronzek, S., Cramer, W., and Sykes, M.: Projecting the future distribution of European potential natural vegetation with a generalized tree species-based dynamic vegetation model, *Global Ecol. Biogeogr.*, 21, 50–63, 2012.

Hinzman, L. D., Bettez, N. D., Bolton, W. R., Chapin, F. S., Dyurgerov, M. B., Fastie, C. L., Griffith, B., Hollister, R. D., Hope, A., Huntington, H. P., Jensen, A. M., Jia, G. J., Jorgenson, T., Kane, D. L., Klein, D. R., Kofinas, G., Lynch, A. H., Lloyd, A. H., McGuire, A. D., Nelson, F. E., Nolan, M., Oechel, W. C., Osterkamp, T. E., Racine, C. H., Romanovsky, V. E., Stone, R. S., Stow, D. A., Sturm, M., Tweedie, C. E., Vourlitis, G. L., Walker, M. D., Walker, D. A., Webber, P. J., Welker, J. M., Winker, K. S., and Yoshikawa, K.: Evidence and implications of recent climate change in northern Alaska and other Arctic regions, *Climatic Change*, 72, 251–298, 2005.

Karlsen, S. R., Høgda, K. A., Wielgolaski, F. E., Tolvanen, A., Tømmervik, H., Poikolainen, J., and Kubin, E.: Growing-season trends in Fennoscandia 1982–2006, determined from satellite and phenology data, *Clim. Res.*, 39, 275–286, 2009.

Kim, Y., Kimball, J. S., Zhang, K., and McDonald, K. C.: Satellite detection of increasing northern hemisphere non-frozen seasons from 1979 to 2008: Implications for regional vegetation growth, *Remote Sens. Environ.*, 12, 472–487, 2012.

Kimball, J. S., Jones, L. A., Zhang, K., Heinsch, F. A., McDonald, K. C., and Oechel, W. C.: A satellite approach to estimate land-atmosphere CO₂ exchange for Boreal and Arctic biomes using MODIS and AMSR-E, *IEEE T. Geosci. Remote Sens.*, 47, 569–587, doi:10.1109/TGRS.2008.2003248, 2009.

Koven, C., Friedlingstein, P., Ciais, P., Khvorostyanov, D., Krinner, G., and Tarnocai, C.: On the formation of high-latitude soil carbon stocks: Effects of cryoturbation and insulation by organic matter in a land surface model, *Geophys. Res. Lett.*, 36, L21501, doi:10.1029/2009GL040150, 2009.

Koven, C. D., Ringeval, B., Friedlingstein, P., Ciais, P., Cadule, P., Khvorostyanov, D., Krinner, G., and Tarnocai, C.: Permafrost carbon-climate feedbacks accelerate global warming, *P. Natl. Acad. Sci.*, 108, 14769–14774, doi:10.1073/pnas.1103910108, 2011.

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, Z.-L., Levis, S., Sakaguchi, K., Bonan, G. B., and Slater, A. G.: Parameterization improvements and functional and structural advances in version 4 of the Community Land Model, *Journal of Advances in Modeling Earth Systems*, 3, M03001, doi:10.1029/2011MS000045, 2011.

Levy, P. E., Cannell, M. G. R., and Friend, A. D.: Modelling the impact of future changes in climate, CO₂ concentration and land use on natural ecosystems and the terrestrial carbon sink, *Global Environ. Chan.*, 14, 21–30, 2004.

McDonald, K. C., Kimball, J. S., Njoku, E., Zimmermann, R., and Zhao, M.: Variability in spring-time thaw in the terrestrial high latitudes: Monitoring a major control on the biospheric assimilation of atmospheric CO₂ with spaceborne microwave remote sensing, *Earth Interact.*, 8, 23 pp., 2004.

McGuire, A. D., Melillo, J. M., Joyce, L. A., Kicklighter, D. W., Grace, A. L., Moore III, B., and Vörösmarty, C. J.: Interactions between carbon and nitrogen dynamics in estimating net primary productivity for potential vegetation in North America, *Global Biogeochem. Cy.*, 6, 101–124, 1992.

McGuire, A. D., Melillo, J. M., Kicklighter, D. W., Pan, Y., Xiao, X., Helfrich, J., Moore III, B., Vorosmarty, C. J., and Schloss, A. L.: Equilibrium responses of global net primary production and carbon storage to doubled atmospheric carbon dioxide: Sensitivity to changes in vegetation nitrogen concentration, *Global Biogeochem. Cy.*, 11, 173–189, 1997.

McGuire, A. D., Clein, J. S., Melillo, J. M., Kicklighter, D. W., Meier, R. A., Vorosmarty, C. J., and Serreze, M. C.: Modelling carbon responses of tundra ecosystems to historical and projected climate: sensitivity of pan-Arctic carbon storage to temporal and spatial variation in climate, *Glob. Change Biol.*, 6, S141–S159, 2000.

McGuire, A. D., Anderson, L. G., Christensen, T. R., Dallimore, S., Guo, L., Hayes, D. J., Heimann, M., Lorenson, T. D., Macdonald, R. W., and Roulet, N.: Sensitivity of the carbon cycle in the Arctic to climate change, *Ecol. Monogr.*, 79, 523–555, 2009.

McGuire, A. D., Hayes, D. J., Kicklighter, D. W., Manizza, M., Zhuang, Q., Chen, M., Follows, M. J., Gurney, K. R., McClelland, J. W., Melillo, J. M., Peterson, B. J., and Prinn, R.: An analysis of the carbon balance of the Arctic Basin from 1997 to 2006, *Tellus*, 62B, 455–474, doi:10.1111/j.1600-0889.2010.00497.x, 2010.

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nemani, R. R., Keeling, C. D., Hashimoto, H., Jolly, W. M., Piper, S. C., Tucker, C. J., Myneni, R. B., and Running, S. W.: Climate-driven increases in global terrestrial net primary production from 1982 to 1999, *Science*, 300, 1560–1563, 2003.

New, M., Hulme, M., and Jones, P. D.: Representing twentieth-century space-time climate variability. Part I: development of a 1961–90 mean monthly terrestrial climatology, *J. Climate*, 12, 829–856, 1999.

Oechel, W. C., Hastings, S. J., Vourlitis, G., Jenkins, M., Riechers, G., and Grulke, N.: Recent change of Arctic tundra ecosystems from a net carbon dioxide sink to a source, *Nature*, 361, 520–523, 1993.

Parmesan, C.: Influence of species, latitudes, and methodologies on estimates of phenological response to global warming, *Glob. Change Biol.*, 13, 1860–1872, 2007.

Piao, S., Ciais, P., Friedlingstein, P., Peylin, P., Reichstein, M., Luysaert, S., Margolis, H., Fang, J., Barr, A., Chen, A., Grelle, A., Hollinger, D. Y., Laurilla, T., Lindroth, A., Richardson, A. D., and Vesala, T.: Net carbon dioxide losses of northern ecosystems in response to autumn warming, *Nature*, 451, 49–52, doi:10.1038/nature06444, 2008.

Piao, S., Wang, X., Ciais, P., Zhu, B., Wang, T., and Liu, J.: Changes in satellite-derived vegetation growth trend in temperate and boreal Eurasia from 1982–2006, *Glob. Change Biol.*, 17, 3228–3239, doi:10.1111/j.1365-2486.2011.02419.x, 2011.

Ping, C. L., Michaelson, G. J., Jorgenson, M. T., Kimble, J. M., Epstein, H., Romanovsky, V. E., and Walker, D. A.: High stocks of soil organic carbon in the North American Arctic region, *Nat. Geosci.*, 1, 615–619, 2008.

Post, E., Forchhammer, M. C., Bret-Harte, S., Callaghan, T. V., Christensen, T. R., Elberling, B., Fox, A. D., Gilg, O., Hik, D. S., Hoyer, T. T., Ims, R. A., Jeppesen, E., Klein, D. R., Madsen, J., McGuire, A. D., Rysgaard, S., Schindler, D. E., Stirling, I., Tamstorf, M. P., Tyler, N. J. C., van der Wal, R., Welker, J., Wookey, P. A., and Aastrup, P.: Ecological dynamics across the Arctic associated with recent climate change, *Science*, 325, 1355–1358, 2009.

Poutou, E., Krinner, G., Genthon, C., and de Noblet-Ducoudre, N.: Role of soil freezing in future boreal climate change, *Clim. Dynam.*, 23, 621–639, 2004.

Qian, H., Renu, J., and Zeng, N.: Enhanced terrestrial carbon uptake in the northern high latitudes in the 21st century from the coupled carbon cycle climate model intercomparison project model projections, *Glob. Change Biol.*, 16, 641–656, doi:10.1111/j.1365-2486.2009.01989.x, 2010.

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rawlins, M. A., Steele, M., Holland, M. M., Adam, J. C., Cherry, J. E., Francis, J. A., Groisman, P. Y., Hinzman, L. D., Huntington, T. G., Kane, D. L., Kimball, J. S., Kwok, R., Lammers, R. B., Lee, C. M., Lettenmaier, D. P., McDonald, K. C., Podest, E., Pundsack, J. W., Rudels, B., Serreze, M. C., Shiklomanov, A., Skagseth, Ø., Troy, T. J., Vörösmarty, C. J., Wensnahan, M., Wood, E. F., Woodgate, R., Yang, D., Zhang, K., and Zhang, T.: Analysis of the Arctic system for freshwater cycle intensification: Observations and expectations, *J. Climate*, 23, 5716–5737; doi:10.1175/2010JCLI3421.1, 2010.

Rowland, J. C., Jones, C. E., Altmann, G., Bryan, R., Crosby, B. T., Geernaert, G. L., Hinzman, L. D., Kane, D. L., Lawrence, D. M., Mancino, A., Marsh, P., McNamara, J. P., Romanovsky, V. E., Toniolo, H., Travis, J., Trochim, E., and Wilson, C. J.: Arctic landscapes in transition: Responses to thawing permafrost, *Eos*, 91, 229–230, 2010.

Running, S. W., Nemani, R. R., Heinsch, F. A., Zhao, M., Reeves, M., and Hashimoto, H.: A continuous satellite-derived measure of global terrestrial primary productivity: Future science and applications, *Bioscience*, 56, 547–560, 2004.

Schaefer, K., Zhang, T., Bruhwiler, L., and Barrett, A. P.: Amount and timing of permafrost carbon release in response to climate warming, *Tellus* 63B, 165–180, 2011.

Schneider von Deimling, T., Meinshausen, M., Levermann, A., Huber, V., Frieler, K., Lawrence, D. M., and Brovkin, V.: Estimating the near-surface permafrost-carbon feedback on global warming, *Biogeosciences*, 9, 649–665, doi:10.5194/bg-9-649-2012, 2012.

Schuur, E., Bockheim, J., Canadell, J., Euskirchen, E., Field, C., Goryachkin, S., Hagemann, S., Kuhry, P., Lafleur, P., and Lee, H.: Vulnerability of permafrost carbon to climate change: Implications for the global carbon cycle, *Bioscience*, 58, 701–714, 2008.

Schuur, E. A. G., Abbott, B. W., Bowden, W. B., Brovkin, V., Camill, P., Canadell, J. P., Chapin III, F. S., Christensen, T. R., Chanton, J. P., Ciais, P., Crill, P. M., Crosby, B. T., Czimczik, C. I., Grosse, G., Hayes, D. J., Hugelius, G., Jastrow, J. D., Kleinen, T., Koven, C. D., Krinner, G., Kuhry, P., Lawrence, D. M., Natali, S. M., O'Donnell, J. A., Ping, C. L., Rinke, A., Riley, W. J., Romanovsky, V. E., Sannel, A. B. K., Schädel, C., Schaefer, K., Subin, Z. M., Tarnocai, C., Turetsky, M., Walter-Anthony, K. M., Wilson, C. J., and Zimov, S. A.: High risk of permafrost thaw, *Nature*, 480, 32–33, 2011.

Serreze, M. C., Walsh, J. E., Chapin III, F. S., Osterkamp, T., Dyurgerov, M., Romanovsky, V., Oechel, W. C., Morison, J., Zhang, T., and Barry, R. G.: Observational evidence of recent change in the northern high-latitude environment, *Climatic Change*, 46, 159–207, 2000.

- Serreze, M. C., Bromwich, D. H., Clark, M. P., Etringer, A. J., Zhang, T., and Ammers, R.: Large-scale hydro-climatology of the terrestrial Arctic drainage system, *J. Geophys. Res.*, 108, 8160, doi:10.1029/2001JD000919, 2003.
- Shaver, G. R., Billings, W. D., Chapin, F. S., Giblin, A. E., Nadelhoffer, K. J., Oechel, W. C., and Rastetter, E. B.: Global change and the carbon balance of Arctic ecosystems, *Bioscience*, 42, 433–441, 1992.
- Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J., Levis, S., Lucht, W., Sykes, M., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ Dynamic Global Vegetation Model, *Glob. Change Biol.*, 9, 161–185, 2003.
- Sitch, S., McGuire, A. D., Kimball, J., Gedney, N., Gamon, J., Engstrom, R., Wolf, A., Zhuang, Q., Clein, J. S., and McDonald, K. C.: Assessing the carbon balance of circumpolar arctic tundra using remote sensing and process modeling, *Ecol. Appl.*, 17, 213–234, 2007.
- Sitch, S., Huntingford, C., Gedney, N., Levy, P. E., Lomas, M., Piao, S. L., Betts, R., Ciais, P., Cox, P., and Friedlingstein, P.: Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs), *Glob. Change Biol.*, 14, 2015–2039, 2008.
- Smith, B., Prentice, I. C., and Sykes, M. T.: Representation of vegetation dynamics in modelling of terrestrial ecosystems: Comparing two contrasting approaches within European climate space, *Global Ecol. Biogeogr.*, 10, 621–637, 2001.
- Sörlin, S. and Danell, K.: 1st IPY Workshop on Sustaining Arctic Observing Networks: Workshop Report, available at: http://www.arcticobserving.org/images/stories/workshop_report/SAON1_report_total.pdf, 2008.
- Snow, Water, Ice, and Permafrost in the Arctic (SWIPA) Assessment: Executive Summary, Arctic Monitoring and Assessment Program (AMAP) Secretariat, Oslo, Norway, 16 pp., available at: www.amap.no, 2011.
- Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil organic carbon pools in the northern circumpolar permafrost region, *Global Biogeochem. Cy.*, 23, GB2023, doi:10.1029/2008GB003327, 2009.
- Thornton, P. E., Lamarque, J. F., Rosenbloom, N. A., and Mahowald, N. M.: Influence of carbon-nitrogen cycle coupling on land model response to CO₂ fertilization and climate variability, *Global Biogeochem. Cy.*, 21, GB4018, doi:10.1029/2006GB002868, 2007.

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Thornton, P. E., Doney, S. C., Lindsay, K., Moore, J. K., Mahowald, N., Randerson, J. T., Fung, I., Lamarque, J.-F., Feddes, J. J., and Lee, Y.-H.: Carbon-nitrogen interactions regulate climate-carbon cycle feedbacks: results from an atmosphere-ocean general circulation model, *Biogeosciences*, 6, 2099–2120, doi:10.5194/bg-6-2099-2009, 2009.

5 Vlassova, T. K.: Human Impacts on the Tundra-Taiga Zone Dynamics: The case of the Russian Lesotundra, *Ambio Special Report (Tundra-Taiga Treeline Research)*, 12, 30–36, 2002.

Wania, R., Ross, I., and Prentice, I. C.: Integrating peatlands and permafrost into a dynamic global vegetation model: I. Evaluation and sensitivity of physical land surface processes, *Global Biogeochem. Cy.*, 23, GB3014, doi:10.1029/2008GB003412, 2009a.

10 Wania, R., Ross, I., and Prentice, I. C.: Integrating peatlands and permafrost into a dynamic global vegetation model: II. Evaluation and sensitivity of vegetation and carbon cycle processes, *Global Biogeochem. Cy.*, 23, GB015, doi:10.1029/2008GB003413, 2009b.

Wania, R., Ross, I., and Prentice, I. C.: Implementation and evaluation of a new methane model within a dynamic global vegetation model: LPJ-WHyMe v1.3.1, *Geosci. Model Dev.*, 3, 565–584, doi:10.5194/gmd-3-565-2010, 2010.

15 Woodward, F. I. and Lomas, M. R.: Vegetation dynamics: simulating responses to climate change, *Biol. Rev.*, 79, 643–670, 2004.

Woodward, F. I., Smith, T. M., and Emanuel, W. R.: A global land primary productivity and phytogeography model, *Global Biogeochem. Cy.*, 9, 471–490, 1995.

20 Zaehle, S. and Friend, A. D.: Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model description, site-scale evaluation and sensitivity to parameter estimates, *Global Biogeochem. Cy.*, 24, GB1005, doi:10.1029/2009GB003521, 2010.

Zhang, K., Kimball, J. S., Hogg, E. H., Zhao, M., Oechel, W. C., Cassano, J. J., and Running, S. W.: Satellite-based model detection of recent climate driven changes in northern high latitude vegetation productivity, *J. Geophys. Res.*, 113, G03033, doi:10.1029/2007JG000621, 2008.

25 Zhao, M. and Running, S. W.: Drought-induced reduction in global terrestrial net primary production from 2000–2009, *Science*, 329, 940–943, 2010.

Zhuang, Q., Romanovsky, V. E., and McGuire, A. D.: Incorporation of a permafrost model into a large-scale ecosystem model: Evaluation of temporal and spatial scaling issues in simulating soil thermal dynamics, *J. Geophys. Res.-Atmos.*, 106, 33649–33670, 2001.

30 Zhuang, Q., McGuire, A. D., Melillo, J. M., Clein, J. S., Dargaville, R. J., Kicklighter, D. W., Myneni, R. B., Dong, J., Romanovsky, V. E., Harden, J., and Hobbie, J. E.: Carbon cycling in

extratropical terrestrial ecosystems of the Northern Hemisphere during the 20th century: a modeling analysis of the influences of soil thermal dynamics, *Tellus*, 55B, 751–776, 2003.

Zhuang, Q., Melillo, J. M., Kicklighter, D. W., Prinn, R. G., McGuire, A. D., Steudler, P. A., Felzer, B. S., and Hu, S.: Methane fluxes between terrestrial ecosystems and the atmosphere at northern high latitudes during the past century: A retrospective analysis with a process-based biogeochemistry model, *Global Biogeochem. Cy.*, 18, GB3010, doi:10.1029/2004GB002239, 2004.

Zhuang, Q., Melillo, J. M., McGuire, A. D., Kicklighter, D. W., Prinn, R. G., Steudler, P. A., Felzer, B. S., and Hu, S.: Net emissions of CH₄ and CO₂ in Alaska: Implications for the region's greenhouse gas budget, *Ecol. Appl.*, 17, 203–212, 2007.

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. A summary of technical performance in various categories of different flux measurement techniques. Modified from Drösler et al. (unpublished data).

	Eddy covariance	Automatic chambers	Manual chambers
Steady state, undisturbed measurement	++	±	–
Integration over spatial variability	++	– (quantity of chambers vs. mosaic)	–(quantity of chambers vs. mosaic)
Direct measurement of small scale spatial variability	–	+	++
Tracking temporal variability	++	++	– (campaigns as basis for modelling)
Costs	–	–	++
Workload	++	+	–
Performance under all climate conditions	±	±	++

Table 2. Description of process-based models compared in this study. See Appendix B in the Supplement for additional details.

	Model Description	LPJ-Guess WWhyMe	Orchidee	TCF	TEM6
	Recent Arctic Applications	Smith et al. (2001) Wania et al. (2009a) Wania et al. (2009b) Wania et al. (2010)	Koven et al. (2009) Koven et al. (2011)	Kimball et al. (2009)	Hayes et al. (2011) McGuire et al. (2010)
Scale					
Temporal	Resolution Application	Daily Daily to Century	30 min Diurnal to multi-seasonal (Century application)	Daily Daily to Decadal	Month Seasonal to Century
Spatial	Resolution Application	0.5° Plot – Globe	2° Plot – Globe	25 km Plot – Region	0.5° Plot – Globe
Structure					
Vegetation		Dynamic	Prescribed	Prescribed	Prescribed
Litter/Soil Pools		1/3	4/3	3	2
Microbial Pools		No	No	No	No
Processes					
General	Photosynthesis	Farquhar et al. (1980) modified by Haxeltine and Prentice (1996)	Farquhar et al. 1980	MODIS MOD17 production efficiency model (Running et al., 2004)	GPP based on multiple limiting factors (see McGuire et al., 1997)
	Heterotrophic Respiration (RH)	Dependent on soil carbon, soil temperature and moisture (Smith et al., 2001). Follows Wania et al. (2009a, b) for peatlands.	Q_{10} of 2 with respect to each soil layer temperature for unfrozen soil layers; linear drop off to 0 RH between 0 and -2°C	Dependent on surface (<10cm) soil organic carbon, surface soil temperature and moisture (Kimball et al., 2009)	Dependent on soil organic carbon, soil moisture, soil temperature (see Zhuang et al., 2003)
	Fire	Yes. On upland soils only, not in peatlands.	No	No	Yes
	C:N dynamics	Optimal N allocation to canopy assumed (Haxeltine and Prentice, 1996)	No	No	Yes
	Especially Relevant to Arctic Tundra Ecosystems	Non-Vascular Plants (Mosses/Lichens)	Yes, but mosses only represented in peatlands	No	No
	Permafrost	Freeze-thaw processes as described in Wania et al. (2009a, b)	Freeze/thaw processes described in Poutou et al. (2004); permafrost carbon pools described in Koven et al. (2009)	No	Yes, see Zhuang et al. (2001, 2003) and Hayes et al. (2011)
	Lateral Hydrology	No	No	No	No
	Methane	Yes, as described in Wania et al. (2010)	No	No	Yes, see Zhuang et al. (2004, 2007)

BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 3. Summary of Observationally Based Estimates of Mean Net CO₂-C and CH₄-C Exchange from Arctic Tundra to the Atmosphere (g C m⁻² season⁻¹) for Different Subregions.

Time Period	North America	North Atlantic	Northern Europe	Eurasia
CO ₂ Exchange				
Summer				
1990–1999	-7 (52 ¹ ; -22 to 7 ²)	-32 (9; -58 to -5)	-98 (4; -127 to -68)	-25 (18; -50 to -1)
2000–2009	-18 (28; -39 to 3)	-53 (12; -90 to -16)	-92 (5; -220 to 36)	-73 (12; -141 to -4)
1990–2009	-11 (80; -23 to 1)	-44 (21; -66 to -21)	-94 (9; -151 to -38)	-44 (30; -74 to -14)
Winter				
1990–2009	31 (9; 15 to 47)	-	41 (3; 11 to 71)	29 (2; -13 to 73)
Annual				
1990–1999	29 (9; 2 to 57)	-	-25 (6; -63 to -14)	-
2000–2009	-3 (14; -28 to 21)	-	-19 (33; -30 to -8)	-
1990–2009	10 (23; -10 to 28)	-68 (4; -213 to 78)	-20 (39; -30 to -10)	-82 (6; -134 to -30)
CH ₄ Exchange				
Summer				
1990–1999	2.4 (10; 0.7 to 4.2)	-	7.8 (13; 1.4 to 14.2)	3.0 (15; -0.4 to 6.5)
2000–2009	1.4 (9; -0.2 to 2.9)	-	12.5 (10; 1.9 to 23.0)	5.1 (29; 1.6 to 8.6)
1990–2009	1.9 (19; 0.8 to 3.0)	3.1 (3; 2.3 to 3.8)	9.9 (23; 4.5 to 15.2)	4.4 (44; 1.9 to 6.9)
Annual				
1990–1999	4.4 (24; 1.4 to 7.4)	-	15.0 (3; -18.3 to 48.3)	-
2000–2009	16.9 (2; -12.0 to 45.7)	-	11.3 (27; 6.6 to 16.0)	8.2 (7; -1.8 to 18.1)
1990–2009	5.4 (26; 1.9 to 8.9)	-	11.7 (30; 7.2 to 16.1)	8.2 (7; -1.8 to 18.1)

¹ Number of site-year estimates, ² 95 % confidence interval



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Table 4. Summary of Observationally Based Estimates of Mean Net CO₂-C and CH₄-C Exchange from Arctic Tundra to the Atmosphere (g C m⁻² season⁻¹) for Different Tundra Types.

Time Period	Wet Tundra	Dry/Mesic Tundra
CO₂ Exchange		
Summer	-43 (45 ¹ ; -27 to -59 ²)	5 (46; -11 to 21)
Winter	31 (3; 1 to 61)	31 (7; 11 to 51)
Annual	-26 (27; -15 to -37)	10 (12; -27 to 47)
CH₄ Exchange		
Summer	9.2 (38; 5.4 to 13.0)	0.8 (25; 0.3 to 1.4)
Annual	14.6 (22; 8.5 to 20.2)	2.3 (24; 0.3 to 4.3)

¹ Number of site-year estimates, ² 95% confidence interval

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Table 5. Mean C Budget of Arctic Tundra Simulated by Process Models for 1990–1999.

Model	GPP	NPP	RH	NEP	FIREC	CO ₂ *	BIOCH4	OTHER	NCB
g C m⁻² yr⁻¹									
Regional Apps.									
LPJ-G WHyMe	191	124	102	22	1	-21	4	-	17
Orchidee	577	350	322	28	-	-28	-	-	28
TEM6	261	105	95	10	4	-6	2	2	2
Global Apps.									
CLM4C	101	31	30	1	1	0	-	-	0
CLM4CN	-	-	-	1	-	-1	-	-	1
Hyland	23	11	11	0	-	0	-	-	0
LPJ	464	291	253	38	18	-20	-	-	20
LPJ-Guess	490	306	265	41	20	-21	-	-	21
O-CN	64	28	27	1	-	-1	-	-	1
SDGVM	-	-	-	23	5	-18	-	-	18
TRIFFID	332	208	200	8	-	-8	-	-	-8
Tg C yr⁻¹									
Regional Apps.									
LPJ-G WHyMe	1755	1133	934	199	10	-189	34	-	155
Orchidee	5295	3209	2954	255	-	-255	-	-	255
TEM6	2391	960	875	85	30	-55	15	20	20
Global Apps.									
CLM4C	929	289	275	14	8	-6	-	-	6
CLM4CN	-	-	-	5	-	-5	-	-	5
Hyland	210	105	105	0	-	0	-	-	0
LPJ	4255	2671	2323	348	169	-179	-	-	179
LPJ-Guess	4497	2804	2433	371	183	-188	-	-	188
O-CN	585	256	252	4	-	-4	-	-	4
SDGVM	-	-	-	212	44	-168	-	-	168
TRIFFID	3053	1907	1837	70	-	-70	-	-	70

* Net CO₂-C Exchange, positive sign indicates source to the atmosphere and negative sign indicates tundra sink.

Table 6. Mean C Budget of Arctic Tundra Simulated by Process Models for 2000–2006.

Model	GPP	NPP	RH	NEP	FIREC	CO ₂ ¹	BIOCH4	OTHER	NCB
gC m ⁻² yr ⁻¹									
Regional Apps.									
LPJ-G WHyMe	209	135	110	25	1	-24	4	-	20
Orchidee	615	371	337	34	-	-34	-	-	34
TEM6	270	109	98	11	8	-3	2	2	-1
TCF	307	181	183	2	-	-	-	-	-
Global Apps.									
CLM4C	108	34	32	0 ²	1	-1	-	-	1
CLM4CN	-	-	-	1	-	-1	-	-	1
Hyland	27	14	14	0	-	0	-	-	0
LPJ	502	315	274	22 ²	19	-3	-	-	3
LPJ-Guess	562	349	305	44	20	-24	-	-	24
O-CN	74	32	31	3 ²	-	-3	-	-	3
SDGVM	-	-	-	21	5	-16	-	-	16
TRIFFID	361	227	210	17	-	-17	-	-	17
TgCyr ⁻¹									
Regional Apps.									
LPJ-G WHyMe	1918	1239	1009	230	9	-221	37	-	184
Orchidee	5643	3404	3092	312	-	-312	-	-	312
TEM6	2476	1000	899	101	73	-28	18	18	-8
TCF	2817	1661	-	-	-	-	-	-	-
Global Apps.									
CLM4C	994	314	296	18	9	-9	-	-	9
CLM4CN	-	-	-	8	-	-8	-	-	8
Hyland	248	124	123	1	-	-1	-	-	1
LPJ	4608	2887	2518	369	171	-198	-	-	198
LPJ-Guess	5160	3204	2797	407	185	-222	-	-	222
O-CN	682	298	287	11	-	-11	-	-	11
SDGVM	-	-	-	193	47	-146	-	-	146
TRIFFID	3310	2079	1927	152	-	-152	-	-	152

¹ Net CO₂-C Exchange, positive sign indicates source to the atmosphere and negative sign indicates tundra sink.

² Reported NEP does not equal reported NPP – reported RH.

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Table 7. Mean C Budget of Arctic Tundra Simulated by Process Models for 1990–2006.

Model	GPP	NPP	RH	NEP	FIREC	CO ₂ ¹	BIOCH4	OTHER	NCB
gC m ⁻² yr ⁻¹									
Regional Apps.									
LPJ-G WHyMe	200	130	106	24	1	-23	4	-	19
Orchidee	596	361	330	31	-	-31	-	-	31
TEM6	266	107	97	10	6	-4	2	2	0
Global Apps.									
CLM4C	105	32	31	1	1	0	-	-	0
CLM4CN	-	-	-	1	-	-1	-	-	1
Hyland	25	13	13	0	-	0	-	-	0
LPJ	483	303	264	30 ²	19	-11	-	-	11
LPJ-Guess	526	328	285	43	20	-23	-	-	23
O-CN	69	30	29	2 ²	-	-2	-	-	2
SDGVM	-	-	-	22	5	-17	-	-	17
TRIFFID	347	218	205	13	-	-13	-	-	13
Tg C yr ⁻¹									
Regional Apps.									
LPJ-G WHyMe	1837	1186	972	214	9	-205	35	-	170
Orchidee	5469	3307	3023	284	-	-284	-	-	284
TEM6	2434	980	887	93	52	-41	16	19	6
Global Apps.									
CLM4C	962	302	286	16	9	-7	-	-	7
CLM4CN	-	-	-	8	-	-8	-	-	8
Hyland	229	115	114	1	-	-1	-	-	1
LPJ	4432	2779	2421	359	170	-189	-	-	189
LPJ-Guess	4829	3004	2615	389	184	-205	-	-	205
O-CN	634	277	270	7	-	-7	-	-	7
SDGVM	-	-	-	203	46	-157	-	-	157
TRIFFID	3182	1993	1882	111	-	-111	-	-	111

¹ Net CO₂-C Exchange, positive sign indicates source to the atmosphere and negative sign indicates tundra sink.

² Reported NEP does not equal reported NPP – reported RH.

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Table 8. Inferred climate effect of NPP, RH, and NEP for 1990–2006 as the difference between simulations of the regional applications of process models driven by transient climate and constant climate.

Model	Transient Climate	Constant Climate	Inferred Climate Effect
NPP ($\text{g C m}^{-2} \text{ yr}^{-1}$)			
LPJ-Guess WHyMe	128	101	27
Orchidee	359	326	33
TEM6	106	90	16
RH ($\text{g C m}^{-2} \text{ yr}^{-1}$)			
LPJ-Guess WHyMe	105	86	19
Orchidee	328	291	37
TEM6	96	81	15
NEP ($\text{g C m}^{-2} \text{ yr}^{-1}$)			
LPJ-Guess WHyMe	23	15	8
Orchidee	31	35	–4
TEM6	10	9	1

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Table 9. Mean Annual Net CO₂-C Exchange Estimates of Inversion Models of Arctic Tundra for the time periods 1990–1999 and 2000–2006.

Model	1990–1999 Tg C yr ⁻¹	2000–2006 Tg C yr ⁻¹	1990–2006 Tg C yr ⁻¹	1990–1999 g C m ⁻² yr ⁻¹	2000–2006 g C m ⁻² yr ⁻¹	1990–2006 g C m ⁻² yr ⁻¹
C13_CCAM_law	–	243		–	26	
C13_MATCH_rayner	–	–283		–	–31	
JENA_s96_v3.3	–	–117		–	–13	
JMA_2010	–321	–341	–331	–35	–37	–36
LSCE_an_v2.1	–	–124		–	–14	
LSCE_var_v1.0	140	206	173	15	22	19
NICAM_niwa_woaia	–178	–80	–129	–19	–9	
rigc_Patra	–	–439		–	–48	–14

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 10. Comparison of Estimates of Mean Net CO₂-C and CH₄-C Exchange from Arctic Tundra to the Atmosphere (Tg C yr⁻¹) among Observations, Process-Based Models, and Inversion Models.

Time Period	Observations	Regional Process-Based Models	Global Process-Based Models	Inversion Models
CO ₂ Exchange				
1990–1999				
Central Estimate	138	–166	–78	–13
Uncertainty	–102 to 378	–255 to –55	–188 to 0	–321 to 140
2000–2006*				
Central Estimate	–202	–187	–93	–117
Uncertainty	–628 to 224	–312 to –28	–222 to –1	–439 to 243
1990–2006*				
Central Estimate	–103	–177	–86	–96
Uncertainty	–297 to 89	–284 to –41	–205 to –1	–331 to 173
CH ₄ Exchange				
1990–1999				
Central Estimate	10	25	–	–
Uncertainty	–1 to 22	15 to 34	–	–
2000–2006*				
Central Estimate	20	28	–	–
Uncertainty	–11 to 51	18 to 37	–	–
1990–2006*				
Central Estimate	11	26	–	–
Uncertainty	0 to 22	16 to 35	–	–

* The estimates for the observations reported for the 2000–2006 and the 1990–2006 periods may include information after 2006.

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



BGD

9, 4543–4594, 2012

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.




Fig. 1. The Arctic Tundra RECCAP Region.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩


◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.




Fig. 2. A summary of the data presented in Appendix A in the Supplement; the summary includes observations prior to 1990. The synthesis of observed NEE for different geographical regions is shown in the first five panels. The mean ($\text{g C m}^{-2} \text{day}^{-1}$) \pm standard deviation and the median are shown for summer, winter, and annual analyses. The number of studies used from Appendix A in the Supplement to estimate each mean/median is shown above the bars. Methane emissions for the circumpolar North are shown in the sixth panel in a similar fashion.




Fig. 3. Mean monthly fluxes of gross primary production (GPP), net primary production (NPP), heterotrophic respiration (RH), and net ecosystem production (NEP) simulated by LPJ-Guess WHyMe (panel A), Orchidee (panel B), and TEM6 (panel D) between 1990 and 2006 and by TCF (panel C) between 2000 and 2009.

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶


◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.




Fig. 4. Difference in mean monthly NEP between the 2000s (2000–2006) and the 1990s (1990–1999) for LPJ-Guess WHyMe, Orchidee, and TEM6.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.




Fig. 5. Comparisons among LPJ-Guess WHyMe, Orchidee, TCF, and TEM of inter-annual variability between 1990 and 2010 for anomalies of gross primary production (GPP, panel **A**), net primary production (NPP, panel **B**), heterotrophic respiration (NPP, panel **C**), and net ecosystem production (NEP, panel **D**).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. Comparisons among LPJ-Guess WhyMe, Orchidee, TCF, and TEM of inter-annual variability between 1990 and 2010 for anomalies of fire emissions (panel **A**), biogenic CH₄ emissions (panel **B**), Other fluxes (harvest and DOC exports, panel **C**), and net carbon balance (NCB, panel **D**).




Fig. 7. Comparison among inversion model estimates of the mean net monthly exchange of CO₂ between arctic tundra and the atmosphere.

An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶


◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An assessment of the carbon balance of arctic tundra

A. D. McGuire et al.




Fig. 8. Comparison among inversion model estimates of the interannual variability between 1985 and 2009 for anomalies of the net annual exchange of CO₂ between arctic tundra and the atmosphere.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

