

AN EFFECTIVE ALGORITHM TO COMPUTE MANDELBROT SETS IN PARAMETER PLANES

ANTONIO GARIJO, XAVIER JARQUE, AND JORDI VILLADELPRAT

ABSTRACT. In 2000 McMullen [McM00] proved that copies of generalized Mandelbrot set are dense in the bifurcation locus for generic families of rational maps. We develop an algorithm to an effective computation of the location and size of these generalized Mandelbrot sets in parameter space. We illustrate the effectiveness of the algorithm by applying it to concrete families of rational and entire maps.

Keywords: Holomorphic dynamics, Julia and Fatou sets, bifurcation locus, Misiurewicz bifurcation, Mandelbrot set, Algorithm.

In non linear science two fruitful ideas has been considered over the years to have a better understanding of the dynamics of mathematical models. On the one hand, the decomposition of the phase space of a particular dynamical system into several pieces, so that the restricted dynamics is more tractable; and on the other hand, the exploration of the changes that occur in the phase space under small perturbation of the original dynamical system. Under the general framework of smooth dynamical systems these ideas lead, respectively, to the study of invariant manifolds and bifurcation diagrams.

Accordingly, when working on mathematical models it is of high interest to know in advance, before studding a concrete dynamical space, the parameters for which the model may present undesirable dynamics. In other words, we want to characterize in the parameter space the set of parameters for which in dynamical plane contains open sets of initial conditions incompatible with the dynamics we are interested with.

In holomorphic dynamics, i.e. discrete dynamical systems generated by the iterates of holomorphic maps, the phase space is the complex plane and it is decomposed into two completely invariant sets: the Fatou and the Julia set. Moreover inside the Julia set of many rational and entire maps is very common to observe copies of the Julia set of the quadratic family. In a similar way in the parameter space we distinguish between the hyperbolic and the bifurcation parameters, and is very common to observe copies of the Mandelbrot set in the parameter space of many families of rational and entire maps.

In this sense the building block in holomorphic dynamics is the *Quadratic family* given by $Q_c(z) = z^2 + \xi$ where ξ is a complex parameter. The phase space or dynamical plane is divided into two completely invariant subsets: the *Fatou set* and the *Julia set*. The Fatou set, denoted by $\mathcal{F}(Q_\xi)$, is defined as the set of points z_0 where the family of iterates $\{Q_\xi^n(z_0), n \in \mathbb{N}\}$ is normal in some neighbourhood of z_0 , and its complement is the Julia set, denoted by $\mathcal{J}(Q_\xi)$. The Fatou set is formed by points with tame dynamics while its complement, the Julia set, is fulfilled by points with chaotic dynamics.

For the quadratic family, the central object of the parameter plane, or ξ -plane, is the Mandelbrot set (see Figure 1(a)), denoted by M_2 , defined as the set of ξ -parameters such

The second author is partially supported by the Polish NCN grant decision DEC-2012/06/M/ST1/00168 and the MDM-2014-445 Maria de Maeztu. The first and second authors are supported by the Spanish project MTM2014-52209-C2-2-P and the third author by the Spanish project MTM2014-52209-C2-1-P.

that the critical orbit $\{Q_\xi^n(0)\}_{n \geq 0}$ remains bounded as n tends to ∞ (see also the definition of M_d below). Easily, one can show that the Mandelbrot set is a compact set. Each one of the black bubbles is called a *hyperbolic component* of M_2 and corresponds to ξ -parameters for which the dynamical plane has an attracting periodic orbit of the same period. Moreover, in each bubble the Julia set moves continuously with respect to ξ , having a (unique) *center* for which the corresponding attracting periodic orbit has zero multiplier, i.e., it is super-attracting. In contrast, the *bifurcation parameters*, consists of the set of parameters ξ such that the Julia set does not move continuously respect to ξ . The *Misiurewicz parameters*, i.e., ξ -parameters such the critical orbit (the orbit of $z = 0$) lands in a repelling periodic orbit, form a dense subset of ∂M_2 . Proving that the interior of the Mandelbrot set consists only on hyperbolic parameters is still a challenged open question. For an introduction to complex dynamics and the quadratic family we refer to [DH84, DH85a, McM94, Mil06, CG93, Tan00] and references therein.

For more general (uni-parametric) families of holomorphic maps a similar approach has been considered to split the dynamical plane into the Fatou and the Julia sets as well as to define the bifurcation locus. In this situation is very common to observe in the dynamical plane copies of the Julia set of the quadratic family and in the parameter plane copies of the Mandelbrot set. One of the cornerstone tools for this assertion is the theory of polynomial-like mappings, developed by A. Douady and J.H. Hubbard in the landmark paper [DH85b]. They rigorously explained why it is so common to observe, in computer experiments, homeomorphic copies of polynomial Julia set in dynamical planes of rational or transcendental families, and small homeomorphic copies of (generalized) Mandelbrot set in the corresponding parameter planes. Remember that for a given $d \geq 2$ the *generalized Mandelbrot set* is defined as

$$M_d = \{\xi \in \mathbb{C}, \mathcal{J}(z^d + \xi) \text{ is connected}\}.$$


In Figure 1(a)-(b) we show the parameter planes of $z^d + \xi$ for $d = 2$ and $d = 3$, respectively. The black bubbles, as before, are hyperbolic components and the boundary of M_d is the bifurcation locus.

This central contribution due to A. Douady and J.H. Hubbard has been, and it is still being, applied widely. In 2000 C. McMullen [McM00] push forward the arguments above by showing that the existence of *polynomial dynamics* in families of rational maps is, in fact, *universal*. More precisely, he proved that small generalized Mandelbrot sets are dense in the bifurcation locus for generic families of rational maps, and so pieces of the Julia set for those non-polynomial maps resembles the Julia set of polynomials.

This result has important theoretical implications. For instance, let P be a polynomial of degree $d \geq 2$. A natural root finding algorithm to solve the equation $P(z) = 0$ is given by the Newton map

$$N_P(z) := z - \frac{P(z)}{P'(z)}.$$

It is well known that the zeros of P correspond bijectively to the fixed points of N_P which are all (locally) attracting. Roughly speaking, if there are no other stable dynamics for N_P different from the attracting basins of the attracting fixed points, we can ensure that $N_P^n(z_0)$ will numerically converge to one of the roots of P for *almost all* initial conditions $z_0 \in \mathbb{C}$. However, if other stable dynamics coexist in the dynamical plane this will no longer be true. Precisely the existence of these *bad* initial conditions in dynamical plane is in direct correspondence with the existence of generalized Mandelbrot sets in parameter plane. So,

(a) The parameter plane of $z^2 + \xi$.(b) The parameter plane of $z^3 + \xi$.FIGURE 1. Mandelbrot sets of degree $d = 2$ and $d = 3$.

the natural question is *where are they?* Where, in parameter plane, are located the universal Mandelbrot sets predicted by McMullen?

The main goal of this article is to give an effective algorithm to compute hyperbolic components (location and size) of the copies of the Mandelbrot set lying in the parameter plane of a uni-parametric holomorphic family (see section 2). It is worth to be noticed that to implement the algorithm it is not required to run over McMullen's paper (see section 1), and, more interesting, can be applied to general holomorphic, not necessary rational, maps. We illustrate, by means of different sort of examples, the use of the algorithm in Section 3.

1. SMALL MANDELBROT SETS IN THE BIFURCATION LOCUS

To provide a solid theoretical content to the algorithm we will describe in next section, we firstly discuss the key results in [McM00]. Even though we will apply the algorithm to transcendental maps we use a rational framework, as it is done in [McM00], for the theoretical discussion. At the end of the section we will briefly argue how to deal with the general scenario.

We consider a holomorphic family of rational maps over the unit disc \mathbb{D} as a map

$$f : \mathbb{D} \times \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}} \\ (t, z) \rightarrow f_t(z)$$

requiring that for each parameter $t \in \mathbb{D}$, the rational map $f_t : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ satisfies $\deg(f_t) \geq 2$. The *Bifurcation locus*, denoted in what follows by $B(f) \subset \mathbb{D}$, it is defined as a set of t -parameters such that the Julia set $\mathcal{J}(f_t)$ does not vary continuously (in the Hausdorff topology) over any neighbourhood of t . Without lost of generality we assume that $0 \in B(f)$. Indeed, we can always arrange the base point using the change of parameters $t := t_0 + t$. See Subsection 3.2. A *local bifurcation* is a holomorphic family of rational maps f_t over the unit disc such that $t = 0 \in B(f)$, and for simplicity we say a local bifurcation f . A point $c_t \in \mathbb{C}$ such that

$f'_t(c_t) = 0$ it is called a *critical point of f_t* , and its image $v_t := f_t(c_t)$ it is called a *critical value of f_t* .

The following definitions and proposition play a crucial role in the description of $B(f)$ (see details in [McM00]).

Definition 1. *Let f be a local bifurcation. A marked critical point c is a holomorphic map $c : \mathbb{D} \rightarrow \hat{\mathbb{C}}$ such that $f'_t(c_t) = 0$. To fix the notation we will write (f, c) as the holomorphic family of rational maps with a marked critical point. We say that the marked critical point c of f is active if the sequence of iterates $\{f_t^n(c_t)\}_{n \geq 0}$ fails to be a normal family in any neighbourhood of $t = 0$.*

Definition 2. *Let f be a local bifurcation. We say that (f, c) is a (degree d) Misiurewicz bifurcation if the following conditions hold.*

- M1. $f_0(c_0)$ is a repelling fixed-point of f_0 , with multiplier $\lambda_0 := f'_0(f_0(c_0))$;
- M2. c_0 is unramified for f_0 (that is, c_0 has infinitely backward images eventually mapping to c_0 with degree 1);
- M3. $f_t(c_t)$ is not a fixed point of f_t for some t (c active); and
- M4. $\deg(f_t, c_t) = d$ for all t sufficiently small (the degree of the critical point is locally constant).

Proposition 1.1. *For any local bifurcation (f, c) with c active and c_0 unramified, there is a base change and an $n > 0$ such that (f^n, c) is a (degree d) Misiurewicz bifurcation.*

In [McM00] McMullen deeply studies the (degree d) Misiurewicz bifurcations, by means of the theory of polynomial-like maps, to show that the bifurcation locus $B(f) \subset \mathbb{D}$ of any local bifurcation is either empty or contains the quasiconformal image of ∂M_d for some d . Moreover, since arguments are of *local* character and Misiurewicz points are dense in the bifurcation locus, he concludes that small generalized Mandelbrot set are dense in the bifurcation locus (Theorem 1.1 and Corollary 1.2 in [McM00]). See Figure 2 where t_n are the center of the main hyperbolic component of baby Mandelbrot sets converging to 0.

He shows that any Misiurewicz bifurcation produces a cascade of generalized Mandelbrot sets (corresponding to values of t_n tending to $t = 0$). The precise statement is given below. We use $\text{diam}(A)$ to denote the diameter of a set $A \subset \mathbb{C}$ and $a_n = O(b_n)$ if $|a_n| < C|b_n|$ for some constant $C > 0$.




FIGURE 2. Parameter plane of the Newton map N_t (Equation 8). Misiurewicz bifurcation at $t = 0$ of degree 2 for the Newton map N_t . For details see subsection 3.1.

Theorem 1.2 (Theorem 4.1 in [McM00]). *Let (f, c) be a (degree d) Misiurewicz bifurcation. Then, the parameter space \mathbb{D} contains quasiconformal copies \mathcal{M}_d^n of the degree d Mandelbrot*




FIGURE 3. Sketch of the construction of a superattracting orbit of period n near a Misiurewicz bifurcation.

set M_d , converging to the origin, with $\partial\mathcal{M}_d^n$ contained in the bifurcation locus $B(f)$. More precisely, for all $n \gg 0$ there are homeomorphisms

$$\phi_n : M_d \rightarrow \mathcal{M}_d^n \subset \mathbb{D}$$

such that:

- (a) f_t^n is hybrid equivalent to $z^d + \xi$ whenever $t = \phi_n(\xi)$;
- (b) $d(0, \mathcal{M}_d^n) \asymp |\lambda_0|^{-n/r}$;
- (c) $\text{diam}(\mathcal{M}_d^n)/d(0, \mathcal{M}_d^n) \asymp |\lambda_0|^{-n/(d-1)}$;
- (d) ϕ_n extends to a quasiconformal map of the plane with dilatation bounded by $1 + O(\epsilon_n)$;
and
- (e) $\psi_n^{-1} \circ \phi_n(\xi) = \xi + O(\epsilon_n)$, where $\psi_n(\xi) = t_n(1 + \gamma_n \xi)$.

Although details in the proof of the above result are far from being straightforward, the idea of the construction is quite simple and useful to define the algorithm in the next section.

Assume that we have a Misiurewicz bifurcation. For $t = 0$, the critical point c_0 lands into a repelling fixed point, so $c_0 \mapsto f_0(c_0) \mapsto f_0(c_0)$ (see Figure 3(a)). Besides the dynamics of f_0 near $f_0(c_0)$ is conformally conjugated to $z \mapsto f_0(c_0) + \lambda_0(z - f_0(c_0))$, with $|\lambda_0| > 1$ (see [CG93] for details on linearizing coordinates).

For small values of t we have two crucial dynamical facts. On the one hand, the fixed point $p_0 := f_0(c_0)$ moves continuously, that is, there exists p_t such that $p_0 := f_0(c_0)$, $f_t(p_t) = p_t$ and $|\lambda_t| := |f_t'(p_t)| > 1$. On the other hand, because of condition M3, we can assume that $f_t(c_t)$ is not anymore a fixed point but, by continuity, $f_t(c_t)$ is close to the repelling fixed point p_t . See Figure 3(b). We claim that for n large enough there exists a small t such that $f_t^n(c_t) = c_t$ and so we have created a periodic point with multiplier 0, so a parameter corresponding to the center of the main cardioid of a generalized Mandelbrot set (degree depending on the multiplicity of the critical point c_t). To see the claim we observe that c_0 is unramified (infinitely many backward images) and p_0 belongs to the Julia set, so there must be a point b_0 near p_0 such that $f_0^n(b_0) = c_0$ for n large. The value of the parameter r in Theorem 1.2 (b) is equal to the multiplicity, in terms of the variable t , of the map $f_t(c_t)$ at $t = 0$; or in other words $f_t(c_t) = f_0(c_0) + O(t^r)$. For a complete proof see [McM00].

Remark 1. Since all arguments used in [McM00] are local we can extend those results to other holomorphic families of non-rational maps, like transcendental entire.

2. THE ALGORITHM

In this section we present an algorithm to compute the location and size of the small generalized Mandelbrot sets in the bifurcation locus $B(f)$ coming from the Misiurewicz bifurcation using the estimations obtained in Theorem 1.2. Next we will effectively use the algorithm in some examples.

We assume that (f, c) is a (degree d) Misiurewicz bifurcation. In order to be more precise we denote the holomorphic family by $f(t, z) := f_t(z)$ to emphasize the dependence on the parameter t and the variable z . In a similar way the marked critical point c_t is given by the holomorphic map $c : \mathbb{D} \rightarrow \hat{\mathbb{C}}$, so we write $c_t = c(t)$. We define the following auxiliary maps,

$$(1) \quad G_0(t) = c(t) \quad \text{and} \quad G_k(t) = f^k(t, c(t)) \quad \text{for } k > 0,$$

that is G_k , $k \geq 0$, is a function depending on t giving the k -th iterate of the critical point $c(t)$, for the corresponding map f_t . We have the following recursive expression, for $k \geq 0$,

$$(2) \quad G_{k+1}(t) = f^{k+1}(t, c(t)) = f\left(t, f^k(t, c(t))\right) = f(t, G_k(t)).$$

Some computations show that

$$(3) \quad G'_0(t) = c'(t) \quad \text{and} \quad G'_{k+1}(t) = \frac{\partial f}{\partial t}(t, G_k(t)) \cdot 1 + \frac{\partial f}{\partial z}(t, G_k(t)) \cdot G'_k(t), \quad k > 0.$$

The *center* of the generalized Mandelbrot set M_d of the family $z^d + \xi$ is the parameter $\xi_0 = 0$, so the center of M_d corresponds to the polynomial z^d having a superattracting fixed point at the origin. See Figure 1. Similarly, we define the *size* of M_d measuring the distance between two concrete parameters in M_d . In order to do this, we consider the distance, denoted by L , between the center ξ_0 of M_d and a parameter ξ_1 such that $z^d + \xi_1$ exhibits a superattracting two cycle; thus $L = |\xi_0 - \xi_1|$. See also Figure 1. Easy computations show that there are $d - 1$ possible choices for the parameter ξ_1 given by $d^{-1}\sqrt{-1}$. However, these $d - 1$ values are symmetrically distributed around the origin with equal distance $L = 1$ to the origin. By definition the center and the size of the generalized Mandelbrot set M_d is zero and one, respectively.

Since the definition of the center and the size of M_d are in dynamical terms, we can easily transfer these two definition to any small copy of M_d .

Definition 3. *Let \mathcal{M}_d^n be a homeomorphic copy of M_d (see Theorem 1.2). We define $t_n := \phi_n(0)$, the center of \mathcal{M}_d^n ; or equivalently, the parameter such that the marked critical point $c(t_n)$ has a superattracting orbit of minimal period n , and $s_n := \phi_n(d^{-1}\sqrt{-1})$ (we choose one of the roots), a parameter such that the critical marked point $c(s_n)$ exhibits a superattracting orbit of minimal period $2n$. Then the size \mathcal{M}_d^n is defined by $L_n = |t_n - s_n|$.*

Under this notation the main concern is to numerically compute t_n (location) and L_n (size) of the small generalized Mandelbrot sets in the parameter plane of a holomorphic family. However we notice that, at least theoretically, the computations of those values using the auxiliary maps G_k 's is immediate.

Lemma 2.1. *Assume the notation of Definition 3. The values of t_n and s_n are solutions of the following system of equations*

$$(4) \quad \begin{cases} G_n(t_n) & = c(t_n) \\ G_{2n}(s_n) & = c(s_n). \end{cases}$$

Of course $L_n = |t_n - s_n|$.

Proof. The proof is straightforward since $G_n(t_n) = f^n(t_n, c(t_n))$ and $G_{2n}(s_n) = f^{2n}(s_n, c(s_n))$. \square

We will use Newton's method to numerically solve these equations. Nonetheless remember that the small copies of the Mandelbrot set we are looking for are dense in the bifurcation locus which implies that it is crucial to have an *arbitrarily good* accuracy on the initial condition in order to find the correct parameters. In particular we notice that if t_n solves the first equation, then it also solves the second equation. To attain this goal we will use a continuation method.

Computing the location t_n of \mathcal{M}_d^n . We will compute t_n recursively. Assume that we know with high accuracy the location t_{n-1} . From Theorem 1.2(b) we get

$$|t_n| \approx |\lambda_0|^{-n/r}, \quad |t_{n-1}| \approx |\lambda_0|^{-(n-1)/r}$$

and we conclude $|t_n| \approx |t_{n-1}| \cdot |\lambda_0|^{-1/r}$. So, we numerically compute t_n as a result of the iterative process (Newton's method applied to the equation $G_n(t) - c(t) = 0$) given by

$$(5) \quad \begin{aligned} \tau_0 &= t_{n-1} \cdot \lambda_0^{-1/r} \\ \tau_k &= \tau_{k-1} - (G_n(\tau_{k-1}) - c(\tau_{k-1})) / (G'_n(\tau_{k-1}) - c'(\tau_{k-1})), \quad k \geq 1. \end{aligned}$$

We can numerically compute $G_n(t)$ and $G'_n(t)$ using the recursive expressions obtained in Equations (2) and (3).

Computing the size L_n of \mathcal{M}_d^n . In this case we also assume that t_{n-1} and L_{n-1} are known values and we will compute the size $L_n = |t_n - s_n|$ of \mathcal{M}_d^n . As before we can obtain s_n solving (4) using the Newton's method from an initial condition close to s_n .

We firstly find an initial condition close to s_n . In this case we identify the diameter of \mathcal{M}_d^n with L_n and using Theorem 1.2 estimations (b) and (c), we have that

$$\frac{L_n}{L_{n-1}} \approx \frac{|t_n|}{|t_{n-1}|} \cdot \frac{|\lambda_0|^{-n/(d-1)}}{|\lambda_0|^{-(n-1)/(d-1)}} = \frac{|t_n|}{|t_{n-1}|} \cdot |\lambda_0|^{-\frac{1}{d-1}},$$

so, we conclude that $s_n \approx t_n \left(1 + \frac{L_{n-1}}{t_{n-1}} \cdot |\lambda_0|^{-\frac{1}{d-1}} \cdot {}^{d-1}\sqrt{-1}\right)$.

Finally, we use this initial condition to solve numerically the equation $G_{2n}(t) - c(t) = 0$. We construct the following sequence of iterates $(\eta_k)_{k \geq 0}$ tending to s_n as k tends to ∞ ,

$$(6) \quad \begin{aligned} \eta_0 &= t_n \left(1 + \frac{L_{n-1}}{t_{n-1}} \cdot \lambda_0^{-\frac{1}{d-1}} \cdot {}^{d-1}\sqrt{-1}\right) \\ \eta_k &= \eta_{k-1} + (G_{2n}(\eta_{k-1}) - c(\eta_{k-1})) / (G'_{2n}(\eta_{k-1}) - c'(\eta_{k-1})). \end{aligned}$$

We remark that in the above algorithm to compute t_n and L_n it is crucial to have an accurate initial condition. Assuming that the rational map f_t has degree d , we have that the n -th iterate f_t^n has degree d^n . Hence the number of solutions of (4) grows exponentially.

From the numerical computation point of view of the location t_n and the size L_n we can also visualize the results obtained in Theorem 1.2(b-c). Thus for example simple computations show a linear relation between $\log(t_n)$ and $\log(L_n)$ with respect to n . More precisely,

$$(7) \quad \begin{aligned} \log(t_n) &\approx \alpha + \beta \cdot n && \text{where } \beta = -\log|\lambda_0|/r \\ \log(L_n) &\approx \gamma + \delta \cdot n && \text{where } \delta = -\log|\lambda_0| \left(\frac{1}{d-1} + \frac{1}{r}\right). \end{aligned}$$

In the next section we illustrate how to implement the above algorithm to compute the location t_n and the size L_n of a cascade of small copies of the Mandelbrot set M_d for different families covering rational as well as transcendental scenarios.

3. APPLICATIONS

3.1. Example 1. The Newton method. In this example we consider the Newton's method applied to the family of polynomials

$$p_t(z) = z^{d+1} - tz + 1,$$

where $d \geq 2$ and t is a complex parameter. To simplify the notation we will assume that d is fixed; so, we erase the dependence on d . In fact the parameter d coincides with the degree of the Newton method applied to the polynomial p_t at the unique active critical point.

The expression of the Newton's map applied to p_t writes as

$$(8) \quad N_t(z) = z - \frac{p_t(z)}{p_t'(z)} = z - \frac{z^{d+1} - tz + 1}{(d+1)z^d - t} = \frac{dz^{d+1} - 1}{(d+1)z^d - t}.$$

It is easy to check that the zeros of the polynomial p_t are finite fixed points of N_t which are always attracting. Moreover the point at ∞ is a repelling fixed point of N_t with multiplier $\lambda_0 = (d+1)/d$ (independently of t). We remark that given any holomorphic family of polynomials q_t of degree $d+1$ and their associated family of Newton's method N_{q_t} , then the point of ∞ is the unique repelling fixed point of N_{q_t} and its multiplier is always equal to $\lambda_0 = (d+1)/d$ (independently of q_t).

The critical points of N_t correspond to the zeros of the rational map

$$N_t'(z) = \frac{p_t(z)p_t''(z)}{(p_t'(z))^2}.$$

So the critical points of N_t are the zeros of p_t and $z = 0$ since $p_t''(z) = (d+1)dz^{d-1}$. The zeros of the polynomial are fixed points of the Newton map, so there are not active critical points. Therefore the origin is the unique active critical point of N_t and the degree of N_t at the origin is equal to d , so $\deg(N_t, 0) = d$. We observe that the marked critical point is $c(t) = 0$ for all the values of t ; and we can interpret the parameter d as the degree of this active critical point. The corresponding critical value, $N_t(0)$, is equal to $1/t$. Furthermore at the parameter $t = 0$ we have a Misiurewicz bifurcation of degree d (see Definition 4). In Figure 4 we show the parameter plane of the Newton map for different values of d .

The hyperbolic components in the t -parameter plane are the open subsets of \mathbb{C} in which the unique active critical point $c(t) = 0$ either eventually map to one of the immediate basin of attraction corresponding to one of the roots of p_t or it has its own hyperbolic dynamics associated to an attracting periodic point of period greater than one. Of course, the bifurcation locus corresponds to the union of all boundaries of those components and possible accumulating points, see Figure 4. For a detailed study of the dynamical system given by (8) we cite [CGJV14].

We choose this example for several reasons. Firstly, there is only one active critical point, $c(t) = 0$, and changing the value of the parameter $d \geq 2$ the active critical point has local degree d . Thus, changing the value of the parameter d we find out generalized Mandelbrot sets \mathcal{M}_d densely appearing in the bifurcation locus of N_t (see Figure 4). Moreover, N_t has a Misiurewicz bifurcation at $t = 0$ of degree d (see Figure 2). Secondly, and more important, we claim that the family N_t can be considered as a model for the study of Misiurewicz bifurcations for Newton's method applied to arbitrary families of polynomials. To see the claim, suppose that q_t is a family of polynomials of degree $d+1$ and we denote by N_{q_t} the Newton's method associated to q_t . Assume, without loss of generality, that $t = 0$ is a Misiurewicz bifurcation of some degree. As we already explain the unique repelling fixed point of N_{q_t} is located at ∞




FIGURE 4. Different parameter planes of N_t for two values of the parameter d .

and its multiplier, $\lambda_0 = (d + 1)/d$, coincides with the multiplier of the family N_t . But from Theorem 1.2(b-c) this *universal* multiplier controls the location and the size of the cascade of the generalized Mandelbrot sets tending to $t = 0$ that appears in the parameter plane. This phenomenon explain why the parameter plane of two Newton's method applied to *different* families of polynomials of the *same* degree look like so similar.

Going back to N_t , remember that it has a Misiurewicz bifurcation at $t = 0$ of degree d . So we have in the parameter plane of N_t a cascade of generalized Mandelbrot sets, denoted by \mathcal{M}_d^n where n denotes the period of the attracting cycle in the main cardioid, tending to the Misiurewicz bifurcation parameter, $t = 0$. We denote t_n and L_n the location and size of the baby Mandelbrot set \mathcal{M}_d^n of period n . Applying the algorithm developed in section 2 we can compute numerically the centers and the size of these baby Mandelbrot sets. In Table 3.1 we show these values for $n \leq 25$ when $d = 2$ and $d = 3$. In Figure 2 we also plot the first terms of this sequence for $d = 2$.

	$d = 2$	$d = 2$	$d = 3$	$d = 3$
Period n	Location t_n	Size L_n	Location t_n	Size L_n
2	1.259921050	$1.82397728 \cdot 10^{-2}$	1.316074013	$2.40997622 \cdot 10^{-2}$
3	0.765955195	$1.17992525 \cdot 10^{-2}$	0.916566576	$2.26459898 \cdot 10^{-2}$
4	0.481413938	$5.91059457 \cdot 10^{-3}$	0.653403073	$1.65675678 \cdot 10^{-2}$
5	0.309653924	$2.77393767 \cdot 10^{-3}$	0.474076799	$1.14015617 \cdot 10^{-2}$
6	0.201928044	$1.26947908 \cdot 10^{-3}$	0.347781941	$7.66436441 \cdot 10^{-3}$
7	0.132762229	$5.73984187 \cdot 10^{-4}$	0.256914343	$5.09179360 \cdot 10^{-3}$
8	$8.772559252 \cdot 10^{-2}$	$2.57817423 \cdot 10^{-4}$	0.190650421	$3.35928347 \cdot 10^{-3}$
9	$5.814830777 \cdot 10^{-2}$	$1.15357132 \cdot 10^{-4}$	0.141908798	$2.20632049 \cdot 10^{-3}$
10	$3.862008787 \cdot 10^{-2}$	$5.14925041 \cdot 10^{-5}$	0.105850492	$1.44463194 \cdot 10^{-3}$
11	$2.568314590 \cdot 10^{-2}$	$2.29504099 \cdot 10^{-5}$	$7.907129190 \cdot 10^{-2}$	$9.43859243 \cdot 10^{-4}$
12	$1.709415373 \cdot 10^{-2}$	$1.02191826 \cdot 10^{-5}$	$5.912954709 \cdot 10^{-2}$	$6.15720590 \cdot 10^{-4}$
13	$1.138377583 \cdot 10^{-2}$	$4.54744711 \cdot 10^{-6}$	$4.425102165 \cdot 10^{-2}$	$4.01208952 \cdot 10^{-4}$
14	$7.583732763 \cdot 10^{-3}$	$2.02273525 \cdot 10^{-6}$	$3.313488345 \cdot 10^{-2}$	$2.61215381 \cdot 10^{-4}$
15	$5.053407225 \cdot 10^{-3}$	$8.99480984 \cdot 10^{-7}$	$2.482142214 \cdot 10^{-2}$	$1.69966087 \cdot 10^{-4}$
16	$3.367867386 \cdot 10^{-3}$	$3.99912962 \cdot 10^{-7}$	$1.859945698 \cdot 10^{-2}$	$1.10542637 \cdot 10^{-4}$
17	$2.244769739 \cdot 10^{-3}$	$1.77781615 \cdot 10^{-7}$	$1.394029959 \cdot 10^{-2}$	$7.18706781 \cdot 10^{-5}$
18	$1.496302176 \cdot 10^{-3}$	$7.90266424 \cdot 10^{-8}$	$1.045001811 \cdot 10^{-2}$	$4.67159515 \cdot 10^{-5}$
19	$9.974410761 \cdot 10^{-4}$	$3.51266814 \cdot 10^{-8}$	$7.834593604 \cdot 10^{-3}$	$3.03597177 \cdot 10^{-5}$
20	$6.649190879 \cdot 10^{-4}$	$1.56129631 \cdot 10^{-8}$	$5.874306375 \cdot 10^{-3}$	$1.97273977 \cdot 10^{-5}$
21	$4.432608954 \cdot 10^{-4}$	$6.93942194 \cdot 10^{-9}$	$4.404809479 \cdot 10^{-3}$	$1.28173032 \cdot 10^{-5}$
22	$2.954990445 \cdot 10^{-4}$	$3.08428448 \cdot 10^{-9}$	$3.303090086 \cdot 10^{-3}$	$8.32702141 \cdot 10^{-6}$
23	$1.969957106 \cdot 10^{-4}$	$1.37082183 \cdot 10^{-9}$	$2.477027012 \cdot 10^{-3}$	$5.40950304 \cdot 10^{-6}$
24	$1.313288506 \cdot 10^{-4}$	$6.09262656 \cdot 10^{-10}$	$1.857606938 \cdot 10^{-3}$	$3.51403497 \cdot 10^{-6}$
25	$8.755184568 \cdot 10^{-5}$	$2.70776582 \cdot 10^{-10}$	$1.393113383 \cdot 10^{-3}$	$2.28265653 \cdot 10^{-6}$

TABLE 1. Location t_n and size L_n of the Mandelbrot sets \mathcal{M}_2^n tending to $t = 0$ for the family N_t with $d = 2$ and $d = 3$. See Figure 2.

Another way to visualise the output of these numerical computations is to use Theorem 1.2 and (7). From there we have

$$\begin{aligned} |t_n| &\simeq C |\lambda_0|^{-n/r} \\ |L_n| &\simeq D |\lambda_0|^{-n/r - n/(d-1)}, \end{aligned}$$

or equivalently,

$$\begin{aligned} \log |t_n| &\simeq \log(C) - \frac{\log |\lambda_0|}{r} \cdot n \\ \log |L_n| &\simeq \log(D) - \log |\lambda_0| \left(\frac{1}{r} + \frac{1}{d-1} \right) \cdot n. \end{aligned}$$

The above expressions can be expressed as a linear regressions between the logarithm of the location and size of \mathcal{M}_d^n and n , respectively. We plot in Figure 5 the results of the numerical computation of t_n and L_n for different values of the parameter d . In both cases, subfigures (a) and (b), the x -axis correspond to the period n of the small Mandelbrot set \mathcal{M}_d^n , while the y -axis corresponds to $\log |t_n|$ and $\log |L_n|$, respectively.

3.2. Example 2. An entire transcendental family. Due to the existence of an essential singularity at infinity, the global dynamics of transcendental entire maps is quite different to the dynamics of rational maps. For instance new types of Fatou components may exist like Baker or wandering domains (see [Dev94] for a brief introduction to transcendental dynamics).




FIGURE 5. Linear regression between $\log(t_n)$ and n (left) and $\log(L_n)$ and n (right) for the map N_t and several values of d .

However, according to Remark 1, since the theory of Misiurewicz bifurcations is of local character, it can be easily extended to families of transcendental entire maps. The main goal is to show that the above algorithm runs with no significant modifications.

We consider the family of entire transcendental maps given by

$$(9) \quad f_t(z) = tz^2 \exp(z),$$

where $t \neq 0$ is a complex parameter. For all the values of the parameter t we have a superattracting fixed point at the origin, i.e., $f_t(0) = f'_t(0) = 0$, which is also an asymptotic value (a singularity of the inverse map, see [Dev94], pages 181-182). Computing the derivative we have that $f'_t(z) = tz(z+2)\exp(z)$ and thus there also exists a unique (active) critical point located at $c(t) \equiv -2$. The local degree of both critical points is 2 since $f''_t(0) \neq 0$ and $f''_t(-2) \neq 0$. The Fatou set of f_t always contains the basin of attraction of 0, that is the points in \mathbb{C} that under iteration tends to 0.

Although ∞ is an essential singularity of the map f_t and so, the global dynamics is far from being polynomial or rational, the existence of a critical point at -2 of degree 2 makes some similarities between the family f_t and the quadratic family. Such similarities are explained when plotting the parameter plane of f_t where we can see some copies of small Mandelbrot sets. In this parameter plane we plot three different behaviours, the first one is the set of parameters such that the orbit of the active critical point -2 tends to zero (blue), the second one when the orbit escapes to ∞ (red) and finally when the orbit accumulates in an attracting cycle (black). See Figure 6.

In order to study a Misiurewicz bifurcation we need to find parameter values such that the active critical point lands in a repelling fixed point. Thus, we first compute the critical value $v(t) = f_t(-2) = 4te^{-2}$, and imposing that $v(t)$ is a fixed point $f_t(v(t)) = v(t)$ we arrive to the equation

$$f_t(4te^{-2}) = 4te^{-2}.$$

Instead of solve numerically this equation we can take advantage to write it down in term of the function $g(z) = z \exp(z)$ and its (multi-valued) function $W(z)$ known as Lambert function.




FIGURE 6. Parameter plane of the entire transcendental map f_t . Misiurewicz bifurcation at $t = \bar{\tau}$ of degree 2 for the map f_t .

Using this approach the above equation writes as

$$[g(z)]^2 = \frac{1}{e^2} \quad \text{where } z = 2te^{-2},$$

obtaining two real solutions $W(\pm 1/e)e^2/2$. Only one of those $\bar{\tau} = W(1/e)e^2/2 \approx 1.028795064$ satisfies the dynamical condition that the critical point is prefix. The other value correspond to the case that the critical point is a fixed point. For this parameter value $\bar{\tau}$ the active critical point $c(\bar{\tau}) = -2$ lands in a repelling fixed point located at $4\bar{\tau}e^{-2}$ with multiplier $\lambda_0 \approx 2.556929$. We can redefine the parameter $t := t + \bar{\tau}$ and considering the family of functions

$$(10) \quad g_t(z) := f_{t+\bar{\tau}}(z) = (t + \bar{\tau})z^2 \exp(z)$$

concluding that g_t has a Misiurewicz bifurcation of degree 2 at the parameter $t = 0$. Applying the algorithm developed in Section 2 we have computed the location t_n and size L_n of the cascade of Mandelbrot sets \mathcal{M}_2^n tending to $t = 0$. To illustrate these computations we use, as before, the regression between the period n of the small Mandelbrot set \mathcal{M}_2^n with $\log |t_n|$ and $\log |L_n|$, in Figure 7(a) or Figure 7(b), respectively.

We end up this section by considering a slight modification of the Misiurewicz bifurcation described above (where, remember, the active critical point $c(t)$ lands in one iterate at the repelling fixed point). Following [McM00], we introduce a new parameter k controlling the number of iterates that the active critical point needs to land at the repelling fixed point




FIGURE 7. Linear regression between $\log(t_n)$ and n (left) and $\log(L_n)$ and n (right) for the entire map f_t .

Definition 4. Let f be a local bifurcation. We say that (f, c) is a degree d generalized Misiurewicz bifurcation if the following conditions hold.

- M1. There exists $k \geq 1$ such that $f_0^k(c_0)$ is a repelling fixed-point of f_0 , with multiplier $\lambda_0 = f_0'(f_0^k(c_0))$;
- M2. c_0 is unramified for f_0 ;
- M3. $f_t^k(c_t)$ is not a fixed point of f_t , for some t ; and
- M4. $\deg(f_t, c_t) = d$ and $\deg(f_t^k, c_t) = d$ for all t sufficiently small.

Since the active critical point takes k iterates to arrive to the repelling fixed point, in the last condition we require that this critical point does not collide with another critical point of the map until it reaches the repelling fixed point.

It is worth to be noticed that this approach is different than the one consisting of applying the theory of Misiurewicz bifurcation developed in Section 1 to the map $g_t = f_t^k$. Indeed if the active critical point c_0 needs k iterates to land at the repelling fixed point $f^k(c_0)$, it turns out that the multiplier of such fixed point is $\lambda_0 = f_0'(f_0^k(c_0))$. In contrast if we take $g_t := f_t^k$, the fixed point under g_0 is $g_0(c_0)$ and its multiplier is $g_0'(g_0(c_0)) = \lambda_0^k$; and on the other hand the periods of the baby Mandelbrot sets are multiples of k .

Assuming that we have a generalized Misiurewicz bifurcation at $t = 0$ we obtain the same estimations of the location t_n and size L_n of the generalized copies of the Mandelbrot set \mathcal{M}_2^n tending to $t = 0$. We recall that the key ingredient in these estimations is the local behavior near the repelling fixed point $f^k(c_0)$, the only difference is that the critical point takes more iterates to reach this point. As we mention before we exclude the case when this critical point heats another critical point and thus increasing the degree of the map.

We apply this generalization to the family of maps f_t (Equation 9) for $k = 2$. Thus, the first step is to find a concrete value of the parameter, denoted by $\hat{\tau}$, such that the active critical point $c(\hat{\tau}) = -2$ lands into two iterates in a repelling fixed point. This is equivalent to solve the equation

$$f_t^3(-2) = f_t^2(-2).$$

We solve numerically the above equation obtaining the parameter $\hat{\tau} \approx -19.63162088$. For this concrete parameter value, the critical point -2 lands into two iterates to the repelling fixed point $f_{\hat{\tau}}^2(-2) \approx -5.37511380$ with multiplier $\lambda_0 \approx 1.9462489$. In Figure 8 we show in the parameter plane of the map f_t the location of the parameter $\hat{\tau}$ and the sequence of Mandelbrot sets \mathcal{M}_2^n tending to this generalized Misiurewicz bifurcation.




FIGURE 8. Parameter plane of the entire transcendental map f_t (Equation 9). Generalized Misiurewicz bifurcation at $t = \hat{\tau}$ of degree 2 for the map f_t . Observe that the smallest period is $n = k + 1 = 3$.

We observe that by construction the sequence of copies of Mandelbrot set \mathcal{M}_2^n have $n \geq 3$, since we need two iterates to be close to the repelling fixed and at least another iterate to return to the critical point. In general, when the critical point takes k iterates to land into a repelling fixed point, the small copies of Mandelbrot set have period $n \geq k + 1$. Applying again the algorithm developed in the Section 2 we obtain the location \hat{t}_n and size \hat{L}_n of \mathcal{M}_2^n of degree n . Finally, in Figure 9 we show the linear regression between the period and the logarithm of the location and size of \mathcal{M}_2^n as expected.




FIGURE 9. Linear regression between $\log(\hat{t}_n)$ and n (left) and $\log(\hat{L}_n)$ and n (right) for the entire map f_t .

REFERENCES

- [CG93] Lennart Carleson and Theodore W. Gamelin. *Complex dynamics*. Universitext: Tracts in Mathematics. Springer-Verlag, New York, NY, 1993.

- [CGJV14] Beatriz Campos, Antonio Garijo, Xavier Jarque, and Pura Vindel. Newton's method on Bring-Jerrard polynomials. *Publ. Mat.*, 58(suppl.):81–109, 2014.
- [Dev94] Robert L. Devaney. *Complex dynamical systems (Cincinnati, OH, 1994)*, volume 49, chapter Complex dynamics and entire functions, pages 181–206. Amer. Math. Soc., Providence, RI, 1994.
- [DH84] Adrien Douady and John H. Hubbard. *Étude dynamique des polynômes complexes. Partie I*, volume 84 of *Publications Mathématiques d'Orsay*. Université de Paris-Sud, Orsay, 1984.
- [DH85a] Adrien Douady and John H. Hubbard. *Étude dynamique des polynômes complexes. Partie II*, volume 85 of *Publications Mathématiques d'Orsay*. Université de Paris-Sud, Orsay, 1985.
- [DH85b] Adrien Douady and John Hamal Hubbard. On the dynamics of polynomial-like mappings. *Ann. Sci. École Norm. Sup. (4)*, 18(2):287–343, 1985.
- [McM94] Curtis T. McMullen. *Complex dynamics and renormalization*, volume 135 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1994.
- [McM00] Curtis T. McMullen. The Mandelbrot set is universal. In *The Mandelbrot set, theme and variations*, volume 274 of *London Math. Soc. Lecture Note Ser.*, pages 1–17. Cambridge Univ. Press, Cambridge, 2000.
- [Mil06] John Milnor. *Dynamics in one complex variable*, volume 160 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, third edition, 2006.
- [Tan00] Tan Lei. *The Mandelbrot set is universal*, volume 274 of *London Math. Soc. Lecture Note Ser.* Cambridge Univ. Press, Cambridge, 2000.

DEPARTAMENT D'ENGINYERIA INFORMÀTICA I MATEMÀTIQUES, UNIVERSITAT ROVIRA I VIRGILI, 43007 TARRAGONA, CATALONIA, SPAIN
E-mail address: `antonio.garijo@urv.cat`

DEPARTAMENT DE MATEMÀTICA I INFORMÀTICA AND GRADUATE SCHOOL OF MATHEMATICS, UNIVERSITAT DE BARCELONA, 08007 BARCELONA, CATALONIA, SPAIN
E-mail address: `xavier.jarque@ub.edu`

DEPARTAMENT D'ENGINYERIA INFORMÀTICA I MATEMÀTIQUES, UNIVERSITAT ROVIRA I VIRGILI, 43007 TARRAGONA, CATALONIA, SPAIN
E-mail address: `jordi.villadelprat@urv.cat`