

An epigenetic biomarker of aging for lifespan and healthspan

Morgan E. Levine¹, Ake T. Lu¹, Austin Quach¹, Brian H. Chen², Themistocles L. Assimes³, Stefania Bandinelli⁴, Lifang Hou⁵, Andrea A. Baccarelli⁶, James D. Stewart⁷, Yun Li⁸, Eric A. Whitsel^{7,9}, James G Wilson¹⁰, Alex P Reiner¹¹, Abraham Aviv¹², Kurt Lohman¹³, Yongmei Liu¹⁴, Luigi Ferrucci^{2,*}, Steve Horvath^{1,15,*}

¹Department of Human Genetics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA 90095, USA

²Longitudinal Studies Section, Translational Gerontology Branch, National Institute on Aging, National Institutes of Health, USA. Baltimore, MD 21224, USA

³Department of Medicine, Stanford University School of Medicine, Stanford, CA 94305, USA

⁴Geriatric Unit, Azienda Toscana Centro, Florence, Italy

⁵Center for Population Epigenetics, Robert H. Lurie Comprehensive Cancer Center and Department of Preventive Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL 60611, USA

⁶Laboratory of Environmental Epigenetics, Departments of Environmental Health Sciences and Epidemiology, Columbia University Mailman School of Public Health, New York, NY 10032, USA

⁷Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC 27599, USA

⁸Department of Genetics, Department of Biostatistics, Department of Computer Science, University of North Carolina, Chapel Hill, NC 27599, USA

⁹Department of Medicine, School of Medicine, University of North Carolina, Chapel Hill, NC 27599, USA

¹⁰Department of Physiology and Biophysics, University of Mississippi Medical Center, Jackson, MS 39216, USA

¹¹Public Health Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, WA 98109, USA

¹²Center of Human Development and Aging, New Jersey Medical School, Rutgers State University of New Jersey, Newark, NJ 07103, USA

¹³Department of Biostatistics, Division of Public Health Sciences, Wake Forest School of Medicine, Winston-Salem, NC 27157, USA

¹⁴Department of Epidemiology & Prevention, Division of Public Health Sciences, Wake Forest School of Medicine, Winston-Salem, NC 27157, USA

¹⁵Department of Biostatistics, Fielding School of Public Health, University of California Los Angeles, Los Angeles, CA 90095, USA

*Co-senior authors

Correspondence to: Steve Horvath; email: shorvath@mednet.ucla.edu

Keywords: epigenetic clock, DNA methylation, biomarker, healthspan

Received: March 20, 2018

Accepted: April 8, 2018

Published: April 17, 2018

Copyright: Levine et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Identifying reliable biomarkers of aging is a major goal in geroscience. While the first generation of epigenetic biomarkers of aging were developed using chronological age as a surrogate for biological age, we hypothesized that incorporation of composite clinical measures of phenotypic age that capture differences in lifespan and healthspan may identify novel CpGs and facilitate the development of a more powerful epigenetic biomarker of

aging. Using an innovative two-step process, we develop a new epigenetic biomarker of aging, DNAm PhenoAge that strongly outperforms previous measures in regards to predictions for a variety of aging outcomes, including all-cause mortality, cancers, healthspan, physical functioning, and Alzheimer's disease. While this biomarker was developed using data from whole blood, it correlates strongly with age in every tissue and cell tested. Based on an in-depth transcriptional analysis in sorted cells, we find that increased epigenetic, relative to chronological age, is associated with increased activation of pro-inflammatory and interferon pathways, and decreased activation of transcriptional/translational machinery, DNA damage response, and mitochondrial signatures. Overall, this single epigenetic biomarker of aging is able to capture risks for an array of diverse outcomes across multiple tissues and cells, and provide insight into important pathways in aging.

INTRODUCTION

One of the major goals of geroscience research is to define 'biomarkers of aging'[1, 2], which can be thought of as individual-level measures of aging that capture inter-individual differences in the timing of disease onset, functional decline, and death over the life course. While chronological age is arguably the strongest risk factor for aging-related death and disease, it is important to distinguish chronological time from biological aging. Individuals of the same chronological age may exhibit greatly different susceptibilities to age-related diseases and death, which is likely reflective of differences in their underlying biological aging processes. Such biomarkers of aging will be crucial to enable evaluation of interventions aimed at promoting healthier aging, by providing a measurable outcome, which unlike incidence of death and/or disease, does not require extremely long follow-up observation.

One potential biomarker that has gained significant interest in recent years is DNA methylation (DNAm). Chronological time has been shown to elicit predictable hypo- and hyper-methylation changes at many regions across the genome [3-7], and as a result, the first generation of DNAm based biomarkers of aging were developed to predict chronological chronological age [8-13]. The blood-based algorithm by Hannum [10] and the multi-tissue algorithm by Horvath [11] produce age estimates (DNAm age) that correlate with chronological age well above $r=0.90$ for full age range samples. Nevertheless, while the current epigenetic age estimators exhibit statistically significant associations with many age-related diseases and conditions [14-27], the effect sizes are typically small to moderate. One explanation is that using chronological age as the reference, by definition, may exclude CpGs whose methylation patterns don't display strong time-dependent changes, but instead signal the departure of biological age from chronological age. Thus, it is important to not only capture CpGs that display changes with chronological time, but also those that account for differences in risk and physiological status among individuals of the same chronological age.

Previous work by us and others have shown that "phenotypic aging measures", derived from clinical biomarkers [28-32], strongly predict differences in the risk of all-cause mortality, cause-specific mortality, physical functioning, cognitive performance measures, and facial aging among same-aged individuals. What's more, in representative population data, some of these measures have been shown to be better indicators of remaining life expectancy than chronological age [28], suggesting that they may be approximating individual-level differences in biological aging rates. As a result, we hypothesize that a more powerful epigenetic biomarker of aging could be developed by replacing prediction of chronological age with prediction of a surrogate measure of "phenotypic age" that, in and of itself, differentiates morbidity and mortality risk among same-age individuals.

RESULTS

Overview of the statistical model and analysis

Our development of the new epigenetic biomarker of aging proceeded along three main steps (Fig. 1). In step 1, a novel measure of 'phenotypic age' was developed using clinical data from the third National Health and Nutrition Examination Survey (NHANES). Details on the phenotypic age estimator can be found in Table 1 and in Supplement 1. In step 2, DNAm from whole blood was used to predict phenotypic age, such that:

$$\begin{aligned} \text{DNAm PhenoAge} &= \text{intercept} + \text{CpG}_1 \times \beta_1 + \text{CpG}_2 \\ &\quad \times \beta_2 + \dots \text{CpG}_{513} \times \beta_{513} \end{aligned}$$

The coefficient values of this model can be found in Supplement 2 (Table S6). Predicted estimates from this model represent a person's epigenetic age, which we refer to as 'DNAm PhenoAge'. Using multiple independent datasets, we then tested whether DNAm PhenoAge was associated with a number of aging-related outcomes. We also tested whether it differed as a function of social, behavioral, and demographic characteristics, and whether it was applicable to tissues other than whole blood.

Finally, in step 3, we examine the underlying biology of the 513 CpGs in the DNAm PhenoAge measure by examining differential expression, GO and pathway enrichment, chromosomal locations, and heritability.

Estimating phenotypic age from clinical biomarkers

For step 1, NHANES III was used to generate a measure of phenotypic age. NHANES III is a nationally-representative sample, with over twenty-three years of mortality follow-up, from which our analytical sample included 9,926 adults with complete biomarker data. A Cox penalized regression model—where the hazard of mortality was regressed on forty-two clinical markers and chronological age—was used to select variables for inclusion in our phenotypic age score. The forty-two biomarkers considered represent

those that were available in both NHANES III and IV. Based on 10-fold cross-validation, ten variables (including chronological age) were selected for the phenotypic age predictor (Table 1, Table S1). These nine biomarkers and chronological age were then combined in a phenotypic age estimate (in units of years) as detailed in Methods.

Validation data for phenotypic age came from NHANES IV, and included up to 12 years of mortality follow-up for n=6,209 national representative US adults. In this population, phenotypic age is correlated with chronological age at $r=0.94$. Results from all-cause and cause-specific (competing risk) mortality predictions, adjusting for chronological age (Table 2), show that a one year increase in phenotypic age is associated with a 9% increase in the risk of all-cause




Figure 1. Roadmap for developing DNAm PhenoAge. The roadmap depicts our analytical procedures. In step 1, we developed an estimate of 'Phenotypic Age' based on clinical measure. Phenotypic age was developed using the NHANES III as training data, in which we employed a proportional hazard penalized regression model to narrow 42 biomarkers to 9 biomarkers and chronological age. This measure was then validated in NHANES IV and shown to be a strong predictor of both morbidity and mortality risk. In step 2, we developed an epigenetic biomarker of phenotypic age, which we call DNAm PhenoAge, by regressing phenotypic age (from step 1) on blood DNA methylation data, using the InCHIANTI data. This produced an estimate of DNAm PhenoAge based on 513 CpGs. We then validated our new epigenetic biomarker of aging, DNAm PhenoAge, using multiple cohorts, aging-related outcomes, and tissues/cells. In step 3, we examined the underlying biology of the 513 CpGs and the composite DNAm PhenoAge measure, using a variety of complementary data (gene expression, blood cell counts) and various genome annotation tools including chromatin state analysis and gene ontology enrichment.

mortality (HR=1.09, p=3.8E-49), a 9% increase in the risk of mortality from aging-related diseases (HR=1.09, p=4.5E-34), a 10% increase in the risk of CVD mortality (HR=1.10, p=5.1E-17), a 7% increase in the risk of cancer mortality (HR=1.07, p=7.9E-10), a 20% increase in the risk of diabetes mortality (HR=1.20, p=1.9E-11), and a 9% increase in the risk of chronic lower respiratory disease mortality (HR=1.09, p=6.3E-4). Further, phenotypic age is highly associated with comorbidity count (p=3.9E-21) and physical functioning measures (p=2.1E-10, Supplement 1: Fig. S1).

An epigenetic biomarker of aging (DNAm PhenoAge)

For step 2 (Fig. 1), data from n=456 participants at two time-points in the Invecchiare in Chianti (InCHIANTI) study was used to relate blood DNAm levels to phenotypic age. InCHIANTI was used as training data for the new epigenetic biomarker because the study assessed all clinical measures needed to estimate phenotypic age, contained data on DNAm, and had a large age range population (21-100 years). A total of 20,169 CpGs were considered when generating the new

DNAm measure. They represented those CpGs available on all three chips (27k, 450k, EPIC), so as to facilitate usability across platforms. Elastic net regression, with 10-fold cross-validation, produced a model in which phenotypic age is predicted by DNAm levels at 513 of the 20,169 CpGs. The linear combination of the weighted 513 CpGs yields a DNAm based estimator of phenotypic age that we refer to as ‘DNAm PhenoAge’ (mean=58.9, s.d.=18.2, range=9.1-106.1), in contrast to the previously published Hannum and Horvath ‘DNAm Age’ measures.

While our new clock was trained on cross-sectional data in InCHIANTI, we capitalized on the repeated time-points to test whether changes in DNAm PhenoAge are related to changes in phenotypic age. As expected, between 1998 and 2007, mean change in DNAm PhenoAge was 8.51 years, whereas mean change in clinical phenotypic age was 8.88 years. Moreover, participants’ clinical phenotypic age (adjusting for chronological age) at the two time-points was correlated at r=0.50, whereas participants’ DNAm PhenoAge (adjusting for chronological age) at the two time-points was correlated at r=0.68 (Supplement 1: Fig. S2). We also find that the change in phenotypic age between

Table 1. Phenotypic aging measures and Gompertz coefficients.

Variable		Units	Weight
Albumin	Liver	g/L	-0.0336
Creatinine	Kidney	umol/L	0.0095
Glucose, serum	Metabolic	mmol/L	0.1953
C-reactive protein (log)	Inflammation	mg/dL	0.0954
Lymphocyte percent	Immune	%	-0.0120
Mean (red) cell volume	Immune	fL	0.0268
Red cell distribution width	Immune	%	0.3306
Alkaline phosphatase	Liver	U/L	0.0019
White blood cell count	Immune	1000 cells/uL	0.0554
Age		Years	0.0804

Table 2. Mortality validations for phenotypic age.

Mortality Cause	Cases	HR	P-Value
All-Cause	1052	1.09	3.8E-49
Aging-Related	661	1.09	4.5E-34
CVD	272	1.10	5.1E-17
Cancer	265	1.07	7.9E-10
Alzheimer's	30	1.04	2.6E-1
Diabetes	41	1.20	1.9E-11
Chronic lower respiratory diseases	53	1.09	6.3E-4

1998 and 2007 is highly correlated with the change in DNAm PhenoAge between these two time-points ($r=0.74$, $p=3.2E-80$, Supplement 1: Fig. S2).

DNAm PhenoAge strongly relates to all-cause mortality

In step 2 (Fig. 1), the epigenetic biomarker, DNAm PhenoAge, was calculated in five independent large-scale samples—two samples from Women’s Health Initiative (WHI) ($n=2,016$; and $n=2,191$), the Framingham Heart Study (FHS) ($n=2,553$), the Normative Aging Study (NAS) ($n=657$), and the Jackson Heart Study (JHS) ($n=1,747$). The first four studies used the Illumina 450K array while the JHS employed the latest Illumina EPIC array platform. In these studies, DNAm PhenoAge correlated with chronological age at $r=0.66$ in WHI (Sample 1), $r=0.69$ in WHI (Sample 2), $r=0.78$ in FHS, $r=0.62$ in the NAS, and $r=0.89$ in JHS. The five validation samples were then used to assess the effects of DNAm PhenoAge on mortality in comparison to the Horvath and Hannum DNAm Age measures. DNAm PhenoAge was significantly associated with subsequent mortality risk in all studies (independent of chronological age), such that, a one year increase in DNAm PhenoAge is associated with a 4.5% increase in the risk of all-cause mortality (Meta(FE)=1.045, Meta $p=7.9E-47$, Fig. 2). To better conceptualize what this increase represents, we compared the predicted life expectancy and mortality risk for person’s representing the top 5% (fastest agers), the average, and the bottom 5% (slowest agers). Results suggest that those in the top 5% of fastest agers have a mortality hazard of death that is about 1.62 times that of the average person, i.e. the hazard of death is 62% higher than that of an average person. Further, contrasting the 5% fastest agers with the 5% slowest agers, we find that the hazard of death of the fastest agers is 2.58 times higher than that of the bottom 5% slowest agers ($HR=1.045^{11.0}/1.045^{-10.5}$). Additionally, both observed and predicted Kaplan-Meier survival estimates showed that faster agers had much lower life expectancy and survival rates compared to average and/or slow agers (Fig. 2).

As shown in Fig. 2, the DNAm age based measures from Hannum and Horvath also related to all-cause mortality, consistent with what has been reported previously [15, 19, 23, 33, 34]. To directly compare the three epigenetic measures, we contrasted their accuracy in predicting 10-year and 20-year mortality risk, using receiver operating characteristics (ROC) curves. DNAm PhenoAge (adjusted for age) predicts both 10-year mortality and 20-year mortality significantly better than the Horvath and Hannum DNAmAge measures (Supplement 1: Table S2). When examining a model that

includes all three measures (Supplement 1: Table S3), we find that only DNAm PhenoAge is positively associated with mortality ($HR=1.04$, $p=1.33E-8$), whereas Horvath DNAm Age is now negatively associated ($HR=0.98$, $p=2.72E-2$), and Hannum DNAm Age has no association ($HR=1.01$, $p=4.66E-1$).

DNAm PhenoAge strongly relates to aging-related morbidity

Given that aging is believed to also influence disease incidence/prevalence, we examined whether DNAm PhenoAge relates to diverse age-related morbidity outcomes. We observe strong associations between DNAm PhenoAge and a variety of other aging outcomes using the same five validation samples (Table 3). For instance, independent of chronological age, higher DNAm PhenoAge is associated with an increase in a person’s number of coexisting morbidities ($\beta=0.008$ to 0.031 ; Meta P -value= $1.95E-20$), a decrease in likelihood of being disease-free ($\beta=-0.002$ to -0.039 ; Meta P -value= $2.10E-10$), an increase in physical functioning problems ($\beta=-0.016$ to -0.473 ; Meta P -value= $2.05E-13$), an increase in the risk of coronary heart disease (CHD) risk ($\beta=0.016$ to 0.073 ; Meta P -value= $3.35E-11$).

DNAm PhenoAge and smoking

Cigarette exposure has been shown to have an epigenetic fingerprint[35-37], which has been reflected in previous DNAm risk predictors[38]. Similarly, we find that DNAm PhenoAge significantly differs between never ($n=1,097$), current ($n=209$), and former smokers ($n=710$) ($p=0.0033$) (Supplement 1, Fig. S3A); however, conversely, we do not find a robust association between pack-years and DNAm PhenoAge (Supplement 1, Fig. S3B-D). Given the association between DNAm PhenoAge and smoking, we re-evaluated the morbidity and mortality associations (fully-adjusted) in our four samples, stratifying by smoking status (Supplement 1: Fig. S4 and Table S4). We find that DNAm PhenoAge is associated with mortality among both smokers (adjusted for pack-years) (Meta(FE)=1.050, Meta $p=7.9E-31$), and non-smokers (Meta(FE)=1.033, Meta $p=1.2E-10$). DNAm PhenoAge relates to the number of coexisting morbidities, physical functioning status, disease free status, and CHD for both smokers and non-smokers (Supplement 1: Table S4). In previous work we showed that Horvath DNAm age of blood predicts lung cancer risk in the first WHI sample [20]. Using the same data, we find that a one year increase in DNAm PhenoAge (adjusting for chronological age, race/ethnicity, pack-years, and smoking status) is associated with a 5% increase in the risk of lung cancer incidence and/or


mortality (HR=1.05, p=0.031). Further, when restricting the model to current smokers only, we find that the effect of DNAm PhenoAge on future lung cancer incidence and/or mortality is even stronger (HR=1.10, p=0.014).

DNAm PhenoAge in other tissues


One advantage of developing biological aging estimates based on molecular markers (like DNAm), rather than clinical risk measures (e.g. those in the phenotypic age variable), is that they may lend themselves to measuring

tissue/cell specific aging. Although DNAm PhenoAge was developed using samples from whole blood, our empirical results show that it strongly correlates with chronological age in a host of different tissues and cell types (Fig. 3). For instance, when examining all tissues concurrently, the correlation between DNAm PhenoAge and chronological age was 0.71. Age correlations in brain tissue ranged from 0.54 to 0.92, while correlations were also found in breast ($r=0.47$), buccal cells ($r=0.88$), dermal fibroblasts ($r=0.87$), epidermis ($r=0.84$), colon ($r=0.88$), heart ($r=0.66$), kidney ($r=0.64$), liver ($r=0.80$), lung ($r=0.55$), and saliva ($r=0.81$).

A



B



C




Figure 2. Mortality Prediction by DNAm PhenoAge. (A) Using five samples from large epidemiological cohorts—two samples from the Women's health Initiative, the Framingham Heart Study, the Normative Aging Study, and the Jackson Heart Study—we tested whether DNAm PhenoAge was predictive of all-cause mortality. The Fig. displays a forest plot for fixed-effect meta-analysis, based on Cox proportional hazard models, and adjusting for chronological age. Results suggest that DNAm PhenoAge is predictive of mortality in all samples, and that overall, a one year increase in DNAm PhenoAge is associated with a 4.5% increase in the risk of death ($p=9.9E-47$). This is contrasted against the first generation of epigenetic biomarkers of aging by Hannum and Horvath, which exhibit less significant associations with lifespan ($p=1.7E-21$ and $p=4.5E-5$, respectively). (B and C) Using the WHI sample 1, we plotted Kaplan-Meier survival estimates using actual data from the fastest versus the slowest agers (panel B). We also applied the equation from the proportional hazard model to predict remaining life expectancy and plotted predicted survival assuming a chronological age of 50 and a DNAm PhenoAge of either 40 (slow ager), 50 (average ager), or 60 (fast ager) (panel C). Median life expectancy at age 50 was predicted to be approximately 81 years for the fastest agers, 83.5 years for average agers, and 86 years for the slowest agers.

Table 3. Morbidity validation for DNAm PhenoAge.

Sample	Comorbidity		Disease Free		CHD Risk		Physical Functioning	
	Coefficient	P-value	Coefficient	P-value	Coefficient	P-value	Coefficient	P-value
DNAm PhenoAge								
WHI BA23 White	0.008	2.38E-01	-0.002	3.82E-01	0.016	5.36E-02	-0.396	1.04E-04
WHI BA23 Black	0.013	6.15E-02	-0.006	2.40E-02	0.021	2.02E-02	-0.423	4.50E-04
WHI BA23 Hispanic	0.024	1.64E-02	-0.004	3.67E-01	0.033	5.07E-02	-0.329	7.37E-02
WHI EMPC White	0.031	2.95E-07	-0.026	1.63E-02	0.023	1.89E-01	-0.361	3.81E-05
WHI EMPC Black	0.014	7.67E-02	-0.023	6.98E-02	0.048	2.27E-02	-0.473	3.75E-04
WHI EMPC Hispanic	0.003	7.83E-01	0.002	9.28E-01	0.073	1.98E-01	-0.377	6.54E-02
FHS	0.022	3.93E-07	-0.034	1.59E-03	0.028	5.47E-06	-0.016	4.60E-01
NAS	0.023	7.59E-06	-0.062	2.00E-04	0.030	2.27E-02	NA	NA
JHS	0.018	1.86E-08	-0.039	5.92E-05	0.033	4.73E-02	NA	NA
Meta P-value (Stouffer)	1.95E-20		2.14E-10		3.35E-11		2.05E-13	
DNAmAge Hannum								
WHI BA23 White	0.007	3.90E-01	-0.003	3.48E-01	0.013	2.36E-01	-0.399	2.90E-03
WHI BA23 Black	0.022	2.72E-02	-0.007	6.03E-02	0.015	2.67E-01	-0.345	4.29E-02
WHI BA23 Hispanic	0.010	4.33E-01	-0.010	6.24E-02	0.011	6.10E-01	-0.599	1.16E-02
WHI EMPC White	0.025	1.53E-03	-0.020	1.55E-01	0.022	3.30E-01	-0.284	1.43E-02
WHI EMPC Black	0.022	6.34E-02	-0.008	6.62E-01	0.055	6.12E-02	-0.323	9.56E-02
WHI EMPC Hispanic	-0.012	4.17E-01	0.035	2.09E-01	-0.012	8.85E-01	-0.345	2.54E-01
FHS	0.019	5.94E-04	-0.030	2.55E-02	0.022	1.55E-02	0.040	1.32E-01
NAS	0.009	2.19E-01	-0.026	2.26E-01	0.025	1.83E-01	NA	NA
JHS	0.020	2.09E-05	-0.036	9.91E-03	0.086	1.64E-04	NA	NA
Meta P-value (Stouffer)	1.50E-08		1.64E-04		1.40E-05		2.03E-05	
DNAmAge Horvath								
WHI BA23 White	0.007	3.49E-01	-0.004	1.69E-01	0.001	9.12E-01	-0.440	5.10E-04
WHI BA23 Black	0.018	3.96E-02	-0.006	6.25E-02	0.009	4.07E-01	-0.305	4.52E-02
WHI BA23 Hispanic	0.012	3.65E-01	-0.007	1.86E-01	-0.001	9.78E-01	-0.204	4.12E-01
WHI EMPC White	0.031	1.99E-04	-0.043	5.56E-03	0.000	9.88E-01	-0.288	1.74E-02
WHI EMPC Black	0.016	1.93E-01	-0.003	8.56E-01	0.033	2.87E-01	-0.144	4.68E-01
WHI EMPC Hispanic	-0.025	8.99E-02	-0.016	5.70E-01	-0.064	4.63E-01	-0.012	9.70E-01
FHS	0.011	5.82E-02	-0.021	8.34E-02	0.007	5.19E-01	0.027	3.16E-01
NAS	0.011	7.90E-02	-0.039	4.53E-02	0.006	7.14E-01	NA	NA
JHS	0.014	2.03E-03	-0.040	1.78E-03	0.049	3.93E-02	NA	NA
Meta P-value (Stouffer)	3.26E-06		6.36E-07		1.49E-01		1.43E-03	

Alzheimer's disease and brain samples

Based on the accuracy of the age prediction in other tissues/cells, we examined whether aging in a given tissue was associated with tissue-associated outcomes. For instance, using data from approximately 700 postmortem samples from the Religious Order Study (ROS) and the Memory and Aging Project (MAP) [39, 40] we tested the association between pathologically diagnosed Alzheimer's disease and DNAm PhenoAge in dorsolateral-

prefrontal cortex (DLPFX). Results suggest (Fig. 4) that those who are diagnosed with Alzheimer's disease (AD), based on postmortem autopsy, have DLPFX that appear more than one year older than same aged individuals who are not diagnosed with AD postmortem ($p=4.6E-4$). Further, age adjusted DNAm PhenoAge was found to be positively associated with neuropathological hallmarks of Alzheimer's disease, such as amyloid load ($r=0.094$, $p=0.012$), neuritic plaques ($r=0.11$, $p=0.0032$), and neurofibrillary tangles ($r=0.10$, $p=0.0073$).




Figure 3. Chronological age versus DNAm PhenoAge in a variety of tissues and cells. Although DNAm PhenoAge was developed using methylation data from whole blood, it also tracks chronological age in a wide variety of tissues and cells. **(A)** The correlation across all tissues/cells we examined is $r=0.71$, $p<1e-200$. **(B-ZJ)** report results in different sources of DNA as indicated in panel headings. The numbers correspond to the data sets from (Horvath 2013). Overall, correlations range from $r=0.35$ (breast, panel O) to $r=0.92$ (temporal cortex in brain, panel L).

Lifestyle and demographic variables

In evaluating the relationship between DNAm PhenoAge in blood and additional characteristics we observe significant differences between racial/ethnic groups ($p=5.1E-5$), with non-Hispanic blacks having the highest DNAm PhenoAge on average, and non-Hispanic whites having the lowest (Supplement 1: Fig. S5). We also find evidence of social gradients in DNAm PhenoAge, such that those with higher education ($p=6E-9$) and higher income ($p=9E-5$) appear younger (Figure 5). DNAm PhenoAge relates to exercise and dietary habits, such that increased exercise ($p=7E-5$) and markers of fruit/vegetable consumption (such as carotenoids, $p=2E-27$) are associated with lower DNAm PhenoAge (Figure 5, Supplement 1: Fig. S6). Cross-sectional studies in the WHI also revealed that DNAmPhenoAge acceleration is positively correlated with C-reactive protein ($r=0.18$, $p=5E-22$, Figure 5), insulin ($r=0.15$, $p=2E-20$), glucose ($r=0.10$, $p=2E-10$), triglycerides ($r=0.09$, $p=5E-9$), waist to hip ratio ($r=0.15$, $p=5E-22$) but it is negatively correlated with HDL cholesterol ($r=-0.09$, $p=7E-9$).

DNAm PhenoAge and Immunosenescence

To test the hypothesis that DNAm PhenoAge captures aspects of age-related decline of the immune system, we correlated DNAm PhenoAge with estimated blood cell count (Supplement 1, Fig. S7). After adjusting for age, we find that DNAm PhenoAgeAccel is negatively correlated with naïve CD8+ T cells ($r=-0.35$, $p=9.2E-65$), naïve CD4+ T cells ($r=-0.29$, $p=4.2E-42$), CD4+ helper T cells ($r=-0.34$, $p=3.6E-58$), and B cells ($r=-0.18$, $p=8.4E-17$). Further, DNAm PhenoAgeAccel is positively correlated with the proportion of granulocytes ($r=0.32$, $p=2.3E-51$), exhausted CD8+ (defined as CD28-CD45RA-) T cells ($r=0.20$, $p=1.9E-20$), and plasma blast cells ($r=0.26$, $p=6.7E-34$). These results are consistent with age related changes in blood cells [41] and suggest that DNAm PhenoAge may capture aspects of immuno-senescence in blood. However, three lines of evidence suggest that DNAm PhenoAge is not simply a measure of immunosenescence. First, another measure of immunosenescence, leukocyte telomere length, is only weakly correlated with DNAm PhenoAgeAccel ($r=-0.13$ $p=0.00019$ in the WHI; $r=$

0.087, $P=7.6E-3$ in Framingham Heart study; JHS $p=7.83E-7$, Supplement 1, Fig. S8). Second, the strong association between DNAm PhenoAge and mortality does not simply reflect changes in blood cell composition, as can be seen from the fact that in Supplement 1, Fig. S9 the robust association remains even after adjusting for estimates of seven blood cell count measures ($\text{Meta}(FE)=1.036$, $\text{Meta } p=5.6E-21$). Third, DNAmPhenoAge correlates with chronological age in non-blood tissue.

DNA sequence characteristics of the 513 CpGs in DNAm PhenoAge

Of the 513 CpGs in DNAm PhenoAge, we find that, 41 CpGs were also in the Horvath DNAm age measure (Supplement 2: Table S6). This represents a 4.88-fold increase over what would be expected by chance ($p=8.97E-15$). Of the 41 overlapping CpGs, the average absolute value for their age correlations was $r=0.40$, and 31 had age correlations with absolute values in the top 20% of what is found among the 513 CpGs in the DNAm PhenoAge score. We also observed 6 CpGs that

overlapped between the Hannum DNAm Age score and the DNAm PhenoAge score—five of which were also found in the Horvath DNAm Age measure. All six CpGs had extremely high age correlations (half positive, half negative), with absolute values between $r=0.49$ and $r=0.76$. The five CpGs that are found in all three epigenetic aging measures were: cg05442902 (*P2RXLI*), cg06493994 (*SCGN*), cg09809672 (*EDARADD*), cg19722847 (*IPO8*), and cg22736354 (*NHLRC1*).

Several additional DNAm biomarkers have been described in the literature [12, 13]. A direct comparison of 6 DNAm biomarkers (including DNAm PhenoAge) reveals that DNAm PhenoAge stands out in terms of its predictive accuracy for lifespan, its relationship with smoking status, its relationship with leukocyte telomere length, naïve CD8+ T cells and CD4+ T cells (Supplement 1: Table S5).

Next, we conducted a functional enrichment analysis of the chromosomal locations of the 513 CpGs and found that 149 CpGs whose age correlation exceeded 0.2 tended to be located in CpG islands ($p=0.0045$, Supple-




Figure 4. DNAm PhenoAge measured in dorsolateral prefrontal cortex relates to Alzheimer's disease and related neuropathologies. Using postmortem data from the Religious Order Study (ROS) and the Memory and Aging Project (MAP), we find a moderate/high correlation between chronological age and DNAm PhenoAge (panel A). We also estimate the proportion of neurons via the CETS algorithm and show that it correlates with DNAm PhenoAge (B). Further, the correlation between chronological age and DNAm PhenoAge is increased after adjusting for the estimated proportion of neurons in each sample (panel C). We also find that DNAm PhenoAge is significantly higher ($p=0.00046$) among those with Alzheimer's disease versus controls (panel D), and that it positively correlates with amyloid load ($p=0.012$, panel E), neuritic plaques ($p=0.0032$, panel F), diffuse plaques ($p=0.036$, panel G), and neurofibrillary tangles ($p=0.0073$, panel H).

ment 1: Fig. S10) and were significantly enriched with polycomb group protein targets ($p=8.7E-5$, Supplement 1: Fig. S10), which in line with results of epigenome wide studies of aging effects [4, 5, 42].

Transcriptional and genetic studies of DNAm PhenoAge

Using the genome-wide data from FHS and WHI, we estimated the heritability of DNAm PhenoAge. The heri-

tability estimated by the SOLAR polygenic model for those of European ancestry in the FHS was $h^2=0.33$, while the heritability estimated for those of European ancestry in WHI, using GCTA-GREML analysis [43] was $h^2=0.51$.

Using the monocyte data mentioned above, as well as PBMC expression data on 2,188 persons from the FHS, we conducted a transcriptional analysis to identify differential expression associated with DNAm PhenoAge-

Pooled WHI samples Adjusted for ethnicity and dataset			
	n	μ	PhenoAgeAccel
			bicor p
Diet	log2(Total energy)	3700	10.58 0.01 0.40
	Carbohydrate	3700	48.64 -0.02 0.23
	Protein	3700	16.39 -0.01 0.45
	Fat	3700	35.05 0.03 0.12
	log2(1+Red meat)	3700	0.69 0.05 2E-3
	log2(1+Poultry)	3700	0.39 -0.02 0.14
	log2(1+Fish)	3700	0.25 -0.03 0.10
	log2(1+Dairy)	3700	1.21 -0.01 0.76
	log2(1+Whole grains)	3700	1.00 -0.02 0.19
	log2(1+Nuts)	3700	0.10 -0.04 0.03
Dietary biomarkers	log2(Fruits)	3700	0.57 -0.02 0.13
	log2(Vegetables)	3700	0.71 -0.03 0.09
	Retinol	2267	0.58 0.03 0.17
	Mean carotenoids	2266	0.06 -0.22 2E-27
	Lycopene	2267	0.36 -0.11 3E-7
	log2(alpha-Carotene)	2267	-4.21 -0.19 5E-20
	log2(beta-Carotene)	2266	-2.15 -0.18 2E-17
	log2(Lutein+Zeaxanthin)	2267	-2.38 -0.17 2E-16
Measurements	log2(beta-Cryptoxanthin)	2267	-3.76 -0.17 2E-15
	log2(alpha-Tocopherol)	2267	3.87 -0.03 0.19
	log2(gamma-Tocopherol)	2267	0.88 0.07 6E-4
	log2(C-reactive protein)	2809	1.57 0.18 5E-22
	log2(Insulin)	4042	5.78 0.15 2E-20
	log2(Glucose)	4144	6.57 0.10 2E-10
	log2(Triglyceride)	4148	7.02 0.09 5E-9
	Total cholesterol	4148	224.00 -0.04 4E-3
	LDL cholesterol	4084	140.00 -0.05 2E-3
	HDL cholesterol	4145	53.00 -0.09 7E-9
Socio-behavioral	log2(Creatinine)	2748	-0.43 0.01 0.52
	Systolic blood pressure	4177	129.00 0.08 1E-6
	Diastolic blood pressure	4178	76.00 0.02 0.12
	log2(Waist / hip ratio)	4037	-0.28 0.15 5E-22
	BMI	4145	28.88 0.13 5E-16
	Education	4143	7.00 -0.09 6E-9
	Income	4054	3.00 -0.06 9E-5

Figure 5. Lifestyle factors versus DNAm PhenoAge acceleration in blood in the WHI. In this cross sectional analysis, the correlation test analysis (bicor, biweight midcorrelation) between select variables and DNAm PhenoAgeAccel reveals that education, income, exercise, proxies of fruit/vegetable consumption, and HDL cholesterol are negatively associated (blue) with DNAm PhenoAge, i.e. younger epigenetic age. Conversely, CRP, insulin, glucose, triglycerides, BMI, waist-to-hip ratio, systolic blood pressure, and smoking have a positive association (red) with DNAm PhenoAge. All results have been adjusted for ethnicity and batch/data set. Similar results based on multivariate regression models can be found in Supplementary Figure 6B.

Accel (Supplement 3: Table S7). Overall, we find that genes show similar associations with chronological age and DNAm PhenoAgeAccel. DNAm PhenoAgeAccel represents aging differences among same-aged individuals and is adjusted so as to exhibit a correlation of $r=0.0$ with chronological age. Thus, this observation suggests that genes whose transcription increases with age are upregulated among epigenetically older compared to epigenetically younger persons of the same chronological age (Supplement 1: Fig. S11); the same applies for genes that show decreases with chronological age being downregulated in epi-genetically older versus younger persons of the same age.

Using the transcriptional data from monocytes described above (adjusting for array, sex, race/ethnicity, age, and imputed cell counts), we tested for GO enrichment among genes that are positively associated with DNAm PhenoAge and those that are negatively associated with DNAm PhenoAge (Supplement 4: Table S8). Among those with positive aging associations (over-expression among epigenetically older individuals), we observed enrichment for a number of pro-inflammatory signaling pathways. These pathways included, but are not limited to: multiple toll-like receptor signaling pathways (7,9,3,2), regulation of inflammatory response, JAK-STAT cascade, response to lipopolysaccharide, tumor necrosis factor-mediated signaling pathway, and positive regulation of NF-kappaB transcription factor activity. Additionally, positively associated genes were also enriched for a number anti-viral response pathways—type I interferon signaling, defense response to virus, interferon-gamma-mediated signaling pathway, cellular response to interferon-alpha, etc. Other interesting GO terms enriched among positively associated genes included: response to nutrient, JAK-STAT cascade involved in growth hormone signaling pathway, multicellular organism growth, and regulation of DNA methylation.

When testing for enrichment among genes that were negatively associated with DNAm PhenoAgeAccel (decreased expression among epigenetically older persons) we observed that many were implicated in processes involving transcriptional and translational machinery, as well as damage recognition and repair. These included: translational initiation; regulation of translational initiation; ribosomal large subunit assembly; ribosomal small subunit assembly; translational elongation; transcription initiation from RNA polymerase I promoter; transcription-coupled nucleotide-excision repair; nucleotide-excision repair, DNA incision, 5'-to lesion; nucleotide-excision repair, DNA damage recognition; DNA damage response, detection of DNA damage; and regulation of DNA damage checkpoint.

DISCUSSION

Using a novel two-step method, we were successful in developing a DNAm based biomarker of aging that is highly predictive of nearly every morbidity and mortality outcome we tested. Training an epigenetic predictor of phenotypic age instead of chronological age led to substantial improvement in mortality/healthspan predictions over the first generation of DNAm based biomarkers of chronological age from Hannum[10], Horvath[11] and other published DNAm biomarkers. In doing so, this is the first study to conclusively demonstrate that DNAm biomarkers of aging are highly predictive of cardiovascular disease and coronary heart disease. DNAm PhenoAge also tracks chronological age and relates to disease risk in samples other than whole blood. Finally, we find that an individual's DNAm PhenoAge, relative to his/her chronological age, is moderately heritable and is associated with activation of pro-inflammatory, interferon, DNAm damage repair, transcriptional/ translational signaling, and various markers of immuno-senescence: a decline of naïve T cells and shortened leukocyte telomere length (Supplementary Information).

The ability of our measure to predict multifactorial aging conditions is consistent with the fundamental underpinnings of Geroscience research [1, 44], which posits that aging mechanisms give rise to multiple pathologies and thus, differences in the rate of aging will have implications for a wide array of diseases and conditions. Further, these results answer a fundamental biological question of whether differences in multi-system dysregulation (estimated using clinical phenotypic age measures), healthspan, and lifespan are reflected at the epigenetic level, in the form of differential DNAm at specific CpG sites.

The improvement over previous epigenetic biomarkers, likely comes down to the types of CpGs selected for the various measures. Only 41 of the 513 CpGs in DNAm PhenoAge were shared with the Horvath clock, while only five CpGs were shared between all three clocks (DNAm PhenoAge, Horvath, and Hannum). In general, these CpGs did not tend to be drivers of the DNAm PhenoAge score, and instead represented those with large age correlations. This may explain the improvements of DNAm PhenoAge over previous epigenetic biomarkers of aging. While the previous DNAm age estimators selected CpGs to optimize prediction of chronological age, the CpGs in DNAm PhenoAge were optimized to predict a multi-system proxy of physiological dysregulation (phenotypic age). In doing so, we were able to not only capture CpGs that exhibited strong correlations with age, but also those that captured variations in risk of death and disease

among same aged individuals. In general, the CpGs with the highest weights in the new clock did not correlate with chronological age (Supplement 1: Fig. S12), but instead were related to the difference between phenotypic and chronological age—i.e. divergence in the rate of aging.

While DNAm PhenoAge greatly outperformed all previous DNAm biomarkers of aging (Supplement 1: Table S5), the utility of DNAm PhenoAge for estimating risk does not imply that it should replace clinical biomarkers when it comes to informing medical and health-related decisions. In fact, but perhaps not surprisingly, the phenotypic age measure used to select CpGs is a better predictor of morbidity and mortality outcomes than DNAm PhenoAge. While the addition of error in performing a two-step process, rather than training a DNAm predictor directly on mortality may contribute, we don't believe this accounts for the difference in predictive performance. In fact, a recent DNAm measure by Zhang et al. [38] was trained to directly predict mortality risk, yet it appears to be a weaker predictor than both our DNAm PhenoAge measure and our clinical phenotypic age measure (Supplement 1: Table S9). The first generation of DNAm age estimators only exhibit weak associations with clinical measures of physiological dysregulation [24, 45]. Physiological dysregulation, which is more closely related to our clinical age measure “phenotypic age” than to chronological age, is not only the result of exogenous/endogenous stress factors (such as obesity, infections) but also a result of age related molecular alterations, one example of which are modifications to the epigenome. Over time, dysregulation within organ systems leads to pathogenesis of disease (age-related molecular changes → physiological dysregulation → morbidity → mortality)[46]. However, stochasticity and variability exist at each of these transitions. Therefore, measures of physiological dysregulation, will be better predictors of transition to the next stage in the aging trajectory (i.e. morbidity and mortality) than will measures of age related molecular alterations, like DNAm PhenoAge. Similarly, quantification of disease pathogenesis (cancer stage, Alzheimer's stage) is likely a better predictor of mortality risk than clinical phenotypic aging measures. As a result, clinical phenotypic aging measures may be preferable to epigenetic measures when the goal is risk prediction, and samples come from blood.

That being said, when the aim is to study the mechanisms of the aging process, DNAm measures have advantages over clinical measures. First, they may better capture “pre-clinical aging” and thus may be more suited for differentiating aging in children, young adults, or extremely healthy individuals, for whom

measures like CRP, albumin, creatinine, glucose, etc. are still fairly homogenous. Second, as demonstrated, these molecular measures can capture cell and/or tissue specific aging rates and therefore may also lend themselves to in vitro studies of aging, studies for which blood is not available, studies using postmortem samples, and/or studies comparing aging rates between tissues/cells. While the fundamental drivers of aging are believed to be shared across cells/tissues, that is not to say that all the cells and tissues within an individual will age at the same rate. In fact, it is more likely that individuals will vary in their patterning of aging rates across tissues, and that this will have implications for death and disease risk. Relatedly, it is not known how predictions based on DNAm PhenoAge measures from non-blood samples will compare to phenotypic age predictions. It may be the case that various outcomes will be more tightly related to aging in specific cells/tissues, rather than blood. Finally, examination of DNAm based aging rates facilitates the direct study of the proposed mechanisms of aging, of which “epigenetic alterations” is one of the seven hypothesized “pillars of aging” [1].

While more work needs to be done to model the biology linking DNAm PhenoAge and aging outcomes, we began to explore this using differential expression, functional enrichment, and heritability estimates. Overall, we found that CpGs that had larger increases with aging tended to be located in CpG islands and enriched with polycomb group protein targets, consistent with what has been reported in previous epigenome wide studies of aging effects [4-7, 42]. While typically DNAm of CpG islands and/or polycomb recruitment is linked to transcriptional silencing [47], for the most part, we did not observe associations between DNAm and expression for co-locating CpG-gene pairs—this was also true when only considering CpGs located in islands. These findings may suggest that the genes annotated to the CpGs in our score are not part of the link between changes in DNAm and aging. Nevertheless, we also recognize that these null results could stem from the fact that 1) associations were only tested in monocytes, 2) DNAm and expression represents what is present globally for each sample, rather than on a cell-by-cell basis, and 3) stronger associations between DNAm and gene expression levels may only exist early in life.

Nevertheless, we do identify potentially promising transcriptional pathways when considering DNAm PhenoAge as a whole. For instance, we observe that higher DNAm PhenoAge is associated with increases in the activation of proinflammatory pathways, such as NF-kappaB; increased interferon (IFN) signaling; decreases in ribosomal-related and translational

machinery pathways; and decreases in damage recognition and repair pathways. These findings are consistent with previous work describing aging associated changes, comprising increases in dysregulated inflammatory activation, increased DNA damage, and loss of translational fidelity. For instance, there exists a large body of literature highlighting the importance of an increased low-grade pro-inflammatory status as a driver of the aging process, termed inflammaging [41, 48, 49]. IFN signaling pathways have been shown to be markers of DNA damage and mediators of cellular senescence[50]. Additionally, it has been shown that breakdown of the transcriptional and translational machinery may play a central role in the aging process [51, 52]. For instance, the ribosome is believed to be a key regulator of proteostasis, and in turn, aging [51, 53]. Relatedly, loss of integrity in DNA damage repair pathways is considered another hallmark of the aging process [54-56].

In general, many of these pathways will have implications for adaptation to exogenous and endogenous stressors. Factors related to stress resistance and response have repeatedly been shown to be drivers of differences in lifespan and aging [49, 57-61]. This may partially account for our findings related to smoking. In general, it is not surprising that a biomarker of aging and mortality risk relates to smoking, given that life expectancies of smokers are on average ten years shorter than never smokers, and smoking history is associated with a drastic increase in the risk of a number of age-related conditions. However, perhaps more interestingly, we find that the effects of DNAm PhenoAge on mortality appear to be higher for smokers than non-smokers, which could suggest that DNAm PhenoAge represent differences in innate resilience/ vulnerability to pro-aging stressors, such as cigarette smoke.

Interestingly, we observed moderately high heritability estimates for DNAm PhenoAge. For instance, we estimated that genetic differences accounted for one-third to one-half of the variance in DNAm PhenoAge, relative to chronological age. In moving forward, it will be useful to identify the genetic architecture underlying differences in epigenetic aging. Finally, we reported that individuals' DNAm PhenoAges—relative to their chronological ages—remained fairly stable over a nine-year period. However, it is unclear whether it is attributable to genetic influences, or the fact that social and behavioral characteristics tend to also remain stable for most individuals.

If the goal is to utilize accurate quantifiable measures of the rate of aging, such as DNAm PhenoAge, to assess the efficacy of aging interventions, more work will be needed to evaluate the dynamics of DNAmPhenoAge

following various treatments. For instance, it remains to be seen whether interventions can reverse DNAmPhenoAge in the short term. Along these lines, it will be essential to determine causality—does DNAm drive the aging process, or is it simply a surrogate marker of organismal senescence? If the former is true, DNAm PhenoAge could provide insight into promising targets for therapies aimed at lifespan, and more importantly, healthspan extension.

Overall, DNAm PhenoAge is an attractive composite biomarker that captures organismal age and the functional state of many organ systems and tissues, above and beyond what is explained by chronological time. Our validation studies in multiple large and independent cohorts demonstrate that DNAm PhenoAge is a highly robust predictor of both morbidity and mortality outcomes, and represents a promising biomarker of aging, which may prove to be beneficial to both basic science and translational research.

METHODS

Using the NHANES training data, we applied a Cox penalized regression model—where the hazard of aging-related mortality (mortality from diseases of the heart, malignant neoplasms, chronic lower respiratory disease, cerebrovascular disease, Alzheimer's disease, Diabetes mellitus, nephritis, nephrotic syndrome, and nephrosis) was regressed on forty-two clinical markers and chronological age to select variables for inclusion in our phenotypic age score. Ten-fold cross-validation was employed to select the parameter value, lambda, for the penalized regression. In order to develop a sparse parsimonious phenotypic age estimator (fewer biomarker variables preferred to produce robust results) we selected a lambda of 0.0192, which represented a one standard deviation increase over the lambda with minimum mean-squared error during cross-validation (Supplement 1, Fig. S13). Of the forty-two biomarkers included in the penalized Cox regression model, this resulted in ten variables (including chronological age) that were selected for the phenotypic age predictor.

These nine biomarkers and chronological age were then included in a parametric proportional hazards model based on the Gompertz distribution. Based on this model, we estimated the 10-year (120 months) mortality risk of the j-the individual. Next, the mortality score was converted into units of years (Supplement 1). The resulting phenotypic age estimate was regressed on DNA methylation data using an elastic net regression analysis. The penalization parameter was chosen to minimize the cross validated mean square error rate (Supplement 1, Fig. S14), which resulted in 513 CpGs.

Estimation of blood cell counts based on DNAm levels

We estimate blood cell counts using two different software tools. First, Houseman's estimation method [62] was used to estimate the proportions of CD8+ T cells, CD4+ T, natural killer, B cells, and granulocytes (mainly neutrophils). Second, the Horvath method, implemented in the advanced analysis option of the epigenetic clock software [11, 18], was used to estimate the percentage of exhausted CD8+ T cells (defined as CD28-CD45RA-), the number (count) of naïve CD8+ T cells (defined as CD45RA+CCR7+) and plasmablasts. We and others have shown that the estimated blood cell counts have moderately high correlations with corresponding flow cytometric measures [62, 63].

Additional descriptions of methods and materials can be found in Supplement 1.

Ethics approval

This study was reviewed by the UCLA institutional review board (IRB#13-000671, IRB#15-000697, IRB#16-001841, IRB#15-000682).

Availability of data and materials

The WHI data are available at dbGaP under the accession numbers phs000200.v10.p3. The FHS data are available at dbGaP under the accession numbers phs000342 and phs000724. The Normative Aging data are available from dbGAP phs000853.v1.p1. The Jackson Heart Study data are available from <https://www.jacksonheartstudy.org/Research/Study-Data>.

AUTHOR CONTRIBUTIONS

ML and SH developed the DNAmPhenoAge estimator and wrote the article. SH, ML, LF conceived of the study. ML, SH, AL, AQ carried out the statistical analysis. The remaining authors contributed data and participated in the interpretation of the results.

ACKNOWLEDGEMENTS

We would like to acknowledge The WHI Investigators listed below:

Program Office: (National Heart, Lung, and Blood Institute, Bethesda, Maryland) Jacques Rossouw, Shari Ludlam, Dale Burwen, Joan McGowan, Leslie Ford, and Nancy Geller.

Clinical Coordinating Center: (Fred Hutchinson Cancer Research Center, Seattle, WA) Garnet Anderson, Ross Prentice, Andrea LaCroix, and Charles Kooperberg.

Investigators and Academic Centers: (Brigham and Women's Hospital, Harvard Medical School, Boston, MA) JoAnn E. Manson; (MedStar Health Research Institute/Howard University, Washington, DC) Barbara V. Howard; (Stanford Prevention Research Center, Stanford, CA).

Marcia L. Stefanick; (The Ohio State University, Columbus, OH) Rebecca Jackson; (University of Arizona, Tucson/Phoenix, AZ) Cynthia A. Thomson; (University at Buffalo, Buffalo, NY).

Jean Wactawski-Wende; (University of Florida, Gainesville/Jacksonville, FL) Marian Limacher; (University of Iowa, Iowa City/Davenport, IA) Robert Wallace; (University of Pittsburgh, Pittsburgh, PA) Lewis Kuller; (Wake Forest University School of Medicine, Winston-Salem, NC) Sally Shumaker.

The Framingham Heart Study is funded by National Institutes of Health contract N01-HC-25195 and HHSN268201500001I. The laboratory work for this investigation was funded by the Division of Intramural Research, National Heart, Lung, and Blood Institute, National Institutes of Health. The analytical component of this project was funded by the Division of Intramural Research, National Heart, Lung, and Blood Institute, and the Center for Information Technology, National Institutes of Health, Bethesda, MD. The Framingham Heart Study is conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with Boston University. This manuscript was not prepared in collaboration with investigators of the Framingham Heart Study and does not necessarily reflect the opinions or views of the Framingham Heart Study, Boston University, or the NHLBI.

The United States Department of Veterans Affairs (VA) Normative Aging Study (NAS) is supported by the Cooperative Studies Program/ERIC and is a research component of the Massachusetts Veterans Epidemiology Research and Information Center (MAVERIC), Boston Massachusetts.

The MESA Epigenomics and Transcriptomics Studies were funded by R01HL101250, R01 DK103531-01, R01 DK103531, R01 AG054474, and R01 HL135009-01 to Wake Forest University Health Sciences.

We thank the Jackson Heart Study (JHS) participants and staff for their contributions to this work. The JHS is supported by contracts HHSN268201300046C,

HHSN268201300047C, HHSN268201300048C, HHSN268201300049C, HHSN268201300050C from the National Heart, Lung, and Blood Institute and the National Institute on Minority Health and Health Disparities. Dr. Wilson is supported by U54GM115428 from the National Institute of General Medical Sciences.

CONFLICTS OF INTEREST

The Regents of the University of California is the sole owner of a provisional patent application directed at this invention for which MEL, SH are named inventors.

FUNDING

This study was supported by NIH/NIA U34AG051425-01 (Horvath) and NIH/NIA K99AG052604 (Levine). The WHI epigenetic studies were supported by NIH/NHLBI 60442456 BAA23 (Assimes, Absher, Horvath) and by the National Institute of Environmental Health Sciences R01-ES020836 WHI-EMPC (Whitsel, Baccarelli, Hou). The WHI program is funded by the National Heart, Lung, and Blood Institute, National Institutes of Health, U.S. Department of Health and Human Services through contracts HHSN268201100046C, HHSN268201100001C, HHSN268201100002C, HHSN268201100003C, HHSN268201100004C, and HHSN271201100004C. The authors thank the WHI investigators and staff for their dedication, and the study participants for making the program possible. A full listing of WHI investigators can be found at: www.whi.org/researchers/Documents%20%20Write%20a%20Paper/WHI%20Investigator%20Short%20List.pdf.

The InCHIANTI study baseline (1998-2000) was supported as a "targeted project" (ICS110.1/RF97.71) by the Italian Ministry of Health and in part by the U.S. National Institute on Aging (Contracts: 263 MD 9164 and 263 MD 821336).

Funding for the DNA methylation in JHS provided by NHLBI R01HL116446.

The funding bodies played no role in the design, the collection, analysis, or interpretation of the data.

REFERENCES

1. Kennedy BK, Berger SL, Brunet A, Campisi J, Cuervo AM, Epel ES, Franceschi C, Lithgow GJ, Morimoto RI, Pessin JE, Rando TA, Richardson A, Schadt EE, et al. *Geroscience: linking aging to chronic disease*. Cell. 2014; 159:709–13.
<https://doi.org/10.1016/j.cell.2014.10.039>
2. Burch JB, Augustine AD, Frieden LA, Hadley E, Howcroft TK, Johnson R, Khalsa PS, Kohanski RA, Li XL, Macchiarini F, Niederehe G, Oh YS, Pawlyk AC, et al. *Advances in geroscience: impact on healthspan and chronic disease*. J Gerontol A Biol Sci Med Sci. 2014 (Suppl 1); 69:S1–3.
<https://doi.org/10.1093/gerona/glu041>
3. Fraga MF, Esteller M. *Epigenetics and aging: the targets and the marks*. Trends Genet. 2007; 23:413–18. <https://doi.org/10.1016/j.tig.2007.05.008>
4. Rakyan VK, Down TA, Maslau S, Andrew T, Yang TP, Beyan H, Whittaker P, McCann OT, Finer S, Valdes AM, Leslie RD, Deloukas P, Spector TD. *Human aging-associated DNA hypermethylation occurs preferentially at bivalent chromatin domains*. Genome Res. 2010; 20:434–39.
<https://doi.org/10.1101/gr.103101.109>
5. Teschendorff AE, Menon U, Gentry-Maharaj A, Ramus SJ, Weisenberger DJ, Shen H, Campan M, Noushmehr H, Bell CG, Maxwell AP, Savage DA, Mueller-Holzner E, Marth C, et al. *Age-dependent DNA methylation of genes that are suppressed in stem cells is a hallmark of cancer*. Genome Res. 2010; 20:440–46.
<https://doi.org/10.1101/gr.103606.109>
6. Jung M, Pfeifer GP. *Aging and DNA methylation*. BMC Biol. 2015; 13:7. <https://doi.org/10.1186/s12915-015-0118-4>
7. Zheng SC, Widswendter M, Teschendorff AE. *Epigenetic drift, epigenetic clocks and cancer risk*. Epigenomics. 2016; 8:705–19.
<https://doi.org/10.2217/epi-2015-0017>
8. Bocklandt S, Lin W, Sehl ME, Sánchez FJ, Sinsheimer JS, Horvath S, Vilain E. *Epigenetic predictor of age*. PLoS One. 2011; 6:e14821.
<https://doi.org/10.1371/journal.pone.0014821>
9. Garagnani P, Bacalini MG, Pirazzini C, Gori D, Giuliani C, Mari D, Di Blasio AM, Gentilini D, Vitale G, Collino S, Rezzi S, Castellani G, Capri M, et al. *Methylation of ELOVL2 gene as a new epigenetic marker of age*. Aging Cell. 2012; 11:1132–34.
<https://doi.org/10.1111/ace.12005>
10. Hannum G, Guinney J, Zhao L, Zhang L, Hughes G, Sadda S, Klotzle B, Bibikova M, Fan JB, Gao Y, Deconde R, Chen M, Rajapakse I, et al. *Genome-wide methylation profiles reveal quantitative views of human aging rates*. Mol Cell. 2013; 49:359–67.
<https://doi.org/10.1016/j.molcel.2012.10.016>
11. Horvath S. *DNA methylation age of human tissues and cell types*. Genome Biol. 2013; 14:R115.
<https://doi.org/10.1186/gb-2013-14-10-r115>
12. Weidner CI, Lin Q, Koch CM, Eisele L, Beier F, Ziegler P, Bauerschlag DO, Jöckel KH, Erbel R, Mühlleisen TW,

- Zenke M, Brümmendorf TH, Wagner W. Aging of blood can be tracked by DNA methylation changes at just three CpG sites. *Genome Biol.* 2014; 15:R24. <https://doi.org/10.1186/gb-2014-15-2-r24>
13. Lin Q, Weidner CI, Costa IG, Marioni RE, Ferreira MR, Deary IJ, Wagner W. DNA methylation levels at individual age-associated CpG sites can be indicative for life expectancy. *Aging (Albany NY)*. 2016; 8:394–401. <https://doi.org/10.1863/aging.100908>
14. Horvath S, Erhart W, Brosch M, Ammerpohl O, von Schönfels W, Ahrens M, Heits N, Bell JT, Tsai PC, Spector TD, Deloukas P, Siebert R, Sipos B, et al. Obesity accelerates epigenetic aging of human liver. *Proc Natl Acad Sci USA*. 2014; 111:15538–43. <https://doi.org/10.1073/pnas.1412759111>
15. Marioni RE, Shah S, McRae AF, Chen BH, Colicino E, Harris SE, Gibson J, Henders AK, Redmond P, Cox SR, Pattie A, Corley J, Murphy L, et al. DNA methylation age of blood predicts all-cause mortality in later life. *Genome Biol.* 2015; 16:25. <https://doi.org/10.1186/s13059-015-0584-6>
16. Marioni RE, Shah S, McRae AF, Ritchie SJ, Muniz-Terrera G, Harris SE, Gibson J, Redmond P, Cox SR, Pattie A, Corley J, Taylor A, Murphy L, et al. The epigenetic clock is correlated with physical and cognitive fitness in the Lothian Birth Cohort 1936. *Int J Epidemiol.* 2015; 44:1388–96. <https://doi.org/10.1093/ije/dyu277>
17. Horvath S, Garagnani P, Bacalini MG, Pirazzini C, Salvioli S, Gentilini D, Di Blasio AM, Giuliani C, Tung S, Vinters HV, Franceschi C. Accelerated epigenetic aging in Down syndrome. *Aging Cell*. 2015; 14:491–95. <https://doi.org/10.1111/acel.12325>
18. Horvath S, Levine AJ. HIV-1 Infection Accelerates Age According to the Epigenetic Clock. *J Infect Dis*. 2015; 212:1563–73. <https://doi.org/10.1093/infdis/jiv277>
19. Horvath S, Pirazzini C, Bacalini MG, Gentilini D, Di Blasio AM, Delledonne M, Mari D, Arosio B, Monti D, Passarino G, De Rango F, D'Aquila P, Giuliani C, et al. Decreased epigenetic age of PBMCs from Italian semi-supercentenarians and their offspring. *Aging (Albany NY)*. 2015; 7:1159–70. <https://doi.org/10.1863/aging.100861>
20. Levine ME, Hosgood HD, Chen B, Absher D, Assimes T, Horvath S. DNA methylation age of blood predicts future onset of lung cancer in the women's health initiative. *Aging (Albany NY)*. 2015; 7:690–700. <https://doi.org/10.1863/aging.100809>
21. Levine ME, Lu AT, Bennett DA, Horvath S. Epigenetic age of the pre-frontal cortex is associated with neuritic plaques, amyloid load, and Alzheimer's disease related cognitive functioning. *Aging (Albany NY)*. 2015; 7:1198–211. <https://doi.org/10.1863/aging.100864>
22. Levine ME, Lu AT, Chen BH, Hernandez DG, Singleton AB, Ferrucci L, Bandinelli S, Salfati E, Manson JE, Quach A, Kusters CD, Kuh D, Wong A, et al. Menopause accelerates biological aging. *Proc Natl Acad Sci USA*. 2016; 113:9327–32. <https://doi.org/10.1073/pnas.1604558113>
23. Chen BH, Marioni RE, Colicino E, Peters MJ, Ward-Caviness CK, Tsai PC, Roetker NS, Just AC, Demerath EW, Guan W, Bressler J, Fornage M, Studenski S, et al. DNA methylation-based measures of biological age: meta-analysis predicting time to death. *Aging (Albany NY)*. 2016; 8:1844–65. <https://doi.org/10.1863/aging.101020>
24. Quach A, Levine ME, Tanaka T, Lu AT, Chen BH, Ferrucci L, Ritz B, Bandinelli S, Neuhouser ML, Beasley JM, Snetselaar L, Wallace RB, Tsao PS, et al. Epigenetic clock analysis of diet, exercise, education, and lifestyle factors. *Aging (Albany NY)*. 2017; 9:419–46. <https://doi.org/10.1863/aging.101168>
25. Dugue PA, Bassett JK, Joo JE, Jung CH, Ming Wong E, Moreno-Betancur M, Schmidt D, Makalic E, Li S, Severi G, Hodge AM, Buchanan DD, English DR, et al. DNA methylation-based biological aging and cancer risk and survival: pooled analysis of seven prospective studies. *Int J Cancer*. 2018; 142:1611–19. <https://doi.org/10.1002/ijc.31189>
26. Simpkin AJ, Howe LD, Tilling K, Gaunt TR, Lyttleton O, McArdle WL, Ring SM, Horvath S, Smith GD, Relton CL. The epigenetic clock and physical development during childhood and adolescence: longitudinal analysis from a UK birth cohort. *Int J Epidemiol*. 2017; 46:549–58. <https://doi.org/10.1093/ije/dyw307>
27. Maierhofer A, Flunkert J, Oshima J, Martin GM, Haaf T, Horvath S. Accelerated epigenetic aging in Werner syndrome. *Aging (Albany NY)*. 2017; 9:1143–52. <https://doi.org/10.1863/aging.101217>
28. Levine ME. Modeling the rate of senescence: can estimated biological age predict mortality more accurately than chronological age? *J Gerontol A Biol Sci Med Sci*. 2013; 68:667–74. <https://doi.org/10.1093/gerona/gls233>
29. Belsky DW, Caspi A, Houts R, Cohen HJ, Corcoran DL, Danese A, Harrington H, Israel S, Levine ME, Schaefer JD, Sugden K, Williams B, Yashin AI, et al. Quantification of biological aging in young adults. *Proc Natl Acad Sci USA*. 2015; 112:E4104–10. <https://doi.org/10.1073/pnas.1506264112>
30. Li S, Wong EM, Joo JE, Jung CH, Chung J, Apicella C, Stone J, Dite GS, Giles GG, Southey MC, Hopper JL. Genetic and Environmental Causes of Variation in the Difference Between Biological Age Based on DNA

- Methylation and Chronological Age for Middle-Aged Women. *Twin Res Hum Genet.* 2015; 18:720–26. <https://doi.org/10.1017/thg.2015.75>
31. Sebastiani P, Thyagarajan B, Sun F, Schupf N, Newman AB, Montano M, Perls TT. Biomarker signatures of aging. *Aging Cell.* 2017; 16:329–38. <https://doi.org/10.1111/acel.12557>
32. Ferrucci L, Hesdorffer C, Bandinelli S, Simonsick EM. Frailty as a Nexus Between the Biology of Aging, Environmental Conditions and Clinical Geriatrics. *Public Health Rev.* 2010; 32:475–88. <https://doi.org/10.1007/BF03391612>
33. Christiansen L, Lenart A, Tan Q, Vaupel JW, Aviv A, McGue M, Christensen K. DNA methylation age is associated with mortality in a longitudinal Danish twin study. *Aging Cell.* 2016; 15:149–54. <https://doi.org/10.1111/acel.12421>
34. Perna L, Zhang Y, Mons U, Holleczek B, Saum KU, Brenner H. Epigenetic age acceleration predicts cancer, cardiovascular, and all-cause mortality in a German case cohort. *Clin Epigenetics.* 2016; 8:64. <https://doi.org/10.1186/s13148-016-0228-z>
35. Breitling LP, Yang R, Korn B, Burwinkel B, Brenner H. Tobacco-smoking-related differential DNA methylation: 27K discovery and replication. *Am J Hum Genet.* 2011; 88:450–57. <https://doi.org/10.1016/j.ajhg.2011.03.003>
36. Joehanes R, Just AC, Marioni RE, Pilling LC, Reynolds LM, Mandaviya PR, Guan W, Xu T, Elks CE, Aslibekyan S, Moreno-Macias H, Smith JA, Brody JA, et al. Epigenetic Signatures of Cigarette Smoking. *Circ Cardiovasc Genet.* 2016; 9:436–47. <https://doi.org/10.1161/CIRGENETICS.116.001506>
37. Lee KW, Pausova Z. Cigarette smoking and DNA methylation. *Front Genet.* 2013; 4:132. <https://doi.org/10.3389/fgene.2013.00132>
38. Zhang Y, Wilson R, Heiss J, Breitling LP, Saum KU, Schöttker B, Holleczek B, Waldenberger M, Peters A, Brenner H. DNA methylation signatures in peripheral blood strongly predict all-cause mortality. *Nat Commun.* 2017; 8:14617. <https://doi.org/10.1038/ncomms14617>
39. Bennett DA, Schneider JA, Arvanitakis Z, Wilson RS. Overview and findings from the religious orders study. *Curr Alzheimer Res.* 2012; 9:628–45. <https://doi.org/10.2174/156720512801322573>
40. Bennett DA, Schneider JA, Buchman AS, Barnes LL, Boyle PA, Wilson RS. Overview and findings from the Rush Memory and Aging Project. *Curr Alzheimer Res.* 2012; 9:646–63. <https://doi.org/10.2174/156720512801322663>
41. Franceschi C, Bonafè M, Valensin S, Olivieri F, De Luca M, Ottaviani E, De Benedictis G. Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann NY Acad Sci.* 2000; 908:244–54. <https://doi.org/10.1111/j.1749-6632.2000.tb06651.x>
42. Horvath S, Zhang Y, Langfelder P, Kahn RS, Boks MP, van Eijk K, van den Berg LH, Ophoff RA. Aging effects on DNA methylation modules in human brain and blood tissue. *Genome Biol.* 2012; 13:R97. <https://doi.org/10.1186/gb-2012-13-10-r97>
43. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet.* 2011; 88:76–82. <https://doi.org/10.1016/j.ajhg.2010.11.011>
44. Sierra F, Kohanski R. Geroscience and the trans-NIH Geroscience Interest Group, GSIG. *Geroscience.* 2017; 39:1–5. <https://doi.org/10.1007/s11357-016-9954-6>
45. Belsky DW, Moffitt TE, Cohen AA, Corcoran DL, Levine ME, Prinz JA, Schaefer J, Sugden K, Williams B, Poulton R, Caspi A. Eleven Telomere, Epigenetic Clock, and Biomarker-Composite Quantifications of Biological Aging: Do They Measure the Same Thing? *Am J Epidemiol.* 2017. <https://doi.org/10.1093/aje/kwx346>
46. Kubben N, Misteli T. Shared molecular and cellular mechanisms of premature ageing and ageing-associated diseases. *Nat Rev Mol Cell Biol.* 2017; 18:595–609. <https://doi.org/10.1038/nrm.2017.68>
47. Merlo A, Herman JG, Mao L, Lee DJ, Gabrielson E, Burger PC, Baylin SB, Sidransky D. 5' CpG island methylation is associated with transcriptional silencing of the tumour suppressor p16/CDKN2/MTS1 in human cancers. *Nat Med.* 1995; 1:686–92. <https://doi.org/10.1038/nm0795-686>
48. Finch CE. CHAPTER 1 - Inflammation and Oxidation in Aging and Chronic Diseases. *The Biology of Human Longevity.* (Burlington: Academic Press), 2007. pp. 1–112.
49. Finch CE. Evolution in health and medicine Sackler colloquium: Evolution of the human lifespan and diseases of aging: roles of infection, inflammation, and nutrition. *Proc Natl Acad Sci USA.* 2010 (Suppl 1); 107:1718–24. <https://doi.org/10.1073/pnas.0909606106>
50. Moiseeva O, Mallette FA, Mukhopadhyay UK, Moores A, Ferbeyre G. DNA damage signaling and p53-dependent senescence after prolonged beta-interferon stimulation. *Mol Biol Cell.* 2006; 17:1583–92. <https://doi.org/10.1091/mbc.E05-09-0858>
51. Gonskikh Y, Polacek N. Alterations of the translation apparatus during aging and stress response. *Mech*

- Ageing Dev. 2017 (Suppl C); 168:30–36.
<https://doi.org/10.1016/j.mad.2017.04.003>
52. Wolters S, Schumacher B. Genome maintenance and transcription integrity in aging and disease. *Front Genet.* 2013; 4:19.
<https://doi.org/10.3389/fgene.2013.00019>
53. Steffen KK, Dillin A. A Ribosomal Perspective on Proteostasis and Aging. *Cell Metab.* 2016; 23:1004–12. <https://doi.org/10.1016/j.cmet.2016.05.013>
54. Garinis GA, Uittenboogaard LM, Stachelscheid H, Fousteri M, van Ijcken W, Breit TM, van Steeg H, Mullenders LH, van der Horst GT, Brüning JC, Niessen CM, Hoeijmakers JH, Schumacher B. Persistent transcription-blocking DNA lesions trigger somatic growth attenuation associated with longevity. *Nat Cell Biol.* 2009; 11:604–15.
<https://doi.org/10.1038/ncb1866>
55. Schumacher B, Garinis GA, Hoeijmakers JH. Age to survive: DNA damage and aging. *Trends Genet.* 2008; 24:77–85. <https://doi.org/10.1016/j.tig.2007.11.004>
56. McMurray CT, Vijg J. Editorial overview: Molecular and genetic bases of disease: the double life of DNA. *Curr Opin Genet Dev.* 2014; 26:v–vii.
<https://doi.org/10.1016/j.gde.2014.09.002>
57. Levine ME, Crimmins EM. A Genetic Network Associated With Stress Resistance, Longevity, and Cancer in Humans. *J Gerontol A Biol Sci Med Sci.* 2016; 71:703–12.
<https://doi.org/10.1093/gerona/glv141>
58. Zhou KI, Pincus Z, Slack FJ. Longevity and stress in *Caenorhabditis elegans*. *Aging (Albany NY)*. 2011; 3:733–53. <https://doi.org/10.18632/aging.100367>
59. Lin YJ, Seroude L, Benzer S. Extended life-span and stress resistance in the *Drosophila* mutant methuselah. *Science.* 1998; 282:943–46.
<https://doi.org/10.1126/science.282.5390.943>
60. Yu BP, Chung HY. Stress resistance by caloric restriction for longevity. *Ann N Y Acad Sci.* 2001; 928:39–47. <https://doi.org/10.1111/j.1749-6632.2001.tb05633.x>
61. Martin GM, Austad SN, Johnson TE. Genetic analysis of ageing: role of oxidative damage and environmental stresses. *Nat Genet.* 1996; 13:25–34.
<https://doi.org/10.1038/ng0596-25>
62. Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, Wiencke JK, Kelsey KT. DNA methylation arrays as surrogate measures of cell mixture distribution. *BMC Bioinformatics.* 2012; 13:86.
<https://doi.org/10.1186/1471-2105-13-86>
63. Horvath S, Gurven M, Levine ME, Trumble BC, Kaplan H, Allayee H, Ritz BR, Chen B, Lu AT, Rickabaugh TM, Jamieson BD, Sun D, Li S, et al. An epigenetic clock analysis of race/ethnicity, sex, and coronary heart disease. *Genome Biol.* 2016; 17:171.
<https://doi.org/10.1186/s13059-016-1030-0>

SUPPLEMENTARY MATERIAL

Please browse the Full Text version of this manuscript
to see Supplementary Methods, Tables, and Figures
presented in Supplements 1-4.