

 Open access • Journal Article • DOI:10.1086/301542

An Extragalactic H I Cloud with No Optical Counterpart — [Source link](#)

[Virginia A. Kilborn](#), [Lister Staveley-Smith](#), [M. Marquarding](#), [Rachel L. Webster](#) ...+28 more authors

Institutions: [University of Melbourne](#), [Australia Telescope National Facility](#), [Cardiff University](#), [University of Western Sydney](#) ...+5 more institutions

Published on: 01 Sep 2000 - [The Astronomical Journal](#) (IOP Publishing)

Topics: [HIPASS](#), [Star formation](#) and [Galaxy](#)

Related papers:

- [An Extragalactic HI Cloud with No Optical Counterpart](#)
- [HIPASS detection of an intergalactic gas cloud in the NGC 2442 group](#)
- [The HI Parkes All Sky Survey: southern observations, calibration and robust imaging](#)
- [Where are the missing galactic satellites](#)
- [Discovery of a large intergalactic H I cloud in the M96 group](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/an-extragalactic-h-i-cloud-with-no-optical-counterpart-4yw5wyc2j>

AN EXTRAGALACTIC H I CLOUD WITH NO OPTICAL COUNTERPART?

V. A. KILBORN,¹ L. STAVELEY-SMITH,² M. MARQUARDING,³ R. L. WEBSTER,³ D. F. MALIN,⁴ G. D. BANKS,⁵ R. BHATHAL,⁶ W. J. G. DE BLOK,² P. J. BOYCE,⁵ M. J. DISNEY,⁵ M. J. DRINKWATER,³ R. D. EKERS,² K. C. FREEMAN,⁷ B. K. GIBSON,⁸ P. A. HENNING,^{2,9} H. JERJEN,⁷ P. M. KNEZEK,^{10,11} B. KORIBALSKI,² R. F. MINCHIN,^{5,11} J. R. MOULD,⁷ T. OOSTERLOO,¹² R. M. PRICE,^{2,9} M. E. PUTMAN,⁷ S. D. RYDER,¹³ E. M. SADLER,¹⁴ I. STEWART,² F. STOOTMAN,⁶ AND A. E. WRIGHT²

Received 1999 December 27; accepted 2000 May 11

ABSTRACT

We report the discovery, from the H I Parkes All-Sky Survey (HIPASS), of an isolated cloud of neutral hydrogen, which we believe to be extragalactic. The H I mass of the cloud (HIPASS J1712–64) is very low, $1.7 \times 10^7 M_{\odot}$, using an estimated distance of ~ 3.2 Mpc. Most significantly, we have found no optical companion to this object to very faint limits [$\mu(B) \sim 27$ mag arcsec⁻²]. HIPASS J1712–64 appears to be a binary system similar to, but much less massive than, H I 1225+01 (the Virgo H I cloud) and has a size of at least 15 kpc. The mean velocity dispersion measured with the Australia Telescope Compact Array (ATCA) is only 4 km s⁻¹ for the main component and, because of the weak or non-existent star formation, possibly reflects the thermal line width ($T < 2000$ K) rather than bulk motion or turbulence. The peak column density for HIPASS J1712–64, from the combined Parkes and ATCA data, is only 3.5×10^{19} cm⁻², which is estimated to be a factor of 2 below the critical threshold for star formation. Apart from its significantly higher velocity, the properties of HIPASS J1712–64 are similar to the recently recognized class of compact high-velocity clouds. We therefore consider the evidence for a Local Group or Galactic origin, although a more plausible alternative is that HIPASS J1712–64 was ejected from the interacting Magellanic Cloud–Galaxy system at perigalacticon $\sim 2 \times 10^8$ yr ago.

Key words: galaxies: formation — galaxies: irregular — radio emission lines

1. INTRODUCTION

Until recently, there has been no extensive survey of the extragalactic sky in neutral hydrogen (H I). The mass distribution in the universe has, therefore, been traced by galaxy surveys in other wavebands, particularly in the optical, but also in the infrared using *IRAS*. Because of the limited nature of existing H I surveys, it is unclear whether there is a significant local population of H I-rich objects not seen in optical and infrared surveys (away from the Galactic

plane). In particular, the existence of H I clouds without optical counterparts, or “H I protogalaxies,” might significantly add to the local H I mass density of the universe. Recent observations by Zwaan et al. (1997) and Schneider, Spitzak, & Rosenberg (1998) arrive at somewhat differing conclusions regarding the low H I mass galaxy population, leading to some uncertainty in the contribution of these objects to the H I mass in the local universe.

Limits to the presence of a population of H I-rich clouds and “protogalaxies” have been set by a number of blind or semiblind H I surveys conducted in a range of environments, such as clusters (Barnes et al. 1997; Dickey 1997), groups (Kraan-Korteweg et al. 1999; Banks et al. 1999; Fisher & Tully 1981a), voids (Krumm & Brosch 1984), and the general field (Henning 1995; Sorar 1994; Spitzak & Schneider 1998; Shostak 1977). In addition, there have been numerous observations in off-source calibration regions for galaxies that have been optically selected. For example, Fisher & Tully (1981b) serendipitously detected a couple of dozen galaxies in the calibration scans from their survey, all with optical counterparts.

A number of H I clouds without optical counterparts have been found. However, many appear to be closely associated with massive galaxies (Williams & van Gorkom 1988; Giovanelli et al. 1995) and perhaps the product of tidal interactions (Hibbard & van Gorkom 1996). A number of potential H I protogalaxies have now been classified as high-velocity clouds (Mathewson, Cleary, & Murray 1975), and argument continues over the extragalactic nature of a specific class of compact high-velocity clouds (Braun & Burton 1999; Zwaan & Briggs 2000). One of the larger blind H I surveys to date (Spitzak & Schneider 1998) detected 75 galaxies in 55 deg² of sky. One detection was found not to have any obvious optical counterpart, though a bright star just near the peak of the H I emission may have obscured a low surface brightness optical companion.

¹ School of Physics, University of Melbourne, Parkville, Victoria 3052, Australia; vkilborn@physics.unimelb.edu.au.

² Australia Telescope National Facility, CSIRO, P.O. Box 76, Epping, NSW 2121, Australia.

³ School of Physics, University of Melbourne, Parkville, Victoria 6052, Australia.

⁴ Anglo-Australian Observatory, P.O. Box 296, Epping, NSW 2121, Australia.

⁵ Department of Physics and Astronomy, University of Wales, Cardiff, P.O. Box 913, Cardiff CF2 3YB, UK.

⁶ Department of Physics, University of Western Sydney Macarthur, P.O. Box 555, Campbelltown, NSW 2560, Australia.

⁷ Research School of Astronomy and Astrophysics, ANU, Weston Creek P.O., Weston, ACT 2611, Australia.

⁸ Center for Astrophysics and Space Astronomy, University of Colorado, Boulder, CO 80309-0389.

⁹ Department of Physics and Astronomy, University of New Mexico, 800 Yale Boulevard Northeast, Albuquerque, NM 87131.

¹⁰ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD, 21218.

¹¹ Visiting Astronomer, Cerro Tololo Inter-American Observatory, National Optical Astronomy Observatories, which is operated by the Association of Universities for Research in Astronomy (AURA), Inc., under cooperative agreement with the National Science Foundation (NSF).

¹² Istituto di Fisica Cosmica, via Bassini 15, I-20133, Milano, Italy.

¹³ Joint Astronomy Center, 660 North Aohoku Place, Hilo, HI 96720.

¹⁴ Astrophysics Department, School of Physics, University of Sydney, A28, Sydney, NSW 2006, Australia.

The closest and best examples of H I clouds without prominent counterparts are the Virgo Cloud H I 1225+01 (Giovanelli & Haynes 1989), and the Leo ring (Schneider et al. 1983). Both of these H I clouds were later found to be associated with optical galaxies. H I 1225+01 was found serendipitously during routine calibration observations. It comprises two regions of H I emission. The largest has a Magellanic-type dwarf irregular galaxy at the position of the highest H I column density, while the smaller, possibly an infalling cloud, shows no optical emission to a faint limiting magnitude (Chengular, Giovanelli, & Haynes 1995). The larger component has a total dynamical mass of $\sim 1.0 \times 10^{10} d_{20} M_{\odot}$, and a gas mass of $\sim 3 \times 10^9 d_{20}^2 M_{\odot}$, where d_{20} is the distance in units of 20 Mpc. The heliocentric velocity of the cloud is 1275 km s^{-1} and the extent of the H I emission is about $\sim 200 d_{20}$ kpc. The Leo ring is an intergalactic cloud found during H I observations of the Leo group of galaxies. It comprises a large ring of H I emission surrounding NGC 3384 and M105, with an H I mass of $\sim 10^9 M_{\odot}$. The velocity structure suggests that it is probably not a tidal tail from the nearby galaxies, but a primordial gas cloud that has not started forming stars (Schneider 1989). The cloud is gravitationally bound to the optical galaxies in the group. Both of these objects have been used in H α studies to determine limits on the local ionizing background (Donahue, Aldering, & Stocke 1995), finding limits which suggest that quasar light, not galactic light, dominates the local ionizing background at low redshift.

The H I Parkes¹⁵ All-Sky Survey (HIPASS) has been underway since 1997 and, when complete, will be the largest blind H I survey to date, surveying a volume at least 2 orders of magnitude larger than any previous survey. In early HIPASS observations of 600 deg^2 in the Centaurus region, Banks et al. (1999) found 10 new members of the nearby Centaurus A group. However, all of these have optical companions, although five are faint and were previously uncataloged. Other early observations include the south celestial cap (Putman et al. 1998; Kilborn, Webster, & Staveley-Smith 1999). During routine inspection of this data, a resolved H I detection, HIPASS J1712–64, was selected for follow-up observations as it showed no cataloged galaxy in the vicinity and no optical counterpart on the Digitized Sky Survey. It also had a low systemic velocity (though well-separated from Galactic gas and high-velocity clouds) and a significant angular size.

The observations of HIPASS J1712–64 are discussed in § 2, while the H I structure, dynamics, and optical limits are discussed in § 3. Alternative high-velocity cloud and Local Group hypotheses are discussed in § 4, the nature of HIPASS J1712–64 is discussed in § 5, and the results are summarized in § 6. We assume a Hubble constant of $75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ throughout.

2. OBSERVATIONS AND DATA REDUCTION

2.1. HIPASS

HIPASS is a blind H I survey of the sky $\delta < 2^{\circ}$. The survey uses the Parkes 64 m telescope, which is equipped with a 13 beam receiver (Staveley-Smith et al. 1996). The observations began in 1997 February and were completed

in 2000 May. A northern extension is now underway at Parkes, and parts of a complementary northern survey are currently being conducted with Jodrell Bank Observatory's Lovell Telescope. The velocity range covered is -1200 to $12,700 \text{ km s}^{-1}$. Observations are taken by scanning the multibeam receiver in declination strips of length $\sim 8^{\circ}$. Each declination scan is separated by $35'$ in R.A., and each area of sky is scanned five times, resulting in a final scan separation of $7'$. The final integration time is approximately 460 s beam^{-1} . The mean telescope FWHM beam is 14.3 , although the gridding process increases this to ~ 15.5 (Barnes et al. 2000).

Bandpass calibration, spectral smoothing, and Doppler correction of the data are applied in real time at the telescope (Barnes et al. 1998). Once all data are collected, the spectra are gridded into data cubes, which have an rms noise level of $\sim 13 \text{ mJy beam}^{-1}$. The channel spacing in the final cubes is 13.2 km s^{-1} , the FWHM resolution is 18.0 km s^{-1} , and the pixel size is $4' \times 4'$.

2.2. H I Observations

HIPASS J1712–64 was discovered during a routine visual inspection of data cubes. The detection is very significant, with a peak flux of $\sim 150 \text{ mJy beam}^{-1}$, corresponding to a signal-to-noise ratio (S/N) of about 12. It is also significantly resolved in the beam of the Parkes telescope, extending over at least two beamwidths. On 1999 June 2, a further 2.5 hr observation was made at the Parkes telescope, using the multibeam receiver, to confirm the detection. The observation was of a $3^{\circ} \times 4^{\circ}$ field centered on the original detection. This observation also resulted in a clear positive detection. The rms noise in the follow-up observation was 19 mJy beam^{-1} . The combined and spatially integrated Parkes spectrum is shown in Figure 1. The total flux density peaks at 250 mJy at a heliocentric velocity of $\sim 455 \text{ km s}^{-1}$.

Higher resolution observations in the H I line were made at the Australia Telescope Compact Array (ATCA) on 1999 June 30. Approximately 12 hr of data were obtained using the 375 m array, and the pointing center for the observations was R.A. $17^{\text{h}}12^{\text{m}}13^{\text{s}}$, decl. $-64^{\circ}39'12''$ (J2000.0). The




FIG. 1.—Spatially integrated H I spectra of HIPASS J1712–64 from the Parkes-HIPASS data (solid line) and the ATCA (dashed line). The ATCA appears to recover $\sim 80\%$ of the total flux density. Both spectra are derived by spatially integrating over a box centered on the midpoint of J1712–64. The size of the box was $44'$ and $16'$ for the Parkes and the ATCA data, respectively. The ATCA data has been smoothed to the spectral resolution of the Parkes data. No clipping of the data points was used.

¹⁵ The Parkes telescope is part of the Australia Telescope, which is funded by the Commonwealth of Australia for operation as a national facility managed by CSIRO.

secondary calibrator PKS 1814-637 was observed every 30 minutes for 4 minutes, to calibrate the amplitude and phase of the visibility data. The primary flux scale was set using the primary calibrator PKS 1934-638. The bandwidth of 8 MHz was divided into 512 channels, resulting in a channel spacing of 3.3 km s^{-1} and a FWHM resolution, before any smoothing, of 4.0 km s^{-1} .

The data were reduced using the MIRIAD reduction package. The data were then edited, calibrated in amplitude and phase, and bandpass corrected. Several bright continuum sources were removed by fitting spectral baselines to the line-free channels in the visibility domain. Natural weighting was used to form the data cube, which was cleaned until the absolute maximum residual in each plane fell below 10 mJy. The final FWHM beam is 2.0×1.9 , and the rms noise is $3.7 \text{ mJy beam}^{-1}$. No correction for the primary beam pattern was applied.

J1712-64 was again detected in the ATCA data. The spatially integrated spectrum (smoothed to the same velocity resolution) was overlaid onto the Parkes-HIPASS data

in Figure 1. All except $\sim 20\%$ of the Parkes flux density (mainly at higher velocities) was recovered. This implies that some diffuse or low-level emission was being missed by the ATCA. The ATCA channel images (Hanning-smoothed to 6.6 km s^{-1} resolution) are shown in Figure 2.

A combined cube of the Parkes and ATCA data set was made by spectrally smoothing the ATCA data to match the HIPASS data, then spatially resampling the HIPASS data to match the ATCA data. The data were then combined using the MIRIAD task IMMERGE. It was assumed that both the ATCA and HIPASS flux density scales were accurate to a few percent. The final column density image, with contours overlaid, is shown in Figure 3. The ATCA velocity field, derived from the GIPSY task MOMENTS, is shown in Figure 4 overlaid onto the column density image.

2.3. Optical Observations

A search for an optical counterpart using blue and red film copies of the UK Schmidt Telescope (UKST)/ESO southern sky survey revealed no optical counterpart. Subse-




FIG. 2.—HI channel maps for HIPASS J1712-64 from the Hanning-smoothed ATCA data. The mean heliocentric velocity is given in the top left-hand corner of each map. The separation of plotted channels is 6.6 km s^{-1} . The contours plotted are $-5, 5, 10, 15, 20, 25,$ and 30 mJy beam^{-1} .




FIG. 3.—H I column density map of HIPASS J1712–64 from the combined ATCA-Parkes data set. The contours represent increments of 1×10^{19} atoms cm^{-2} to a maximum value of 3.5×10^{19} atoms cm^{-2} . The gray scale is a linear representation of the column density. The FWHM of the final beam shown in the lower right-hand corner is 2.0×1.9 .

quently, two deep original plates were obtained from the UKST plate archive and subjected to a photographic enhancement technique (Malin 1978) that is capable of revealing extended features over 5 mag fainter than the night sky (Malin & Hadley 1997). At the site of the UKST (Siding Spring) this is typically $\mu(B) = 22.5 \text{ mag arcsec}^{-2}$.

Only two suitable plates were available, and an image was made by combining photographically-enhanced derivatives from both of them, J1659 (IIIa-J/GG 395, 395–530

nm) and OR 15092 (IIIa-F/RG 590, 590–690 nm). Both were plates of excellent quality, but in both cases the object center was close to the vignetting region toward the edge of the UKST focal plane.

The IIIa-J passband is the most sensitive to extended stellar light (assuming a solar spectrum) and would reveal extended features at least as faint as $27.5 \text{ mag arcsec}^{-2}$. The derivative from the red-sensitive plate achieves a detection limit of $\sim 26.8 \text{ mag arcsec}^{-2}$ because of the brighter night

TABLE 1
MEASURED H I PARAMETERS FOR HIPASS J1712–64 AND ITS COMPONENTS

Parameters	NE Component	SW Component	J1712–64
ATCA and Parkes:
R.A. (J2000.0)	$17^{\text{h}}12^{\text{m}}35^{\text{s}}$	$17^{\text{h}}11^{\text{m}}35^{\text{s}}$...
Decl. (J2000.0)	$-64^{\circ}38'12''$	$-64^{\circ}45'11''$...
l	$326^{\circ}65$	$326^{\circ}48$...
b	$-14^{\circ}63$	$-14^{\circ}61$...
Size	11.4×8.2	4.9×4.1	15.9×8.2
P.A.	32°	125°	32°
Peak column density	$3.5 \times 10^{19} \text{ cm}^{-2}$	$2.8 \times 10^{19} \text{ cm}^{-2}$...
Total flux density	4.7 Jy km s^{-1}	1.3 Jy km s^{-1}	7.0 Jy km s^{-1}
ATCA:
Heliocentric velocity, v_{sys}	451 km s^{-1}	464 km s^{-1}	...
50% velocity width, W_{50}	11 km s^{-1}	20 km s^{-1}	...
Velocity dispersion, σ	4 km s^{-1}




FIG. 4.—H I velocity field of HIPASS J1712–64 from the ATCA data superposed on the column density image (from the combined ATCA and Parkes data). The contours are at intervals of 5 km s^{-1} and range from 450 km s^{-1} in the northeast to 475 km s^{-1} in the southwest. The FWHM of the final beam shown in the lower right-hand corner is 2.0×1.9 .

sky in the red. Nothing unusual is seen on deep derivatives from either plate or on the combined image. However, it should be mentioned that large, featureless, low surface brightness objects are particularly difficult to detect in crowded fields. For an optical companion of $1'$ in diameter and lying just beneath our optical limit, these limits suggest an absolute magnitude $M_B > -8.8$, and an H I mass-to-light ratio $M_{\text{H I}}/L_B > 24$ (Table 2).

The source was also imaged in the optical I band at the ANU 40 inch (1 m) telescope at Siding Springs Observatory on 1999 April 15. It was also imaged in the B and R bands by the Cerro Tololo Inter-American Observatory 40 inch (1 m) telescope on 1999 May 4. The data were reduced using the IRAF¹⁶ package, and calibration for the B - and R -band images used photometric observations of the Landolt Standard field 104 (Landolt 1983). The optical point source magnitude limits are $B \sim 19.8$ and $R \sim 19.2$ from the CCD observations. None of the images show any evidence of extended optical emission near the location of the H I detection. Figure 5 shows the H I contours of HIPASS J1712–64 overlaid onto the combined B - and R -band CCD images, showing a dense star field but no extended objects.

3. PHYSICAL PARAMETERS

3.1. H I Structure

HIPASS J1712–64 is well-resolved in each of the velocity channels with significant emission (see Fig. 2). The object appears to have two major components. The prominent one (hereafter the NE component) is centered at R.A. $17^{\text{h}}12^{\text{m}}35^{\text{s}}$, decl. $-64^{\circ}38'12''$ (J2000.0) and has a velocity spread from 438 to 464 km s^{-1} in the ATCA data. The systemic velocity is 451 km s^{-1} and the velocity width is $W_{50} = 11 \text{ km s}^{-1}$ (Table 1). The velocities appear to increase from southeast to northwest across this component, but the difference is only a few kilometers per second at the most (Fig. 4). The other component (hereafter the SW component) is centered at R.A. $17^{\text{h}}11^{\text{m}}35^{\text{s}}$, decl. $-64^{\circ}45'11''$ (J2000.0) and has a velocity spread from 451 to 477 km s^{-1} . The systemic velocity is 464 km s^{-1} and the velocity width is $W_{50} = 20 \text{ km s}^{-1}$ (Table 1). Any gradient across this component is again only a few kilometers per second (Fig. 4).

A bridge of emission appears to join the NE and SW components in a manner reminiscent of the Virgo cloud H I 1225+01 (Chengalur et al. 1995). Velocities appear to increase smoothly from the NE to the SW components. The overall H I structure appears to be that of two separate, but interacting, components.

¹⁶ IRAF is distributed by the National Optical Astronomy Observatories, operated by the AURA, under contract with the NSF.




FIG. 5.—Combined *B*- and *R*-band CCD optical images for HIPASS J1712–64, with Parkes and ATCA H I contours superposed.

3.2. Optical Limits

The source is located close to the Galactic plane at $(l, b) = (326^\circ.6, -14^\circ.6)$, and therefore extinction may be important. We have estimated this using three methods. The local H I column density from the HIPASS data (reprocessed to avoid any spatial filtering) is 484 K km s^{-1} , which corresponds to $N_{\text{H I}} = 8.8 \times 10^{20} \text{ atoms cm}^{-2}$. Following the conversion formula of Burstein & Heiles (1978), this implies $E(B - V) = 0.12 \text{ mag}$. The extinction determined from the *COBE* DIRBE and *IRAS* ISSA maps based on dust tem-

perature is essentially identical: $E(B - V) = 0.11 \text{ mag}$ (Schlegel, Finkbeiner, & Davis 1998). Finally, the Burstein & Heiles (1982) extinction maps suggest a similar value $E(B - V) = 0.09 \text{ mag}$. Taking the mean value, $E(B - V) = 0.11$, this implies a blue-light absorption of $A_B = 4.0E(B - V) = 0.44 \text{ mag}$. Thus although the stellar density in the optical images is quite high (see Fig. 5), there is no evidence for large optical extinction in the direction of the H I detection.

An optical counterpart could be overlooked if J1712–64 is very close to the Milky Way and resolved into stars. One

TABLE 2
DERIVED PARAMETERS FOR HIPASS J1712–64 AND ITS COMPONENTS

Parameters	NE Component	SW Component	J1712–64
Heliocentric velocity, v_{sys}	451 km s ⁻¹	464 km s ⁻¹	...
Dynamical LSR velocity, v_{lsr}	449 km s ⁻¹	462 km s ⁻¹	...
Galactocentric velocity, ^a v_{gsr}	332 km s ⁻¹	344 km s ⁻¹	...
Local Group velocity, ^b v_{LG}	239 km s ⁻¹	251 km s ⁻¹	...
Distance ^c	3.2 Mpc	3.2 Mpc	3.2 Mpc
Spatial size	11 × 8 kpc	5 × 4 kpc	15 × 8 kpc
H I mass, $M_{\text{H I}}$	$1.1 \times 10^7 M_{\odot}$	$3 \times 10^6 M_{\odot}$	$1.7 \times 10^7 M_{\odot}$
Dynamical mass, ^d M_T	$\sim 3 \times 10^7 M_{\odot}$...	$> 1.5 \times 10^8 M_{\odot}$
Blue luminosity, L_B	$< 7 \times 10^5 L_{\odot}$
$M_{\text{H I}}/L_B$	$> 24 M_{\odot}/L_{\odot}$

^a $v_{\text{gsr}} = v_{\text{lsr}} + 220 \sin l \cos b$.
^b Using the formula of Yahil et al. 1977.
^c Assuming an identical distance for the NE and SW components and using $d = v_{\text{LG}}/H_0$ with $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$.
^d See text.

method of detecting a resolved galaxy in a crowded field of stars is to search a color-color plot of the stars in a field to look for a separate stellar population. This method was used to detect the Sagittarius dwarf galaxy (Ibata, Gilmore, & Irwin 1995). We applied this method to the *B*- and *R*-band CCD images. These images cover an area of about 1 deg^2 , which is substantially larger than the H I detection. Fig. 6 shows a histogram of stellar magnitudes for a $20'$ field centered on the H I detection, and other nearby fields of the same size. These fields are statistically equivalent, and thus there is no evidence for a resolved galaxy in the direction of the H I detection.

3.3. Distance and Mass

The systemic velocity of the H I detection is $v_{\text{sys}} = 451 \text{ km s}^{-1}$ (NE component). The velocity with respect to the Local Group can be determined using $v_{\text{LG}} = v_{\text{sys}} - 79 \cos l \cos b + 296 \sin l \cos b - 36 \sin b$ (Yahil, Tammann, & Sandage 1977). This gives $v_{\text{LG}} = 239 \text{ km s}^{-1}$, which, if we assume HIPASS J1712–64 to be extragalactic and $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$, corresponds to a distance of 3.2 Mpc (see Table 2). This would place HIPASS J1712–64 at some distance beyond the Local Group. This distance is subject to considerable uncertainty. Alternate Local Group velocities range from $v_{\text{LG}} = 291 \text{ km s}^{-1}$ (using $300 \sin l \cos b$) to $v_{\text{LG}} = 172 \text{ km s}^{-1}$ (POTENT, Bertschinger et al. 1990), giving distances of 3.9 Mpc and 2.3 Mpc, respectively. Without the aid of optical distance measures, subsequent calculations involving the distance to J1712–64 will therefore be based on the value of $3.2 (\pm 1) \text{ Mpc}$. At this distance, the projected separation of the NW and SW components is $\sim 9 \text{ kpc}$, the overall H I size is $\sim 15 \text{ kpc}$, and the H I mass is $\sim 1.7 \times 10^7 M_{\odot}$. For a substantially smaller distance, say 100 kpc (see § 4), the separation, size, and mass are $\sim 300 \text{ pc}$, $\sim 500 \text{ pc}$, and $\sim 1.7 \times 10^4 M_{\odot}$, respectively.

The velocities within J1712–64 are very low. The velocity separation of the NE and SW components is only 12 km s^{-1} , and the velocity dispersions within each component are

small. For the NE component, the mean velocity dispersion from the ATCA data is $\sigma \sim 4 \text{ km s}^{-1}$ (see Table 1), possibly slightly more for the SW component, although the S/N is too low for a reliable estimate.

If the velocity difference between the NE and SW components is indicative of the rotation of a bound, binary system at 3.2 Mpc, the minimum system mass is $1.5 \times 10^8 M_{\odot}$ (Faber & Gallagher 1979), 1 order of magnitude above the H I mass. The “virial” distance, at which the total mass is 1.3 times the H I mass (to account for He), is implausibly distant ($> 20 \text{ Mpc}$) implying that the system contains substantial dark matter ($\sim 9 M_{\text{HI}}$, which would be a reasonable value for disk galaxies), or else is not bound. The orbital period of the two components corresponding to the minimum total mass is $\sim 4 (d/3.2 \text{ Mpc}) \text{ Gyr}$.

The NE component alone can be modeled as a pressure-supported sphere of gas of total enclosed mass

$$M_T = \alpha \sigma^2 r / G, \quad (1)$$

where σ is the line-of-sight velocity dispersion ($\sim 4 \text{ km s}^{-1}$), r is the radius ($\sim 4 \text{ kpc}$), and $\alpha \sim 1\text{--}3$ depending on the mass distribution. For an isothermal sphere, $\alpha \approx 2$, which gives $M_T(\text{NE}) = 3.0 \times 10^7 M_{\odot}$, only ~ 1.8 times higher than the H I mass for this component. Again, for a distance of 100 kpc , the mass drops to $M_T(\text{NE}) = 10^6 M_{\odot}$, but this is now ~ 60 times greater than the H I mass if in hydrostatic equilibrium.

3.4. Stability

The critical column density, above which local axisymmetric instabilities can occur in a uniformly rotating gaseous disk, is given by the Toomre stability criterion:

$$\Sigma_c = \frac{2v_s \Omega}{\pi G}, \quad (2)$$

where v_s is the sound speed and Ω is angular frequency (Binney & Tremaine 1987). Kennicutt (1989) has shown that similar relations for differentially rotating stellar disks approximately predict the column density above which star formation occurs in spiral galaxies. If we use $\Omega = v/r$ with $v \approx \sigma \approx v_s$ and $r = 4 \text{ kpc}$ for the NE component of J1712–64 (assuming it to be extragalactic), then we obtain $\Sigma_c \approx 8 \times 10^{19} \text{ atoms cm}^{-2}$. This is a factor of 2 above the peak measured column density of just $3.5 \times 10^{19} \text{ atoms cm}^{-2}$ (Fig. 3), consistent with the absence of visible star formation.

4. ALTERNATIVES

4.1. A Local Group Galaxy?

Nearby, faint galaxies that are resolved into stars can be especially hard to spot even far from the crowding in the Galactic plane (e.g., Sextans, Irwin et al. 1990). In § 3.2, we have shown that there is no evidence in color-magnitude diagrams for this. In addition, the Local Group velocity, $v_{\text{LG}} = 239 \text{ km s}^{-1}$ is well beyond the $\pm 60 \text{ km s}^{-1}$ that appears to encompass most known Local Group members (Grebel 1997; van den Bergh 1999). This is demonstrated in Figure 7, where J1712–64 has the highest heliocentric and Local Group velocity of all the probable and possible Local Group members plotted. It also has a higher velocity than the class of “Local Group outliers,” which have distances of $\sim 1\text{--}2 \text{ Mpc}$ and are regarded by Grebel (1997) as only potential members.




FIG. 6.—Number counts of stars in the central region of the *B*-band image and two surrounding regions. The stars show star number counts in the region where we would expect to see an excess of stars. No systematic excess is seen.




FIG. 7.—Velocities of Local Group members and outliers plotted against the cosine of the angle θ from the apex of solar motion relative to the Local Group barycenter (Yahil et al. 1977). HIPASS J1712–64 has the highest heliocentric and Local Group velocity, placing it beyond the Local Group outliers, which are themselves only considered to be distant potential members (Grebel 1997).

In addition, the irregular galaxy IC 4662 lies $3^{\circ}7'$ in the sky from HIPASS J1712–64 and has a velocity of $v_{\text{sys}} = 302 \pm 1 \text{ km s}^{-1}$, almost 150 km s^{-1} lower, yet is not classified as belonging to the Local Group (van den Bergh 1999). Therefore, from its velocity, there is no evidence for J1712–64 being a galaxy lying within the Local Group.

4.2. A High-Velocity Cloud?

The most likely alternative explanation for the classification of HIPASS J1712–64 is that it is an unusual high-velocity cloud (HVC) lying at the edge of our own Galaxy. HVCs are H I objects that do not fit into a simple model of Galactic rotation and do not have optical counterparts (Wakker & van Woerden 1997; Braun & Burton 1999). They are widespread across the sky, with a covering factor estimated to be 0.2–0.4 for $N_{\text{HI}} > 7 \times 10^{17} \text{ atoms m}^{-2}$ (Wakker & van Woerden 1997). There are two good reasons to exclude the possibility of J1712–64 being a normal HVC. The first is its velocity, which is 40 km s^{-1} higher than any currently cataloged HVC (e.g., those around the Large Magellanic Cloud, Morras et al. 2000). The second is J1712–64's H I structure. Most HVCs have a core-halo morphology with only 10%–50% of the total flux able to be recovered by interferometry (Wakker & van Woerden 1997). As shown in Figure 1, however, the ATCA observations were able to recover 80% of the total flux. It may, however, be an extreme version of the compact HVCs studied in detail by Braun & Burton (2000). These objects exhibit narrow line widths and also show apparent velocity gradients.

J1712–64 lies in a 2400 deg^2 region around the south celestial cap (Putman et al. 1998), where many of the HVCs appear to have originated from the interaction of the Magellanic Clouds with the Milky Way. It is possible that a compact HVC, similar to J1712–64, could have originated from gravitational scattering off the three-body Magellanic Clouds–Galaxy system. However, HIPASS J1712–64 does not lie close to the Clouds (it is 46° from the LMC), and the Leading Arm, which includes HVCs 334 and 352 of Wakker & van Woerden (1991, see Fig. 3 of Putman et al. 1998), is $\sim 20^{\circ}$ away. A velocity separation of $\sim 300 \text{ km s}^{-1}$ and a spatial separation from the LMC of $\sim 60 \text{ kpc}$ (or $\sim 80 \text{ kpc}$

from the Galaxy) would be consistent with an ejection event coinciding with the last LMC–SMC encounter, which probably occurred $\sim 2 \times 10^8 \text{ yr}$ ago (Gardiner & Noguchi 1996). An examination of the spatial and velocity distribution of future objects, similar to HIPASS J1712–64, will ultimately demonstrate their relationship, if any, with the Magellanic System.

5. DISCUSSION

Although there have been many previous searches for extragalactic H I in optically blank fields (Fisher & Tully 1981a; Briggs 1990; Barnes et al. 1997; Schneider et al. 1998), there has been little success, implying that massive H I “protogalaxies” are rare at the present epoch. However, HIPASS J1712–64 has a very low H I mass, $1.7 \times 10^7 M_{\odot}$, implying that similar objects may exist in abundance and may have escaped previous detection through lack of sensitivity. Some limits exist from the Cen A HIPASS results of Banks et al. (1999), who found 10 new group members, all with corresponding optical counterparts, in the 600 deg^2 they surveyed around the Cen A group. However, their sensitivity limit is $\sim 10^7 M_{\odot}$, so it is possible that slightly deeper observations (Disney et al. 1999) may reveal more such objects. The HIPASS observations of the south celestial cap, decl. $< -62^{\circ}$ (Kilborn et al. 1999), comprise 2400 deg^2 of sky, but contains only a few other candidate “protogalaxies,” which are awaiting follow-up CCD data. However, HIPASS J1712–64 is the most significant detection in terms of flux density and angular size. Nevertheless, for there to exist large numbers of similar objects would require them to have H I masses $< 10^7 M_{\odot}$.

Interestingly, a system with a similar velocity and H I morphology, ZOA J1616-55, was found in a search for galaxies in the zone of avoidance (ZOA) near $l = 325^{\circ}$ (Staveley-Smith et al. 1998). This system has a similar distance and H I column density to HIPASS J1712–64 but a higher H I mass ($9 \times 10^7 M_{\odot}$) and diameter (86 kpc). Although lying in the ZOA, the estimated blue absorption was only $A_B = 2.7 \text{ mag}$, so the lack of an optical or IR counterpart led Staveley-Smith et al. (1998) to conclude it was a galaxy pair of low optical surface brightness. J1712–64 is only 11° (projected separation 600 kpc) from J1616-55 and 19° (projected separation 1.1 Mpc) from the massive Circinus Galaxy. Possibly, all three are part of a loose southern extension to the Cen A group. In the HVC hypothesis, the alternative is that Circinus is unrelated and both J1712–64 and J1616-55 have a Magellanic origin.

From a study of the Ly α forest lines and a VLA H I study, Shull, Penton, & Stocke (1999) suggest that, at $z = 0$, there may exist clouds, or sheets, of ionized gas $\sim 1 \text{ Mpc}$ in extent that contain ~ 30 times more baryons than the neutral gas confined to known galaxies. Objects such as J1712–64 may be the most easily visible manifestations of these large primordial clouds. So, although the neutral gas content in J1712–64 is hardly enough to make a star cluster, let alone a galaxy, there may be significantly more baryons in the region available to form a galaxy if they were able to cool. The possibility that the dynamical mass of the J1712–64 system is ~ 10 times greater than the H I mass suggests there may be dark matter of some sort associated with the system.

Isolated H I clouds can be used to set stringent limits on the local metagalactic background radiation. Deep narrow-band CCD images of the H α line in H I 1225+01 and the

Leo ring place limits of $J_0 < 3-8 \times 10^{-23}$ ergs s $^{-1}$ cm $^{-1}$ sr $^{-2}$ Hz $^{-1}$ for quasar-like light (Donahue et al. 1995). H α observations of J1712–64 may also help provide strong constraints on this background. Such observations would also be useful in providing a lower limit to the distance to J1712–64 in light of the observation that many nearby HVCs have faint but detectable H α emission probably resulting from the Galactic ionization field (Bland-Hawthorn & Maloney 1999).

The mean velocity dispersion of the NE component of J1712–64 appears to be low, $\sigma = 4$ km s $^{-1}$ (Table 1). This gives an upper limit to the kinetic temperature of $T < 2000$ K. The existence of cool gas would be interesting as this requires that there are densities high enough to allow cooling and may therefore indicate that star formation is about to commence. In order to search for cool gas, we examined the ATCA H I spectrum of PKS 1708-648, which is 19.4 from the center of the J1712–64. However, we were unable to detect any unambiguous evidence for absorption. This is probably not significant as the continuum source is well beyond the last measured contour of HIPASS J1712–64.

6. SUMMARY

An isolated H I cloud, HIPASS J1712–64, has been found during the course of the HIPASS survey. It appears to be a nearby extragalactic cloud lying beyond the outer fringes of the Local Group at a distance of ~ 3.2 Mpc, though an alternative explanation is that of an HVC with unique properties. Optical limits imply a surface brightness,

corrected for obscuration, of $\mu(B) > 27$ mag arcsec $^{-2}$ and an H I mass-to-light ratio $M_{\text{HI}}/L_B > 24 M_{\odot}/L_{\odot}$ for a putative optical counterpart of diameter 1'. The system appears to have two components, separated by 9 kpc, with a bridge of gas between them. The mean velocity dispersion of the NE component is low, ~ 4 km s $^{-1}$ implying that HIPASS J1712–64 contains cool gas ($T < 2000$ K). However, it appears to be dynamically stable against star formation.

Although the H I mass is low, $\sim 1.7 \times 10^7 M_{\odot}$ (for a distance of 3.2 Mpc), the velocity differences within the system suggest that there may be an extended dark matter halo if the system is bound. Is HIPASS J1712–64 important from a cosmological point of view? This depends on whether similar objects exist in great numbers, whether they represent the cool, high density baryonic peaks of low-amplitude overdensities, whether such objects are the building blocks of galaxy and star cluster formation, whether they are simply the leftovers of these processes, and whether they are indeed extragalactic. Future HIPASS observations will cast some light on the frequency of their occurrence, and their distribution in the sky.

We are grateful to Michael Brown for assistance with analysis of the CCD data, using SExtractor, to Emma Ryan for assistance with the ATCA observing, to the staff at the Parkes telescope, and to members of the ZOA observing team for assistance with observations. P. K. acknowledges partial support by a grant from NASA administered by the American Astronomical Society.

REFERENCES

- Banks, G. D., et al. 1999, *ApJ*, 524, 612
 Barnes, D. G., Staveley-Smith, L., Walsh, W., & Webster, R. L. 1997, *MNRAS*, 288, 307
 Barnes, D. G., Staveley-Smith, L., Ye, T., & Oosterloo, T. 1998, in *ASP Conf. Ser. 145, Astronomical Data Analysis Software and Systems VII*, ed. R. Albrecht, R. N. Hook, & H. A. Bushouse (San Francisco: ASP), 89
 Barnes, D. G., et al. 2000, *MNRAS*, submitted
 Bertschinger, E., Dekel, A., Faber, S. M., Dressler, A., & Burstein, D. 1990, *ApJ*, 364, 370
 Binney, J., & Tremaine, S. 1987, *Galactic Dynamics* (Princeton: Princeton Univ. Press), 310
 Bland-Hawthorn, J., & Maloney, P. R. 1999, *ApJ*, 510, L33
 Braun, R., & Burton, W. B. 1999, *A&A*, 341, 437
 ———, 2000, *A&A*, 354, 853
 Briggs, F. H. 1990, *AJ*, 100, 999
 Burstein, D., & Heiles, C. 1978, *ApJ*, 225, 40
 ———, 1982, *AJ*, 87, 1165
 Chengular, J. N., Giovanelli, R., & Haynes, M. P. 1995, *AJ*, 109, 2415
 Dickey, J. M. 1997, *AJ*, 113, 1939
 Disney, M. J., Boyce, P. J., Banks, G. D., Minchin, R. F., & Wright, A. E. 1999, *PASA*, 16, 66
 Donahue, M., Aldering, G., & Stocke, J. T. 1995, *ApJ*, 450, L45
 Faber, S. M., & Gallagher, J. 1979, *ARA&A*, 17, 135
 Fisher, J. R., & Tully, R. B. 1981a, *ApJ*, 243, L23
 ———, 1981b, *ApJS*, 47, 139
 Gardiner, L. T., & Noguchi, M. 1996, *MNRAS*, 278, 191
 Giovanelli, R., & Haynes, M. P. 1989, *ApJ*, 346, L5
 Giovanelli, R., Scodreggio, M., Solanes, J. M., Haynes, M. P., Arce, H., & Sakai, S. 1995, *AJ*, 109, 1451
 Grebel, E. K. 1997, *Rev. Mod. Astron.*, 10, 29
 Henning, P. A. 1995, *ApJ*, 450, 578
 Hibbard, J. E., & van Gorkom, J. H. 1996, *AJ*, 111, 655
 Ibata, R. A., Gilmore, G., & Irwin, M. J. 1995, *MNRAS*, 277, 781
 Irwin, M. J., Bunclark, P. S., Bridgeland, M. T., & McMahon, R. G. 1990, *MNRAS*, 244, 16P
 Kennicutt, R. C. 1989, *ApJ*, 344, 685
 Kilborn, V., Webster, R. L., & Staveley-Smith, L. 1999, *PASA*, 16, 8
 Kraan-Korteweg, R. C., van Driel, W., Briggs, F., Bingelli, B., & Mostefaoui, T. I. 1999, *A&AS*, 135, 255
 Krumm, N., & Brosch, N. 1984, *AJ*, 89, 1461
 Landolt, A. U. 1983, *AJ*, 88, 439
 Malin, D., & Hadley, B. 1997, in *ASP Conf. Ser. 116, The Nature of Elliptical Galaxies*, 2d Stromlo Symposium, ed. M. Arnaboldi, G. S. Da Costa, & P. Saha (San Francisco: ASP), 460
 Malin, D. F. 1978, *Nature*, 276, 591
 Mathewson, D. S., Cleary, M. N., & Murray, J. D. 1975, *ApJ*, 195, L97
 Morras, R., Bajaja, E., Arnal, E. M., & Pöppel, W. G. L. 2000, *A&AS*, 142, 25
 Putman, M. E., et al. 1998, *Nature*, 394, 752
 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
 Schneider, S. E. 1989, *ApJ*, 343, 94
 Schneider, S. E., Helou, G., Salpeter, E. E., & Terzian, Y. 1983, *ApJ*, 273, L1
 Schneider, S. E., Spitzak, J. G., & Rosenberg, J. L. 1998, *ApJ*, 507, L9
 Shostak, G. S. 1977, *A&A*, 54, 919
 Shull, J. M., Penton, S. V., & Stocke, J. T. 1999, *PASA*, 16, 95
 Sorar, E. 1994, Ph.D. thesis, Univ. Pittsburgh
 Spitzak, J. G., & Schneider, S. E. 1998, *ApJS*, 119, 159
 Staveley-Smith L., et al. 1996, *PASA*, 13, 243
 ———, 1998, *AJ*, 116, 2717
 van den Bergh, S. 1999, *A&A Rev.*, 9, 273
 Wakker B. P., & van Woerden, H. 1991, *A&A*, 250, 509
 ———, 1997, *ARA&A*, 35, 217
 Williams B. A., & van Gorkom J. H. 1988, *AJ*, 95, 352
 Yahil, A., Tammann, G. A., & Sandage, A. 1977, *ApJ*, 217, 903
 Zwaan, M. A., & Briggs, F. H., 2000, *ApJ*, 530, L61
 Zwaan, M. A., Briggs, F. H., Sprayberry, D., & Sorar, E. 1997, *ApJ*, 490, 173