



An improved medium for the anaerobic growth of *Paracoccus denitrificans* Pd1222

Stefanie M. Hahnke^{1*}, Philipp Moosmann², Tobias J. Erb² and Marc Strous^{1,3,4}

¹ Microbial Fitness Group, Max Planck Institute for Marine Microbiology, Bremen, Germany

² Microbial Physiology, Institute of Microbiology, Eidgenössische Technische Hochschule (ETH) Zürich, Zürich, Switzerland

³ Microbiology of Sustainable Energy Production, Center for Biotechnology, Institute for Genome Research and Systems Biology, University of Bielefeld, Bielefeld, Germany

⁴ Department of Geoscience, Energy Bioengineering, University of Calgary, Calgary, AB, Canada

Edited by:

Rich Boden, University of Plymouth, UK

Reviewed by:

Rich Boden, University of Plymouth, UK

Michael Cunliffe, Marine Biological Association of the United Kingdom, UK

Hendrik Schäfer, University of Warwick, UK

Lesley Robertson, Delft University of Technology, Netherlands

*Correspondence:

Stefanie M. Hahnke, Microbial Fitness Group, Max Planck Institute for Marine Microbiology, Celsiusstrasse 1, 28359 Bremen, Germany
e-mail: smueller@mpi-bremen.de

Paracoccus denitrificans is a well studied model organism with respect to its aerobic and anaerobic respiratory enzymes. However, until now, the growth medium for this organism has not been optimized for anaerobic growth. In particular, the requirements of *P. denitrificans* for trace elements (TEs) are not well known. In the present study we aimed to improve growth rates of *P. denitrificans* Pd1222 on a defined medium under anoxic conditions. We designed media containing different combinations of TEs at various concentrations, and tested their performance against previously reported media. Our results suggest that growth rate and yield depend on the availability and concentration of TEs in the medium. A chelated TE solution was more suitable than an acidified TE solution. Highest growth rates were achieved with medium comprising the TEs iron, manganese, molybdenum, copper and zinc ranging from 0.1 to 9 μ M. On this medium, *P. denitrificans* Pd1222 grew with a generation time of 4.4 h under anoxic conditions and 2.8 h under oxic conditions. Diauxic growth was clearly shown with respect to nitrate and nitrite reduction under anoxic conditions.

Keywords: *Paracoccus denitrificans*, anaerobic growth, cultivation, defined medium, trace elements, copper, chelator

INTRODUCTION

The first strain of *Paracoccus denitrificans* (synonym *Micrococcus denitrificans*) was isolated from soil more than one century ago by Beijerinck and Minkman (1910). It was shown to grow aerobically and anaerobically performing complete or partial denitrification. Although *P. denitrificans* is an important model organism to study the electron transfer chain and energy conservation (for review see Stouthamer, 1991), there is still a lack of detailed information about the requirements of this organism for optimal growth. However, cultivation conditions have an important influence onto the physiological phenotype of an organism. Prominent examples for physiological processes that are strongly influenced by cultivation are the aerobic oxidation of methane (Stanley et al., 1983; Prior and Dalton, 1985) and methanol, the fixation of nitrogen gas (Lehman and Roberts, 1991), and the assimilation of CO₂, which make use of different enzymes (or even pathways) depending on the composition of the growth medium [most notably the presence or absence of vitamins and trace elements (TEs)]. Consequently, it is important to improve cultivation conditions of an organism and prevent its growth inhibition to avoid misinterpretation of the observed phenotypes.

Several approaches have been developed for medium optimization. Approaches based on evolutionary algorithms have been used lately to develop and optimize media for the isolation of novel bacterial strains without making assumptions about the individual components of a medium (Heylen et al., 2006). In contrast, rational approaches rely on the systematic improvement of existing media by focusing on a particular subset of medium

components, such as vitamin or TE supplements. In this study, we followed a rational strategy focusing on TE composition to improve the medium for the anaerobic growth of *P. denitrificans* Pd1222 on acetate.

P. denitrificans grows aerobically with maximum growth rate at pH 7.6 and at 36°C and can tolerate salt concentrations of at least 3% (Nokhal and Schlegel, 1983). Whereas the suitability of different carbon sources of *P. denitrificans* has been characterized in detail (Nokhal and Schlegel, 1983; Kelly et al., 2006), its requirements for TEs have not been investigated extensively. Different kinds of TEs, as well as large ranges of concentrations have been previously used in different studies (an overview of common TE solutions used for the cultivation of *P. denitrificans* and related organisms is given in Table 1). These solutions have been frequently used by many researchers (Meijer et al., 1979; Stouthamer and Bettenhausen, 1980; Van Spanning et al., 1990; Moir and Ferguson, 1994; Sears et al., 1997). TEs are essential for the correct function of enzymes, such as those of the respiratory chain; however, at higher concentrations they can impair growth and even be toxic. Here we present results on improvement of anaerobic growth of *P. denitrificans* Pd1222 with focus on TE requirements.

MATERIALS AND METHODS

ORGANISMS

Paracoccus denitrificans Pd1222 (16S rRNA gene accession number NR_074152), a derivative of DSM 413^T (De Vries et al., 1989), was obtained from Prof. Dr. R. van Spanning, Vrije Universiteit Amsterdam, faculty of Earth and Life Sciences. The organism was

Table 1 | Compositions of frequently used trace element solutions in the literature (final concentrations in μM in the medium).

Property and composition of solution	References							
	Robertson and Kuenen (1983)	Strohm et al. (2007)	Lawford (1978)	Widdel and Pfennig (1981)	Widdel and Bak (1992)	Chang and Morris (1962)	Nokhal and Schlegel (1983)	Harms et al. (1985)
Metal dissolution	EDTA	Acidic	Acidic	Acidic	EDTA	Iron citrate	Iron citrate	EDTA, citric acid
Na ₂ -EDTA	342.2				14.0			
Iron (II)	36.0 ^a	7.5 ^b		7.5 ^b	7.5 ^a	19.8 ^a		20.0 ^a
Iron (III)			90.0 ^b				4.6 ^c	
Manganese (II)	51.1 ^b	0.5 ^b	50.0 ^b	0.5 ^b	0.5 ^b	4.5 ^a	0.2 ^b	13.2 ^a
Copper (II)	12.6 ^a	0.01 ^b	5.0 ^b	0.1 ^b	0.01 ^b		0.1 ^b	
Molybdenum (VI)	1.8 ^d	0.2 ^e	10.0 ^e	0.2 ^e	0.2 ^e	728.4 ^e	0.1 ^e	599.0 ^e
Cobalt (II) ^b	13.5	0.8	10.0	0.8	0.8		0.8	
Zinc (II)	27.3 ^a	0.5 ^b	25.0 ^b	0.5 ^b	0.5 ^a		0.4 ^a	
Boron (III) ^f		0.1		1.0	0.5		4.9	
Nickel (II) ^b		0.1		0.1	0.1		0.1	
Relevant organism studied	<i>P. denitrificans</i> DSM 413 ^T	<i>P. pantotrophus</i> DSM 65 ^T	<i>P. denitrificans</i> ATCC 13543	<i>Desulfobacter postgatei</i>	Sulfate-reducing bacteria	<i>Micrococcus denitrificans</i>	<i>P. denitrificans</i> DSM 413 ^T and other strains	<i>P. denitrificans</i> various strains
Comments	Modified after Vishniac and Santer (1957)	According to Widdel (1983)	Modified after Light and Garland (1997)				Modified after Pfennig (1974)	
Reproduced in this study	TE-1, Figure 1A	TE-2, Figure 1B	Not shown	Not shown	Figures 2A,B	n.a.	n.a.	n.a.

TE, trace element solution; n.a., not applicable.

^aElements supplied as sulphates.

^bElements supplied as chlorides.

^cIron supplied as Fe(III)NH₄-citrate.

^dMolybdenum supplied as (NH₄)₆Mo₇O₂₄.

^eMolybdenum supplied as Na₂MoO₄.

^fBoron supplied as H₃BO₃.

maintained aerobically on solid LB medium, containing 15 g/L agar, and transferred every three months. For long-term storage the cells were frozen at -80°C in 30% glycerol and revived by spreading frozen cells on LB agar plates and incubating aerobically at 30°C for 2–3 days.

MINERAL SALT MEDIA

The following chemicals were purchased from AppliChem, Darmstadt, Germany: MgSO₄ · 7 H₂O, CaCl₂ · 2 H₂O, K₂HPO₄ and sodium acetate. All other chemicals were received from Carl Roth GmbH, Karlsruhe, Germany. The purity was at least 99% for most chemicals, except for ZnCl₂, 97%, MnCl₂ · 4 H₂O, 98% and NiCl₂ · 6 H₂O, 98%, whereas major impurities comprised sulphates or chlorides.

The experiments presented here were grouped into series 1–5. The mineral salt medium of series 1 was prepared after Taylor and Hoare (1969). The medium was supplemented with two different

TE solutions which are described below. For series 2–5, a freshwater medium modified after Widdel and Bak (1992) was used, containing (in g/L): NH₄Cl (0.5), MgSO₄ · 7 H₂O (0.5), CaCl₂ · 2 H₂O (0.1), KH₂PO₄ (0.04), K₂HPO₄ (0.12), and HEPES (6.0). Phosphate was added from a separately prepared and autoclaved stock solution. This medium was used to test three different TE solutions previously described (see below).

Materials used for medium preparation were rinsed with ultra pure water (Aquintus system, membraPure, Germany) prior to usage. All media were prepared with ultra pure water and the pH was adjusted to 7.2 with 1 M HCl or 1 M NaOH if necessary. The media were supplemented with 60 mM sodium acetate as carbon and energy source and 30 mM KNO₃ as the electron acceptor were added to both anaerobic and aerobic cultures to ensure identical salt concentrations, unless otherwise stated. For anaerobic cultures, 10 mL or 30 mL mineral salt solutions including electron donor and acceptor were filled into Hungate tubes

or 50 mL serum flasks using a volumetric pipette, and capped with a butyl stopper. The headspace was exchanged by applying vacuum, supplying argon at a pressure of 1.5 bar, followed by rigorous shaking (Widdel and Bak, 1992). This procedure was repeated three times. Finally, overpressure was released through a second needle. For aerobic cultures, 50 mL medium were filled into 250 mL Erlenmeyer flasks. All media and stock solutions were autoclaved at 121°C for 25 min.

TRACE ELEMENT SOLUTIONS

The final TE concentrations in the media for all growth experiments in this study are listed in **Tables 1** and **2**. Five TE stock solutions that have been previously presented, were prepared, three acidic solutions and two solutions containing a chelator (**Table 1**). Solution TE-1 was prepared according to Vishniac and Santer (1957) as described by Robertson and Kuenen (1983) and contained (in mg/L): Na₂-EDTA (5000), FeSO₄ · 7 H₂O (500), CaCl₂ · 2 H₂O (728), MnCl₂ · 4 H₂O (506), CuCl₂ · 2 H₂O (107), Na₂MoO₄ · 2 H₂O (22), CoCl₂ · 6 H₂O (161), and ZnCl₂ (185). The acidic solution TE-2 was prepared according to Widdel (1983). The medium after Taylor and Hoare was supplemented with 2 mL/L solution TE-1, a combination which was previously used for the cultivation of *P. denitrificans* DSM 413^T (Robertson and Kuenen, 1992). Alternative growth experiments were performed with the same medium amended with 1 mL/L solution TE-2. Three more TE solutions were prepared (Lawford, 1978; Widdel and Pfennig, 1981; Widdel and Bak, 1992) and tested with medium after Widdel and Bak (1992). The solution after Lawford was previously used for the cultivation of *P. denitrificans* ATCC 13543 (Lawford, 1978). Series 2 of the growth curves was performed with a set of TE solutions, designed in this study (**Table 2**), with increasing Cu²⁺ concentrations that contained (in mg/L): Na₂-EDTA (7300), FeSO₄ · 7 H₂O (2500), MnCl₂ ·

4 H₂O (20), Na₂MoO₄ · 2 H₂O (242), CuCl₂ · 2 H₂O (17; 38; 85; 128; 170; 426). TE solutions for the growth experiments of series 3–5 (**Table 2**) comprised the components of the solutions of series 2 (with 85 mg/L CuCl₂ · 2 H₂O) and optional (in mg/L): CoCl₂ · 6 H₂O (238), ZnCl₂ (340), H₃BO₃ (30), NiCl₂ · 6 H₂O (24). Depending on the final TE concentrations, the errors caused by impurities of chemicals were: 0.7–7.4% (iron), 0.5–4.6% (Co), 0.2–8.4% (Zn), 0.7–3.6% (Mn), and 0.1–2.6% (Mo). The impact of impurities of copper and nickel was higher; therefore the concentrations given in the results were corrected for this error.

TE solution TE-1 and all solutions of series 2–5 were prepared as chelated stock solutions and the pH was adjusted to 6 with 5 M NaOH. Stock solutions were autoclaved at 121°C for 25 min and added to the media with syringes. To reduce the volumetric error, the TE stock solutions were diluted ten or one hundred times, allowing the addition of a larger volume to the mineral media in Hungate tubes.

ANAEROBIC AND AEROBIC GROWTH EXPERIMENTS

Depending on the medium to be tested, the inoculum was prepared in different ways: (1) several colonies from solid LB medium were suspended in 1 mL of the respective mineral salt solution described above. (2) Cells were grown aerobically on the medium to be tested to a final optical density (OD₆₀₀) of 0.2. Both aerobic and anaerobic cultures were inoculated with 1% cell suspension. The cultures were incubated at 30°C. Aerobic cultures were shaken constantly at 120 rpm. Anoxic cultivation at different temperatures (11–45°C) was performed in Hungate tubes in a temperature gradient metal incubator.

SAMPLE WITHDRAWING AND ANALYSES

From cultures with an initial volume of 30 mL or more, samples of 1–1.5 mL were taken aseptically in different time intervals

Table 2 | Compositions of trace element solutions tested in different series of growth experiments in this study (final concentrations in μM in the medium).

Compound	Series 1		Series 2 ^c		Series 3 ^d				Series 4 ^e	Series 5 ^f
	TE-1 ^a	TE-2 ^b	TE-3	TE-3-Co	TE-3-Zn	TE-3-B	TE-3-Ni	TE-4	TE-3-Zn	TE-3-Zn
EDTA	342.2		19.6	19.6	19.6	19.6	19.6	19.6	19.6	19.6
Iron (II)	36.0	7.5	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0
Manganese (II)	51.1	0.5	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Copper (II)	12.6	0.01	0.4 – 2.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Molybdenum	1.8	0.2	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Cobalt (II)	13.5	0.8		1.0				1.0		
Zinc (II)	27.3	0.5			2.5			2.5	2.5	2.5
Boron		0.1				0.5		0.5		
Nickel (II)		0.1					0.07	0.07		
pH	6	1	6	6	6	6	6	6	6	6

^a Trace element solution after Vishniac and Santer (1957)

^b Trace element solution after Widdel (1983).

^c Test of the impact of copper on growth.

^d Test of the impact of other trace elements on growth (also tested at ten times higher concentrations).

^e Aerobic growth on the improved medium.

^f Temperature dependence of anaerobic growth on the improved medium.


depending on medium and growth conditions. Bacterial growth was followed by measuring the OD₆₀₀ at 600 nm (cuvette path length 1 cm) with a spectrophotometer (Genesis 10S UV-VIS, Thermo Scientific). Three replicate growth curves were additionally followed by protein determination (Lowry et al., 1951), revealing a linear relationship between OD₆₀₀ and protein concentration. Subsequently, the sample was centrifuged at 14500 × g at 4°C for 5 min. The supernatant was removed and stored at -20°C for nitrate, nitrite and acetate determination. The cultures in Hungate tubes were not sampled. Instead, the optical density was measured at 600 nm directly in the culture tube with a spectrophotometer (UV-1201V, Shimadzu) in different time intervals depending on medium and growth conditions. The OD₆₀₀ values were corrected corresponding to measurements in cuvettes with 1 cm path length to allow comparison between all measured OD₆₀₀ values.

Nitrate was determined colorimetrically after reaction with salicylic acid (Cataldo et al., 1975). Nitrite was determined colorimetrically using the Griess-Romijn reagent (Griess-Romijn-van Eck, 1966). Acetate was determined with an HPLC system (Sykam GmbH) at a retention time of 13.8 min and a detection limit of 25 μM acetate. With 5 mM H₂SO₄ as eluent and a flow rate of 0.6 ml min⁻¹, acetate was detected using a refractive index (RI) or UV detector (210 nm).

RESULTS

Although *P. denitrificans* is a model organism for denitrification, we found that the growth media and TE solutions described for this organism enabled only very poor anaerobic growth of *P. denitrificans* Pd1222. This phenomenon was most apparent when growing this strain on medium (Taylor and Hoare, 1969) that was amended with either TE solution TE-1 (Vishniac and Santer, 1957; Robertson and Kuenen, 1983) or TE-2 (Widdel, 1983; Strohm et al., 2007). Both TE solutions have been used before for the cultivation of *P. denitrificans* or *P. pantotrophus* and are listed as final concentrations in the medium in **Table 1**. TE-1 is characterized by high metal concentrations between 1.8 and 36 μM and contains ethylenediaminetetraacetic acid (EDTA) as a chelator to keep the metals dissolved, while metals in TE-2 are 4 to more than 1000 times lower concentrated and metal dissolution is achieved by lowering the pH instead of adding EDTA. Under anoxic, denitrifying conditions, very slow growth was detected with a generation time of 12.3 h in medium supplemented with TE-1 and 9.3 h in medium supplemented with TE-2 (**Figure 1**; for specific growth rates, please also refer to **Table 3**). Thus, we sought for a defined medium which was more suitable for the cultivation of *P. denitrificans* Pd1222 under anoxic, denitrifying conditions.

Anaerobic growth of *P. denitrificans* Pd1222 was investigated with a different freshwater medium that was modified after Widdel and Bak (1992) and three different TE compositions as previously described (Lawford, 1978; Widdel and Pfennig, 1981; Widdel and Bak, 1992). The acidic TE solution after Lawford (**Table 1**) contained up to 100 times higher metal concentrations than the two other TE solutions (**Table 1**) that were prepared either as an acidic or a chelated (EDTA containing) solution. All media supplemented with acidic TE solutions supported only limited growth under anoxic, denitrifying conditions after one



week of incubation. In contrast, good growth was observed in the medium supplemented with chelated TE solution according to Widdel and Bak (1992), with a generation time of 8.1 h (**Figure 2A** and **Table 3**, at 10 mM acetate). This indicated that at least one essential compound was insufficiently available to the cells in the absence of a chelator. Based on these results all subsequent growth experiments were performed with TE solutions that contained EDTA for dissolution. The suitability of media subsequently tested, was estimated based on growth rates that are summarized in **Table 3**, and described in more detail below.

Trace metals such as iron and copper are required for the biosynthesis of essential enzymes, such as the membrane bound protein complexes of the aerobic and anaerobic respiratory chain. Thus we investigated whether the absence of an essential TE might have been the reason for growth limitation. For this purpose, we cultured *P. denitrificans* Pd1222 with increasing acetate and nitrate concentrations in the medium. In the absence of any (trace) element limitation, higher substrate concentrations should lead to proportionally higher growth yields. **Figure 2B** shows the growth curves of cultures supplied with 10–100 mM acetate (5–50 mM nitrate). With 10 mM acetate and 5 mM nitrate the maximum optical density (maxOD₆₀₀)

Table 3 | Growth characteristics of *Paracoccus denitrificans* Pd1222, grown in batch cultures under anoxic, denitrifying conditions unless otherwise stated.

Growth experiment	Parameter	Growth μ (h^{-1})	Generation time (h)
TRACE ELEMENT SOLUTION			
Medium according to the literature (Figure 1)	TE-1	0.056 \pm 0.013	12.3
	TE-2	0.075 \pm 0.002	9.3
ACETATE CONCENTRATION (mM)			
Trace elements according to Widdel and Bak (1992); increasing substrate concentrations (Figure 2)	10	0.086 \pm 0.002	8.1
	20	0.092 \pm 0.011	7.6
	40	0.093 \pm 0.001	7.4
	60	0.095 \pm 0.004	7.1
	100	0.097 \pm 0.006	7.6
Cu²⁺ CONCENTRATION (μM)			
Increasing Cu ²⁺ concentrations, anaerobic	0.4	0.123 \pm 0.004	5.7
	0.5	0.134 \pm 0.026	5.2
	0.8	0.133 \pm 0.012	5.2
	1.0	0.129 \pm 0.044	5.4
	1.3	0.157 \pm 0.003	4.4
	2.8	0.144 \pm 0.010	4.8
Increasing Cu ²⁺ concentrations, aerobic	0.5	0.159 \pm 0.008	4.4
	0.8	0.089 \pm 0.008	7.8
	1.0	0.054 \pm 0.002	12.8
	1.3	0.029 \pm 0.003	18.6
	2.8	0.021 \pm 0.007	32.4
TRACE ELEMENT SOLUTION			
Different trace element compositions (Figure 3A)	TE-3	0.129 \pm 0.002	5.4
	TE-3-Co	0.154 \pm 0.011	5.2
	TE-3-Zn	0.159 \pm 0.002	4.4
	TE-3-B	0.135 \pm 0.004	5.1
	TE-3-Ni	0.141 \pm 0.017	4.9
	TE-4	0.168 \pm 0.001	4.7
10 \times concentrated trace elements (Figure 3B)	TE-3	No growth (<0.008)	No growth (>84 h)
	TE-3-Co	0.104 \pm 0.007	6.7
	TE-3-Zn	0.125 \pm 0.033	5.5
	TE-3-B	No growth (<0.008)	No growth (>84 h)
	TE-3-Ni	No growth (<0.008)	No growth (>84 h)
	TE-4	0.129 \pm 0.035	5.4
Aerobic growth (Figure 4)	TE-3-Zn	0.251 \pm 0.032	2.8
TEMPERATURE ($^{\circ}\text{C}$)			
Increasing temperature (Figure 5)	11.2	0.010 \pm 0.001	67.7
	16.1	0.023 \pm 0.001	30.3
	19.3	0.044 \pm 0.001	15.9
	21.0	0.056 \pm 0.002	12.5
	22.5	0.063 \pm 0.003	11.0
	24.2	0.066 \pm 0.004	10.4
	25.9	0.061 \pm 0.008	11.3


(Continued)

Table 3 | Continued


Growth experiment	Parameter	Growth μ (h^{-1})	Generation time (h)
	27.4	0.084 \pm 0.006	8.3
	29.0	0.093 \pm 0.009	7.5
	30.5	0.104 \pm 0.004	6.7
	32.1	0.152 \pm 0.004	4.6
	33.7	0.188 \pm 0.007	3.7
	35.3	0.201 \pm 0.008	3.4
	36.9	0.214 \pm 0.015	3.3
	40.0	0.161 \pm 0.008	4.3

reached 0.23, whereas at 20 mM acetate and 10 mM nitrate the maxOD₆₀₀ reached 0.37, which is only a 1.6-fold increase instead of the expected 2-fold increase in maxOD₆₀₀. At ten times higher acetate and nitrate concentrations, the effect was even more pronounced, as the maxOD₆₀₀ was 0.96, which corresponded only a 4-fold increase instead of the 10-fold maxOD₆₀₀ which would be expected in the absence of any (trace) element limitation. In short, the increase in OD₆₀₀ was never proportional to the increase of substrate concentration. We may explain this observation in two ways. First, the limited availability of at least one TE might have limited growth. Second, non-optimal growth conditions might have promoted the accumulation of toxic metabolic intermediates that inhibit growth. In this case, the organism would require additional energy as biochemical response to the stress conditions at the cost of growth yield. The latter could also be caused by an inappropriate TE composition, leading to the accumulation of toxic intermediates (Sullivan et al., 2013). Thus, we examined whether growth rates could be improved by providing different concentrations of TEs.

Copper was the first TE that was investigated in more detail. It was supplied to different batch cultures in concentrations from 0.4 to 2.8 μM . In this series we used a TE solution containing (besides copper) only iron, manganese, and molybdenum, as these elements were sufficient to sustain aerobic and anaerobic growth of *P. denitrificans* (Chang and Morris, 1962). Our results did not suggest any trend in generation times with increasing copper concentration (between 4.4 and 5.7 h, **Table 3**). However the reproducibility was better at lower copper concentrations; at higher copper concentrations (1.0 μM and higher) at least one out of three replicate cultures did not grow. Overall a copper concentration of 0.8 μM was shown to be high enough for growth and low enough to prevent observed negative effects on growth. For comparison, we grew *P. denitrificans* Pd1222 with increasing copper concentrations under oxic conditions (**Table 3**). In this case, generation times increased with increasing copper concentrations, with a minimum of 4.4 h in medium supplemented with 0.4 μM copper and a generation time of 7.8 h at 0.8 μM copper. Growth was severely affected above 1.0 μM copper (generation time 12.8 h). Thus, a copper concentration of 0.8 μM was provided in the minimal TE solution TE-3 (**Table 2**) that was used for subsequent cultivation of *P. denitrificans* Pd1222 under anoxic conditions.





Next, we investigated the effect of other TEs on growth. For that purpose, TE solution TE-3 that contained only the four essential elements as described above, was amended with one of four additional TEs (cobalt, zinc, boron, and nickel) to obtain four modified TE-3 solutions as listed in **Table 2**. Solution TE-4 was obtained by adding all four TEs to solution TE-3. Growth of *P. denitrificans* Pd1222 was examined at two different concentrations of each modified TE-3 solution and TE-4. *P. denitrificans* Pd1222 grew in all cultures containing low concentrations of TEs (**Tables 2 and 3; Figure 3A**). In medium supplied with TE-3 the cultures grew at a generation time of 5.4 h. The shortest generation time of 4.4 h was achieved with the addition of TE-3-Zn, indicating that zinc positively affected growth of *P. denitrificans* Pd1222. Cultures supplemented with TE-3-Co and TE-3-B resulted in growth with a prolonged lag phase of approximately 22 h (10 h longer than in all other treatments). When combining all individual tested TEs into one TE solution TE-4, medium complemented with TE-4 enabled growth at a generation time of 4.7 h and did not show any improvement over the medium complemented with TE-3-Zn.



In cultures supplied with ten times higher concentrations of TE-3, modified TE-3 solutions and TE-4 solution growth was impaired as shown in **Figure 3B** (note that most of the final metal concentrations in these media correspond to concentrations that were reached with the TE solution after Lawford). No growth was observed in cultures supplied with ten times concentrated TE-3, TE-3-B, or TE-3-Ni after one week of incubation, and only two out of three replicate cultures grew on medium supplied with ten times concentrated TE-3-Co, TE-3-Zn, or TE-4. In this case, generation times were generally higher than in medium containing low concentrations of TEs as given in **Table 2**.

In conclusion, TE solution TE-3-Zn was the most suitable TE solution among all solutions tested in this study for the anaerobic, denitrifying growth of *P. denitrificans* Pd1222. With this solution, a highly reproducible and short generation time of 4.4 h was achieved. Compared to aerobic growth with a generation time of 2.8 h (**Figure 4**), the generation time was only 1.6 times longer (**Table 3**).

Besides TEs, temperature has a substantial influence on growth (**Table 3**). Anaerobic growth of *P. denitrificans* Pd1222 in medium




containing TE solution TE-3-Zn was observed between 11 and 40°C (**Figure 5**), with maximum growth rates at 37°C, similar to what was reported by Nokhal and Schlegel (1983). The generation time was 3.2 h. As compared to previous experiments in this study, these were the highest growth rates observed for anaerobic denitrifying growth of *P. denitrificans* Pd1222.

To study the performance of *P. denitrificans* Pd1222 during growth on the improved medium in more detail, the conversion of substrates and production of intermediates were analyzed (**Figure 6**). Nitrate was completely converted to nitrite (the first intermediate in denitrification) in the first half of the exponential growth and nitrite reached a maximum concentration of 62 mM, which corresponded to the initial nitrate concentration. At this point, growth was slowed down, which can be explained by the time required by *P. denitrificans* Pd1222 to induce the expression of nitrite reductase. After approximately 2 h, the growth rate was recovered and nitrite was reduced within 21 h. Accordingly, acetate was consumed with a lower rate during the transition from nitrate to nitrite respiration. This diauxic growth was previously observed (Błaszczuk, 1993; Saéz et al., 2003), but lower concentrations of nitrite were accumulated and consumed than in this study. The depletion of both nitrate and nitrite confirms that the electron acceptor limited growth, implying that trace and macro element concentrations were sufficiently high.

DISCUSSION

This study aimed at improving cultivation conditions for anaerobic, denitrifying growth of *P. denitrificans* Pd1222, to achieve high growth rates on a minimal medium that are comparable to growth rates in the presence of oxygen. Although *P. denitrificans* has been a long-standing model organism for the study of respiratory enzymes and energy conservation, it was surprising that *P. denitrificans* Pd1222 grew poorly on published media. Thus we sought to test and improve different media that had been used in the past to cultivate this and closely related organisms (Lawford, 1978; Robertson and Kuenen, 1983; Strohm et al., 2007).

Supplementation of media with TEs (and vitamins) in suitable concentrations is known to have an important influence on the



successful cultivation of microorganisms (Widdel, 1980; Peters et al., 2004; Pol et al., 2014). Consequently, this study focused on improving the composition of the TE solution. Our experiments suggested that for anaerobic growth of *P. denitrificans* Pd1222, chelator-based TE solutions are superior to solutions in which trace metals are solubilized and kept in their redox state by lowering the pH. Media that were used by many researchers to study *P. denitrificans* contain citrate as chelator (Chang and Morris, 1962; Lawford, 1978; Nokhal and Schlegel, 1983). Since *P. denitrificans* DSM 413^T has been reported to be able to utilize citrate under oxic, but not under anoxic conditions (Davis et al., 1970; Robertson and Kuenen, 1983), we note that citrate might become a potential carbon source under anoxic conditions, particularly during long-time cultivations. Similar observations have been recently made with *E. coli* (Blount et al., 2008, 2012). Therefore, we used EDTA instead of citrate as the chelator. One previously published medium for *P. denitrificans* DSM 413^T (Robertson and Kuenen, 1983) also featured EDTA, but

in this medium the concentrations of TEs ranged between 13 and 51 μM , notably higher than the 0.1–1 μM which are usually used for TEs (Overmann, 2010, 2013). Higher concentrations of TEs are known to exert toxic effects (Overmann, 2010), and our experiments indeed indicated growth-inhibiting effects of higher concentrations of TEs.

We achieved good growth of *P. denitrificans* Pd1222 with the TEs molybdenum, manganese and copper in concentrations between 0.1 and 1 μM , zinc at 2.5 μM and iron(II) at 9 μM . Three of these elements represent metals required as cofactors by the enzymes of the denitrification pathway of *P. denitrificans* (Stouthamer, 1991). Iron is present in all four respiratory enzymes as Fe-S clusters or as part of a cytochrome domain. The membrane-bound nitrate reductase (Nar) additionally requires molybdenum (Stouthamer, 1991) and the nitrous oxide reductase requires copper (Snyder and Hollocher, 1987). Zinc is not essential for the biosynthesis of the enzymes involved in denitrification, but the increased growth rates in presence of zinc might indicate that it stimulates anaerobic growth in a different way.

In summary, we provide a minimal medium that allows anaerobic growth of *P. denitrificans* Pd1222 at a minimum generation time of 4.4 h at 30°C without the need of vitamin addition. The generation times under anoxic conditions were only 1.6 times longer than under oxic conditions. This may suggest that a more intense study of concentration ranges of all TEs would only result in minor improvements of growth. Our medium is easy to prepare and allows physiological studies of the model organism *P. denitrificans* Pd1222 under anoxic conditions. With this medium, we show that *P. denitrificans* Pd1222 strictly grows diauxically with respect to nitrate and nitrite respiration. Nitrate was completely converted to nitrite before nitrite was reduced to dinitrogen.

We expect that our findings are also relevant to other strains of *P. denitrificans* since previous physiological studies of various strains of *P. denitrificans* showed identical physiological characteristics and similar carbon utilization patterns among the tested strains (Nokhal and Schlegel, 1983). These features specifically relate to major metabolic pathways that constitute the most trace metal dependent enzymes. Therefore, it is likely that the TE composition suggested here represents a suitable composition for more strains of *P. denitrificans*.

AUTHOR CONTRIBUTIONS

Stefanie M. Hahnke, Tobias J. Erb, and Marc Strous conceived the project, Stefanie M. Hahnke and Philipp Moosmann designed and performed the experiments, Stefanie M. Hahnke and Tobias J. Erb analyzed the data, Stefanie M. Hahnke, Tobias J. Erb, and Marc Strous wrote and edited the manuscript.

ACKNOWLEDGMENTS

We thank Maike Kortmann and Patrick de Jonge for help with monitoring growth curves and for nutrient analyses. We are grateful to Prof. Dr. Jens Harder, Prof. Dr. Friedrich Widdel, and Prof. Dr. Rob van Spanning for the scientific discussions and advice on cultivation. We thank Prof. Dr. Rob van Spanning for providing the bacteria. This research was funded by the ERC grant “MASEM” (242635, granted to Marc Strous), the German Federal State of Nordrhein-Westfalen, and the Swiss National Science

Foundation (SNF-Ambizione grant PZ00P3_136828/1, granted to Tobias J. Erb).

REFERENCES

- Beijerinck, M. W., and Minkman, D. C. J. (1910). Bildung und Verbrauch von Stickoxydul durch Bakterien. *Zentralbl. Bakteriol.* 25, 30–63.
- Blaszczak, M. (1993). Effect of medium composition on the denitrification of nitrate by *Paracoccus denitrificans*. *Appl. Environ. Microbiol.* 59, 3951–3953.
- Blount, Z. D., Barrick, J. E., Davidson, C. J., and Lenski, R. E. (2012). Genomic analysis of a key innovation in an experimental *Escherichia coli* population. *Nature* 489, 513–518. doi: 10.1038/nature11514
- Blount, Z. D., Borland, C. Z., and Lenski, R. E. (2008). Historical contingency and the evolution of a key innovation in an experimental population of *Escherichia coli*. *Proc. Natl. Acad. Sci. U.S.A.* 105, 7899–7906. doi: 10.1073/pnas.0803151105
- Cataldo, D. A., Maroon, M., Schrader, L. E., and Youngs, V. L. (1975). Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. *Commun. Soil Sci. Plant Anal.* 6, 71–80. doi: 10.1080/00103627509366547
- Chang, J. P., and Morris, J. G. (1962). Studies on the utilization of nitrate by *Micrococcus denitrificans*. *J. Gen. Microbiol.* 29, 301–310. doi: 10.1099/00221287-29-2-301
- Davis, D. H., Stanier, R. Y., Doudoroff, M., and Mandel, M. (1970). Taxonomic studies on some gram negative polarly flagellated “hydrogen bacteria” and related species. *Arch. Mikrobiol.* 70, 1–13.
- De Vries, G. E., Harms, N., Hoogendijk, J., and Stouthamer, A. H. (1989). Isolation and characterization of *Paracoccus denitrificans* mutants with increased conjugation frequencies and pleiotropic loss of a (nGATcN) DNA-modifying property. *Arch. Microbiol.* 152, 52–57. doi: 10.1007/BF00447011
- Griess-Romijn-van Eck (1966). *Physiological and Chemical Tests for Drinking Water*. Rijswijk: Nederlands Normalisatie Instituut.
- Harms, N., de Vries, G. E., Maurer, K., Veltkamp, E., and Stouthamer, A. H. (1985). Isolation and characterization of *Paracoccus denitrificans* mutants with defects in the metabolism of one-carbon compounds. *J. Bacteriol.* 164, 1064–1070.
- Heylen, K., Vanparys, B., Wittebolle, L., Verstraete, W., Boon, N., and Vos, P. D. (2006). Cultivation of denitrifying bacteria: optimization of isolation conditions and diversity study. *Appl. Environ. Microbiol.* 72, 2637–2643. doi: 10.1128/AEM.72.4.2637
- Kelly, D. P., Rainey, F. A., and Wood, A. P. (2006). “The Genus *Paracoccus*,” in *The Prokaryotes: A Handbook on the Biology of Bacteria*, eds M. Dworkin, S. Falkow, E. Rosenberg, K.-H. Schleifer, and E. Stackebrandt (New York, NY: Springer), 232–249.
- Lawford, H. G. (1978). Energy transduction in the mitochondrionlike bacterium *Paracoccus denitrificans* during carbon- or sulphate-limited aerobic growth in continuous culture. *Can. J. Biochem.* 56, 13–22. doi: 10.1139/o78-003
- Lehman, L. J., and Roberts, G. P. (1991). Identification of an alternative nitrogenase system in *Rhodospirillum rubrum*. *J. Bacteriol.* 173, 5705–5711.
- Light, P. A., and Garland, P. B. (1997). A comparison of mitochondria from *Torulopsis utilis* grown in continuous culture with glycerol, iron, ammonium, magnesium or phosphate as the growth-limiting nutrient. *Biochem. J.* 193, 265–275.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L., and Randall, R. J. (1951). Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193, 265–275.
- Meijer, E. M., Zwaan, J. W. V. D., Wever, R., and Stouthamer, A. H. (1979). Anaerobic respiration and energy conservation in *Paracoccus denitrificans*. *Eur. J. Biochem.* 96, 69–76. doi: 10.1111/j.1432-1033.1979.tb13014.x
- Moir, J. W. B., and Ferguson, S. J. (1994). Properties of a *Paracoccus denitrificans* mutant deleted in cytochrome c_{550} indicate that a copper protein can substitute for this cytochrome in electron transport to nitrite, nitric oxide and nitrous oxide. *Microbiol.* 140, 389–397. doi: 10.1099/13500872-140-2-389
- Nokhal, T.-H., and Schlegel, H. G. (1983). Taxonomic study of *Paracoccus denitrificans*. *Int. J. Syst. Bacteriol.* 33, 26–37. doi: 10.1099/00207713-33-1-26
- Overmann, J. (2010). “Novel cultivation strategies for environmentally important microorganisms,” in *Geomicrobiology: Molecular and environmental perspective*, eds L. L. Barton, M. Mandl, and A. Loy (Heidelberg: Springer), 69–89. doi: 10.1007/978-90-481-9204-5_3
- Overmann, J. (2013). “Principles of enrichment, isolation, cultivation, and preservation of prokaryotes,” in *The Prokaryotes*, eds E. Rosenberg, E. F. Delong, S. Lory, E. Stackebrandt, and F. Thompson (Berlin: Springer), 149–207.

- Peters, F., Rother, M., and Boll, M. (2004). Selenocysteine-containing proteins in anaerobic benzoate metabolism of *Desulfococcus multivorans*. *J. Bacteriol.* 186, 2156–2163. doi: 10.1128/JB.186.7.2156
- Pfennig, N. (1974). *Rhodospseudomonas globiformis*, sp. n., a new species of the Rhodospirillaceae. *Arch. Microbiol.* 100, 197–206. doi: 10.1007/BF00446317
- Pol, A., Barends, T. R. M., Dietl, A., Khadem, A. F., Eygensteyn, J., Jetten, M. S. M., et al. (2014). Rare earth metals are essential for methanotrophic life in volcanic mudpots. *Environ. Microbiol.* 16, 255–264. doi: 10.1111/1462-2920.12249
- Prior, S. D., and Dalton, H. (1985). The effect of copper ions on membrane content and methane monooxygenase activity in methanol-grown cells of *Methylococcus capsulatus* (Bath). *J. Gen. Microbiol.* 131, 155–163. doi: 10.1099/00221287-131-1-155
- Robertson, L. A., and Kuenen, G. (1983). *Thiosphaera pantotropha* gen. nov. sp. nov., a facultatively anaerobic, facultatively autotrophic sulphur bacterium. *J. Gen. Microbiol.* 129, 2847–2855. doi: 10.1099/00221287-129-9-2847
- Robertson, L. A., and Kuenen, J. G. (1992). The effect of electron acceptor variations on the behaviour of *Thiosphaera pantotropha* and *Paracoccus denitrificans* in pure and mixed cultures. *FEMS Microbiol. Lett.* 86, 221–228. doi: 10.1016/0378-1097(92)90785-M
- Saéz, F., Pozo, C., Gómez, M. A., Rodelas, B., and González-López, J. (2003). Growth and nitrite and nitrous oxide accumulation of *Paracoccus denitrificans* ATCC 19367 in the presence of selected pesticides. *Environ. Toxicol. Chem.* 22, 1993–1997. doi: 10.1897/02-351
- Sears, H. J., Spiro, S., and Richardson, D. (1997). Effect of carbon substrate and aeration on nitrate reduction and expression of the periplasmic and membrane-bound nitrate reductases in carbon-limited continuous cultures of *Paracoccus denitrificans* Pd1222. *Microbiology* 143, 3767–3774. doi: 10.1099/00221287-143-12-3767
- Snyder, S. W., and Hollocher, T. C. (1987). Purification and some characteristics of nitrous oxide reductase from *Paracoccus denitrificans*. *J. Biol. Chem.* 262, 6515–6525.
- Stanley, S. H., Prior, S. D., Leak, D. J., and Dalton, H. (1983). Copper stress underlies the fundamental change in intracellular location of methane mono-oxygenase in methane-oxidizing organisms: studies in batch and continuous cultures. *Biotechnol. Lett.* 5, 487–492. doi: 10.1007/Bf00132233
- Stouthamer, A. H. (1991). Metabolic regulation including anaerobic metabolism in *Paracoccus denitrificans*. *J. Bioenerg. Biomembr.* 23, 163–185. doi: 10.1007/BF00762216
- Stouthamer, A. H., and Bettenhausen, C. W. (1980). Growth and physiology of potassium-limited chemostat cultures of *Paracoccus denitrificans*. *Arch. Microbiol.* 125, 239–244. doi: 10.1007/BF00446883
- Strohm, T. O., Griffin, B., Zumft, W. G., and Schink, B. (2007). Growth yields in bacterial denitrification and nitrate ammonification. *Appl. Environ. Microbiol.* 73, 1420–1424. doi: 10.1128/AEM.02508-06
- Sullivan, M. J., Gates, A. J., Appia-Ayme, C., Rowley, G., and Richardson, D. (2013). Copper control of bacterial nitrous oxide emission and its impact on vitamin B12-dependent metabolism. *Proc. Natl. Acad. Sci. U.S.A.* 110, 19926–19931. doi: 10.1073/pnas.1314529110
- Taylor, B. F., and Hoare, D. S. (1969). New facultative *Thiobacillus* and a reevaluation of the heterotrophic potential of *Thiobacillus novellus*. *J. Bacteriol.* 100, 487–497.
- Van Spanning, R. J. M. V., Wansell, C., Harms, N., Oltmann, L. F., and Stouthamer, A. H. (1990). Mutagenesis of the gene encoding cytochrome *c*₅₅₀ of *Paracoccus denitrificans* and analysis of the resultant physiological effects. *J. Bacteriol.* 172, 986–996.
- Vishniac, W., and Santer, M. (1957). The Thiobacilli. *Bacteriol. Rev.* 21, 195–213.
- Widdel, F. (1980). *Anaerobier Abbau von Benzoesäure durch neu isolierte Arten Sulfat-reduzierender Bakterien*. Ph.D. thesis, Universität Göttingen.
- Widdel, F. (1983). Methods for enrichment and pure culture isolation of filamentous gliding sulfate-reducing bacteria. *Arch. Microbiol.* 134, 282–285. doi: 10.1007/BF00407803
- Widdel, F., and Bak, F. (1992). “Gram-negative mesophilic sulfate-reducing bacteria,” in *The Prokaryotes*, eds A. Balows, H. G. Trüper, M. Dworkin, W. Harder, and K.-H. Schleifer (New York, NY: Springer), 3352–3378.
- Widdel, F., and Pfennig, N. (1981). Studies on dissimilatory sulfate-reducing bacteria that decompose fatty acids. *Arch. Microbiol.* 129, 395–400. doi: 10.1007/BF00406470

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 15 October 2013; accepted: 12 January 2014; published online: 31 January 2014.

Citation: Hahnke SM, Moosmann P, Erb TJ and Strous M (2014) An improved medium for the anaerobic growth of *Paracoccus denitrificans* Pd1222. *Front. Microbiol.* 5:18. doi: 10.3389/fmicb.2014.00018

This article was submitted to *Microbial Physiology and Metabolism*, a section of the journal *Frontiers in Microbiology*.

Copyright © 2014 Hahnke, Moosmann, Erb and Strous. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.