



An *In-Vivo* Study of Effects of Platelet-Rich Plasma on Transforming Growth Factor- β 1 and Matrix Metalloprotein 9 Expression in Traumatic Ulcers with Diabetes Mellitus

Desiana Radithia¹  Yuliana Yuliana² Yeni Puspitasari² Rossy Sismiyanti² Aulya Setyo Pratiwi²

¹Department of Oral Medicine, Faculty of Dental Medicine, Universitas Airlangga, Indonesia

²Oral Medicine Residency Program, Faculty of Dental Medicine, Universitas Airlangga, Indonesia

Address for correspondence Desiana Radithia, DDS, PhD, Department of Oral Medicine, Faculty of Dental Medicine, Universitas Airlangga, Jalan Prof. Dr. Moestopo No. 47 Surabaya 60132, Indonesia (e-mail: deisy.radithia@fkg.unair.ac.id).

Eur J Dent

Abstract

Objective Diabetes mellitus is not only characterized by alterations in the wound healing process but also during oral ulcer healing. The platelet-rich plasma (PRP) can be used to stimulate the healing process. This study was performed to analyze the effect of PRP on traumatic ulcers with diabetes mellitus in an animal model by analyzing the expression of transforming growth factor β 1 (TGF- β 1) and matrix metalloprotein 9 (MMP-9).

Materials and Methods The diabetes mellitus model was developed using streptozotocin that was administered to *Rattus norvegicus*. The traumatic ulcer model was obtained by placing a heated tip of a ball burnisher for 5 seconds on the lower mucosa labial. Then, the traumatic ulcer was treated with PRP for 3, 5, and 7 days. The expression of TGF- β 1 and MMP-9 was analyzed with indirect immunohistochemistry, and differences between each marker were analyzed with statistical analysis.

Results All animals showed clinical oral ulceration as a yellow base during the experiment. The application of PRP showed a higher level of TGF- β 1 expression than the controls for 3, 5, and 7 days ($p < 0.05$). In contrast, the MMP-9 expression was lower than the control for 5 and 7 days ($p < 0.05$).

Conclusion The PRP affected traumatic ulcers with diabetes mellitus by promoting healing through TGF- β 1 expression and suppressing the MMP-9 expression. This material can serve to develop a promising topical therapy for traumatic ulcers, especially with an underlying disease such as diabetes mellitus.

Keywords

- ▶ diabetes mellitus
- ▶ oral ulcer
- ▶ platelet-rich plasma
- ▶ TGF- β 1
- ▶ MMP-9

Introduction

Diabetes mellitus has a higher prevalence of oral mucosal disorders that has an association with chronic immunosuppression,¹ such as delayed wound healing and salivary hypofunction.² The most common mucosal wound is a traumatic

ulcer; it is characterized by damage to the epithelial tissue and underlying connective tissue³ due to mechanical injury, thermal, electrical or chemical burn.⁴ The primary healing of traumatic ulcers is similar to wound healing in the skin, which causes delayed healing in patients with diabetes mellitus.

DOI <https://doi.org/10.1055/s-0043-1764429>.
ISSN 1305-7456.

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

The topical therapy commonly used in traumatic ulcers is saline mouthwash, antiseptic mouthwash such as chlorhexidine gluconate 0.2%, nonsteroidal anti-inflammatory mouthwash such as benzydamine hydrochloride 0.15%, or steroids such as hydrocortisone sodium succinate in the form of a mucoadhesive buccal patch.⁵ All these materials were unable to interfere with the healing pathways of traumatic ulcers, especially in diabetes mellitus. The other option as therapy is to use self-material-like platelet-rich plasma (PRP). This material is considered to stimulate the intrinsic pathway of healing. The PRP contains many growth factors, which are highly important for wound healing. These growth factors promote cell migration, proliferation, and differentiation, which are essential for wound healing. After activation in the wound area, platelets will release growth factors to regulate cell migration, proliferation, and matrix deposition and reduce the risk of infection from the wound, increasing the quality of healing and tissue repair.⁶

During the healing process, the transforming growth factor β 1 (TGF- β 1) is the primary growth factor for re-epithelialization, angiogenesis, and fibroblast proliferation. In contrast, the matrix metalloproteinase 9 (MMP-9) is needed to balance the fibroblast proliferation and collagen deposition to prevent fibrosis. In diabetes mellitus patients, TGF- β 1 and MMP-9 activity alteration occurs in delayed healing.⁷

PRP preparations are available in the form of injections,⁸ powder,⁹ and gel.¹⁰ Several studies showed that the PRP injection in non-healing oral ulcers achieved complete epithelialization after 8 weeks.⁸ In the case of oral ulcers caused by oral pemphigus vulgaris, the healing process took 8 weeks to 24 months.^{11,12} Due to a lack of information about the PRP's effects on traumatic ulcer healing, especially in diabetes mellitus, the current study analyzed the PRP influence on TGF- β 1 and MMP-9 expression in traumatic ulcers and considered it a good material for traumatic ulcer therapy.

Materials and Methods

The Ethical Approval

The protocol of this study and the protocol for Animal Care and Use were approved by the Ethics Committee of health and research with registration number 776/HRECC.-FODM/XII/2019.

Experimental Materials

Thirty Wistar rats (*Rattus norvegicus*) were used that were randomly divided into six groups. Each group contained five animals. The animal adapted for 7 days with free access to food and drink. The rats were selected using criteria of 2 to 3 months, gender only male, the body weight of 150 to 250 grams, and health status (measured by the rat's liveliness, shiny hair, and glowing hair).

PRP Preparation

The PRP was obtained from 35 Wistar rats by collecting blood after general anesthesia. Then, the blood was centrifuged

two times at the speed of 4000 rpm. The plasma was collected after centrifugation.

The plasma was subjected to freeze-drying. Before this process, the plasma was frozen at -83°C for 24 hours. Then, PRP was lyophilized using a sublimation drying machine. The product was then mixed with 2% carboxymethylcellulose (CMC) in a ratio of 1:1 (w/w) to form a gel preparation.

Diabetes Mellitus Model

The diabetes mellitus model was obtained by fasting the rats for 4 hours, and then administering streptozotocin (Bio-World with CAS number 18883-66-4, USA; 50 mg/kg body weight) diluted in a citric acid buffer (pH 4.5) intraperitoneally. After the administration of streptozotocin, all animals were given 10% dextrose peroral to avoid sudden hypoglycemia.¹³ Lastly, the rats were observed for 3 days to develop diabetes mellitus; it was confirmed that fasting blood glucose resulted in ≥ 126 mg/dL glucose level.

Traumatic Ulcer Model

After the animals were confirmed to have diabetes mellitus, the traumatic ulcer model was developed using the heated tip of a 3 mm ball burnisher. Before the procedure, the animals were anesthetized by intramuscularly injecting ketamine.^{14,15}

A 3-mm diameter ball burnisher was heated for ± 45 seconds. Then, the tip was gently placed in the lower oral mucosa as deep as the diameter of the burnisher tip for 1 second. The formation of a traumatic ulcer was confirmed if the ulcer showed damaged epithelium, the middle area showed yellowed whitish color, had a well-defined wound, irregular edge, and was surrounded by a red area.^{14,15}

PRP Administration

After the traumatic ulcer developed, PRP gel was applied, while the control group was given 2% CMC only. After the treatment, the experimental animals were not given food and drink for 30 minutes so that the drug could be absorbed maximally. The PRP gel was applied once a daily for 3, 5 and 7 days. After the treatment, the animal was sacrificed by anesthesia using ketamine, and a biopsy of lower labial mucosa of the animals was performed. Next, the tissue was immersed in a 10% formalin buffer solution with 10 times the tissue volume for 1 day.¹⁶

TGF- β 1 and MMP-9 Expression

The expression of TGF- β 1 and MMP-9 was studied by indirect immunohistochemistry. The monoclonal antibodies used were anti-TGF- β 1 (TGF- β 1 3C11, Santa Cruz Biotechnology Inc., USA) and anti-MMP-9 2C3 (MMP-9 2C3, Santa Cruz Biotechnology Inc., USA). The counterstaining was done using hematoxylin-eosin. A single oral pathologist studied each expression with a Nikon E100 microscope under 1000 magnification in five different fields of view.

Statistics Analysis

The independent *t*-test was used to determine the difference in the expression of TGF- β 1 and MMP-9 between the




Fig. 1 Clinical images of traumatic ulcers on the lower labial mucosa. (A and C) Three days; (B and D) 5 days and (C and E) 7 days.

intervention and control groups on each day of observation. The significance value for all analyses was $p < 0.05$. Data were analyzed using Statistical Product and Service Solution (SPSS) version 24 (IBM SPSS Statistic 24 for Mac, New York, NY, USA).

Results

Traumatic Ulcer Observation

The traumatic ulcer was observed in all groups at 3 days, 5 days, and 7 days. A yellow-colored traumatic ulcer appeared within 3 days of both control and PRF application (► **Figs. 1A** and **1D**). After 5 days, the traumatic ulcer partly healed with a reduced yellow area (► **Figs. 1B** and **1E**). After 7 days, the traumatic ulcer appeared to be completely healed, and the mucosa looked normal (► **Fig. 1C** and **1F**).

TGF- β 1 Expression

The TGF- β 1 expression was observed in fibroblast cells (► **Fig. 2**). The application of PRF in TGF- β 1 expression at 3 days, 5 days, and 7 days was higher compared with that in the control group ($p = 0.001$; $p = 0.003$ and $p = 0.002$; respectively) (► **Fig. 3A**).

MMP-9 Expression

The MMP-9 expression was observed in fibroblast cells (► **Fig. 4**). The application of PRF in MMP-9 expression at 5 days and 7 days was lower compared with that in the control group ($p = 0.001$; $p = 0.001$, respectively). While 3 days resulted in no difference in MMP-9 expression compared with the control group ($p = 0.285$) (► **Fig. 3B**).

Discussion

Diabetes mellitus results in delayed healing due to the formation of increased advanced glycation end-products (AGEs) due to hyperglycemia. The binding process of AGE to the AGE receptor (RAGE) increased reactive oxygen species (ROS). Excessively increasing ROS exceeds the antioxidant capacity, resulting in oxidative stress. Oxidative stress




Fig. 2 Microscopic of immunohistochemical examination results showing TGF- β 1 with a light microscope with 1000x magnification.




Fig. 3 The expression marker in traumatic ulcer tissue from mucosa labial. (A) TGF-β1 (B) MMP-9 expression.

triggers the activation of nuclear factor kappa B (NF-κB), followed by an increase in the production of proinflammatory cytokines such as tumor necrosis factor α (TNF-α), interleukin 1β (IL-1β), interleukin 6 (IL-6), and MMP-9. All these cytokines shift the macrophage polarization to the M1 type. The imbalanced M1/M2 ratio imbalance results in a prolonged inflammatory phase so that the ulcer becomes chronic, and the healing process is delayed.¹⁷




Fig. 4 Microscopic immunohistochemical examination results show MMP-9 with a light microscope with 1000x magnification.

After the administration of topical PRP, improvement was noticed until 7 days when the ulcer healed maximally, as compared with the control group, which consisted of traumatic ulcers. Topical administration of PRP provides an environment of growth factors and cytokines that can stimulate and initiate the healing process, including platelet-derived growth factor (PDGF) from platelets, transforming growth factor-β (TGF-β), epithelial cell growth factor (ECGF), insulin-like growth factor (IGF), and vascular endothelial growth factor (VEGF). These growth factors will stimulate the macrophages to shift into the M2 type, thus causing the proliferation of fibroblasts and the synthesis of collagen.¹¹ The PRP contains 50 to 80% alpha granules protein (adhesive protein), one of which is TGF-β1.¹⁸ This research showed an increased level of TGF-β1, which was in line with the study from Boakye *et al.* Besides containing TGF-β, PRP increased the expression of TGF-β.¹⁹ PRP activates new platelets when it encounters tissue activators (calcium, collagen)²⁰ and releases various bioactive proteins that include growth factors, one of which is TGF-β1. Platelets contain 95% of TGF-β1 from total platelets. Administration of PRP increases the amount of TGF-β1 in macrophage cells, which in turn transcribes NF-κB in the macrophage cell nucleus, decreasing the production of proinflammatory cytokines such as TNF-α, IL-1β, IL-6, and MMP-9. TGF-β1 in wound healing regulates not only epithelialization but also inflammation, angiogenesis, and granulation tissue formation. TGF-β1 is shown to be an important player in all phases of wound healing by regulating the functions of keratinocytes, fibroblasts, endothelial cells, monocytes, and other cell types.²¹

TGF-β1 is an essential player in all phases of wound healing by regulating the functions of keratinocytes, fibroblasts, and endothelial cells. Although many growth factors also play a role in keratinocyte migration during wound healing, TGF-β1 has more roles than other growth factors. After the injury, TGF-β1 is rapidly regulated and released by keratinocytes, platelets, monocytes, macrophages, and fibroblasts. In the process of angiogenesis, TGF-β1 will increase the expression of VEGF for the formation of new blood vessels so that the supply of nutrients for wound healing is fulfilled.¹⁸ In addition, TGF-β1 is able to inhibit MMP-9, resulting the inhibition of collagen degradation.^{12,22,23}

Applying PRP significantly decreased MMP-9 expression after 5 and 7 days. In the normal ulcer healing process, MMP-9 plays a role in the inflammatory and proliferative phases.²¹ MMP-9 is a significant marker of delayed wound healing in diabetic conditions. Shin and Oh, 2010 reported that diabetic ulcers are characterized by elevated expression of MMP-9. This study was in line with a study from Shin and Oh in 2010. Shin and Oh also reported that after giving PRP, the MMP-9 expression was decreased on day 5 and day 7.²⁴ In diabetic conditions, MMP-9 was found to be overexpressed while TIMP-1 was under-secreted, The imbalance in the expression of MMP-9 and TIMP-1 inhibited tissue regeneration and revascularization.^{21,22} In contrast, PRP containing TGF-β1 indirectly reduces the MMP-9 expression. TGF-β1 regulates TIMP-1 so that the levels of TIMP-1 increase and the

overexpressed MMP-9 is inhibited by TIMP-1.²² The clinical use of PRP in traumatic ulcers can be advantageous and disadvantageous during the ulcer healing process. The advantages of PRP are that it can be easily obtained from human blood and able to support hemostasis and wound healing process.²⁵ In contrast, the disadvantages of PRP are that it is not cost-effective and its use is partially successful. Due to these reasons, the material to be selected for treatment needs to be carefully considered to get the maximum results at an effective cost.²⁵ The limitation of this research is that although hyperglycemia is one of the factors that induce diabetic wounds, there are several factors in a chronic diabetic condition that can worsen a diabetic wound. Further studies are needed to analyze the effect of PRP in other marker-related chronic diabetic conditions.

Conclusion

PRP heals traumatic ulcers in patients with diabetes mellitus by promoting the expression of TGF- β 1 and suppressing that of MMP-9. This material can be a promising topical therapy for traumatic ulcers, especially with an underlying disease such as diabetes mellitus.

Conflict of Interest

None declared.

Reference

- Al-Maskari AY, Al-Maskari MY, Al-Sudairy S. Oral manifestations and complications of diabetes mellitus. *Sultan Qaboos Univ Med J* 2011;11(02):179–186
- Mauri-Obradors E, Estrugo-Devesa A, Jané-Salas E, Viñas M, López-López J. Oral manifestations of diabetes mellitus. A systematic review. *Med Oral Patol Oral Cir Bucal* 2017;22(05):e586–e594
- Veiga N, Tiago M, Ana Sofia M, et al. Oral manifestations and diabetes. *Biomed J Sci Tech Res* 2018;7(05):6168–6171
- Mortazavi H, Safi Y, Baharvand M, Rahmani S. Diagnostic features of common oral ulcerative lesions: an updated decision tree. *Int J Dent* 2016;2016:7278925
- Lewis MAO, Wilson NHF. Oral ulceration: causes and management. *Pharm J* 2019;302(7923):1–13
- Chicharro-Alcántara D, Rubio-Zaragoza M, Damiá-Giménez E, et al. Platelet-rich plasma: new insights for cutaneous wound healing management. *J Funct Biomater* 2018;9(01):10
- Tripathi R, Tripathi K. Management of non healing oral ulcer in diabetic patient using topical application of epidermal growth factor: a case report. *SAJB* 2015;3:640–643
- Tsachiridi M, Galyfos G, Andreou A, et al. Autologous platelet-rich plasma for nonhealing ulcers: a comparative study. *Vasc Spec Int* 2019;35(01):22–27
- Shiga Y, Orita S, Kubota G, et al. Freeze-dried platelet-rich plasma accelerates bone union with adequate rigidity in posterolateral lumbar fusion surgery model in rats. *Sci Rep* 2016;6(01):36715
- Mazzucco L, Balbo V, Cattana E, Borzini P. Platelet-rich plasma and platelet gel preparation using Plateltext. *Vox Sang* 2008;94(03):202–208
- EL-Komy MH, Hassan AS, Abdel Raheem HM, et al. Platelet-rich plasma for resistant oral erosions of pemphigus vulgaris: a pilot study. *Wound Repair Regen* 2015;23(06):953–955
- Huang YY, Lin CW, Cheng NC, et al. Effect of a novel macrophage-regulating drug on wound healing in patients with diabetic foot ulcers: a randomized clinical trial. *JAMA Netw Open* 2021;4(09):e2122607–e2122607
- Furman BL. Streptozotocin-induced diabetic models in mice and rats. *Curr Protocols Pharmacol* 2015;70(01):47.1, 20
- Sari RP, Wahjuningsi E, Soewondo IK. Modulation of FGF2 after topical application of *Stichopus hermannii* gel on traumatic ulcer in Wistar rats. *Dent J* 2014;47(03):126–129
- Cavalcante GM, Sousa de Paula RJ, Souza LP, Sousa FB, Mota MRL, Alves APNN. Experimental model of traumatic ulcer in the cheek mucosa of rats. *Acta Cir Bras* 2011;26(03):227–234
- Rahman MA, Sultana N, Ayman U, et al. Alcoholic fixation over formalin fixation: A new, safer option for morphologic and molecular analysis of tissues. *Saudi J Biol Sci* 2022;29(01):175–182
- Sireesh D, Dhamodharan U, Ezhilarasi K, Vijay V, Ramkumar KM. Association of NF-E2 Related Factor 2 (Nrf2) and inflammatory cytokines in recent onset Type 2 Diabetes Mellitus. *Sci Rep* 2018;8(01):5126
- Andia I, Perez-Valle A, Del Amo C, Maffulli N. Freeze-drying of platelet-rich plasma: The quest for standardization. *Int J Mol Sci* 2020;21(18):1–20
- Boakye LA, Ross KA, Pinski JM, et al. Platelet-rich plasma increases transforming growth factor-beta1 expression at graft-host interface following autologous osteochondral transplantation in a rabbit model. *World J Orthop* 2015;6(11):961–969
- Nakajima Y, Kawase T, Kobayashi M, Okuda K, Wolff LF, Yoshie H. Bioactivity of freeze-dried platelet-rich plasma in an adsorbed form on a biodegradable polymer material. *Platelets* 2012;23(08):594–603
- Huang Y, Kyriakides TR. The role of extracellular matrix in the pathophysiology of diabetic wounds. *Matrix Biol Plus* 2020;6–7:100037
- Li L, Chen D, Wang C, Liu G, Ran X. The effect of autologous platelet-rich gel on the dynamic changes of the matrix metalloproteinase-2 and tissue inhibitor of metalloproteinase-2 expression in the diabetic chronic refractory cutaneous ulcers. *J Diabetes Res* 2015;2015:954701
- Everts PA, Knape JT, Weibrich G, et al. Platelet-rich plasma and platelet gel: a review. *J Extra Corpor Technol* 2006;38(02):174–187
- Shin HS, Oh HY. The effect of platelet-rich plasma on wounds of OLETF rats using expression on matrix metalloproteinase-2 and -9 mRNA. *Arch Plast Surg* 2012;39(08):110
- Albanese A, Licata ME, Polizzi B, Campisi G. Platelet-rich plasma (PRP) in dental and oral surgery: from the wound healing to bone regeneration. *Immun Ageing* 2013;10(01):23