

An Insect Molecular Clock Dates the Origin of the Insects and Accords with Palaeontological and Biogeographic Landmarks

Michael W. Gaunt and Michael A. Miles

Pathogen Molecular Biology and Biochemistry Unit, Department of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine

A unified understanding of >390 Myr of insect evolution requires insight into their origin. Molecular clocks are widely applied for evolutionary dating, but clocks for the class Insecta have remained elusive. We now define a robust nucleotide and amino acid mitochondrial molecular clock encompassing five insect orders, including the Blattaria (cockroaches), Orthoptera (crickets and locusts), Hemiptera (true bugs), Diptera, and Lepidoptera (butterflies and moths). Calibration of the clock using one of the earliest, most extensive fossil records for insects (the early ancestors of extant Blattaria) was congruent with all available insect fossils, with biogeographic history, with the Cambrian explosion, and with independent dating estimates from Lepidopteran families. In addition, dates obtained from both nucleotide and amino acid clocks were congruent with each other.

Of particular interest to vector biology is the early date of the emergence of triatomine bugs (99.8–93.5 MYA), coincident with the formation of the South American continent during the breakup of Gondwanaland. More generally, we reveal the insects arising from a common ancestor with the Anostraca (fairy shrimps) at around the Silurian-Ordovician boundary (434.2–421.1 MYA) coinciding with the earliest plant megafossil. We explore Tillyard's theory proposing that the terrestrial transition of the aquatic arthropod ancestor to the insects is associated with a particular plant group (early vascular plants). The major output of the study is a comprehensive series of dates for deep-branching points within insect evolution that can act as calibration points for further dating studies within insect families and genera.

Introduction

The molecular clock hypothesis is the basis of the modern molecular phylogenetic approach for dating and resolving evolutionary divergence. Major studies include estimates for the ages of mammals (Hedges et al. 1996) and birds (Cooper and Penny 1997). Nevertheless, deviations from a molecular clock, caused by rate heterogeneity between lineages, have been observed for most genetic markers (Hedges et al. 1996; Bromham, Rambaut, and Harvey 2000; Yoder and Yang 2000). However, recent approaches to molecular clock calculations provide the potential to obtain robust temporal reconstructions by assuming any number of local molecular clocks within a global phylogeny (Yoder and Yang 2000). More deterministic approaches for relaxing the molecular clock have been based on the assumption that nucleotide substitutions along the length of a branch approximate to lognormal or Poisson distributions. The assumptions underlying how these molecular clocks are relaxed differ, with one approach assuming there is a correlation of evolutionary rates among closely related lineages, usually referred to as "autocorrelation" (Sanderson 1997; Thorne, Kishino and Painter 1998) and the

second approach allowing rates to change anywhere on a phylogenetic tree (Huelsenbeck, Larget, and Swofford 2000). An earlier approach to molecular dating involved the selective removal of lineages that deviated from a molecular clock "backbone," identified using a relative rates test (Li and Tanimura 1987), also known as the two-cluster test (Takezaki, Rzhetsky, and Nei 1995).

Although insects demonstrate the greatest biodiversity in the animal kingdom, no molecular clock has previously been identified either for the class as a whole or for any order therein. Moreover, few robust examples of rate uniformity have been observed even at the family level. In addition, unlike studies of other animals, few molecular dating methods for insects have been calibrated using the available fossil record; calibrations have predominantly assumed population isolation following tectonic movements (Russo, Takezaki, and Nei 1995; Pellmyr and Leebens-Mack 1999). The use of geographic calibration points for molecular clocks could be strongly criticized because by taking the most prominent example, the breakup of Gondwanaland, few molecular or palaeontological studies (e.g., Hedges et al. 1996) substantiate that this is a major driving force for global speciation; therefore, assuming that continental breakup is the single source of diversity would appear to be open to error.

Methods

Complete amino acid and nucleotide data sets were obtained from GenBank, aligned using Clustal X, and edited by hand (Thompson et al. 1997). Selection of genetic loci was based on three factors: the most commonly sequenced loci throughout the Insecta, that is, 16S, 18S, cytochrome *b* (*cob*), cytochrome oxidase I (*cox1*), and elongation factor 1 α (EF-1 α) (Caterino, Cho, and Sperling 2000); the availability of nucleotide

Abbreviations: *atp6* and *atp8*, adenosine triphosphate synthases 6 and 8; bp, base pair; CI, confidence interval; *cob*, cytochrome *b*; *cox1*, *cox2*, and *cox3*, cytochrome oxidases I, II, and III; EF-1 α , elongation factor-1 α ; ML, maximum likelihood; *nad1*, *nad2*, *nad3*, *nad4*, *nad4L*, *nad5*, and *nad6*, nicotinamide adenine dinucleotide dehydrogenases 1–6; POLL II, polymerase II.

Key words: insecta, insect evolution, molecular clock, maximum likelihood, triatomine bugs.

Address for correspondence and reprints: Michael W. Gaunt, Pathogen Molecular Biology and Biochemistry Unit, Department of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT. E-mail: michael.gaunt@lshtm.ac.uk.

Mol. Biol. Evol. 19(5):748–761. 2002

© 2002 by the Society for Molecular Biology and Evolution. ISSN: 0737-4038

sequence data from the insect orders *Blattaria* or *Odonata* which both offer the earliest and most extensive fossil records for extant insects to allow for calibration of the clock; existing sequence data for the triatomine bugs; and, finally, availability of sequence data for Brachiopoda as the closest ancestor to the insects. Partial length sequence data were used in each instance, comprising 477, 350, and 133 amino acid residues for *cox1*, EF-1 α , and *cob* data sets, respectively, 1,396 nucleotide base pairs (bp) for 18S, and 258 bp for 16S. All alignments are available on request. 16S and 18S alignments comprised sequence data for all available hematophagous insect vectors. For protein loci two examples of each species were included, and for *cox1* data particular attention was paid to the extensive molecular dating available for the two Lepidopteran families, Proxidae and Incurvidae, to enable a direct comparison with previous date estimates (Pellmyr and Leebens-Mack 1999).

cox1 Phylogeny

An initial assessment of the above loci suggested that *cox1* provided the most suitable locus for a molecular clock study (see *Molecular Clock* section later for details of this initial assessment). The *cox1* data set was then supplemented with a new *cox1* sequence from triatomine bugs of the tribes Rhodniini (*Rhodnius pictipes* [GenBank accession number AF449136], *R. robustus* [AF449137], and *R. prolixus* [AF449138]) and Triatomini (*Triatoma maculata* [AF449139], *Eratyrum mucronatus* [AF449140], and *Panstrongylus herreri* [AF449141]). Amplicons were obtained using a two-round, or "nested," PCR, initially using primers 5' TGT ATT TTM TRT TCG GGG CYT GA 3' and 5' TCG TGG AAG AAG ATA AGT TGT T 3' and then using primers 5' GAG CTG GAA TAA TAG GAA CAT C 3' and 5' TGC ATT AAT CTG CCA YAT TA 3'. The second-round primers had the standard M13 promoter site sequences linked to their 5' terminus (5' TGT AAA ACG ACG GCC AGT 3' and 5' CAG GAA ACA GCT ATG ACC 3'). The second-round amplicons were custom sequenced using M13 primers to ensure high-quality sequence data (MWG-Biotech, U.K.). All phylogenies were outgrouped using all available sequences from the Crustacea.

The robustness of the data sets was examined for deviations in nucleotide or amino acid base composition between taxa because nucleotide or amino acid base homogeneity is a prerequisite for the maximum likelihood (ML) mutation models used throughout this study. We felt this criterion to be particularly important to observe for molecular dating purposes. Mitochondrial *Apis mellifera ligustica* sequence (NC_001566) was not included in the analysis because of a low level of amino acid homogeneity ($P = 0.1$ and >0.99 in all other cases) using the ML quartet puzzling software Puzzle (Strimmer and von Haeseler 1996). For nucleotide data sets, first- and second-nucleotide codon positions were used for evolutionary reconstructions because of the significant deviation of the third base from nucleotide base homogeneity.

The ability of the *cox1* locus in comparison with the other mitochondrial protein-encoding genes to demonstrate a robust molecular clock was reanalyzed for representative insects for which a complete mitochondrial genome had previously been deposited in GenBank. Each mitochondrial protein-encoding gene (adenosine triphosphate synthase 6 and 8 [*atp6* and *atp8*] combined, *cox1*, *cox2*, *cox3*, *cob*, nicotinamide adenine dinucleotide dehydrogenases 1–3, 5, 6 [*nad1*, *nad2*, *nad3*, *nad5*, *nad6*], with *nad4* and *nad4L* combined) was separately aligned and assessed for a molecular clock using amino acid ML quartet puzzling (Puzzle) and second-codon-position nucleotide ML, using a seven-parameter model. The insects used represented the Diptera (*Anopheles quadrimaculatus* [NC_000875]; *A. gambiae* [NC_002084]; *Cochliomyia hominivorax* [AF260826]; *Ceratitis capitata* [NC_000857]; *Drosophila yakuba* [NC_001322]; *D. melanogaster* [NC_001709]), the Lepidoptera (*Bombyx mori* [NC_002355]), and the Hemiptera (*T. dimidiata* [AF301594]), the Orthoptera (*Locusta migratoria* [X80245]), and the phylogeny was rooted using the Brachiopoda (*Daphnia pulex* [NC_000844]; *Artemia franciscana* [X69067]).

A prerequisite of the study was to reconstruct constrained molecular clock ML phylogenies using nucleotide and amino acid data and compare them with the unconstrained models. The basic model of nucleotide evolution was determined using MODELTEST (Posada and Crandall 1998). Seven ML model parameters, including a four-category discrete gamma distribution (one parameter, Γ), the general time-reversible model of nucleotide substitution (six parameters, GTR) and invariant rate parameter (one parameter, PINVAR), were estimated during a heuristic tree search for a reconstruction limit of 1 using PAUP*4.0b (Swofford 2000). The ML amino acid phylogeny (described later) was then used as a backbone to resolve the nucleotide ML phylogeny because nucleotide data could not adequately resolve the interorder topology, and the seven parameters were reestimated. All parameter estimates were incorporated into a full ML heuristic search for nucleotide data (PAUP*4.0b). All parameter estimates are available on request.

ML amino acid phylogenies were constructed by a heuristic search using the JTT (Jones, Taylor, and Thornton 1992) and mtREV24 (Adachi and Hasegawa 1996a) models of amino acid substitution where appropriate by PROTML in the MOLPHY package (Adachi and Hasegawa 1996b). Nucleotide and amino acid parsimony trees were calculated using a heuristic search with Tree bisection-reconnection (TBR) branch swapping for 1,000 random sequence additions (PAUP*4.0b). Minimum distance phylogenies were built using an ML model for nucleotide data with the parameters defined by MODELTEST and using p -distances for amino acid data (PAUP* 4.0). The *cox1* topology was initially assessed using parsimony and minimum distance nonparametric bootstrapping. Further topological differences between the *cox1* ML, parsimony, and minimum distance nucleotide and amino acid phylogenies were assessed by the Shimodaira-Hasegawa test (Shimodaira and Hasegawa

1999), using the full optimization (100 replications) or the RELL approximation (1,000 replications) to the ML bootstrap with the seven-parameter ML model for nucleotide data previously described (Γ , GTR and PINVAR).

Tenebrio molitor cox1 sequence (X88966) was omitted because of a topological discrepancy with insect morphologically-based insect phylogenies. For morphologically-based insect phylogenies, *Tenebrio* sp. (Coleoptera) is an out-group to Diptera and Lepidoptera (Whiting et al. 1997; Wheeler et al. 2001), whereas in the amino acid phylogeny of *cox1*, Diptera is an out-group to Coleoptera and Lepidoptera; therefore, the phylogenetic reconstructions were repeated. The suitability of EF-1 α analysis for dating within Insecta was questionable when a gene duplication event was observed from the quartet puzzling ML phylogeny, resulting in EF-1 α F1 and EF-1 α F2 of *Drosophila melanogaster* disrupting the Brachycera suborder and possibly causing splits in the dipteran monophyly (data not shown). This observation is supported by previous studies on EF-1 α (Danforth and Ji 1998).

Molecular Clock

Molecular clock tests were conducted both before any loci were sequenced and after the sequence data were incorporated to ensure that a molecular clock was still observed. Initial screening of the 16S, 18S, *cob*, *cox1*, and EF-1 α loci for mutation rate consistency between different lineages was performed for nucleotide and for amino acid data using the likelihood ratio test $2\Delta L = n - 2$, where n is the number of taxa and L the negative log likelihood score; if a constant rate of evolution occurs between lineages, then sequence data represented in an ML additive tree, where branch length differences represent sequence divergence, should not be significantly different from an ML ultrametric tree, where the tips of the tree are equidistant from the root. All molecular clocks for 16S, 18S, *cob*, and EF-1 α were significantly rejected, even for small numbers of taxa, using amino acid ML quartet puzzling (Puzzle; Strimmer and von Haeseler 1996) with JTT (Jones, Taylor, and Thornton 1992) or mtREV24 (Adachi and Hasegawa 1996) mutation models and for second-codon positions for nucleotide data, using a seven-parameter model (discussed elsewhere; PAUP*4.0b). However, *cox1* locus robust molecular clocks were obtained for some, but not all, taxa using ML quartet puzzling for amino acid data (not shown) and, in particular, for second-codon positions for nucleotide data using a seven-parameter ML model (discussed later in *Methods*), indicating the suitability of this locus for dating studies.

The calibration point for the *cox1* nucleotide and amino acid molecular clock phylogenies was set at the Blattaria (cockroaches)-Orthoptera (crickets and locusts) divergence. Early ancestors of extant Blattaria, initially called Pinnule Insects (Henning 1981), have more recently been reclassified as members of this order (Jarzembowski 1994) which represents one of the earliest, most extensive fossil records of all extant insects (La-

bandeira and Sepkoski 1993; Ross and Jarzembowski 1993). Furthermore, the onset of their radiation coincided with the radiation seen in the extensive fossil record of an extinct member of the neopteran clade, the Prothoptera, which are more closely related to Orthoptera than to Blattaria. It is noteworthy that Blattaria are one of three closely related orders that include Mantoidea (mantids) and Isoptera (termites), and at least Isoptera are believed to have evolved from wood-eating Blattaria (Lo et al. 2000).

The *cox1* phylogeny of nucleotide sequence, including the triatomine bug sequences, was resubjected to a likelihood ratio test for a global molecular clock, i.e., assuming the same rate for all lineages, using second-codon-position nucleotide data under the seven-parameter model using a four-category discrete Γ distribution, GTR, PINVAR (PAUP 4.0*b). Model parameters for the molecular clock test were estimated for the unconstrained phylogeny and used for the global clock.

The *cox1* phylogeny of amino acid sequence was tested for a molecular clock with a 12-category discrete gamma distribution using PAML, written by Z. Yang (Yoder and Yang 2000). This statistical method allows for variable evolutionary rates among lineages to address complex evolutionary models, where some branches have the same rate, a "local molecular clock," whereas different parts of the tree have different rates. The approach is attractive because it allows the user to specify any given evolutionary model and, in addition, allows a likelihood ratio test to assess the robustness of the clock. One feature of this approach is that by assigning discrete rate categories to given lineages, it often results in abrupt changes in evolutionary rate between species boundaries.

Amino acid ML models for the *cox1* phylogeny assuming a global molecular clock or a local molecular clock for each insect order and different rates for lineages between each order were significantly rejected using a likelihood ratio test $2\Delta L = n - k - 2$ parameters, where k is the number of evolutionary rates in the model, and n and L are as previously described (PAML; Yoder and Yang 2000). These models and all subsequent models discarded ambiguous amino acid sites. The local molecular clock method (PAML) gives the user the potential to estimate the local rate of molecular evolution for each branch in the tree. Given this potential, we adopted a philosophy similar to that of Sanderson (1997) by calculating the local rate of molecular evolution for separate branches, then minimized the rate difference between branches, and incorporated a method of smoothing the differences between rates. However, we sought to avoid reliance on any one assumption of mutation rate heterogeneity. To simplify the calculation, we assumed that autocorrelation of rates occurred within each insect genus, i.e., a local molecular clock was assigned to each genus, but separate mutation rates could occur between genera, with the exception of the separate rates that were assigned for *R. pictipes* and *T. vitticeps* because of their very short branch lengths; in addition, mutation rates were calculated for all intergenera branches. This resulted in a molecular clock phylogeny




FIG. 1.—The ranked distribution of mutation rates from the 85-rate category model of the amino acid *cox1* phylogeny described in *Methods*, which assumes local molecular clocks for most insect genera to define, and examples of using rate 0.2, 0.3, 0.4, and 0.5 interval bins to reduce the total number of rates in the model. In this instance, the interval bins are based on boundary values of 0⁺ to 0.1 as defined in *Methods*. * denotes *Artemia* reference lineage (rate = 1).

with 85 ML rates, having almost as many mutation rates as branches, with a maximum and minimum mutation rate of 0.093 and a 3.59-fold difference in substitutions per site relative to *A. franciscana*, the reference lineage (rate = 1; fig. 1). To reduce the number of rate parameters, we defined a series of bin ranges, or “rate categories,” for the frequency distribution of ranked mutation rates of the 85-ML parameter model, effectively pooling lineages that showed similar mutation rates regardless of where they occurred on the phylogeny. There was no objective way of defining any given bin range, and categorizing what appeared to be continuous data (fig. 1) could result in problems; therefore, in order to smooth the differences between mutation rates a large number of different bin ranges was defined, resulting in a series of mutation rate histograms for the *cox1* phylogeny. All histogram bin ranges were arbitrarily based on two different boundary values, 0⁺ to 0.1 and 0.05⁺ to 0.15 relative to the mutation rate of *A. franciscana*, where ⁺ is an incremental increase on the stated value. Four bin ranges comprising 0.2, 0.3, 0.4, and 0.5 intervals relative to the *A. franciscana* mutation rate were constructed (fig. 1), in addition to a 0-rate category, defining 8–16 different mutation rates for the phylogeny. To further minimize rate interval “edge effects,” each combination of 0.1-unit boundary values within each rate interval was described; for example, a 0.5-unit interval size has five possible 0.1-unit boundary value combinations and therefore five different bin ranges, whereas a 0.2 interval size had two different bin ranges. In total, 28 different models describing the evolutionary rates of the *cox1* phylogeny were calculated (PAML); 14 rate models were derived from different bin intervals




FIG. 2.—Hypothetical example of a nonautocorrelation model (fig. 1) where identical mutation rates occur at different parts of the phylogeny (rate in branches A and B = rate in branches C and D) and the construction of two different groups of models assuming autocorrelation where the mutation rate at different parts of the phylogeny is not assumed to be identical (rate in clade 1 \neq rate in clade 2). Approach 1: mutation rate in branches A and B \neq that in branches C and D, and the model is recalculated. Approach 2: mutation rate in branches A, B, and E is assumed to be identical, as is the mutation rate in branches C, D, and F, but the rate in branches A, B, E \neq rate in branches C, D, E, and the model is recalculated.

(5 + 4 + 3 + 2) based on two different boundary values. Models that did not support the unconstrained model, by a collapse in the branch length of the ancestral lineage to the Branchipoda and Insecta presumably reflecting instability in the malacostracan root, were discarded (17 models were retained). In summary, 17 date estimates were obtained for each node based on the hypothesis that identical rates occur at any part of the phylogeny, although this hypothesis may not represent the true behavior of local clocks within the phylogeny. We therefore employed a second assumption that autocorrelation of mutation rates occurs between insect genera, families, or suborders (fig. 2). Two approaches to autocorrelation were adopted using the 17 models of rate estimates derived from the previous hypothesis; when this hypothesis described two or more clades at different parts of the tree as having identical mutation rates, the assumption of rate identity between unrelated clades was removed, and the model was recalculated (approach 1; 17 models), and in addition, when two “sister” branches showed identical mutation rates, then the mutation rate of the “parental” branch was also assumed to be identical regardless of the value previously assigned to it, and the model was recalculated (approach 2; 17 models).

A weakness of the approach described here, in common with that of Sanderson (1997), is that no external criteria are used for defining which lineages are autocorrelated, such as the comparative reproductive rate between species. Moreover, the rate hypothesis is derived from the data and is also tested against the same data, which is effectively the situation with the “tree pruning” methods and is only relevant if it is compared with an approach where this assumption is removed or minimized, i.e., compared with the nucleotide clock. Ideally, the nucleotide data set should be bootstrapped to obtain a better estimate of the confidence limits, but in conjunction with the models described so far this

would result in an unrealistically large number of models.

Root Stability

Concerns for the stability of the malacostracan root were further investigated for the autocorrelation-based approaches (fig. 2), because of a further collapse in the branch length of the ancestral lineage to the Branchiopoda and Insecta in 10 of the 34 models. The collapsed branch length occurred in models where each of the five branches in the malacostracan root was assigned an independent mutation rate. The remaining models showed identical rates between the parental branch to the *Penaeus monodon* and *P. notialis* sister groups and the parental branch to the *Pagurus longicarpus* and *Penaeus* branches. To test root stability, a final group of models was made assigning independent rates for the *Penaeus* and *Pagurus-Penaeus* parental branches (nine models for approach 1 and nine models for approach 2; fig. 2). A direct comparison of the difference in date estimates could then be made for each model, with and without the extra rate category in the root, and the stability of the root assessed. The maximum difference between date estimates from models with the extra rate category in the root compared with those without it for all 18 models was less than 5% for all insect nodes (branching points), but the nodes within the Brachycera suborder (i.e., the *Drosophila-Ceratitis* split) showed a difference of up to 17%. The maximum difference in date estimates for basal lineages was 7% for the nodes of Branchiopoda, 10.5% for the Malacostraca-Branchiopoda ancestral node, and over 70% variation for the nodes within Malacostraca. Although the root was unstable, and the date estimates for Malacostraca were not considered further, this did not greatly affect the divergence of most lineages, except for the Brachycera suborder which was unusually unstable.

To examine the robustness of date estimates to model assumptions, we obtained the 95% confidence interval (CI) using median bootstrapping for 1,000 replications with the STATA package (<http://www.stata.com>). We did not attempt to determine which is the correct model but simply bootstrapped all dates for each node obtained from the nonautocorrelated- and autocorrelated-based approaches (3×17 models) and the additional models obtained from analyzing the stability of the malacostracan root (2×9 models), giving 69 models in total. Median bootstrapping was used to minimize outlier effects and a priori assumptions of the CI estimation. In addition, this approach sought to minimize a priori assumptions of rate heterogeneity between insect lineages and rate changes only at speciation boundaries (Huelsenbeck, Larget, and Swofford 2000).

Stratigraphic Dating

Permian stratigraphic ranges were taken from a single source (Jin et al. 1997); all other stratigraphic ranges used were taken from Gradstein and Ogg (1996). This geological time scale incorporates several improvements over previous studies (Harland et al. 1989, p. 265) which

includes more accurate Mesozoic chronology (Triassic-Cretaceous) using radiometric dating rather than interpolations from marine magnetic anomaly profiles (Gradstein et al. 1995) and relatively precise isotope datings for Paleozoic chronology (pre-Triassic) (Roberts, Claoue-Long, and Jones 1995; Tucker and McKerrow 1995).

Data deposition

The new nucleotide sequence data described here are available at GenBank under accession numbers AF449136–AF449141. All amino acid and nucleotide alignments and ML parameter estimates are available on request.

Results

As a route to estimating the evolutionary dates for deep divergences within insect evolution, we used near full-length *cox1* nucleotide and amino acid data using ML global and local molecular clocks, respectively. Several lines of evidence suggested that *cox1* provided the most suitable locus for this study. Firstly, it was the only locus that recovered global molecular clocks for some, but not all, of the taxa examined particularly for nucleotide data (fig. 3). Secondly, in comparison with all other protein-encoding mitochondrial genes, using representatives of four of the five insect orders (described in *Methods*) for which there was a complete mitochondrial genome sequence available and including the out-groups of Branchiopoda, *cox1* provided the smallest difference between the global molecular clock and the unconstrained models for amino acid data without Branchiopoda as the root ($2\Delta L = 15.0$ at $df = 7$; $P = 0.036$) and nucleotide data with Branchiopoda as the root ($2\Delta L = 11.1$ at $df = 9$; $P = 0.27$). In contrast, all other protein-encoding mitochondrial genes produced a substantial deviation between the unconstrained and clock models ($P < 0.003$ amino acid and $P < 0.05$ nucleotide data), with the exception of *cox3* nucleotide data ($2\Delta L = 15.2$; $P = 0.084$ at $df = 9$). Furthermore, *cox1*, *cox2*, *cox3*, *nad1*, and *nad4* were the only loci that did not show significant deviation of the amino acid composition between lineages using the mtREV24 mutation model, i.e., the model derived from the empirical mitochondrial sequence.

Robustness of *cox1* Phylogeny

Unequivocal congruence in branching order was obtained within each order for amino acid phylogenies by both parsimony and ML methods, with the exception of the placement of *C. capitata* and small differences within *Triatoma* spp., and provided the basic phylogeny. Moreover, regardless of the method used, either ML or parsimony using amino acids or nucleotides, very similar topologies were recovered within each order. Nucleotide data using first- and second-codon-position nucleotides, however, did not recover the interorder topology (*Methods*). All family and superfamily relationships within each order were strongly supported using neigh-




Fig. 3.—ML global molecular clock phylogeny using second-codon-position *coxI* nucleotides of five insect orders and two classes of Crustacea, calibrated using the Blattaria/Orthoptera divergence (0.022% nucleotide substitutions per million year). Numbers within brackets denote dates obtained when the fastest and slowest lineages within Hemiptera and Lepidoptera were removed.

bor-joining bootstrap analysis for nucleotide data sets, with the exception of taxonomic relationships within Lepidoptera (Appendix 1). Bootstrap analysis of *coxI* amino acid data sets generally resulted in lower bootstrap values; however, unlike nucleotide data, amino acid data recovered the correct interorder topology (Appendix 2). The topology describing Proxidiidae, Incurvidae, Ditrysia, Adelidae, and Helozelidae within Lepidoptera in figures 3 and 4 was recovered for distance, parsimony, and ML tree-building methods using both amino acid and nucleotide data sets. In addition, the robustness of the Adelidae and Helozelidae outgroup within the Lepidoptera was supported by the Shimodaira-Hasegawa test using full optimization to the ML bootstrap for nucleotide data, against the alternative topology of Ditryia forming an out-group to this order ($\Delta L = 11.3$; $P = 0.04$; Appendix 1). Alternative branching orders were obtained by examining topological differences between amino acid and nucleotide data sets among the ML, parsimony, and distance methods and involved the

placement of *C. capitata* within the dipteran suborder Brachycera and lineages within *Triatoma* spp. (although topological comparisons were not exhaustive in this instance) and *Drosophila* spp. using the Shimodaira-Hasegawa test with the RELL approximation to the ML bootstrap for nucleotide data as a quicker alternative to the full-optimization test. The phylogeny was rooted using Malacostraca as previously described (Friedrich and Tautz 1995; Regier and Shultz 1997, 1998; Boore, Lacro, and Brown 1998).

Nucleotide Molecular Clock for *coxI*

Figure 3 describes the molecular clock for *coxI* nucleotide data for second-codon positions for all insect taxa and including Branchiopoda. Although the mutation rate heterogeneity between lineages is significant using a likelihood ratio test, the vast majority of lineages do obey a molecular clock. For instance, if the lineages that demonstrate the fastest or slowest evolutionary rates




FIG. 4.—ML local molecular clock phylogeny using near-full-length *coxI* amino acids of five insect orders and two classes of Crustacea. The model here represents the median value of the average date per node for each of the 69 models (104.5 Myr; average rate = 0.048% amino acid substitutions per million year) and is presented for diagrammatic purposes only. CI (95%) of date estimates were obtained by median bootstrapping (small triangles). Numbers in brackets denote minimum date estimates from a previous study obtained by the two-cluster test using nucleotide data (Pellmyr and Leebens-Mack 1999). Large triangles denote stratigraphic boundaries (Gradstein and Ogg 1996) and are drawn to scale, where Ca, Cambrian; O, Ordovician; S, Silurian; D, Devonian; C₁, Early Carboniferous; C₂, Late Carboniferous.

in Hemiptera and Lepidoptera are excluded under a seven-parameter ML nucleotide model (Γ , GTR and PINVAR; *Methods*), namely, *R. pictipes*, *P. megistus*, *Colias eurytheme*, and *Argyrotaenia citrana*, along with the fastest or slowest evolutionary rates in Incurvidae, *Incurvaria mascullella* and Proxididae, *Prodoxus y-inversus*, a robust molecular clock is observed, although there is a collapse in the root ($2\Delta L = 65.9$ at $df = 55$; $P = 0.15$). The purpose of this analysis was not to perform a “relative rates” test (e.g., the two-cluster test; Russo, Takezaki, and Nei 1995) but primarily to demonstrate that most lineages conform to a molecular clock, and this provides an indication that the locus is suitable for further analysis to account for rate heterogeneity between lineages. Moreover, not even first- and second-codon positions were capable of recovering the phylogeny and raised doubts about an exclusive reliance on second-codon positions for evolutionary dating. Figure 4 presents an amino acid molecular clock phylogeny, accounting for rate heterogeneity between lineages. The date estimates from this analysis arise from 69 models resulting from the assumptions of insect amino acid mutation, including the models that examined root stability, using near-full-length *coxI* sequence for five orders of

insect with all the available crustacean *coxI* sequences (figs. 1 and 2; see *Methods*).

Amino Acid Molecular Clock for *coxI*

Each of the 69 local clock models was compared with the unconstrained model. All amino acid clock models demonstrated a robust molecular clock using a likelihood ratio test described previously ($P > 0.05$ for all models [minimum $2\Delta L = 16.4$ at $df = 9$], $P > 0.225$ for 96% models [$2\Delta L = 12.89$ at $df = 10$], and $P > 0.7$ for 62% models [$2\Delta L = 15.32$ at $df = 19$], where $L = \text{ML log likelihood value}$).

The nucleotide data were primarily used to assess the ability of a locus to form a molecular clock. However, a striking congruence in date estimation was observed between the *coxI* nucleotide and amino acid data sets for all deep nodes in the five orders of Insecta. Thus, the *Daphnia pulex* and *Artemia franciscana* divergence is 498.0–485.4 and 434.2–421.1 MYA, respectively, for the amino acid clock and 420.1/423.9 MYA for the nucleotide clock, where 420.1 MYA is the date from the global nucleotide clock, and 423.9 MYA is the global clock with lineages removed; the nucleotide clock gives

single dates because of a collapse in the root. The divergence between the Blattaria/Orthoptera lineage and Hemiptera is 382.9–373.0 and 375.6–368.2 MYA for the amino acid clock, respectively, and 351.1/351.7 MYA for the nucleotide clock, because of a collapse in the bifurcating topology. Furthermore, congruence is also observed for the divergence between the dipteran suborders Nematocera containing the *Anopheles* lineages and Brachycera containing the *Drosophila* lineages (282.8–247.7 MYA amino acid clock and 259.9/261.1 MYA nucleotide clock), *Triatomini* and *Rhodniini* divergence within the Hemiptera (99.8–93.5 MYA amino acid clock and 112.7/96.3 MYA nucleotide clock), *Chorthippus parallelus* and *L. migratoria* within the Orthoptera (95.8–88.5 MYA amino acid clock and 97.6/99.4 MYA nucleotide clock) as well as for the deepest split in Lepidoptera (255.7–240.4 MYA amino acid clock and 231.6/233.1 MYA nucleotide clock). Date estimates for other nodes within the Lepidoptera were also generally congruent between the amino acid and nucleotide clocks, although for the family Proxidiidae the nucleotide clock gave dates no more than 10% older than did the amino acid clock. The major exception was the deepest nodes within the Brachycera suborder that produced substantially different dates between the amino acid clock (165.9–129.0 MYA) and the nucleotide clock (65.9/65.1 MYA); the dates of the amino acid clock are congruent with the fossil record (table 1). It is worth noting that the dates of the nodes within the Brachycera suborder were affected by instability in the malacostracan root (*Methods*). However, the second deepest node within the Brachycera root, the *Scahophaga stercoraria*-*Drosophila* split, is more congruent between the amino acid (78.7–66.4 MYA) and nucleotide clocks (60.2/59.6 MYA).

Robustness of Insect Dating Estimates

The evolutionary dates obtained in this study provide robust date estimates when compared with the earliest dates for the stratigraphic ranges of all available insect fossils known to us and for the latest date of the Cambrian explosion (table 1). In addition to the fossil record, relevant divergence dates are also robust against proposed biogeographic events affecting speciation in the phylogeny (table 1). In particular, the continental breakup of Gondwanaland (105–95 MYA) provides the earliest date for the isolation of triatomines in South America (95.5–93.6 MYA) and the latest date for the divergence between the globally distributed Lepidoptera Proxidiidae and Incurvariidae (123.1–116.1 MYA). Moreover, the breakup of Gondwanaland was used in a previous study to provide a minimum calibration point for the divergence between moths of the families Proxidiidae and Incurvariidae (Pellmyr and Leebens-Mack 1999), which enabled the minimum dates of nodes within Proxidiidae to be estimated using the two-cluster test (fig. 4, numbers in brackets). Excluding a single date obtained by the two-cluster test because it was incongruent with the chronology for all other phylogenies (Pellmyr and Leebens-Mack 1999), no significant dif-

ference using the chi-square test (χ^2) is observed between the dates for Proxidiidae derived using either the two-cluster test calibrated from the breakup of Gondwanaland (expected result) or the minimum bootstrap values obtained from the ML insect local clocks calibrated from the divergence of Blattaria (observed results) (χ^2 : $P = 0.09$ at 10 df values given in fig. 4). This result is strengthened by removing the two-cluster–test calibration point (χ^2 : $P = 0.23$ at df = 9). However, although the dates in this study fit with those of Pellmyr and Leebens-Mack (1999), it must be noted that generally biogeographic calibration points for molecular clocks are subject to error. In summary, virtually all the evidence of the date estimates obtained here converges to suggest a robust basis for reconstructing over 400 Myr of insect evolution. The temporal congruence verifies the rationale behind using Blattaria fossils as the calibration point of the clock.

The molecular clock phylogenies describe the aquatic-terrestrial transition of the insect-*Artemia* (Anostraca, fairy shrimp) ancestor during the early/late Silurian (428.0–419.0 MYA). The date coincides with the earliest plant megafossils, notably *Cooksonia cambrensis* (423.0–419.0 MYA) (Edwards, Feehan, and Smith 1983). The radiation of the deepest insect order lineages under investigation, notably the Blattaria/Orthoptera lineage, occurred from the middle to late Devonian (approx. Eifelian-Famennian; 380.0–354.0 MYA), or perhaps very early Carboniferous, and provides the latest date for the evolution of insect flight. The date obtained here for the divergence of the Hemiptera lineage also occurred in the middle to late Devonian and denotes the divergence between hemimetabolous and holometabolous insects (Whiting et al. 1997).

Discussion

There are significant weaknesses in the insect fossil record for Lepidoptera and for fossil insects in general, particularly before 323 MYA (i.e., for the first 70–60 Myr of insect evolution). Insect palaeontology depends on assigning fossils to extant taxa usually on the basis of wing characters (Shear and Kukalova-Peck 1990). Consequently, no fossil has been identified that could represent a “missing link” between insect orders, although more recent ancestors of extant insects are seldom classified with certainty. In addition, insect fossilization occurs almost exclusively in water-deposited sediments (Shear and Kukalova-Peck 1990).

Molecular clock approaches enable a temporal reconstruction of ancestral nodes regardless of palaeontological difficulties (Hedges et al. 1996; Cooper and Penny 1997), and this study provides a vital tool to complement the insect fossil record. Most previous molecular dating studies in insects have almost exclusively focused on one gene or two adjacent mitochondrial loci (Brown et al. 1994; Takezaki, Rzhetsky, and Nei 1995; Essegir et al. 1997; Foley et al. 1998; Pellmyr and Leebens-Mack 1999). The *cox1* locus used in this study provided the closest example to a global molecular clock gene against all the other candidate loci examined both

Table 1
Temporal Robustness of the *coxI* Local Molecular Clock Against the Earliest and Latest Dates of Relevant Stratigraphic, Biogeographic, and Fossil Records

Node	Earliest Limits: Oldest Fossils (F) or Biogeography (MYA)	Latest Limits and Significant Events (MYA)	Reconstructed Divergence from this Study (MYA)	References
Triatominae: Rhodniini (palm dwelling)–Triatomini		Biogeographic isolation of family: 105–95. Event: first palms (F) 83.5–71.3; 85.8–83.5?	99.8–93.5; 112.7/96.3 ^a	Palm, megafossils; Collinson, Boulter, and Holmes 1993
Orthoptera: <i>Locusta</i> sp.– <i>Chorthippus</i> sp.	Acrididae (F): 150.7–144.2		95.8–88.5; 97.6/96.3 ^a	Labandeira 1994
Muscomorpha	Drosophilidae (F): 37.0–33.7	Basal muscomorphan fossils below	78.7–66.4	Evenhuis 1994
	Calliphoridae (F): ^b 83.5–71.3		60.2/59.6 ^a	McAlpine 1970
Basal Muscomorpha	Platyezidae (F): 93.5–89.0	Basal brachyceran fossils below	165.9–129.0; 65.9/65.1	Grimaldi, Shedrinsky, and Wampler 2000
Basal Brachycera	Alinkidae (F): 227.4–220.7		282.8–247.7; 259.9/261.1 ^a	Evenhuis 1994
Basal Nematocera	e.g., Empididae (F): 241.7–234.3		282.8–247.7; 259.9/261.1 ^a	Shcherbakov, Lukashovich, and Blagoderov 1995
Basal Diptera	Protodiptera, Permotanyderidae, and Permotipulidae (F): 256.0–253.0		282.8–247.7; 259.9/261.1 ^a	Evenhuis 1994
Incurvidae–Prodoxidae	Biogeographic separation of later family; Cecidosidae: 105–95		123.1–116.1; 121.5/141.7 ^a	Pellmyr and Leebens-Mack 1999
Prodoxidae	Estimated divergence of 10 deep nodes		No difference; <i>P</i> = 0.09 (10 nodes); <i>P</i> = 0.23 (9 nodes)	Pellmyr and Leebens-Mack 1999
<i>Papilio</i> sp.– <i>Pachiopta</i> sp.	Papilionidae (F): 49.0–41.3		100.2–94.1; 89.1/82.5 ^a	Wehr 1998
<i>Manduca</i> sp.– <i>Feltia</i> sp.	Noctuidae (F): 49.0–41.3		113.4–98.2; 57.1/60.7 ^a	Wehr 1998
Basal Incurvariodea	Undescribed Incurvarioid (F): 132.0–127.0		123.1–116.1; 121.5/141.7 ^a	Whalley 1986
Basal Ditrysia	Gracillariidae (F): 99.9–93.5		174.4–166.8; 157.6/191.9 ^a	Labandeira 1994
Basal Lepidoptera–Diptera	Mecoptera (F): 281.5–290		351.4–338.0; 333.1/339.1 ^a	Evenhuis 1994
Pterygota (winged insects)		Hexapoda or Insecta fossils as above	387.7–372.8; 351.1/351.7 ^a	
Insecta–Anostraca	Hexapoda or Insecta (F): 390.4–401, 390–392, 376–379	Event: first land plant fossils: 428	434.2–421.1; 420.1/423.9 ^a	Whalley and Jarzembowski 1981; Shear et al. 1984; Labandeira, Beall, and Hueber 1988
Malacostraca–Branchiopoda	Branchiopoda (F): 536–517.2, 570–536	Event: Cambrian explosion; 545.0–495.0	532.9–508.4	Benton 1993

^a Nucleotide molecular clock as described in the figure 3.

^b A fossil of Calliphoridae from the Triassic proved erroneous (Evenhuis 1994).

at the nucleotide and the amino acid level for the Insecta by providing the most consistent rates of evolution between different lineages. The *coxI* locus also showed similar amino acid base compositions between most orders, a prerequisite for the amino acid mutation models used in this study. EF-1 α amino acid data were also constructed as a series of local molecular clocks as described for *coxI* amino acid data but resulted in unrealistically large values for the divergence between Decapoda and Branchiopoda (>1,000 MYA). It is noteworthy that many studies have identified the ability of *coxI* to form a robust molecular clock in insects, including the butterfly *Heliconius* sp. (Brower 1994), chewing lice

(Phthiraptera; Hafner et al. 1994), *Tetraopes* sp. (Coleoptera; Farrell 2001), and *Alpheus* sp. (Decapoda; Knowlton 1993).

The *coxI* phylogenies obtained were congruent with the insect molecular phylogenies of previous studies, although we do not attempt to resolve the phylogeny with certainty (Russo, Takezaki, and Nei 1995; Caterino and Sperling 1999; Lyman et al. 1999; Pellmyr and Leebens-Mack 1999). In addition, the *coxI* phylogeny of the insect orders was identical with the 18S and 28S phylogeny and a morphologically based insect phylogeny (Whiting et al. 1997; Wheeler et al. 2001). Furthermore, our phylogeny is supported by previous phy-

logenetic analyses that described Malacostraca and Branchiopoda (including Anostraca) forming a sister group to Insecta, using combined EF-1 α and RNA polymerase II (POL II) amino acid sequence data (Regier and Shultz 1997, 1998; Shultz and Regier 2000), combined 18S and 28S data (Friedrich and Tautz 1995), and mitochondrial gene translocations (Boore, Lavrov, and Brown 1998). Finally, our use of the Malacostraca as an out-group to the Anostraca-Insecta clade is also supported by the EF-1 α and POL II analyses (Regier and Shultz 1997).

Molecular clock methods are particularly susceptible to the problems of mutation rate heterogeneity between lineages. We addressed this issue for amino acid data using a large number of local molecular clocks to represent the large range possible of temporal estimates each model could generate and to compare this method with a global molecular clock using second-codon-position nucleotides. The congruence in date estimates between these methods was surprising and particularly strong for deep nodes, despite the nucleotide data containing less phylogenetic information. Moreover, the temporal estimates from both the amino acid local clock models and the global nucleotide clocks were compatible with all known insect fossils, with the breakup of Gondwanaland and the Cambrian explosion. In addition, the use of a clock method that does not rely on a global molecular clock, the use of ML techniques, and the use of models that do not assume any one mechanism of rate variation should minimize or remove dating errors caused by incomplete taxon sampling. We recognize that the sample size defined in this study is far from complete, representing 5 of the 26 orders of extant insects, although the nodal dates described provide a series of potential calibration points for other insect molecular clocks where their fossil record is sparse (most insects) and would provide a more suitable method of calibration than using biogeographic assumptions. The nodal date for the earliest extant hexapod, Collembola, is desirable because it would provide a direct comparison with the fossil record, but the nodal dates from the surrounding lineages were compatible with a date of around 400 MYA.

The calibration point is crucial in all molecular clock calculations. One piece of data validating the calibration point used in this study is the lack of significant difference between the temporal estimates observed for the lepidopteran families Proxodidae and Incurvariidae despite two contrasting calibration points being used, the fossil record of the early ancestors of extant Blattaria used in this study for calibrating amino acid local clock models and the breakup of Gondwanaland used previously for calibrating first- and second-codon-position nucleotide data using the relative rates test (Pellmyr and Leebens-Mack 1999), although in this instance the second-codon-position nucleotide clock described here gives older dates for both these families. Although fossils of the earliest extant family of Blattaria, *Blaberidae*, are found in the Triassic period, five families of ancestral Blattaria (Pinnule Insects), namely, Adeloblattidae, Cobaloblattidae, Mylacridae, Necymylacridae, and Phyllob-

lattidae, have been unequivocally assigned to the late Carboniferous (Bashkirian epoch, 323 MYA; reviewed by Ross and Jarzembowski 1993; Labandeira 1994). Both the ancestors of extant Blattaria and the Blattaria fossils are particularly well preserved, perhaps because they favor damp habitats that favor fossilization, and in addition, entire specimens fossilize, reducing the reliance on wing fragments for identification (Wootton 1981). Apart from morphological analysis (Jarzembowski 1994), support for the classification of the extinct ancestors of Blattaria as members of this order is observed in one phylogenetic study where extant Blattaria are ancestral to Isoptera, their closest relative (Lo et al. 2000). In addition, it is possible that predatory behavior in the closest relative to the Blattaria/Isoptera lineage, the Mantoidea (mantids), was recently derived because it is a rare characteristic within Neoptera, and this would be consistent with the much later appearance of their fossils in Cretaceous and late Triassic strata (Labandeira 1994; Grimaldi 1997). Other members of this clade include Orthoptera and a paraphyletic clade of diverse Protorthoptera now extinct. The early radiation of Protorthoptera also coincides with the earliest Blattaria fossils (Labandeira 1994) and suggests that this date was close to the Blattaria/Orthoptera divergence.

Finally, the reliability of a molecular clock is also dependent on branch lengths of the root lineages (Yoder and Yang 2000). Although the stability of the amino acid clock was affected by the instability of the root lineages of the *coxI* phylogeny, i.e., small fluctuations in the root lineages affected the temporal estimates of all other nodes, this was only a major problem for dates within the root. The nucleotide clock also showed an unstable root, although there was no means of correcting for this using a global molecular clock.

Applications

The primary focus of this study is the resolution of deep lineages within the Insecta. The dating of the deepest nodes within the Insecta is particularly relevant for providing calibration points for dating studies within relevant families and genera and is particularly noteworthy given the congruence between the date estimates obtained and the fossil record. The haematophagous triatomine bugs are a case in question because their poor fossil record has thwarted an accurate understanding of their age. The bugs are the sole vectors of *Trypanosoma cruzi* (the agent of Chagas disease), and the *Rhodniini* (palm dwelling) and *Triatomini* (terrestrial and arboreal dwelling) genera describe the deepest evolutionary divergence within the family (Lyman et al. 1999; Gaunt and Miles 2000; Marcilla et al. 2001). The vast majority of triatomine bugs are restricted to Central and South America, and the date of divergence between the tribes *Rhodniini* and *Triatomini* (99.7–93.6 MYA) coincides with the breakup of Gondwanaland (≥ 95 MYA). The date most importantly provides a calibration point for current studies on the *Rhodnius* genus, linking the evolutionary age of bugs with their habitat. For example, the deepest split within triatomine bugs is close to the

date for the earliest palm megafossils (83.5–71.3 MYA), opening up the possibility of either *Rhodnius*-palm co-evolution or the *Rhodniini-Triatomini* split resulting from adaptation of one genera to a specific habitat.

A more fundamental date for insects is the origin of the insect-anostracan common ancestor. The emergence of the Insecta-Anostraca ancestor obtained here, 434.0–421.0 MYA, coincides with the earliest megafossil of vascular plants, *C. cambrensis*, found in the Wenlock epoch of the early Silurian 428–423 MYA (Edwards, Feehan, and Smith 1983). Cladistic studies of *C. caledonica* and *C. pertonii* besides several other early tracheophytes (vascular plants) suggest that numerous plant species were present in this epoch (Kenrick and Crane 1997, pp. 226–270). The correlation is compatible with the hypothesis that the true insects evolved from an aquatic arthropod that formed an ecological association with a specific group of terrestrial plants and subsequently coevolved with that plant group, a hypothesis that dates back almost three quarters of a century (Tilley 1928). Insect-pteridophyte coemergence from an aquatic-terrestrial anostracan ancestor was also compatible with the palaeontological flora from the three aquatic Devonian localities yielding insect or hexapod fossils, at Rhynie in Scotland (Whalley and Jarzembowski 1981; Ross and Jarzembowski 1993), and at Gaspé (Labandeira, Beall, and Hueber 1988) and Gilboa (Shear et al. 1984) in northeastern North America. For example, an early vascular plant *Zosterophylloids* is found at Gaspé, and one of the earliest vascular plants *Rhynopsida* is found at Rhynie. The Rhynie plants show three types of surface lesions inflicted while the plant was alive, of which two types could be attributed to penetration by an arthropod (Kevan, Chaloner, and Savile 1975). Reexamination of the Rhynie hexapods showed that one specimen was associated with an early Rhynie plant (Whalley and Jarzembowski 1981).

The date obtained for the origin of the neopteran clade comprising Blattaria, Orthoptera, and Hemiptera at 382.9–373.0 MYA for amino acid data and at 350.1 for nucleotide data to some extent coincides with the early pteridophytes (mostly ferns and horsetails) during the late Devonian at around 380 MYA (Kenrick and Crane 1997). For example, early macrofossils of the two largest groups of pteridophytes, the *Lycopsidea* and *Filicopsida*, occur in the early Devonian (417 MYA) and early Carboniferous (354 MYA), respectively. It is also interesting to note that this period immediately precedes the early date obtained for divergence of the Diptera and Lepidoptera lineages that occurred during the early Carboniferous and provides calibration points for further studies to estimate the number of insect orders that were established during the Carboniferous.

The origin of the neopteran insects investigated in this study also provides the latest date for the evolution of insect flight. Kevan, Chaloner, and Savile (1975) hypothesized that insect flight arose as an adaptation to the increasing height of vascular plants during the early evolution of trees. The date obtained here for the ancestor to the neopteran clade, which mostly coincides with the early pteridophytes, i.e., plants with stems tall enough

to prevent herbivory with ground-borne insects, is compatible with this hypothesis.

Acknowledgments


We gratefully acknowledge the financial support from the Wellcome Trust. We thank the two anonymous referees and the subeditor, as well as Z. Yang (University College London), G. Clark, I. Mauricio, S. Solis Mena (LSHTM, U.K.), and the staff at the Instituto Oswaldo Cruz and the Instituto Evando Chagas (Brazil). We are extremely grateful to the relevant authorities who kindly provided current information and detailed advice on the paleontological science described here.

LITERATURE CITED


- ADACHI, J., and M. HASEGAWA 1996a. Model of amino acid substitution in proteins encoded by mitochondrial DNA. *J. Mol. Evol.* **42**:459–468.
- . 1996b. Molphy: MOlecular PHYlogenetics. Available at <http://ftp.ism.ac.jp/pub/ISMLIB/MOLPHY>.
- BENTON, M. J., ed. 1993. *The fossil record 2*. Chapman & Hall, London.
- BOORE, J. L., D. V. LAVROV, and W. M. BROWN. 1998. Gene translocation links insects and crustaceans. *Nature* **392**:667–668.
- BROMHAM, L. D., A. E. RAMBAUT, and P. H. HARVEY. 2000. Determinants of rate variation in DNA sequence evolution of mammals. *J. Mol. Evol.* **43**:610–621.
- BROWER, A. V. Z. 1994. Rapid morphological radiation and convergence among races of the butterfly *Heliconius erato* inferred from patterns of mitochondrial DNA evolution. *Proc. Natl. Acad. Sci. USA* **91**:6491–6495.
- BROWN, J. M., O. PELLMYR, J. N. THOMPSON, and R. G. HARRISON. 1994. Phylogeny of *Greya* (Lepidoptera: Prodoxidae), based on nucleotide sequence variation in mitochondrial cytochrome oxidase I and II: congruence with morphological data. *Mol. Biol. Evol.* **11**:128–141.
- CATERINO, M. S., S. CHO, and F. A. H. SPERLING. 2000. The current state of insect molecular systematics: a thriving Tower of Babel. *Annu. Rev. Entomol.* **45**:1–54.
- CATERINO, M. S., and F. A. H. SPERLING. 1999. *Papilio* phylogeny based on mitochondrial cytochrome oxidase I and II genes. *Mol. Phylogenet. Evol.* **11**:122–137.
- COLLINSON, M. E., A. BOULTER, and P. L. HOLMES. 1993. Magnoliophyta ('Angiospermae'). Pp. 809–841 in M. J. BENTON, ed. *The fossil record 2*. Chapman & Hall, London.
- COOPER, A., and D. PENNY. 1997. Mass survival of birds across the Cretaceous-Tertiary boundary: molecular evidence. *Science* **275**:1109–1113.
- DANFORTH, B. N., and S. Q. JI. 1998. Elongation factor-1 alpha occurs in two copies in bees: implications for phylogenetic analysis of EF-1 alpha sequences in insects. *Mol. Biol. Evol.* **14**:381–390.
- EDWARDS, D., J. FEEHAN, and D. G. SMITH. 1983. A late Wenlock flora from Co. Tipperary, Ireland. *Bot. J. Linn. Soc.* **92**:19–36.
- ESSEGHIR, S., P. D. READY, R. KILLICK-KENDRICK, and R. BEN-ISMAIL. 1997. Mitochondrial haplotypes and phylogeography of *Phlebotomus* vectors of *Leishmania major*. *Insect Mol. Biol.* **6**:211–225.
- EVENHUIS, N. L. 1994. Catalogue of the fossil flies of the world (Insecta: Diptera). Backhuys, Leiden. Available at <http://www.bishopmuseum.org/bishop/ento/fossilcat/>.

- FARRELL, B. D. 2001. Evolutionary assembly of the milkweed fauna: cytochrome oxidase I and the age of *Tetraopes* beetles. *Mol. Phylogenet. Evol.* **18**:467–478.
- FOLEY, D. H., J. H. BRYAN, D. YEATES, and A. SAUL. 1998. Evolution and systematics of *Anopheles*: insights from a molecular phylogeny of Australian mosquitoes. *Mol. Phylogenet. Evol.* **9**:262–275.
- FRIEDRICH, M., and D. TAUTZ. 1995. Ribosomal DNA phylogeny of the major extant arthropod classes and the evolution of myriapods. *Nature* **376**:165–167.
- GAUNT, M., and M. A. MILES. 2000. The ecotopes and evolution of triatomine bugs (triatominae) and their associated trypanosomes. *Mem. Inst. Oswaldo Cruz* **95**:557–565.
- GRADSTEIN, F. M., F. P. AGTERBERG, J. G. OGG, J. HARDENBOL, P. VAN VEEN, J. THIERRY, and Z. HUANG. 1995. A Triassic, Jurassic, and Cretaceous chronostratigraphy charts. In W. A. BERGGREN, ed. *Geochronology, time scales and global stratigraphic correlation*. SEPM Spec. Publ., Tulsa **54**:95–126.
- GRADSTEIN, F. M., and J. OGG. 1996. A Phanerozoic time scale. *Episodes* **19**:3–5.
- GRIMALDI, D. 1997. A fossil mantis (Insecta: Mantodea) in Cretaceous amber of New Jersey, with comments on the early history of Dictyoptera. *Am. Mus. Novit.* **3024**:1–11.
- GRIMALDI, D., A. SHEDRINKSKY, and T. P. WAMPLER. 2000. A remarkable deposit of fossiliferous amber from the Upper Cretaceous (Turonian) of New Jersey. Pp. 1–76 in D. GRIMALDI, ed. *Studies on fossils in amber with particular reference to the Cretaceous*. Backhuys, Leiden.
- HAFNER, M. S., P. D. SUDMAN, F. X. VILLABLANCA, T. A. SPRADLING, J. W. DEMASTES, and S. A. NALDER. 1994. Disparate rates of molecular evolution in cospeciating hosts and parasites. *Science* **265**:1087–1090.
- HARLAND, W. B., R. L. ARMSTRONG, A. V. COX, L. E. CRIAG, A. G. SMITH, and D. G. SMITH. 1990. *A geologic time scale 1989*. Cambridge University Press, Cambridge.
- HEDGES, S. B., P. H. PARKER, C. G. SIBLEY, and S. KUMAR. 1996. Continental breakup and the ordinal diversification of birds and mammals. *Nature* **381**:226–229.
- HENNING, W. 1981. *Insect phylogeny*. Wiley, Chichester, U.K.
- HUELSENBECK, J. P., B. LARGET, and D. SWOFFORD. 2000. A compound Poisson process for relaxing the molecular clock. *Genetics* **154**:1879–1892.
- JARZEMBOWSKI, E. A. 1994. Fossil cockroaches or Pinnule Insects. *Proc. Geol. Assoc.* **105**:305–311.
- JIN, Y. G., B. R. WARDLAW, B. F. GLENISTER, and G. V. KOTLYAR. 1997. Permian chronostratigraphic subdivisions. *Episodes* **20**:10–15.
- JONES, D. T., W. R. TAYLOR, and J. M. THORNTON. 1992. The rapid generation of mutation data matrices from protein sequences. *CABIOS* **8**:275–282.
- KENRICK, P., and P. R. CRANE. 1997. *The origin and early diversification of land plants: a cladistic study*. Smithsonian Institution Press, Washington.
- KEVAN, P. G., W. G. CHALONER, and D. B. O. SAVILE. 1975. Interrelationships of early terrestrial arthropods and plants. *Palaeontology* **18**:391–417.
- KNOWLTON, N., L. A. WEIGT, L. A. SOLORZANO, D. K. MILLS, and E. BERMINGHAM. 1993. Divergence in proteins, mitochondrial-DNA, and reproductive compatibility across the Isthmus of Panama. *Science* **260**:1629–1632.
- LABANDEIRA, C. C. 1994. A compendium of fossil insect families. Pp. 15–71 in R. WATKINS, ed. *Contributions in Biology and Geology*, Vol. 88. Milwaukee Public Museum, Wisconsin.
- LABANDEIRA, C. C., B. S. BEALL, and F. M. HUEBER. 1988. Early insect diversification—evidence from a lower Devonian Bristletail from Quebec. *Science* **242**:913–916.
- LABANDEIRA, C. C., and J. J. SEPKOSKI. 1993. Insect diversity in the fossil record. *Science* **261**:310–315.
- LI, W.-H., and M. TANIMURA. 1987. The molecular clock runs more slowly in man than in apes and monkeys. *Nature* **326**:93–96.
- LO, N., G. TOKUDA, H. WATANABE, H. ROSE, M. SLAYTOR, K. MAEKAWA, C. BANDI, and H. NODA. 2000. Evidence from multiple gene sequences indicates that termites evolved from wood-feeding cockroaches. *Curr. Biol.* **10**:801–804.
- LYMAN, D. E., F. A. MONTEIRO, A. A. ESCALANTE, C. CORDON-ROSALES, D. M. WESSON, J. P. DUJARDIN, and C. B. BEARD. 1999. Mitochondrial DNA sequence variation among triatomine vectors of Chagas' disease. *Am. J. Trop. Med. Hyg.* **60**:377–386.
- MARCILLA, A., M. D. BARGUES, J. M. RAMSEY, E. MAGALLON-GASTELUM, P. M. SALAZAR-SCHETTINO, F. ABAD-FRANCH, J. P. DUJARDIN, C. J. SCHOFIELD, and S. MAS-COMA. 2001. The ITS-2 of the nuclear rDNA as a molecular marker for populations, species, and phylogenetic relationships in Triatominae (Hemiptera: Reduviidae), vectors of Chagas disease. *Mol. Phylogenet. Evol.* **18**:136–142.
- MCALPINE, J. F. 1970. First record of calypterate flies in the Mesozoic Era (Diptera: Calliphoridae). *Can. Entomol.* **102**:342–346.
- PELLMYR, O., and J. LEEBENS-MACK. 1999. Forty million years of mutualism: evidence for Eocene origin of the yucca-yucca moth association. *Proc. Natl. Acad. Sci. USA* **96**:9178–9183.
- POSADA, D., and K. A. CRANDALL. 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics* **14**:817–818.
- REGIER, J. C., and J. W. SHULTZ. 1997. Molecular phylogeny of the major arthropod groups indicates polyphyly of Crustaceans and a new hypothesis for the origin of hexapods. *Mol. Biol. Evol.* **14**:902–913.
- . 1998. Molecular phylogeny of arthropods and the significance of the Cambrian “explosion” for molecular systematics. *Am. Zool.* **38**:918–928.
- ROBERTS, J., J. CLAOUE-LONG, and P. J. JONES. 1995. Australian Early Carboniferous time. In W. A. BERGGREN, ed. *Geochronology, time scales and global stratigraphic correlation*. SEPM Spec. Publ., Tulsa **54**:24–40.
- ROSS, A. J., and E. A. JARZEMBOWSKI. 1993. Arthropoda (Hexapoda; Insecta). Pp. 363–426 in M. J. BENTON, ed. *The fossil record*. Chapman & Hall, London.
- RUSSO, C. A. M., N. TAKEZAKI, and M. NEI. 1995. Molecular phylogeny and divergence times of Drosophilid species. *Mol. Biol. Evol.* **12**:391–404.
- SANDERSON, M. J. 1997. A nonparametric approach to estimating divergence times in the absence of rate constancy. *Mol. Biol. Evol.* **14**:1218–1231.
- SHCHERBAKOV, D. E., E. D. LUKASHEVICH, and V. A. BLAGODEROV. 1995. Triassic Diptera and initial radiation of the order Dipterolog. *Res.* **6**:75–115.
- SHEAR, W. A., P. M. BONAMO, J. D. GRIERSON, W. D. I. ROLFE, E. L. SMITH, and R. A. NORTON. 1984. Early land animals in North America—evidence from Devonian age arthropods from Gilboa, New York. *Science* **224**:492–494.
- SHEAR, W. A., and J. KUKALOVA-PECK. 1990. The ecology of Paleozoic terrestrial arthropods: the fossil evidence. *Can. J. Zool.* **68**:1807–1834.

- SHIMODAIRA, H., and M. HASEGAWA. 1999. Multiple comparisons of log-likelihoods with applications to phylogenetic inference. *Mol. Biol. Evol.* **16**:1114–1116.
- SHULTZ, J. W., and J. C. REGIER. 2000. Phylogenetic analysis of arthropods using two nuclear protein-encoding genes supports a crustacean plus hexapod clade. *Proc. R. Soc. Lond. Ser. B* **267**:1011–1019.
- STRIMMER, K., and A. VON HAESLER. 1996. Quartet puzzling: a quartet maximum-likelihood method for reconstructing tree topologies. *Mol. Biol. Evol.* **13**:964–969.
- SWOFFORD, D. L. 2000. PAUP*: phylogenetic analysis using parsimony (* and other methods). Version 4.0. Sinauer Associates, Sunderland, Mass.
- TAKEZAKI, N., A. RZHETSKY, and M. NEI. 1995. Phylogenetic test of the molecular clock and linearized trees. *Mol. Biol. Evol.* **12**:823–833.
- THOMPSON, J. D., T. J. GIBSON, F. PLEWNIAC, F. JEANMOUGIN, and D. G. HIGGINS. 1997. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **25**:4876–4882.
- THORNE, J. L., H. KISHINO, and I. S. PAINTER. 1998. Estimating the rate of evolution of the rate of molecular evolution. *Mol. Biol. Evol.* **15**:1647–1657.
- TILYARD, R. J. 1928. Some remarks on the Devonian fossil insects from the Rhynie Chert beds, old red sandstone. *Trans. Entomol. Soc. Lond.* **76**:65–71.
- TUCKER, R. D., and W. S. MCKERROW. 1995. Early Paleozoic chronology: a review in light of new U-Pb zircon ages from Newfoundland and Britain. *Can. J. Earth Sci.* **32**:368–379.
- WEHR, W. C. 1998. Middle Eocene insects and plants of the Okanogan Highlands. Thomas Burke memorial. *Wash. State Mus. Res. Rep.* **6**:99–109.
- WHALLEY, P. 1986. A review of the current fossil evidence of Lepidoptera in the Mesozoic. *Biol. J. Linn. Soc.* **28**:253–271.
- WHALLEY, P., and E. A. JARZEMBOWSKI. 1981. A new assessment of Rhyniella, the earliest known insect, from the Devonian of Rhynie, Scotland. *Nature* **291**:317.
- WHEELER, W. C., M. WHITING, Q. D. WHEELER, and J. M. CARPENTER. 2001. The phylogeny of the extant hexapod orders. *Cladistics* **17**:113–169.
- WHITING, M. F., J. C. CARPENTER, Q. D. WHEELER, and W. C. WHEELER. 1997. The strepsiptera problem: phylogeny of the holometabolous insect orders inferred from 18S and 28S ribosomal DNA sequences and morphology. *Syst. Biol.* **46**:1–68.
- YODER, A. D., and Z. H. YANG. 2000. Estimation of primate speciation dates using local molecular clocks. *Mol. Biol. Evol.* **17**:1081–1090.
- RICHARD THOMAS, reviewing editor
- Accepted January 15, 2002



APPENDIX 1.—ML amino acid phylogeny of *coxI* showing neighbor-joining and parsimony bootstrap analysis for nucleotide data after 1,000 replications. Key: * Prodoxidae/Incurvariidae, Ditrystia, Adelidae/Helozelidae topology supported by the Shimodaira-Hasegawa test using the full optimization ML bootstrap analysis, with alternative branching order of Ditrystia forming an out-group to Prodoxidae/Incurvariidae and Adelidae/Helozelidae.



APPENDIX 2.—ML amino acid *coxI* phylogeny showing neighbor-joining and parsimony bootstrap analysis for amino acid data after 1,000 replications.