



# 1 **An intercomparison of approaches for improving predictability** 2 **in operational seasonal streamflow forecasting**

3 Pablo A. Mendoza<sup>1,\*</sup>, Andrew W. Wood<sup>1</sup>, Elizabeth Clark<sup>2</sup>, Eric Rothwell<sup>3</sup>, Martyn P. Clark<sup>1</sup>, Bart  
4 Nijssen<sup>2</sup>, Levi D. Brekke<sup>4</sup> and Jeffrey R. Arnold<sup>5</sup>

5  
6 <sup>1</sup>Hydrometeorological Applications Program, National Center for Atmospheric Research, Boulder, Colorado, USA

7 <sup>2</sup>Department of Civil and Environmental Engineering, University of Washington, USA

8 <sup>3</sup>Bureau of Reclamation, Boise, USA

9 <sup>4</sup>Bureau of Reclamation, Denver, USA

10 <sup>5</sup>Climate Preparedness and Resilience Programs, U.S. Army Corps of Engineers, Seattle, USA

11 *Correspondence to:* Pablo A. Mendoza (pmendoza@colorado.edu)

12 **Abstract.** For much of the last century, forecasting centers around the world have offered seasonal streamflow  
13 predictions to support water management. Recent work suggests that the two major avenues to advance seasonal  
14 predictability are improvements in the estimation of initial hydrologic conditions (IHCs) and the incorporation of  
15 climate information. This study investigates the marginal benefits of a variety of methods using IHC and/or climate  
16 information, focusing on seasonal water supply forecasts (WSFs) in five case study watersheds located in the U.S.  
17 Pacific Northwest region. We specify two benchmark methods that mimic standard operational approaches – statistical  
18 regression against IHCs, and model-based ensemble streamflow prediction (ESP) – and then systematically inter-  
19 compare WSFs across a range of lead times. Additional methods include: (i) statistical techniques using climate  
20 information either from standard indices or from climate reanalysis variables; and (ii) several hybrid/hierarchical  
21 approaches harnessing both land surface and climate predictability. In basins where atmospheric teleconnection  
22 signals are strong, and when watershed predictability is low, climate information alone provides considerable  
23 improvements. For those basins showing weak teleconnections, custom predictors from reanalysis fields were more  
24 effective in forecast skill than standard climate indices. ESP predictions tended to have high correlation skill but  
25 greater bias compared to other methods, and climate predictors failed to substantially improve these deficiencies  
26 within a trace weighting framework. Lower complexity techniques were competitive with more complex methods,  
27 and the hierarchical expert regression approach introduced here (HESP) provided a robust alternative for skillful and  
28 reliable water supply forecasts at all initialization times. Three key findings from this effort are: (1) objective  
29 approaches supporting methodologically consistent hindcasts open the door to a broad range of beneficial forecasting  
30 strategies; (2) the use of climate predictors can add to the seasonal forecast skill available from IHCs; and (3) sample  
31 size limitations must be handled rigorously to avoid over-trained forecast solutions. Overall, the results suggest that  
32 despite a rich, long heritage of operational use, there remain a number of compelling opportunities to improve the skill  
33 and value of seasonal streamflow predictions.



## 34 1 Introduction

35 The operational hydrology community has long grappled with the challenge of producing skillful seasonal  
36 streamflow forecasts to support water supply operations and planning. Proactive water management has become  
37 critical for many regions in the world that are susceptible to water stress associated with the intensification of the  
38 water cycle. Paradoxically, although we have seen important technological advances – including increased computing  
39 power, the broader availability to climate reanalysis, forecasts and reforecasts, and more complex process-based  
40 hydrologic models (Pagano et al., 2016), the skill of operational seasonal runoff predictions in the US, termed water  
41 supply forecasts (WSFs), has shown little or no improvement over time (e.g., Pagano et al., 2004; Harrison and Bales,  
42 2016). Hence, there is both a scientific and practical need to understand the potential of new datasets, modeling  
43 resources and methods to accelerate progress towards more skillful and reliable operational seasonal streamflow  
44 forecasts.

45 There is general consensus in the research community on the main opportunities to improve seasonal streamflow  
46 prediction skill (e.g., Maurer et al., 2004; Wood and Lettenmaier, 2008; Yossef et al., 2013). These include improving  
47 knowledge of: (i) the amount of water stored in the catchment – hereinafter referred to as initial hydrologic conditions  
48 (IHCs), and (ii) weather and climate outcomes during the forecast period. Our ability to leverage the first predictability  
49 source (i.e., hydrologic predictability) depends on the accuracy of watershed observations and models, including  
50 model input forcings (e.g., precipitation and temperature), process representations, and the effectiveness of hydrologic  
51 data assimilation (DA) methods. Our ability to leverage the second source (climate predictability) depends both on  
52 how well we can characterize and predict the state of the climate and on how effectively we can incorporate this  
53 information into streamflow forecasting methods. This idea has been explored in different frameworks using standard  
54 indices – e.g., Niño3.4, the Pacific Decadal Oscillation (PDO) – and/or custom (i.e., watershed-specific) climate  
55 indices derived from climate reanalyses (e.g., Grantz et al., 2005; Bradley et al., 2015), or using seasonal climate  
56 forecasts to run hydrologic model simulations (e.g., Wood et al., 2005; Yuan et al., 2013).

57 Despite generally promising findings from this body of work and from a number of agency development efforts  
58 (Weber et al., 2012; Demargne et al., 2014), current operational practice in the US still takes little to no advantage of  
59 large-scale climate information for real-time seasonal streamflow forecasting. Clear examples can be found in the  
60 western United States, a large snowmelt dominated region where official WSFs are produced via two main approaches:  
61 (i) statistical models leveraging in situ watershed moisture measurements such as snow water equivalent (SWE),  
62 accumulated precipitation and streamflow (Garen, 1992; Pagano et al., 2004); and (ii) outputs from the National  
63 Weather Service (NWS) Ensemble Streamflow Prediction method (ESP; Day, 1985; Crochemore et al., 2016), based  
64 on watershed modeling. These approaches rely solely on the predictability from IHCs and do not leverage any type of  
65 large-scale current or future climate state information that might influence the forecasted hydrologic outcomes.

66 This paper presents an assessment of several seasonal streamflow prediction approaches in harnessing both  
67 watershed and climate related predictability. The methods are applied to seasonal WSFs and span a range of  
68 complexity, from purely statistical to purely dynamical and hybrid statistical/dynamical approaches. In this paper,  
69 ‘increased complexity’ indicates a gradient from purely data-driven techniques (e.g., linear regression) to the use of  
70 dynamical watershed models (Plummer et al., 2009), the outputs of which may be further processed using additional



71 statistical approaches. Although most of the techniques evaluated here are not new, the intercomparison offers new  
72 insights for researchers and developers in the operational community because: (1) the experiment is broader than prior  
73 efforts and benchmarks alternative methods against current operational ones; and (2) the methods are chosen to be  
74 operationally feasible, avoiding the use of data that cannot be obtained in real-time. In addition, the work uses a  
75 hindcast/verification framework and follows more rigorous standards for cross-validation than were used in some of  
76 the prior studies.

77 The remainder of this paper is organized as follows. Section 2 describes prior methodological work and context  
78 for statistical, dynamical and hybrid approaches to seasonal streamflow forecasting. The study domain is described in  
79 Section 3. Datasets, experimental design, individual methods, and forecast verification measures are detailed in  
80 Section 4. Results and discussion are presented in Section 5, followed by the main conclusions of this study (Section  
81 6).

## 82 2 Background

83 Seasonal streamflow forecasting methods can be categorized as dynamical, statistical, or hybrid, and span  
84 different degrees of complexity and information requirements. Dynamical methods use time-stepping simulation  
85 models to represent hydrologic processes. They describe future climate using either historical meteorology or inputs  
86 derived from seasonal climate forecasts (e.g., Beckers et al., 2016). On the other hand, statistical or purely data-driven  
87 methods rely on empirical relationships between seasonal streamflow volumes, and large-scale climate variables  
88 and/or in situ watershed observations. Several statistical approaches can be found in the literature, encompassing  
89 different degrees of complexity (e.g., Garen, 1992; Piechota et al., 1998; Grantz et al., 2005; Tootle et al., 2007; Wang  
90 et al., 2009; Moradkhani and Meier, 2010). Other studies have tested multi-model combination techniques for purely  
91 statistical seasonal forecasts, using objective performance criteria (e.g., Regonda et al., 2006), both performance and  
92 predictor state information (Devineni et al., 2008), and Bayesian model averaging (e.g., Mendoza et al., 2014), among  
93 others.

94 Hybrid methods strive to combine the strengths from both dynamical and statistical techniques. For instance,  
95 uncertainties in dynamical predictions indicate that dynamical forecasts can benefit from statistical post-processing  
96 (e.g., Wood and Schaake, 2008). One line of research has examined the potential benefits of using simulated watershed  
97 state variables – either from hydrologic or land surface models – as predictors for statistical models (e.g., Rosenberg  
98 et al., 2011; Robertson et al., 2013). Another popular technique consists in incorporating climate information within  
99 ESP frameworks, either deriving input sequences of mean areal precipitation and temperature from current climate or  
100 climate forecast considerations (e.g., Werner et al., 2004; Wood and Lettenmaier, 2006; Luo and Wood, 2008; Gobena  
101 and Gan, 2010; Yuan et al., 2013) – referred to as *pre-ESP* –, or ESP weighting (also referred to as *post-ESP*) based  
102 on climate information (e.g., Smith et al., 1992; Werner et al., 2004; Najafi et al., 2012; Bradley et al., 2015). Werner  
103 et al. (2004) found that the post-ESP method (termed ‘trace weighting’) was more effective than pre-ESP to improve  
104 forecast skill.

105 The combination of outputs from different models has also been shown to benefit seasonal hydroclimatic  
106 forecasting (e.g., Hagedorn et al., 2005). Although several studies have demonstrated that statistical multimodel



107 techniques applied on dynamical models tend to outperform the ‘best’ single model (e.g., Georgakakos et al., 2004;  
108 Duan et al., 2007), fewer insights have been gained on combining statistical or dynamical models in seasonal  
109 streamflow forecasting. Recently, Najafi and Moradkhani (2015) tested multimodel combination techniques of  
110 different complexities from both statistical and dynamical forecasts, concluding that model combination generally  
111 outperforms the best individual forecast model. Many sophisticated seasonal forecasting frameworks can be found in  
112 the literature, some of which incorporate DA techniques (e.g., Dechant and Moradkhani, 2011), a topic not discussed  
113 here. For this reason, the hydrology community may benefit from a broad assessment of the marginal benefits of  
114 choices made in a range of seasonal streamflow forecasting frameworks.

### 115 3 Study Domain

116 Our test domain is the U.S. Pacific Northwest (PNW) region (Figure 1), which relies heavily on winter snow  
117 accumulation and spring snowmelt to fulfill water needs during spring and summer (e.g., Mote, 2003; Maurer et al.,  
118 2004; Wood et al., 2005). We select catchments contributing to five reservoirs: Dworshak (DWRI1), Howard Hanson  
119 (HHDW1), Hungry Horse (HHWM8), Libby (LYDM8) and Prineville (PRVO). Two of them – Hungry Horse and  
120 Prineville reservoirs – are owned and operated by the U.S. Bureau of Reclamation (USBR), while the rest are operated  
121 by the U.S. Army Corps of Engineers (USACE).

122 The main physical and hydroclimatic characteristics of the case study basins are summarized in Table 1. These  
123 basins cover a wide range of runoff efficiencies (from 0.13 at Prineville to 0.78 at Howard Hanson) and dryness indices  
124 (from 0.63 at Howard Hanson to 3.83 at Prineville). Relatively high basin-averaged elevations condition a pronounced  
125 seasonal temperature pattern, with minimum values below the freezing point between December and February, and  
126 maximum temperatures during June-September (not shown). These topographic and hydroclimatic features favor  
127 snowpack development in the months October-April, stressing the seasonal behavior of other water storages and  
128 fluxes. This is illustrated in Figure 2, including model precipitation (i.e., observed precipitation with a snow correction  
129 factor, SCF) and monthly averages of hydrologic variables simulated with the Sacramento Soil Moisture Accounting  
130 (SAC-SMA, Burnash et al., 1973) and SNOW-17 (Anderson, 1973) watershed models (see Section 4). Although  
131 seasonal precipitation patterns may differ, water starts accumulating in October as snow water equivalent (SWE)  
132 and/or soil moisture (SM) in all basins. Increases in SM and runoff in most basins are driven by snowmelt at the  
133 beginning of spring with the exception of Howard Hanson, where the bulk of annual streamflow occurs in November-  
134 May. Among these basins, Dworshak, Hungry Horse and Libby share similar SWE, soil moisture, and runoff cycles,  
135 although precipitation is relatively uniform in the last one throughout the year.

136 The hydroclimatology of the PNW region is affected by a number of large-scale climate teleconnections. The  
137 warm (cold) phase of El Niño Southern Oscillation (ENSO) is typically associated with above (below) average  
138 temperatures and below (above) average precipitation during winter (e.g., Redmond and Koch, 1991), and therefore  
139 decreased (increased) snowpack (Clark et al., 2001) and spring/summer runoff (e.g., Piechota et al., 1997). The Pacific  
140 Decadal Oscillation (PDO; Mantua et al., 1997) – which reflects the dominant mode in decadal variability of SSTs –  
141 has also been found a relevant driver for the hydroclimatology of the PNW (e.g., McCabe and Dettinger, 2002). The  
142 joint influence of ENSO and PDO on North American climate conditions, snowpack and spring/summer runoff has



143 been also well recognized and documented (e.g., Hamlet and Lettenmaier, 1999). As a consequence, many authors  
144 have explored the incorporation of large-scale climate information for seasonal streamflow forecasting in the PNW –  
145 using either standard indices (e.g., Hamlet and Lettenmaier, 1999; Maurer et al., 2004), custom indices from reanalysis  
146 fields (e.g., Opitz-Stapleton et al., 2007; Tootle et al., 2007), both (Moradkhani and Meier, 2010), or downscaled  
147 climate forecasts (Wood et al., 2005) – finding improved predictability for lead times longer than 2 months, and  
148 particularly in years of strong anomalies in climate oscillations such as ENSO.

## 149 **4 Approach**

### 150 **4.1 Experimental Design**

151 We use several decades of seasonal streamflow hindcasts to assess a suite of methods (Figure 3), focusing on  
152 April-July streamflow (runoff) volume, the most common western US water supply forecast predictand. Probabilistic  
153 (ensemble) WSFs for this period are generated the first day of each month from October to April, in every year of the  
154 hindcast period 1981-2015. For the methods involving statistical prediction, we use a leave-three-out cross validation  
155 at all stages of the forecast process. This procedure is repeated for consecutive 3-year periods (e.g., 1984-1986, 1987-  
156 1989, 1990-1992, etc.), except for the last time window (2014-2015).

157 The techniques assessed here are categorized as follows. The first group, *IHC-based* methods, includes two  
158 approaches (referred to as *benchmark methods*) – ESP and IHC-based statistical – currently used operationally in the  
159 western U.S. (both harnessing only IHC information), and a very simple ESP post-processor to reduce systematic  
160 biases. A second group, *climate-only* methods, includes statistical techniques harnessing climate information from  
161 two different sources – standard indices (e.g., Niño3.4, PDO, AMO), or variables extracted from the Climate System  
162 Forecast Reanalysis (CFSR; Saha et al., 2010). A third group of *hybrid* or *hierarchical* methods includes subgroups  
163 of techniques that: (i) combine watershed predictors (IHCs) and climate predictors (either indices or CFSR variables)  
164 within a statistical framework, (ii) use climate information to post-process outputs from a dynamical method (i.e.,  
165 ESP), or (iii) combine purely climate-based ensembles with purely watershed-based ensembles.

166 In operational practice, ESP produces an ensemble of streamflow estimates whereas statistical water supply  
167 forecasting yields a statistical distribution. In this study, we generate ensembles of the final predictand for all methods.  
168 An ensemble size 500 is used – wherein the members are generated through a resampling (in some cases weighted)  
169 of the predictive distributions – except for the ESP and bias-corrected ESP methods, for which 32 members are  
170 generated (i.e., 35 total historical years less the three out of sample test years). In the statistical approaches, seasonal  
171 flows are log-transformed and predictor and predictand data are normalized before training statistical method  
172 parameters or weights. The statistical models were then applied in log-standard-normal space for forecast generation,  
173 and predictands are transformed back to streamflow space.



## 174 4.2 Forecasting Methods

### 175 4.2.1 IHC-based methods

#### 176 Ensemble Streamflow Prediction (ESP)

177 The traditional ESP method (Day, 1985) relies on deterministic hydrologic model simulations forced with  
178 observed meteorological inputs up to the initialization time of the forecast. The approach assumes that meteorological  
179 data and model are perfect – i.e., there are no errors in IHCs, and that historical meteorological conditions during the  
180 simulation period can be used to represent climate forecast conditions. For hindcast verification purposes, the  
181 meteorological input traces associated with forecast years must be excluded.

182 The hydrology models used in this study were the NWS Snow-17, SAC-SMA and a unit-hydrograph routing  
183 model, all implemented in lumped fashion with 2-3 snow elevation zones per watershed. The models were calibrated  
184 via an automated multi-objective parameter estimation to reproduce observed daily streamflow. Hydrologic model  
185 forcings were drawn from a 1/16 degree real-time implementation of the ensemble forcing generation method  
186 described in Newman et al. (2015). Naturalized flow data was obtained from a combination of sources, including the  
187 Bonneville Power Administration (BPA, 2011), the USBR Hydromet historical data access system, and the USACE  
188 Data Query System.

189 Figure 4 shows simulated and observed monthly time series of streamflow for the period Oct/1990 – Sep/2000.  
190 With the exception of Prineville, where neither meteorology nor flow are well measured, all basins show values of  
191 NSE and  $r$  higher than 0.76 and 0.87, respectively. Further, the climatological seasonality of streamflow is reproduced  
192 well in all basins.

#### 193 Statistical forecasting using initial hydrologic conditions (Stat-IHC)

194 This method mimics the approach of the U.S. Natural Resources Conservation Service (NRCS), but differs in  
195 using model-simulated basin-averaged SWE and SM as surrogates for ground-based observations of SWE,  
196 precipitation and streamflow used operationally by the NWS and NRCS (as demonstrated in Rosenberg et al., 2011).  
197 A linear regression equation is developed between log-transformed seasonal runoff and IHCs represented by the sum  
198 of simulated basin-averaged SWE and SM. The training period equations are used to issue a deterministic runoff  
199 volume prediction for each year left out, and ensembles are generated by adding 500 Gaussian random numbers with  
200 zero mean and a standard deviation equal to the standard error of the individual prediction. The predictions are then  
201 exponentiated.

#### 202 Bias Corrected Ensemble Streamflow Prediction (BC-ESP)

203 ESP predictions often exhibit a systematic bias due to inadequate model parameters and/or other sources or error  
204 (e.g., input forcing selection, model structure). If the ESP approach provides a consistent hindcast, as it does here,  
205 post-processing in the form of a simple bias-correction (BC-ESP) can be applied. This is achieved by multiplying the  
206 raw ESP forecasts by a mean scaling factor that is obtained by computing the ratio between the mean of observed  
207 seasonal runoff volumes (i.e., the predictand) and the mean of ESP forecast median volumes, for each initialization  
208 time. Each scaling factor calculation and application is cross-validated.



#### 209 4.2.2 Statistical forecasting harnessing only climate information

##### 210 **Multiple linear regression (MLR) using standard climate indices (Stat-Ind)**

211 This method evaluates 12 standard climate indices as candidate predictors (Table 2). For each initialization time  
212 (e.g., November 1) and climate index (e.g., Niño3.4), the 3-month time window that maximizes the correlation  
213 coefficient between a preceding seasonal (e.g., August-October) predictor average and seasonal streamflow volume  
214 over the training period is selected. Once this procedure is repeated for all potential predictors, the best possible time  
215 series are obtained for the 12 climate indices, and ensemble forecasts are produced for a given initialization through  
216 the following steps:

- 217 1. Several combinations of predictors are selected subject to the constraint that no pairs of predictors with an  
218 inter-correlation larger than  $C_{\text{thresh}} = 0.3$  should be included.
- 219 2. Stepwise MLR models are fit for all combinations of predictors identified in Step 1, and the set of predictors  
220 that minimizes the Bayesian Information Criterion (BIC) score (Akaike, 1974) over the training period is  
221 selected.
- 222 3. An ensemble forecast is generated (as for Stat-IHC) with the MLR model from Step 2.

223 We choose MLR over more parameterized regression methods (e.g., local polynomial regression) since these  
224 were found to perform poorly in cross-validation, mainly due to the limited samples sizes available in the seasonal  
225 hydrologic prediction context.

##### 226 **Partial Least Squares Regression using reanalysis fields (Stat-CFSR)**

227 The teleconnections captured in off-the-shelf climate indices are not influential everywhere. Therefore, we also  
228 assess the potential of custom climate predictor indices derived from reanalysis data. Following Tootle et al. (2007),  
229 we use Partial Least Squares Regression (PLSR; Wold, 1966) to extract information from climate fields. PLSR  
230 decomposes a set of independent variables  $X$  and dependent variables  $Y$  into a small number of principal components  
231 that explain as much covariance as possible between the two variable sets (Abdi, 2010). PLSR components are formed  
232 from CFSR 700 mb geopotential height ( $Z700$ ) and sea surface temperatures (SSTs) over the domain  $20^{\circ}\text{S}$ – $80^{\circ}\text{N}$ ;  
233  $130^{\circ}\text{E}$ – $10^{\circ}\text{W}$ . For dates beyond 2010, we merged the 1979–2010 CFSR data with monthly analysis fields from the  
234 Climate Forecast System version 2 (CFSv2; Saha et al., 2014), aggregating the latter product to  $2.0^{\circ} \times 2.0^{\circ}$  horizontal  
235 resolution. Similar to the Stat-Ind method, we use 3-month averages of these variables. The seasonal forecasts are  
236 generated for each initialization by following these steps:

- 237 1. Compute principal components from the combined SST and  $Z700$  gridded values for each training sample  
238 and the left-out prediction years.
- 239 2. Fit a regression model to the resulting PLSR components (predictors), accepting each additional component  
240 only when its mean partial correlation with volume runoff is above a threshold. We used a threshold of 0.30  
241 throughout the study after finding that nearby values – e.g., 0.25, 0.35 – did not substantially change the  
242 results. The small sample size and low predictability supported at most two components.
- 243 3. Compute a mean runoff volume forecast using the regression model obtained in Step 2, and generate an  
244 ensemble by adding 500 Gaussian random numbers with zero mean and a standard deviation equal to the





245 root mean squared error of prediction (RMSEP) obtained from leave-three-out cross validation within the  
246 training period.

247 The main implication of developing PLSR components and the subsequent estimation of regression coefficients  
248 in cross validation – as conducted here – is that climate information from the target prediction period is not used at  
249 all, as is the case in real-time systems. This is a key methodological difference versus past studies that used all  
250 historical available information to define custom reanalysis predictor fields (e.g., Grantz et al., 2005; Regonda et al.,  
251 2006; Bracken et al., 2010; Mendoza et al., 2014), yielding a moderate yet erroneous boost in predictability.

#### 252 **4.2.3 Hybrid/hierarchical methods combining watershed and climate information**

##### 253 **Stepwise MLRs using IHCs and climate predictors**

254 We applied two statistical methods that combine climate and dynamical watershed model predictors: Stat-Ind-  
255 IHC (which uses climate indices and IHCs), and Stat-CFSR-IHC (which uses CFSR-based PLSR components and  
256 IHCs). These approaches are implemented in identical fashion to Stat-Ind, except that IHCs are added to the potential  
257 suite of climate predictors.

##### 258 **Hierarchical Ensemble Streamflow Prediction (HESP)**

259 The underlying idea of HESP is that the two main sources of predictability – watershed IHCs and climate – may  
260 best be addressed sequentially to ensure that only climate uncertainty is related to climate predictors. This may not be  
261 the case if a climate variable enters first into a regression model that attempts to explain streamflow variance from both  
262 IHCs and climate, possibly leading to a sub-optimal predictor selection. HESP is thus a hierarchical regression  
263 approach in which streamflow is first related to IHCs by fitting  $Q = f(\text{IHC predictors}) + \mathcal{E}_{climate}$ , given sufficient IHC  
264 predictor strength. The residual uncertainty is then related to climate predictors (again if possible) by fitting  $\mathcal{E}_{climate}$   
265  $= g(\text{climate predictors}) + \mathcal{E}_{residual}$ , such that the final forecast equation takes the form:

$$266 \quad Q = f(\text{IHC predictors}) + g(\text{climate predictors}) + \mathcal{E}_{residual} \quad (1)$$

267 Here the predictor pool used to explain  $\mathcal{E}_{climate}$  may include standard climate indices or reanalysis PLSR  
268 components, depending on the performance obtained during the training period. Absent IHC predictability, HESP is  
269 equivalent to Stat-Ind or Stat-CFSR; whereas without climate predictability, it defaults to Stat-IHC. Lacking both IHC  
270 and climate predictability, HESP defaults to climatology – i.e., an ensemble forecast is issued by resampling from  
271 historical observations over the training period.

##### 272 **ESP Trace Weighting Scheme (TWS)**

273 A well-known strategy for incorporating climate information into ESP forecasts is called ‘trace weighting’  
274 (Smith et al., 1992; Werner et al., 2004), where forecasted flow probabilities are corrected by weighting each ensemble  
275 member according to the similarity between a climate-related feature of the current year (e.g., PDO) and the  
276 meteorological year used to generate that member. Here, for a given basin and forecast period, either climate indices  
277 or CFSR-based components are selected based on their training period performance (i.e., RMSE) and used to weight  
278 each trace obtained from BC-ESP (see Section 7.1 for further details).





### 279 **Equally weighted ensembles (EWE) and RMSE-weighted ensembles (RWE)**

280 EWE combines the best-performing climate-only hindcast (i.e., Stat-Ind or Stat-CFSR, based on RMSE over the  
281 training period) with the best watershed-only hindcast (either Stat-IHC or BC-ESP), resampling ensemble members  
282 equally from each source to form a new 500-member ensemble forecast. A variation of this combination approach  
283 (RWE) instead performs a weighted resampling from the two forecast sources according to their skill during the  
284 training period: i.e., the weights equal  $1/\text{RMSE}$ , where RMSE the root mean squared error of the ensemble median.

### 285 **Bayesian Model Averaging (BMA) and Quantile model averaging (QMA)**

286 These methods combine the best-performing climate-only hindcast with the best performing watershed-only  
287 hindcast. While BMA (Raftery et al., 2005) attempts to provide a weighted average of forecast probability densities,  
288 QMA (Schepen and Wang, 2015) applies a weighted average to forecast values (quantiles) for a given cumulative  
289 probability. A notable difference between the two approaches is that QMA produces smoother and consistently  
290 unimodal distributions compared to potentially bimodal BMA outputs (Schepen and Wang, 2015). More details on  
291 these techniques are provided in section 7.2.

## 292 **4.3 Forecast evaluation**

293 Forecast method performance was evaluated using the metrics listed in Table 3. These include some standard  
294 metrics used in hydrology, such as correlation coefficient ( $r$ ), root mean squared error ( $RMSE$ ), and percent bias, and  
295 also probabilistic measures to assess skill and reliability. Skill is obtained using the continuous ranked probability  
296 score (CRPS; Hersbach, 2000), which measures the temporal average error between forecast CDF with that from the  
297 observation. Forecast reliability – i.e., adequacy of the forecast ensemble spread to represent the uncertainty in  
298 observations – is evaluated using an index from the predictive quantile-quantile (QQ) plot (Renard et al., 2010). QQ  
299 plots compare the empirical CDF of forecast  $p$ -values (i.e.  $P_i(o_i)$ , where  $P_i$  and  $o_i$  are the forecast CDF and observation  
300 at year  $i$ ) with that from a uniform distribution  $U[0,1]$  (Laio and Tamea, 2007).

301 Confidence intervals for the verification statistics are created using bootstrapping with replacement. In each  
302 resampling step,  $N$  pairs of ensemble forecasts and observations were resampled from the original joint distribution  
303 ( $N$  is the total number of events for which probabilistic forecasts are available). This process is repeated 1000 times,  
304 and all statistics are then computed for each realization and ranked in order to obtain 95 % confidence limits.

## 305 **5 Results and discussion**

### 306 **5.1 Deterministic evaluation**

307 We first compare methods using the WSF median, a critical predictand for many water decisions (e.g., Lake  
308 Powell releases on the Colorado River in the western US). Figure 5 displays correlation coefficients ( $r$ ) between  
309 forecast median and observed April-July runoff volumes for the five case study basins. As expected, near-zero or  
310 negative  $r$  values were obtained for October 1 and November 1 WSFs with the IHC-based methods. Negative  
311 correlation scores arise in very low-skill situations as an artifact of cross-validation (e.g., leaving a high predictand  
312 out of a training sample biases the resulting prediction in the opposite direction). The seasonality of SM and SWE in



313 the basins of interest (Figure 2) does not yield watershed moisture accumulations with predictive power until  
314 December or January. In contrast,  $r$  values as high as 0.48 for Dworshak and 0.49 for Hungry Horse could be attained  
315 on October 1 using only information from climate indices (Stat-Ind). Generally, but not everywhere, methods  
316 harnessing predictability from the climate (with the exception of TWS) enhance skill in comparison to IHC-based  
317 methods at initializations early in the water year. TWS is unable to shift the parent ESP distribution sufficiently to  
318 impart much climate skill at this time of year.

319 After January, the hydrologic model begins to capture a useful moisture variability signal from the watershed,  
320 thus IHCs start to become a dominant source of predictability in all basins. Indeed, watershed information is  
321 particularly relevant at Libby and Prineville (Figure 5d and 5e), where correlations within the range 0.39-0.47 are  
322 achieved as early as December 1 with the three IHC-based techniques. In these basins, standard climate indices do not  
323 provide useful long-lead predictability, although CFSR-based predictors do support a consistent improvement. For  
324 example, the correlation from Stat-Ind for Libby (Prineville) on December 1 is -0.23 (0.02), while the  $r$  value from  
325 Stat-CFSR is 0.19 (0.30). These differences between Stat-Ind and Stat-CFSR remain at these basins for subsequent  
326 monthly initializations.

327 Figure 5 reveals several notable outcomes that are evident in many of the results plots. First, a linear regression  
328 against IHCs can provide similar  $r$  values than the more computationally expensive ESP method, especially at late  
329 initializations (i.e. March 1 or April 1). Likewise, straightforward ensemble combination techniques (e.g., EWE or  
330 RWE) may outperform more complex methods such as BMA (e.g., February 1 – April 1) at all basins. From a  
331 correlation skill perspective, on the other hand, ESP generally outperforms the rest of the methods in late winter and  
332 spring. For example, ESP provides the highest  $r$  values for Dworshak (0.82) and Howard Hanson (0.67) on April 1.  
333 Notably, EWE was found to be the best method on April 1 for Hungry Horse ( $r = 0.88$ ) and Prineville ( $r = 0.79$ ) based  
334 on correlation. This indicates that, although simple post-processing can provide substantial forecast improvement, the  
335 small sample size available for training during the cross-validation process results in noisy parameter estimates that  
336 can undermine the potential correlation skill achievable with techniques that are more complex.

337 Root mean squared errors (RMSE) for ensemble forecast medians (Figure 6) show that despite some  
338 discrepancies between techniques, inter-method differences are not as large as for correlation. In most basins, errors  
339 can be reduced at earlier initializations (i.e., Oct 1 and Nov 1) by introducing climate information. For instance, on  
340 October 1, Stat-Ind and Stat. Ind+IHC generate respective reductions in RMSE of 10% and 13% at Dworshak, 23%  
341 and 16% at Howard Hanson, and 14% and 12% at Hungry Horse, relative to the best IHC-based method in each basin.  
342 These benefits are seen in most initializations and catchments except at Libby, where the best results were mostly  
343 achieved using ESP (Oct 1) and Stat-IHC (Dec 1, and Feb 1 – Apr 1). In agreement with Beckers et al. (2016), this  
344 study was unable to find encouraging climate teleconnections at Libby, despite its relative proximity to Hungry Horse.

345 Figure 6 underscores that from a median error perspective, intuitive ensemble combinations approaches (i.e.,  
346 EWE and RWE, shown in dark green) can be effective for reducing forecast errors once the watershed begins to  
347 provide useful predictability (i.e. after January 1). For instance, EWE was the best performing method in Hungry  
348 Horse and Prineville for forecasts initialized on March 1 and Apr 1. Further, Figure 6 illustrates that the best (or worst)  
349 techniques when looking at RMSE vary with each basin, although it is clear that TWS and only-climate methods



350 perform poorly at early and late initializations, respectively. The joint inspection of Figures 5 and 6 shows that inter-  
351 method agreement in correlation does not necessarily translate into similar forecast median errors. For example, while  
352 ESP and HESP provide close  $r$  values at Dworshak (0.74 and 0.73) on March 1, larger discrepancies are obtained in  
353 RMSE, with values of 0.58 million-acre-feet (MAF) and 0.79 MAF for ESP and HESP, respectively.

354 Another interesting result is that no substantial reductions in RMSE were achieved at Howard Hanson between  
355 October 1 and April 1, in contrast to the gradual growth of hydrologic predictability to support forecast skill in other  
356 basins. Indeed, the best performing techniques for October 1 (Stat-Ind) and April 1 (BC-ESP) forecasts provide similar  
357 RMSE values ( $\sim 0.064$  and  $0.065$  MAF, respectively). This outcome can be attributed to the relatively more rainfall-  
358 dominated hydrograph of Howard Hanson in comparison to the rest of the catchments (Table 1; Figure 2), and  
359 sustained runoff variability generated by seasonally high SM and fall-winter precipitation.

360 Figure 7 (forecast median bias) shows that raw ESP outputs have the largest biases through most initializations  
361 at Howard Hanson, Libby and Prineville. In particular, absolute biases at Prineville – which is the worst simulated  
362 basin in the group – increase to 53% on October 1 before decreasing to 20% on April 1. Further, relatively large biases  
363 (in comparison to the rest of techniques) were obtained at late initializations in Dworshak and Hungry Horse.  
364 Excepting Prineville, inter-method differences were not substantial, and none of the methods exceeded a 16% bias at  
365 any initialization. The simple bias correction applied in this study was able to reduce absolute biases to less than  $\pm$   
366 3% at Prineville, and less than  $\pm 1\%$  at the rest of the basins. Hence, from a bias reduction perspective, BC-ESP was  
367 the best technique for most basins/initializations, with the exceptions of Dworshak on Feb 1 and Prineville on Mar 1  
368 and Apr 1, for which Stat. CFSR+IHC and TWS provided the best results.

## 369 5.2 Probabilistic verification

370 Figure 8 displays continuous ranked probability skill scores computed with mean climatology as a reference  
371 (CRPSS<sub>clim</sub>). Consistent with the correlation analysis results (Figure 5), better skill values are obtained for long lead  
372 times (i.e. Oct 1 and Nov 1) if climate predictors are incorporated in the forecasting framework. For example, Stat.  
373 (Ind+IHC) augments skill by 56% in HHDm1 and 7% in Hungry Horse with respect to Stat-IHC (i.e., the best  
374 benchmark in terms of CRPSS<sub>clim</sub>) when forecasts are initialized on November 1. The skill of IHC-based methods  
375 generally increases from October 1 to April 1. Nevertheless, at late initializations it is still possible to outperform these  
376 techniques in some basins (e.g., Stat (CFSR+IHC) and EWE in Hungry Horse provide skill increases of 7% and 5%  
377 in April 1 forecasts over the best IHC-based technique). For late season initializations – when IHC predictability is  
378 strong –, it is expected that climate-only forecasts underperform other methods.

379 The results from Figure 8 corroborate several findings alluded to in Section 5.1. Climate predictors applied to  
380 low-skilled (BC-)ESP forecasts in a TWS framework are less effective than when applied in a separate statistical  
381 method. Additionally, less complex multi-model schemes can perform better than more complex approaches (e.g.,  
382 BMA), supporting previous findings by Najafi and Moradkhani (2015). Among the three hybrid regression methods  
383 (Figure 3), Stat-CFSR-IHC was in most cases the worst performer. This result may be determined by the relative  
384 strength of standard (in particular ENSO) indices for the PNW region. Namely, there is less of an opportunity for  
385 custom predictor components to fill a climate influence gap, and the parameter estimation cost of the CFSR-PLSR



386 relative to an off-the-shelf index may be more exposed. It should also be noted that skill results – especially those  
387 making use of ESP output – are subject to large uncertainties due to limited sample size (i.e., only 35 years for forecast  
388 generation and verification).

389 Overall, the results presented in Figures 5 and 8 suggest a division of the study basins into two groups showing  
390 different relative predictabilities – i.e., driven by watershed conditions versus climate – from October to January. The  
391 first group is formed by Dworshak, Howard Hanson and Hungry Horse, where the state of the climate is the dominant  
392 source of predictability from Oct 1 to Dec 1, and IHCs start providing useful information on Jan 1. The second group  
393 is formed by Libby and Prineville, where little or no skill can be found from any source until Dec 1, when some  
394 predictability can be harnessed from IHCs. This is illustrated in Figure 9, where time series with cross-validated  
395 seasonal streamflow forecasts – initialized on December 1, period 1981-2015 – are shown for two IHC-based methods  
396 (BC-ESP and Stat-IHC), and two climate-based statistical methods (i.e. Stat-Ind and Stat-CFSR). At such  
397 initialization, there is not enough information in the watershed (IHCs) to predict interannual variations in April-July  
398 streamflow at Dworshak (Figure 9a) or Howard Hanson (Figure 9b); nevertheless, climate predictors are more  
399 successful, a result that is also reflected in positive correlation results (Figure 5) and skill scores (e.g., CRPSS<sub>clim</sub>  
400 increases from 0.23 with Stat-IHC to 0.39 with Stat-Ind at Howard Hanson). For the particular case of Hungry Horse  
401 (Figure 9c), some predictability is provided by watershed information alone (i.e., BC-ESP), although with smaller  
402 correlation and skill than Stat-Ind or Stat-CFSR. Finally, the ensemble forecast time series displayed for Libby (Figure  
403 9d) and Prineville (Figure 9e) portray the relative predictive power of IHCs in these basins compared to climate  
404 predictors alone. Indeed, at the December 1 initialization in these basins, watershed information alone supports  $r$   
405 values of 0.43 (Libby) and 0.39 (Prineville) from BC-ESP, and  $r$  values of 0.47 from Stat-IHC.

406 Forecast reliability can be critical to support risk-based decision making, in which actions may be tied to the  
407 forecast distribution rather than the median. The reliability index  $\alpha$  (Figure 10), which measures the closeness between  
408 the empirical CDF of forecast  $p$ -values with a theoretical CDF of  $U[0,1]$  (Table 3) shows that – although (BC-)ESP  
409 forecast correlation (Figure 5) and skill (Figure 8) generally increase during the year, forecast reliability from the ESP  
410 methods degrades (i.e. toward lower  $\alpha$ ) as the initializations approach Apr 1. Because TWS is constrained by ESP  
411 spread, it cannot provide substantial enhancements to poor late-season reliability indices obtained with (BC-)ESP.

412 In general, forecasts involving statistical calibration (which helps to improve spread and bias) are most reliable.  
413 Indeed, regression-based forecasting methods (e.g., Stat-IHC, Stat-Ind, Stat. Ind+IHC) stand out in all basins,  
414 suggesting that the ensemble generation approach used in this paper (based on the standard error of the cross-validated  
415 hindcasts) is capable of providing statistically consistent ensembles. Multi-model techniques appear to inherit this  
416 characteristic, with only small discrepancies apparent between them (green lines in Figure 10). Similar inter-method  
417 differences across multiple initializations were found when looking at the  $\epsilon$  reliability index (not shown) defined by  
418 Renard et al. (2010).

419 Although HESP was only found to be the ‘most reliable’ method in a limited number of cases (e.g.,  $\alpha = 0.95$  at  
420 Dworshak on Oct 1;  $\alpha = 0.96$  at Libby on Apr 1), relatively high  $\alpha$  values were consistently attained in all basins and  
421 forecast lead times. This suggests – in conjunction with the results shown in Figures 5-8 – that HESP has strong  
422 potential for operational streamflow forecasting at all initialization dates, since it is capable of flexibly harnessing



423 seasonally varying sources of predictability. Figure 11 illustrates this idea through time series of cross-validated  
424 ensemble forecasts obtained with HESP for three initialization times (Oct 1, Jan 1, and Apr 1). Forecasts issued on  
425 Oct 1 provide positive skill with respect to climatology in Dworshak, Howard Hanson and Hungry Horse, and although  
426 CRPSS relative to ESP does not necessarily improve, the associated correlation coefficients (0.42, 0.37 and 0.47,  
427 respectively) are a clear enhancement over negative  $r$  values obtained from IHC-based methods. The lower  
428 probabilistic skill and near-zero correlation in Libby and Prineville reflect the lack of predictability from either the  
429 watershed or climate conditions at such a long lead time. Higher values of  $CRPSS_{clim}$  for ensemble forecasts initialized  
430 on Jan 1 and Apr 1 reflect the increasing power of IHCs, while smaller (and sometimes negative)  $CRPSS_{esp}$  values in  
431 some basins reflect the increasing difficulty to outperform ESP as IHCs provide more forecast signal. Overall, HESP  
432 provides positive skill with respect to mean climatology in all cases, relatively high  $r$  values, and statistically consistent  
433 forecast ensembles.

### 434 5.3 Wet/dry year forecasts

435 Summary statistics provide an overview of forecast performance, but additional insights can be gained from  
436 exploring extreme years in the record – in which forecasts can have disproportionate value to help water managers  
437 negotiate atypical challenges – and from visualizing the behavior of the forecasting methods as individual seasons  
438 progress. We therefore performed a retrospective comparison of all techniques for two regionally wet (1997 and 2011)  
439 and dry (1987 and 2001) water years at Hungry Horse (Figure 12), one of the most teleconnected basins in our study  
440 domain. Figure 12 illustrates how SWE and SM, the primary sources of predictability for IHC-based methods,  
441 progressively gain influence on ensemble forecasts (e.g., HESP and TWS outputs) as the beginning of the snowmelt  
442 season approaches (i.e. April 1). These single-year forecast evolution plots highlight the contrast for late season (i.e.  
443 Feb 1 onwards) between overconfident predictions exhibiting poor reliability (e.g., ESP, BC-ESP, TWS), and under-  
444 confident forecasts (e.g. EWE and RWE).

445 Figure 12a,b show that climate information is required to reduce forecast errors in wet years at very long lead  
446 times (i.e., Oct 1 and Nov 1), either alone or combined with watershed information through hybrid approaches. For  
447 example, the technique that provided the smallest forecast median error on Oct. 1 1997 was TWS. For shorter lead  
448 times (i.e., forecasts initialized on March 1 or Apr 1) and WY 1997, the incorporation of IHCs helps to provide a  
449 better match with observations compared to methods that only use climate information. Interestingly, reanalysis fields  
450 at Hungry Horse provide considerable predictive power for WY 2011 (Figure 12b) at short lead times (e.g., Stat-CFSR  
451 provides a forecast median error of 2.7 % on March 1).

452 In the two dry years, Figure 12c illustrates that climate predictors alone had considerable predictive power at  
453 long lead times (i.e., Oct 1 and Nov 1) in WY 1987. However, this was not the case for WY 2001 (Figure 12d), when  
454 the method providing smallest forecast median volume errors at all initialization times (i.e., either BC-ESP or TWS)  
455 always required knowledge on watershed moisture conditions. This was also the case for other pilot study basins (not  
456 shown).

457 The above results suggest that despite the value of large-scale climate information for this study domain,  
458 enhanced hydrologic predictability is critical for accurate streamflow volumes in snowmelt-dominated regions under



459 extreme climatic conditions, especially during dry years. Past and ongoing efforts aimed to improve basin-scale  
460 meteorological forcing datasets, pursue realistic process representations in hydrologic models, advance parameter  
461 calibration, and improve DA techniques for better IHC estimates have built a robust platform to accelerate progress  
462 in this area. However, a long-term retrospective implementation (that is consistent with the real-time deployment) of  
463 these various modeling decisions and sources of information is critical to understand their performance, and  
464 benchmark methodological choices.

## 465 6 Conclusions

466 Generating accurate water supply forecasts is an ongoing challenge for improving water resources operations  
467 and planning. Despite substantial work on seasonal streamflow forecasting methods applied worldwide, the marginal  
468 value of increased complexity and combining different sources of information via different strategies has not been  
469 systematically assessed. In this paper, we compare a range of techniques that leverage predictability from watershed  
470 hydrologic conditions and/or large-scale climate information. The forecast intercomparison showed that hybrid  
471 techniques that leverage hindcasts to combine both sources of predictability could lead to improved skill compared to  
472 current operational approaches. Additional key findings that may be relevant beyond the study domain – due to the  
473 inclusion of both teleconnected and non-teleconnected basins – are as follows:

- 474 • In basins showing strong teleconnections between large-scale climate and local meteorology, the use of large-  
475 scale climate information can be an effective strategy to improve seasonal streamflow predictability,  
476 potentially providing skillful forecasts at times when watershed predictability is limited.
- 477 • Standard climate indices provide useful information, and custom climate predictors from reanalyses were  
478 also an effective complementary strategy for extracting the signal from climate fields (e.g., SST and  
479 geopotential height).
- 480 • The relative importance of watershed IHC versus climate information to predict streamflow was found to  
481 vary even within a small region, depending on sub-domain catchment hydroclimatological characteristics.
- 482 • The ESP trace weighting method only provided promising results at forecast lead times where ESP raw  
483 forecasts contained moderate skill, indicating that climate information cannot adequately shift the prior ESP  
484 forecast if it lacks forecast resolution or contains significant bias.
- 485 • Increasing methodological complexity does not necessarily translate into better ensemble forecast quality  
486 (e.g., Stat-IHC versus BC-ESP; EWE versus BMA), in part because the small sample sizes associated with  
487 seasonal hindcasts preclude reliable parameter estimation for more elaborate methods. There can be a trade-  
488 off between improving one forecast characteristic (e.g., bias) and degrading another (e.g., correlation skill).
- 489 • Cross-validation is an essential part of seasonal forecast development and implementation, particularly where  
490 multiple predictions may be combined based on their purported relative strengths and predictive uncertainty  
491 must be accurately estimated. In the small-sample context of seasonal streamflow prediction, cross-validation  
492 reveals significant limitations in the supportable complexity of statistical forecasting elements.





493 The often equivocal comparison of methods through multiple verification metrics (e.g., correlation, reliability)  
494 for individual wet and dry years, and for different basins, starkly illustrated the challenge of selecting a single method  
495 that will provide optimal results for all forecast initialization dates. There is a significant tension between optimizing  
496 forecast qualities through a mixture of methods and data sources that vary seasonally and across basins, and an oft-  
497 stated preference from forecasters and users for a consistent forecasting methodology. With this in mind, we developed  
498 HESP as a flexible data-driven framework to harness skill across varying predictability regimes, although it admittedly  
499 departs from the constraint of predictor uniformity.

500 A notable omission from this intercomparison study is the derivation of climate predictors from global climate  
501 model forecasts, a strategy that has also been pursued in this context (e.g., see Crochemore et al. 2016). The experiment  
502 summarized here did assess the skill of CFSv2 9-month climate forecasts at an earlier stage, but such evaluation has  
503 been excluded from this paper because the results did not show significantly higher skill from the CFSv2 forecasts  
504 than the CFSR-based empirical predictions, as is consistent with prior skill assessments (e.g., Yuan et al., 2011).  
505 Nonetheless, the topic of augmenting hydrologic predictability from dynamical climate forecasts remains an appealing  
506 area for future study and comparison, as does the potential for including IHC data assimilation to enhance watershed  
507 model-based predictability (e.g., Dechant and Moradkhani, 2011; Huang et al., 2016). Future work can also explore  
508 alternative methodological choices such as multiple hydrological models, different climate datasets or smaller details  
509 such as alternative variable transformations in statistical approaches (e.g., Wang et al., 2012).

510 Finally, this work is part of a larger project that explores the potential of an automated (i.e. ‘over-the-loop’)  
511 forecasting workflow as a viable strategy for operational streamflow prediction that can open the door to potential  
512 scientific and technical advances in streamflow forecasting (Pagano et al., 2016). In this context, a critical lesson is  
513 that the entire study, in particular the assessment of approach alternatives, depends on the automation of the forecast  
514 workflow to enable the generation of hindcasts that are consistent with real-time forecasts. Demonstrating that such  
515 over-the-loop methods – all of which were implemented in real-time by the authors during the study period (2015-  
516 2017) – can yield credible predictions should be regarded as a strong argument for exploring this objective paradigm  
517 in real-world operational agency settings.

## 518 7 Appendix

### 519 7.1 ESP trace weighting

520 The trace weighting scheme used here involves the following steps (Werner et al., 2004):

- 521 1. Compute a vector **D** of distances between the vector with climate predictors for the target water year ( $x_t$ ),  
522 and the vectors with predictors for the training period ( $x_i$ ):

$$523 \quad \mathbf{D} = (d_1, d_2, \dots, d_n) \quad (\text{A1})$$

$$524 \quad d_i = \|x_t - x_i\| \quad (\text{A2})$$

- 525 2. Sort the vector **D** from lowest to highest:

$$526 \quad \tilde{\mathbf{D}} = (d_{(1)}, d_{(2)}, \dots, d_{(n)}), \quad d_{(1)} \leq d_{(2)} \leq \dots \leq d_{(n)} \quad (\text{A3})$$

- 527 3. Compute weights using the following equation:





528 
$$w_i = \left[1 - \frac{d_{(i)}}{d_{(k)}}\right]^\lambda, \quad d_{(i)} \leq d_{(k)} \quad (\text{A4})$$

529 
$$w_i = 0, \quad d_{(i)} > d_{(k)} \quad (\text{A5})$$

530 
$$k = \text{NINT} \left(\frac{n}{\alpha}\right) \quad (\text{A6})$$

531 where  $\lambda$  is a distance-sensitive weighting parameter,  $\alpha$  is a parameter that influences the  $k$  nearest neighbors  
532 used, and NINT refers to the nearest integer operator. In this paper, we set  $\lambda = 2$  and  $\alpha = 1$  after conducting  
533 several experiments (not shown).

534 4. Normalize weights and construct a cumulative distribution function (CDF) based on these values and the  
535 ESP hindcast.

536 5. Resample from the CDF obtained in step 4 using 500 uniform random numbers.

## 537 7.2 BMA and QMA

538 The principle of BMA (Raftery et al., 2005) is that given an ensemble forecast with  $M$  members, each ensemble  
539 member  $f_i$  ( $i = 1, 2, \dots, M$ ) is associated with a conditional PDF  $h_i(y|f_i)$ , which can be interpreted as the PDF of the  
540 variable  $y$  given  $f_i$ . Thus, the BMA predictive model is:

541 
$$p(y|f_1, \dots, f_M) = \sum_{i=1}^M w_i h_i(y|f_i) \quad (\text{A7})$$

542 where the BMA weight  $w_i$  is the posterior probability of forecast  $i$  and is obtained based on its relative  
543 performance during the training period. Therefore, the weights  $w_i$ 's are nonnegative and add up to 1, i.e.  $\sum_{i=1}^M w_i = 1$   
544 (Raftery et al., 2005).

545 In this paper, the weights for the two models (best climate-based and best watershed-based) are estimated by  
546 maximum likelihood, assuming that the conditional PDFs of  $\log(Q)$  are approximated by a normal distribution. The  
547 likelihood is maximized using the expectation-maximization (EM) algorithm (Dempster et al., 1977) which is  
548 implemented in the R package ensembleBMA ([https://cran.r-](https://cran.r-project.org/web/packages/ensembleBMA/ensembleBMA.pdf)  
549 [project.org/web/packages/ensembleBMA/ensembleBMA.pdf](http://www.rproject.org/)) at the public domain statistical software R  
550 (<http://www.rproject.org/>). Prior information (i.e., initial weights) is provided by weights computed as  $1/\text{RMSE}$ .  
551 Finally, the BMA forecast ensemble is obtained by sampling a fraction of members from each model equal to the  
552 weight  $w_i$ .

553 The quantile model averaging (QMA) forecast values are obtained from the weighted average of forecast  
554 quantiles from all models. Schepen and Wang (2015) recently found that nearly identical skill results can be obtained  
555 with BMA and QMA, and that very similar performance can be achieved either by calibrating QMA weights or by  
556 using BMA weights within a QMA framework. Therefore, we obtain the QMA forecast using the same weights  
557 obtained from the BMA calibration, by sorting the ensemble members from the best climate and best watershed  
558 forecast approaches, and computing the weighted average of equally ranked ensemble members from the two sources.



559 **8 Acknowledgments**

560 This work was supported through a contract with the U.S. Army Corps of Engineers, and through a Cooperative  
561 Agreement with the U.S. Bureau of Reclamation.

562 **9 References**

- 563 Abdi, H.: Partial least squares regression and projection on latent structure regression, *Wiley Interdiscip. Rev. Comput.*  
564 *Stat.*, 2, 97–106, doi:10.1002/wics.051, 2010.
- 565 Akaike, H.: A new look at the statistical model identification, *IEEE Trans. Automat. Contr.*, 19(6), 716–723,  
566 doi:10.1109/TAC.1974.1100705, 1974.
- 567 Anderson, E.: National Weather Service River Forecast system - snow accumulation and ablation model, NOAA Tech.  
568 Memo. NWS HYDRO-17, 217, 1973.
- 569 Beckers, J. V. L. V. L., Weerts, A. H. H., Tjeldeman, E. and Welles, E.: ENSO-conditioned weather resampling method  
570 for seasonal ensemble streamflow prediction, *Hydrol. Earth Syst. Sci.*, 20(8), 3277–3287, doi:10.5194/hess-20-3277-  
571 2016, 2016.
- 572 BPA: 2010 Level Modified Streamflow: 1928-2008., 2011.
- 573 Bracken, C., Rajagopalan, B. and Prairie, J.: A multisite seasonal ensemble streamflow forecasting technique, *Water*  
574 *Resour. Res.*, 46, W03532, doi:10.1029/2009WR007965, 2010.
- 575 Bradley, A. A., Habib, M. and Schwartz, S. S.: Climate index weighting of ensemble streamflow forecasts using a  
576 simple Bayesian approach, *Water Resour. Res.*, 51(9), 7382–7400, doi:10.1002/2014WR016811, 2015.
- 577 Burnash, R., Ferral, R. and McGuire, R.: A generalized streamflow simulation system - Conceptual modeling for  
578 digital computers, Sacramento, California., 1973.
- 579 Clark, M. P., Serreze, M. C. and McCabe, G. J.: Historical effects of El Nino and La Nina events on the seasonal  
580 evolution of the montane snowpack in the Columbia and Colorado River Basins, *Water Resour. Res.*, 37(3), 741–757,  
581 doi:10.1029/2000WR900305, 2001.
- 582 Crochemore, L., Ramos, M.-H. and Pappenberger, F.: Bias correcting precipitation forecasts to improve the skill of  
583 seasonal streamflow forecasts, *Hydrol. Earth Syst. Sci.*, 20(2002), 3601–3618, doi:10.5194/hess-20-3601-2016, 2016.
- 584 Day, G. N.: Extended Streamflow Forecasting Using NWSRFS, *J. Water Resour. Plan. Manag.*, 111(2), 157–170,  
585 doi:10.1061/(ASCE)0733-9496(1985)111:2(157), 1985.
- 586 Dechant, C. M. and Moradkhani, H.: Improving the characterization of initial condition for ensemble streamflow  
587 prediction using data assimilation, *Hydrol. Earth Syst. Sci.*, 15(11), 3399–3410, doi:10.5194/hess-15-3399-2011,  
588 2011.
- 589 Demargne, J., Wu, L., Regonda, S. K., Brown, J. D., Lee, H., He, M., Seo, D. J., Hartman, R., Herr, H. D., Fresch,  
590 M., Schaake, J. and Zhu, Y.: The science of NOAA’s operational hydrologic ensemble forecast service, *Bull. Am.*  
591 *Meteorol. Soc.*, 95(1), 79–98, doi:10.1175/BAMS-D-12-00081.1, 2014.
- 592 Dempster, A., Laird, N. and Rubin, D.: Maximum likelihood from incomplete data via the EM algorithm, *J. R. Stat.*  
593 *Soc.*, 39(1), 1–38, 1977.
- 594 Devineni, N., Sankarasubramanian, A. and Ghosh, S.: Multimodel ensembles of streamflow forecasts: Role of



- 595 predictor state in developing optimal combinations., *Water Resour. Res.*, 44, W09404, doi:10.1029/2006WR005855,  
596 2008.
- 597 Duan, Q., Ajami, N. K., Gao, X. and Sorooshian, S.: Multi-model ensemble hydrologic prediction using Bayesian  
598 model averaging, *Adv. Water Resour.*, 30(5), 1371–1386, doi:10.1016/j.advwatres.2006.11.014, 2007.
- 599 Garen, D. C.: Improved Techniques in Regression-Based Streamflow Volume Forecasting, *J. Water Resour. Plan.*  
600 *Manag.*, 118(6), 654–670, doi:10.1061/(ASCE)0733-9496(1992)118:6(654), 1992.
- 601 Georgakakos, K. P., Seo, D.-J., Gupta, H., Schaake, J. and Butts, M. B.: Towards the characterization of streamflow  
602 simulation uncertainty through multimodel ensembles, *J. Hydrol.*, 298, 222–241, doi:10.1016/j.jhydrol.2004.03.037,  
603 2004.
- 604 Gobena, A. K. and Gan, T. Y.: Incorporation of seasonal climate forecasts in the ensemble streamflow prediction  
605 system, *J. Hydrol.*, 385(1–4), 336–352, doi:10.1016/j.jhydrol.2010.03.002, 2010.
- 606 Grantz, K., Rajagopalan, B., Clark, M. and Zagona, E.: A technique for incorporating large-scale climate information  
607 in basin-scale ensemble streamflow forecasts, *Water Resour. Res.*, 41, W10410, doi:10.1029/2004WR003467, 2005.
- 608 Hagedorn, R., Doblas-Reyes, F. and Palmer, T.: The rationale behind the success of multi-model ensembles in seasonal  
609 forecasting - I. Basic concept, *Tellus A*, 57(3), 219–233, doi:10.1111/j.1600-0870.2005.00103.x, 2005.
- 610 Hamlet, A. F. and Lettenmaier, D. P.: Columbia River Streamflow Forecasting Based on ENSO and PDO Climate  
611 Signals, *J. Water Resour. Plan. Manag.*, 125(6), 333–341, doi:10.1061/(ASCE)0733-9496(1999)125:6(333), 1999.
- 612 Harrison, B. and Bales, R.: Skill Assessment of Water Supply Forecasts for Western Sierra Nevada Watersheds, *J.*  
613 *Hydrol. Eng.*, 21(4), 04016002, doi:10.1061/(ASCE)HE.1943-5584.0001327, 2016.
- 614 Hersbach, H.: Decomposition of the Continuous Ranked Probability Score for Ensemble Prediction Systems, *Weather*  
615 *Forecast.*, 15(5), 559–570, doi:10.1175/1520-0434(2000)015<0559:DOTCRP>2.0.CO;2, 2000.
- 616 Huang, C., Newman, A. J., Clark, M. P., Wood, A. W. and Zheng, X.: Evaluation of snow data assimilation using the  
617 Ensemble Kalman Filter for seasonal streamflow prediction in the Western United States, *Hydrol. Earth Syst. Sci.*  
618 *Discuss.*, (May), 1–29, doi:10.5194/hess-2016-185, 2016.
- 619 Laio, F. and Tamea, S.: Verification tools for probabilistic forecasts of continuous hydrological variables, *Hydrol.*  
620 *Earth Syst. Sci.*, 11(4), 1267–1277, doi:10.5194/hess-11-1267-2007, 2007.
- 621 Luo, L. and Wood, E. F.: Use of Bayesian Merging Techniques in a Multimodel Seasonal Hydrologic Ensemble  
622 Prediction System for the Eastern United States, *J. Hydrometeorol.*, 9(5), 866–884, doi:10.1175/2008JHM980.1,  
623 2008.
- 624 Mantua, N. J., Hare, S. R., Zhang, Y., Wallace, J. M. and Francis, R. C.: A Pacific Interdecadal Climate Oscillation  
625 with Impacts on Salmon Production, *Bull. Am. Meteorol. Soc.*, 78(6), 1069–1079, doi:10.1175/1520-  
626 0477(1997)078<1069:APICOW>2.0.CO;2, 1997.
- 627 Maurer, E. P., Lettenmaier, D. P. and Mantua, N. J.: Variability and potential sources of predictability of North  
628 American runoff, *Water Resour. Res.*, 40(9), W09306, doi:10.1029/2003WR002789, 2004.
- 629 McCabe, G. J. and Dettinger, M. D.: Primary Modes and Predictability of Year-to-Year Snowpack Variations in the  
630 Western United States from Teleconnections with Pacific Ocean Climate, *J. Hydrometeorol.*, 3(1), 13–25,  
631 doi:10.1175/1525-7541(2002)003<0013:PMAPOY>2.0.CO;2, 2002.
- 632 Mendoza, P. A., Rajagopalan, B., Clark, M. P., Cortés, G. and McPhee, J.: A robust multimodel framework for  
633 ensemble seasonal hydroclimatic forecasts, *Water Resour. Res.*, 50(7), 6030–6052, doi:10.1002/2014WR015426,



- 634 2014.
- 635 Moradkhani, H. and Meier, M.: Long-Lead Water Supply Forecast Using Large-Scale Climate Predictors and  
636 Independent Component Analysis, *J. Hydrol. Eng.*, 15(10), 744–762, doi:10.1061/(ASCE)HE.1943-5584.0000246,  
637 2010.
- 638 Mote, P. W.: Trends in snow water equivalent in the Pacific Northwest and their climatic causes, *Geophys. Res. Lett.*,  
639 30(12)(L1601), 1–4, doi:10.1029/2003GL017258, 2003.
- 640 Najafi, M. and Moradkhani, H.: Ensemble Combination of Seasonal Streamflow Forecasts, *J. Hydrol. Eng.*, (2001),  
641 4015043, doi:10.1061/(ASCE)HE.1943-5584.0001250, 2015.
- 642 Najafi, M. R., Moradkhani, H. and Piechota, T. C.: Ensemble Streamflow Prediction: Climate signal weighting  
643 methods vs. Climate Forecast System Reanalysis, *J. Hydrol.*, 442-443, 105–116, doi:10.1016/j.jhydrol.2012.04.003,  
644 2012.
- 645 Newman, A. J., Clark, M. P., Craig, J., Nijssen, B., Wood, A., Gutmann, E., Mizukami, N., Brekke, L. and Arnold, J.  
646 R.: Gridded Ensemble Precipitation and Temperature Estimates for the Contiguous United States, *J. Hydrometeorol.*,  
647 16(6), 2481–2500, doi:10.1175/JHM-D-15-0026.1, 2015.
- 648 Opitz-Stapleton, S., Gangopadhyay, S. and Rajagopalan, B.: Generating streamflow forecasts for the Yakima River  
649 Basin using large-scale climate predictors, *J. Hydrol.*, 341(3-4), 131–143, doi:10.1016/j.jhydrol.2007.03.024, 2007.
- 650 Pagano, T., Garen, D. and Sorooshian, S.: Evaluation of Official Western U.S. Seasonal Water Supply Outlooks,  
651 1922–2002, *J. Hydrometeorol.*, 5(5), 896–909, doi:10.1175/1525-7541(2004)005<0896:EOOWUS>2.0.CO;2, 2004.
- 652 Pagano, T. C., Pappenberger, F., Wood, A. W., Ramos, M., Persson, A. and Anderson, B.: Automation and human  
653 expertise in operational river forecasting, *Wiley Interdiscip. Rev. Water*, (June), doi:10.1002/wat2.1163, 2016.
- 654 Piechota, T. C., Dracup, J. A. and Fovell, R. G.: Western US streamflow and atmospheric circulation patterns during  
655 El Niño-Southern Oscillation, *J. Hydrol.*, 201, 249–271, doi:10.1016/s0022-1694(97)00043-7, 1997.
- 656 Piechota, T. C., Chiew, F. H. S., Dracup, J. A. and McMahon, T. A.: Seasonal streamflow forecasting in eastern  
657 Australia and the El Niño-Southern Oscillation, *Water Resour. Res.*, 34(11), 3035–3044, doi:10.1029/98WR02406,  
658 1998.
- 659 Plummer, N., Tuteja, N., Wang, Q., Wang, E., Robertson, D., Zhou, S., Schepen, A., Alves, O., Timbal, B. and Puri,  
660 K.: A Seasonal Water Availability Prediction Service: Opportunities and Challenges, in 18th World IMACS /  
661 MODSIM Congress, pp. 1–15, Cairns, Australia., 2009.
- 662 Raftery, A. E., Gneiting, T., Balabdaoui, F. and Polakowski, M.: Using Bayesian Model Averaging to Calibrate  
663 Forecast Ensembles, *Mon. Weather Rev.*, 133(5), 1155–1174, doi:10.1175/MWR2906.1, 2005.
- 664 Redmond, K. T. and Koch, R. W.: Surface Climate and Streamflow Variability in the Western United States and Their  
665 Relationship to Large-Scale Circulation Indices, *Water Resour. Res.*, 27(9), 2381–2399, doi:10.1029/91WR00690,  
666 1991.
- 667 Regonda, S. K., Rajagopalan, B., Clark, M. and Zagona, E.: A multimodel ensemble forecast framework: Application  
668 to spring seasonal flows in the Gunnison River Basin, *Water Resour. Res.*, 42, W09404, doi:10.1029/2005WR004653,  
669 2006.
- 670 Renard, B., Kavetski, D., Kuczera, G., Thyer, M. and Franks, S. W.: Understanding predictive uncertainty in  
671 hydrologic modeling: The challenge of identifying input and structural errors, *Water Resour. Res.*, 46(5), W05521,  
672 doi:10.1029/2009WR008328, 2010.



- 673 Robertson, D. E., Pokhrel, P. and Wang, Q. J.: Improving statistical forecasts of seasonal streamflows using  
674 hydrological model output, *Hydrol. Earth Syst. Sci.*, 17(2), 579–593, doi:10.5194/hess-17-579-2013, 2013.
- 675 Rosenberg, E. A., Wood, A. W. and Steinemann, A. C.: Statistical applications of physically based hydrologic models  
676 to seasonal streamflow forecasts, *Water Resour. Res.*, 47(3), W00H14, doi:10.1029/2010WR010101, 2011.
- 677 Saha, S., Moorthi, S., Pan, H.-L., Wu, X., Wang, J., Nadiga, S., Tripp, P., Kistler, R., Woollen, J., Behringer, D., Liu,  
678 H., Stokes, D., Grumbine, R., Gayno, G., Wang, J., Hou, Y.-T., Chuang, H.-Y., Juang, H.-M. H., Sela, J., Iredell, M.,  
679 Treadon, R., Kleist, D., Van Delst, P., Keyser, D., Derber, J., Ek, M., Meng, J., Wei, H., Yang, R., Lord, S., Van Den  
680 Dool, H., Kumar, A., Wang, W., Long, C., Chelliah, M., Xue, Y., Huang, B., Schemm, J.-K., Ebisuzaki, W., Lin, R.,  
681 Xie, P., Chen, M., Zhou, S., Higgins, W., Zou, C.-Z., Liu, Q., Chen, Y., Han, Y., Cucurull, L., Reynolds, R. W.,  
682 Rutledge, G. and Goldberg, M.: The NCEP Climate Forecast System Reanalysis, *Bull. Am. Meteorol. Soc.*, 91(8),  
683 1015–1057, doi:10.1175/2010BAMS3001.1, 2010.
- 684 Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y.-T., Chuang, H., Iredell, M., Ek,  
685 M., Meng, J., Yang, R., Mendez, M. P., van den Dool, H., Zhang, Q., Wang, W., Chen, M. and Becker, E.: The NCEP  
686 Climate Forecast System Version 2, *J. Clim.*, 27(6), 2185–2208, doi:10.1175/JCLI-D-12-00823.1, 2014.
- 687 Schepen, A. and Wang, Q. J.: Model averaging methods to merge operational statistical and dynamic seasonal  
688 streamflow forecasts in Australia, *Water Resour. Res.*, 6(4), 1–16, doi:10.1002/2014WR016163, 2015.
- 689 Smith, J. A., Day, G. N. and Kane, M. D.: Nonparametric Framework for Long-range Streamflow Forecasting, *J.*  
690 *Water Resour. Plan. Manag.*, 118(1), 82–92, doi:10.1061/(ASCE)0733-9496(1992)118:1(82), 1992.
- 691 Tootle, G. A., Singh, A. K., Piechota, T. C. and Farnham, I.: Long Lead-Time Forecasting of U.S. Streamflow Using  
692 Partial Least Squares Regression, *J. Hydrol. Eng.*, 12(5), 442–451, doi:10.1061/(ASCE)1084-0699(2007)12:5(442),  
693 2007.
- 694 Wang, Q. J., Robertson, D. E. and Chiew, F. H. S.: A Bayesian joint probability modeling approach for seasonal  
695 forecasting of streamflows at multiple sites, *Water Resour. Res.*, 45(5), 1–18, doi:10.1029/2008WR007355, 2009.
- 696 Wang, Q. J., Shrestha, D. L., Robertson, D. E. and Pokhrel, P.: A log-sinh transformation for data normalization and  
697 variance stabilization, *Water Resour. Res.*, 48(5), 1–7, doi:10.1029/2011WR010973, 2012.
- 698 Weber, F., Garen, D. and Gobena, A.: Invited commentary: themes and issues from the workshop “Operational river  
699 flow and water supply forecasting,” *Can. Water Resour. J.*, 37(3), 151–161, doi:10.4296/cwrj2012-953, 2012.
- 700 Werner, K., Brandon, D., Clark, M. and Gangopadhyay, S.: Climate Index Weighting Schemes for NWS ESP-Based  
701 Seasonal Volume Forecasts, *J. Hydrometeorol.*, 5(6), 1076–1090, doi:10.1175/JHM-381.1, 2004.
- 702 Wold, H.: Estimation of principal components and related models by iterative least squares, in *Multivariate Analysis*,  
703 edited by P. R. Krishnaia, pp. 391–420, Academic Press, New York., 1966.
- 704 Wood, A. W. and Lettenmaier, D. P.: A Test Bed for New Seasonal Hydrologic Forecasting Approaches in the  
705 Western United States, *Bull. Am. Meteorol. Soc.*, 87(12), 1699–1712, doi:10.1175/BAMS-87-12-1699, 2006.
- 706 Wood, A. W. and Lettenmaier, D. P.: An ensemble approach for attribution of hydrologic prediction uncertainty,  
707 *Geophys. Res. Lett.*, 35(14), L14401, doi:10.1029/2008GL034648, 2008.
- 708 Wood, A. W. and Schaake, J. C.: Correcting Errors in Streamflow Forecast Ensemble Mean and Spread, *J.*  
709 *Hydrometeorol.*, 9(1), 132–148, doi:10.1175/2007JHM862.1, 2008.
- 710 Wood, A. W., Kumar, A. and Lettenmaier, D. P.: A retrospective assessment of National Centers for Environmental  
711 prediction climate model-based ensemble hydrologic forecasting in the western United States, *J. Geophys. Res. D*



- 712 Atmos., 110(4), 1–16, doi:10.1029/2004JD004508, 2005.
- 713 Yossef, N. C., Winsemius, H., Weerts, A., Van Beek, R. and Bierkens, M. F. P.: Skill of a global seasonal streamflow  
714 forecasting system, relative roles of initial conditions and meteorological forcing, *Water Resour. Res.*, 49(8), 4687–  
715 4699, doi:10.1002/wrcr.20350, 2013.
- 716 Yuan, X., Wood, E. F., Luo, L. and Pan, M.: A first look at Climate Forecast System version 2 (CFSv2) for  
717 hydrological seasonal prediction, *Geophys. Res. Lett.*, 38(13), 1–7, doi:10.1029/2011GL047792, 2011.
- 718 Yuan, X., Wood, E. F., Roundy, J. K. and Pan, M.: CFSv2-Based seasonal hydroclimatic forecasts over the  
719 conterminous United States, *J. Clim.*, 26(13), 4828–4847, doi:10.1175/JCLI-D-12-00683.1, 2013.
- 720
- 721



722 **10 List of Figures**

723 Figure 1: Location map with the pilot basins included in this study. ....26

724 Figure 2: Corrected precipitation P (i.e. observed precipitation multiplied by a snow correction factor SCF) and  
 725 simulated water balance variables—active SM, SWE, and runoff (RO)—for the five study basins: (a) Dworshak  
 726 Reservoir inflow (DWRI1), (b) Howard Hanson reservoir inflow (HHDW1), (c) Hungry Horse reservoir inflow  
 727 (HHWM8), (d) Libby dam inflow (LYDM8), and (e) Prineville reservoir inflows (PRVO). For model SM, we subtract  
 728 the lowest mean monthly value of the year so that the plotted values show only the active range of variation. ....27

729 Figure 3: Schematic figure showing all seasonal streamflow forecasting methods included in the inter-comparison  
 730 framework. The benchmark methods are operationally implemented in the Western United States, and they are solely  
 731 based on hydrologic predictability.....28

732 Figure 4: Monthly streamflow simulations (red) and observations (black) for the period Oct/1980 – Sep/2000. Left  
 733 panels display monthly time series, with NSE and  $r$  denoting the Nash-Sutcliffe efficiency and correlation,  
 734 respectively. Right panels show simulated and observed seasonal streamflow cycles. Results are displayed for (a)  
 735 Dworshak Reservoir inflow (DWRI1); (b) Howard Hanson reservoir inflow (HHDW1); (c) Hungry Horse reservoir  
 736 inflow (HHWM8); (d) Libby dam inflow (LYDM8); and (e) Prineville reservoir inflows (PRVO). ....29

737 Figure 5: Correlation coefficients of forecast ensemble medians versus observations obtained from all methods at  
 738 different initialization dates. The error bars define 95% confidence limits obtained through bootstrapping with  
 739 replacement. Results are displayed for (a) Dworshak Reservoir inflow (DWRI1); (b) Howard Hanson reservoir inflow  
 740 (HHDW1); (c) Hungry Horse reservoir inflow (HHWM8); (d) Libby dam inflow (LYDM8); and (e) Prineville  
 741 reservoir inflows (PRVO). ....30

742 Figure 6: Same as in Figure 5, but for root mean squared error (RMSE) of ensemble forecast medians versus  
 743 observations. See text for further details. ....31

744 Figure 7: Same as in Figure 5, but for percent bias (% bias) in forecast ensemble medians versus observations. See  
 745 text for further details. ....32

746 Figure 8: Continuous Ranked Probability Skill Score of the forecast ensembles with respect to mean observed  
 747 climatology ( $CRPSS_{clim}$ ). See text for further details. ....33

748 Figure 9: Time series with cross-validated hindcasts initialized on December 1, obtained with two watershed-based  
 749 methods (BC-ESP and Stat-IHC) and two climate-based techniques (Stat-Ind and Stat-CFSR) for the five case study  
 750 locations (a-e). The verification metrics  $CRPSS_{clim}$  and  $CRPSS_{esp}$  denote continuous ranked probability skill scores  
 751 using the mean climatology and raw ESP output as the reference, respectively. Black dashed lines represent 10%, 50%  
 752 and 90% flows from the observed climatology, and boxplots show the 10<sup>th</sup>, 30<sup>th</sup>, 50<sup>th</sup>, 70<sup>th</sup> and 90<sup>th</sup> hindcast percentiles.  
 753 .....34

754 Figure 10: The  $\alpha$  reliability index for the hindcast ensembles for five case study locations. See text for further details.  
 755 .....35

756 Figure 11: Time series with cross-validated hindcasts obtained with the Hierarchical Ensemble Streamflow Prediction  
 757 (HESP) approach, initialized on (left) October 1, (center) January 1, and (right) April 1. Results are displayed for the  
 758 five case study locations: (a) Dworshak Reservoir inflow (DWRI1); (b) Howard Hanson reservoir inflow (HHDW1);





759	(c) Hungry Horse reservoir inflow (HHWM8); (d) Libby dam inflow (LYDM8); and (e) Prineville reservoir inflows	
760	(PRVO). Black dashed lines represent 10%, 50% and 90% flows from the observed climatology, and boxplots show	
761	the 10 <sup>th</sup> , 30 <sup>th</sup> , 50 <sup>th</sup> , 70 <sup>th</sup> and 90 <sup>th</sup> hindcast percentiles. ....	36
762	Figure 12: April-July water supply forecasts obtained at the Hungry Horse reservoir (HHWM8) with different methods	
763	for two wet years – (a) 1997, and (b) 2011 – and two dry years – (c) 1987, and (d) 2001. The red dashed line represents	
764	the observed flow, while black dashed lines represent 10%, 50% and 90% flows from observed climatology, and	
765	boxplots show the 10 <sup>th</sup> , 30 <sup>th</sup> , 50 <sup>th</sup> , 70 <sup>th</sup> and 90 <sup>th</sup> hindcast percentiles. ....	37
766		



767 **Table 1: List of basin characteristics. Hydrologic variables correspond to the period October 1980 to September 2015. P,**  
 768 **R, PE, RR, and DI denote basin-averaged mean annual values of precipitation, runoff, potential evapotranspiration, runoff**  
 769 **ratio, and dryness index, respectively.**

	Dworshak	Howard Hanson	Hungry Horse	Libby	Prineville
Symbol	DWRI1	HHDW1	HHWM8	LYDM8	PRVO
Area (km <sup>2</sup> )	6300	570	4200	23270	6825
Basin average elevation (m.a.s.l.)	1290	905	1773	1648	1301
Mean annual precipitation, P (mm/yr)	1182	1890	1043	813	349
Mean annual runoff, R (mm/yr)	761	1483	676	408	47
Mean annual PE* (mm/yr)	1362	1191	1272	990	1338
Mean annual RE (R/P)	0.64	0.78	0.65	0.50	0.13
Mean annual DI (PE/P)	1.15	0.63	1.22	1.22	3.83

770 \*Potential evapotranspiration using the Priestley-Taylor method

771

772

773

774 **Table 2: List of climate indices included as potential predictors**

Index	Pattern
Niño 3.4	East Central Tropical Pacific sea surface temperature (SST)
Niño 1+2	Extreme Eastern Tropical Pacific SST
Niño 3	Eastern Tropical Pacific SST
Niño 4	Central Tropical Pacific SST
AMO	Atlantic Multidecadal Oscillation
NAO	North Atlantic Oscillation
PDO	Pacific Decadal Oscillation
PNA	Pacific North American Index
SOI	Southern Oscillation Index
MEI	Multivariate ENSO index
WP	Western Pacific Index
TNA	Tropical Northern Atlantic Index

775



776 **Table 3: Performance metrics used to assess and compare seasonal streamflow forecasting methods.**

Notation	Name	Equation	Description
$r$	Correlation coefficient	$r = \frac{\sum_{i=1}^N (q_{m,i} - \bar{q}_m)(o_i - \bar{o})}{\sqrt{\sum_{i=1}^N (q_{m,i} - \bar{q}_m)^2} \sqrt{\sum_{i=1}^N (o_i - \bar{o})^2}}$	Deterministic metric that varies [-1,1] with a perfect score of 1. It measures the linear association between forecasts and observations independent of the mean and variance of the marginal distributions.
%Bias	Percent bias	$\%Bias = \frac{\sum_{i=1}^N (q_{m,i} - o_i)}{\sum_{i=1}^N o_i} \times 100$	Deterministic metric that varies $(-\infty, \infty)$ , with perfect score of 0. It measures the difference between the mean of the forecasts and the mean of observations.
RMSE	Root mean squared error	$RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (q_{m,i} - o_i)^2}$	Deterministic metric that varies $[0, \infty)$ , with perfect score of 0.
CRPSS	Continuous ranked probability skill score	$CRPSS = 1 - \frac{CRPS_{fcst}}{CRPS_{ref}}$ $CRPS = \frac{1}{N} \sum_{i=1}^N \int_{-\infty}^{\infty} [F(q) - F_o(q)]^2 dq$ $F_o(q) = \begin{cases} 0, & q < o \\ 1, & q \geq o \end{cases}$	Probabilistic metric that varies $(-\infty, 1]$ , with perfect score of 1. It measures the skill of CRPS relative to a reference forecast (Hersbach, 2000). CRPS quantifies the difference between the cumulative distribution (CDF) function of a forecast ( $F$ ), and the corresponding CDF of the observations ( $F_o$ ).
$\alpha$	$\alpha$ reliability index	$\alpha = 1 - 2 \left[ \frac{1}{N} \sum_{i=1}^N  P_i(o_i) - U(o_i)  \right]$	Probabilistic metric that varies $[0, 1]$ . It quantifies the closeness between the empirical CDF of sample p-values with the CDF of a uniform distribution. A value of 0 is the worst, and 1 reflects perfect reliability (Renard et al., 2010).

777  $q_{m,i}$ : Forecast ensemble median for year  $i$ .

778  $\bar{q}_m$ : Temporal average over forecast ensemble medians.

779  $o_i$ : Observation for year  $i$ .


780  $\bar{o}$ : Temporal average of observations.

781  $P_i(o_i)$ : Non-exceedance probability of  $o_i$  using ensemble forecasts at year  $i$ .


782  $U_i(o_i)$ : Non-exceedance probability of  $o_i$  using the uniform distribution  $U[0,1]$ .

783

784




785  
786 **Figure 1: Location map with the pilot basins included in this study.**  
787  
788



789

790 Figure 2: Corrected precipitation P (i.e. observed precipitation multiplied by a snow correction factor SCF) and simulated  
 791 water balance variables—active SM, SWE, and runoff (RO)—for the five study basins: (a) Dworshak Reservoir inflow  
 792 (DWRI1), (b) Howard Hanson reservoir inflow (HDDW1), (c) Hungry Horse reservoir inflow (HHWM8), (d) Libby dam  
 793 inflow (LYDM8), and (e) Prineville reservoir inflows (PRVO). For model SM, we subtract the lowest mean monthly value  
 794 of the year so that the plotted values show only the active range of variation.




795

796 **Figure 3: Schematic figure showing all seasonal streamflow forecasting methods included in the inter-comparison**  
 797 **framework. The benchmark methods are operationally implemented in the Western United States, and they are solely**  
 798 **based on hydrologic predictability.**

799

800




801

802 **Figure 4: Monthly streamflow simulations (red) and observations (black) for the period Oct/1980 – Sep/2000. Left panels**  
 803 **display monthly time series, with NSE and  $r$  denoting the Nash-Sutcliffe efficiency and correlation, respectively. Right**  
 804 **panels show simulated and observed seasonal streamflow cycles. Results are displayed for (a) Dworshak Reservoir inflow**  
 805 **(DWRI1); (b) Howard Hanson reservoir inflow (HHDW1); (c) Hungry Horse reservoir inflow (HHWM8); (d) Libby dam**  
 806 **inflow (LYDM8); and (e) Prineville reservoir inflows (PRVO).**


807





808  
 809  
 810  
 811  
 812


**Figure 5: Correlation coefficients of forecast ensemble medians versus observations obtained from all methods at different initialization dates. The error bars define 95% confidence limits obtained through bootstrapping with replacement. Results are displayed for (a) Dworshak Reservoir inflow (DWR11); (b) Howard Hanson reservoir inflow (HHDW1); (c) Hungry Horse reservoir inflow (HHWM8); (d) Libby dam inflow (LYDM8); and (e) Prineville reservoir inflows (PRVO).**



813

814 **Figure 6:** Same as in Figure 5, but for root mean squared error (RMSE) of ensemble forecast medians versus observations.  
 815 See text for further details.

816




817

818 **Figure 7: Same as in Figure 5, but for percent bias (% bias) in forecast ensemble medians versus observations. See text for**  
 819 **further details.**


820





821

822 **Figure 8: Continuous Ranked Probability Skill Score of the forecast ensembles with respect to mean observed climatology**  
 823 **(CRPSS<sub>clim</sub>). See text for further details.**



824


825

826

827

828


**Figure 9:** Time series with cross-validated hindcasts initialized on December 1, obtained with two watershed-based methods (BC-ESP and Stat-IHC) and two climate-based techniques (Stat-Ind and Stat-CFSR) for the five case study locations (a-e). The verification metrics CRPSS<sub>clim</sub> and CRPSS<sub>esp</sub> denote continuous ranked probability skill scores using the mean climatology and raw ESP output as the reference, respectively. Black dashed lines represent 10%, 50% and 90% flows from the observed climatology, and boxplots show the 10<sup>th</sup>, 30<sup>th</sup>, 50<sup>th</sup>, 70<sup>th</sup> and 90<sup>th</sup> hindcast percentiles.



829

830 **Figure 10: The  $\alpha$  reliability index for the hindcast ensembles for five case study locations. See text for further details.**





831


832

833 **Figure 11: Time series with cross-validated hindcasts obtained with the Hierarchical Ensemble Streamflow Prediction**  
 834 **(HESP) approach, initialized on (left) October 1, (center) January 1, and (right) April 1. Results are displayed for the five**  
 835 **case study locations: (a) Dworshak Reservoir inflow (DWR11); (b) Howard Hanson reservoir inflow (HHDW1); (c) Hungry**  
 836 **Horse reservoir inflow (HHWM8); (d) Libby dam inflow (LYDM8); and (e) Prineville reservoir inflows (PRVO). Black**  
 837 **dashed lines represent 10%, 50% and 90% flows from the observed climatology, and boxplots show the 10<sup>th</sup>, 30<sup>th</sup>, 50<sup>th</sup>, 70<sup>th</sup>**  
**and 90<sup>th</sup> hindcast percentiles.**

838

839





840

841 **Figure 12: April-July water supply forecasts obtained at the Hungry Horse reservoir (HHWM8) with different methods for**  
 842 **two wet years – (a) 1997, and (b) 2011 – and two dry years – (c) 1987, and (d) 2001. The red dashed line represents the**  
 843 **observed flow, while black dashed lines represent 10%, 50% and 90% flows from observed climatology, and boxplots show**  
 844 **the 10<sup>th</sup>, 30<sup>th</sup>, 50<sup>th</sup>, 70<sup>th</sup> and 90<sup>th</sup> hindcast percentiles.**

845