

Article

An $M^{[X]}/G(a, b)/1$ Queueing System with Breakdown and Repair, Stand-By Server, Multiple Vacation and Control Policy on Request for Re-Service

G. Ayyappan  and S. Karpagam * 

Department of Mathematics, Pondicherry Engineering College, Puducherry 605014, India; ayyappan@pec.edu

* Correspondence: karpagammaths19@gmail.com

Received: 29 March 2018; Accepted: 29 May 2018; Published: 14 June 2018

Abstract: In this paper, we discuss a non-Markovian batch arrival general bulk service single-server queueing system with server breakdown and repair, a stand-by server, multiple vacation and re-service. The main server's regular service time, re-service time, vacation time and stand-by server's service time are followed by general distributions and breakdown and repair times of the main server with exponential distributions. There is a stand-by server which is employed during the period in which the regular server remains under repair. The probability generating function of the queue size at an arbitrary time and some performance measures of the system are derived. Extensive numerical results are also illustrated.

Keywords: non-Markovian queue; general bulk service; multiple vacation; breakdown and repair; stand-by server; re-service

Mathematics Subject Classification: 60K25; 90B22; 68M20

1. Introduction

Queueing systems with general bulk service and vacations have been studied by many researchers because they deal with effective utilization of the server's idle time for secondary jobs. Such queueing systems have a wide range of application in many real-life situations such as production line systems, inventory systems, digital communications and computer networks. Doshi [1] and Takagi [2] have made a comprehensive survey of queueing systems with vacations. A batch arrival $M^{[X]}/G/1$ queueing system with multiple vacations was first studied by Baba [3]. Krishna Reddy et al. [4] have discussed an $M^{[X]}/G(a, b)/1$ model with an N-policy, multiple vacations and setup times. Jeyakumar and Senthilnathan [5] analyzed the bulk service queueing system with multiple working vacations and server breakdown.

The first work on re-service was done by Madan [6]. He consider an $M/G/1$ queueing model, in which the server performs the first essential service for all arriving customers. As soon as the first service is executed, they may leave the system with probability $(1 - \theta)$, and the second optional service is provided with θ . Madan et al. [7] considered a bulk arrival queue with optional re-service. Jeyakumar and Arumuganathan [8] discussed a bulk queue with multiple vacation and a control policy on request for re-service. Recently, Haridass and Arumuganathan [9] analyzed a batch service queueing system with multiple vacations, setup times and server choice of admitting re-service.

No system is found to be perfect in the real world, since all the devices fail more or less frequently. Thus, the random failures and systematic repair of components of a machining system have a significant impact on the output and the productivity of the machining system. A detailed survey on queues with interruptions was undertaken by Krishnamoorthy et al. [10]. Ayyappan and Shyamala [11] derived the transient solution to an $M^{[X]}/G/1$ queueing system with feedback, random

breakdowns, Bernoulli schedule server vacation and random setup time. An M/G/1 queue with two phases of service subject to random breakdown and delayed repair was examined by Choudhury and Tadj [12]. Senthilnathan and Jeyakumar [13] studied the behavior of the server breakdown without interruption in an $M^{[X]}/G(a,b)/1$ queueing system with multiple vacations and closedown time. An M/G/1 two-phase multi-optional retrial queue with Bernoulli feedback, non-persistent customers and breakdown and repair was analyzed by Lakshmi and Ramanath [14]. Recently, a discrete time queueing system with server breakdowns and changes in the repair times was investigated by Atencia [15].

The operating machine may fail in some cases, but due to the standby machines of the queueing machining system, it remains operative and continues to perform the assigned job. The provision of stand-by and repairmen support to the queueing system maintains the smooth functioning of the system. In the field of computer and communications systems, distribution and service systems, production/manufacturing systems, etc., the applications of queueing models with standby support may be noted.

This paper is organized as follows. A literature survey is given in Section 2. In Section 3, the queueing problem is defined. The system equations are developed in Sections 4. The Probability Generating Function (PGF) of the queue length distribution in the steady state is obtained in Section 5. Various performance measures of the queueing system are derived in Section 6. A computational study is illustrated in Section 7. Conclusions are given in Section 8.

2. Literature Survey

Various authors have analyzed queueing problems of server vacation with several combinations. A batch arrival queue with a vacation time under a single vacation policy was analyzed by Choudhury [16]. Jeyakumar and Arumuganathan [17] have discussed steady state analysis of an $M^{[X]}/G/1$ queue with two service modes and multiple vacation, in which they obtained PGF of the queue size and some performance measures. Balasubramanian et al. [18] discussed steady state analysis of a non-Markovian bulk queueing system with overloading and multiple vacations. Haridass and Arumuganathan [19] discussed a batch arrival general bulk service queueing system with a variant threshold policy for secondary jobs. Recently, Choudhury and Deka [20] discussed a batch arrival queue with an unreliable server and delayed repair, with two phases of service and Bernoulli vacation under multiple vacation policy.

Queueing systems, where the service discipline involves more than one service, have been receiving much attention recently. They are said to have an additional service channel, or to have feedback, or to have optional re-service, or to have two phases of heterogeneous service. Madan [21] analyzed a queueing system with feedback. Madan [22], generalized his previous model by incorporating server vacation. Medhi [23] discussed a single server Poisson input queue with a second optional channel. Arumuganathan and Maliga [24] also examined a bulk queue with re-service of the service station and setup time. Baruah et al. [25] studied a batch arrival queue with two types of service, balking, re-service and vacation. Ayyappan and Sathiya [11] derived the PGF of the non-Markovian queue with two types of service and optional re-service with a general vacation distribution.

One can find an enormous amount of work done on queueing systems with breakdowns. For some papers on random breakdowns in queueing systems, the reader may see Aissani et al. [26], Maraghi et al. [27] and Fadhil et al. [28]. Rajadurai et al. [29] analyzed an $M^{[X]}/G/1$ retrial queue with two phases of service under Bernoulli vacation and random breakdown. Jiang et al. [30] have made a computational analysis of a queue with working breakdown and delayed repair.

The operating system may fail in some cases, but due to stand-by machines, it remains operative and continuous to perform the assigned job. Madan [31] studied the steady state behavior of a queueing system with a stand-by server to serve customers only during the repair period. In that work, repair times were assumed to follow an exponential distribution. Khalaf [32] examined the queueing system

with four different main servers' interruption and a stand-by server. Jain et al. [33] have made a cost analysis of the machine repair problem with standby, working vacation and server breakdown. Kumar et al. [34] discussed a bi-level control of a degraded machining system with two unreliable servers, multiple standbys, startup and vacation. Murugeswari et al. [35] analyzed the bulk arrival queueing model with a stand-by server and compulsory server vacation. Recently, we provided an excellent survey on standby by Kolledath et al. [36].

3. Model Description

This paper deals with a queueing model whose arrival follows a compound Poisson process with intensity rate λ . The main server and stand-by servers serve the customers under the general bulk service rule. The general bulk service rule was first introduced by Neuts [37]. The general bulk service rule states that the server will start to provide service only when at least 'a' units are present in the queue, and the maximum service capacity is 'b' ($b > a$). On completion of a batch service, if less than 'a' customers are present in the queue, then the server has to wait until the queue length reaches the value 'a'. If less than or equal to 'b' and greater than or equal to 'a' customers are in the queue, then all the existing customers are taken into service. If greater than or equal to 'b' customers are in the queue, then 'b' customers are taken into service. The main server may breakdown at any time during regular service with exponential rate α , and in such cases, the main server immediately goes for a repair, which follows an exponential distribution with rate η , while the service to the current batch is interrupted. Such a batch of customers is changed to the stand-by server, which starts service to that batch afresh. The stand-by server remains in the system until the main server's repair is completed. At the instant of repair completion, if the stand-by server is busy, then the current batch of customers is exchanged to the main server, which starts that batch service afresh. At the completion of a regular service (by the main server), the leaving batch may request for a re-service with probability π . However, the re-service is rendered only when the number of customers waiting in the queue is less than a . If no request for re-service is made after the completion of a regular service and the number of customers in the queue is less than a , then the server will avail itself of a vacation of a random length. The server takes a sequence of vacations until the queue size reaches at least a . In addition, we assume that the service time of the main server and stand-by server, re-service and vacation time of the main server are independent of each other and follow a general (arbitrary) distribution.

Notations

Let X be the group size random variable of arrival, g_k be the probability of 'k' customers arriving in a batch and $X(z)$ be its PGF. $S_b(\cdot)$, $R(\cdot)$, $S_s(\cdot)$ and $V(\cdot)$ represent the Cumulative Distribution Functions (CDF) of the regular service and re-service time of the main server, the service time of the stand-by server and the vacation time of the main server with corresponding probability density functions of $s_b(x)$, $r(x)$, $s_s(x)$ and $v(x)$, respectively. $S_b^0(t)$, $R^0(t)$, $S_s^0(t)$ and $V^{(0)}(t)$ represent the remaining regular service and re-service time of service given by the main server, the remaining service time of service given by the stand-by server and the remaining vacation time of the main server at time 't', respectively. $\tilde{S}_b(\theta)$, $\tilde{R}(\theta)$, $\tilde{S}_s(\theta)$ and $\tilde{V}(\theta)$ represent the Laplace–Stieltjes Transform (LST) of S_b , R , S_s and V , respectively.

For further development of the queueing system, let us define the following:

$\varepsilon(t) = 1, 2, 3, 4, 5$ and 6 at time t ; the main server is in regular service, re-service and vacation, and at time t , the stand-by server is in service and idle, respectively.

$Z(t) = j$, if the server is on the j -th vacation.

$N_s(t)$ = number of customers in service station at time t .

$N_q(t)$ = number of customers in the queue at time t .

Define the probabilities:

$$\begin{aligned}
 T_n(t)\Delta t &= Pr\{N_q(t) = n, \varepsilon(t) = 5\}, 0 \leq n \leq a - 1, \\
 P_{m,n}(x,t)\Delta t &= Pr\{N_s(t) = m, N_q(t) = n, x \leq S_b^0(t) \leq x + \Delta t, \varepsilon(t) = 1\}, \\
 &\qquad\qquad\qquad a \leq m \leq b, n \geq 0, \\
 R_n(x,t)\Delta t &= Pr\{N_s(t) = m, N_q(t) = n, x \leq R^0(t) \leq x + \Delta t, \varepsilon(t) = 2\}, \\
 &\qquad\qquad\qquad a \leq m \leq b, n \geq 0, \\
 B_{m,n}(x,t)\Delta t &= Pr\{N_s(t) = m, N_q(t) = n, x \leq S_s^0(t) \leq x + \Delta t, \varepsilon(t) = 4\}, \\
 &\qquad\qquad\qquad a \leq m \leq b, n \geq 0, \\
 Q_{l,j}(x,t)\Delta t &= Pr\{Z(t) = l, N_q(t) = j, x \leq V^0(t) \leq x + \Delta t, \varepsilon(t) = 3\}, \\
 &\qquad\qquad\qquad l \geq 1, j \geq 0.
 \end{aligned}$$

4. Queue Size Distribution

From the above-defined probabilities, we can easily construct the following steady state equations:

$$(\lambda + \eta)T_0 = \sum_{m=a}^b B_{m,0}(0), \tag{1}$$

$$(\lambda + \eta)T_n = \sum_{m=a}^b B_{m,n}(0) + \sum_{k=1}^n T_{n-k}\lambda g_k, 1 \leq n \leq a - 1, \tag{2}$$

$$\begin{aligned}
 -P'_{i,0}(x) &= -(\lambda + \alpha)P_{i,0}(x) + \sum_{m=a}^b P_{m,i}(0)s_b(x) + \eta \int_0^\infty B_{i,0}(y) dy s_b(x) \\
 &\quad + R_i(0)s_b(x) + \sum_{l=1}^\infty Q_{l,i}(0)s_b(x), a \leq i \leq b,
 \end{aligned} \tag{3}$$

$$\begin{aligned}
 -P'_{i,j}(x) &= -(\lambda + \alpha)P_{i,j}(x) + \eta \int_0^\infty B_{i,j}(y) dy s_b(x) + \sum_{k=1}^j P_{i,j-k}(x)\lambda g_k \\
 &\qquad\qquad\qquad j \geq 1, a \leq i \leq b - 1,
 \end{aligned} \tag{4}$$

$$\begin{aligned}
 -P'_{b,j}(x) &= -(\lambda + \alpha)P_{b,j}(x) + \sum_{m=a}^b P_{m,b+j}(0)s_b(x) + \eta \int_0^\infty B_{b,j}(y) dy s_b(x) \\
 &\quad + R_{b+j}(0)s_b(x) + \sum_{l=1}^\infty Q_{l,b+j}(0)s_b(x) + \sum_{k=1}^j P_{b,j-k}(x)\lambda g_k, j \geq 1,
 \end{aligned} \tag{5}$$

$$\begin{aligned}
 -B'_{i,0}(x) &= -(\lambda + \eta)B_{i,0}(x) + \sum_{m=a}^b B_{m,i}(0)s_s(x) + \alpha \int_0^\infty P_{i,0}(y) dy s_s(x) \\
 &\quad + \sum_{k=0}^{a-1} T_k \lambda g_{i-k} s_s(x), a \leq i \leq b,
 \end{aligned} \tag{6}$$

$$\begin{aligned}
 -B'_{i,j}(x) &= -(\lambda + \eta)B_{i,j}(x) + \alpha \int_0^\infty P_{i,j}(y) dy s_s(x) + \sum_{k=1}^j B_{i,j-k}(x)\lambda g_k \\
 &\qquad\qquad\qquad j \geq 1, a \leq i \leq b - 1,
 \end{aligned} \tag{7}$$

$$\begin{aligned}
 -B'_{b,j}(x) &= -(\lambda + \eta)B_{b,j}(x) + \sum_{m=a}^b B_{m,b+j}(0)s_s(x) + \sum_{k=1}^j B_{b,j-k}(x)\lambda g_k \\
 &\quad + \alpha \int_0^\infty P_{b,j}(y) dy s_s(x) + \sum_{k=0}^{a-1} T_k \lambda g_{b+j-k} s_s(x), j \geq 1,
 \end{aligned} \tag{8}$$

$$-R'_0(x) = -\lambda R_0(x) + \pi \sum_{m=a}^b P_{m,0}(0)r(x), \tag{9}$$

$$-R'_n(x) = -\lambda R_n(x) + \pi \sum_{m=a}^b P_{m,n}(0)r(x) + \sum_{k=1}^n R_{n-k}(x)\lambda g_k, \quad 1 \leq n \leq a-1, \tag{10}$$

$$-R'_n(x) = -\lambda R_n(x) + \sum_{k=1}^n R_{n-k}(x)\lambda g_k, \quad n \geq a, \tag{11}$$

$$-Q'_{1,0}(x) = -\lambda Q_{1,0}(x) + (1-\pi) \sum_{m=a}^b P_{m,0}(0)v(x) + R_0(0)v(x) + \eta T_0v(x), \tag{12}$$

$$-Q'_{1,n}(x) = -\lambda Q_{1,n}(x) + (1-\pi) \sum_{m=a}^b P_{m,n}(0)v(x) + R_n(0)v(x) + \eta T_nv(x) + \sum_{k=1}^n Q_{1,n-k}(x)\lambda g_k, \quad 1 \leq n \leq a-1, \tag{13}$$

$$-Q'_{1,n}(x) = -\lambda Q_{1,n}(x) + \sum_{k=1}^n Q_{1,n-k}(x)\lambda g_k, \quad n \geq a, \tag{14}$$

$$-Q'_{j,0}(x) = -\lambda Q_{j,0}(x) + Q_{j-1,0}(0)v(x), \quad j \geq 2, \tag{15}$$

$$-Q'_{j,n}(x) = -\lambda Q_{j,n}(x) + Q_{j-1,n}(0)v(x) + \sum_{k=1}^n Q_{j,n-k}(x)\lambda g_k, \quad j \geq 2, \tag{16}$$

$$1 \leq n \leq a-1,$$

$$-Q'_{j,n}(x) = -\lambda Q_{j,n}(x) + \sum_{k=1}^n Q_{j,n-k}(x)\lambda g_k, \quad j \geq 2, \quad n \geq a. \tag{17}$$

Taking the LST on both sides of Equations (3)–(17), we get,

$$\theta \tilde{P}_{i,0}(\theta) - P_{i,0}(0) = (\lambda + \alpha) \tilde{P}_{i,0}(\theta) - \sum_{m=a}^b P_{m,i}(0) \tilde{S}_b(\theta) - \eta \int_0^\infty B_{i,0}(y) dy \tilde{S}_b(\theta) - R_i(0) \tilde{S}_b(\theta) - \sum_{l=1}^\infty Q_{l,i}(0) \tilde{S}_b(\theta), \quad a \leq i \leq b, \tag{18}$$

$$\theta \tilde{P}_{i,j}(\theta) - P_{i,j}(0) = (\lambda + \alpha) \tilde{P}_{i,j}(\theta) - \eta \int_0^\infty B_{i,j}(y) dy \tilde{S}_b(\theta) - \sum_{k=1}^j \tilde{P}_{i,j-k}(\theta) \lambda g_k, \tag{19}$$

$$a \leq i \leq b-1, \quad j \geq 1,$$

$$\theta \tilde{P}_{b,j}(\theta) - P_{b,j}(0) = (\lambda + \alpha) \tilde{P}_{b,j}(\theta) - \sum_{m=a}^b P_{m,b+j}(0) \tilde{S}_b(\theta) - \eta \int_0^\infty B_{b,j}(y) dy \tilde{S}_b(\theta) - R_{b+j}(0) \tilde{S}_b(\theta) - \sum_{l=1}^\infty Q_{l,b+j}(0) \tilde{S}_b(\theta) - \sum_{k=1}^j \tilde{P}_{b,j-k}(\theta) \lambda g_k, \quad j \geq 1, \tag{20}$$

$$\theta \tilde{B}_{i,0}(\theta) - B_{i,0}(0) = (\lambda + \eta) \tilde{B}_{i,0}(\theta) - \sum_{m=a}^b B_{m,i}(0) \tilde{S}_s(\theta) - \alpha \int_0^\infty P_{i,0}(y) dy \tilde{S}_s(\theta) - \sum_{k=0}^{a-1} T_k \lambda g_{i-k} \tilde{S}_s(\theta), \quad a \leq i \leq b, \tag{21}$$

$$\theta \tilde{B}_{i,j}(\theta) - B_{i,j}(0) = (\lambda + \eta) \tilde{B}_{i,j}(\theta) - \alpha \int_0^\infty P_{i,j}(y) dy \tilde{S}_s(\theta) - \sum_{k=1}^j \tilde{B}_{i,j-k}(\theta) \lambda g_k, \tag{22}$$

$$j \geq 1, \quad a \leq i \leq b-1,$$

$$\begin{aligned} \theta \tilde{B}_{b,j}(\theta) - B_{b,j}(0) &= (\lambda + \eta) \tilde{B}_{b,j}(\theta) - \sum_{m=a}^b B_{m,b+j}(0) \tilde{S}_s(\theta) - \sum_{k=1}^j \tilde{B}_{b,j-k}(\theta) \lambda g_k \\ &\quad - \alpha \int_0^\infty P_{b,j}(y) dy \tilde{S}_s(\theta) - \sum_{k=0}^{a-1} T_k \lambda g_{b+j-k} \tilde{S}_s(\theta), \quad j \geq 1, \end{aligned} \tag{23}$$

$$\theta \tilde{R}_0(\theta) - R_0(0) = \lambda \tilde{R}_0(\theta) - \pi \sum_{m=a}^b P_{m,0}(0) \tilde{R}(\theta), \tag{24}$$

$$\begin{aligned} \theta \tilde{R}_n(\theta) - R_n(0) &= \lambda \tilde{R}_n(\theta) - \pi \sum_{m=a}^b P_{m,n}(0) \tilde{R}(\theta) - \sum_{k=1}^n \tilde{R}_{n-k}(\theta) \lambda g_k, \\ &1 \leq n \leq a - 1, \end{aligned} \tag{25}$$

$$\theta \tilde{R}_n(\theta) - R_n(0) = \lambda \tilde{R}_n(\theta) - \sum_{k=1}^n \tilde{R}_{n-k}(\theta) \lambda g_k, \quad n \geq a, \tag{26}$$

$$\begin{aligned} \theta \tilde{Q}_{1,0}(\theta) - Q_{1,0}(0) &= \lambda \tilde{Q}_{1,0}(\theta) - (1 - \pi) \sum_{m=a}^b P_{m,0}(0) \tilde{V}(\theta) - R_0(0) \tilde{V}(\theta) \\ &\quad - \eta T_0 \tilde{V}(\theta), \end{aligned} \tag{27}$$

$$\begin{aligned} \theta \tilde{Q}_{1,n}(\theta) - Q_{1,n}(0) &= \lambda \tilde{Q}_{1,n}(\theta) - (1 - \pi) \sum_{m=a}^b P_{m,n}(0) \tilde{V}(\theta) - R_n(0) \tilde{V}(\theta) \\ &\quad - \eta T_n \tilde{V}(\theta) - \sum_{k=1}^n \tilde{Q}_{1,n-k}(\theta) \lambda g_k, \quad 1 \leq n \leq a - 1, \end{aligned} \tag{28}$$

$$\theta \tilde{Q}_{1,n}(\theta) - Q_{1,n}(0) = \lambda \tilde{Q}_{1,n}(\theta) - \sum_{k=1}^n \tilde{Q}_{1,n-k}(\theta) \lambda g_k, \quad n \geq a, \tag{29}$$

$$\theta \tilde{Q}_{j,0}(\theta) - Q_{j,0}(0) = \lambda \tilde{Q}_{j,0}(\theta) - Q_{j-1,0}(0) \tilde{V}(\theta), \quad j \geq 2, \tag{30}$$

$$\begin{aligned} \theta \tilde{Q}_{j,n}(\theta) - Q_{j,n}(0) &= \lambda \tilde{Q}_{j,n}(\theta) - Q_{j-1,n}(0) \tilde{V}(\theta) - \sum_{k=1}^n \tilde{Q}_{j,n-k}(\theta) \lambda g_k, \quad j \geq 2, \\ &1 \leq n \leq a - 1, \end{aligned} \tag{31}$$

$$\theta \tilde{Q}_{j,n}(\theta) - Q_{j,n}(0) = \lambda \tilde{Q}_{j,n}(\theta) - \sum_{k=1}^n \tilde{Q}_{j,n-k}(\theta) \lambda g_k, \quad j \geq 2, \quad n \geq a. \tag{32}$$

To find the Probability Generating Function (PGF) for the queue size, we define the following PGFs:

$$\begin{aligned} \tilde{P}_i(z, \theta) &= \sum_{j=0}^\infty \tilde{P}_{i,j}(\theta) z^j, \quad P_i(z, 0) = \sum_{j=0}^\infty P_{i,j}(0) z^j, \quad a \leq i \leq b, \\ \tilde{R}(z, \theta) &= \sum_{j=0}^\infty \tilde{R}_j(\theta) z^j, \quad R(z, 0) = \sum_{j=0}^\infty R_j(0) z^j, \\ \tilde{B}_i(z, \theta) &= \sum_{j=0}^\infty \tilde{B}_{i,j}(\theta) z^j, \quad B_i(z, 0) = \sum_{j=0}^\infty B_{i,j}(0) z^j, \quad a \leq i \leq b, \\ \tilde{Q}_l(z, \theta) &= \sum_{j=0}^\infty \tilde{Q}_{l,j}(\theta) z^j, \quad Q_l(z, 0) = \sum_{j=0}^\infty Q_{l,j}(0) z^j, \quad l \geq 1. \end{aligned} \tag{33}$$

By multiplying Equations (18)–(32) with suitable power of z^n and summing over n ($n = 0$ to ∞) and using Equation (33), we get:

$$(\theta - u(z))\tilde{P}_i(z, \theta) = P_i(z, 0) - \tilde{S}_b(\theta) \left[\sum_{m=a}^b P_{m,i}(0) + R_i(0) + \sum_{j=0}^{\infty} Q_{l,i}(0) + \eta \tilde{B}_i(z, 0) \right],$$

$$a \leq i \leq b - 1, \tag{34}$$

$$z^b(\theta - u(z))\tilde{P}_b(z, \theta) = (z^b - \tilde{S}_b(\theta))P_b(z, 0) - \tilde{S}_b(\theta) \left[\sum_{m=a}^{b-1} P_m(z, 0) + R(z, 0) + \sum_{l=1}^{\infty} Q_l(z, 0) + z^b \eta \tilde{B}_b(z, 0) - \sum_{j=0}^{b-1} \left(\sum_{m=a}^b P_{m,j}(0)z^j + R_j(0)z^j + \sum_{j=0}^{\infty} Q_{l,j}(0)z^j \right) \right], \tag{35}$$

$$(\theta - v(z))\tilde{B}_i(z, \theta) = B_i(z, 0) - \tilde{S}_s(\theta) \left[\alpha \tilde{P}_i(z, 0) + \sum_{m=a}^b B_{m,i}(0) + \sum_{k=0}^{a-1} T_k \lambda g_{i-k} \right],$$

$$a \leq i \leq b - 1, \tag{36}$$

$$z^b(\theta - v(z))\tilde{B}_b(z, \theta) = (z^b - \tilde{S}_s(\theta))B_b(z, 0) - \tilde{S}_s(\theta) \left[\sum_{m=a}^{b-1} B_m(z, 0) + z^b \alpha \tilde{P}_b(z, 0) + \lambda \sum_{k=0}^{a-1} \sum_{j=b}^{\infty} T_k z^k g_{j-k} z^{j-k} - \sum_{j=0}^{b-1} \sum_{m=a}^b B_{m,j}(0)z^j \right], \tag{37}$$

$$(\theta - w(z))\tilde{R}(z, \theta) = R(z, 0) - \pi \tilde{R}(\theta) \sum_{n=0}^{a-1} \sum_{m=a}^b P_{m,n}(0)z^n, \tag{38}$$

$$(\theta - w(z))\tilde{Q}_1(z, \theta) = Q_1(z, 0) - \tilde{V}(\theta) \sum_{n=0}^{a-1} \left[(1 - \pi) \sum_{m=a}^b P_{m,n}(0)z^n + R_n(0)z^n + \eta T_n z^n \right], \tag{39}$$

$$(\theta - w(z))\tilde{Q}_j(z, \theta) = Q_j(z, 0) - \tilde{V}(\theta) \sum_{n=0}^{a-1} Q_{j-1,n}(0)z^n, \quad j \geq 2, \tag{40}$$

where

$$u(z) = \lambda + \alpha - \lambda X(z), \quad v(z) = \lambda + \eta - \lambda X(z), \quad w(z) = \lambda - \lambda X(z).$$

Substitute $\theta = u(z)$ in (34) and (35), we get,

$$P_i(z, 0) = \tilde{S}_b(u(z)) \left[\sum_{m=a}^b P_{m,i}(0) + R_i(0) + \sum_{j=0}^{\infty} Q_{l,i}(0) + \eta \tilde{B}_i(z, 0) \right],$$

$$a \leq i \leq b - 1, \tag{41}$$

$$P_b(z, 0) = \frac{\tilde{S}_b(u(z))}{(z^b - \tilde{S}_b(u(z)))} \left[\sum_{m=a}^{b-1} P_m(z, 0) + R(z, 0) + \sum_{l=1}^{\infty} Q_l(z, 0) + z^b \eta \tilde{B}_b(z, 0) - \sum_{j=0}^{b-1} \left(\sum_{m=a}^b P_{m,j}(0)z^j + R_j(0)z^j + \sum_{j=0}^{\infty} Q_{l,j}(0)z^j \right) \right], \tag{42}$$

substitute $\theta = v(z)$ in (36) and (37), we get,

$$B_i(z, 0) = \tilde{S}_s(v(z)) \left[\alpha \tilde{P}_i(z, 0) + \sum_{m=a}^b B_{m,i}(0) + \sum_{k=0}^{a-1} T_k \lambda g_{i-k} \right], \quad a \leq i \leq b-1, \tag{43}$$

$$B_b(z, 0) = \frac{\tilde{S}_s(v(z))}{(z^b - \tilde{S}_s(v(z)))} \left[\sum_{m=a}^{b-1} B_m(z, 0) + z^b \alpha \tilde{P}_b(z, 0) + \lambda \sum_{k=0}^{a-1} \sum_{j=b}^{\infty} T_k z^k g_{j-k} z^{j-k} - \sum_{j=0}^{b-1} \sum_{m=a}^b B_{m,j}(0) z^j \right], \tag{44}$$

substitute $\theta = w(z)$ in (38) to (40), we get

$$R(z, 0) = \pi \tilde{R}(w(z)) \sum_{n=0}^{a-1} \sum_{m=a}^b P_{m,n}(0) z^n, \tag{45}$$

$$Q_1(z, 0) = \tilde{V}(w(z)) \sum_{n=0}^{a-1} \left[(1 - \pi) \sum_{m=a}^b P_{m,n}(0) z^n + R_n(0) z^n + \eta T_n z^n \right], \tag{46}$$

$$Q_j(z, 0) = \tilde{V}(w(z)) \sum_{n=0}^{a-1} Q_{j-1,n}(0) z^n, \quad j \geq 2. \tag{47}$$

Substitute Equations (41)–(47) in Equations (34)–(40) after simplification, and we get,

$$(\theta - u(z)) \tilde{P}_i(z, \theta) = (\tilde{S}_b(u(z)) - \tilde{S}_b(\theta)) \left[\sum_{m=a}^b P_{m,i}(0) + R_i(0) + \sum_{j=0}^{\infty} Q_{l,i}(0) + \eta \tilde{B}_i(z, 0) \right], \quad a \leq i \leq b-1, \tag{48}$$

$$(\theta - u(z)) \tilde{P}_b(z, \theta) = \frac{(\tilde{S}_b(u(z)) - \tilde{S}_b(\theta))}{(z^b - \tilde{S}_b(u(z)))} \left[\sum_{m=a}^{b-1} P_m(z, 0) + R(z, 0) + \sum_{l=1}^{\infty} Q_l(z, 0) + z^b \eta \tilde{B}_b(z, 0) - \sum_{j=0}^{b-1} \left(\sum_{m=a}^b P_{m,j}(0) z^j + R_j(0) z^j + \sum_{j=0}^{\infty} Q_{l,j}(0) z^j \right) \right], \tag{49}$$

$$(\theta - v(z)) \tilde{B}_i(z, \theta) = (\tilde{S}_s(v(z)) - \tilde{S}_s(\theta)) \left[\alpha \tilde{P}_i(z, 0) + \sum_{m=a}^b B_{m,i}(0) + \sum_{k=0}^{a-1} T_k \lambda g_{i-k} \right], \quad a \leq i \leq b-1, \tag{50}$$

$$(\theta - v(z)) \tilde{B}_b(z, \theta) = \frac{(\tilde{S}_s(v(z)) - \tilde{S}_s(\theta))}{(z^b - \tilde{S}_s(v(z)))} \left[\sum_{m=a}^{b-1} B_m(z, 0) + z^b \alpha \tilde{P}_b(z, 0) + \lambda \sum_{k=0}^{a-1} \sum_{j=b}^{\infty} T_k z^k g_{j-k} z^{j-k} - \sum_{j=0}^{b-1} \sum_{m=a}^b B_{m,j}(0) z^j \right], \tag{51}$$

$$(\theta - w(z)) \tilde{R}(z, \theta) = (\tilde{R}(w(z)) - \tilde{R}(\theta)) \pi \sum_{n=0}^{a-1} \sum_{m=a}^b P_{m,n}(0) z^n, \tag{52}$$

$$(\theta - w(z)) \tilde{Q}_1(z, \theta) = (\tilde{V}(w(z)) - \tilde{V}(\theta)) \sum_{n=0}^{a-1} \left[(1 - \pi) \sum_{m=a}^b P_{m,n}(0) z^n + R_n(0) z^n + \eta T_n z^n \right], \tag{53}$$

$$(\theta - w(z)) \tilde{Q}_j(z, \theta) = (\tilde{V}(w(z)) - \tilde{V}(\theta)) \sum_{n=0}^{a-1} Q_{j-1,n}(0) z^n, \quad j \geq 2. \tag{54}$$

5. Probability Generating Function of the Queue Size

5.1. The PGF of the Queue Size at an Arbitrary Time Epoch

Let $P(z)$ be the PGF of the queue size at an arbitrary time epoch. Then,

$$P(z) = \sum_{i=a}^b \tilde{P}_i(z, 0) + \sum_{i=a}^b \tilde{B}_i(z, 0) + \tilde{R}(z, 0) + \sum_{l=1}^{\infty} \tilde{Q}_l(z, 0) + T(z). \tag{55}$$

By substituting $\theta = 0$ in Equations (48)–(54), then Equation (55) becomes:

$$\begin{aligned} & K_1(z) \sum_{i=a}^{b-1} (z^b - z^i) c_i + (1 - \tilde{V}(w(z))) K_3(z) \sum_{n=0}^{a-1} c_n z^n \\ & + K_2(z) \sum_{i=a}^{b-1} (z^b - z^i) d_i + (\tilde{V}(w(z)) - \tilde{R}(w(z))) K_3(z) \sum_{n=0}^{a-1} \pi p_n z^n \\ & + \left[\eta [Y_1(z) - \tilde{V}(w(z)) K_3(z)] - v(z) K_2(z) + w(z) Y_1(z) \right] \sum_{k=0}^{a-1} T_k z^k \end{aligned} \tag{56}$$

$$P(z) = \frac{\hspace{10em}}{w(z) Y_1(z)}$$

where $p_i = \sum_{m=a}^b P_{m,i}(0)$, $v_i = \sum_{l=1}^{\infty} Q_{l,i}(0)$, $q_i = \sum_{m=a}^b B_{m,i}(0)$, $R_i(0) = r_i$, $c_i = p_i + v_i + r_i$ and $d_i = q_i + \sum_{k=0}^{a-1} T_k \lambda g_{i-k}$ and the expressions for $K_1(z)$, $K_2(z)$, $K_3(z)$ and $Y_1(z)$ are defined in Appendix A.

5.2. Steady State Condition

The probability generating function has to satisfy $P(1) = 1$. In order to satisfy this condition, applying L'Hopital's rule and evaluating $\lim_{z \rightarrow 1} P(z)$, then equating the expression to one, we have, $H = (-\lambda X_1) F_1$, where the expressions H and F_1 are defined in Appendix B.

Since p_i , c_i , d_i and T_i are probabilities of ' i ' customers being in the queue, it follows that H must be positive. Thus, $P(1) = 1$ is satisfied iff $(-\lambda X_1) F_1 > 0$. If:

$$\rho = \frac{\lambda X_1 (\alpha + \eta) (1 - \tilde{S}_b(\alpha)) (1 - \tilde{S}_s(\eta))}{b \alpha \eta [\tilde{S}_b(\alpha) (1 - \tilde{S}_s(\eta)) + \tilde{S}_s(\eta) (1 - \tilde{S}_b(\alpha))]}$$

then $\rho < 1$ is the condition for the existence of the steady state for the model under consideration.

5.3. Computational Aspects

Equation (56) has $2b + a$ unknowns $c_0, c_1, \dots, c_{b-1}, d_a, \dots, d_{b-1}, p_0, p_1, \dots, p_{a-1}$ and T_0, T_1, \dots, T_{a-1} . Now, Equation (56) gives the PGF of the number of customers involving only $2b + a$ unknowns. We can express $c_i (0 \leq i \leq a - 1)$ in terms of p_i and T_i in such a way that the numerator has only $2b$ constants. Now, Equation (56) gives the PGF of the number of customers involving only $2b$ unknowns. By Rouché's theorem, it can be proven that $Y_1(z)$ has $2b - 1$ zeros inside and one on the unit circle $|z| = 1$. Since $P(z)$ is analytic within and on the unit circle, the numerator must vanish at these points, which gives $2b$ equations in $2b$ unknowns. We can solve these equations by any suitable numerical technique.

5.4. Result 1

The probability that $n (0 \leq n \leq a - 1)$ customers are in queue during the main server's re-service completion r_n can be expressed as the probability of n customers in the queue during the main server's regular busy period p_n as,

$$r_n = \pi \sum_{k=0}^n \gamma_{n-k} p_k, \quad n = 0, 1, 2, \dots, a - 1 \tag{57}$$

where γ_n are the probabilities of n customers arriving during the main server’s re-service time.

5.5. Result 2

The probability that $n(0 \leq n \leq a - 1)$ customers are in queue c_n can be expressed as the sum of the probability of n customers in the queue during the main server’s busy period and the stand-by server’s idle time p_n and T_n as,

$$c_n = \sum_{k=0}^n \tau_{n-k}^{(1)} p_k + \tau_{n-k}^{(2)} T_k, \quad 0 \leq n \leq a - 1. \tag{58}$$

where:

$$\begin{aligned} \tau_n^{(1)} &= \frac{\pi(\gamma_n - \beta_n) + \sum_{k=1}^n \beta_k \tau_{n-k}^{(1)}}{1 - \beta_0} \\ \tau_n^{(2)} &= \frac{\eta \beta_n + \sum_{k=1}^n \beta_k \tau_{n-k}^{(2)}}{1 - \beta_0} \end{aligned} \tag{59}$$

γ_n, β_n are the probabilities of n customers arriving during the main server’s re-service and vacation time, respectively.

5.6. Particular Case

Case 1:

When there is no breakdown and re-service, then Equation (65) reduces to:

$$P(z) = \frac{(\tilde{S}_b(w(z)) - 1) \sum_{i=a}^{b-1} (z^b - z^i) c_i + (z^b - 1)(\tilde{V}(w(z)) - 1) \sum_{n=0}^{a-1} c_n z^n}{(-w(z))(z^b - \tilde{S}_b(w(z)))} \tag{60}$$

which coincides with the PGF of Senthilnathan et al. [19] without closedown.

Case 2:

When there is no breakdown, then Equation (65) reduces to:

$$\begin{aligned} & (1 - \tilde{S}_b(w(z))) \sum_{i=a}^{b-1} (z^b - z^i) c_i + (z^b - 1)(1 - \tilde{V}(w(z))) \sum_{n=0}^{a-1} c_n z^n \\ & + (z^b - 1)(\tilde{V}(w(z)) - \tilde{R}(w(z))) \sum_{n=0}^{a-1} \pi p_n z^n \\ P(z) &= \frac{\hspace{10em}}{(w(z))(z^b - \tilde{S}_b(w(z)))} \end{aligned} \tag{61}$$

which is the PGF of Jeyakumar et al. [38].

5.7. PGF of the Queue Size in Various Epochs

5.7.1. PGF of the Queue Size in the Main Server’s Service Completion Epoch

The probability generating function of the main server’s service completion epoch $M(z)$ is obtained from Equations (48) and (49):

$$\begin{aligned}
 & (1 - \tilde{S}_b(u(z))) \left[v(z)(z^b - \tilde{S}_s(v(z))) \left(\sum_{i=a}^{b-1} (z^b - z^i)c_i - \sum_{k=0}^{a-1} c_k z^k \right) \right. \\
 & \quad \left. + \pi \tilde{R}(w(z)) \sum_{n=0}^{a-1} p_n z^n + \tilde{V}(w(z)) \sum_{n=0}^{a-1} ((1 - \pi)p_n + r_n + \eta T_n + v_n) z^n \right) \\
 & \quad \left. + z^b \eta (1 - \tilde{S}_s(v(z))) \left(\sum_{i=a}^{b-1} (z^b - z^i)(q_i + r_i^{(2)} + \sum_{k=0}^{a-1} T_k \lambda g_{i-k}) - v(z)T(z) \right) \right] \\
 M(z) = & \frac{\hspace{10em}}{Y_1(z)}
 \end{aligned} \tag{62}$$

5.7.2. PGF of the Queue Size in the Vacation Completion Epoch

The PGF of the main server’s vacation completion epoch $V(z)$ is obtained from Equations (53) and (54); we get,

$$V(z) = \frac{(1 - \tilde{V}(w(z))) \sum_{n=0}^{a-1} ((1 - \pi)p_n + r_n + \eta T_n + v_n) z^n}{w(z)} \tag{63}$$

5.7.3. PGF of the Queue Size in the Main Server’s Re-Service Completion Epoch

The PGF of the main server’s re-service completion epoch $R(z)$ is obtained from Equation (52); we get,

$$R(z) = \frac{(1 - \tilde{R}(w(z))) \sum_{n=0}^{a-1} \pi p_n z^n}{w(z)} \tag{64}$$

5.7.4. PGF of the Queue Size in the Stand-by Server’s Service Completion Epoch

The probability generating function of the stand-by server’s service completion epoch $N(z)$ is derived from Equations (50) and (51); we get,

$$\begin{aligned}
 & (1 - \tilde{S}_s(v(z))) \left[z^b \alpha (1 - \tilde{S}_b(u(z))) \left(\sum_{i=a}^{b-1} (z^b - z^i)c_i - \sum_{k=0}^{a-1} c_k z^k \right) \right. \\
 & \quad \left. + \pi \tilde{R}(w(z)) \sum_{n=0}^{a-1} p_n z^n + \tilde{V}(w(z)) \sum_{n=0}^{a-1} ((1 - \pi)p_n + r_n + \eta T_n + v_n) z^n \right) \\
 & \quad \left. + u(z)(z^b - \tilde{S}_b(u(z))) \left(\sum_{i=a}^{b-1} (z^b - z^i)(q_i + \sum_{k=0}^{a-1} T_k \lambda g_{i-k}) - v(z)T(z) \right) \right] \\
 N(z) = & \frac{\hspace{10em}}{Y_1(z)}
 \end{aligned} \tag{65}$$

6. Some Performance Measures

6.1. The Main Server’s Expected Length of Idle Period

Let K be the random variable denoting the ‘idle period due to multiple vacation processes’. Let Y be the random variable defined by:

$$Y = \begin{cases} 0 & \text{if the server finds at least 'a' customers after the first vacation} \\ 1 & \text{if the server finds less than 'a' customers after the first vacation} \end{cases}$$

Now,

$$\begin{aligned}
 E(K) &= E(K/Y = 0)P(Y = 0) + E(K/Y = 1)P(Y = 1) \\
 &= E(V)P(Y = 0) + (E(V) + E(K))P(Y = 1),
 \end{aligned}$$

Solving for $E(K)$, we get:

$$E(K) = \frac{E(V)}{(1 - P(Y = 1))} = \frac{E(V)}{\left(1 - \sum_{n=0}^{a-1} \sum_{i=0}^n \left[\beta_i [p_n + r_n + \eta T_n]\right]\right)}$$

6.2. Expected Queue Length

The mean number of customers waiting in the queue $E(Q)$ in an arbitrary time epoch is obtained by differentiating $P(z)$ at $z = 1$ and is given by:

$$\begin{aligned}
 & f_1(X, S_b, S_s) \left[\sum_{i=a}^{b-1} [b(b-1) - i(i-1)]c_i \right] \\
 & + f_1(X, S_b, S_s) \left[\sum_{i=a}^{b-1} (b(b-1) - i(i-1))d_i \right] \\
 & + f_2(X, S_b, S_s) \left[\sum_{i=a}^{b-1} (b-i)c_i \right] + f_3(X, S_b, S_s) \sum_{i=a}^{b-1} (b-i)d_i \\
 & + f_4(X, S_b, S_s, V) \sum_{n=0}^{a-1} n c_n + f_5(X, S_b, S_s, V) \sum_{n=0}^{a-1} c_n \\
 & + f_6(X, S_b, S_s, R, V) \sum_{n=0}^{a-1} \pi n p_n + f_7(X, S_b, S_s, R, V) \sum_{n=0}^{a-1} \pi p_n \\
 & + f_8(X, S_b, S_s, V) \sum_{n=0}^{a-1} n T_n + f_9(X, S_b, S_s, V) \sum_{n=0}^{a-1} T_n
 \end{aligned}$$

$$E(Q) = \frac{\text{above sum}}{3(F_{12})^2}, \tag{66}$$

the expressions for $f_i (i = 1, 2, \dots, 9)$ are defined in Appendix B.

6.3. Expected Waiting Time

The expected waiting time is obtained using Little’s formula as:

$$E(W) = \frac{E(Q)}{\lambda E(X)} \tag{67}$$

where $E(Q)$ is given in Equation (66).

7. Numerical Example

A numerical example of our model is analyzed for a particular case with the following assumptions:

1. The batch size distribution of the arrival is geometric with mean 2.
2. Take $a = 5$ and $b = 8$, and the service time distribution is Erlang-2 (both servers).
3. The vacation and re-service time of the main server follow an exponential distribution with parameter $\omega = 5, \epsilon = 3$, respectively.
4. Let m_1 be the service rate for the main server.
5. Let m_2 be the service rate for the stand-by server.

The unknown probabilities of the queue size distribution are computed using numerical techniques. The zeros of the function $Y_1(z)$ are obtained (see Figure 1), and simultaneous equations are solved by using MATLAB. The values which are satisfies the stability condition (see Figure 2) are used for calculating the table values.

The expected queue length $E(Q)$ and the expected waiting time $E(W)$ are calculated for various arrival rate sand service rates, and the results are tabulated.

From Tables 1–4, the following observations can be made.

1. As arrival rate λ increases, the expected queue size and expected waiting time are also increase.
2. When the main server's and stand-by server's service rate increases, the expected queue size and expected waiting time decrease.
3. When the main server's vacation rate increases, the expected queue size increases.

Table 1. Arrival rate vs. expected queue length and expected waiting time for the values $m_1 = 10, m_2 = 9.5, \alpha = 1, \eta = 2, \pi = 0.3, \epsilon = 3,$ and $\omega = 5.$

λ	ρ	$E(Q)$	$E(W)$
5.00	0.131407	8.657374	0.865737
5.25	0.137978	9.539724	0.908545
5.50	0.144548	10.375816	0.943256
5.75	0.151119	11.149076	0.969485
6.00	0.157689	11.843026	0.986919
6.25	0.164259	12.441064	0.995285
6.50	0.170830	12.927026	0.994387
6.75	0.177400	13.285663	0.984123
7.00	0.183970	13.502456	0.964461
7.25	0.190541	13.563622	0.935422
7.50	0.197111	13.456834	0.897122

Table 2. Main server's service rate vs. expected queue length and expected waiting time for the values $\lambda = 5, m_2 = 5, \alpha = 1, \eta = 2, \pi = 0.3, \epsilon = 3,$ and $\omega = 5.$

m_1	ρ	$E(Q)$	$E(W)$
5.25	0.257410	26.824821	2.682482
5.50	0.248885	26.082729	2.608273
5.75	0.240905	25.388555	2.538855
6.00	0.233420	24.735574	2.473557
6.25	0.226385	24.117976	2.411798
6.50	0.219761	23.531054	2.353105
6.75	0.213513	22.971202	2.297120
7.00	0.207610	22.435611	2.243561
7.25	0.202024	21.921511	2.192151
7.50	0.196731	21.427009	2.142701
7.75	0.191707	20.950354	2.095035
8.00	0.186934	20.490051	2.049005

Table 3. Stand-by server's service rate vs. expected queue length and expected waiting time for the values $\lambda = 5, m_1 = 10, \alpha = 1, \eta = 2, \pi = 0.3, \epsilon = 3,$ and $\omega = 5.$

m_2	ρ	$E(Q)$	$E(W)$
4.0	0.260103	65.007246	4.062953
4.5	0.254645	59.415276	3.713455
5.0	0.249401	53.808260	3.363016
5.5	0.244361	48.320593	3.020037
6.0	0.239515	43.028238	2.689265
6.5	0.234853	37.970425	2.373152
7.0	0.230366	33.162455	2.072653
7.5	0.226045	28.605406	1.787838
8.0	0.22188	24.292435	1.518277
8.5	0.217865	20.212075	1.263255
9.0	0.213991	16.351005	1.021938

Table 4. The effect of the main server's vacation rate on expected queue length for the values $\lambda = 5$, $m_1 = 10$, $m_2 = 9.5$, $\alpha = 1$, $\eta = 2$, $\pi = 0.3$, and $\epsilon = 3$.

ω	Erlang	Exponential
5.00	8.657374	8.279153
5.25	8.808004	8.448114
5.50	8.950939	8.607757
5.75	9.086640	8.758748
6.00	9.215535	8.901685
6.25	9.338068	9.037158
6.50	9.454625	9.165674
6.75	9.565595	9.287730
7.00	9.671330	9.403767
7.25	9.772165	9.514200
7.50	9.868410	9.619407
7.75	9.960351	9.719734
8.00	10.048247	9.815494

```

1 -   clc
2 -   clear all
3 -   syms z X Y1 Y2;
4 -   f=0.25;
5 -   for Lam = 5:f:7.5
6 -       Lam
7 -       m1 = 10;
8 -       m2 = 9.5;
9 -       a=5;
10 -      b=8;
11 -      Eta = 2;
12 -      Alp = 1;
13 -      X = -3/(10*((7*z)/10 - 1));
14 -      u = Lam - Lam*X+Alp;
15 -      v = Lam - Lam*X+Eta;
16 -      S1 = ((2*m1)/(2*m1+u))^2;
17 -      B1 = ((2*m2)/(2*m2+v))^2;
18 -      Y1=u*v*(z^b-S1)*(z^b-B1)-z^b*z^b*Alp*Eta*(1-S1)*(1-B1);
19 -      Y2 = solve(Y1)
20 -   end

```

Figure 1. MATLAB code for finding the roots.

```

1 -   clc
2 -   clear all
3 -   f=0.25;
4 -   for Lam =5.00:f:7.5
5 -       Lam;
6 -       fprintf('Lam=%g \n', Lam);
7 -       m1 = 10;
8 -       m2 = 9.5;
9 -       a=5;
10 -      b=8;
11 -      pi= 0.3;
12 -      Eta = 2;
13 -      Alp = 1;
14 -      X1=2;
15 -      S1 = ((2*m1)/(2*m1+Alp))^2;
16 -      B1 = ((2*m2)/(2*m2+Eta))^2;
17 -      N=(Lam*X1*(Alp+Eta)*(1-S1)*(1-B1));
18 -      D=b*Alp*Eta*(B1*(1-S1)+S1*(1-B1));
19 -      fprintf('rho=%g \n', (N/D))
20 -   end
21

```

Figure 2. MATLAB code for finding the rho value.

8. Conclusions

In this paper, a batch arrival general bulk service queueing system with breakdown and repair, stand-by server, multiple vacation and control policy on request for re-service is analyzed. The probability generating function of the queue size distribution at an arbitrary time is obtained. Some performance measures are calculated. The particular cases of the model are also deduced. From the numerical results, it is observed that when the arrival rate increases, the expected queue length and waiting time of the customers are also increase; if the service rate increases (for both server's), then the expected queue length and expected waiting time decrease. It is also observed that, if the main server's vacation rate increases, then the expected queue length increases.

Author Contributions: G.A.: To describe the model. S.K.: Convert the theoretical model into mathematical model and solving.

Conflicts of Interest: There is no conflict of interest by the author to publish this paper.

Appendix A

The expressions used in Equation (56) are defined as follows:

$$K_1(z) = w(z)(1 - \tilde{S}_b(u(z)))A_1(z),$$

$$K_2(z) = w(z)(1 - \tilde{S}_s(v(z)))A_2(z),$$

$$K_3(z) = Y_1(z) - w(z)A_1(z)(1 - \tilde{S}_b(u(z))),$$

where

$$A_1(z) = v(z)(z^b - \tilde{S}_s(v(z))) + z^b\alpha(1 - \tilde{S}_s(v(z)))$$

$$A_2(z) = u(z)(z^b - \tilde{S}_b(u(z))) + z^b\eta(1 - \tilde{S}_b(u(z)))$$

$$Y_1(z) = u(z)v(z)(z^b - \tilde{S}_b(u(z)))(z^b - \tilde{S}_s(v(z))) - z^{2b}\alpha\eta(1 - \tilde{S}_b(u(z)))(1 - \tilde{S}_s(v(z))).$$

Appendix B

The expressions for f_i 's in (66) are defined as follows:

$$\begin{aligned}
 f_1(X, S_b, S_s) &= 3E_1 F_{12}, \\
 f_2(X, S_b, S_s) &= 3F_7 F_{12} - 2E_1 F_{13}, \\
 f_3(X, S_b, S_s) &= 3F_9 F_{12} - 2E_1 F_{13}, \\
 f_4(X, S_b, S_s, V) &= 6b\alpha\eta V_1 E_2 F_{12}, \\
 f_5(X, S_b, S_s, V) &= 3F_{12}[b\alpha\eta E_2 V_2 + V_1(F_7 - F_2)] - 2b\alpha\eta E_2 V_1 F_{13}, \\
 f_6(X, S_b, S_s, R_1, V) &= 6b\alpha\eta(R_1 - V_1) E_2 F_{12}, \\
 f_7(X, S_b, S_s, R_1, V) &= 3F_{12}[b\alpha\eta E_2(R_2 - V_2) + (R_1 - V_1)(F_7 - F_2)] - 2b\alpha\eta E_2(R_1 + V_1) F_{13}, \\
 f_8(X, S_b, S_s, V) &= 6F_{11} F_{12}, \\
 f_9(X, S_b, S_s, V) &= 3E_{10} F_{12} - 2F_{11} F_{13},
 \end{aligned}$$

where:

$$\begin{aligned}
 E_1 &= -\lambda X_1(\alpha + \eta)(1 - \tilde{S}_b(\alpha))(1 - \tilde{S}_s(\eta)), \\
 E_2 &= \tilde{S}_b(\alpha)(\tilde{S}_s(\eta) - 1) + \tilde{S}_s(\eta)(\tilde{S}_b(\alpha) - 1), \\
 E_3 &= S_{b1}(1 - \tilde{S}_s(\eta)) + S_{s1}(1 - \tilde{S}_b(\alpha)), \\
 E_4 &= -\lambda X_1(1 - \tilde{S}_s(\eta)), \\
 E_5 &= -\lambda X_2(1 - \tilde{S}_s(\eta)) + 2\lambda X_1 S_{s1}, \\
 E_6 &= -\lambda X_3(1 - \tilde{S}_s(\eta)) + 3\lambda X_2 S_{s1} + 3\lambda X_1 S_{s2}, \\
 E_7 &= -\lambda X_1(1 - \tilde{S}_b(\alpha)), \\
 E_8 &= -\lambda X_2(1 - \tilde{S}_b(\alpha)) + 2\lambda X_1 S_{b1}, \\
 E_9 &= -\lambda X_3(1 - \tilde{S}_b(\alpha)) + 3\lambda X_2 S_{b1} + 3\lambda X_1 S_{b2}, \\
 E_{10} &= \eta[b\alpha\eta E_2 V_2 - V_1(F_2 - F_7)] + \lambda X_2(E_1 - F_1) + \lambda X_1(F_9 - F_2) - (\eta/3)(F_8 + F_{10}), \\
 F_1 &= E_1 - b\alpha\eta E_2, \\
 F_2 &= (1 - \tilde{S}_b(\alpha))(1 - \tilde{S}_s(\eta))[2(\lambda X_1)^2 - \lambda X_2(\alpha + \eta) - 2b(2b - 1)\alpha\eta] \\
 &\quad + [b(b - 1)\alpha\eta - 2b\lambda X_1(\alpha + \eta)][(1 - \tilde{S}_b(\alpha)) + (1 - \tilde{S}_s(\eta))] \\
 &\quad + 2[2b\alpha\eta + \lambda X_1(\alpha + \eta)]E_3 + 2b\alpha\eta(b - S_{b1} - S_{s1}) \\
 F_3 &= (\alpha + \eta)(b - S_{s1}) + E_4 - b\alpha\tilde{S}_s(\eta), \\
 F_4 &= (\alpha + \eta)(b(b - 1) - S_{s2}) - b\alpha[(b - 1)\tilde{S}_s(\eta) + 2S_{s1}] - 2b\lambda X_1 + E_5, \\
 F_5 &= (\alpha + \eta)(b - S_{b1}) + E_7 - b\eta\tilde{S}_b(\alpha), \\
 F_6 &= (\alpha + \eta)(b(b - 1) - S_{b2}) - b\eta[(b - 1)\tilde{S}_b(\alpha) + 2S_{b1}] - 2b\lambda X_1 + E_8, \\
 F_7 &= E_8(\alpha + \eta)(1 - \tilde{S}_s(\eta)) + 2E_7 F_3, \\
 F_8 &= E_9(\alpha + \eta)(1 - \tilde{S}_s(\eta)) + 3E_8 F_3 + 3E_7 F_4, \\
 F_9 &= E_5(\alpha + \eta)(1 - \tilde{S}_b(\alpha)) + 2E_4 F_5, \\
 F_{10} &= E_6(\alpha + \eta)(1 - \tilde{S}_b(\alpha)) + 3E_5 F_5 + 3E_4 F_6, \\
 F_{11} &= b\alpha\eta E_2(\eta V_1 + \lambda X_1) + \eta\lambda X_1[(\alpha + \eta)E_3 - (1 - \tilde{S}_b(\alpha))F_3 + (1 - \tilde{S}_s(\eta))F_5], \\
 F_{12} &= -2\lambda X_1 F_1, \\
 F_{13} &= -3[\lambda X_1 F_2 + \lambda X_2 F_1].
 \end{aligned}$$

$$H = \begin{cases} 2E_1 \sum_{i=a}^{b-1} (b - i)c_i + 2E_1 \sum_{i=a}^{b-1} (b - i)d_i + 2b\alpha\eta E_2 V_1 \sum_{n=0}^{a-1} c_n \\ + 2b\alpha\eta E_2 (R_1 + V_1) \sum_{n=0}^{a-1} \pi p_n + 2F_{11} \sum_{n=0}^{a-1} T_n, \end{cases}$$

where:

$$\begin{aligned}
 S_{b1} &= -\lambda X_1 \tilde{S}'_b(\alpha), \quad S_{b2} = \tilde{S}''_b(\alpha)(-\lambda X_1)^2 - \lambda X_2 \tilde{S}'_b(\alpha), \\
 S_{s1} &= -\lambda X_1 \tilde{S}'_s(\eta), \quad S_{s2} = \tilde{S}''_s(\eta)(-\lambda X_1)^2 - \lambda X_2 \tilde{S}'_s(\eta), \\
 R_1 &= \lambda X_1 E(R), \quad R_2 = \lambda X_2 E(R) + \lambda^2 X_1^2 E(R^2) \\
 V_1 &= \lambda X_1 E(V), \quad V_2 = \lambda X_2 E(V) + \lambda^2 X_1^2 E(V^2), \quad X_1 = E(X), \quad X_2 = E(X^2).
 \end{aligned}$$

References

1. Doshi, B.T. Queueing systems with vacations—A survey. *Queueing Syst.* **1986**, *1*, 29–66. [[CrossRef](#)]
2. Takagi, H. Vacation and Priority Systems. *Queueing Analysis: A Foundation of Performance Evaluation*; North-Holland: Amsterdam, The Netherlands, 1991; Volume I.
3. Baba, Y. On the $M^{[X]}/G/1$ queue with vacation time. *Oper. Res. Lett.* **1986**, *5*, 93–98. [[CrossRef](#)]
4. Krishna Reddy, G.V.; Nadarajan, R.; Arumuganathan, R. Analysis of a bulk queue with N policy multiple vacations and setup times. *Comput. Oper. Res.* **1998**, *25*, 957–967. [[CrossRef](#)]
5. Jeyakumar, S.; Senthilnathan, B. Modelling and analysis of a bulk service queueing model with multiple working vacations and server breakdown. *RAIRO-Oper. Res.* **2017**, *51*, 485–508. [[CrossRef](#)]
6. Madan, K.C. An M/G/1 queue with second optional service. *Queueing Syst.* **2000**, *34*, 37–46. [[CrossRef](#)]
7. Madan, K.C. On $M^{[X]}/(G1, G2)/1$ queue with optional re-service. *Appl. Math. Comput.* **2004**, *152*, 71–88.
8. Jeyakumar, S.; Arumuganathan, R. A Non-Markovian Bulk Queue with Multiple Vacations and Control Policy on Request for Re-Service. *Qual. Technol. Quant. Manag.* **2011**, *8*, 253–269. [[CrossRef](#)]
9. Haridass, M.; Arumuganathan, R. A batch service queueing system with multiple vacations, setup time and server's choice of admitting re-service. *Int. J. Oper. Res.* **2012**, *14*, 156–186. [[CrossRef](#)]
10. Krishnamoorthy, A.; Pramod, P.K.; Chakravarthy, S.R. Queues with interruptions: A survey. *Oper. Res. Decis. Theory* **2014**, *22*, 290–320. [[CrossRef](#)]
11. Ayyappan, G.; Shyamala, S. Transient solution of an $M^{[X]}/G/1$ queueing model with feedback, random breakdowns, Bernoulli schedule server vacation and random setup time. *Int. J. Oper. Res.* **2016**, *25*, 196–211. [[CrossRef](#)]
12. Choudhury, G.; Tadj, L. An M/G/1 queue with two phases of service subject to the server breakdown and delayed repair. *Appl. Math. Model.* **2009**, *33*, 2699–2709. [[CrossRef](#)]
13. Senthilnathan, B.; Jeyakumar, S. A study on the behaviour of the server breakdown without interruption in a $M^{[X]}/G(a, b)/1$ queueing system with multiple vacations and closedown time. *Appl. Math. Comput.* **2012**, *219*, 2618–2633.
14. Lakshmi, K.; Kasturi Ramanath, K. An M/G/1 two phase multi-optional retrial queue with Bernoulli feedback, non-persistent customers and breakdown and repair. *Int. J. Oper. Res.* **2014**, *19*, 78–95. [[CrossRef](#)]
15. Atencia, I. A discrete-time queueing system with server breakdowns and changes in the repair times. *Ann. Oper. Res.* **2015**, *235*, 37–49. [[CrossRef](#)]
16. Choudhury, G. A batch arrival queue with a vacation time under single vacation policy. *Comput. Oper. Res.* **2002**, *29*, 1941–1955. [[CrossRef](#)]
17. Jeyakumar, S.; Arumuganathan, R. Analysis of Single Server Retrial Queue with Batch Arrivals, Two Phases of Heterogeneous Service and Multiple Vacations with N-Policy. *Int. J. Oper. Res.* **2008**, *5*, 213–224.
18. Balasubramanian, M.; Arumuganathan, R.; Senthil Vadivu, A. Steady state analysis of a non-Markovian bulk queueing system with overloading and multiple vacations. *Int. J. Oper. Res.* **2010**, *9*, 82–103. [[CrossRef](#)]
19. Haridass, M.; Arumuganathan, R. Analysis of a batch arrival general bulk service queueing system with variant threshold policy for secondary jobs. *Int. J. Math. Oper. Res.* **2011**, *3*, 56–77. [[CrossRef](#)]
20. Choudhury, G.; Deka, M. A batch arrival unreliable server delaying repair queue with two phases of service and Bernoulli vacation under multiple vacation policy. *Qual. Technol. Quant. Manag.* **2018**, *15*, 157–186. [[CrossRef](#)]
21. Madan, K.C. A cyclic queueing system with three servers and optional two-way feedback. *Microelectron. Reliabil.* **1988**, *28*, 873–875. [[CrossRef](#)]
22. Madan, K.C. On a single server queue with two-stage heterogeneous service and deterministic server vacations. *Int. J. Syst. Sci.* **2001**, *32*, 837–844. [[CrossRef](#)]

23. Medhi, J. A Single Server Poisson Input Queue with a Second Optional Channel. *Queueing Syst.* **2002**, *42*, 239–242. [[CrossRef](#)]
24. Arumuganathan, R.; Malliga, T.J. Analysis of a bulk queue with repair of service station and setup time. *Int. J. Can. Appl. Math. Quart.* **2006**, *13*, 19–42.
25. Baruah, M.; Madan, K.C.; Eldabi, T. A Two Stage Batch Arrival Queue with Reneging during Vacation and Breakdown Periods. *Am. J. Oper. Res.* **2013**, *3*, 570–580. [[CrossRef](#)]
26. Aissani, A.; Artalejo, J. On the single server retrial queue subject to breakdowns. *Queueing Syst.* **1988**, *30*, 309–321. [[CrossRef](#)]
27. Maraghi, F.A.; Madan, K.C.; Darby-Dowman, K. Bernoulli schedule vacation queue with batch arrivals and random system breakdowns having general repair time distribution. *Int. J. Oper. Res.* **2010**, *7*, 240–256. [[CrossRef](#)]
28. Fadhil, R.; Madan, K.C.; Lukas, A.C. An $M(X)/G/1$ Queue with Bernoulli Schedule General Vacation Times, Random Breakdowns, General Delay Times and General Repair Times. *Appl. Math. Sci.* **2011**, *5*, 35–51.
29. Rajadurai, P.; Varalakshmi, M.; Saravananarajan, M.C.; Chandrasekaran, V.M. Analysis of $M^{[X]}/G/1$ retrial queue with two phase service under Bernoulli vacation schedule and random breakdown. *Int. J. Math. Oper. Res.* **2015**, *7*, 19–41. [[CrossRef](#)]
30. Jiang, T.; Xin, B. Computational analysis of the queue with working breakdowns and delaying repair under a Bernoulli-schedule-controlled policy. *J. Commun. Stat. Theory Methods* **2018**, 1–16. [[CrossRef](#)]
31. Madan, K.C. A bulk input queue with a stand-by. *S. Afr. Stat. J.* **1995**, *29*, 1–7.
32. Khalaf, R.F. Queueing Systems With Four Different Main Server's Interruptions and a Stand-By Server. *Int. J. Oper. Res.* **2014**, *3*, 49–54. [[CrossRef](#)]
33. Jain, M.; Preeti. Cost analysis of a machine repair problem with standby, working vacation and server breakdown. *Int. J. Math. Oper. Res.* **2014**, *6*, 437–451. [[CrossRef](#)]
34. Kumar, K.; Jain, M. Bi-level control of degraded machining system with two unreliable servers, multiple standbys, startup and vacation. *Int. J. Oper. Res.* **2014**, *21*, 123–142. [[CrossRef](#)]
35. Murugeswari, N.; Maragatha Sundari, S. A Standby server bulk arrival Queueing model of Compulsory server Vacation. *Int. J. Eng. Dev. Res.* **2017**, *5*, 337–341.
36. Kolledath, S.; Kumar, K.; Pippal, S. Survey on queueing models with standbys support. *Yugoslav J. Oper. Res.* **2018**, *28*, 3–20. [[CrossRef](#)]
37. Neuts, M.F. A general class of bulk queues with poisson input. *The Annals of Mathematical Statistics*, **1967**, *38*, 759–770. [[CrossRef](#)]
38. Ayyappan, G.; Sathiya, S. Non Markovian Queue with Two Types service Optional Re-service and General Vacation Distribution. *Appl. Appl. Math. Int. J.* **2016**, *11*, 504–526.

