



## An Official American Thoracic Society/American Association of Critical-Care Nurses/American College of Chest Physicians/Society of Critical Care Medicine Policy Statement: The Choosing Wisely® Top 5 List in Critical Care Medicine

Scott D. Halpern, Deborah Becker, J. Randall Curtis, Robert Fowler, Robert Hyzy, Lewis J. Kaplan, Nishi Rawat, Curtis N. Sessler, Hannah Wunsch, and Jeremy M. Kahn; on behalf of the Choosing Wisely Taskforce

THIS OFFICIAL STATEMENT WAS APPROVED BY THE AMERICAN THORACIC SOCIETY (ATS), JUNE 2014; THE AMERICAN ASSOCIATION OF CRITICAL-CARE NURSES (AACN), MARCH 2014; THE AMERICAN COLLEGE OF CHEST PHYSICIANS (ACCP), APRIL 2014; AND THE SOCIETY OF CRITICAL CARE MEDICINE (SCCM), MARCH 2014

**Rationale:** The high costs of health care in the United States and other developed nations are attributable, in part, to overuse of tests, treatments, and procedures that provide little to no benefit for patients. To improve the quality of care while also combating this problem of cost, the American Board of Internal Medicine Foundation developed the Choosing Wisely Campaign, tasking professional societies to develop lists of the top five medical services that patients and physicians should question.

**Objectives:** To present the Critical Care Societies Collaborative's Top 5 list in Critical Care Medicine and describe its development.

**Methods:** Each professional society in the Collaborative nominated members to the Choosing Wisely task force, which established explicit criteria for evaluating candidate items, generated lists of items, performed literature reviews on each, and sought external input from content experts. Task force members narrowed the list to the Top 5 items using a standardized scoring system based on each item's likely overall impact and merits on the five explicit criteria.

**Measurements and Main Results:** From an initial list of 58 unique recommendations, the task force proposed a Top 5 list that was ultimately endorsed by each Society within the Collaborative. The five recommendations are: (1) do not order diagnostic tests at regular intervals (such as every day), but rather in response to specific clinical questions; (2) do not transfuse red blood cells in hemodynamically stable, nonbleeding ICU patients with an Hb concentration greater than 7 g/dl; (3) do not use parenteral nutrition in adequately nourished critically ill patients within the first 7 days of an ICU stay; (4) do not deeply sedate mechanically ventilated patients without a specific indication and without daily attempts to lighten sedation; and (5) do not continue life support for patients at high risk for death or severely impaired functional recovery without offering patients and their families the alternative of care focused entirely on comfort.

**Conclusions:** These five recommendations provide a starting point for clinicians and patients to make decisions leading to higher-quality, lower-cost care. Future work is needed to promote adherence to these recommendations and to develop additional ways for intensive care clinicians to take leadership in reining in health-care costs.

**Keywords:** Choosing Wisely; critical care; quality improvement; cost effectiveness

### Introduction

In 2012, the United States spent \$2.87 trillion on health care, accounting for 18.3% of its Gross Domestic Product. In recent years, roughly \$100 billion, or 0.66% of U.S. Gross Domestic Product, has been spent on the provision of critical care (1). For this reason, hospital leaders, payers, and policy

makers interested in opportunities for cost reduction frequently focus on the intensive care unit (ICU).

One factor contributing to the high cost of critical care may be clinician decision making at the bedside. Data indicate that intensivists vary widely in their daily discretionary spending, with no correlation between physicians' spending and their

patients' outcomes (2). More broadly, evidence suggests that physicians' decisions contribute importantly to the high and rising costs of health care in the United States (3–5). Thus, in an effort to improve quality and reduce costs, the American Board of Internal Medicine Foundation launched the Choosing Wisely Campaign in 2011 to identify tests or interventions

Correspondence and requests for reprints should be addressed to Scott D. Halpern, M.D., Ph.D., University of Pennsylvania, 719 Blockley Hall, 423 Guardian Drive, Philadelphia, PA 19104. E-mail: shalpern@exchange.upenn.edu

Am J Respir Crit Care Med Vol 190, Iss 7, pp 818–826, Oct 1, 2014

Copyright © 2014 by the American Thoracic Society

DOI: 10.1164/rccm.201407-1317ST

Internet address: www.atsjournals.org

## Contents

### Introduction

#### Methods

**Origins of the Critical Care Societies Collaborative Choosing Wisely Initiative Development of Draft Recommendations Vetting and Endorsement**

#### Results

**Recommendation #1: Do Not Order Diagnostic Tests at Regular Intervals (Such as Every Day), But Rather in Response to Specific Clinical Questions**

**Recommendation #2: Do Not Transfuse RBCs in Hemodynamically Stable, Nonbleeding ICU Patients with an Hb Concentration Greater Than 7 g/dl**

**Recommendation #3: Do Not Use Parenteral Nutrition in Adequately Nourished Critically Ill Patients within the First 7 Days of an ICU Stay**

**Recommendation #4: Do Not Deeply Sedate Mechanically Ventilated Patients without a Specific Indication and without Daily Attempts to Lighten Sedation**

**Recommendation #5: Do Not Continue Life Support for Patients at High Risk for Death or Severely Impaired Functional Recovery without Offering Patients and Their Families the Alternative of Care Focused Entirely on Comfort**

#### Discussion

“that physicians and patients should question” (6, 7).

The seeds of Choosing Wisely Campaign stem from the Physician Charter for Medical Professionalism produced by the American Board of Internal Medicine Foundation, the American College of Physicians, and the European Federation of Internal Medicine (8). The Charter moved beyond older conceptualizations of medical professionalism by noting that “physicians are *required* to provide health care that is based on the wise and cost-effective management of limited clinical resources [emphasis added].”

Based on this model, medical ethicist Howard Brody subsequently charged physicians to “seize the moral high ground” and take leadership in declaring what tests and interventions should be used less commonly (9). He noted that if physicians failed to lead in this area, they would be forced to follow cost-cutting measures imposed by politicians or payers (9). Brody recommended that each professional society appoint a panel to decide that specialty’s “Top 5” list of “diagnostic tests or treatments that are very commonly ordered by members of that specialty, that are among the most expensive services provided, and that have been shown by the currently available evidence not to provide any meaningful benefit to at least some major categories of patients” (9). This policy statement describes this initiative in critical care medicine, discusses the process and rationale for selecting each of the Top 5 items in critical care, and discusses important next steps for clinicians to lead the way toward the more efficient provision of critical care.

## Methods

### Origins of the Critical Care Societies Collaborative Choosing Wisely Initiative

This initiative was proposed by the first author and championed by the last author in his role as Chair of the American Thoracic Society’s Quality Improvement Committee, in June 2012. A Memorandum of Understanding was signed in September of 2012 by the four societies in the Critical Care Societies Collaborative: the American Association of Critical-Care Nurses, the American College of Chest Physicians, the American Thoracic Society, and the Society of Critical Care Medicine. Each of these societies was invited to nominate members to join a task force charged with developing the critical care Top 5 list. The final task force included the 10 authors, representing all four societies and spanning the clinical disciplines of critical care medicine, including internal medicine, surgery, anesthesiology, emergency medicine, and critical care nursing.

This task force was charged with developing a list of five “low-value” practices in the field of critical care medicine—that is, practices that are unlikely to provide benefits to patients, and even if they did, the benefits would be

insufficient to justify the expenditures (10). The charge was not to develop a clinical practice guideline, because the methods, conducted in accordance with other lists developed as part of the Choosing Wisely campaign, did not adhere to the guideline standards of the Institute of Medicine.

### Development of Draft Recommendations

During an initial meeting in Boston, Massachusetts in September 2012, we established task force goals and criteria for evaluating recommendations. The agreed-on evaluative criteria, which members could weigh at their discretion in proposing items, are listed in Table 1. We selected these criteria to help us focus on recommendations that were likely to be (1) correct (i.e., supported by evidence), (2) important (i.e., have high prevalence, significance for patients, and associated costs), (3) germane to practitioners of critical care medicine (as opposed to medical professionals more generally), and (4) novel.

Next, task force members submitted lists of suggested items. The first author accumulated all submitted items, removed duplicates, and organized the items into five broad categories: diagnostic tests, therapies, procedures, end-of-life care, and structures of care. There were 58 unique items on this “initial” list.

During subsequent teleconferences, members discussed each item in turn, using an iterative consensus process to narrow this list through consensus to 16 items. Task force members then formally prioritized these 16 items, such that items were retained if a majority of task force members believed the item had a reasonable chance of making the final Top 5 list. Nine items were retained, and, for each, one or two members were assigned to conduct in-depth evidence reviews and consult with external content experts.

The nine evidence-informed summaries were reviewed by teleconference in November 2012, during which the group deliberated each item’s performance on the five evaluation criteria (Table 1). Members subsequently rated each item independently using a National Institutes of Health–style scale from 1 to 9, where 1 is the best possible rating. Ratings were provided for each of the five criteria, and an overall impact rating was provided for each item.

**Table 1.** Criteria on Which Each Proposed Item Was Evaluated

1. Strength of evidence: How sure are we that our suggestion is correct?
2. Prevalence: How commonly do we think this issue arises?
3. Aggregate cost: How large are the anticipated cost savings if this suggestion is adhered to?
4. Relevance: To what extent is this either a “core” or “unique” part of the practice of critical care medicine, as opposed to an ancillary activity or part of good medical practice more generally?
5. Innovation: How much does this suggestion represent an advance (i.e., “move the needle”) rather than recapitulate known best practices or previously published clinical guidelines?

Task force members were instructed that the overall rating need not be an average of the five criteria ratings, enabling members to weigh the criteria as they deemed most appropriate.

Members submitted their ratings to the first author, who summed and averaged the five criterion scores and the overall impact scores. The items with the best overall impact scores were retained in the “penultimate” list. All members then contributed textual edits for these five items as well as to the one-paragraph summaries accompanying each item.

### Vetting and Endorsement

This penultimate list was submitted to the four societies’ executive committees for review. The executive committees sought external feedback from other content experts and provided written suggestions to the task force. The task force considered all suggestions and revised items’ descriptions based on this feedback. The first author and members of the societies’ executive committees resolved disagreements by consensus and compromise via teleconference. This process led to the inclusion of new evidence that informed the wording of the five items, but the actual recommendations themselves and the core elements of the rationale supporting each item were retained. The final list was then sent back to four societies’ executive committees for consideration of final endorsement. Each society endorsed the list in February 2013. This final list was submitted to the Choosing Wisely Campaign in March 2013, approved in May 2013, and published by the Campaign in January 2014.

### Results

The five recommendations of the Critical Care Societies Collaborative Choosing

Wisely task force are listed in Table 2. Each recommendation is accompanied by a single paragraph providing a brief rationale for the recommendation. In the following sections we provide more detailed analyses of the reasons we chose each of these items, the evidence supporting the items, and of how each item scored on certain evaluative criteria.

#### **Recommendation #1: Do Not Order Diagnostic Tests at Regular Intervals (Such as Every Day), But Rather in Response to Specific Clinical Questions**

Whether due to convenience or perceived indications, intensivists commonly order diagnostic tests such as complete blood counts, serum chemistries, chest radiographs, and arterial blood gas analyses on a routine basis (e.g., daily) (11). However, all available evidence suggests that routine diagnostic evaluations, rather than tests ordered to answer specific clinical questions, carry measurable burdens and costs for patients, hospitals, and the health-care system (12–14). For example, routine phlebotomy causes iatrogenic anemia, leading ICU patients to receive extra transfusions of red blood cells (RBCs), raising costs and conferring risks of alloimmunization, transfusion-related acute lung injury, transfusion-associated circulatory overload, infection, and organ failure (15). Computer order entry that incorporates decision aids has been shown to reduce duplicative order sets and inappropriate daily laboratory testing, thereby reducing phlebotomy rates and enabling more efficient use of health-care personnel (13, 14).

Similarly, evidence suggests that routine chest radiography and computed tomography scanning are rarely indicated and may be harmful (16, 17). Unexpected findings on chest radiographs were noted in less than 6% of the 2,457 daily routine

radiographs that were ordered in 754 consecutive ICU patients in a mixed medical-surgical ICU (18). In fewer than half of these cases did the finding prompt a change in management (18). Relatively stable patients are particularly unlikely to benefit from routine chest radiography, a practice which accrues increased cost and radiation dose and more interventions (19–21). A protocol to reduce chest radiograph ordering decreased use by up to 36% (22).

Among all items considered by the task force, this recommendation was among the most highly ranked regarding the prevalence of the problem, the potential for aggregate cost savings, and the applicability of the issue to the practice of critical care. This recommendation scored more modestly on the criteria of the strength of evidence and the innovation of the recommendation.

#### **Recommendation #2: Do Not Transfuse RBCs in Hemodynamically Stable, Nonbleeding ICU Patients with an Hb Concentration Greater Than 7 g/dl**

Many critically ill patients receive RBC transfusions during their ICU stay. The largest national study to date, conducted in 2001 in a mixed medical-surgical population, found that 44% of ICU patients received at least one transfusion, with an average of 4.6 units per patient (23). By contrast, a more recent study of medical ICU patients in Maryland found rates of approximately 18% in 2007 (24). Although RBC transfusion is indicated in severe anemia and hemorrhagic shock, the majority of transfusions appear to be given to patients with asymptomatic anemia. For example, in the national study, the average Hb concentration at transfusion was 8.6 g/dl, with 65% of patients receiving transfusions at Hb concentrations above 7.0 g/dl (23).

**Table 2.** Final Top 5 List Approved by Societies' Leadership and Included in Choosing Wisely Campaign

- 1. Do not order diagnostic tests at regular intervals (such as every day), but rather in response to specific clinical questions.**  
Many diagnostic studies (including chest radiographs, arterial blood gases, blood chemistries and counts, and ECGs) are ordered at regular intervals (e.g., daily). Compared with a practice of ordering tests only to help answer clinical questions, or when doing so will affect management, the routine ordering of tests increases health-care costs, does not benefit patients, and may in fact harm patients. Potential harms include anemia due to unnecessary phlebotomy, which may necessitate risky and costly transfusion, and the aggressive work-up of incidental and nonpathological results found on routine studies.
- 2. Do not transfuse RBCs in hemodynamically stable, nonbleeding ICU patients with an Hb concentration greater than 7 mg/dl.**  
Most red blood cell transfusions in the ICU are for benign anemia rather than acute bleeding that causes hemodynamic compromise. For all patient populations in which it has been studied, transfusing RBCs at a threshold of 7 mg/dl is associated with similar or improved survival, fewer complications, and reduced costs compared with higher transfusion triggers. More aggressive transfusion may also limit the availability of a scarce resource. It is possible that different thresholds may be appropriate in patients with acute coronary syndromes, although most observational studies suggest harms of aggressive transfusion even among such patients.
- 3. Do not use parenteral nutrition in adequately nourished critically ill patients within the first 7 d of an ICU stay.**  
For patients who are adequately nourished before ICU admission, parenteral nutrition initiated within the first 7 d of an ICU stay has been associated with harm, or at best no benefit, in terms of survival and length of stay in the ICU. Early parenteral nutrition is also associated with unnecessary costs. These findings are true even among patients who cannot tolerate enteral nutrition. Evidence is mixed regarding the effects of early parenteral nutrition on nosocomial infections. For patients who are severely malnourished directly before their ICU admission, there may be benefits to earlier parenteral nutrition.
- 4. Do not deeply sedate mechanically ventilated patients without a specific indication and without daily attempts to lighten sedation.**  
Many mechanically ventilated ICU patients are deeply sedated as a routine practice despite evidence that using less sedation reduces the duration of mechanical ventilation and ICU and hospital length of stay. Several protocol-based approaches can safely limit deep sedation, including the explicit titration of sedation to the lightest effective level, the preferential administration of analgesic medications before initiating anxiolytics, and the performance of daily interruptions of sedation in appropriately selected patients receiving continuous sedative infusions. Although combining these approaches may not improve outcomes compared to one approach alone, each has been shown to improve patient outcomes compared with approaches that provide deeper sedation for ventilated patients.
- 5. Do not continue life support for patients at high risk for death or severely impaired functional recovery without offering patients and their families the alternative of care focused entirely on comfort.**  
Patients and their families often value the avoidance of prolonged dependence on life support. However, many of these patients receive aggressive life-sustaining therapies, in part due to clinicians' failures to elicit patients' values and goals and to provide patient-centered recommendations. Routinely engaging high-risk patients and their surrogate decision makers in discussions about the option of forgoing life-sustaining therapies may promote patients' and families' values, improve the quality of dying, and reduce family distress and bereavement. Even among patients pursuing life-sustaining therapy, initiating palliative care simultaneously with ongoing disease-focused therapy may be beneficial.

*Definition of abbreviations:* ICU = intensive care unit; RBC = red blood cell.

The Choosing Wisely Top 5 List in Critical Care Medicine originally appeared on the Choosing Wisely website (<http://www.choosingwisely.org/doctor-patient-lists/american-college-of-chest-physicians-and-american-thoracic-society/>).

These transfusion thresholds are not supported by current evidence. Several randomized clinical trials (RCTs) have found that using a transfusion trigger of an Hb concentration of less than 7 g/dl was associated with either equivalent or lower mortality compared with a transfusion trigger of less than 9 g/dl (25–27). Based on this evidence, a recent multisociety clinical practice guideline strongly recommends an Hb concentration transfusion threshold of less than 7 g/dl in hemodynamically stable patients without trauma or acute bleeding (28). Some question remains regarding appropriate transfusion thresholds for patients who are actively bleeding or with acute coronary syndromes. However, a recent RCT showed that even patients with upper gastrointestinal bleeding fare as well or better with restrictive transfusion thresholds (27). Additionally, a number of observational studies (29–31), a recent metaanalysis (32), and a small RCT (33) all

suggest that restrictive transfusion strategies are superior for anemic patients with acute coronary syndromes including myocardial infarction. Given the preponderance of evidence that lower transfusion thresholds are uniformly safe, and the absence of evidence suggesting benefit to a transfusion threshold greater than 7 g/dl in any particular patient population (34, 35), intensivists might reasonably use this threshold in all critically ill patients.

Task force members rated this item extremely highly on the strength of the evidence. Members also rated it highly on its centrality to the practice of critical care medicine. The item was rated less highly on prevalence and innovation, given the perception of many task force members that RBC transfusion use may already be in decline. However, a study published after the task force voted found that despite publication of the landmark RCTs,

transfusion practices have not declined appreciably (24). Such evidence would only have enhanced the task force's rating of this item.

### **Recommendation #3: Do Not Use Parenteral Nutrition in Adequately Nourished Critically Ill Patients within the First 7 Days of an ICU Stay**

Enteral nutrition is the optimal method for delivering nutritional support during critical illness (36–39). For patients who are adequately nourished before ICU admission, early parenteral nutrition—that is, parenteral nutrition initiated within the first 7 days of an ICU stay—is associated with equal or worse survival and length of stay in the ICU compared with no nutrition (36, 37, 40–42). Early parenteral nutrition also worsens adiposity in normally muscle-laden areas (43) and carries unnecessary costs (36, 38, 40, 44).



Recent evidence is mixed regarding the effects of early parenteral nutrition on poorly validated surrogate endpoints such as nosocomial infections and the duration of mechanical ventilation (40–42). Thus, although the optimal timing of parenteral nutrition is likely to be further informed as new data emerge, there appears to be no rationale for administering early parenteral nutrition for patients who are adequately nourished shortly before their ICU admission (45). For patients with preexisting protein-calorie malnutrition, it is possible that earlier parenteral nutrition may be helpful if the delivery of enteral nutrition is insufficient to meet nutritional needs (36, 39, 46). Despite this evidence base, recent data suggest that parenteral nutrition is commonly used early in critical illness (47).

Among the recommendations considered, this item scored highly with respect to its prevalence in both academic and community ICUs and in terms of expected cost savings and innovation. The recommendation scored less well with respect to the strength of evidence and its specificity to the practice of critical care medicine.

**Recommendation #4: Do Not Deeply Sedate Mechanically Ventilated Patients without a Specific Indication and without Daily Attempts to Lighten Sedation**

Ensuring patient safety and comfort during critical illness is an important goal for critical care clinicians. However, accumulating evidence suggests substantial risk from excessively deep or unnecessarily prolonged sedation. Various structured approaches that share the common goal of minimizing sedation have been linked to shorter duration of mechanical ventilation, shorter ICU and hospital length of stay, fewer tracheostomies, fewer tests to investigate altered mental status, and reduced delirium (48, 49).

Basic tenets of sedation and analgesia management applied in many protocols include establishing a specific indication for sedation, setting a target level of consciousness, monitoring sedation and pain using validated scales, and titrating medications to achieve therapeutic goals while maintaining the lightest effective level of sedation (48–62). Because administration of sedative medications by continuous intravenous infusion has been linked to

unnecessarily deep sedation and delayed recovery, some approaches emphasize intermittent dosing (50, 63). Other effective approaches target the treatment of pain before giving sedative agents or a structured evaluation and management of both pain and agitation (64–66). For patients who are more deeply sedated, scheduling daily interruptions of sedative and analgesic infusions until patients are alert may also facilitate earlier extubation (61, 67, 68). Although head-to-head, randomized comparisons of various techniques have not demonstrated the superiority of any specific approach, approaches that minimize sedation are consistently found to be superior to approaches yielding deeper sedation.

Of the items considered by the task force, the recommendation to minimize sedation was considered to be highly germane to the practice of critical care medicine. Additionally, this recommendation scored highly regarding the strength of evidence and potential cost savings, with the latter attributable primarily to decreased lengths of stay. Among the final items considered, this recommendation received average ratings for its prevalence and innovation.

**Recommendation #5: Do Not Continue Life Support for Patients at High Risk for Death or Severely Impaired Functional Recovery without Offering Patients and Their Families the Alternative of Care Focused Entirely on Comfort**

Patients and their families often value the avoidance of prolonged dependence on life support (69, 70). However, many seriously ill patients receive aggressive life-sustaining therapies (71–73), which has been associated with reduced quality (74, 75) and perhaps quantity (76–78) of life near its end. Prolonged aggressive care in the ICU is also associated with long-lasting pathological bereavement among family members (75, 79–81), contravening most patients' strong desires not to burden their loved ones (82–84).

Much of this aggressive care may be attributable to variability among ICU clinicians in their practices of eliciting patients' values and goals and in their provision of recommendations (85–87). Clinician variability, in turn, may contribute to the marked geographic variation across the United States in the

costs of care in the last year of life, without corresponding differences in patients' preferences for the aggressiveness of care (88, 89). Routinely engaging patients at high risk of death and their surrogate decision makers in discussions about the option of forgoing life-sustaining therapies may promote patients' and families' values, improve the quality of dying, and reduce family distress and bereavement (90–94). Even among patients pursuing life-sustaining therapy, initiating palliative care simultaneously with ongoing disease-focused therapy may be beneficial, although the evidence for this is strongest in the outpatient setting (76, 77).

Among all items considered by the task force, this item was the highest rated regarding its centrality to the practice of critical care medicine. This item also scored highly in terms of the prevalence with which this issue arises. Criteria on which it scored less well included the strength of the evidence supporting the recommendation, which is mostly observational in nature, and the anticipated cost savings, which may be modest (95, 96).

## Discussion

When this Top 5 list of common ICU practices to be questioned was published by the Choosing Wisely Campaign in January 2014, it joined more than 50 similar lists developed by other professional societies. In addition to signifying the Critical Care Societies Collaborative's commitment to this national campaign, there are several unique strengths of our process.

First, this Top 5 list emanated from a collaborative effort by four unique professional societies. The shared consensus among multiple societies that these recommendations are appropriate supports the credibility of the product. Second, this is the only list to have involved a professional nursing society in its development, reflecting the intrinsically interprofessional nature of critical care. Third, we began by establishing explicit criteria on which we would evaluate the items, lending transparency and consistency to a year-long process. Finally, we used an evaluative process long promoted by study sections at the National Institutes of Health and elsewhere in which we encouraged task force members to weight the individual criteria as they deemed most appropriate in arriving at their final scores.

Despite these strengths, our work is only a first step in what we hope will be a much broader effort to improve the quality, efficiency, and overall value of critical care delivery. The costs of critical care, like most inpatient services, are relatively fixed in the short term (97) and are primarily driven by the large numbers of operational ICU beds in the United States (98) and the clinicians employed to staff these beds. Thus, although adherence to our recommendations would improve patient outcomes and cut costs, dramatic reductions in the costs of critical care may require strategic contractions of the supply of ICU beds or prominent changes in their use. Although recent studies suggest that such contractions might be undertaken without harming patients (99–102), the task force considered this evidence base too nascent to yield specific recommendations regarding the supply and use of ICU beds.

A second limitation is that Choosing Wisely lists are restricted to five items, a deliberate decision intended to make the lists actionable and to focus on recommendations for which the greatest consensus exists (9). Yet, as a consequence, we necessarily excluded other recommendations that might improve the value of critical care delivery. Among the items that just missed inclusion in the Top 5 list were several related to antimicrobial use, computed tomography, and routine arterial and central venous catheterization. The task force scored each of these items less strongly, typically due to one or two core limitations. For example, task force members agreed that several forms of computed tomography were overused in critical care medicine but discounted scores for these items because they were not believed

to be unique matters for our profession. Such recommendations were also considered less innovative because numerous other professional societies have recommended reducing computed tomography and other imaging modalities as part of the Choosing Wisely Campaign (103).

Thus, although this list is by no means exhaustive, we believe it provides five key ways for intensivist clinicians to begin improving the quality and reining in the costs of their care. Furthermore, by formalizing specific approaches for clinicians to more earnestly adhere to their duties to use resources judiciously, the list may have spillover effects such that critical care clinicians more routinely consider costs and effectiveness in their daily decision making.

A third limitation, which pertains to the Choosing Wisely Campaign more generally, is that in many cases, the factors driving use of low-value services are unknown. Much but not all overuse is attributable to individual decisions made by clinicians and patients. To the extent that service provision is driven by factors outside the control of individual clinicians and patients, interventions other than Choosing Wisely recommendations will be needed. For example, systems-level interventions, such as changes in physician and nurse staffing patterns, may yield substantial improvements in the value of health-care delivery. By contrast, we focused our recommendations on tests or therapies that are largely under clinicians' control, as these are the care processes most likely to be impacted by published lists in publications targeting clinicians.

With the publication of this list, the Critical Care Societies Collaborative must begin the hard work of promoting adherence to these recommendations.

Indeed, our five recommendations all reflect best practices, so clearly a key problem is that of adherence. Several options are available that go beyond traditional dissemination by passive education. For example, there is considerable interest in reforming provider reimbursement structures such that disincentives exist for ordering unnecessary tests or performing unnecessary procedures. Reducing service use through financial disincentives holds considerable promise due to physicians' natural loss aversion, such that, everything else being equal, a reimbursement reduction for ordering an unnecessary test is likely to be more effective than a reimbursement increase for ordering an indicated test (104). However, changing reimbursement structures are unlikely to succeed unless physicians are simultaneously made aware of exactly which low-value tests and procedures to avoid (105). Thus, disseminating the Top 5 lists of the Choosing Wisely Campaign may usefully guide physician reimbursement reforms.

More directly, these five recommendations provide a starting point for physicians, nurses, allied health professionals, patients, and patients' families to make decisions leading to higher-quality, lower-cost critical care. Future work is needed to develop additional ways for clinicians to take leadership in reining in health-care costs and to promote adherence to such recommendations. In the meantime, reducing the low-value care identified in these recommendations is an important first step, enabling our field to help stabilize health-care costs and thereby promote the availability of health care for all patients. ■

---

This statement was prepared by the Choosing Wisely Taskforce

#### Members of the Taskforce:

SCOTT D. HALPERN, M.D., Ph.D. (*Chair*)  
 DEBORAH BECKER, Ph.D., R.N.  
 J. RANDALL CURTIS, M.D., M.P.H.  
 ROBERT FOWLER, M.D.  
 ROBERT HYZY, M.D.  
 LEWIS J. KAPLAN, M.D.  
 NISHI RAWAT, M.D.  
 CURTIS N. SESSLER, M.D.  
 HANNAH WUNSCH, M.D., M.Sc.  
 JEREMY M. KAHN, M.D., M.Sc.

**Author Disclosures:** J.M.K. has received research support from Cerner Corp. (\$1–4,999). S.D.H., D.B., J.R.C., R.F., R.H., L.J.K., N.R., C.N.S., and H.W. reported no relevant financial relationships with commercial interests.

**Acknowledgment:** The published version of the Choosing Wisely Top 5 List in Critical Care Medicine, as well as the other published Choosing Wisely Lists from other disciplines, can be found at [www.choosingwisely.org](http://www.choosingwisely.org). The authors thank

Jessica Wisk, formerly of the American Thoracic Society, for her remarkable efforts in support of this initiative during its formative stages. They also thank Jenny Nemkovich of the American College of Chest Physicians and Judy Corn of the American Thoracic Society for their support throughout this process. They also thank content experts, including Thomas Bleck, M.D., Allan Garland, M.D., M.A., Daren Heyland, M.D., Bernard Lo, M.D., Gordon D. Rubenfeld, M.D., M.Sc., and Douglas B. White, M.D., M.A.S.

## References

- Halpern NA, Pastores SM. Critical care medicine in the United States 2000-2005: an analysis of bed numbers, occupancy rates, payer mix, and costs. *Crit Care Med* 2010;38:65-71.
- Garland A, Shaman Z, Baron J, Connors AF Jr. Physician-attributable differences in intensive care unit costs: a single-center study. *Am J Respir Crit Care Med* 2006;174:1206-1210.
- Berwick DM, Hackbarth AD. Eliminating waste in US health care. *JAMA* 2012;307:1513-1516.
- Campbell EG, Regan S, Gruen RL, Ferris TG, Rao SR, Cleary PD, Blumenthal D. Professionalism in medicine: results of a national survey of physicians. *Ann Intern Med* 2007;147:795-802.
- Kale MS, Bishop TF, Federman AD, Keyhani S. "Top 5" lists top \$5 billion. *Arch Intern Med* 2011;171:1856-1858.
- Cassel CK, Guest JA. Choosing wisely: helping physicians and patients make smart decisions about their care. *JAMA* 2012;307:1801-1802.
- American Board of Internal Medicine Foundation. Choosing Wisely [accessed 2014 Jan 23]. Available from: <http://www.choosingwisely.org/>
- Sox HC, Blank L, Cohen J, et al.; ABIM Foundation. American Board of Internal Medicine; ACP-ASIM Foundation. American College of Physicians-American Society of Internal Medicine; European Federation of Internal Medicine. Medical professionalism in the new millennium: a physician charter. *Ann Intern Med* 2002;136:243-246.
- Brody H. Medicine's ethical responsibility for health care reform—the Top Five list. *N Engl J Med* 2010;362:283-285.
- Baker DW, Qaseem A, Reynolds PP, Gardner LA, Schneider EC; American College of Physicians Performance Measurement Committee. Design and use of performance measures to decrease low-value services and achieve cost-conscious care. *Ann Intern Med* 2013;158:55-59.
- Ezzie ME, Aberegg SK, O'Brien JM Jr. Laboratory testing in the intensive care unit. *Crit Care Clin* 2007;23:435-465.
- Peixoto AA, Meneses FA, Barbosa BP, Pessoa LFP, Melo RHT, Fideles GM. Laboratory routine in the ICU: a practice to be abolished? *Crit Care* 2013;17:1.
- May TA, Clancy M, Critchfield J, Ebeling F, Enriquez A, Gallagher C, Genevro J, Kloo J, Lewis P, Smith R, et al. Reducing unnecessary inpatient laboratory testing in a teaching hospital. *Am J Clin Pathol* 2006;126:200-206.
- Collins SR, Nuzum R, Rustgi SD, Mika S, Schoen C, Davis K. How health care reform can lower the costs of insurance administration. *Issue Brief (Commonw Fund)* 2009;61:1-19.
- Carson JL, Grossman BJ, Kleinman S, Tinmouth AT, Marques MB, Fung MK, Holcomb JB, Illoh O, Kaplan LJ, Katz LM, et al.; Clinical Transfusion Medicine Committee of the AABB. Red blood cell transfusion: a clinical practice guideline from the AABB. *Ann Intern Med* 2012;157:49-58.
- Ganapathy A, Adhikari NK, Spiegelman J, Scales DC. Routine chest x-rays in intensive care units: a systematic review and meta-analysis. *Crit Care* 2012;16:R68.
- Smith-Bindman R, Lipson J, Marcus R, Kim KP, Mahesh M, Gould R, Berrington de González A, Miglioretti DL. Radiation dose associated with common computed tomography examinations and the associated lifetime attributable risk of cancer. *Arch Intern Med* 2009;169:2078-2086.
- Graat ME, Choi G, Wolthuis EK, Korevaar JC, Spronk PE, Stoker J, Vroom MB, Schultz MJ. The clinical value of daily routine chest radiographs in a mixed medical-surgical intensive care unit is low. *Crit Care* 2006;10:R11.
- Strain DS, Kinasewitz GT, Vereen LE, George RB. Value of routine daily chest x-rays in the medical intensive care unit. *Crit Care Med* 1985;13:534-536.
- Krivopal M, Shlobin OA, Schwartzstein RM. Utility of daily routine portable chest radiographs in mechanically ventilated patients in the medical ICU. *Chest* 2003;123:1607-1614.
- Wang ML, Schuster KM, Bhattacharya B, Maung AA, Kaplan LJ, Davis KA. Repositioning endotracheal tubes in the intensive care unit: depth changes poorly correlate with postrepositioning radiographic location. *J Trauma Acute Care Surg* 2013;75:146-149.
- Price MB, Grant MJ, Welkie K. Financial impact of elimination of routine chest radiographs in a pediatric intensive care unit. *Crit Care Med* 1999;27:1588-1593.
- Corwin HL, Gettinger A, Pearl RG, Fink MP, Levy MM, Abraham E, MacIntyre NR, Shabot MM, Duh MS, Shapiro MJ. The CRIT Study: anemia and blood transfusion in the critically ill—current clinical practice in the United States. *Crit Care Med* 2004;32:39-52.
- Murphy DJ, Needham DM, Netzer G, Zeger SL, Colantuoni E, Ness P, Pronovost PJ, Berenholtz SM. RBC transfusion practices among critically ill patients: has evidence changed practice? *Crit Care Med* 2013;41:2344-2353.
- Hébert PC, Wells G, Blajchman MA, Marshall J, Martin C, Pagliarello G, Tweeddale M, Schweitzer I, Yetisir E. A multicenter, randomized, controlled clinical trial of transfusion requirements in critical care. Transfusion Requirements in Critical Care Investigators, Canadian Critical Care Trials Group. *N Engl J Med* 1999;340:409-417.
- Walsh TS, Boyd JA, Watson D, Hope D, Lewis S, Krishan A, Forbes JF, Ramsay P, Pearse R, Wallis C, et al.; RELIEVE Investigators. Restrictive versus liberal transfusion strategies for older mechanically ventilated critically ill patients: a randomized pilot trial. *Crit Care Med* 2013;41:2354-2363.
- Villanueva C, Colomo A, Bosch A, Concepción M, Hernandez-Gea V, Aracil C, Graupera I, Poca M, Alvarez-Urturi C, Gordillo J, et al. Transfusion strategies for acute upper gastrointestinal bleeding. *N Engl J Med* 2013;368:11-21.
- Napolitano LM, Kurek S, Luchette FA, Corwin HL, Barie PS, Tisherman SA, Hebert PC, Anderson GL, Bard MR, Bromberg W, et al.; American College of Critical Care Medicine of the Society of Critical Care Medicine; Eastern Association for the Surgery of Trauma Practice Management Workgroup. Clinical practice guideline: red blood cell transfusion in adult trauma and critical care. *Crit Care Med* 2009;37:3124-3157.
- Rao SV, Jollis JG, Harrington RA, Granger CB, Newby LK, Armstrong PW, Moliterno DJ, Lindblad L, Pieper K, Topol EJ, et al. Relationship of blood transfusion and clinical outcomes in patients with acute coronary syndromes. *JAMA* 2004;292:1555-1562.
- Murphy GJ, Reeves BC, Rogers CA, Rizvi SI, Culliford L, Angelini GD. Increased mortality, postoperative morbidity, and cost after red blood cell transfusion in patients having cardiac surgery. *Circulation* 2007;116:2544-2552.
- Garfinkle M, Lawler PR, Filion KB, Eisenberg MJ. Red blood cell transfusion and mortality among patients hospitalized for acute coronary syndromes: a systematic review. *Int J Cardiol* 2013;164:151-157.
- Chatterjee S, Wetterslev J, Sharma A, Lichstein E, Mukherjee D. Association of blood transfusion with increased mortality in myocardial infarction: a meta-analysis and diversity-adjusted study sequential analysis. *JAMA Intern Med* 2013;173:132-139.
- Cooper HA, Rao SV, Greenberg MD, Rumsey MP, McKenzie M, Alcorn KW, Panza JA. Conservative versus liberal red cell transfusion in acute myocardial infarction (the CRIT Randomized Pilot Study). *Am J Cardiol* 2011;108:1108-1111.
- Desjardins P, Turgeon AF, Tremblay MH, Lauzier F, Zarychanski R, Boutin A, Moore L, McIntyre LA, English SW, Rigamonti A, et al. Hemoglobin levels and transfusions in neurocritically ill patients: a systematic review of comparative studies. *Crit Care* 2012;16:R54.
- Marik PE, Corwin HL. Efficacy of red blood cell transfusion in the critically ill: a systematic review of the literature. *Crit Care Med* 2008;36:2667-2674.
- McClave SA, Martindale RG, Vanek VW, McCarthy M, Roberts P, Taylor B, Ochoa JB, Napolitano L, Cresci G; A.S.P.E.N. Board of Directors; American College of Critical Care Medicine; Society of Critical Care Medicine. Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *JPEN J Parenter Enteral Nutr* 2009;33:277-316.
- Singer P, Berger MM, Van den Berghe G, Biolo G, Calder P, Forbes A, Griffiths R, Kreyman G, Leverve X, Pichard C, et al. ESPEN Guidelines on Parenteral Nutrition: intensive care. *Clin Nutr* 2009;28:387-400.



38. Heyland DK, Dhaliwal R, Drover JW, Gramlich L, Dodek P; Canadian Critical Care Clinical Practice Guidelines Committee. Canadian clinical practice guidelines for nutrition support in mechanically ventilated, critically ill adult patients. *JPEN J Parenter Enteral Nutr* 2003;27:355–373.
39. Kreyman KG, Berger MM, Deutz NE, Hiesmayr M, Jolliet P, Kazandjiev G, Nitenberg G, van den Berghe G, Wernerman J, Ebner C, et al.; DGEM (German Society for Nutritional Medicine); ESPEN (European Society for Parenteral and Enteral Nutrition). ESPEN Guidelines on Enteral Nutrition: intensive care. *Clin Nutr* 2006;25: 210–223.
40. Casaer MP, Mesotten D, Hermans G, Wouters PJ, Schetz M, Meyfroidt G, Van Cromphaut S, Ingels C, Meersseman P, Muller J, et al. Early versus late parenteral nutrition in critically ill adults. *N Engl J Med* 2011;365:506–517.
41. Heidegger CP, Berger MM, Graf S, Zingg W, Darmon P, Costanza MC, Thibault R, Pichard C. Optimisation of energy provision with supplemental parenteral nutrition in critically ill patients: a randomised controlled clinical trial. *Lancet* 2013;381:385–393.
42. Doig GS, Simpson F, Sweetman EA, Finfer SR, Cooper DJ, Heighes PT, Davies AR, O'Leary M, Solano T, Peake S; Early PN Investigators of the ANZICS Clinical Trials Group. Early parenteral nutrition in critically ill patients with short-term relative contraindications to early enteral nutrition: a randomized controlled trial. *JAMA* 2013;309: 2130–2138.
43. Casaer MP, Langouche L, Coudyzer W, Vanbeeckvoort D, De Dobbelaer B, Güiza FG, Wouters PJ, Mesotten D, Van den Berghe G. Impact of early parenteral nutrition on muscle and adipose tissue compartments during critical illness. *Crit Care Med* 2013;41: 2298–2309.
44. Vanderheyden S, Casaer MP, Kesteloot K, Simoons S, De Rijdt T, Peers G, Wouters PJ, Coenegrachts J, Grieten T, Polders K, et al. Early versus late parenteral nutrition in ICU patients: cost analysis of the EPaNIC trial. *Crit Care* 2012;16:R96.
45. Heyland DK, Wischmeyer PE. Does artificial nutrition improve outcome of critical illness? An alternative viewpoint! *Crit Care* 2013;17:324.
46. Buzby GP. Overview of randomized clinical trials of total parenteral nutrition for malnourished surgical patients. *World J Surg* 1993;17: 173–177.
47. Gershengorn HB, Kahn JM, Wunsch H. Temporal trends in the use of parenteral nutrition in critically ill patients. *Chest* 2014;145:508–517.
48. Barr J, Fraser GL, Puntillo K, Ely EW, Gélinas C, Dasta JF, Davidson JE, Devlin JW, Kress JP, Joffe AM, et al.; American College of Critical Care Medicine. Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. *Crit Care Med* 2013;41:263–306.
49. Sessler CN, Pedram S. Protocolized and target-based sedation and analgesia in the ICU. *Crit Care Clin* 2009;25:489–513, viii.
50. Brook AD, Ahrens TS, Schaiff R, Prentice D, Sherman G, Shannon W, Kollef MH. Effect of a nursing-implemented sedation protocol on the duration of mechanical ventilation. *Crit Care Med* 1999;27: 2609–2615.
51. Mascia MF, Koch M, Medicis JJ. Pharmacoeconomic impact of rational use guidelines on the provision of analgesia, sedation, and neuromuscular blockade in critical care. *Crit Care Med* 2000;28: 2300–2306.
52. MacLaren R, Plamondon JM, Ramsay KB, Rocker GM, Patrick WD, Hall RI. A prospective evaluation of empiric versus protocol-based sedation and analgesia. *Pharmacotherapy* 2000;20:662–672.
53. Brattebø G, Hofoss D, Flaatten H, Muri AK, Gjerde S, Plsek PE. Effect of a scoring system and protocol for sedation on duration of patients' need for ventilator support in a surgical intensive care unit. *Qual Saf Health Care* 2004;13:203–205.
54. De Jonghe B, Bastuji-Garin S, Fanguio P, Lacherade JC, Jabot J, Appéré-De-Vecchi C, Rocha N, Outin H. Sedation algorithm in critically ill patients without acute brain injury. *Crit Care Med* 2005;33: 120–127.
55. Elliott R, McKinley S, Aitken LM, Hendrikz J. The effect of an algorithm-based sedation guideline on the duration of mechanical ventilation in an Australian intensive care unit. *Intensive Care Med* 2006;32: 1506–1514.
56. Quenot JP, Ladoire S, Devoucoux F, Doise JM, Cailliod R, Cunin N, Aubé H, Blettery B, Charles PE. Effect of a nurse-implemented sedation protocol on the incidence of ventilator-associated pneumonia. *Crit Care Med* 2007;35:2031–2036.
57. Bucknall TK, Manias E, Presneill JJ. A randomized trial of protocol-directed sedation management for mechanical ventilation in an Australian intensive care unit. *Crit Care Med* 2008;36:1444–1450.
58. Marshall J, Finn CA, Theodore AC. Impact of a clinical pharmacist-enforced intensive care unit sedation protocol on duration of mechanical ventilation and hospital stay. *Crit Care Med* 2008;36: 427–433.
59. Skrobik Y, Ahern S, Leblanc M, Marquis F, Awissi DK, Kavanagh BP. Protocolized intensive care unit management of analgesia, sedation, and delirium improves analgesia and subsyndromal delirium rates. *Anesth Analg* 2010;111:451–463.
60. de Wit M, Gennings C, Jenvey WI, Epstein SK. Randomized trial comparing daily interruption of sedation and nursing-implemented sedation algorithm in medical intensive care unit patients. *Crit Care* 2008;12:R70.
61. Carson SS, Kress JP, Rodgers JE, Vinayak A, Campbell-Bright S, Levitt J, Bourdet S, Ivanova A, Henderson AG, Pohlman A, et al. A randomized trial of intermittent lorazepam versus propofol with daily interruption in mechanically ventilated patients. *Crit Care Med* 2006; 34:1326–1332.
62. Mehta S, Burry L, Cook D, Fergusson D, Steinberg M, Granton J, Herridge M, Ferguson N, Devlin J, Tanios M, et al.; SLEAP Investigators; Canadian Critical Care Trials Group. Daily sedation interruption in mechanically ventilated critically ill patients cared for with a sedation protocol: a randomized controlled trial. *JAMA* 2012; 308:1985–1992.
63. Kollef MH, Levy NT, Ahrens TS, Schaiff R, Prentice D, Sherman G. The use of continuous i.v. sedation is associated with prolongation of mechanical ventilation. *Chest* 1998;114:541–548.
64. Strøm T, Martinussen T, Toft P. A protocol of no sedation for critically ill patients receiving mechanical ventilation: a randomised trial. *Lancet* 2010;375:475–480.
65. Breen D, Karabinis A, Malbrain M, Morais R, Albrecht S, Jarnvig IL, Parkinson P, Kirkham AJ. Decreased duration of mechanical ventilation when comparing analgesia-based sedation using remifentanyl with standard hypnotic-based sedation for up to 10 days in intensive care unit patients: a randomised trial [ISRCTN47583497]. *Crit Care* 2005;9:R200–R210.
66. Chanques G, Jaber S, Barbotte E, Violet S, Sebbane M, Perrigault PF, Mann C, Lefrant JY, Eledjam JJ. Impact of systematic evaluation of pain and agitation in an intensive care unit. *Crit Care Med* 2006;34: 1691–1699.
67. Girard TD, Kress JP, Fuchs BD, Thomason JW, Schweickert WD, Pun BT, Taichman DB, Dunn JG, Pohlman AS, Kinniry PA, et al. Efficacy and safety of a paired sedation and ventilator weaning protocol for mechanically ventilated patients in intensive care (Awakening and Breathing Controlled trial): a randomised controlled trial. *Lancet* 2008;371:126–134.
68. Kress JP, Pohlman AS, O'Connor MF, Hall JB. Daily interruption of sedative infusions in critically ill patients undergoing mechanical ventilation. *N Engl J Med* 2000;342:1471–1477.
69. Fields MJ, Cassel CK. Approaching death: improving care at the end of life. Washington, D.C.: National Academy Press; 1997.
70. Fried TR, Bradley EH, Towle VR, Allore H. Understanding the treatment preferences of seriously ill patients. *N Engl J Med* 2002;346: 1061–1066.
71. Angus DC, Barnato AE, Linde-Zwirble WT, Weissfeld LA, Watson RS, Rickert T, Rubenfeld GD; Robert Wood Johnson Foundation ICU End-Of-Life Peer Group. Use of intensive care at the end of life in the United States: an epidemiologic study. *Crit Care Med* 2004;32: 638–643.
72. Kwok AC, Semel ME, Lipsitz SR, Bader AM, Barnato AE, Gawande AA, Jha AK. The intensity and variation of surgical care at the end of life: a retrospective cohort study. *Lancet* 2011;378:1408–1413.
73. Smith AK, McCarthy E, Weber E, Cenzler IS, Boscardin J, Fisher J, Covinsky K. Half of older Americans seen in emergency department in last month of life; most admitted to hospital, and many die there. *Health Aff (Millwood)* 2012;31:1277–1285.



74. Wright AA, Keating NL, Balboni TA, Matulonis UA, Block SD, Prigerson HG. Place of death: correlations with quality of life of patients with cancer and predictors of bereaved caregivers' mental health. *J Clin Oncol* 2010;28:4457-4464.
75. Wright AA, Zhang B, Ray A, Mack JW, Trice E, Balboni T, Mitchell SL, Jackson VA, Block SD, Maciejewski PK, et al. Associations between end-of-life discussions, patient mental health, medical care near death, and caregiver bereavement adjustment. *JAMA* 2008;300:1665-1673.
76. Temel JS, Greer JA, Muzikansky A, Gallagher ER, Admane S, Jackson VA, Dahlin CM, Blinderman CD, Jacobsen J, Pirl WF, et al. Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 2010;363:733-742.
77. Bakitas M, Lyons KD, Hegel MT, Balan S, Brokaw FC, Seville J, Hull JG, Li Z, Tosteson TD, Byock IR, et al. Effects of a palliative care intervention on clinical outcomes in patients with advanced cancer: the Project ENABLE II randomized controlled trial. *JAMA* 2009;302:741-749.
78. Connor SR, Pyenson B, Fitch K, Spence C, Iwasaki K. Comparing hospice and nonhospice patient survival among patients who die within a three-year window. *J Pain Symptom Manage* 2007;33:238-246.
79. Azoulay E, Pochard F, Kentish-Barnes N, Chevret S, Aboab J, Adrie C, Annane D, Bleichner G, Bollaert PE, Darmon M, et al.; FAMIREA Study Group. Risk of post-traumatic stress symptoms in family members of intensive care unit patients. *Am J Respir Crit Care Med* 2005;171:987-994.
80. Siegel MD, Hayes E, Vanderwerker LC, Loseth DB, Prigerson HG. Psychiatric illness in the next of kin of patients who die in the intensive care unit. *Crit Care Med* 2008;36:1722-1728.
81. Wendler D, Rid A. Systematic review: the effect on surrogates of making treatment decisions for others. *Ann Intern Med* 2011;154:336-346.
82. Terry PB, Vettese M, Song J, Forman J, Haller KB, Miller DJ, Stallings R, Sulmasy DP. End-of-life decision making: when patients and surrogates disagree. *J Clin Ethics* 1999;10:286-293.
83. Singer PA, Martin DK, Lavery JV, Thiel EC, Kelner M, Mendelssohn DC. Reconceptualizing advance care planning from the patient's perspective. *Arch Intern Med* 1998;158:879-884.
84. Steinhauser KE, Christakis NA, Clipp EC, McNeilly M, McIntyre L, Tulskey JA. Factors considered important at the end of life by patients, family, physicians, and other care providers. *JAMA* 2000;284:2476-2482.
85. Curtis JR, Engelberg RA, Wenrich MD, Shannon SE, Treece PD, Rubenfeld GD. Missed opportunities during family conferences about end-of-life care in the intensive care unit. *Am J Respir Crit Care Med* 2005;171:844-849.
86. Brush DR, Rasinski KA, Hall JB, Alexander GC. Recommendations to limit life support: a national survey of critical care physicians. *Am J Respir Crit Care Med* 2012;186:633-639.
87. Schenker Y, Tiver GA, Hong SY, White DB. Association between physicians' beliefs and the option of comfort care for critically ill patients. *Intensive Care Med* 2012;38:1607-1615.
88. Barnato AE, Herndon MB, Anthony DL, Gallagher PM, Skinner JS, Bynum JP, Fisher ES. Are regional variations in end-of-life care intensity explained by patient preferences?: a study of the US Medicare population. *Med Care* 2007;45:386-393.
89. Anthony DL, Herndon MB, Gallagher PM, Barnato AE, Bynum JP, Gottlieb DJ, Fisher ES, Skinner JS. How much do patients' preferences contribute to resource use? *Health Aff (Millwood)* 2009;28:864-873.
90. Scheunemann LP, Arnold RM, White DB. The facilitated values history: helping surrogates make authentic decisions for incapacitated patients with advanced illness. *Am J Respir Crit Care Med* 2012;186:480-486.
91. Quill TE. Perspectives on care at the close of life. Initiating end-of-life discussions with seriously ill patients: addressing the "elephant in the room". *JAMA* 2000;284:2502-2507.
92. Gries CJ, Engelberg RA, Kross EK, Zatzick D, Nielsen EL, Downey L, Curtis JR. Predictors of symptoms of posttraumatic stress and depression in family members after patient death in the ICU. *Chest* 2010;137:280-287.
93. Kross EK, Engelberg RA, Gries CJ, Nielsen EL, Zatzick D, Curtis JR. ICU care associated with symptoms of depression and posttraumatic stress disorder among family members of patients who die in the ICU. *Chest* 2011;139:795-801.
94. Lautrette A, Darmon M, Megarbane B, Joly LM, Chevret S, Adrie C, Barnoud D, Bleichner G, Bruel C, Choukroun G, et al. A communication strategy and brochure for relatives of patients dying in the ICU. *N Engl J Med* 2007;356:469-478.
95. Luce JM, Rubenfeld GD. Can health care costs be reduced by limiting intensive care at the end of life? *Am J Respir Crit Care Med* 2002;165:750-754.
96. Curtis JR, Engelberg RA, Bensink ME, Ramsey SD. End-of-life care in the intensive care unit: can we simultaneously increase quality and reduce costs? *Am J Respir Crit Care Med* 2012;186:587-592.
97. Roberts RR, Frutos PW, Ciavarella GG, Gussow LM, Mensah EK, Kampe LM, Straus HE, Joseph G, Rydman RJ. Distribution of variable vs fixed costs of hospital care. *JAMA* 1999;281:644-649.
98. Wunsch H, Angus DC, Harrison DA, Collange O, Fowler R, Hoste EA, de Keizer NF, Kersten A, Linde-Zwirble WT, Sandiumenge A, et al. Variation in critical care services across North America and Western Europe. *Crit Care Med* 2008;36:2787-2793, e1-e9.
99. Chen LM, Render M, Sales A, Kennedy EH, Wiitala W, Hofer TP. Intensive care unit admitting patterns in the Veterans Affairs health care system. *Arch Intern Med* 2012;172:1220-1226.
100. Seymour CW, Kahn JM. Addressing the growth in intensive care: comment on "Intensive care unit admitting patterns in the Veterans Affairs health care system". *Arch Intern Med* 2012;172:1226-1228.
101. Wagner J, Gabler NB, Ratcliffe SJ, Brown SES, Strom BL, Halpern SD. Outcomes among patients discharged from busy intensive care units. *Ann Intern Med* 2013;159:447-455.
102. Wunsch H, Wagner J, Herlim M, Chong DH, Kramer AA, Halpern SD. ICU occupancy and mechanical ventilator use in the United States. *Crit Care Med* 2013;41:2712-2719.
103. Mahesh M, Durand DJ. The choosing wisely campaign and its potential impact on diagnostic radiation burden. *J Am Coll Radiol* 2013;10:65-66.
104. Volpp KG, Loewenstein G, Asch DA. Choosing wisely: low-value services, utilization, and patient cost sharing. *JAMA* 2012;308:1635-1636.
105. Chien AT, Rosenthal MB. Waste not, want not: promoting efficient use of health care resources. *Ann Intern Med* 2013;158:67-68.