

Published in final edited form as:

Nat Chem Biol. 2007 April ; 3(4): 218–221. doi:10.1038/nchembio864.

## An RNA G-quadruplex in the 5' UTR of the *NRAS* proto-oncogene modulates translation

Sunita Kumari<sup>1</sup>, Anthony Bugaut<sup>1</sup>, Julian L Huppert<sup>2</sup>, and Shankar Balasubramanian<sup>1</sup>

<sup>1</sup>University Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW, UK.

<sup>2</sup>Wellcome Trust Sanger Institute, Hinxton, Cambridge CB10 1SA, UK.

### Abstract

Guanine-rich nucleic acid sequences can adopt noncanonical four-stranded secondary structures called guanine (G)-quadruplexes<sup>1</sup>. Bioinformatics analysis suggests that G-quadruplex motifs are prevalent in genomes<sup>2</sup>, which raises the need to elucidate their function. There is now evidence for the existence of DNA G-quadruplexes at telomeres with associated biological function<sup>3</sup>. A recent hypothesis supports the notion that gene promoter elements contain DNA G-quadruplex motifs that control gene expression at the transcriptional level<sup>4</sup>. We discovered a highly conserved, thermodynamically stable RNA G-quadruplex in the 5' untranslated region (UTR) of the gene transcript of the human *NRAS* proto-oncogene. Using a cell-free translation system coupled to a reporter gene assay, we have demonstrated that this *NRAS* RNA G-quadruplex modulates translation. This is the first example of translational repression by an RNA G-quadruplex. Bioinformatics analysis has revealed 2,922 other 5' UTR RNA G-quadruplex elements in the human genome. We propose that RNA G-quadruplexes in the 5' UTR modulate gene expression at the translational level.

The existence of RNA G-quadruplexes *in vivo* is more inevitable than the existence of DNA G-quadruplexes, given that (i) the former are generally more thermodynamically stable in the folded form than their DNA counterparts<sup>5</sup>, and (ii) RNA is single-stranded, which implies that quadruplex formation does not have to compete with hybridization to a complementary strand. In this study we have focused on the 5' UTRs of mRNA, which are known to be involved in translational regulation, particularly for growth factors, transcription factors and oncoproteins<sup>6</sup>. The neuroblastoma RAS viral oncogene homolog (*NRAS*)-encoded protein p21 mediates both signal transduction across the plasma membrane and the intracellular signaling pathways responsible for cell proliferation and differentiation<sup>7</sup>. Activating mutations in the coding region of *NRAS* are responsible for increased cell proliferation<sup>8</sup>. The suppression of oncogenic *NRAS* by small interfering RNA causes apoptosis of tumor cells<sup>9</sup>, which suggests that inhibiting the expression of oncogenic *NRAS* is a potential therapeutic strategy. Using a computational search algorithm we developed for locating quadruplex sequence motifs<sup>2</sup>, we identified a putative G-quadruplex

© 2007 Nature Publishing Group

Correspondence should be addressed to S.B. (sb10031@cam.ac.uk).

#### AUTHOR CONTRIBUTIONS

S.K., A.B. and S.B. conceived and designed the experiments. S.K. performed all the experiments. S.K. and A.B. analyzed the experimental data. J.L.H. conceived, designed and carried out the bioinformatics studies. All four authors discussed the results, and the manuscript was written by S.K. and S.B. with contributions from A.B. and J.L.H.

#### COMPETING INTERESTS STATEMENT

The authors declare no competing financial interests.

**Accession codes.** EMBL Nucleotide Sequence Database: DNA sequences for *UTRQ*, *DelQ* and *MutQ* have been deposited under accession numbers AM489486, AM489487 and AM489488, respectively.

forming sequence in the 5' UTR of the human *NRAS* proto-oncogene mRNA. This 254-nucleotide-long *NRAS* 5' UTR10 contains the *NRAS* RNA G-quadruplex (NRQ) motif (5'-GGGAGGGGCGGGUCUGGG-3'), which is located 14 nucleotides downstream of the 5' cap and 222 nucleotides upstream of the translation start site (Supplementary Fig. 1 online). This motif is highly conserved, in both its sequence and its position relative to the translation start site, across the 5' UTRs of human, chimpanzee, macaque, mouse, rat and dog genes orthologous to *NRAS* (Table 1 and Supplementary Table 1 online).

To confirm that the putative RNA G-quadruplex NRQ folds into a stable quadruplex, we carried out biophysical experiments on the synthetic oligonucleotide 5'-UGUGGGAGGGGCGGGUCUGGG-3'. Circular dichroism (CD) spectroscopy has been widely used to characterize the structure of folded nucleic acid quadruplexes<sup>11</sup>. At pH 7.4, 100 mM KCl, the CD spectrum of NRQ showed a positive peak at 263 nm and a negative peak at 241 nm (Fig. 1a), which is the characteristic CD signature of a parallel quadruplex structure<sup>11,12</sup>. The thermal melting of quadruplex nucleic acids can be characterized by an inverse UV transition at 295 nm<sup>13</sup>, and has generally been found to have a significant and characteristic cation dependence<sup>14</sup>. At pH 7.4 and low K<sup>+</sup> (1 mM), the UV-melting profile of NRQ at 295 nm showed a hypochromic shift (Fig. 1b), with a  $T_m$  value of  $63 \pm 1$  °C (error here and below is s.e.m.); this result was corroborated by CD melting at 260 nm under comparable conditions, which also gave a  $T_m$  of  $63 \pm 1$  °C (Supplementary Fig. 2 online). At 100 mM KCl, the folded NRQ G-quadruplex could not be unfolded, even at 95 °C, which is indicative of a very stable quadruplex. The NRQ RNA G-quadruplex was reasonably stable ( $T_m = 43 \pm 1$  °C) even in the absence of added stabilizing cations. Studies over a ten-fold strand concentration range (from 1 to 10  $\mu$ M) showed no change in the  $T_m$  value (Supplementary Table 2 online), which is consistent with the melting of an intramolecular quadruplex. The monovalent ion dependence for stabilization of folded NRQ, as judged by  $T_m$ , was in the order K<sup>+</sup> > Na<sup>+</sup> > Li<sup>+</sup> (Supplementary Table 3 online), which is characteristic of G-quadruplex nucleic acids<sup>14</sup>. Thus, NRQ folds into a very stable, parallel intramolecular G-quadruplex under near-physiological pH and salt conditions.

To specifically evaluate the influence of the NRQ RNA G-quadruplex on the efficiency of translation, we cloned the 254-base-pair 5' UTR of *NRAS* from human genomic DNA upstream of the firefly luciferase reporter gene and immediately downstream of the minimal T7 promoter. Whereas this plasmid (pSKC11) encodes the transcript *UTRQ* (Fig. 2a), which contains the native 5' UTR of *NRAS* including the intact NRQ G-quadruplex motif, we also generated two controls. We obtained the plasmid pSKC12, which encodes the transcript *DelQ* (Fig. 2a), by deleting the first 29 base pairs of the *NRAS* 5' UTR (that comprises the RNA G-quadruplex-forming sequence). The second control plasmid, pSKC13, which encodes the transcript *MutQ* (Fig. 2a), was derived from pSKC11 by a GGG-to-AAA substitution that was made at base pairs 15-17 of the *NRAS* 5' UTR to disrupt RNA G-quadruplex formation but maintain the natural length of the 5' UTR. For each system the corresponding 5'-capped RNA transcript was generated by *in vitro* transcription using T7 RNA polymerase, and the transcripts were subjected to *in vitro* translation using the rabbit reticulocyte lysate. We evaluated the efficiency of translation by the standard luminescence assay for luciferase catalytic activity<sup>15</sup>. Figure 2b describes the relative translation efficiency for each system. Deletion of the G-quadruplex motif (*DelQ*) resulted in a 3.7-fold increase in translation efficiency relative to the native sequence system, *UTRQ*. This indicates that the RNA G-quadruplex NRQ suppresses translation. The GGG-to-AAA quadruplex-destabilizing triple mutation (*MutQ*) caused a 3.6-fold increase in translation efficiency relative to *UTRQ*. Taken together, these data indicate that the natural 5' UTR NRQ RNA G-quadruplex has an inhibitory effect on translation. Translational control of proto-oncogenes mediated via the 5' UTR is generally complex<sup>6</sup>, and we postulate that this

RNA G-quadruplex is one regulatory element within the 5' UTR of *NRAS*. Factors that disrupt NRQ RNA G-quadruplex formation, such as mutations, may lead to oncogenesis.

On the basis of this study, we reasoned that analogous RNA G-quadruplexes may exist within the 5' UTRs of other genes. To investigate this, we carried out a bioinformatics search for RNA G-quadruplexes in 5' UTRs in the human genome. We examined 38,915 5' UTRs from all known *Homo sapiens* genes in Ensembl (version 40, National Center for Biotechnology Information (NCBI) build 36). 2,922 of the 5' UTRs contained one or more G-quadruplex motifs, giving a total of 3,321 5' UTR G-quadruplex motifs. The density of G-quadruplex motifs in 5' UTRs as a whole is 0.305 per kilobase, compared with a genome average of 0.063 per kilobase (for one strand only)<sup>2</sup>, which is a 4.8-fold enrichment of G-quadruplex motifs. It is noteworthy that the genes containing 5' UTR G-quadruplex motifs include several other proto-oncogenes, including *BCL2*, *JUN* and *FGR* (Table 2).

Stable RNA hairpin secondary structures in the 5' UTR of mRNAs inhibit the process of translation<sup>16</sup>. The intramolecular RNA G-quadruplex motif (NRQ) that we have identified in the 5' UTR of *NRAS* is the first example to be elucidated of a quadruplex that inhibits translation. Detailed elucidation of the quadruplex structure and also of the quadruplex-mediated inhibitory mechanism will be the subject of future work. The presence of G-quadruplex motifs in the 5' UTRs of 2,922 human genes suggests that the functional role of the *NRAS* RNA G-quadruplex may be a paradigm for other genes. Substituted flat polyaromatic heterocyclic molecular scaffolds that complement the surface of a G-tetrad have provided many small quadruplex-binding ligands<sup>17</sup>. Given the emerging interest in small molecules that target and stabilize nucleic acid quadruplexes with specificity<sup>18</sup>, this RNA G-quadruplex has obvious potential as a molecular target for small-molecule therapeutic agents that act by inhibiting mRNA translation.

## METHODS

### Bioinformatics

Genomic data was taken from Ensembl using version 40, corresponding to NCBI build 36, and was acquired using the EnsMart tool. It was then analyzed using quadparser2 and custom-written perl utilities.

### Spectroscopy

For spectroscopic studies, we prepared RNA samples at 4  $\mu$ M strand concentration in RNase-free water and degassed buffers containing 10 mM Tris-HCl, pH 7.4, and varying KCl concentrations from 1 mM to 100 mM. The samples were annealed by heating at 90 °C for 10 min and then slow cooling to 5 °C at a controlled rate of 0.2 °C min<sup>-1</sup>.

CD experiments were performed using a Jasco J-810 spectropolarimeter equipped with a Peltier temperature controller. Typically, a 200  $\mu$ l sample was placed in a quartz cuvette with an optical path length of 1 mm, transferred to the spectropolarimeter and allowed to equilibrate at 20 °C for 10 min. Five CD scans, over the wavelength range of 220 to 320 nm, were performed at 50 nm min<sup>-1</sup> with a 2-s response time, 1-nm pitch and 1-nm bandwidth, and their average was taken. For each experiment, a CD spectrum of the buffer was recorded and subtracted from the spectrum obtained for the RNA-containing solution. Data were zero-corrected at 320 nm.

UV-melting studies were carried out on a Varian Cary 1E UV-visible spectrophotometer equipped with a Peltier temperature controller. Typically, a 100  $\mu$ l sample was transferred to a quartz cuvette with an optical path length of 1 cm and was covered with approximately 100  $\mu$ l of mineral oil to prevent sample evaporation. This sample was transferred to the

spectrophotometer and then heated to 90 °C and cooled to 5 °C, twice consecutively at 0.25 °C min<sup>-1</sup> temperature gradient, and absorption data recorded at 295 nm were collected every 0.5 min on both annealing and melting steps.  $T_m$  values were determined using the van't Hoff method<sup>19</sup>.

### Plasmid construction

To construct the plasmid pSKC11, which encodes the 254-nucleotide *UTRQ* transcript, we PCR-amplified the 237 base pairs of exon-I of the *NRAS* 5' UTR from human genomic DNA using *Pfu* DNA polymerase (Promega) and two sequence-specific primers: a forward primer tailed with *Hind*III (underlined) and a minimal T7 promoter (5'-AAGCTTTAATACGACTCACTATAGGGGACGTCCCCTGTGGGAGGGGCG-3'); bold GG indicates positions that were mutated from AA in the natural 5' UTR sequence to yield an efficient transcription template), and a restriction site *Sma*I-tailed (naturally present at the end of exon -I) reverse primer (5'-CTCAAGCTCAAGCTCCACTGCCTCTGC-3'). This amplified product (A) was digested with *Hind*III and *Sma*I, and purified by gel electrophoresis. We amplified the firefly luciferase gene from the pGL3 basic vector (Promega) using a forward primer tailed with *Sma*I (underlined) and the 17 base pairs of the 5' UTR from exon I (5'-CTTGAGGTTCTTGCTGGTGTGAAATGGAAGACGCCAAAAACATAAAG-3'), and an *Eco*RI-tailed reverse primer (5'-GAATTCTTACACGGCGATCTTTCC-3'). This amplified product (B) was digested with *Sma*I and *Eco*RI and purified by gel electrophoresis. In a three-piece ligation, we ligated the two gel-purified fragments (A and B) at the *Sma*I site and inserted into the *Hind*III- and *Eco*RI-digested linear pUC18 vector (Invitrogen) to obtain the full-length (254 base pairs) *NRAS* 5' UTR containing the G-quadruplex-forming sequence at its natural position (14 base pairs downstream of the 5' cap). We confirmed positive clones for the full length of insert by DNA sequencing.

To construct pSKC12, the control plasmid encoding the transcript *DelQ*, we deleted the first 29 base pairs of the 5' UTR, including the G-quadruplex-forming sequence, from the pSKC11 plasmid using the forward primer (5'-GGGTGCGGCCTGCCGCATGACTCG-3') and an *Eco*RI-tailed reverse primer (5'-GAATTCTTACACGGCGATCTTTCC-3'). We gel-purified this amplified product and reamplified with a forward primer tailed with *Hind*III and a minimal T7 promoter (5'-AAGCTTTAATACGACTCACTATAGGGTGCGGCCTGCCGCATGACTCG-3') and the above-mentioned *Eco*RI-tailed reverse primer. This reamplified product was inserted into the *Hind*III- and *Eco*RI-digested linear pUC18 vector. The second control plasmid, pSKC13, which encodes the *MutQ* transcript, was derived from the pSKC11 plasmid by a GGG-to-AAA substitution at base pairs 15-17 of the *NRAS* 5' UTR using the QuikChange site-directed mutagenesis kit (Stratagene) according to the manufacturer's protocol. We sequenced all the constructs to confirm the presence of the intended changes.

### *In vitro* transcription

The plasmids pSKC11, pSKC12 and pSKC13, which encode the transcripts *UTRQ*, *DelQ* and *MutQ*, respectively, were linearized at the 3' end of the region to be transcribed using *Eco*RI. 5'-capped transcripts were synthesized *in vitro* using the mMessage mMachine T7 kit (Ambion). We purified all transcripts on 1% agarose gel, after DNase treatment. The RNA concentration was determined by UV spectroscopy. We confirmed the integrity and size of each transcript on 1% agarose gel.

### *In vitro* translation and luciferase assay

We carried out *in vitro* translation of the mRNAs in a cell-free translation system consisting of extracts from nuclease-treated rabbit reticulocytes lysate (Promega) according to the

manufacturer's protocol. We measured firefly luciferase activity using luciferase assay reagent (Promega) according to the manufacturer's protocol on a Varian Cary Eclipse fluorescence spectrophotometer working in bio/chemiluminescence mode.

## Acknowledgments

We thank Cancer Research UK, the Cambridge Commonwealth Trust and Trinity College, Cambridge for funding. S.B. is a Biotechnology and Biological Sciences Research Council career development fellow. J.L.H. is a research fellow at Trinity College, Cambridge. We thank Z. Jawad-Alami for useful discussions and S. Sewitz for critically reading this manuscript.


## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## References


1. Neidle, S.; Balasubramanian, S. *Quadruplex Nucleic Acids*. Cambridge: RSC Biomolecular Sciences; 2006.
2. Huppert JL, Balasubramanian S. Prevalence of quadruplexes in the human genome. *Nucleic Acids Res.* 2005; 33:2908–2916. [PubMed: 15914667]
3. Paeschke K, Simonsson T, Postberg J, Rhodes D, Lipps HJ. Telomere end-binding proteins control the formation of G-quadruplex DNA structures *in vivo*. *Nat. Struct. Mol. Biol.* 2005; 12:847–854. [PubMed: 16142245]
4. Siddiqui-Jain A, Grand CL, Bearss DJ, Hurley LH. Direct evidence for a G-quadruplex in a promoter region and its targeting with a small molecule to repress c-*MYC* transcription. *Proc. Natl. Acad. Sci. USA.* 2002; 99:11593–11598. [PubMed: 12195017]
5. Sacca B, Lacroix L, Mergny J-L. The effect of chemical modifications on the thermal stability of different G-quadruplex-forming oligonucleotides. *Nucleic Acids Res.* 2005; 33:1182–1192. [PubMed: 15731338]
6. Willis AE. Translational control of growth factor and proto-oncogene expression. *Int. J. Biochem. Cell Biol.* 1999; 31:73–86. [PubMed: 10216945]
7. Downward J. Targeting Ras signalling pathways in cancer therapy. *Nat. Rev. Cancer.* 2003; 3:11–22. [PubMed: 12509763]
8. Barbacid M. *Ras* genes. *Annu. Rev. Biochem.* 1987; 56:779–827. [PubMed: 3304147]
9. Eskandarpour M, et al. Suppression of oncogenic *NRAS* by RNA interference induces apoptosis of human melanoma cells. *Int. J. Cancer.* 2005; 115:65–73. [PubMed: 15688405]
10. Hall A, Brown R. Human *N-ras*: cDNA cloning and gene structure. *Nucleic Acids Res.* 1985; 13:5255–5268. [PubMed: 2991860]
11. Balagurumorthy P, Brahmachari SK, Mohanty D, Bansal M, Sasisekharan V. Hairpin and parallel quartet structure for telomeric sequences. *Nucleic Acids Res.* 1992; 20:4061–4067. [PubMed: 1508691]
12. Tang C-F, Shafer RH. Engineering the quadruplex fold: nucleoside conformation determines both folding topology and molecularity in guanine quadruplexes. *J. Am. Chem. Soc.* 2006; 128:5966–5973. [PubMed: 16637665]
13. Mergny J-L, Phan A-T, Lacroix L. Following G-quartet formation by UV-spectroscopy. *FEBS Lett.* 1998; 435:74–78. [PubMed: 9755862]
14. Hardin CC, Watson T, Corregan M, Bailey C. Cation-dependent transition between the quadruplex and Watson-Crick hairpin forms of d(CGCG3GCG). *Biochemistry.* 1992; 31:833–841. [PubMed: 1731941]
15. DiLella AG, et al. Utility of firefly luciferase as a reporter gene for promoter activity in transgenic mice. *Nucleic Acids Res.* 1988; 16:4159. [PubMed: 3375079]
16. Kozak M. Structural features in eukaryotic mRNAs that modulate the initiation of translation. *J. Biol. Chem.* 1991; 266:19867–19870. [PubMed: 1939050]

17. Searle, MS.; Balkwill, GD. Quadruplex Nucleic Acids. Neidle, S.; Balasubramanian, S., editors. Cambridge: RSC Biomolecular Sciences; 2006. p. 131-153.
18. Neidle S, Parkinson G. Telomere maintenance as a target for anticancer drug discovery. *Nat. Rev. Drug Discov.* 2002; 1:383–393. [PubMed: 12120414]
19. Mergny JL, Lacroix L. Analysis of thermal melting curves. *Oligonucleotides.* 2003; 13:515–537. [PubMed: 15025917]



**Figure 1.** Biophysical analysis of the NRQ RNA G-quadruplex. **(a)** CD spectrum of NRQ at 4  $\mu$ M strand concentration in 10 mM Tris-HCl, pH 7.4, and 100 mM KCl, at 20 °C. **(b)** Melting and annealing UV profiles of NRQ at 4  $\mu$ M strand concentration in 10 mM Tris-HCl, pH 7.4, and 1 mM KCl.





**Figure 2.** Effect of the *NRAS* 5' UTR on the translational efficiency of chimeric RNA. **(a)** Schematic representation of mRNAs of luciferase reporter constructs: *UTRQ*, full-length (254 nucleotides) *NRAS* 5' UTR; *DelQ*, in which the first 29 nucleotides of *UTRQ*, containing the G-quadruplex sequence, are deleted; and *MutQ*, full-length *NRAS* 5' UTR with a GGG-to-AAA mutation (shown in blue) at nucleotides 15-17 to disrupt the RNA G-quadruplex formation. **(b)** Relative translation efficiency of the three constructs, as judged by quantitation of luciferase enzyme activity. Results were normalized relative to data for the *UTRQ* system. Error bars represent the s.d. of three independent experiments.



**Table 1**Conservation of the *NRAS* 5' UTR G-quadruplex

Species	Sequence <sup>a</sup>	Position <sup>b</sup>
Human	<b>GGGAGGGGCGGG</b> --U---C <b>UGGG</b>	-222
Chimpanzee	<b>GGGAGGGGCGGG</b> --U---C <b>UGGG</b>	-222
Macaque	<b>GGGAGGGGCGGG</b> --U---C <b>UGGG</b>	-222
Mouse <sup>c</sup>	<b>GGGGGCGGGGCGGGG</b> CUGGAC <b>UGGG</b>	-220
Rat	<b>GGGUGGGGAGGGGCGGGG</b> -UG---- <b>GGG</b>	-221
Dog <sup>c</sup>	<b>GGGAGGGGCGGG</b> -----AA <b>UGGG</b>	-730 <sup>d</sup>
Consensus	<b>GGGAGGGGCGGG</b> --U---C <b>UGGG</b>	

<sup>a</sup>Dashes represent gaps in the alignment. Nucleotides in bold are runs of guanines capable of forming G-quartets.

<sup>b</sup>The position of the last G of the putative G-quadruplex sequence is given relative to the translation start site.

<sup>c</sup>Mouse and dog each have a second G-quadruplex sequence motif in their 5' UTRs (Supplementary Table 1).

<sup>d</sup>The difference in position for dog is likely to be due to the incomplete annotation of introns, as all other *NRAS* ortholog sequences have an intron in the unspliced 5' UTR (480 nucleotides in humans), whose removal reduces the distance to approximately 220 nucleotides in the spliced form.

Table 2

## Human proto-oncogenes containing 5' UTR G-quadruplex motifs

Gene description	HGNC code	5' UTR length	Motif position <sup>d</sup>	Motif sequence	Ensembl gene ID
GTPase NRAS	<i>NRAS</i>	254	-222	GGGAGGGGGGGGUCUGGG	ENSG00000168638 <sup>b</sup>
Proto-oncogene tyrosine-protein kinase	<i>FGR</i>	186	-156	GGGUGAGAGGGGCAAGGGGGCUAGGG	ENSG00000000938
Thyroid hormone receptor- $\alpha$	<i>THRA</i>	483	-85	GGGCCUGGGUGGCAAGGGGUGGG	ENSG00000126351
Transcription factor Maf	<i>MAF</i>	812	-170	GGGGGAGGGAGGGCGGGCCGGG GGGCCGGGGCAGGGCGGGGGG	ENSG00000178573
Friend leukemia integration 1 transcription factor	<i>FLI1</i>	165	-95	GGGAGGGCCCCAGGGCCCGGG	ENSG00000151702
Proto-oncogene C-crk	<i>CRK</i>	141	-44	GGGCCGGGGCCCGGGGGCCGGAGGGG	ENSG00000167193 <sup>c</sup>
Apoptosis regulator	<i>BCL2</i>	493	-42	GGGGCCCGUGGGGUGGGAGCUGGGG	ENSG00000171791
Mitogen-activated protein kinase kinase 8	<i>MAP3K8</i>	696	-647	GGGCAAUAGGGGGCGGGGUGGGCGGG UGGGGGGG	ENSG00000107968 <sup>d</sup>
Proto-oncogene tyrosine-protein kinase	<i>ABL1</i>	381	-270	GGGAACGCCAGGGGCCCCUGGGUCGGACGGG	ENSG00000097007
Proto-oncogene tyrosine-protein kinase	<i>ABL1</i>	381	-212	GGGCCGGGGCGGGCCUGGGCGGG	ENSG00000097007
Proto-oncogene tyrosine-protein kinase	<i>ABL1</i>	381	-126	GGGCCGGGGCGGGCGGGCGGGGG	ENSG00000097007
Proto-oncogene tyrosine-protein kinase	<i>ABL1</i>	381	-66	GGGGCCGGGGCGGGCGGGGGCGGGGGCC GAGCCGGG	ENSG00000097007
Transcription factor API	<i>JUN</i>	1,257	-1,231	GGGGAGGGGACCGGGGAAGAGAGGG	ENSG00000177606

<sup>a</sup>The position of the last G of the G-quadruplex motif is given relative to the translation start site.

<sup>b</sup>Because of an annotation error in version 40 of Ensembl, whereby the *NRAS* gene was accidentally fused with the neighboring *CSDE1* gene, we have used data from the original publication 10, which is identical to that in previous versions of Ensembl and the manually curated vertebrate genome annotation (Vega) database.

<sup>c</sup>There is also an alternative transcript (ENST00000300574) that has eight fewer bases in the 5' UTR, upstream of the G-quadruplex motif.

<sup>d</sup>There is also an alternative transcript (ENST00000375328) that has seven fewer bases in the 5' UTR, downstream of the G-quadruplex motif. HGNC, Hugo Gene Nomenclature Committee.