

An Update on Adenosine A_{2A}-Dopamine D₂ Receptor Interactions: Implications for the Function of G Protein-Coupled Receptors

S. Ferré^{1,*}, C. Quiroz¹, A.S. Woods¹, R. Cunha², P. Popoli³, F. Ciruela⁴, C. Lluís⁵, R. Franco⁵, K. Azdad⁶ and S. N. Schiffmann⁶

¹National Institute on Drug Abuse, I.R.P., N.I.H., D.H.H.S., Baltimore, MD 21224, USA; ²Center for Neurosciences of Coimbra, Faculty of Medicine, Institute of Biochemistry, University of Coimbra, 3004-504 Coimbra, Portugal; ³Istituto Superiore di Sanita, 00161 Rome, Italy; ⁴Unitat de Farmacologia, Departament de Patologia i Terapèutica Experimental, Facultat de Medicina, Universitat de Barcelona, 08907 Barcelona, Spain; ⁵Institut d'Investigació Biomedica August Pi I Sunyer, CIBERNED, University of Barcelona, 08028 Barcelona, Spain and ⁶Universite Libre de Bruxelles, 1070 Brussels, Belgium

Abstract: Adenosine A_{2A}-dopamine D₂ receptor interactions play a very important role in striatal function. A_{2A}-D₂ receptor interactions provide an example of the capabilities of information processing by just two different G protein-coupled receptors. Thus, there is evidence for the coexistence of two reciprocal antagonistic interactions between A_{2A} and D₂ receptors in the same neurons, the GABAergic enkephalinergic neurons. An antagonistic A_{2A}-D₂ intramembrane receptor interaction, which depends on A_{2A}-D₂ receptor heteromerization and G_{q/11}-PLC signaling, modulates neuronal excitability and neurotransmitter release. On the other hand, an antagonistic A_{2A}-D₂ receptor interaction at the adenylyl-cyclase level, which depends on G_{s/olf}- and G_{i/o}- type V adenylyl-cyclase signaling, modulates protein phosphorylation and gene expression. Finally, under conditions of upregulation of an activator of G protein signaling (AGS3), such as during chronic treatment with addictive drugs, a synergistic A_{2A}-D₂ receptor interaction can also be demonstrated. AGS3 facilitates a synergistic interaction between G_{s/olf}- and G_{i/o}-coupled receptors on the activation of types II/IV adenylyl cyclase, leading to a paradoxical increase in protein phosphorylation and gene expression upon co-activation of A_{2A} and D₂ receptors. The analysis of A_{2A}-D₂ receptor interactions will have implications for the pathophysiology and treatment of basal ganglia disorders and drug addiction.

Key Words: Adenosine A_{2A} Receptor, Dopamine D₂ Receptor, G Protein-Coupled Receptors, Receptor Heteromers, Striatum, Basal Ganglia Disorders, Drug Addiction.

LOCALIZATION OF THE A_{2A}-D₂ RECEPTOR HETEROMER

Applying a broad definition of “neurotransmitter” [1], adenosine can be considered as an important neurotransmitter in the CNS, which acts through different subtypes of G protein-coupled receptors (GPCRs). From the four cloned adenosine receptors (adenosine A₁, A_{2A}, A_{2B} and A₃ receptors), A₁ and A_{2A} receptors are the main targets for the physiological effects of adenosine in the brain [2]. A₁ receptor is widely distributed in the brain, including the striatum, while A_{2A} receptor is mostly concentrated in the striatum [2,3]. It is becoming increasingly obvious that the modulatory role of adenosine in the striatum is related to the ability of A₁ and A_{2A} receptors to heteromerize with themselves and with other GPCRs, such as dopamine, glutamate, cannabinoid and ATP receptors [4-14]. The present review focuses on the role of one particular adenosine receptor heteromer, the one constituted by the A_{2A} and the dopamine D₂ receptor, which is already having important implications for the treatment of neuropathologies involving the striatum (see below).

Striatal medium spiny neurons are GABAergic efferent neurons which constitute more than 95% of the striatal neuronal population. They receive two main afferents, cortical-limbic-thalamic glutamatergic inputs and dopaminergic mesencephalic inputs, from the substantia nigra pars compacta and the VTA. These inputs converge in the dendritic spine, with the glutamatergic input making synaptic contact with the head of the dendritic spine and the dopaminergic input making synaptic contact with the neck of the dendritic spine [15,16]. The dendritic spine, the glutamatergic terminal, the dopaminergic terminal and astroglial processes that wrap the glutamatergic synapse constitute the most common local module in the striatum, which we have recently called striatal spine module [16]. In the striatal spine module adenosine plays a very important role in

the modulation of both glutamatergic and dopaminergic neurotransmission [14,16,17].

It was initially thought that most extracellular adenosine came from intracellular adenosine as a product of ATP, due to an increased metabolic demand of the cell [17]. However, recent studies suggest that astroglia plays a very important role in the production of extracellular adenosine. Astrocytes express glutamate and ATP receptors, which when activated induce astrocytes to release glutamate and ATP, which can then be converted to adenosine by means of ectonucleotidases [18-20]. This adds more relevance to the already known key role of astrocytes in the computation of information in the striatal spine module [16,18,19]. Finally, increasing evidence suggests that ATP is co-released with glutamate by the glutamatergic terminals and converted to adenosine by ectonucleotidases [14,16].

There are two subtypes of GABAergic striatal efferent neurons, the GABAergic striopallidal neuron, which can be called GABAergic enkephalinergic neuron, since it expresses the peptide enkephalin, and the GABAergic striatonigral-striatoentopeduncular neuron, which can be called GABAergic dynorphinergic neuron, since it expresses the peptide dynorphin (and also substance P). The GABAergic enkephalinergic neuron predominantly expresses dopamine and adenosine receptors of the D₂ and A_{2A} receptor subtype [3,9,14-17,21-23], while the GABAergic dynorphinergic neuron expresses dopamine and adenosine receptors of the D₁ and A₁ subtype [3,9,23].

We found evidence for the existence of A_{2A}-D₂ receptor interactions that modulate the function of the GABAergic enkephalinergic neuron and A₁-D₁ receptor interactions that modulate the function of GABAergic dynorphinergic neuron [23]. We and other authors also found evidence for the existence of selective heteromerization of A_{2A} and D₂ receptors and A₁ and D₁ receptors in transfected cells [4,8,10,11] and found biochemical characteristics of these receptor heteromers, called “intramembrane receptor-receptor interactions” [9], which could also be identified in the striatum (reviewed in ref.

*Address correspondence to this author at the National Institute on Drug Abuse, I.R.P., N.I.H., D.H.H.S., 5500 Nathan Shock Dr., Baltimore, MD 21224, USA; E-mail: sferré@intra.nida.nih.gov

22), therefore demonstrating the existence of A_{2A}-D₂ and A₁-D₁ receptor heteromers in the brain [14,24,25].

Then, going back to the striatal spine module, we have to differentiate between two types of modules, the one centered at the dendritic spine of the GABAergic enkephalinergic neuron, which contains A_{2A}-D₂ receptor heteromers, and the one centered at the dendritic spine of the dynorphinergic neuron, which contains A₁-D₁ receptor heteromers. In addition to the A_{2A}-D₂ and A₁-D₁ receptor heteromers, we have identified A₁-A_{2A} receptor heteromers in the glutamatergic terminals of, most probably, both striatal spine modules [12]. Furthermore, there is functional evidence for the existence of presynaptic interactions between A_{2A} and D₂ (or maybe D₄) receptors that modulate striatal glutamate release (see below).

THE ANTAGONISTIC A_{2A}-D₂ INTRAMEMBRANE RECEPTOR INTERACTION

A_{2A}-D₂ receptor heteromerization was first demonstrated in mammalian transfected cells with co-immunoprecipitation, and fluorescence and bioluminescence resonance energy transfer techniques (FRET and BRET, respectively) [10,11] (Fig. 1). By using computerized modeling, pull-down and mass spectrometry techniques, it was shown that this heteromerization depends on an electrostatic interaction between an arginine-rich epitope of the N-terminal segment of the third intracellular loop (NI3L) of the D₂ receptor and a phosphate group in the C-terminus of the A_{2A} receptor [10,26,27]. FRET and BRET, however, are difficult techniques to implement in tissues and the demonstration of the A_{2A}-D₂ receptor heteromer was demonstrated by indirect means, by identifying a biochemical characteristic, what we have called a “biochemical fingerprint” of the heteromer [14,24,25].

The concept of “intramembrane receptor interactions” was first described by Luigi Agnati and Kjell Fuxe more than 20 years ago (reviewed in ref. 9). In these interactions, stimulation of one receptor changes the binding characteristics of an adjacent receptor in membrane preparations from brain tissue or transfected cells. It is now recognized that intramembrane receptor interactions constitute a common biochemical characteristic of receptor heteromers

[14,23-25]. The antagonistic A_{2A}-D₂ intramembrane receptor interaction has been repeatedly reported by different research groups in membrane preparations from different transfected cell lines and from human and rat striatum [28-34]. In these membrane preparations, the addition of a selective A_{2A} receptor agonist decreases the ability of dopamine (or a D₂ receptor agonist) to displace the binding of a selective D₂ receptor radioligand [28-34] (Fig. 2a).

The antagonistic A_{2A}-D₂ intramembrane receptor interaction seems to determine the ability of A_{2A} receptors to control the inhibitory role of D₂ receptors on neuronal excitability and neurotransmitter release in the GABAergic enkephalinergic neuron. It was demonstrated by means of *in vivo* microdialysis experiments that the perfusion of a D₂ receptor agonist in the dorsal striatum, localization of the cell bodies of the GABAergic enkephalinergic neuron, leads to a decrease in the extracellular levels of GABA in the ipsilateral globus pallidus, localization of the nerve terminals of the enkephalinergic neurons [35]. On the other hand, the striatal perfusion of an A_{2A} receptor agonist did not produce any significant effect, but it completely counteracted the effect of the D₂ receptor agonist [35]. Thus, in this experimental setting, an A_{2A} receptor agonist behaves as a D₂ receptor antagonist. In the nucleus accumbens (ventral striatum), there is a more tonic effect of D₂ receptor stimulation by endogenous dopamine on neurotransmitter release by the GABAergic enkephalinergic neuron [36]. Also in this experimental setting the striatal perfusion of an A_{2A} receptor agonist produced the same effects than a D₂ receptor antagonist, i.e., an increase in the extracellular levels of GABA in the ipsilateral ventral pallidum [36].

The striatal output is determined by the bursting activity of the GABAergic striatal efferent neurons. These bursts are driven by cortico-striatal inputs that depolarize the GABAergic enkephalinergic and dynorphinergic neurons from their resting hyperpolarized membrane potential around -80 mV, the down-state, to a more depolarized level near -55 mV, the up-state [37]. These down- to up-state transitions require channels that can be regulated by striatal transmitters acting through GPCRs, such as the interacting A_{2A} and D₂ receptors in the GABAergic enkephalinergic neurons. In a re-




Fig. (1). Bioluminescence and fluorescence resonance energy transfer (BRET and FRET, respectively) techniques allow the demonstration of neurotransmitter receptor heteromers in the natural environment of the living cell. FRET can be obtained when two fluorescent proteins, one acting as a donor and the other acting as acceptor, are close enough (within 10 nm). Therefore, to demonstrate receptor dimerization, cDNA constructs of one receptor fused to the fluorescent donor and another receptor fused to the fluorescent acceptor are prepared and transfected in a heterologous cell system. Different versions of the green fluorescent protein (GFP) are currently used as donors whereas yellow fluorescent proteins (YFPs) are used as acceptors. If the two receptors are forming dimers, the acceptor fluorescent signal is detected after donor excitation. In BRET, instead of a fluorescent donor one of the receptors is fused to the luminescent protein, *Renilla luciferase* (*Rluc*), which upon addition of a substrate (coelenterazine or Deep Blue C) allows energy transfer and excitation of the fluorescent acceptor.

cent study, perforated patch clamp recordings on acute brain slices, together with the loading of competitive peptides to block specific protein-protein interactions, were used to characterize the role of A_{2A}-D₂ receptors interactions in the modulation of down- to up-state transitions, modeled *in vitro* by the application of NMDA. A D₂ receptor agonist abolished the firing in up-state and inhibited the down/up-state transition in the GABAergic enkephalinergic neurons by a mechanism involving the regulation of L-type calcium channel Ca_v1.3 through protein-protein interactions with scaffold proteins Shank1/3 (Azdad *et al.*, Society for Neuroscience Abstracts, 2007). On the other hand, the A_{2A} receptor agonist CGS 21680 did not induce any modification in state transition or in the firing frequency, but it totally reversed the effects of D₂ receptor activation (Azdad *et al.*). This action was blocked by the selective A_{2A} receptor antagonist SCH 58261 (1 μM) and was absent in A_{2A} receptor knock-out mice (Azdad *et al.*). The application of peptides containing the same aminoacid sequence than the epitopes involved in A_{2A}-D₂ receptor heteromerization counteracted the ability of A_{2A} receptor activation to antagonize the effect of D₂ receptor activation (Azdad *et al.*). This demonstrates that A_{2A}-D₂ receptors heteromerization is strictly mandatory for the A_{2A} receptor-mediated control of D₂ receptor-mediated modulation of the excitability of GABAergic enkephalinergic neurons.

These effects on neurotransmitter release and neuronal excitability are paralleled by effects on motor activity and other behavioral responses, where selective A_{2A} receptor agonists or antagonists respectively counteract or potentiate the motor activation induced by dopamine D₂ receptor agonists [38-42]. Consequently, we predicted 15 years ago that A_{2A} receptor antagonists could be useful in Parkinson's disease, especially potentiating the effects of L-dopa or D₂ receptor agonists [43]. In fact, in different experimental models of Parkinson's disease, A_{2A} receptor antagonists potentiate the motor activating effects of L-DOPA or D₂ receptor agonists (for review see ref. 42). Also in agreement, in the rodent dopamine-denervated striatum, local application of a D₂ receptor agonist potently inhibits the increased neuronal activity (compared with the non-denervated striatum) and this effect is counteracted or potentiated with application of A_{2A} agonists or antagonists, respectively [41]. Importantly, the A_{2A} receptor ligands did not have any significant effects on their own [41]. On the other hand, in patients with Parkinson's disease the association of L-DOPA and an A_{2A} receptor antagonist has already given promising therapeutic results (reviewed in ref. 44).

THE ANTAGONISTIC A_{2A}-D₂ RECEPTOR INTERACTION AT THE SECOND MESSENGER LEVEL

A_{2A} receptor, through its coupling to G_{oif} proteins, can potentially stimulate adenylyl-cyclase and activate the cAMP-PKA signaling pathway, with phosphorylation of several PKA substrates, such as DARPP-32, CREB and AMPA receptors and the consequent increase in the expression of different genes, such as *c-fos* or *preproenkephalin* in the GABAergic enkephalinergic neuron [3,9,14,16,23,24]. For instance, in CHO cells stably transfected with A_{2A} receptors, the addition of an A_{2A} receptor agonist produced cAMP accumulation, CREB phosphorylation and increase in *c-fos* expression [31]. In the same cell line we could demonstrate the existence of an antagonistic A_{2A}-D₂ intramembrane receptor interaction with radioligand binding experiments [31]. Furthermore, in the same cell line, we found a reciprocal antagonistic A_{2A}-D₂ receptor interaction by which the D₂ receptor, which can couple to G_{i/o} proteins, inhibits the effects of A_{2A} receptor stimulation at the level of adenylyl cyclase [31] (Fig. 2b). A D₂ receptor agonist did not produce a significant effect on its own, but it completely counteracted the effect induced by A_{2A} receptor stimulation on cAMP accumulation, CREB phosphorylation and *c-fos* expression [31].

The two kind of reciprocal antagonistic A_{2A}-D₂ receptor interactions could also be demonstrated in another cell line, a human

SH-SY5Y neuroblastoma cell line that constitutively expresses A_{2A} receptors and with transfected D₂ receptors [32]. In this cell line, D₂ receptor stimulation completely counteracted cAMP accumulation induced by an A_{2A} receptor agonist [8]. But, at the same time, an antagonistic A_{2A}-D₂ intramembrane receptor interaction with functional consequences could be demonstrated with radioligand binding experiments and intracellular Ca²⁺ responses. Thus, D₂ receptor activation inhibited a KCL-induced increase in intracellular concentration of Ca²⁺, which was counteracted by A_{2A} receptor stimulation [32]. This is most probably the same mechanism by which the antagonistic A_{2A}-D₂ intramembrane receptor interaction controls the excitability of the GABAergic enkephalinergic neurons (see above).

It is intriguing that both types of reciprocal antagonistic A_{2A}-D₂ receptor interactions coexist in the same cells and, in fact, they do coexist in the brain. Under normal conditions, there is a strong tonic activation of D₂ receptors that blocks the ability of A_{2A} receptors to signal through the cAMP-PKA pathway. For instance, in the rodent striatum, the *in vivo* administration of D₂ receptor antagonists produces a significant increase in the PKA-dependent phosphorylation of DARPP-32 or the AMPA receptor and an increase in the expression of *c-fos* and *preproenkephalin* genes, which depends on the ability of D₂ receptor blockade to liberate A_{2A} receptor signaling activated by endogenous adenosine [45,46]. Thus, the effect of the D₂ receptor antagonists was counteracted by the previous administration of an A_{2A} receptor antagonist, which did not have a significant effect on its own [45,46].

COEXISTENCE OF THE RECIPROCAL ANTAGONISTIC A_{2A}-D₂ RECEPTOR INTERACTIONS

There is therefore evidence for the coexistence of two reciprocal antagonistic interactions between A_{2A} and D₂ receptors in the GABAergic enkephalinergic neurons. There is an antagonistic A_{2A}-D₂ intramembrane receptor interaction, which depends on A_{2A}-D₂ receptor heteromerization, which modulates neuronal excitability and neurotransmitter release; and there is an antagonistic A_{2A}-D₂ receptor interaction at the level of adenylyl-cyclase that modulates protein phosphorylation and gene expression. These results provide a clear example of a functional dissociation between neuronal excitability and gene expression. Thus, co-stimulation of A_{2A} and D₂ receptors implies a simultaneous A_{2A} receptor-mediated inhibition of the D₂ receptor-mediated modulation of neuronal excitability and a D₂ receptor-mediated inhibition of the A_{2A} receptor-mediated modulation of gene expression.

There are at least two possible, but not exclusive, mechanisms that could explain this apparently incompatible coexistence of reciprocal antagonistic A_{2A}-D₂ receptor interactions. First, one possibility is a different G-protein coupling between different sets of D₂ receptors. Thus, it has been shown that D₂ receptor couples to G_{i/o}, and therefore negatively to adenylyl-cyclase, when not forming heteromers, and that it couples to G_{q/11}-PLC signaling when forming heteromers with D₁ receptors [47]. In fact, A_{2A} receptor is morphologically and functionally very similar to the D₁ receptor. They both have a short third intracellular loop and a long acidic C-terminus and they both couple to G_{oif} proteins in the striatum. Also, they most probably use the same epitope (phosphorylated serine in the C-terminus) for their physical interaction with the D₂ receptor [27]. Furthermore, the inhibitory role of D₂ receptors in the excitability of GABAergic enkephalinergic neurons depends mostly on the suppression of Ca²⁺ currents through L-type voltage-dependent calcium channels, which depends on activation of the G_{q/11}-PLC signaling pathway [37].

As shown in Fig. (2), the most probable scenario is the one that considers that, when not forming heteromers, the most common basic composition of A_{2A} and D₂ receptors and any GPCR is as homodimers [25,48-52], and that only one (heterotrimeric) G protein binds to a receptor dimer [48-50]. The selectivity of G protein recognition is determined by multiple intracellular regions, with the

most critical regions being the second intracellular loop (I2L), the N13L and the C-terminal segment of the third intracellular loop [53]. The relative contribution of these intracellular receptor domains to the selectivity of G protein recognition varies among different classes of GPCRs [53]. For D₂ receptors, the arginine-rich epitope of the N13L has been shown to be fundamental for the coupling to G_{i/o} proteins [54]. The same epitope has been demonstrated to bind to calmodulin and, as mentioned before, to the C-terminus of the A_{2A} receptor [10,26,27,55,56]. These findings would agree with the inability of D₂ receptor heteromers to signal through G_{i/o} proteins when bound to A_{2A} receptors (see above) or to calmodulin [55].

Another possibility for the coexistence of reciprocal antagonistic A_{2A}-D₂ receptor interactions is the existence of an additional partner for A_{2A} and D₂ receptors to interact at the adenylyl-cyclase level. A_{2A} receptors have also been found to form receptor heteromers with metabotropic glutamate mGlu₅ receptors both in transfected cells and in the striatum [7]. In transfected cells and in the striatum, co-stimulation of A_{2A} and mGlu₅ receptors produced a very synergistic effect on *c-fos* expression, which depends on interactions between both receptors at the adenylyl-cyclase and at the MAPK levels [7,57]. *In vivo* experiments demonstrated that co-stimulation of A_{2A} and mGlu₅ receptors, with the central administration of selective agonists, allows A_{2A} receptor to get rid of the tonic inhibitory effect of D₂ receptor and signal through cAMP-PKA pathway [7]. Since these A_{2A}-mGlu₅-D₂ receptor interactions can be demonstrated in animal models of Parkinson's disease [58,59], we postulated that co-administration of A_{2A} and mGlu₅ receptor antagonists could be used as a therapeutic strategy in this disease [58]. Similarly, *in vivo* microdialysis experiments have shown that A_{2A}-mGlu₅-D₂ receptor interactions modulate the function of the GABAergic enkephalinergic neurons of the nucleus accumbens [60], which can have implications for schizophrenia and drug addiction.

THE SYNERGISTIC A_{2A}-D₂ RECEPTOR INTERACTION AT THE G PROTEIN LEVEL

The antagonistic A_{2A}-D₂ receptor interaction at the adenylyl cyclase level just described depends on the ability of activated D₂ receptors to counteract A_{2A} receptor-mediated type V adenylyl-cyclase (ACV) activation [61]. Under some conditions, a synergistic A_{2A}-D₂ receptor interaction can also be detected [34,62], which seems to depend on the presence of an activator of G protein signaling (AGS3), which facilitates a synergistic interaction between G_{s/o1f} and G_{i/o}-coupled receptors on the activation of types II/IV adenylyl cyclase (ACII/IV) (reviewed in ref. 63). AGS3 binds preferentially to G_iα, and stabilizes the GDP-bound conformation of G_i, thereby dampening the signaling of the receptor through G_i-GTP, while simultaneously increasing the activity of Gβγ-regulated effectors [64]. Upon co-activation of the G_i-coupled D₂ receptor, unbound βγ subunits released in the presence of AGS3 are free to transiently stimulate ACII/IV upon co-activation of the G_{o1f}-coupled A_{2A} receptor, leading to a paradoxical increase in cAMP-PKA signaling [63] (Fig. 2e).

The antagonistic A_{2A}-D₂ receptor interaction at the adenylyl-cyclase is however predominant in most conditions, since ACV is the most expressed type of adenylyl-cyclase in the striatum [65]. However, the synergistic interaction can become particularly important during conditions of upregulation of AGS3, such as during chronic treatment with addictive drugs. It has recently been shown that withdrawal from repeated treatment with cocaine (self- or non-self-administered) up-regulates AGS3 in the prefrontal cortex and in the core region of the nucleus accumbens [66]. In rats, knocking down AGS3 expression in the prefrontal cortex or the nucleus accumbens core (with antisense oligonucleotides) counteracts reinstatement of cocaine- or heroin-seeking behaviour, respectively [66,67]. Therefore, upregulation of AGS3, with the consequent

dampening of G_iα signaling while simultaneously promoting βγ-dependent signaling of G_s-coupled receptors, such as D₁ in the prefrontal cortex or in the nucleus accumbens [66], or A_{2A} in the nucleus accumbens [63], can be an important mechanism responsible for the pathophysiologic changes associated with different addictive drugs. Thus, we have postulated that A_{2A} receptor antagonists could be useful in the treatment of drug addiction and relapse during drug withdrawal [63].

PRESYNAPTIC ANTAGONISTIC A_{2A}-D₂ RECEPTOR INTERACTIONS

The GABAergic enkephalinergic neuron does not only express A_{2A} and D₂ receptors in the somatodendritic area, but both receptors are also colocalized in the nerve terminals [16]. Stimulation of A_{2A} receptors in the globus pallidus, localization of the nerve terminals of the GABAergic enkephalinergic neurons, has been shown to stimulate GABA release by using *in vivo* microdialysis and slice preparations [68-70]. This effect of A_{2A} receptor stimulation is dependent on the activation of the cAMP-PKA pathway [70], which depends on the already reported ability of PKA to phosphorylate different elements of the machinery involved in vesicular fusion [71]. In fact, the ability of A_{2A} receptors to stimulate neurotransmitter release through cAMP-PKA signaling has also been demonstrated for acetylcholine in the striatum [72] and serotonin in the hippocampus [73]. In the globus pallidus, stimulation of D₂ receptors produces a strong counteraction of A_{2A} receptor-mediated GABA release [68,70]. Altogether, this has all the characteristics of the antagonistic A_{2A}-D₂ receptor interaction at the second messenger level described above, which does not seem to depend on A_{2A}-D₂ receptor heteromerization. Recent studies in an animal model of Parkinson's disease suggest that this pallidal A_{2A}-D₂ receptor interaction may contribute to the antiparkinsonian effects of the co-administration of A_{2A} antagonists and L-DOPA or D₂ receptor agonists [74].

Finally, there is still another presynaptic localization where A_{2A}-D₂ receptor interactions seem to have an important functional role. We have recently demonstrated that A_{2A} receptors localized in striatal glutamatergic terminals play a very important control of striatal glutamate release [12]. These receptors form heteromers with A₁ receptors, and the A₁-A_{2A} receptor heteromer constitutes a "concentration-dependent switch" that regulates glutamate release depending on the extracellular concentration of adenosine. Thus, low concentrations of adenosine inhibit glutamate release by stimulating the A₁ receptor, while higher concentrations induce glutamate release by also stimulating A_{2A} receptors, which shuts down A₁ receptor signaling by means of an antagonistic A₁-A_{2A} intramembrane receptor interaction [3,12,14,16,17]. However, the A_{2A} receptor-dependent modulation of glutamate release seems to be under an inhibitory control by a co-localized D₂ receptor. Thus, dopamine denervation strongly potentiates A_{2A} receptor agonist-mediated stimulation of striatal glutamate release [71]. Again, this has the biochemical characteristics of an antagonistic A_{2A}-D₂ receptor interaction at the second messenger level, which might not depend on A_{2A}-D₂ receptor heteromerization or which might depend on A_{2A}-mGlu₅-D₂ receptor interactions (see above). In fact, A_{2A} and mGlu₅ receptors have been found to be co-localized in a large proportion of striatal glutamatergic terminals, where they facilitate glutamate release in a synergistic manner [76]. This presynaptic antagonistic A_{2A}-D₂ receptor interaction can also have implications for the treatment of Parkinson's disease with the co-administration of A_{2A} antagonists and L-DOPA or D₂ receptor agonists. Thus, in the striatum of parkinsonian animals there is overactivity of striatal glutamatergic transmission, which is restored with L-DOPA treatment. In recent electrophysiological experiments in cortico-striatal slices, concomitant activation of D₂ receptors and inactivation of A_{2A} receptors has shown to produce a very significant decrease in striatal glutamate release [77].




Fig. (2). Different types of A_{2A} - D_2 receptor interactions: **a.** antagonistic A_{2A} - D_2 intramembrane receptor interaction which depends on A_{2A} - D_2 receptor heteromerization and $G_{q/11}$ -PLC signaling; **b.** antagonistic A_{2A} - D_2 receptor interaction at the adenylyl-cyclase level, which depends on $G_{s/olf}$ - and G_i - type V adenylyl-cyclase signaling; **c.** synergistic A_{2A} - D_2 receptor interaction at the adenylyl-cyclase level, which depends on upregulation of AGS3 and $G_{s/olf}$ - and G_i - type II/IV adenylyl-cyclase signaling.

CONCLUSIONS

In conclusion, we are beginning to understand the complexities of A_2 - D_2 receptor interactions, which play a very important role in basal ganglia physiology. Furthermore, it is obvious that the analysis of the different A_{2A} - D_2 receptor interactions will have important general implications about the processing of information of GPCRs. The present review stresses the fact that multiple and functionally different interactions can occur between just two different GPCRs, sometimes involving receptor heteromerization. Particularly important is the example of the coexistence of two apparently incompatible reciprocal antagonistic interactions between A_{2A} and D_2 receptors, which allows a segregated control of neuronal excitability and gene expression in the GABAergic enkephalinergic neurons by the same receptors. The analysis of A_2 - D_2 receptor interactions will

also have implications for the pathophysiology and treatment of basal ganglia disorders and drug addiction, in view of their key processing role in the computation of information by the striatal spine module [16].

ACKNOWLEDGEMENTS

Supported by the NIDA IRP funds and grants from FRS-FNRS and FMRE (to SNS), Spanish "Ministerio de Educación y Ciencia" (SAF2005-00903 to FC), "Ministerio de Ciencia y Tecnología" (SAF2006-05481) and "Fundació La Marató TV3 (060110).

REFERENCES

- [1] Snyder SH, Ferris CD. Novel neurotransmitters and their neuropsychiatric relevance. *Am J Psychiatry* 2000; 157: 1738-51.

- [2] Fredholm BB, IJzerman AP, Jacobson KA, Klotz KN, Linden J. International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol Rev* 2001; 53: 527-52.
- [3] Schiffmann SN, Fisone G, Moresco R, Cunha R, Ferré S. Adenosine A_{2A} receptors and basal ganglia physiology. *Prog Neurobiol* 2007; 83: 277-92.
- [4] Ginés S, Hillion J, Torvinen M, Le Crom S, Casado V, Canela EI, *et al.* Dopamine D₁ and adenosine A₁ receptors form functionally interacting heteromeric complexes. *Proc Natl Acad Sci USA* 2000; 97: 8606-11.
- [5] Ciruela F, Escriche M, Burgueno J, Angulo E, Casado V, Soloviev MM, *et al.* Metabotropic glutamate 1alpha and adenosine A₁ receptors assemble into functionally interacting complexes. *J Biol Chem* 2001; 276: 18345-51.
- [6] Yoshioka K, Saitoh O, Nakata H. Heteromeric association creates a P_{2Y}-like adenosine receptor. *Proc Natl Acad Sci USA* 2001; 98: 7617-22.
- [7] Ferré S, Karcz-Kubicha M, Hope BT, Popoli P, Burgueno J, Gutierrez MA, *et al.* Synergistic interaction between adenosine A_{2A} and glutamate mGlu₅ receptors: implications for striatal neuronal function. *Proc Natl Acad Sci USA* 2002; 99: 11940-5.
- [8] Hillion J, Canals M, Torvinen M, Casado V, Scott R, Terasmaa A, *et al.* Coaggregation, cointernalization, and codesensitization of adenosine A_{2A} receptors and dopamine D₂ receptors. *J Biol Chem* 2002; 277: 18091-7.
- [9] Agnati LF, Ferré S, Lluís C, Franco R, Fuxe K. Molecular mechanisms and therapeutical implications of intramembrane receptor/receptor interactions among heptahelical receptors with examples from the striatopallidal GABA neurons. *Pharmacol Rev* 2003; 55: 509-50.
- [10] Canals M, Marcellino D, Fanelli F, Ciruela F, de Benedetti P, Goldberg SR, *et al.* Adenosine A_{2A}-dopamine D₂ receptor-receptor heteromerization: qualitative and quantitative assessment by fluorescence and bioluminescence energy transfer. *J Biol Chem* 2003; 278: 46741-9.
- [11] Kamiya T, Saitoh O, Yoshioka K, Nakata H. Oligomerization of adenosine A_{2A} and dopamine D₂ receptors in living cells. *Biochem Biophys Res Commun* 27; 306: 544-9.
- [12] Ciruela F, Casado V, Rodrigues RJ, Lujan R, Burgueno J, Canals M, *et al.* Presynaptic control of striatal glutamatergic neurotransmission by adenosine A₁-A_{2A} receptor heteromers. *J Neurosci* 2006; 26: 2080-7.
- [13] Carriba P, Ortiz O, Patkar K, Justinova Z, Stroik J, Themann A, *et al.* Striatal adenosine A_{2A} and cannabinoid CB₁ receptors form functional heteromeric complexes that mediate the motor effects of cannabinoids. *Neuropsychopharmacology* 2007; 32: 2249-59.
- [14] Ferré S, Ciruela F, Woods AS, Lluís C, Franco R. Functional relevance of neurotransmitter receptor heteromers in the central nervous system. *Trends Neurosci* 2007; 30: 440-6.
- [15] Gerfen CR. Basal Ganglia. In: Paxinos G Ed, *The rat nervous system*. Amsterdam, Elsevier Academic Press 2004; 445-508.
- [16] Ferré S, Agnati LF, Ciruela F, Lluís C, Woods AS, Fuxe K, *et al.* Neurotransmitter receptor heteromers and their integrative role in 'local modules': The striatal spine module. *Brain Res Rev* 2007; 55: 55-67.
- [17] Ferré S, Borycz J, Goldberg SR, Hope BT, Morales M, Lluís C, *et al.* Role of adenosine in the control of homosynaptic plasticity in striatal excitatory synapses. *J Integr Neurosci* 2005; 4: 445-64.
- [18] Newman EA. New roles for astrocytes: Regulation of synaptic transmission. *Trends Neurosci* 2003; 26: 536-42.
- [19] Hertz L, Zielke R. Astrocytic control of glutamatergic activity: astrocytes as stars of the show. *Trends Neurosci* 2004; 27: 735-43.
- [20] Pascual O, Casper KB, Kubera C, Zhang J, Revilla-Sanchez R, Sul JY, *et al.* Astrocytic purinergic signaling coordinates synaptic networks. *Science* 2005; 310: 113-6.
- [21] Schiffmann SN, Jacobs O, Vanderhaeghen JJ. Striatal restricted adenosine A₂ receptor (RDC8) is expressed by enkephalin but not by substance P neurons: an *in situ* hybridization histochemistry study. *J Neurochem* 1991; 57: 1062-7.
- [22] Schiffmann SN, Vanderhaeghen JJ. Adenosine A₂ receptors regulate the gene expression of striatopallidal and striatonigral neurons. *J Neurosci* 1993; 13: 1080-7.
- [23] Ferré S, Fredholm BB, Morelli M, Popoli P, Fuxe K. Adenosine-dopamine receptor-receptor interactions as an integrative mechanism in the basal ganglia. *Trends Neurosci* 1997; 20: 482-7.
- [24] Ferré S, Ciruela F, Quiroz C, Lujan R, Popoli P, Cunha RA, *et al.* Adenosine receptor heteromers and their integrative role in striatal function. *ScientificWorldJournal* 2007; 7: 74-85.
- [25] Franco R, Casado V, Cortes A, Ferrada C, Mallol J, Woods A, *et al.* Basic concepts in G-protein-coupled receptor homo- and heteromerization. *ScientificWorldJournal* 2007; 7: 48-57.
- [26] Ciruela F, Burgueno J, Casado V, Canals M, Marcelino D, Goldberg SR, *et al.* Combining mass spectrometry and pull-down techniques for the study of receptor heteromerization. Direct epitope-epitope electrostatic interactions between adenosine A_{2A} and dopamine D₂ receptors. *Anal Chem* 2004; 76: 5354-63.
- [27] Woods AS, Ferré S. The amazing stability of the arginine-phosphate electrostatic interaction. *J Proteome Res* 2005; 4: 1397-402.
- [28] Ferré S, von Euler G, Johansson B, Fredholm BB, Fuxe K. Stimulation of high affinity adenosine A-2 receptors decreases the affinity of dopamine D-2 receptors in rat striatal membranes. *Proc Natl Acad Sci USA* 1991; 88: 7238-41.
- [29] Dasgupta S, Ferré S, Kull B, Hendlund P, Finnman U-B, Ahlberg S, *et al.* Adenosine A_{2A} receptors modulate the binding characteristics of dopamine D₂ receptors in stably cotransfected fibroblast cells. *Eur J Pharmacol* 1996; 316: 325-31.
- [30] Dixon AK, Widdowson L, Richardson PJ. Desensitisation of the adenosine A₁ receptor by the A_{2A} receptor in the rat striatum. *J Neurochem* 1997; 69: 315-21.
- [31] Kull B, Ferré S, Arslan G, Svenningsson P, Fuxe K, Owman C, *et al.* Reciprocal interactions between adenosine A_{2A} and dopamine D₂ receptors in CHO cells co-transfected with the two receptors. *Biochem Pharmacol* 1999; 58: 1035-45.
- [32] Salim H, Ferré S, Dalal A, Peterfreund RA, Fuxe K, Vincent JD, *et al.* Activation of adenosine A₁ and A_{2A} receptors modulates dopamine D₂ receptor-induced responses in stably cotransfected human neuroblastoma cells. *J Neurochem* 2000; 74: 432-9.
- [33] Diaz-Cabiale Z, Hurd Y, Guidolin D, Finnman U-B, Zoli M, Vanderhaeghen J-J, *et al.* Adenosine A_{2A} agonist CGS 21680 decreases the affinity of dopamine D₂ receptors for dopamine in human striatum. *NeuroReport* 2001; 12: 1831-4.
- [34] Kudlacek O, Just H, Korkhov VM, Vartian N, Klinger M, Pankevych H, *et al.* The human D₂ dopamine receptor synergizes with the A_{2A} adenosine receptor to stimulate adenylyl cyclase in PC12 cells. *Neuropsychopharmacology* 2003; 28: 1317-27.
- [35] Ferré S, O'Connor WT, Fuxe K, Ungerstedt U. The striopallidal neuron: a main locus for adenosine-dopamine interactions in the brain. *J Neurosci* 1993; 13: 5402-6.
- [36] Ferré S, O'Connor WT, Snaprud P, Ungerstedt U, Fuxe K. Antagonistic interaction between adenosine A_{2A} and dopamine D₂ receptors in the ventral striopallidal system. Implications for the treatment of schizophrenia. *Neuroscience* 1994; 63: 765-73.
- [37] Surmeier DJ, Ding J, Day M, Wang Z, Shen W. D₁ and D₂ dopamine-receptor modulation of striatal glutamatergic signaling in striatal medium spiny neurons. *Trends Neurosci* 2007; 30: 228-35.
- [38] Ferré S, Herrera-Marschitz M, Grabowska-Andén M, Ungerstedt U, Casas M, Andén N-E. Postsynaptic dopamine/adenosine interaction: I. Adenosine analogues inhibit a D₂-mediated behaviour in short-term reserpinized mice. *Eur J Pharmacol* 1991; 192: 30-5.
- [39] Ferré S, Herrera-Marschitz M, Grabowska-Andén M, Ungerstedt U, Casas M, Andén N-E. Postsynaptic dopamine/adenosine interaction: II. Postsynaptic dopamine agonism and adenosine antagonism of methylxanthines in short-term reserpinized mice. *Eur J Pharmacol* 1991; 192: 36-42.
- [40] Rimondini R, Ferré S, Giménez-Llort L, Ögren SO, Fuxe K. Differential effects of selective adenosine A₁ and A_{2A} receptor agonists on dopamine receptor agonist-induced behavioural responses in rats. *Eur J Pharmacol* 1998; 347: 153-8.
- [41] Stromberg I, Popoli P, Müller CE, Ferré S, Fuxe K. Electrophysiological and behavioural evidence for an antagonistic modulatory role of adenosine A_{2A} receptors in dopamine D₂ receptor regulation in the rat dopamine-denervated striatum. *Eur J Neurosci* 2000; 12: 4033-7.
- [42] Ferré S, Popoli P, Gimenez-Llort L, Rimondini R, Müller CE, Stromberg I, *et al.* Adenosine/dopamine interaction: implications

- for the treatment of Parkinson's disease. *Parkinsonism Relat Disord* 2001; 7: 235-41.
- [43] Ferré S, Fuxe K, von Euler G, Johansson B, Fredholm BB. Adenosine-dopamine interactions in the brain. *Neuroscience* 1992; 51: 501-12.
- [44] Muller CE, Ferré S. Blocking striatal adenosine A2A receptors: A new strategy for basal ganglia disorders. *Recent Pat CNS Drug Disc* 2007; 2: 1-21.
- [45] Svenningsson P, Lindskog M, Ledent C, Parmentier M, Greengard P, Fredholm BB, *et al.* Regulation of the phosphorylation of the dopamine- and cAMP-regulated phosphoprotein of 32 kDa *in vivo* by dopamine D1, dopamine D2, and adenosine A2A receptors. *Proc Natl Acad Sci USA* 2000; 97: 1856-60.
- [46] Hakansson K, Galdi S, Hendrick J, Snyder G, Greengard P, Fisone G. Regulation of phosphorylation of the GluR1 AMPA receptor by dopamine D2 receptors. *J Neurochem* 2006; 96: 482-8.
- [47] Rashid AJ, So CH, Kong MM, Furtak T, El-Ghundi M, Cheng R, *et al.* D1-D2 dopamine receptor heterooligomers with unique pharmacology are coupled to rapid activation of Gq/11 in the striatum. *Proc Natl Acad Sci USA* 2007; 104: 654-9.
- [48] Banères JL, Parelo J. Structure-based analysis of GPCR function: evidence for a novel pentameric assembly between the dimeric leukotriene B4 receptor BLT1 and the G-protein. *J Mol Biol* 2003; 329: 815-29.
- [49] Liang Y, Fotiadis D, Filipek S, Saperstein DA, Palczewski K, Engel A. Organization of the G protein-coupled receptors rhodopsin and opsin in native membranes. *J Biol Chem* 2003; 278: 21655-62.
- [50] Herrick-Davis K, Grinde E, Harrigan TJ, Mazurkiewicz JE. Inhibition of serotonin 5-hydroxytryptamine2c receptor function through heterodimerization: receptor dimers bind two molecules of ligand and one G-protein. *J Biol Chem* 2005; 280: 40144-51.
- [51] Franco R, Casado V, Mallol J, Ferrada C, Ferré S, Fuxe K, *et al.* The "two-state dimer receptor model". A general model for receptor dimers. *Mol Pharmacol* 2006; 69: 1905-12.
- [52] Casado V, Cortes A, Ciruela F, Mallol J, Ferré S, Lluís C, *et al.* Old and new ways to calculate the affinity of agonists and antagonists interacting with G-protein-coupled monomeric and dimeric receptors: The receptor-dimer cooperativity index. *Pharmacol Ther* 2007; 116: 343-54.
- [53] Wess J. Molecular basis of receptor/G-protein-coupling selectivity. *Pharmacol Ther* 1998; 80: 231-64.
- [54] Voss T, Wallner E, Czernilofsky AP, Freissmuth M. Amphipathic alpha-helical structure does not predict the ability of receptor-derived synthetic peptides to interact with guanine nucleotide-binding regulatory proteins. *J Biol Chem* 1993; 268: 4637-42.
- [55] Bofill-Cardona E, Kudlacek O, Yang Q, Ahorn H, Freissmuth M, Nanoff C. Binding of calmodulin to the D2-dopamine receptor reduces receptor signaling by arresting the G protein activation switch. *J Biol Chem* 2000; 275: 32672-80.
- [56] Liu Y, Buck DC, Macey TA, Lan H, Neve KA. Evidence that calmodulin binding to the dopamine D2 receptor enhances receptor signaling. *J Recept Signal Transduct Res* 2007; 27: 47-65.
- [57] Nishi A, Liu F, Matsuyama S, Hamada M, Higashi H, Nairn AC, *et al.* Metabotropic mGlu5 receptors regulate adenosine A2A receptor signaling. *Proc Natl Acad Sci USA* 2003; 100: 1322-7.
- [58] Popoli P, Pezzola A, Torvinen M, Reggio R, Pintor A, Scarchili L, *et al.* The selective mGlu5 receptor agonist CHPG inhibits quinpirole-induced turning in 6-hydroxydopamine-lesioned rats and modulates the binding characteristics of dopamine D2 receptors in the rat striatum: Interactions with adenosine A2A receptors. *Neuropharmacology* 2001; 25: 505-13.
- [59] Kachroo A, Orlando LR, Grandy DK, Chen JF, Young AB, Schwarzschild MA. Interactions between metabotropic glutamate 5 and adenosine A2A receptors in normal and parkinsonian mice. *J Neurosci* 2005; 25: 10414-19.
- [60] Díaz-Cabiale Z, Vivó M, Del Arco A, O'Connor WT, Harte MK, Müller CE, *et al.* Metabotropic glutamate mGlu5 receptor-mediated modulation of the ventral striopallidal GABA pathway. Interactions with adenosine A2A and dopamine D2 receptors. *Neurosci Lett* 2002; 324: 154-8.
- [61] Lee KW, Hong JH, Choi IY, Che Y, Lee JK, Yang SD, *et al.* Impaired D2 dopamine receptor function in mice lacking type 5 adenylyl cyclase. *J Neurosci* 2002; 22: 7931-40.
- [62] Yao L, Arolfo MP, Dohrman DP, Jiang Z, Fan P, Fuchs S, *et al.* Betagamma dimers mediate synergy of dopamine D2 and adenosine A2 receptor-stimulated PKA signaling and regulate ethanol consumption. *Cell* 2002; 109: 733-43.
- [63] Ferré S, Diamond I, Goldberg SR, Yao L, Hourani SMO, Huang ZL, *et al.* Adenosine A2A receptors in ventral striatum, hypothalamus and nociceptive circuitry. Implications for drug addiction, sleep and pain. *Prog Neurobiol* 2007; 83: 332-47.
- [64] Blumer JB, Cismowski MJ, Sato M, Lanier SM. AGS proteins: receptor-independent activators of G-protein signaling. *Trends Pharmacol Sci* 2005; 26: 470-6.
- [65] Chern Y. Regulation of adenylyl cyclase in the central nervous system. *Cell Signal* 2000; 12: 195-204.
- [66] Bowers MS, McFarland K, Lake RW, Peterson YK, Lapish CC, Gregory ML, *et al.* Activator of G protein signaling 3: a gatekeeper of cocaine sensitization and drug seeking. *Neuron* 2004; 42: 269-81.
- [67] Yao L, McFarland K, Fan P, Jiang Z, Inoue Y, Diamond I. Activator of G protein signaling 3 regulates opiate activation of protein kinase A signaling and relapse of heroin-seeking behavior. *Proc Natl Acad Sci USA* 2005; 102: 8746-51.
- [68] Mayfield RD, Suzuki F, Zahniser NR. Adenosine A2a receptor modulation of electrically evoked endogenous GABA release from slices of rat globus pallidus. *J Neurochem* 1993; 60: 2334-7.
- [69] Ochi M, Koga K, Kurokawa M, Kase H, Nakamura J, Kuwana Y. Systemic administration of adenosine A(2A) receptor antagonist reverses increased GABA release in the globus pallidus of unilateral 6-hydroxydopamine-lesioned rats: a microdialysis study. *Neuroscience* 2000; 100: 53-62.
- [70] Shindou T, Nonaka H, Richardson PJ, Mori A, Kase H, Ichimura M. Presynaptic adenosine A(2A) receptors enhance GABAergic synaptic transmission via a cyclic AMP dependent mechanism in the rat globus pallidus. *Br J Pharmacol* 2002; 136: 296-302.
- [71] Leenders AG, Sheng ZH. Modulation of neurotransmitter release by the second messenger-activated protein kinases: implications for presynaptic plasticity. *Pharmacol Ther* 2005; 105: 69-84.
- [72] Gubituz AK, Widdowson L, Kurokawa M, Kirkpatrick KA, Richardson PJ. Dual signalling by the adenosine A2a receptor involves activation of both N- and P-type calcium channels by different G proteins and protein kinases in the same striatal nerve terminals. *J Neurochem* 1996; 67: 374-81.
- [73] Okada M, Nutt DJ, Murakami T, Zhu G, Kamata A, Kawata Y, *et al.* Adenosine receptor subtypes modulate two major functional pathways for hippocampal serotonin release. *J Neurosci* 2001; 21: 628-40.
- [74] Simola N, Fenu S, Baraldi PG, Tabrizi MA, Morelli M. Involvement of globus pallidus in the antiparkinsonian effects of adenosine A(2A) receptor antagonists. *Exp Neurol* 2006; 202: 255-7.
- [75] Tanganelli S, Sandager Nielsen K, Ferraro L, Antonelli T, Kehr J, Franco R, *et al.* Striatal plasticity at the network level. Focus on adenosine A2A and D2 interactions in models of Parkinson's disease. *Parkinsonism Relat Disord* 2004; 10: 273-80.
- [76] Rodrigues RJ, Alfaro TM, Rebola N, Oliveira CR, Cunha RA. Colocalization and functional interaction between adenosine A and metabotropic group 5 receptors in glutamatergic nerve terminals of the rat striatum. *J Neurochem* 2005; 92: 433-41.
- [77] Tozzi A, Tschertner A, Belcastro V, Tantucci M, Costa C, Picconi B, *et al.* Interaction of A2A adenosine and D2 dopamine receptors modulates corticostriatal glutamatergic transmission. *Neuropharmacology* 2007; 53: 783-9.